Cochrane
Library

Cochrane Database of Systematic Reviews

Pharmacological treatments for fatigue associated with palliative care (Review)

Mücke M, Mochamat M, Cuhls H, Peuckmann-Post V, Minton O, Stone P, Radbruch L

Mücke M, Mochamat m, Cuhls H, Peuckmann-Post V, Minton O, Stone P, Radbruch L.
Pharmacological treatments for fatigue associated with palliative care.
Cochrane Database of Systematic Reviews 2015, Issue 5. Art. No.: CD006788.
DOI: [10.1002/14651858.CD006788.pub3](https://doi.org/10.1002/14651858.CD006788.pub3).

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
BACKGROUND	3
OBJECTIVES	4
METHODS	4
RESULTS	7
Figure 1.	8
Figure 2.	9
Figure 3.	10
Figure 4.	13
Figure 5.	14
Figure 6.	14
DISCUSSION	14
AUTHORS' CONCLUSIONS	15
ACKNOWLEDGEMENTS	16
REFERENCES	17
CHARACTERISTICS OF STUDIES	24
DATA AND ANALYSES	58
Analysis 1.1. Comparison 1: Methylphenidate versus placebo in cancer, Outcome 1: FACIT-F score change	58
Analysis 2.1. Comparison 2: Modafinil versus placebo in multiple sclerosis, Outcome 1: Fatigue score change	58
Analysis 3.1. Comparison 3: Pemoline versus placebo in multiple sclerosis, Outcome 1: Fatigue score change	59
ADDITIONAL TABLES	59
APPENDICES	64
FEEDBACK	73
WHAT'S NEW	74
HISTORY	74
CONTRIBUTIONS OF AUTHORS	75
DECLARATIONS OF INTEREST	75
SOURCES OF SUPPORT	75
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	75
NOTES	76
INDEX TERMS	76

[Intervention Review]

Pharmacological treatments for fatigue associated with palliative care

Martin Mücke^{1,2}, mochatat Mochamat^{1,3}, Henning Cuhls¹, Vera Peuckmann-Post⁴, Ollie Minton⁵, Patrick Stone⁶, Lukas Radbruch¹

¹Department of Palliative Medicine, University Hospital of Bonn, Bonn, Germany. ²Department of General Practice and Family Medicine, University Hospital of Bonn, Bonn, Germany. ³Department of Anesthesiology and Intensive Care, University of Diponegoro/Kariadi Hospital, Semarang, Indonesia. ⁴Palliative Medicine, Medical Faculty RWTH Aachen University, Aachen, Germany. ⁵Division of Mental Health, St George's University of London, London, UK. ⁶Marie Curie Palliative Care Research Department, Division of Psychiatry, University College London, London, UK

Contact address: Martin Mücke, martin.muecke@ukb.uni-bonn.de.

Editorial group: Cochrane Pain, Palliative and Supportive Care Group.

Publication status and date: Stable (no update expected for reasons given in 'What's new'), published in Issue 4, 2020.

Citation: Mücke M, Mochamat m, Cuhls H, Peuckmann-Post V, Minton O, Stone P, Radbruch L. Pharmacological treatments for fatigue associated with palliative care. *Cochrane Database of Systematic Reviews* 2015, Issue 5. Art. No.: CD006788. DOI: [10.1002/14651858.CD006788.pub3](https://doi.org/10.1002/14651858.CD006788.pub3).

Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

This review updates the original review, 'Pharmacological treatments for fatigue associated with palliative care' and also incorporates the review 'Drug therapy for the management of cancer-related fatigue'.

In healthy individuals, fatigue is a protective response to physical or mental stress, often relieved by rest. By contrast, in palliative care patients' fatigue can be severely debilitating and is often not counteracted with rest, thereby impacting daily activity and quality of life. Fatigue frequently occurs in patients with advanced disease (e.g. cancer-related fatigue) and modalities used to treat cancer can often contribute. Further complicating issues are the multidimensionality, subjective nature and lack of a consensus definition of fatigue. The pathophysiology is not fully understood and evidence-based treatment approaches are needed.

Objectives

To evaluate the efficacy of pharmacological treatments for fatigue in palliative care, with a focus on patients at an advanced stage of disease, including patients with cancer and other chronic diseases.

Search methods

For this update, we searched the Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE, PsycINFO and EMBASE, and a selection of cancer journals up to 28 April 2014. We searched the references of identified articles and contacted authors to obtain unreported data. To validate the search strategy we selected sentinel references.

Selection criteria

We considered randomised controlled trials (RCTs) concerning adult palliative care with a focus on pharmacological treatment of fatigue compared to placebo, application of two drugs, usual care or a non-pharmacological intervention. The primary outcome had to be non-specific fatigue (or related terms such as asthenia). We did not include studies on fatigue related to antineoplastic treatment (e.g. chemotherapy, radiotherapy, surgical intervention). We also included secondary outcomes that were assessed in fatigue-related studies (e.g. exhaustion, tiredness).

Data collection and analysis

Two review authors (MM and MC) independently assessed trial quality and extracted data. We screened the search results and included studies if they met the selection criteria. If we identified two or more studies that investigated a specific drug with the same dose in

a population with the same disease and using the same assessment instrument or scale, we conducted meta-analysis. In addition, we compared the type of drug investigated in specific populations, as well as the frequent adverse effects of fatigue treatment, by creating overview tables.

Main results

For this update, we screened 1645 publications of which 45 met the inclusion criteria (20 additional studies to the previous reviews). In total, we analysed data from 18 drugs and 4696 participants. There was a very high degree of statistical and clinical heterogeneity in the trials and we discuss the reasons for this in the review. There were some sources of potential bias in the included studies, including a lack of description of the methods of blinding and allocation concealment, and the small size of the study populations. We included studies investigating pemoline and modafinil in participants with multiple sclerosis (MS)-associated fatigue and methylphenidate in patients suffering from advanced cancer and fatigue in meta-analysis. Treatment results pointed to weak and inconclusive evidence for the efficacy of amantadine, pemoline and modafinil in multiple sclerosis and for carnitine and donepezil in cancer-related fatigue. Methylphenidate and pemoline seem to be effective in patients with HIV, but this is based only on one study per intervention, with only a moderate number of participants in each study. Meta-analysis shows an estimated superior effect for methylphenidate in cancer-related fatigue (standardised mean difference (SMD) 0.49, 95% confidence interval (CI) 0.15 to 0.83). Therapeutic effects could not be described for dexamphetamine, paroxetine or testosterone. There were a variety of results for the secondary outcomes in some studies. Most studies had low participant numbers and were heterogeneous. In general, adverse reactions were mild and had little or no impact.

Authors' conclusions

Based on limited evidence, we cannot recommend a specific drug for the treatment of fatigue in palliative care patients. Fatigue research in palliative care seems to focus on modafinil and methylphenidate, which may be beneficial for the treatment of fatigue associated with palliative care although further research about their efficacy is needed. Dexamethasone, methylprednisolone, acetylsalicylic acid, armodafinil, amantadine and L-carnitine should be further examined. Consensus is needed regarding fatigue outcome parameters for clinical trials.

PLAIN LANGUAGE SUMMARY

Pharmacological treatments for fatigue associated with advanced disease

In an advanced disease such as cancer, fatigue can be described as tiredness, weakness or lack of energy. Fatigue can affect daily activity and quality of life, and it is frequently reported by palliative care patients. The underlying causes of fatigue are not very well understood and fatigue is difficult to treat.

We searched the literature in April 2014 and found 45 randomised controlled trials for this update of the review. We analysed data from 4696 participants who received treatment for their fatigue. The trials dealt with neurological diseases (such as multiple sclerosis (753 participants), post-polio syndrome (58) and Parkinson's disease (19)), different types of cancer (3223), HIV/AIDS (514), end-stage renal disease (56), multi-type advanced disease in hospice patients (30), amyotrophic lateral sclerosis (28) and end-stage chronic lung disease (15).

There was weak evidence for the efficacy of amantadine, pemoline and modafinil in reducing fatigue in patients with multiple sclerosis. There was also weak evidence for the efficacy of carnitine and donepezil for cancer-related fatigue. One small trial showed that people with HIV/AIDS and fatigue seemed to benefit from treatment with methylphenidate or pemoline. There was some low-quality evidence from small trials that methylphenidate, a stimulant drug that improves concentration, is effective for the management of cancer-related fatigue. There was no information about dexamphetamine, paroxetine or testosterone.

Previous studies have shown that erythropoietin and darbepoetin, drugs that improve anaemia (a shortage of red cells or haemoglobin in the blood), are also effective for cancer-related fatigue. However, due to safety concerns and side effects shown by more recent studies, erythropoietin and darbepoetin should no longer be used. Therefore, we excluded these drugs from this review update.

Overall, most side effects of the investigated drugs seemed to be mild.

Based on limited evidence from small studies, the evidence does not support the use of a specific drug for the treatment of fatigue in palliative care. Future trials should measure fatigue in advanced disease using comparable and standardised measures.

BACKGROUND

This review is not only an update of a previously published review in the Cochrane Database of Systematic Reviews (Peuckmann-Post 2011). We have also conducted it with a new search strategy to incorporate another review on drug therapy for the management of cancer-related fatigue (Minton 2010), in order to increase its scope.

In healthy individuals, fatigue serves as a protective response to physical or mental stress. By contrast, in patients with chronic disease, fatigue can be severely debilitating and thereby have an impact on quality of life and daily activities (Morrow 2005). Fatigue is a common symptom in palliative care patients and virtually every intervention used to treat cancer, as well as the primary disease itself, may cause or contribute to fatigue. In a study of 1000 patients in an American palliative care programme, fatigue, weakness and lack of energy were three of the five most frequently reported symptoms with a prevalence of 84%, 66% and 61%, respectively (Walsh 2000). Fatigue is also commonly reported in non-cancer patients with progressive life-threatening diseases, such as multiple sclerosis and amyotrophic lateral sclerosis (progressive degeneration of motor neurons leading to cumulative paralysis), as well as chronic heart, kidney or lung diseases (Jhamb 2013; Tang 2010). More than half of patients with multiple sclerosis describe fatigue as one of their most troubling symptoms (Bakshi 2003; Krupp 2003; NMSS 2002). Fatigue has also been reported by the majority of patients with chronic obstructive pulmonary disease (COPD) (Elkington 2005; Stridsman 2014; Trendall 2001) and heart failure (Goodlin 2005). Approximately half of HIV patients suffer from fatigue (Breitbart 1998; Norval 2004). High levels of tumour necrosis factor (TNF) and interleukin-1 from HIV infections may cause fatigue (Darko 1995). Cancer-related fatigue is one of the most common symptoms experienced by cancer patients (Cuhls 2014; Morrow 2003). It can be problematic at the time of diagnosis, during and after treatment and in patients with advanced disease (Cuhls 2014; Morrow 2003). Most studies have reported prevalence figures in excess of 60% (Stone 2002). The subjective sensations attributed to cancer-related fatigue are characterised by a pervasive and persistent sense of tiredness, which is not relieved by sleep or rest. Several drugs, such as the new anti-neoplastic therapies, may be associated with novel causes of secondary fatigue. Drugs regularly used in palliative care have sedative properties, for example opioid analgesics, benzodiazepines, antidepressants or anticonvulsants can cause fatigue load (EAPC 2008).

Description of the condition

The pathophysiology of fatigue in palliative care patients is not fully understood. 'Primary fatigue' has been said to be related to the high cytokine load (release of high amounts of cytokines from the tumour or antineoplastic therapy). Associated disease-related symptoms, such as sleep disturbances, infections, malnutrition, hypothyroidism and anaemia, may also account for fatigue and may be termed 'secondary fatigue'. Synonyms for fatigue are asthenia, neuromuscular weakness and tiredness. There seems to be a considerable overlap between fatigue and depression. Weakness and tiredness are among the predominant symptoms of depression and feeling depressed is often part of the affective dimension of fatigue. However, there are some symptoms that are associated with depression (such as sustained feelings of worthlessness, recurrent thoughts of death) and some symptoms

that are considered specific to fatigue (such as post-exertional malaise).

Different definitions have been proposed for fatigue. In partnership with the American Cancer Society, the National Comprehensive Cancer Network (NCCN) defines cancer-related fatigue as "a distressing persistent, subjective sense of tiredness or exhaustion related to cancer or cancer treatment that is not proportional to recent activity and interferes with usual functioning" (NCCN 2014). A similar definition has been used for multiple sclerosis: "a subjective lack of physical and/or mental energy that is perceived by the individual or caregiver to interfere with usual and desired activities" (NMSS 2002). The Fatigue Coalition has suggested the use of the International Classification of Diseases – 10 (ICD-10) criteria for the definition of cancer-related fatigue, which is "significant fatigue, diminished energy or an increased need to rest, disproportionate to any recent change in activity level" and has to be present every day or nearly every day for two consecutive weeks out of the last month (Cella 2001). Five out of 10 additional symptoms, such as generalised weakness, diminished concentration or unrestorative sleep, are required for the diagnosis. However, the symptom thresholds and time span have been chosen arbitrarily (Cella 1998), and this has been criticised. Considering the lack of an internationally acknowledged definition, we identified and chose the following working definition for this review: "Fatigue is a subjective feeling of tiredness, weakness, or lack of energy"; this definition has been suggested by an expert working group of the European Association of Palliative Care (EAPC 2008).

Description of the intervention

Assessment of fatigue will depend on subjective self evaluation by the patient, substituted by caregiver or medical staff estimations only where self assessment is not possible. Single-item scales (e.g. 'do you get tired for no reason?') have been proposed and a multitude of checklists and questionnaires with multiple dimensions (such as physical, affective and cognitive) have been validated (Dittner 2004).

Lack of consensus on the definition of fatigue and its subjective and multidimensional nature, as well as culture and language differences, have challenged research approaches to fatigue. Further, standard research studies, such as double-blinded randomised controlled trials (RCTs), can often not be performed in this debilitated patient population due to fluctuating symptom intensity, declining performance status, rapid disease trajectories and short prognosis, resulting in weak evidence for treatment strategies. Several instruments exist to evaluate fatigue, but no consensus on significant cut-off levels has yet been achieved to establish a model of clinically meaningful improvement of fatigue. Only a few scales and instruments have been evaluated for different languages (Cantarero-Villanueva 2014; Kummer 2011). A structured approach with assessment and treatment steps is lacking. As a result, fatigue continues to be underestimated and undertreated. Consequently, a structured approach, including treatment options, is needed.

Treatment options for fatigue should address the causal mechanism if possible. However, the underlying mechanism of action is often not known and may be complex. Most patients will require symptomatic treatment of fatigue with pharmacological and non-pharmacological therapies. Non-pharmacological treatment options include patient education with

provision of information on fatigue and its treatment, keeping a diary, energy expenditure planning and physical exercise (review in [Mock 2004](#); [NCCN 2014](#); [Schmitz 2005](#)). Recent studies support the use of resistance training or 'anabolic' exercise ([Galvao 2005](#)). Most patients will try to counteract exhaustion and fatigue with prolonged periods of rest ([Richardson 1997](#)). However, rest will often not restore energy and persistent reduction of physical activity may even promote fatigue ([Evans 2007](#)).

Pharmacological treatment of fatigue may work through interaction with cytokine load and the patient's host reaction to the underlying disease, restoring peripheral energy depletion, or by treating metabolic disorders and supplementing other apparent physiological deficiencies such as decreased haemoglobin concentration ([Morrow 2005](#)). There is a growing body of evidence that gives examples of effective pharmacological treatments for fatigue ([Barak 2014](#); [Lawrence 2004](#); [Morrow 2005](#); [Patrick 2004](#); [Rao 2004](#); [Rosenberg 2005](#); [Wagner 2004](#); [Yennurajalingam 2013](#); [Zifko 2004](#)). Drug treatment for non-specific fatigue will be required by many patients in addition to specific measures against deficiencies or comorbid conditions. Concerning drug-induced fatigue, for example with opioid treatment, symptomatic pharmacological treatment of fatigue may functionally counteract this adverse effect, probably by enhancing excitatory mechanisms as observed following treatment with amphetamines ([Sood 2006](#)).

Why it is important to do this review

The latest systematic review about pharmacological treatments for fatigue associated with palliative care was published several years ago ([Peuckmann-Post 2011](#)). Other systematic reviews have covered the use of particular drugs for fatigue in multiple sclerosis, such as amantadine ([Pucci 2007](#)), and carnitine ([Tejani 2012](#)). Treatment of fatigue in cancer patients, including erythropoietic agents, was the subject of another Cochrane review with meta-analysis ([Minton 2010](#)). In contrast to this meta-analysis our review aims to cover a broader scope of palliative care patients with cancer and non-cancer diseases, but it is restricted to advanced-stage diseases. We have incorporated the main topic of [Minton 2010](#) into this review, so that cancer-related fatigue will be reflected.

OBJECTIVES

To evaluate the efficacy of pharmacological treatments for fatigue in palliative care, with a focus on patients at an advanced stage of disease, including patients with cancer and other chronic diseases.

METHODS

Criteria for considering studies for this review

Types of studies

We only included randomised controlled trials (RCTs). We considered full reports concerning fatigue in palliative care with a focus on pharmacological treatment. The primary outcome of these studies had to be fatigue (or related terms such as asthenia). We searched for diseases requiring palliative care or diseases at an advanced, life-threatening stage.

Types of participants

- Age 18 years or more.
- Participants of both sexes.

- Palliative care patients with fatigue, i.e. patients with an incurable disease (terminal illness) such as advanced cancer, HIV/AIDS, multiple sclerosis, amyotrophic lateral sclerosis, or cardiac, lung or kidney failure. Participants could receive anticancer treatment. We considered terminal illness to be when the estimated life expectancy is six months or less, under the assumption that the disease will run its normal course.

Types of interventions

- Identified studies had to evaluate and report the effect of pharmacological treatment on fatigue with the following drugs: psychostimulants (amphetamines, modafinil, armodafinil, methylphenidate, pemoline), amantadine, corticosteroids (dexamethasone, prednisone, methylprednisolone), donepezil, antidepressants such as selective serotonin reuptake inhibitors (SSRIs; paroxetine), acetylsalicylic acid, megestrol acetate, alfacalcidol and acetyl-L-carnitine.
- If we identified further agents used for the treatment of unspecific fatigue, we added these studies.
- Studies should compare fatigue with drug treatment versus no drug treatment or versus alternative drug treatment, or both.
- We did not include studies on the pharmacological treatment of fatigue with a primary target of clinical conditions such as depression or anxiety.
- We included antidepressants only if used for the treatment of fatigue as the primary outcome.
- We did not focus on physiological deficiencies such as lack of haemoglobin (erythropoietic agents, blood transfusion), nor did we focus on drugs targeting specific cytokines, e.g. for reduction of tumour necrosis factor alpha, as these treatments target specific aetiologies of fatigue. Erythropoietic agents have been covered in the review by Minton et al ([Minton 2010](#)).
- We did not include studies comparing different types of cancer-modifying treatment and the effect on prognosis and quality of life. We also excluded those studies which did not focus on pharmacological treatment.
- We did not include studies on fatigue related to antineoplastic treatment (e.g. chemotherapy, radiotherapy, surgical intervention).

Types of outcome measures

Primary outcomes

1. Patient-reported fatigue (self reported measures or validated self assessment tools, or both), substituted by caregiver or medical staff estimations only where self assessment was not possible, and measurement using reliable and valid assessment instruments (single-item scales or questionnaire instruments).
2. Improvement of fatigue. Since no gold standard for the treatment or improvement of fatigue exists, we suggested an improvement of fatigue intensity by 33% related to the range of the assessment instrument to be clinically significant. This is congruent with the improvement of pain intensity, where a 33% reduction has been described as significant from the patient's point of view ([Farrar 2003](#)).

Secondary outcomes

1. Asthenia (lack of strength) assessed with quality of life instruments such as SF-36.

2. Weakness assessed with scales, e.g. visual analogue scale - fatigue (VAS-F).
3. Tiredness, sedation assessed with scales such as the Chalder Fatigue Scale (CFS) or Center for Epidemiologic Studies Depression (CESD) scale.
4. Exhaustion, assessed with e.g. the European Organization for Research and Treatment of Cancer Quality of Life Questionnaire (EORTC-QLQ-C30).
5. Treatment-related burden: adverse events (including cardiac arrhythmia and thromboembolic events), morbidity or mortality measured as percentages of participants, adverse events also measured as a percentage of participants with moderate/severe intensity, probability of a causal relationship between adverse events and treatment.

Search methods for identification of studies

For this update, we have re-engineered the search strategy (filter) of the previous review (Peuckmann-Post 2011) to facilitate the combination with another review (Minton 2010). To identify studies for inclusion in this updated review, we developed a detailed search strategy for each electronic database and other resources. To validate the search strategy, we selected sentinel references. There were no language or date restrictions.

Electronic searches

We searched the following electronic databases:

- Cochrane Central Register of Controlled Trials (CENTRAL) (2014, Issue 3); search strategy as detailed in [Appendix 1](#);
- MEDLINE (OVID) from inception to 28 April 2014; search strategy as detailed in [Appendix 2](#);
- EMBASE (OVID) from inception to 28 April 2014; search strategy as detailed in [Appendix 3](#);
- PsycINFO (OVID) from inception to 28 April 2014; search strategy as detailed in [Appendix 4](#).

Searching other resources

We screened the reference lists of identified articles for additional studies.

We handsearched standard textbooks on palliative medicine (*Oxford Textbook of Palliative Medicine*, Oxford; *Textbook of Nursing, Textbook of Palliative Medicine*).

We also obtained unpublished literature through searches of conference proceedings, such as all meetings of the American Society of Clinical Oncology (ASCO) from 2000 to 2013, the 2013 meeting of the European Cancer Congress (ECCO) and the European Association of Palliative Care (EAPC) databases of all abstracts registered online from 2003 to 2014.

We contacted experts in the field of fatigue in palliative care in order to identify research awaiting publication.

Data collection and analysis

Selection of studies

We retrieved in full all studies in which the abstract refers to drug intervention aimed at treating fatigue in palliative care. Eligible

studies had to define fatigue as a primary outcome and at least one treatment arm had to be a drug intervention.

Data extraction and management

We organised data using the software Review Manager 5.3 (RevMan 2014). Two review authors extracted data (MM and MC) using a standard data extraction form. Two authors (MM and MC) reviewed the data from the included studies and two other authors (LR and HC) cross-checked a sub-sample. We resolved disagreement by consensus.

Assessment of risk of bias in included studies

Two authors (MM and MC) independently assessed risk of bias for each study, using the criteria outlined in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011) and adapted from those used by the Cochrane Pregnancy and Childbirth Group, with any disagreements resolved by discussion or by involving other review authors (LR, HC). We assessed the following for each study:

Random sequence generation (checking for possible selection bias)

We assessed the method used to generate the allocation sequence as: low risk of bias (any truly random process, e.g. random number table; computer random number generator); unclear risk of bias (method used to generate sequence not clearly stated). We carefully considered studies using a non-random process.

Allocation concealment (checking for possible selection bias)

The method used to conceal allocation to interventions prior to assignment determines whether intervention allocation could have been foreseen in advance of, or during recruitment, or changed after assignment. We assessed the methods as: low risk of bias (e.g. telephone or central randomisation; consecutively numbered, sealed, opaque envelopes); unclear risk of bias (method not clearly stated). We excluded studies that did not conceal allocation (e.g. open list).

Blinding of outcome assessment (checking for possible detection bias)

We assessed the methods used to blind study participants and outcome assessors from the knowledge of which intervention a participant received. We assessed the methods as: low risk of bias (study states that it was blinded and describes the method used to achieve blinding, e.g. identical tablets, matched in appearance and smell); unclear risk of bias (study states that it was blinded but does not provide an adequate description of how this was achieved). We excluded studies that were not double-blind.

Incomplete outcome data (checking for possible attrition bias due to the amount, nature and handling of incomplete outcome data)

We assessed the methods used to deal with incomplete data as: low risk (less than 10% of participants did not complete the study and/or used 'baseline observation carried forward' analysis); unclear risk of bias (used 'last observation carried forward' analysis); high risk of bias (used 'completer' analysis).

Size of study (checking for possible biases confounded by small size)

We assessed studies as being at low risk of bias (200 participants or more per treatment arm); unclear risk of bias (50 to 199 participants per treatment arm); high risk of bias (fewer than 50 participants per treatment arm).

Measures of treatment effect

We intended to evaluate the methodological quality of each study, as well as the evidence for drug interventions to treat fatigue, using the GRADE system (Atkins 2004). We aimed to further evaluate studies by setting 'clinically relevant improvement' equal to improvement by one-third compared to baseline fatigue intensity and to calculate a number needed to treat to benefit (NNTB) accordingly (NNTB to achieve at least a 33% improvement of fatigue intensity compared to the baseline fatigue level). We aimed to calculate numbers needed to treat to harm (NNTH) for adverse events with moderate or severe intensity.

If calculation of the NNTB was not possible, 'clinically relevant improvement' would be defined as a change of 5% or more of the primary outcome instrument used in the study. This would be in accordance with the study of Cella et al, identifying changes between 3.7% and 5.8% of the aggregated summary scores as 'clinically important' (Cella 2002).

In addition, for each identified study, we aimed to extract:

- number of participants in each arm;
- type of control group;
- quality of the study (randomisation, blinding, per protocol analysis, intention-to-treat (ITT) analysis, number of withdrawals described);
- demographic characteristics, including age and sex;
- type of primary disease;
- type and stage of treatment, if applicable;
- type of drug used for the pharmacological intervention;
- duration and pharmacological regimen of drug treatment with the drug of interest;
- outcome measures employed, including means and standard deviations.

We documented outcomes of fatigue in different ways. If we identified two or more studies that investigated a specific drug with the same dose in a population with the same disease and same assessment instrument or scale, we conducted a meta-analysis. We aimed to calculate the standardised mean difference (SMD) in fatigue intensity, stating whether it had reached a potentially significant cut-off level and we aimed to compare these data to other interventions and control groups, wherever available. Wherever appropriate, we calculated standard deviations (SD) for standardisation of the mean difference (MD) as the square root of the average of the variances before and after the intervention. Similarly, if only 95% confidence intervals (CI) were described, we converted these values to SDs by dividing the difference of the CI and mean by a factor of 1.96.

A NNTB of 50% reduction has been suggested as significant for an exercise intervention in cancer-related fatigue (Cramp 2012). Since only limited information on the cut-off level for clinically significant fatigue has been identified to date, we suggested a 33% improvement as a basis for NNTB calculation, since palliative care patients are more debilitated than the population that was

examined in the Cramp 2012 review. A 33% reduction of pain intensity has been described as significant from the patients' point of view in studies on breakthrough pain management (Farrar 2003).

Unit of analysis issues

The unit of randomisation was the individual patient.

Dealing with missing data

We contacted the original investigators to request missing data. We used intention-to-treat (ITT) analysis where possible. The ITT population consisted of participants who were randomised, took the assigned study medication and provided at least one post-baseline assessment. We assigned missing participants zero improvement (baseline observation carried forward (BOCF), where this could be done. We were aware that imputation methods might be problematic and examined trial reports for information about them.

Assessment of heterogeneity

We explored the homogeneity of the results of the various endpoints of interest using I^2 statistic values. We regarded heterogeneity in the results as a result of many potential factors (postulated a priori), and we made efforts to identify subgroups for sensitivity analysis. We undertook meta-analysis. As a result of high statistical heterogeneity, we used a random-effects model for analysis. If possible, and where applicable, we conducted subgroup analyses to explore possible sources of heterogeneity due to participants, interventions or methods.

Potential sources of heterogeneity:

- disease entities;
- performance status;
- quality of studies;
- medication dose and frequency;
- duration of treatment;
- duration of follow-up;
- rate of attrition;
- outcome measures used;
- case mix/stage of disease accessed.

Assessment of reporting biases

To decrease the influence of potential publication bias, we conducted manual and electronic searches of multiple databases, without imposing any language restriction, to check for published or registered study protocols and to verify whether results from these studies have been published subsequently. We contacted the trial authors by email if there was insufficient information to assess reporting bias. We also contacted the authors to clarify information, if there were mismatches between study protocols and reports.

Data synthesis

For data synthesis, we used Review Manager 5.3 (RevMan 2014), as provided by The Cochrane Collaboration. We grouped meta-analyses of the data from all included studies using the I^2 statistic. If the I^2 value was greater than or equal to 50% (substantial or considerable heterogeneity), we used a random-effects model. If the I^2 value was less than 50%, we used a fixed-effect model. We reported the results from both models.

Subgroup analysis and investigation of heterogeneity

We intended to conduct a subgroup analysis where data were available. We performed separate analysis for different kinds of primary diseases (such as cancer or multiple sclerosis). If it was not possible to carry out a quantitative analysis, then we considered a qualitative review and a synthesis of the study results.

In this review, we performed subgroup analyses for the following:

- different types of drug treatment;
- different types of disease;
- different types of assessment tools.

Sensitivity analysis

We carried out sensitivity analysis for the primary outcome measurements in order to explore effect size differences and the robustness of our conclusions. We planned sensitivity analysis determined a priori based on:

- studies without study limitations with regard to a) allocation concealment; b) blinding of participants and investigators; c) recruitment bias; d) baseline imbalance between groups; e) loss to follow-up of clusters; f) adequate analysis;
- method of analysis: results of a) studies using number of patients analysed; b) studies using number of patients randomised.

RESULTS

Description of studies

This review updates and combines two existing reviews: [Minton 2010](#) and [Peuckmann-Post 2011](#). The previous review, [Minton 2010](#), found 31 studies involving 7140 participants and the review [Peuckmann-Post 2011](#) identified 22 eligible studies with a total of 1632 participants. We used a new search strategy for this update, providing 1645 results that we screened. Most of these articles did not focus on fatigue as a primary outcome but on the drug treatment of underlying disease. The updated search strategy identified 20 additional studies suitable for inclusion. Only seven of the studies included in the [Minton 2010](#) review, but all 22 of the previously included studies in [Peuckmann-Post 2011](#), matched our results. The small number of matched study results between this updated review and the [Minton 2010](#) review is due to the fact that we did not include haemopoietic growth factors in our search strategy. Safety concerns have been raised regarding the erythropoiesis-stimulating agents (erythropoietin and darbepoetin) since the last publication of the original review, therefore the use of these drugs is no longer recommended in practice ([Bennet 2008](#); [Bohlius 2009](#); [EAPC 2008](#); [Glaspy 2009](#); [Glaspy 2010](#); [Tonelli 2009](#)). We found and removed 186 duplicate studies ([Figure 1](#)). We checked the retrieved articles against the inclusion criteria and included 45 studies. We identified no additional unpublished data by contacting experts in palliative care and we retrieved no additional studies from the handsearched reference lists or abstract databases on the Internet.

Figure 1. Study flow diagram.

Meta-analysis of data was possible for modafinil in multiple sclerosis (two studies, $n = 136$ participants), pemoline in multiple sclerosis (two studies, $n = 103$ participants) and methylphenidate in cancer patients (two studies, $n = 146$ participants).

See also [Characteristics of included studies](#) and [Characteristics of excluded studies](#).

Results of the search

We screened 1645 publications, of which 45 met the inclusion criteria and we included them for further analysis. In total, we analysed data from 18 drugs and 4696 participants. We used studies investigating pemoline and modafinil in participants with multiple sclerosis-associated fatigue and methylphenidate in participants suffering from cancer and fatigue for meta-analysis.

Included studies

See [Characteristics of included studies](#) table.

The 45 studies retrieved for the review yielded a broad spectrum of diseases associated with fatigue as well as different drugs ([Table 1](#)).

Fatigue was typically examined in association with diseases as follows: multiple sclerosis (13 studies), HIV/AIDS (including HIV and hypogonadism) (seven studies), cancer (including advanced cancer, cancer of different origins and brain tumours) (18 studies), post-polio (two studies), Parkinson's disease (one study), end-stage renal disease (ESRD) (one study), amyotrophic lateral sclerosis (one study), multi-type advanced disease (hospice patients) (one study) and end-stage COPD (one study).

Drug studied were as follows: amantadine (nine studies), pemoline (three studies), megestrol acetate (one study), methylphenidate (seven studies), dexamphetamine (two studies), paroxetine (two studies), acetyl-L-carnitine (four studies), testosterone (three studies), donepezil (one study), modafinil (eight studies), dexamethasone (one study), fluoxetine (two studies), alfalcidol (one study), armodafinil (one study), mistletoe extract PS76A2 (one study).

study), methylprednisolone (one study), medroxyprogesterone acetate (one study) and acetylsalicylic acid (two studies).

In contrast, [Morrow 2003](#) examined fatigue related to both cancer and treatment. [Wagner 2000](#) and [Rabkin 2004](#) examined fatigue in patients with depression, where it may be difficult to differentiate whether fatigue was primary or secondary to depression.

Most studies reported data from a relatively small number of participants: for the 45 studies included, only 17 reported a participant number greater than 100. Participant numbers, including controls, varied from six ([Rosenberg 1988](#)) to 544 ([Jean-Pierre 2010](#)). The total number of participants who completed the studies included in our analysis was 4696. Issues associated with inconclusive findings were as follows: differences in data reporting, heterogenous populations, inconsistent symptom assessment (the use of instrument differed greatly) and lack of a consistent definition for a clinically significant reduction in fatigue. Therefore it was not possible to calculate the NNTB or NNTH, nor could we calculate a 5% reduction of fatigue as consistently as intended and described above.

Some trials in this review only used single-item fatigue assessment. However, it is now considered by European palliative care experts that a single-item tool could be as good as a comprehensive assessment. Single-item assessments of fatigue may have drawbacks, but more complex assessment methods may also be problematic in patients with advanced disease as they require some cognitive capacity, which is often lacking in a fatigue patient group. Excluding these studies would cause a new bias, as would excluding studies in end-of-life care and focused on patients in oncological care. Hence, we included and analysed these trials in our review.

Excluded studies

The previous review, [Peuckmann-Post 2011](#), excluded 23 studies and the review by [Minton 2010](#) excluded 14 studies. For this update,

we had to exclude 49 studies in the final selection process after reading the manuscript. Reasons for exclusion were typically a lack of inclusion criteria match (e.g. fatigue defined as a secondary outcome; case series; not a RCT), investigation of fatigue attributed to narcotic-induced sedation, data were a subset of another study which had been already included in the analysis, or data were not from patients with advanced disease. We listed these studies with the reason for exclusion in the [Characteristics of excluded studies](#) table.

Risk of bias in included studies

Some studies specifically described the randomisation and blinding procedure and we therefore considered them to have a minimal risk of bias concerning allocation concealment (e.g. [Cruciani 2012](#); [Roth 2010](#)). However, some studies did not specify these procedures, but used the terminology "double-blind, randomised controlled trial", allowing the assumption of a low risk of bias (e.g. [Barak 2014](#); [Lange 2009](#)).

Considering the relatively small number of participants in most of the included studies, as well as the variety of instruments used, the results have to be interpreted with caution. The small numbers of participants may be a reason for a lack of stratification (for example, for relevant health conditions such as depression or sociodemographic data such as age or sex), which may alter the outcome as demonstrated by some investigators ([Eriksen 2003](#); [Tomassini 2004](#)).

We assessed each study using the Cochrane 'Risk of bias' tool. The findings are presented in the 'Risk of bias' graph ([Figure 2](#)), which reviews the authors' judgements about each risk of bias item shown as percentages across all included studies, and the 'Risk of bias' summary ([Figure 3](#)), which reviews the authors' judgements about each risk of bias item for each included study.

Figure 2. 'Risk of bias' graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.

Figure 3. 'Risk of bias' summary: review authors' judgements about risk of bias items for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of outcome assessment (detection bias): All outcomes	Incomplete outcome data (attrition bias): All outcomes	Size of study
Ashtari 2009	?	?	+	+	-
Auret 2009	?	?	+	+	-
Barak 2014	?	?	?	+	?
Brass 2001	?	?	?	+	-
Breitbart 2001	+	?	+	+	-
Bruera 2006	+	+	?	+	?
Bruera 2007	+	+	?	+	?
Butler 2007	?	?	+	?	-
Canadian MSRG 1987	?	?	?	+	?
Cohen 1989	?	?	?	+	-
Cruciani 2009	+	?	+	+	-
Cruciani 2012	+	?	+	+	?
Della Cuna 1989	?	?	?	+	?
Escalante 2014	?	?	?	+	-
Fisch 2003	+	?	?	+	?
Jean-Pierre 2010	+	?	?	+	+
Kerr 2012	+	?	?	+	-
Knapp 2008	+	?	?	+	-
Krupp 1995	?	?	+	?	-
Lacasse 2004	?	+	?	+	-
Lange 2009	?	?	?	+	-
Lou 2009	?	?	+	+	-
Moraska 2010	+	+	?	+	?
Morrow 2003	+	?	+	+	+
Murray 1985	?	?	?	+	-
Rabkin 2000	+	?	+	+	-

Figure 3. (Continued)

Murray 1985	?	?	?	+	+
Rabkin 2000	+	?	+	+	+
Rabkin 2004	+	?	+	+	+
Rabkin 2009	+	+	+	+	+
Rabkin 2010	+	?	+	+	?
Rabkin 2011	+	?	+	+	+
Rosenberg 1988	?	?	?	+	+
Roth 2010	?	?	+	+	+
Semiglazov 2006	+	+	+	+	?
Shaygannejad 2012	+	+	?	+	+
Simons 1996	+	?	?	+	?
Spathis 2014	+	+	+	+	?
Stankoff 2005	?	?	?	+	?
Stein 1995	?	?	?	+	+
Tomassini 2004	?	?	?	+	+
Vasconcelos 2007	+	+	+	+	+
Wagner 2000	?	?	?	+	+
Weinshenker 1992	?	?	?	?	+
Westman 1999	+	?	+	+	?
Wingerchuk 2005	+	+	+	+	+
Yennurajalingam 2013	?	?	?	?	+

Allocation

All studies reported randomisation, but no more than 23 included a proper description of the methods used (Breitbart 2001; Bruera 2006; Bruera 2007; Cruciani 2009; Cruciani 2012; Fisch 2003; Jean-Pierre 2010; Kerr 2012; Knapp 2008; Moraska 2010; Morrow 2003; Rabkin 2000; Rabkin 2004; Rabkin 2009; Rabkin 2010; Rabkin 2011; Semiglazov 2006; Shaygannejad 2012; Simons 1996; Spathis 2014; Vasconcelos 2007; Westman 1999; Wingerchuk 2005). Only nine described the method used to conceal the allocation appropriately (Bruera 2006; Bruera 2007; Lacasse 2004; Moraska 2010; Rabkin 2009; Semiglazov 2006; Shaygannejad 2012; Spathis 2014; Vasconcelos 2007; Wingerchuk 2005).

Blinding

Twenty studies were double-blind (Ashtari 2009; Auret 2009; Breitbart 2001; Butler 2007; Cruciani 2009; Cruciani 2012; Krupp 1995; Lou 2009; Morrow 2003; Rabkin 2000; Rabkin 2004; Rabkin 2009; Rabkin 2010; Rabkin 2011; Roth 2010; Semiglazov 2006; Spathis 2014; Vasconcelos 2007; Westman 1999; Wingerchuk 2005). The remainder did not provide sufficient information to assess risk of bias (unclear risk).

Incomplete outcome data

There were only four studies lacking sufficient information to assess risk (unclear risk of bias) (Butler 2007; Krupp 1995; Weinshenker 1992; Yennurajalingam 2013). We judged the remaining 41 studies to meet the criteria for low risk of bias.

Other potential sources of bias

Treatment group size was an issue. Small studies are thought to be at increased risk of bias, probably because the conduct of small studies is more likely to be less rigorous, allowing critical criteria to be compromised. Only two of the treatment groups in this review were large enough to give a low risk of bias (Jean-Pierre 2010; Morrow 2003). We judged 14 studies to have an unclear risk (Barak 2014; Bruera 2006; Bruera 2007; Canadian MSRG 1987; Cruciani 2012; Della Cuna 1989; Fisch 2003; Moraska 2010; Rabkin 2010; Semiglazov 2006; Simons 1996; Spathis 2014; Stankoff 2005; Westman 1999). We judged the remainder to have a high risk of bias due to size.

Effects of interventions

Most studies reported some benefit of the active treatment. However, there was often a substantial and very similar placebo effect. Further, reviewing only fatigue-specific instruments (e.g. the Chalder Fatigue Scale (CFS), Fatigue Severity Scale (FSS), visual analogue scale (VAS) for fatigue), these benefits could not be confirmed in many cases. See also: [Types of outcome measures](#). In general, adverse reactions were mild and had little or no impact (Table 2).

Acetyl-L-carnitine versus placebo

Acetyl-L-carnitine was compared with placebo in a study of 29 participants with cancer-related fatigue, with no significant effect (Cruciani 2009). FACT-fatigue mean scores were 15.7 (SD 10.6) at baseline and 22.2 (10.4) at week two (P value = 0.97). A recent study of 209 participants did not show statistically significant improvement in fatigue compared to placebo (Cruciani 2012).

The primary outcome, fatigue, measured using the Brief Fatigue Inventory (BFI), improved in both arms in comparison to baseline (L-carnitine: -0.96, 95% confidence interval (CI) -1.32 to -0.60; placebo: -1.11, 95% CI -1.44 to -0.78). There were no statistically significant differences between the arms (P value = 0.57), while the secondary outcomes, including fatigue measured on the Functional Assessment for Chronic Illness Therapy - Fatigue (FACIT-F) scale, did not show a significant difference between arms, with a change in mean of 31.4 (standard deviation (SD) 9.21; L-carnitine) versus 23.67 (SD 11.24; placebo) (P value = 0.61).

Acetyl-L-carnitine versus amantadine

In another study of 36 participants, acetyl-L-carnitine was compared with amantadine in patients with multiple sclerosis (Tomassini 2004). Improvement assessed with the FSS was superior with L-carnitine compared with amantadine. In addition, L-carnitine was better tolerated than amantadine. However, differences in the absolute changes in the FSS did not reach statistical significance and the number of responders was similar for both drugs. There was no significant change for the secondary outcome of clinical scale scores (Fatigue Impact Scale, Beck Depression Inventory and Social Experience Checklist). The fatigue domain of the Kidney Disease Questionnaire (KDQ) also significantly improved after 12 weeks and 24 weeks of carnitine therapy compared with placebo in 56 patients with end-stage renal disease, but did not significantly affect the total score (Brass 2001).

Acetylsalicylic acid versus placebo

Wingerchuk 2005 was able to show significantly better relief of fatigue with 1300 mg/day of acetylsalicylic acid compared to placebo in 26 patients with multiple sclerosis. Wingerchuk et al used the Modified Fatigue Impact Scale (MFIS) and Kurtzke Expanded Disability Status Scale (KEDSS) as instruments. In another study of 52 patients with multiple sclerosis (Shaygannejad 2012), 500 mg/day showed a significant decrease in FSS scores. We did not perform meta-analysis due to the different doses and assessment instruments.

Alfacalcidol versus placebo

A recent study using alfacalcidol 1 µg/day compared with placebo was conducted to treat fatigue in 158 multiple sclerosis patients. The Fatigue Impact Scale (FIS) total score decreased significantly. 'RAYS' quality of life assessments also improved significantly in the psychological and social subscale (Barak 2014).

Amantadine versus placebo

Amantadine was used in seven studies of 370 participants with multiple sclerosis with heterogeneous outcomes, showing a tendency towards improved outcomes with amantadine with different fatigue instruments and scales (Ashtari 2009; Canadian MSRG 1987; Krupp 1995; Murray 1985; Rosenberg 1988; Shaygannejad 2012; Tomassini 2004). Amantadine was significantly better than placebo on the MS-Fatigue scale. One study of 25 participants used amantadine post-polio using the Fatigue Severity Scale (FSS) without proving efficacy (Stein 1995).

Amantadine versus other drugs

Krupp 1995 tested amantadine against pemoline (93 participants). No difference between amantadine and pemoline was seen on the FSS. On the other hand, Tomassini 2004 performed a

comparison of amantadine against acetyl-L-carnitine (ALCAR). ALCAR showed superior effect to amantadine. Shaygannejad 2012 tested amantadine against acetylsalicylic acid. Both groups showed a significant decrease in the FSS.

Armodafinil versus placebo

Armodafinil was tested in 70 patients with HIV to evaluate its efficacy and safety for fatigue and depressive symptoms. In intention-to-treat analyses, the fatigue response rate to armodafinil was 75% and to placebo 26% (Rabkin 2011). Armodafinil appeared to be effective in alleviating fatigue and was well tolerated. For the secondary endpoint depression, measured with the Hamilton Depression Rating Scale (HDRS) and Beck Depression Inventory (BDI), a significant effect was not shown. However, further multicentre studies with larger samples are needed.

Dexamethasone versus placebo

Dexamethasone 4 mg was compared with placebo in a recent study to treat fatigue in 84 participants with advanced cancer. The study showed that dexamethasone was significantly superior to placebo (Yennurajalingam 2013). However, there was no significant difference in the improvement of individual symptoms on the Edmonton Symptom Assessment Scale (ESAS), psychological distress, anxiety scores on the Hospital Anxiety and Depression Scale (HADS) or HADS depression scores in the dexamethasone group compared with placebo. Further dexamethasone studies are needed.

Dextroamphetamine versus placebo

Dextroamphetamine was compared with placebo to treat fatigue in a study of 39 cancer patients (Auret 2009) and in 22 patients with HIV (Wagner 2000). Neither study showed significant effects. The secondary outcome quality of life, which was measured using the McGill Quality of Life Questionnaire (MQOL), also showed no significant effect (Auret 2009).

Donepezil versus placebo

In a large study with 142 participants with cancer, donepezil significantly relieved fatigue intensity in both the intervention and control arms, with no significant difference as measured by the FACIT-F scale. Edmonton Symptom Assessment Scale (ESAS) and sleep pattern assessment were also used as supplemental scales. Neither showed significant effects. Thus, donepezil did not appear to be superior to placebo (Bruera 2007).

Fluoxetine versus placebo

Another study of 129 patients with multiple sclerosis showed that 20 mg of fluoxetine over 12 weeks appeared to be superior to placebo as measured by the change in Functional Assessment of Cancer Therapy - General (FACT-G) scores. The fluoxetine group also showed significant improvement on the depression scale (P value = 0.0005) compared with placebo, which was measured using the Brief Zung Self-Rating Depression Scale (BZSDS) (Fisch 2003).

Fluoxetine versus testosterone

Fluoxetine was inferior to testosterone in 90 patients with HIV and fatigue, while the difference in the effect on depression, as measured with the HDRS, was non-significant (P value = 0.38) (Rabkin 2004).

Medroxyprogesterone versus placebo

Medroxyprogesterone was tested in only one study of 134 participants using the European Organization for Research and Treatment of Cancer Quality of Life Questionnaire (EORTC-QLQ-C30) (Simons 1996). The use of 500 mg twice a day over 12 weeks showed no significant effect.

Megestrol acetate versus placebo

Megestrol acetate was tested in 255 patients with cancer. The administration of 320 mg/day for 12 weeks was performed to investigate the effect of megestrol acetate on quality of life, appetite, weight and survival. However, megestrol acetate does not appear to improve global quality of life as measured by the EORTC QLQ-C30 (Westman 1999).

Methylphenidate versus placebo

Five studies tested methylphenidate in 318 cancer patients (Bruera 2006; Butler 2007; Escalante 2014; Moraska 2010; Roth 2010). Meta-analysis was possible only for two studies (Bruera 2006; Butler 2007), which used FACIT-F as the assessment tool in fatigue, comparing methylphenidate with placebo. The studies showed a slightly superior effect of methylphenidate compared to placebo (standardised mean difference (SMD) 0.49, 95% confidence interval (CI) 0.15 to 0.83; Analysis 1.1; Figure 4). Moraska 2010 used the Brief Fatigue Inventory (BFI) instrument, and could not show significant effects of methylphenidate (18 mg to 54 mg) compared with placebo. Escalante 2014 used a stable dose (18 mg) for two weeks. They reported that methylphenidate improved cancer-related fatigue according to the BFI scores. The smallest dose was used by Roth 2010 (5 mg to 30 mg). In this study, clinically significant improvement was seen on the BFI.

Figure 4. Forest plot of comparison: 1 Methylphenidate vs Placebo in cancer, outcome: 1.1 FACIT-F score change.

The other study of 30 participants using methylphenidate concerned the treatment of fatigue in multi-type advanced disease (hospice) patients (Kerr 2012). The study showed that methylphenidate was superior to placebo, although the improvement of the fatigue score was dose-dependent, as measured by Piper Fatigue Scale, Visual Analogue Scale for Fatigue and ESAS. However, secondary outcomes of depressive symptoms, measured with the revised Beck Depression Inventory-II (BDI-II), the ESAS depression score and the Center for Epidemiologic Studies Depression Scale (CESD), showed no statistically significant differences.

Methylphenidate was also investigated in 209 HIV patients and compared with pemoline and placebo (Breitbart 2001). This study demonstrated a significantly higher number of responders using methylphenidate compared with placebo.

Methylprednisolone versus placebo

Methylprednisolone (125 mg/day for eight weeks) was used with significant effect in only one study of 403 participants with cancer-related fatigue (Della Cuna 1989).

Mistletoe extract PS76A2 versus placebo

Mistletoe extract was tested in 337 patients with breast cancer (Semiglazov 2006). It showed a significant positive effect using the FACT-G.

Modafinil versus placebo

Modafinil has been explored for the treatment of fatigue in two studies of 704 patients with cancer (Jean-Pierre 2010; Spathis 2014). The Jean-Pierre 2010 study showed a significant interaction between treatment condition (modafinil 200 mg/day) and baseline fatigue, where patients with severe baseline fatigue benefited from modafinil and patients with mild or moderate fatigue did not. A recent study demonstrated that both modafinil (100 to 200 mg/day) and placebo led to a clinically significant improvement in FACIT-F scores (Spathis 2014). However, there was no significant difference between placebo and modafinil.

Though modafinil has been tested in several studies in multiple sclerosis, we found only two controlled trials (Lange 2009; Stankoff 2005). The Stankoff 2005 study included a larger number of participants (n = 115), but failed to demonstrate the superiority of modafinil versus placebo. Lange 2009 included only 21 participants and stated that they used a subpopulation of a larger trial. Thus, although there was a clear beneficial effect of modafinil, this result must be interpreted with caution due to the small participant numbers. Meta-analysis of these two studies also failed to demonstrate a significant effect, with a SMD of -0.14 (95% CI -0.48 to 0.21; Analysis 2.1; Figure 5).

Figure 5. Forest plot of comparison: 4 Modafinil in multiple sclerosis, outcome: 4.1 Modafinil.

Modafinil was also tested for the treatment of fatigue in 33 participants with post-polio syndrome (Vasconcelos 2007). However, in this study modafinil failed to demonstrate superiority over placebo.

There was also a trial using modafinil for fatigue in 105 HIV/AIDS patients (Rabkin 2010). In the intention-to-treat analyses, the fatigue response rate to modafinil was 73% and to placebo 28%; the attrition rate was 9%. Secondary endpoints (depression), including the Hamilton Depression Rating Scale (HDRS) and Beck Depression Inventory (BDI), did not show a significant effect.

Paroxetine versus placebo

Paroxetine was tested in a study of 479 patients with cancer (Morrow 2003), and a study of 15 patients with end-stage chronic obstructive pulmonary disease (COPD; Lacasse 2004). Neither study showed significant effects. However, the Center for Epidemiological Studies Depression (CESD) score, controlling baseline depression scores, confirmed that the dose of paroxetine provided was more effective than placebo in reducing depression (P value = 0.001).

Pemoline versus placebo

Pemoline was used for the treatment of fatigue in 41 participants with multiple sclerosis. However, in this study pemoline failed to demonstrate superiority over placebo (Weinshenker 1992).

Pemoline versus other drugs

Pemoline was tested in 93 patients with multiple sclerosis and severe fatigue (Krupp 1995). The drug did not appear to be superior to placebo. One study with HIV patients compared pemoline to methylphenidate and to a placebo (Breitbart 2001). The Breitbart 2001 study demonstrated a significantly higher number of responders with pemoline and methylphenidate compared with placebo.

Meta-analysis was possible for two studies in multiple sclerosis patients as they all included a control group with placebo and reported continuous outcome indicators (Krupp 1995; Weinshenker 1992). There was no superior effect of pemoline (SMD -0.02; 95% CI -0.12 to 0.08; Figure 6).

Figure 6. Forest plot of comparison: 3 Pemoline in multiple sclerosis, outcome: 3.1 Fatigue score change.

However, the US Food and Drug Administration (FDA) has decided to withdraw pemoline products (marketed as Cylert) due to the risk of liver toxicity, which outweighs the benefits of the drug. All manufacturers have agreed to stop the sale and marketing of pemoline (FDA 2005).

Testosterone versus placebo

Testosterone cypionate was investigated in 208 patients with HIV in three studies by the same workgroup (Knapp 2008; Rabkin 2000; Rabkin 2004). The older study included HIV patients with symptoms of hypogonadism and found that testosterone relieved these symptoms in patients with fatigue at baseline (Rabkin 2000). However, fatigue intensity on the CFS merely showed a positive trend, but no significant difference from placebo. The newer study did not show any significant benefit of testosterone compared to fluoxetine or placebo (Rabkin 2004). Only Knapp 2008 showed superiority of testosterone over placebo.

DISCUSSION

This review updates the original review 'Pharmacological treatments for fatigue associated with palliative care' (Peuckmann-Post 2011) and also incorporates 'Drug therapy for the management of cancer-related fatigue' (Minton 2010). Peuckmann-Post 2011 and Minton 2010 included 22 and 31 studies, respectively. This updated review includes seven studies from Minton 2010 and 22 studies from Peuckmann-Post 2011.

Summary of main results

This systematic review update identified 45 studies for inclusion, with a wide range of underlying diseases and drug interventions. Treatment results pointed to weak and inconclusive evidence for the efficacy of amantadine, pemoline and modafinil in multiple sclerosis and for carnitine and donepezil in cancer-related fatigue. Methylphenidate and pemoline seem to be effective in patients

with HIV, but this is based only on one study per intervention, with only a moderate number of participants in each study. Meta-analysis shows an estimated superior effect for methylphenidate in cancer-related fatigue, but not for pemoline and modafinil in multiple sclerosis. Therapeutic effects could not be described for dexamphetamine, paroxetine or testosterone.

Acetylsalicylic acid demonstrated surprising efficacy in patients with multiple sclerosis (Shaygannejad 2012; Wingerchuk 2005). However, this result was not supported by other studies. For clinical recommendation, further research confirming this positive effect is needed.

Some studies used other drugs, for example mistletoe extract, megestrol acetate and medroxyprogesterone acetate (Semiglazov 2006; Simons 1996; Westman 1999), but these studies lacked strong evidence and the clinical use of these drugs is rather rare. Further studies need to demonstrate their clinical efficacy.

The available evidence from randomised controlled trials (RCTs) is scarce, even though there is a surplus of case reports and uncontrolled trials. It has to be kept in mind that many of the included studies involved only a small number of participants (Lacasse 2004; Rosenberg 1988), and did not follow a consistent research methodology. In some cases the investigated population was very heterogenous and any outcome may have been associated with depression (Breitbart 2001; Wagner 2000), making it difficult to distinguish from primary fatigue. Fatigue often occurs in clusters, and fatigue and emotional distress are often concurrent factors. The National Comprehensive Cancer Network (NCCN) guidelines discuss studies, some of which show fatigue as an independent factor of depression, while others report moderate correlation. The 2014 NCCN guidelines recommended that the psychostimulant methylphenidate may be considered with caution for the pharmacological treatment of fatigue in selected patients. However, corticosteroids are recommended for short-term but not long-term therapy, due to their toxicity (NCCN 2014).

Treatment of secondary fatigue should be initiated with the treatment of the underlying cause. Some causes of secondary fatigue, such as anaemia, depression, infection, dehydration, malnutrition, hypercalcaemia, hypomagnesaemia, other metabolic disorders or the side effects of treatment with opioids and other sedative drugs, should also be treated, though little evidence from randomised trials is available on the efficacy of these treatments (EAPC 2008).

This review only included studies specifically focusing on palliative care in patients with advanced disease with the aim of relieving fatigue. Studies investigating curative treatment (e.g. treatment of early breast cancer) and fatigue directly related to treatment were not included in this review.

In addition, the results of the literature search indicate that recent research interest focuses on modafinil (eight studies) and its use in fatigue management for palliative care patients. This may be an interesting future perspective.

Potential biases in the review process

None of the authors of this review were involved in any of the excluded or included studies. In addition, none of the authors has any conflict of interest. Our search strategy was as comprehensive as possible. All studies were independently assessed for inclusion

by two review authors so we are confident that we have included all relevant studies and attempted to reduce bias in the review process.

Due to difficulties in conducting these types of trials for any of these drugs, trials may not have been published at all. Therefore, there is the potential for publication bias in this review. However, by contacting experts in this field we have attempted to reduce this bias as much as possible.

Agreements and disagreements with other studies or reviews

There have been two previous Cochrane Reviews (Minton 2010; Peuckmann-Post 2011), which are merged in this systematic review using a new search strategy to increase its coverage. Minton 2010 concluded in their last published version that there is increasing evidence that psychostimulant trials improve cancer-related fatigue at a clinically meaningful level. This review described the need for a large-scale RCT of methylphenidate to confirm the preliminary results from their review. They recommended haemopoietic growth factors for the treatment of cancer-related fatigue. However, with new safety data indicating increased adverse outcomes, these drugs can no longer be recommended in the treatment of cancer-related fatigue (Bennet 2008; Bohlius 2009; EAPC 2008; Glaspy 2009; Tonelli 2009). Peuckmann-Post 2011 concluded that, based on limited evidence, they could not recommend a specific drug for the treatment of fatigue in palliative care patients. The review pointed out that until 2009, corticosteroids had not been the focus of research on fatigue treatment, although these drugs were frequently used in clinical practice. This review also stated that fatigue research seems to focus on modafinil, which may be beneficial, although there is no evidence.

AUTHORS' CONCLUSIONS

Implications for practice

This review update found no evidence to support the use of a specific drug to treat fatigue in palliative care patients. Amantadine seems to be promising in patients with multiple sclerosis and fatigue, while methylphenidate is advantageous in patients with cancer-related fatigue. However, since the number of studies examining the effect of these drugs on fatigue, as well as the participant numbers, were relatively low, the evidence remains weak. Whether amantadine also relieves fatigue in cancer patients and whether methylphenidate relieves fatigue in multiple sclerosis patients has not been shown but should be investigated. Further studies are needed in patients with advanced disease and fatigue to show whether dexamethasone, methylprednisolone, donepezil, carnitine or modafinil may be beneficial, since the available evidence currently does not support these interventions.

Overall, most adverse effects of the investigated drugs seemed to be fairly moderate. Patients with HIV/AIDS and fatigue should be offered treatment with methylphenidate or pemoline. Similarly, other drugs may be tested, if the above-mentioned drugs are not available.

Clinical practice includes corticosteroids as a short-term therapeutic option for relief of fatigue in palliative care (Radbruch 2008; Yennurajalingam 2013). However, only one randomised

controlled trial (RCT) from our literature search matched the inclusion criteria (Yennurajalingam 2013). There are still not enough studies to show the therapeutic benefits of corticosteroids for this indication.

Implications for research

Patient group

Trials with a higher participant numbers may be able to detect small differences between groups. Palliative care studies in patients with advanced diseases, with the aim of relieving primary or secondary fatigue, should remain a focus of research. Further trials investigating subpopulations with different diseases in the end stage may be helpful. Stratification according to sociodemographic variables should be performed since, for example, younger age and male sex have been shown to predict worse fatigue (Auret 2009; Butler 2007).

Interventions

Amantadine, methylphenidate, carnitine, acetylsalicylic acid, dexamethasone, alfacalcidol, armodafinil and modafinil have been used in a few studies in this review with positive results. These drugs should be investigated in more detail to confirm their efficacy and should be examined for potentially similar efficacy in populations with different diseases and related fatigue. For instance, amantadine has been studied exclusively in multiple sclerosis patients and in post-polio studies. Therefore, it may be relevant to examine the effect of amantadine in other populations, such as cancer patients. Also, investigation of corticosteroids in RCTs would be highly relevant.

Comparisons

Future trials should compare one anti-fatigue drug to another anti-fatigue drug, combined with placebo-controlled comparisons.

Outcomes

There is no consensus on threshold values for relief of fatigue or on criteria for the responder. Patient-reported fatigue and variance in the outcome instruments could be used to measure the improvement of fatigue. Outcomes should also clearly define both the response and adverse effects from the treatment. It would be helpful to agree on the use of particular measurement instruments in order to perform better comparisons and analysis (Minton 2009).

Trial design

Further research is needed to identify effective and safe treatment for fatigue in palliative care. Multi-centre RCTs are recommended to assess the value of pharmacological treatments for fatigue. Additional data are required to confirm the results of this review and to provide a more significant estimate of efficacy.

Overall, this review demonstrates a lack of evidence rather than a lack of efficacy of the interventions. With regard to future research, studies with larger participant numbers are needed. However, the difficulties of low recruitment and high attrition rates have been described repeatedly in research in palliative care, as patients with advanced and life-threatening diseases are involved. It would also be helpful to limit the diversity of scales and scores used. Since no standardised recommendations for assessment of fatigue yet exist, it is difficult to determine whether the outcome measures were appropriate. Minton and Stone have suggested using the European Organization for Research and Treatment of Cancer Quality of Life Questionnaire (EORTC-QLQ-C30) fatigue subscale or the Functional Assessment of Cancer Therapy fatigue scale (Minton 2009).

Interestingly, there were no more than two studies with a focus on corticosteroids and fatigue as the primary outcome (Della Cuna 1989; Yennurajalingam 2013), although corticosteroids have been recommended (EAPC 2008) and are frequently used for this indication by clinicians since clinical experience shows a beneficial effect on many of the symptoms experienced by palliative care patients. The impact of corticosteroids should be investigated in future studies.

ACKNOWLEDGEMENTS

We would like to thank Frank Elsner, Norbert Krumm and Peter Trottenburg for their preparatory work. We would like to acknowledge the authors of the previous review: Ollie Minton, Alison Richardson, Michael Sharpe, Matthew Hotopf and Patrick Stone. We thank Anna Mücke, Putri Nastiti, Birgit Jaspers, Johannes Kögler, Grace and Anna Hobson for their helpful support during all stages of the report and their never-ending patience with all delays that occurred, and Roman Rolke who gave valuable feedback for the drafting of the review.

CRG Funding Acknowledgement: The National Institute for Health Research (NIHR) is the largest single funder of the Cochrane PaPaS Group. Disclaimer: The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the NIHR, NHS or the Department of Health.

REFERENCES

References to studies included in this review

Ashtari 2009 {published data only}

Ashtari F, Fatehi F, Shaygannejad V, Chitsaz A. Does amantadine have favourable effects on fatigue in Persian patients suffering from multiple sclerosis? *Neurologia i Neurochirurgia Polska* 2009; **43**(5):428-32. [PMID: 11176767]

Auret 2009 {published data only}

Auret KA, Schug SA, Bremner AP, Bulsara M. A randomized, double-blind, placebo-controlled trial assessing the impact of dexamphetamine on fatigue in patients with advanced cancer. *Journal of Pain and Symptom Management* 2009; **37**(4):613-21. [PMID: 18790598]

Barak 2014 {published data only}

Barak Y, Magalashvili D, Dolgopiat MD, Ladkani D, Nitzani D, Mazor Z, et al. Effect of alfacalcidol on fatigue in ms patients: a randomized, double-blind study. *Neurology* 2014; **82**(10 Suppl):S23.004.

Brass 2001 {published data only}

Brass EP, Adler S, Sietsema KE, Hiatt WR, Orlando AM, Amato A, et al. Intravenous L-carnitine increases plasma carnitine, reduces fatigue, and may preserve exercise capacity in hemodialysis patients. *American Journal of Kidney Diseases* 2001; **37**(5):1018-28. [PMID: 11325685]

Breitbart 2001 {published data only}

Breitbart W, Rosenfeld B, Kaim M, Funesti-Esch J. A randomized, double-blind, placebo-controlled trial of psychostimulants for the treatment of fatigue in ambulatory patients with human immunodeficiency virus disease. *Archives of Internal Medicine* 2001; **161**(3):411-20. [PMID: 11176767]

Bruera 2006 {published data only}

Bruera E, Valero V, Driver L, Shen L, Willey J, Zhang T, et al. Patient-controlled methylphenidate for cancer fatigue: a double-blind, randomized, placebo-controlled trial. *Journal of Clinical Oncology* 2006; **24**(13):2073-8. [PMID: 16648508]

Bruera 2007 {published data only}

Bruera E, El Osta B, Valero V, Driver LC, Pei BL, Shen L, et al. Donepezil for cancer fatigue: a double-blind, randomized, placebo-controlled trial. *Journal of Clinical Oncology* 2007; **25**(23):3475-81. [PMID: 17687152]

Butler 2007 {published data only}

, Case LD, Atkins J, Frizzell B, Sanders G, Griffin P, et al. A phase III, double-blind, placebo-controlled prospective randomized clinical trial of d-threo-methylphenidate HCl in brain tumor patients receiving radiation therapy. *International Journal of Radiation Oncology, Biology, Physics* 2007; **69**(5):1496-501. [PMID: 17869448]

Canadian MSRG 1987 {published data only}

Canadian MS Research Group. A randomized controlled trial of amantadine in fatigue associated with multiple sclerosis.

Canadian Journal of Neurological Sciences 1987; **14**(3):273-8. [PMID: 2889518]

Cohen 1989 {published data only}

Cohen RA, Fisher M. Amantadine treatment of fatigue associated with multiple sclerosis. *Archives of Neurology* 1989; **46**(6):676-80. [PMID: 2730380]

Cruciani 2009 {published data only}

Cruciani RA, Dvorkin E, Homel P, Culliney B, Malamud S, Lapin J, et al. L-carnitine supplementation in patients with advanced cancer and carnitine deficiency: a double-blind, placebo-controlled study. *Journal of Pain and Symptom Management* 2009; **37**(4):622-31. [PMID: 18809275]

Cruciani 2012 {published data only}

Cruciani RA, Zhang JJ, Manola J, Cella D, Ansari B, Fisch MJ. L-carnitine supplementation for the management of fatigue in patients with cancer: an Eastern Cooperative Oncology Group phase III, randomized, double-blind, placebo-controlled trial. *Journal of Clinical Oncology* 2012; **30**(31):3864-9. [PMID: 22987089]

Della Cuna 1989 {published data only}

Della Cuna GR, Pellegrini A, Piazzi M. Effect of methylprednisolone sodium succinate on quality of life in preterminal cancer patients: a placebo-controlled, multicenter study. The Methylprednisolone Preterminal Cancer Study Group. *European Journal of Cancer and Clinical Oncology* 1989; **25**(12):1817-21. [PMID: 2698804]

Escalante 2014 {published data only}

Escalante CP, Meyers C, Reuben JM, Wang X, Qiao W, Manzullo E, et al. A randomized, double-blind, 2-period, placebo-controlled crossover trial of a sustained-release methylphenidate in the treatment of fatigue in cancer patients. *Cancer Journal (Sudbury, Mass.)* 2014; **20**(1):8-14. [PMID: 24445757]

Fisch 2003 {published data only}

Fisch MJ, Loehrer PJ, Kristeller J, Passik S, Jung SH, Shen J, et al. Fluoxetine versus placebo in advanced cancer outpatients: a double-blinded trial of the Hoosier Oncology Group. *Journal of Clinical Oncology* 2003; **21**(10):1937-43. [PMID: 12743146]

Jean-Pierre 2010 {published data only}

Jean-Pierre P, Morrow GR, Roscoe JA, Heckler C, Mohile S, Janelins M, et al. A phase 3 randomized, placebo-controlled, double-blind, clinical trial of the effect of modafinil on cancer-related fatigue among 631 patients receiving chemotherapy: a University of Rochester Cancer Center Community Clinical Oncology Program Research base study. *Cancer* 2010; **116**(14):3513-20. [PMID: 20564068]

Kerr 2012 {published data only}

Kerr CW, Drake J, Milch RA, Brazeau DA, Skretny JA, Brazeau GA, et al. Effects of methylphenidate on fatigue and depression: a randomized, double-blind, placebo-controlled trial. *Journal of Pain and Symptom Management* 2012; **43**(1):68-77. [PMID: 20564068]

Knapp 2008 {published data only}

Knapp PE, Storer TW, Herbst KL, Singh AB, Dzekov C, Dzekov J, et al. Effects of a supraphysiological dose of testosterone on physical function, muscle performance, mood, and fatigue in men with HIV-associated weight loss. *American Journal of Physiology. Endocrinology and Metabolism* 2008; **294**(6):E1135-43. [PMID: 18430965]

Krupp 1995 {published data only}

Geisler MW, Sliwinski M, Coyle PK, Masur DM, Doscher C, Krupp LB. The effects of amantadine and pemoline on cognitive functioning in multiple sclerosis. *Archives of Neurology* 1996; **53**(2):185-8. [PMID: 8639070]

* Krupp LB, Coyle PK, Doscher C, Miller A, Cross AH, Jandorf L, et al. Fatigue therapy in multiple sclerosis: results of a double-blind, randomized, parallel trial of amantadine, pemoline, and placebo. *Neurology* 1995; **45**(11):1956-61. [PMID: 7501140]

Lacasse 2004 {published data only}

Lacasse Y, Beaudoin L, Rousseau L, Maltais F. Randomized trial of paroxetine in end-stage COPD. *Monaldi Archives for Chest Disease* 2004; **61**(3):140-7. [PMID: 15679006]

Lange 2009 {published data only}

Lange R, Volkmer M, Heesen C, Liepert J. Modafinil effects in multiple sclerosis patients with fatigue. *Journal of Neurology* 2009; **256**(4):645-50. [PMID: 19367356]

Lou 2009 {published data only}

Lou JS, Dimitrova DM, Park BS, Johnson SC, Eaton R, Arnold G, et al. Using modafinil to treat fatigue in Parkinson disease: a double-blind, placebo-controlled pilot study. *Clinical Neuropharmacology* 2009; **32**(6):305-10. [PMID: 19620846]

Moraska 2010 {published data only}

Moraska AR, Sood A, Dakhil SR, Sloan JA, Barton D, Atherton PJ, et al. Phase III, randomized, double-blind, placebo-controlled study of long-acting methylphenidate for cancer-related fatigue: North Central Cancer Treatment Group NCCTG-N05C7 trial. *Journal of Clinical Oncology* 2010; **28**(23):3673-9. [PMID: 20625123]

Morrow 2003 {published data only}

Morrow GR, Hickok JT, Roscoe JA, Raubertas RF, Andrews PL, Flynn PJ, et al. Differential effects of paroxetine on fatigue and depression: a randomized, double-blind trial from the University of Rochester Cancer Center Community Clinical Oncology Program. *Journal of Clinical Oncology* 2003; **21**(24):4635-41. [PMID: 14673053]

Murray 1985 {published data only}

Murray TJ. Amantadine therapy for fatigue in multiple sclerosis. *Canadian Journal of Neurological Sciences* 1985; **12**(3):251-4. [PMID: 3902184]

Rabkin 2000 {published data only}

Rabkin JG, Wagner GJ, Rabkin R. A double-blind, placebo-controlled trial of testosterone therapy for HIV-positive men with hypogonadal symptoms. *Archives of General Psychiatry* 2000; **57**(2):141-7; discussion 155-6. [PMID: 10665616]

Rabkin 2004 {published data only}

Rabkin JG, Wagner GJ, McElhiney MC, Rabkin R, Lin SH. Testosterone versus fluoxetine for depression and fatigue in HIV/AIDS: a placebo-controlled trial. *Journal of Clinical Psychopharmacology* 2004; **24**(4):379-85. [PMID: 15232328]

Rabkin 2009 {published data only}

Rabkin JG, Gordon PH, McElhiney M, Rabkin R, Chew S, Mitsumoto H. Modafinil treatment of fatigue in patients with ALS: a placebo-controlled study. *Muscle and Nerve* 2009; **39**(3):297-303. [PMID: 19208404]

Rabkin 2010 {published data only}

Rabkin JG, McElhiney MC, Rabkin R, McGrath P. Modafinil treatment for fatigue in HIV/AIDS: a randomized placebo-controlled study. *Journal of Clinical Psychiatry* 2010; **71**(6):707-15. [PMID: 20492840]

Rabkin 2011 {published data only}

Rabkin JG, McElhiney MC, Rabkin R. Treatment of HIV-related fatigue with armodafinil: a placebo-controlled randomized trial. *Psychosomatics* 2011; **52**(4):328-36. [PMID: 2177715]

Rosenberg 1988 {published data only}

Rosenberg GA, Appenzeller O. Amantadine, fatigue, and multiple sclerosis. *Archives of Neurology* 1988; **45**:1104-6. [PMID: 2972270]

Roth 2010 {published data only}

Roth AJ, Nelson C, Rosenfeld B, Scher H, Slovin S, Morris M, et al. Methylphenidate for fatigue in ambulatory men with prostate cancer. *Cancer* 2010; **116**(21):5102-10. [PMID: 20665492]

Semiglazov 2006 {published data only}

Semiglazov VF, Stepula VV, Dudov A, Schnitker J, Mengs U. Quality of life is improved in breast cancer patients by Standardised Mistletoe Extract PS76A2 during chemotherapy and follow-up: a randomised, placebo-controlled, double-blind, multicentre clinical trial. *Anticancer research* 2006; **26**(2B):1519-29. [PMID: 16619567]

Shaygannejad 2012 {published data only}

Shaygannejad V, Janghorbani M, Ashtari F, Zakeri H. Comparison of the effect of aspirin and amantadine for the treatment of fatigue in multiple sclerosis: a randomized, blinded, crossover study. *Neurological Research* 2012; **34**(9):854-8. [PMID: 22979982]

Simons 1996 {published data only}

Simons JP, Aaronson NK, Vansteenkiste JF, ten Velde GP, Muller MJ, Drenth BM, et al. Effects of medroxyprogesterone acetate on appetite, weight, and quality of life in advanced-stage non-hormone-sensitive cancer: a placebo-controlled multicenter study. *Journal of Clinical Oncology* 1996; **14**(4):1077-84. [PMID: 8648360]

Spathis 2014 {published data only}

Spathis A, Fife K, Blackhall F, Dutton S, Bahadori R, Wharton R, et al. Modafinil for the treatment of fatigue in lung cancer: results of a placebo-controlled, double-blind, randomized trial. *Journal of Clinical Oncology* 2014; **32**:1-8. [PMID: 24778393]

Stankoff 2005 {published data only}

Stankoff B, Waubant E, Confavreux C, Edan G, Debouverie M, Rumbach L, et al. Modafinil for fatigue in MS: a randomized placebo-controlled double-blind study. *Neurology* 2005; **64**(7):1139-43. [PMID: 15824337]

Stein 1995 {published data only}

Stein DP, Dambrosia JM, Dalakas MC. A double-blind, placebo-controlled trial of amantadine for the treatment of fatigue in patients with the post-polio syndrome. *Annals of the New York Academy of Sciences* 1995; **753**:296-302. [PMID: 7611638]

Tomassini 2004 {published data only}

Tomassini V, Pozzilli C, Onesti E, Pasqualetti P, Marinelli F, Pisani A, et al. Comparison of the effects of acetyl L-carnitine and amantadine for the treatment of fatigue in multiple sclerosis: results of a pilot, randomised, double-blind, crossover trial. *Journal of the Neurological Sciences* 2004; **218**(1-2):103-8. [PMID: 14759641]

Vasconcelos 2007 {published data only}

Vasconcelos OM, Prokhorenko OA, Salajegheh MK, Kelley KF, Livornese K, Olsen CH, et al. Modafinil for treatment of fatigue in post-polio syndrome: a randomized controlled trial. *Neurology* 2007; **68**(20):1680-6. [PMID: 17502549]

Wagner 2000 {published data only}

Wagner GJ, Rabkin R. Effects of dextroamphetamine on depression and fatigue in men with HIV: a double-blind, placebo-controlled trial. *Journal of Clinical Psychiatry* 2000; **61**(6):436-40. [PMID: 10901342]

Weinshenker 1992 {published data only}

Weinshenker BG, Penman M, Bass B, Ebers GC, Rice GP. A double-blind, randomized, crossover trial of pemoline in fatigue associated with multiple sclerosis. *Neurology* 1992; **42**(8):1468-71. [PMID: 1641137]

Westman 1999 {published data only}

Westman G, Bergman B, Albertsson M, Kadar L, Gustavsson G, Thaning L, et al. Megestrol acetate in advanced, progressive, hormone-insensitive cancer. Effects on the quality of life: a placebo-controlled, randomised, multicentre trial. *European Journal of Cancer* 1999; **35**(4):586-95. [PMID: 10492632]

Wingerchuk 2005 {published data only}

Wingerchuk DM, Benarroch EE, O'Brien PC, Keegan BM, Lucchinetti CF, Noseworthy JH, et al. A randomized controlled crossover trial of aspirin for fatigue in multiple sclerosis. *Neurology* 2005; **64**(7):1267-9. [PMID: 15824361]

Yennurajalingam 2013 {published data only}

Yennurajalingam S, Frisbee-Hume S, Palmer JL, Delgado-Guay MO, Bull J, Phan AT, et al. Reduction of cancer-related fatigue with dexamethasone: a double-blind, randomized, placebo-controlled trial in patients with advanced cancer. *Journal of Clinical Oncology* 2013; **31**(25):3076-82. [PMID: 23897970]

References to studies excluded from this review
Agteresch 2000 {published data only}

Agteresch HJ, Dagnelie PC, van der Gaast A, Stijnen T, Wilson JH. Randomized clinical trial of adenosine 5'-triphosphate in patients with advanced non-small-cell lung cancer. *Journal of the National Cancer Institute* 2000; **92**(4):321-8. [PMID: 10675381]

Bruera 1987 {published data only}

Bruera E, Chadwick S, Brenneis C, Hanson J, MacDonald RN. Methylphenidate associated with narcotics for the treatment of cancer pain. *Cancer Treatment Reports* 1987; **71**(1):67-70. [PMID: 3791269]

Bruera 1998 {published data only}

Bruera E, Ernst S, Hagen N, Spachynski K, Belzile M, Hanson J, et al. Effectiveness of megestrol acetate in patients with advanced cancer: a randomized, double-blind, crossover study. *Cancer Prevention and Control* 1998; **2**(2):74-8. [PMID: 9765767]

Bruera 2003 {published data only}

Bruera E, Strasser F, Shen L, Palmer JL, Willey J, Driver LC, et al. The effect of donepezil on sedation and other symptoms in patients receiving opioids for cancer pain: a pilot study. *Journal of Pain and Symptom Management* 2003; **26**(5):1049-54. [PMID: 14585555]

Capuron 2002 {published data only}

Capuron L, Gumnick JF, Musselman DL, Lawson DH, Reemsnyder A, Nemeroff CB, et al. Neurobehavioral effects of interferon-alpha in cancer patients: phenomenology and paroxetine responsiveness of symptom dimensions. *Neuropsychopharmacology* 2002; **26**(5):643-52. [PMID: 11927189]

Carter 2005 {published data only}

Carter GT, Weiss MD, Lou JS, Jensen MP, Abresch RT, Martin TK, et al. Modafinil to treat fatigue in amyotrophic lateral sclerosis: an open label pilot study. *American Journal of Hospice and Palliative Care* 2005; **22**(1):55-9. [PMID: 15736608]

Cerchietti 2004 {published data only}

Cerchietti LC, Navigante AH, Peluffo GD, Diamant MJ, Stillitani I, Klein SA, et al. Effects of celecoxib, medroxyprogesterone, and dietary intervention on systemic syndromes in patients with advanced lung adenocarcinoma: a pilot study. *Journal of Pain and Symptom Management* 2004; **27**(1):85-95. [PMID: 14711473]

Cruciani 2004 {published data only}

Cruciani RA, Dvorkin E, Homel P, Culliney B, Malamud S, Shaiova L, et al. L-carnitine supplementation for the treatment of fatigue and depressed mood in cancer patients with carnitine deficiency: a preliminary analysis. *Annals of the New York Academy of Sciences* 2004; **1033**:168-76. [PMID: 15591014]

Cruciani 2006 {published data only}

Cruciani RA, Dvorkin E, Homel P, Malamud S, Culliney B, Lapin J, et al. Safety, tolerability and symptom outcomes associated with L-carnitine supplementation in patients with cancer, fatigue, and carnitine deficiency: a phase I/II study. *Journal*

- of *Pain and Symptom Management* 2006; **32**(6):551-9. [PMID: 17157757]
- Cueva 2012** {published data only}
Cueva JF, Calvo M, Anido U, León L, Gallardo E, Areses C, et al. Methylphenidate in the management of asthenia in breast cancer patients treated with docetaxel: results of a pilot study. *Investigational New Drugs* 2012; **30**(2):688-94. [PMID: 20844926]
- Cullum 2004** {published data only}
Cullum JL, Wojciechowski AE, Pelletier G, Simpson JS. Bupropion sustained release treatment reduces fatigue in cancer patients. *Canadian Journal of Psychiatry. Revue Canadienne de Psychiatrie* 2004; **49**(2):139-44. [PMID: 15065748]
- De Conno 1998** {published data only}
De Conno F, Martini C, Zecca E, Balzarini A, Venturino P, Groff L, et al. Megestrol acetate for anorexia in patients with far-advanced cancer: a double-blind controlled clinical trial. *European Journal of Cancer* 1998; **34**(11):1705-9. [PMID: 9893656]
- Diel 2004** {published data only}
Diel IJ, Body JJ, Lichinitser MR, Kreuser ED, Dornoff W, Gorbunova VA, et al. Improved quality of life after long-term treatment with the bisphosphonate ibandronate in patients with metastatic bone disease due to breast cancer. *European Journal of Cancer* 2004; **40**(11):1704-12. [PMID: 15251160]
- Downer 1993** {published data only}
Downer S, Joel S, Allbright A, Plant H, Stubbs L, Talbot D, et al. A double blind placebo controlled trial of medroxyprogesterone acetate (MPA) in cancer cachexia. *British Journal of Cancer* 1993; **67**(5):1102-5. [PMID: 8494706]
- Dunlop 2007** {published data only}
Dunlop BW, Crits-Christoph P, Evans DL, Hirschowitz J, Solvason HB, Rickels K, et al. Coadministration of modafinil and a selective serotonin reuptake inhibitor from the initiation of treatment of major depressive disorder with fatigue and sleepiness: a double-blind, placebo-controlled study. *Journal of Clinical Psychopharmacology* 2007; **27**(6):614-9. [PMID: 18004129]
- Gehring 2012** {published data only}
Gehring K, Patwardhan SY, Collins R, Groves MD, Etzel CJ, Meyers CA, et al. A randomized trial on the efficacy of methylphenidate and modafinil for improving cognitive functioning and symptoms in patients with a primary brain tumor. *Journal of Neuro-oncology* 2012; **107**(1):165-74. [PMID: 21964738]
- Graziano 2002** {published data only}
Graziano F, Bisonni R, Catalano V, Silva R, Rovidati S, Mencarini E, et al. Potential role of levocarnitine supplementation for the treatment of chemotherapy-induced fatigue in non-anaemic cancer patients. *British Journal of Cancer* 2002; **86**(12):1854-7. [PMID: 12085175]
- Hannestad 2011** {published data only}
Hannestad J, Della Gioia N, Ortiz N, Pittman B, Bhagwagar Z. Citalopram reduces endotoxin-induced fatigue. *Brain, Behavior, and Immunity* 2011; **25**(2):256-9. [PMID: 20955776]
- Hovey 2014** {published data only}
Hovey E, de Souza P, Marx G, Parente P, Rapke T, Hill A, et al. Phase III, randomized, double-blind, placebo-controlled study of modafinil for fatigue in patients treated with docetaxel-based chemotherapy. *Supportive Care in Cancer* 2014; **22**(5):1233-42. [PMID: 24337761]
- Inoue 2003** {published data only}
Inoue A, Yamada Y, Matsumura Y, Shimada Y, Muro K, Gotoh M, et al. Randomized study of dexamethasone treatment for delayed emesis, anorexia and fatigue induced by irinotecan. *Supportive Care in Cancer* 2003; **11**(8):528-32. [PMID: 12844250]
- Lauretti 2013** {published data only}
Lauretti GR, Rizzo CC, Mattos AL, Rodrigues SW. Epidural methadone results in dose-dependent analgesia in cancer pain, further enhanced by epidural dexamethasone. *British Journal of Cancer* 2013; **108**(2):259-64. [PMID: 23322191]
- Laval 2008** {published data only}
Laval G, Paris A. Methylphenidate in palliative care in cancer patient: a double-blind randomised trial versus placebo [Methylphenidate et asthenie en soins palliatifs chez le patient cancéreux: essai multicentrique randomise versus placebo en double aveugle]. *Bulletin du Cancer* 2008; **95**(2):241-6. [PMID: 18304907]
- Lower 2009** {published data only}
Lower EE, Fleishman S, Cooper A, Zeldis J, Faleck H, Yu Z, et al. Efficacy of dexmethylphenidate for the treatment of fatigue after cancer chemotherapy: a randomized clinical trial. *Journal of Pain and Symptom Management* 2009; **38**(5):650-62. [PMID: 19896571]
- Mar Fan 2008** {published data only}
Mar Fan HG, Clemons M, Xu W, Chemerynsky I, Breunis H, Braganza S, et al. A randomised, placebo-controlled, double-blind trial of the effects of d-methylphenidate on fatigue and cognitive dysfunction in women undergoing adjuvant chemotherapy for breast cancer. *Supportive Care in Cancer* 2008; **16**(6):577-83. [PMID: 17972110]
- McElhiney 2010** {published data only}
McElhiney M, Rabkin J, Van Gorp W, Rabkin R. Modafinil effects on cognitive function in HIV+ patients treated for fatigue: a placebo controlled study. *Journal of Clinical and Experimental Neuropsychology* 2010; **32**(5):474-80. [PMID: 19937504]
- McElhiney 2013** {published data only}
McElhiney M, Rabkin J, Van Gorp W, Rabkin R. Effect of armodafinil on cognition in patients with HIV/AIDS and fatigue. *Journal of Clinical and Experimental Neuropsychology* 2013; **35**(7):718-27. [PMID: 23944194]

Mercadante 2001 {published data only}

Mercadante S, Fulfaro F, Casuccio A. The use of corticosteroids in home palliative care. *Supportive Care in Cancer* 2001; **9**(5):386-9. [PMID: 11497394]

Metz 2004 {published data only}

Metz LM, Patten SB, Archibald CJ, Bakker JI, Harris CJ, Patry DG, et al. The effect of immunomodulatory treatment on multiple sclerosis fatigue. *Journal of Neurology, Neurosurgery, and Psychiatry* 2004; **75**(7):1045-7. [PMID: 15201369]

Moertel 1974 {published data only}

Moertel CG, Schutt AJ, Reitemeier RJ, Hahn RG. Corticosteroid therapy of preterminal gastrointestinal cancer. *Cancer* 1974; **33**(6):1607-9. [PMID: 4135151]

Mohr 2003 {published data only}

Mohr DC, Hart SL, Goldberg A. Effects of treatment for depression on fatigue in multiple sclerosis. *Psychosomatic Medicine* 2003; **65**(4):542-7. [PMID: 12883103]

Monk 2006 {published data only}

Monk JP, Phillips G, Waite R, Kuhn J, Schaaf LJ, Otterson GA, et al. Assessment of tumor necrosis factor alpha blockade as an intervention to improve tolerability of dose-intensive chemotherapy in cancer patients. *Journal of Clinical Oncology* 2006; **24**(12):1852-9. [PMID: 16622259]

Moss 2006 {published data only}

Moss EL, Simpson JS, Pelletier G, Forsyth P. An open-label study of the effects of bupropion SR on fatigue, depression and quality of life of mixed-site cancer patients and their partners. *Psycho-Oncology* 2006; **15**(3):259-67. [PMID: 16041840]

Popiela 1989 {published data only}

Popiela T, Lucchi R, Giongo F. Methylprednisolone as palliative therapy for female terminal cancer patients. The Methylprednisolone Female Preterminal Cancer Study Group. *European Journal of Cancer and Clinical Oncology* 1989; **25**(12):1823-9. [PMID: 2483687]

Rabkin 2000a {published data only}

Rabkin JG, Ferrando SJ, Wagner GJ, Rabkin R. DHEA treatment for HIV+ patients: effects on mood, androgenic and anabolic parameters. *Psychoneuroendocrinology* 2000; **25**(1):53-68. [PMID: 10633535]

Rabkin 2011a {published data only}

Rabkin JG, McElhiney MC, Rabkin R. Modafinil and armodafinil treatment for fatigue for HIV-positive patients with and without chronic hepatitis C. *International Journal of STD & AIDS* 2011; **22**(2):95-101. [PMID: 21427431]

Rammohan 2002 {published data only}

Rammohan KW, Rosenberg JH, Lynn DJ, Blumenfeld AM, Pollak CP, Nagaraja HN. Efficacy and safety of modafinil (Provigil) for the treatment of fatigue in multiple sclerosis: a two centre phase 2 study. *Journal of Neurology, Neurosurgery, and Psychiatry* 2002; **72**(2):179-83. [PMID: 11796766]

Romani 2004 {published data only}

Romani A, Bergamaschi R, Candeloro E, Alfonsi E, Callieco R, Cosi V. Fatigue in multiple sclerosis: multidimensional assessment and response to symptomatic treatment. *Multiple Sclerosis (Houndmills, Basingstoke, England)* 2004; **10**(4):462-8. [PMID: 15327047]

Roscoe 2005 {published data only}

Roscoe JA, Morrow GR, Hickok JT, Mustian KM, Griggs JJ, Matteson SE, et al. Effect of paroxetine hydrochloride (Paxil) on fatigue and depression in breast cancer patients receiving chemotherapy. *Breast Cancer Research and Treatment* 2005; **89**(3):243-9. [PMID: 15754122]

Sailer 2000 {published data only}

Sailer M, Heinze HJ, Schoenfeld MA, Hauser U, Smid HG. Amantadine influences cognitive processing in patients with multiple sclerosis. *Pharmacopsychiatry* 2000; **33**(1):28-37. [PMID: 10721881]

Shaw 2006 {published data only}

Shaw EG, Rosdhal R, Lovato J, Naughton MJ, Robbins ME, et al. Phase II study of donepezil in irradiated brain tumor patients: effect on cognitive function, mood, and quality of life. *Journal of Clinical Oncology* 2006; **24**(9):1415-20. [PMID: 16549835]

Shaw 2013 {published data only}

Shaw EG, Case D, Bryant D, Grisell D, Lesser G, Monitto DC, et al. Phase II double-blind placebo-controlled study of armodafinil for brain radiation induced fatigue. *Journal of Clinical Oncology* 2013; **31**(Suppl):Abstr 9505.

Spathis 2009 {published data only}

Spathis A, Dhillan R, Booden D, Forbes K, Vrotsou K, Fife K. Modafinil for the treatment of fatigue in lung cancer: a pilot study. *Palliative Medicine* 2009; **23**(4):325-31. [PMID: 19270033]

Stockler 2007 {published data only}

Stockler MR, O'Connell R, Nowak AK, Goldstein D, Turner J, Wilcken NR, et al. Effect of sertraline on symptoms and survival in patients with advanced cancer, but without major depression: a placebo-controlled double-blind randomised trial. *Lancet Oncology* 2007; **8**(7):603-12. [PMID: 17548243]

Torta 2007 {published data only}

Torta R, Berra C, Binaschi L, Borio R. Amisulpride in the short-term treatment of depressive and physical symptoms in cancer patients during chemotherapies. *Supportive Care in Cancer* 2007; **15**(5):539-46. [PMID: 17406919]

Wade 2002 {published data only}

Wade DT, Young CA, Chaudhuri KR, Davidson DL. A randomised placebo controlled exploratory study of vitamin B-12, lofepramine, and L-phenylalanine (the "Cari Loder regime") in the treatment of multiple sclerosis. *Journal of Neurology, Neurosurgery, and Psychiatry* 2002; **73**(3):246-9. [PMID: 12185153]

Wagner 1997 {published data only}

Wagner GJ, Rabkin JG, Rabkin R. Dextroamphetamine as a treatment for depression and low energy in AIDS patients:

a pilot study. *Journal of Psychosomatic Research* 1997; **42**(4):407-11. [PMID: 9160280]

Weitzner 2002 {published data only}

Weitzner MA, Moncello J, Jacobsen PB, Minton S. A pilot trial of paroxetine for the treatment of hot flashes and associated symptoms in women with breast cancer. *Journal of Pain and Symptom Management* 2002; **23**(4):337-45. [PMID: 11997203]

Wilwerding 1995 {published and unpublished data}

Wilwerding MB, Loprinzi CL, Mailliard JA, O'Fallon JR, Miser AW, van Haelst C, et al. A randomized, crossover evaluation of methylphenidate in cancer patients receiving strong narcotics. *Supportive Care in Cancer* 1995; **3**(2):135-8. [PMID: 7539701]

Zifko 2002 {published data only}

Zifko UA, Rupp M, Schwarz S, Zipko HT, Maida EM. Modafinil in treatment of fatigue in multiple sclerosis. Results of an open-label study. *Journal of Neurology* 2002; **249**(8):983-7. [PMID: 12195441]

Additional references

Atkins 2004

Atkins D, Best D, Briss PA, Eccles M, Falck-Ytter Y, Flottorp S, et al. Grading quality of evidence and strength of recommendations. *BMJ* 2004; **328**(7454):1490.

Bakshi 2003

Bakshi R. Fatigue associated with multiple sclerosis: diagnosis, impact and management. *Multiple Sclerosis* 2003; **9**(3):219-27. [PMID: 12814166]

Bennet 2008

Bennett CL, Silver SM, Djulbegovic B, Samaras AT, Blau CA, Gleason KJ, et al. Venous thromboembolism and mortality associated with recombinant erythropoietin and darbepoetin administration for the treatment of cancer-associated anemia. *JAMA* 2008; **299**(8):914-24. [DOI: [10.1001/jama.299.8.914](https://doi.org/10.1001/jama.299.8.914)] [PMID: 18314434]

Bohlius 2009

Bohlius J, Schmidlin K, Brillant C, Schwarzer G, Trelle S, Seidenfeld J, et al. Recombinant human erythropoiesis-stimulating agents and mortality in patients with cancer: a meta-analysis of randomised trials. *Lancet* 2009; **373**(9674):1532-42. [DOI: [10.1016/S0140-6736\(09\)60502-X](https://doi.org/10.1016/S0140-6736(09)60502-X)] [PMID: 19410717]

Breitbart 1998

Breitbart W, McDonald MV, Rosenfeld B, Monkman ND, Passik S. Fatigue in ambulatory AIDS patients. *Journal of Pain and Symptom Management* 1998; **15**(3):159-67. [PMID: 9564117]

Cantarero-Villanueva 2014

Cantarero-Villanueva I, Fernández-Lao C, Díaz-Rodríguez L, Cuesta-Vargas AI, Fernández-de-las-Peñas C, Piper BF, et al. The Piper Fatigue Scale-Revised: translation and psychometric evaluation in Spanish-speaking breast cancer survivors. *Quality of Life Research* 2014; **23**(1):271-6. [PMID: 23695719]

Cella 1998

Cella D, Peterman A, Passik S, Jacobsen P, Breitbart W. Progress towards guidelines for the management of fatigue. *Oncology (Williston Park)* 1998; **12**(11A):369-77. [PMID: 10028520]

Cella 2001

Cella D, Davis K, Breitbart W, Curt G. Cancer-related fatigue: prevalence of proposed diagnostic criteria in a United States sample of cancer survivors. *Journal of Clinical Oncology* 2001; **19**(14):3385-91. [PMID: 11454886]

Cella 2002

Cella D, Eton DT, Lai J, Peterman SA, Merkel DE. Combining anchor and distribution-based methods to derive minimal clinically important differences on the Functional Assessment of Cancer Therapy (FACT) anemia and fatigue scales. *Journal of Pain and Symptom Management* 2002; **24**:547-61. [PMID: 12551804]

Cramp 2012

Cramp F, Byron-Daniel J. Exercise for the management of cancer-related fatigue in adults. *Cochrane Database of Systematic Reviews* 2012, Issue 11. [DOI: [10.1002/14651858.CD006145.pub3](https://doi.org/10.1002/14651858.CD006145.pub3)] [PMID: 23152233]

Cuhls 2014

Cuhls H, Mücke M, Mochamat M, Radbruch L. Underestimated symptom: frequent and burdensome than pain [Unterschätztes Symptom: Häufiger und belastender als Schmerzen]. *Im Focus Onkologie* 2014; **17**(9):60-7. [DOI: [10.1007/s15015-014-0007-7](https://doi.org/10.1007/s15015-014-0007-7)]

Darko 1995

Darko DF, Mitler MM, Henriksen SJ. Lentiviral infection, immune response peptides and sleep. *Advances in Neuroimmunology* 1995; **5**:57-77. [PMID: 7795894]

Dittner 2004

Dittner AJ, Wessely SC, Brown RG. The assessment of fatigue: a practical guide for clinicians and researchers. *Journal of Psychosomatic Research* 2004; **56**(2):157-70. [PMID: 15016573]

EAPC 2008

Radbruch L, Strasser F, Elsner F, Gonçalves JF, Løge J, Kaasa S, et al. Research Steering Committee of the European Association for Palliative Care (EAPC). Fatigue in palliative care patients -- an EAPC approach. *Palliative Medicine* 2008; **22**:13-32. [DOI: [10.1177/0269216307085183](https://doi.org/10.1177/0269216307085183)] [PMID: 18216074]

Elkington 2005

Elkington H, White P, Addington-Hall J, Higgs R, Edmonds P. The healthcare needs of chronic obstructive pulmonary disease patients in the last year of life. *Palliative Medicine* 2005; **19**(6):485-91. [PMID: 16218161]

Eriksen 2003

Eriksen J, Jensen MK, Sjogren P, Ekholm O, Rasmussen NK. Epidemiology of chronic non-malignant pain in Denmark. *Pain* 2003; **106**(3):221-8. [PMID: 14659505]

Evans 2007

Evans WJ, Lambert CP. Physiological basis of fatigue. *American Journal of Physical Medicine and Rehabilitation* 2007; **86**:S29-46. [PMID: 17370370]

Farrar 2003

Farrar JT, Berlin JA, Strom BL. Clinically important changes in acute pain outcome measures: a validation study. *Journal of Pain and Symptom Management* 2003; **25**(5):406-11. [PMID: 12727037]

FDA 2005

. Information for Healthcare Professionals: Pemoline Tablets and Chewable Tablets (marketed as Cylert). <http://www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/ucm126461.htm> 2005 (accessed 3 September 2014).

Galvao 2005

Galvao DA, Newton RU. Review of exercise intervention studies in cancer patients. *Journal of Clinical Oncology* 2005; **23**(4):899-909. [PMID: 15681536]

Glaspy 2009

Glaspy JA. Erythropoietin in cancer patients. *Annual Review of Medicine* 2009; **60**:181-92. [DOI: [10.1146/annurev.med.60.050307.110718](https://doi.org/10.1146/annurev.med.60.050307.110718)] [PMID: 18980468]

Glaspy 2010

Glaspy J, Crawford J, Vansteenkiste J, Henry D, Rao S, Bowers P, et al. Erythropoiesis-stimulating agents in oncology: a study-level meta-analysis of survival and other safety outcomes. *British Journal of Cancer* 2010; **102**:301-15. [DOI: [10.1038/sj.bjc.6605498](https://doi.org/10.1038/sj.bjc.6605498)] [PMID: 20051958]

Goodlin 2005

Goodlin SJ. Palliative care for end-stage heart failure. *Current Heart Failure Reports* 2005; **2**(3):155-60. [PMID: 16138952]

Higgins 2011

Higgins JPT, Altman DG, . Chapter 8: Assessing risk of bias in included studies. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Jhamb 2013

Jhamb M, Liang K, Yabes J, Steel JL, Dew MA, Shah N, et al. Prevalence and correlates of fatigue in chronic kidney disease and end-stage renal disease: are sleep disorders a key to understanding fatigue? *American Journal of Nephrology* 2013; **38**(6):489-95. [PMID: 24335380]

Krupp 2003

Krupp LB. Fatigue in multiple sclerosis: definition, pathophysiology and treatment. *CNS Drugs* 2003; **17**(4):225-34. [PMID: 12665396]

Kummer 2011

Kummer A, Scalzo P, Cardoso F, Teixeira AL. Evaluation of fatigue in Parkinson's disease using the Brazilian version of

Parkinson's Fatigue Scale. *Acta Neurologica Scandinavica* 2011; **123**(2):130-6. [DOI: [10.1111/j.1600-0404.2010.01364.x](https://doi.org/10.1111/j.1600-0404.2010.01364.x)] [PMID: 20456242]

Lawrence 2004

Lawrence DP, Kupelnick B, Miller K, Devine D, Lau J. Evidence report on the occurrence, assessment, and treatment of fatigue in cancer patients. *Journal of the National Cancer Institute. Monographs* 2004; **32**:40-50. [PMID: 15263040]

Minton 2009

Minton O, Stone P. A systematic review of the scales used for the measurement of cancer-related fatigue (CRF). *Annals of Oncology* 2009; **20**(1):17-25. [PMID: 18678767]

Minton 2010

Minton O, Richardson A, Sharpe M, Hotopf M, Stone P. Drug therapy for the management of cancer-related fatigue. *Cochrane Database of Systematic Reviews* 2010, Issue 7. [DOI: [10.1002/14651858.CD006704.pub3](https://doi.org/10.1002/14651858.CD006704.pub3)] [20614448]

Mock 2004

Mock V. Evidence-based treatment for cancer-related fatigue. *Journal of the National Cancer Institute. Monographs* 2004; **32**:112-8. [PMID: 15263051]

Morrow 2005

Morrow GR, Shelke AR, Roscoe JA, Hickok JT, Mustian K. Management of cancer-related fatigue. *Cancer Investigations* 2005; **23**(3):229-39. [PMID: 15945509]

NCCN 2014

National Comprehensive Cancer Network. Cancer related fatigue. www.nccn.org/professionals/physician_gls/PDF/fatigue.pdf (accessed 30 October 2014).

NMSS 2002

National Multiple Sclerosis Society. Management of MS-related fatigue. www.nationalmssociety.org/PRC.asp 2002 (accessed 17 November 2005).

Norval 2004

Norval DA. Symptoms and sites of pain experienced by AIDS patients. *South African Medical Journal* 2004; **94**(6):450-4. [PMID: 15250459]

Patrick 2004

Patrick DL, Ferketich SL, Frame PS, Harris JJ, Hendricks CB, Levin B, et al. National Institutes of Health state-of-the-science conference statement: symptom management in cancer: pain, depression, and fatigue, July 15-17, 2002. *Journal of the National Cancer Institute. Monographs* 2004; **32**:9-16. [PMID: 15263035]

Pucci 2007

Pucci E, Branäs P, D'Amico R, Giuliani G, Solari A, Taus C, et al. Amantadine for fatigue in multiple sclerosis. *Cochrane Database of Systematic Reviews* 2007, Issue 1. [DOI: [10.1002/14651858.CD002818.pub2](https://doi.org/10.1002/14651858.CD002818.pub2)] [PMID: 17253480]

Radbruch 2008

Radbruch L, Strasser F, Elsner F, Goncalves JF, Loge J, Kaasa S, et al. Fatigue in palliative care patients -- an EAPC approach. *Palliative Medicine* 2008; **22**(1):13-32. [PMID: 18216074]

Rao 2004

Rao A, Cohen HJ. Symptom management in the elderly cancer patient: fatigue, pain, and depression. *Journal of the National Cancer Institute. Monographs* 2004; **32**:150-7. [PMID: 15263059]

RevMan 2014 [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration Review Manager. Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Richardson 1997

Richardson A, Ream EK. Self-care behaviours initiated by chemotherapy patients in response to fatigue. *International Journal of Nursing Studies* 1997; **34**(1):35-43. [PMID: 9055119]

Rosenberg 2005

Rosenberg JH, Shafor R. Fatigue in multiple sclerosis: a rational approach to evaluation and treatment. *Current Neurology and Neuroscience Reports* 2005; **5**(2):140-6. [PMID: 15743552]

Schmitz 2005

Schmitz KH, Holtzman J, Courneya KS, Masse LC, Duval S, Kane R. Controlled physical activity trials in cancer survivors: a systematic review and meta-analysis. *Cancer Epidemiology, Biomarkers and Prevention* 2005; **14**(7):1588-95. [PMID: 16030088]

Sood 2006

Sood A, Barton DL, Loprinzi CL. Use of methylphenidate in patients with cancer. *American Journal of Hospice and Palliative Medicine* 2006; **23**:35-40. [PMID: 16450661]

Stone 2002

Stone P. The measurement, causes and effective management of cancer-related fatigue. *International Journal of Palliative Nursing* 2002; **8**(3):120-8. [PMID: 11923740]

Stridsman 2014

Stridsman C, Lindberg A, Skär L. Fatigue in chronic obstructive pulmonary disease: a qualitative study of people's experiences. *Scandinavian Journal of Caring Sciences* 2014; **28**(1):130-8. [DOI: [10.1111/scs.12033](https://doi.org/10.1111/scs.12033)] [PMID: 23517049]

Tang 2010

Tang WR, Yu CY, Yeh SJ. Fatigue and its related factors in patients with chronic heart failure. *Journal of Clinical Nursing* 2010; **19**(1-2):69-78. [PMID: 20500245]

Tejani 2012

Tejani AM, Wasdell M, Spiwak R, Rowell G, Nathwani S. Carnitine for fatigue in multiple sclerosis. *Cochrane Database of Systematic Reviews* 2012, Issue 5. [DOI: [10.1002/14651858.CD007280.pub3](https://doi.org/10.1002/14651858.CD007280.pub3)] [PMID: 22592719]

Tonelli 2009

Tonelli M, Hemmelgarn B, Reiman T, Manns B, Reaume MN, Lloyd A, et al. Benefits and harms of erythropoiesis-stimulating agents for anemia related to cancer: a meta-analysis. *Canadian Medical Association Journal* 2009; **180**(11):62-71. [DOI: [10.1503/cmaj.090470](https://doi.org/10.1503/cmaj.090470)] [PMID: 19407261]

Trendall 2001

Trendall J. Assessing fatigue in patients with COPD. *Professional Nurse* 2001; **16**(7):1217-20. [PMID: 12026780]

Wagner 2004

Wagner LI, Cella D. Fatigue and cancer: causes, prevalence and treatment approaches. *British Journal of Cancer* 2004; **91**(5):822-8. [PMID: 15238987]

Walsh 2000

Walsh D, Donnelly S, Rybicki L. The symptoms of advanced cancer: relationship to age, gender, and performance status in 1000 patients. *Supportive Care in Cancer* 2000; **8**(3):175-9. [PMID: 10789956]

Zifko 2004

Zifko UA. Management of fatigue in patients with multiple sclerosis. *Drugs* 2004; **64**(12):1295-304. [PMID: 15200345]

References to other published versions of this review

Peuckmann-Post 2011

Peuckmann-Post V, Elsner F, Krumm N, Trottenberg P, Radbruch L. Pharmacological treatments for fatigue associated with palliative care. *Cochrane Database of Systematic Reviews* 2010, Issue 11. [DOI: [10.1002/14651858.CD006788.pub2](https://doi.org/10.1002/14651858.CD006788.pub2)]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Ashtari 2009

Study characteristics	
Methods	Placebo-controlled RCT
Participants	N = 42 patients with multiple sclerosis who were divided randomly into 2 groups

Pharmacological treatments for fatigue associated with palliative care (Review)

Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Ashtari 2009 (Continued)

Interventions	Amantadine 200 mg/day Duration: 2 months
Outcomes	FSS scores at baseline and 2 months later were compared in the 2 groups
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Identical preprinted medication code labels
Incomplete outcome data (attrition bias) All outcomes	Low risk	Withdrawals < 10%
Size of study	High risk	< 50 participants per treatment arm

Auret 2009
Study characteristics

Methods	Placebo-controlled, DB, RCT
Participants	N = 70 pts with advanced cancer and cancer considered, 50 included
Interventions	Dexamphetamine 10 mg/twice a day or placebo Duration: 8 days ITT basis
Outcomes	BFI, McGill Quality of Life Questionnaire, ECOG; side effects monitored. Pts were included if they reported fatigue as at least 4 of 10 on a 0 to 10 NRS. Reduction by 2 points on a 10-point scale (BFI) was considered a minimum important clinical difference
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described

Auret 2009 (Continued)

Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The active drug and the placebo were pre-packed in identical generic capsules
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Barak 2014

Study characteristics

Methods	Double-blind, randomised, placebo-controlled trial
Participants	N = 158 pts with multiple sclerosis (mean age 41.1 + 9.2 years)
Interventions	Patients were randomly assigned to receive alfacalcidol (1 µg) or placebo once daily for 6 months
Outcomes	FIS and QOL
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Baseline observation carried forward
Size of study	Unclear risk	Participants 50 to 199 per treatment arm

Brass 2001

Study characteristics

Pharmacological treatments for fatigue associated with palliative care (Review)

Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Brass 2001 (Continued)

Methods	Placebo-controlled, DB, RCT
Participants	N = 60 pts with end-stage renal disease (ESRD)
Interventions	Patients were randomised and divided into 2 groups 1:1; placebo and treatment Study A: treatment with L-carnitine 20 mg/kg IV for 24 weeks Study B: treatment with L-carnitine dose-ranging 10 to 20 to 40 mg/kg for 24 weeks
Outcomes	KDQ
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis; withdrawals < 10%
Size of study	High risk	< 50 participants per treatment arm

Breitbart 2001

Study characteristics

Methods	Placebo-controlled, DB, RCT
Participants	HIV pts with fatigue receiving ambulatory care
Interventions	Titration up to max. 60 mg methylphenidate or 150 mg pemoline, or 8 capsules placebo daily Duration 6 weeks
Outcomes	PFS, VAS-F; side effects monitored
Notes	The authors concluded that pts responded favourably to both treatments. However, while the PFS total score showed a significant difference, the VAS total fatigue scale (in which the energy score was significant only) did not. Therefore, the authors' conclusion seems to be too positive A confounding factor may have been improvement in fatigue due to an antidepressant effect, according to the authors

Risk of bias

Pharmacological treatments for fatigue associated with palliative care (Review)

Breitbart 2001 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Used block randomisation
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	All medications were prepared in identical capsules
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Bruera 2006
Study characteristics

Methods	Placebo-controlled, patient-controlled, DB, RCT
Participants	N = 112 cancer pts with a fatigue score of at least 4 on a scale of 0 to 10 were included
Interventions	Pts were randomised to methylphenidate 5 mg PO or placebo every 2 hours prn (max 4 times/day) for 1 week, followed by a 4-week open-label trial of methylphenidate
Outcomes	FACIT-F, ESAS, daily diary. The authors proposed to detect a decrease in fatigue in the methylphenidate group over and above the placebo group, of half the SD, or a score of approximately 7 on the FACIT-F scale. They therefore adjusted the sample size to declare this difference statistically significant, assuming a one-sided significance level of 0.05 and 80% power
Notes	The authors stated that a longer study duration of more than 1 week was justified

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Restricted random assignment with random balance points from 1 to 5 blocks
Allocation concealment (selection bias)	Low risk	A list of random assignments was prepared and the next eligible patient was entered on the next available assignment line
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis; withdrawals < 10%
Size of study	Unclear risk	Participants 50 to 199 per treatment arm

Bruera 2007

Study characteristics

Methods	PC, DB, RCT
Participants	Pts with fatigue score of at least 4 on a scale of 0 to 10 were included; 112 of 142 pts with cancer fatigue were assigned to treatment (71 each)
Interventions	Patients were randomised to donepezil 5 mg PO or placebo every morning for 1 week. Second week: open-label
Outcomes	FACIT-F, ESAS
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Restricted random assignment with random balance points from 1 to 5 blocks
Allocation concealment (selection bias)	Low risk	A list of random assignments was prepared and the next eligible patient was entered on the next available assignment line
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	Unclear risk	Participants 50 to 199 per treatment arm

Butler 2007

Study characteristics

Methods	PC, DB, RCT, phase III trial
Participants	N = 68 pts with primary or metastatic brain tumour scheduled to receive radiotherapy; 34 pts included in each group
Interventions	D-threo-methylphenidate HCl 5 mg/twice a day, titrated to max. 15 mg/twice a day, duration 12 weeks
Outcomes	Cognitive function, FACIT-F, Center for Epidemiologic Studies scale, MMSE, KPS Assessed at baseline, end of RT and 4, 8 and 12 weeks after brain RT
Notes	This study examines prophylactic treatment (before and after RT). The population was quite heterogeneous, since approximately half of the pts had metastatic disease and 75% received radiotherapy without chemotherapy. The age range was 28 to 83 years (median 52 for verum and 60 years for placebo group)

Butler 2007 (Continued)

There was a 71% drop-out from baseline (n = 33 methylphenidate group, n = 29 placebo group) until weeks 5 to 12 (n = 9 each group). Further, the post-treatment effect of radiotherapy during a 12-week course may vary concerning fatigue and pain, for example, which in part may be caused or relieved by the treatment

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Participants received a bottle of pills containing either the study drug or a matched placebo
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Canadian MSRG 1987
Study characteristics

Methods	PC, DB, RCT, cross-over; multicentre trial (11 MS centres)
Participants	N = 115 of 159 were randomised to treatment Eligible participants had at least a 6-month history of "definite multiple sclerosis" and a 3-month history of chronic, persistent, moderate to severe daily fatigue
Interventions	Amantadine 100 mg/twice a day or placebo 10 weeks duration 2 x 3-week treatment periods Each treatment period was preceded by a single-blind, 2-week placebo period
Outcomes	KEDSS; BDI; VAS-F
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described

Canadian MSRG 1987 (Continued)

Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	Unclear risk	Participants 50 to 199 per treatment arm

Cohen 1989
Study characteristics

Methods	DB, RCT, cross-over, single centre
Participants	N = 29 eligible participants had "satisfied criteria for a definite/probable diagnosis of multiple sclerosis" at least 6 months before diagnosed; all had daily symptomatic fatigue for at least 3 months
Interventions	Random assignment to 100 mg amantadine Hcl twice a day or placebo for 4 weeks, followed by a 2-week wash-out, then cross-over to an alternate treatment (verum/placebo) for another 4-week-period
Outcomes	Kurtzke rating; mBDI, "Stroop Interference Test" (attentional measure of freedom from distracting information). "Overall fatigue" was averaged across mean diary ratings for 7 indices of fatigue on a 5-point scale; 1 = poor; 5 = excellent)
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Cruciani 2009

Study characteristics

Methods	PC, DB, RCT
Participants	N = 149 eligible pts with carnitine deficiency, cancer and fatigue (n = 27 were excluded due to screen failure)
Interventions	L-carnitine (initial dose 0.5 g/day for 2 days, followed by 1 g/day for 2 days, then 2 g/day for 10 days or placebo), followed by an open-label phase, during which all pts received L-carnitine for 2 weeks
Outcomes	FACT-An, LASA, MMSE, KPS
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer program
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The liquid carnitine and placebo were prepared by a research pharmacist and were identical in appearance and taste
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Cruciani 2012

Study characteristics

Methods	PC, DB, RCT phase III trial
Participants	376 pts with cancer
Interventions	Pts were randomly assigned to either 2 g/day of L-carnitine oral supplementation or matching placebo for 4 weeks
Outcomes	BFI, FACIT-F
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement

Cruciani 2012 (Continued)

Random sequence generation (selection bias)	Low risk	Computerised randomisation system
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Drug and placebo matched in appearance
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	Unclear risk	Participants 50 to 199 per treatment arm

Della Cuna 1989
Study characteristics

Methods	PC, DB, RCT
Participants	403 pts with cancer Average age 62.7
Interventions	Methylprednisolone 125 mg/day IV 8 weeks Matching placebo
Outcomes	NOSIE, LASA, the Physicians' Global
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	Unclear risk	Participants 50 to 199 per treatment arm

Escalante 2014

Study characteristics

Methods	Placebo-controlled, DB, RCT - cross-over
Participants	N = 42 pts were enrolled and were diagnosed with lymphoma, myeloma or breast, gastrointestinal or lung cancers, and either undergoing chemotherapy or hormonal treatment or completed treatment in the previous 12 months
Interventions	The study duration was 4 weeks. Patients were randomised into 1 of 2 arms: Methylphenidate (18 mg/day) for 2 weeks followed by placebo for 2 weeks (arm A) or placebo for 2 weeks followed by methylphenidate (18 mg/day) for 2 weeks (arm B)
Outcomes	BFI at the end of each 2-week period
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Fisch 2003

Study characteristics

Methods	PC, DB, RCT
Participants	163 pts with cancer
Interventions	Fluoxetine 20 mg/day 12 weeks
Outcomes	FACT-G, BZSDS
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement

Fisch 2003 (Continued)

Random sequence generation (selection bias)	Low risk	Preprinted randomisation table
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	Unclear risk	Participants 50 to 199 per treatment arm

Jean-Pierre 2010
Study characteristics

Methods	Placebo-controlled, DB, RCT, phase III
Participants	877 cancer patients enrolled in the study N = 431 (modafinil) and n = 436 (placebo) Majority of sample were Caucasian and reported diverse marital status and educational level. 67% of the participants were females at baseline
Interventions	100 mg of modafinil or placebo on day 10 of the chemotherapy cycle, then increase to the full dose of 200 mg of modafinil or placebo after 3 days and continue on this regimen until day 7 of study cycle 4
Outcomes	BFI, ESS, POMS-DD Fatigue and depression were assessed during cycles 2 to 4 using psychometrically valid measures Group differences (treatment versus control) in the worst level of fatigue during the previous week at cycle 4 were examined using an analysis of covariance (ANCOVA) adjusting for baseline fatigue (cycle 2)
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number generator program
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias)	Low risk	Used 'baseline observation carried forward' analysis

Pharmacological treatments for fatigue associated with palliative care (Review)

Jean-Pierre 2010 (Continued)

All outcomes

Size of study	Low risk	> 200 participants per treatment arm

Kerr 2012

Study characteristics

Methods	Placebo-controlled, DB, RCT
Participants	N = 30 pts with advanced disease in hospice hospital 2 weeks duration
Interventions	Pts were randomly assigned to receive either 5 mg of methylphenidate or placebo at 8 am and 1 pm. Doses of methylphenidate were titrated every 3 days according to response and adverse effects. Home care patients were monitored daily by telephone and visited by a research nurse on study days 0 (baseline), 3, 7 and 14
Outcomes	PFS, VAS-F, ESAS, BDI-II, ESAS depression score, CESD
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number table
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Knapp 2008

Study characteristics

Methods	PC, DB, RCT
Participants	61 patients with HIV
Interventions	300 mg IM of testosterone enanthate or placebo for 16 weeks

Knapp 2008 (Continued)

Outcomes	Body composition, muscle strength, physical function and MOS-30	
Notes	—	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomisation list
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Krupp 1995

Study characteristics		
Methods	PC, DB, RCT	
Participants	N = 119 pts with multiple sclerosis and severe fatigue were eligible (FSS score at least 4 of 7; 9 items, each with potential score of 1 to 7; scoring is done by calculating the average response)	
Interventions	Pts were randomised to treatment with pemoline (18.75 mg/day for 1 week, 37.5 mg/day for week 2, and 56.25 mg weeks 3 to 6) or placebo or amantadine (amantadine 100 mg twice a day) for 6 weeks	
Outcomes	FSS, MS SFS, subjective response (verbal rating)	
Notes	Authors conclude that "amantadine was significantly better than placebo in treating fatigue in MS patients", however, this applies only to ratings on the MS Specific Fatigue Scale, not on the FSS. Interestingly, the authors chose a FSS cut-off score as an inclusion criterion, which highlights this scale as a key instrument. Thus, a clear effect of amantadine on fatigue seems questionable	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias)	Low risk	Drugs including the matching placebo were delivered by the pharmaceutical industry

Pharmacological treatments for fatigue associated with palliative care (Review)

Krupp 1995 (Continued)

All outcomes

Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Lacasse 2004
Study characteristics

Methods	PC, DB, RCT
Participants	N = 23 of 342 pts with end-stage COPD were found eligible (n = 82 refusals) 12-week duration
Interventions	Paroxetine was started at a dose of 5 mg/day, with weekly 5 mg increments up to a maximum of 20 mg/day "or the highest dose not associated with any side effect"
Outcomes	SF-36, "chronic respiratory questionnaire (CRQ)": 1 of 4 domains was fatigue (each domain includes 4 to 7 items and each item is scored on a 7-point-scale.); geriatric depression scale
Notes	342 pts were assessed for eligibility; 319 pts were excluded to ineligibility (237 pts) and refusal (82 pts) Primary outcome was quality of life Very low number of participants. Power analysis was conducted: 80% was reached at sample size of 9 to detect large treatment effects (type I error 0.05). This number has not been reached

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Low risk	Random numbers table was used to allocate participants to treatment
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Lange 2009
Study characteristics
Pharmacological treatments for fatigue associated with palliative care (Review)

Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Lange 2009 (Continued)

Methods	PC, DB, RCT
Participants	N = 21 pts with multiple sclerosis and fatigue 8-week duration
Interventions	Placebo or modafinil was started at a dose of 100 mg/day for the first week and 200 mg/day for the subsequent 7 weeks
Outcomes	FSS, FSS sum score was used (potential scores 4 to 63), alertness test, NHPT, TMS
Notes	The authors state that "the TMS subgroup consisted of 21 consecutive multiple sclerosis pts from a larger, randomised, double-blind trial"

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Lou 2009
Study characteristics

Methods	PC, DB, RCT pilot study
Participants	19 pts with Parkinson's disease Duration 8 weeks
Interventions	Participants took their regular medications and were randomly assigned to the treatment group (9 participants, modafinil 100 mg capsule twice a day PO) or placebo group (10 participants) twice a day for 8 weeks
Outcomes	MFI, ESS, CESD
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement

Lou 2009 (Continued)

Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Placebo group and treatment had the same appearance
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis; withdrawals < 10%
Size of study	High risk	< 50 participants per treatment arm

Moraska 2010

Study characteristics

Methods	PC, DB, RCT phase III study
Participants	148 pts with cancer were enrolled for this study
Interventions	Pts were randomly assigned to receive 1 tablet of methylphenidate or placebo on days 1 through 7, 2 tablets on days 8 through 14, and 3 tablets on days 15 through 28. Each methylphenidate tablet was 18 mg, resulting in the goal dose of 54 mg/day for the final 2 weeks of the study. Tablets were to be taken in the morning
Outcomes	BFI, SED, SF-36 Vitality Subscale, LASA, PSQI, SGIC
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomisation
Allocation concealment (selection bias)	Low risk	Centrally allocated randomisation
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	Unclear risk	Participants 50 to 199 per treatment arm

Morrow 2003

Study characteristics

Methods	PC, DB, RCT
Participants	N = 549 of 704 cancer patients with fatigue (recruited on days 1 to 3 undergoing chemotherapy)
Interventions	Paroxetine 20 mg/day oral or placebo for 8 weeks
Outcomes	FSC, MAF (total score was substituted by score of question no. 1; potential score 1 to 10), MPMM, CESD. Assessments were performed at cycles 3 and 4 of chemotherapy
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated random numbers table
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Medication and placebo were identical
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	Low risk	> 200 participants per treatment arm

Murray 1985

Study characteristics

Methods	PC, DB, CT, cross-over
Participants	N = 40 pts with multiple sclerosis and persistent fatigue of at least 3 months' duration
Interventions	Amantadine 100 mg or placebo twice a day 6 weeks "with one week wash-out period between active drug and placebo"
Outcomes	KDSS; improvement of fatigue measured by percentage of participants who felt improvement
Notes	The authors reported that "In only a few instances the same patients were used in more than one study". Date of study investigation was 1985 and methods may have been different

Risk of bias

Bias	Authors' judgement	Support for judgement

Murray 1985 (Continued)

Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Rabkin 2000
Study characteristics

Methods	PC, DB, RCT
Participants	N = 74 HIV-positive men with hypogonadal symptoms (diminished libido, depressed mood, low energy, depleted muscle mass) and symptomatic HIV illness
Interventions	Testosterone cypionate IM injections (initial dose 200 mg, increased to 400 mg) bi-weekly for 6 weeks, followed by 12-week open-label maintenance
Outcomes	CGI scale ratings for libido, mood, energy and erectile function; CFS (7-item scale measured by response options from 1 = never to 5 = always)
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated list of numbers
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Matching coded vials containing medication or placebo
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis; withdrawals < 10%
Size of study	High risk	< 50 participants per treatment arm

Rabkin 2004

Study characteristics

Methods	PC, DB, RCT
Participants	HIV-positive men N = 123 men were found eligible
Interventions	Testosterone cypionate IM injections (initial dose 200 mg, increased to 400 mg bi-weekly, or fluoxetine up to 60 mg/day (starting dose not mentioned; final mean dose 34 mg, range 20 mg to 40 mg)
Outcomes	CGI scale rating "was expanded to include ratings of energy, as well as mood and a global rating (CGI), significant improvement was defined as score of 1 or 2 on CGI"; HDRS; CFS scores (7-item scale measured by response options from 1 = never to 5 = always)
Notes	Overall, the authors did not recommend testosterone as first-line treatment for depressive disorders in HIV-positive men, but suggested further investigation in pts experiencing fatigue as well as depression

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated list of numbers
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Matching coded vials containing medication or placebo
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis; withdrawals < 10%
Size of study	High risk	< 50 participants per treatment arm

Rabkin 2009

Study characteristics

Methods	PC, DB, RCT
Participants	32 patients with amyotrophic lateral sclerosis (ALS)
Interventions	Modafinil 300 mg/day Duration 4 weeks
Outcomes	CGI improvement scale, FSS, ESS, BDI, RFS and VAS
Notes	—

Risk of bias

Rabkin 2009 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Block randomisation
Allocation concealment (selection bias)	Low risk	Centrally allocated randomisation
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Medication and placebo were identical
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Rabkin 2010
Study characteristics

Methods	PC, DB, RCT
Participants	115 pts with HIV/AIDS
Interventions	Modafinil with maximum dose 200 mg/day for 8 weeks
Outcomes	CGI improvement, FSS, HDRS, BDI. Safety assessment used assays of CD4 cell count and HIV RNA viral load
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated list of numbers
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Medication and placebo were identical in appearance
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	Unclear risk	Participants 50 to 199 per treatment arm

Rabkin 2011

Study characteristics

Methods	PC, DB, RCT
Participants	70 pts with HIV
Interventions	Maximum trial dose of armodafinil was 250 mg/day for 4 weeks
Outcomes	CGI, FSS, HDRS and BDI. Safety was assessed with assays of CD4 cell count and HIV RNA viral load and the SAFTEE side effects rating scale
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated list of numbers
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Medication and placebo were identical in appearance
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis; withdrawals < 10%
Size of study	High risk	< 50 participants per treatment arm

Rosenberg 1988

Study characteristics

Methods	PC, DB, RCT
Participants	N = 10 pts with multiple sclerosis
Interventions	Amantadine 200 mg/day or placebo for 1 week, followed by a 1-week wash-out, followed by a cross-over to another drug for 1 week
Outcomes	KEDSS (potential range 0 to 10; 0 = normal neurological examination, 10 = death), patient preference
Notes	Baseline values of KEDSS were not provided; very small participant number. No information concerning a power analysis given

Risk of bias

Bias	Authors' judgement	Support for judgement

Rosenberg 1988 (Continued)

Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Roth 2010
Study characteristics

Methods	PC, DB, RCT
Participants	N = 32 pts ambulatory with prostate cancer
Interventions	The patients were randomly allocated to receive either methylphenidate or placebo for a period of 6 weeks. Methylphenidate was administered in capsules containing 5 mg each, with a starting dose of 1 capsule in the morning The dose was increased by 1 capsule (5 mg) on day 3, added as a midday dose, if fatigue was not substantially reduced, there was no toxicity from the study treatment and if the patient was willing to increase the dose. Dosage was titrated upwards (or down) every 2 to 3 days to a maximum of 6 capsules daily, divided into morning and midday doses (equivalent to a total maximum daily dose of 30 mg of methylphenidate)
Outcomes	BFI
Notes	The authors suggest the need to monitor pulse and blood pressure during the treatment

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Medication and placebo were identically appearing capsules
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis

Roth 2010 (Continued)

Size of study	High risk	< 50 participants per treatment arm

Semiglazov 2006
Study characteristics

Methods	PC, DB, RCT
Participants	Pts with breast cancer N = 352 Average age 46.2
Interventions	Standardised mistletoe (PS76A) extract for 6 cycles of chemotherapy; matched placebo up to 18 weeks
Outcomes	FACT-G
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomisation list
Allocation concealment (selection bias)	Low risk	Centrally allocated randomisation
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Medication and placebo were identical in terms of appearance, colour and packaging
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	Unclear risk	Participants 50 to 199 per treatment arm

Shaygannejad 2012
Study characteristics

Methods	PC, DB, RCT, cross-over
Participants	N = 52 pts with multiple sclerosis Age range 21 to 53 years old
Interventions	The first group received amantadine (100 mg twice a day) for a total of 4 weeks. The second group received 500 mg/day acetylsalicylic acid (ASA) for 4 weeks. After a 2-week wash-out period, they crossed over to the alternative treatment for 4 weeks
Outcomes	FSS

Shaygannejad 2012 (Continued)

Notes —

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomisation list
Allocation concealment (selection bias)	Low risk	Group assignments were concealed in an opaque, sealed envelope
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis; withdrawals < 10%
Size of study	High risk	< 50 participants per treatment arm

Simons 1996

Study characteristics

Methods	PC, DB, RCT
Participants	N = 206 Male 153, female 53 Average age 64
Interventions	Medroxyprogesterone acetate (MPA) 500 mg twice a day for 12 weeks
Outcomes	EORTC-QLQ-C30 fatigue subscale
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Block randomisation in permutation blocks of 4
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias)	Low risk	Used 'baseline observation carried forward' analysis

Pharmacological treatments for fatigue associated with palliative care (Review)

Simons 1996 (Continued)

All outcomes

Size of study	Unclear risk	Participants 50 to 199 per treatment arm

Spathis 2014

Study characteristics

Methods	PC, DB, RCT
Participants	N = 208 pts with non-small cell lung cancer (NSCLC)
Interventions	Modafinil (100 mg on days 1 to 14; 200 mg on days 15 to 28) or matched placebo
Outcomes	FACIT-F, ESS, HADS, QOL-LAS
Notes	Future trials need to have sufficient power to evaluate the effect in those patients with severe fatigue and could incorporate a placebo wash-in period to minimise the influence of the placebo effect

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated
Allocation concealment (selection bias)	Low risk	Central telephone system
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The over-encapsulated active drugs and placebo capsules were matched
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	Unclear risk	Participants 50 to 199 per treatment arm

Stankoff 2005

Study characteristics

Methods	PC, DB, RCT
Participants	N = 115 pts with multiple sclerosis and "stable disability", minimal baseline score on MFIS of 45 N = 56 modafinil, N = 59 placebo
Interventions	Modafinil initial dose 200 mg for 1 week, increased by 100 mg weekly up to 400 mg/day max., then continued for 2 weeks or placebo 5-week duration

Stankoff 2005 (Continued)

Outcomes	MFIS; score range 0 to 84; lower score indicating less fatigue, VAS-F, ESS	
Notes	—	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Blocks of 4 randomisation
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	Unclear risk	Participants 50 to 199 per treatment arm

Stein 1995

Study characteristics		
Methods	PC, DB, RCT	
Participants	N = 25 pts with post-polio syndrome and fatigue were eligible	
Interventions	Amantadine 100 mg twice a day or placebo for 6 weeks	
Outcomes	FSS; improvement of fatigue as noted by participants given in percentage	
Notes	Very small participant numbers. Power analysis was calculated to detect 50% reduction in fatigue with 80% power and probability of type I error 0.05	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias)	Low risk	Used 'baseline observation carried forward' analysis

Stein 1995 (Continued)

All outcomes

Size of study	High risk	< 50 participants per treatment arm

Tomassini 2004

Study characteristics

Methods	DB, RCT, cross-over pilot trial
Participants	N = 36 pts with multiple sclerosis and fatigue
Interventions	Amantadine 100 mg twice a day or acetyl-L-carnitine (ALCAR) 1 g twice a day for 3 months, followed by a 3-month wash-out period, then followed by a cross-over to alternative treatment for 3 months
Outcomes	FSS, FIS, BDI, SEC; reduction of FSS scores observed in participants given in percentage
Notes	Inclusion of relapsing-remitting and secondary progressive multiple sclerosis

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Vasconcelos 2007

Study characteristics

Methods	DB, RCT, cross-over trial
Participants	N = 36 pts with post-polio syndrome N = 18 (modafinil) and n = 18 (placebo)
Interventions	Treatment with modafinil or equivalent placebo lasted for 6-week periods, being separated by a 14-day wash-out interval. First 3 weeks of treatment, patients received half of the targeted dose (i.e. 200 mg/day supplied as 2 x 100 mg capsules, 1 at breakfast and lunch). At the end of the 3rd week patients doubled their intake (i.e. 400 mg/day, 200 mg at breakfast and lunch)

Pharmacological treatments for fatigue associated with palliative care (Review)

Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Vasconcelos 2007 (Continued)

Outcomes	FSS, VAS-F, FIS, SF-36 QOL	
Notes	—	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated
Allocation concealment (selection bias)	Low risk	Concealed allocations from investigators by securing treatment codes
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Active drug and placebo had the same appearance
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Wagner 2000

Study characteristics		
Methods	PC, DB, RCT	
Participants	N = 23 pts with HIV, depression and debilitating fatigue	
Interventions	Dextroamphetamine initial dose 2.5 mg twice a day, titrated up to maximum of 40 mg/day. Mean dose was 26 +/- 12 mg daily, range 10 to 40 mg/day or placebo for 2 weeks, followed by 24 weeks of open-label trial	
Outcomes	SCID; HDRS, BFI, BHS, VAS for mood, CFS (possible range 0 to 28), VAS for energy level	
Notes	Groups were heterogenous concerning major depression, which was diagnosed in 12 of the 23 participants	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described

Wagner 2000 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis; withdrawals < 10%
Size of study	High risk	< 50 participants per treatment arm

Weinshenker 1992

Study characteristics		
Methods	DB, RCT, cross-over; 2-centre trial	
Participants	N = 46 pts with multiple sclerosis and fatigue (patients with multiple sclerosis exacerbation during study period were excluded)	
Interventions	Pemoline titration in 1 week (18.75 mg to 75 mg/day), continued for 3 weeks or placebo, followed by a cross-over to the other treatment	
Outcomes	VAS-F, KEDSS, mBDI; tolerance of adverse effects and checklist to identify their nature	
Notes	Interestingly, the authors noted increasing benefit from pemoline over the 2-week study period and supposed that there may have been a larger treatment effect if the study period was longer. Of 13 patients who chose to continue taking the drug, 7 were still taking it after 1 year and reported benefit	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Used 'last observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Westman 1999

Study characteristics	
Methods	DB, RCT, cross-over trial
Participants	N = 255 pts with cancer Male 134, female 121

Westman 1999 (Continued)

Average age 70 (not on treatment)

Interventions	Megestrol acetate (MA) 320 mg/day
Outcomes	EORTC-QLQ-C30
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Block randomisation in permutation blocks of four
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Active drugs and placebo had the same shape, size, appearance, colour and taste
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis; withdrawals < 10%
Size of study	Unclear risk	Participants 50 to 199 per treatment arm

Wingerchuk 2005

Study characteristics

Methods	PC, DB, RCT, cross-over
Participants	N = 30 pts with multiple sclerosis and fatigue
Interventions	Acetylsalicylic acid (ASA) 1300 mg/day or placebo for 6 weeks, followed by 2-week wash-out period, followed by cross-over to alternative treatment
Outcomes	MFIS (possible score 0 to 84), KEDSS, CESD, Global Fatigue Change self assessment, VAS
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Generated random number list
Allocation concealment (selection bias)	Low risk	Allocation concealment from investigators by securing the results in the pharmacy
Blinding of outcome assessment (detection bias)	Low risk	Active drug and placebo had the same appearance

Wingerchuk 2005 (Continued)

All outcomes

Incomplete outcome data (attrition bias) All outcomes	Low risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

Yennurajalingam 2013
Study characteristics

Methods	PC, DB, RCT
Participants	N = 84 pts with advanced cancer (dexamethasone = 43, placebo = 41)
Interventions	Patients were randomly assigned to either dexamethasone 4 mg or placebo orally twice per day for 14 days
Outcomes	ESAS, FACIT-F, HADS, FAACT
Notes	—

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not described
Allocation concealment (selection bias)	Unclear risk	Method not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Method not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Used 'baseline observation carried forward' analysis
Size of study	High risk	< 50 participants per treatment arm

BDI = Beck Depression Inventory; BFI = Brief Fatigue Inventory; BHS = Beck Hopelessness Scale; BZSDS = Brief Zung Self-Rating Depression Scale; CESD = Center for Epidemiologic Studies Depression; CFS = Chalder Fatigue Scale; CRQ = Chronic Respiratory Questionnaire; DB = double-blind; ECOG = performance status according to the Eastern Cooperative Oncology Group Scale; EORTC-QLQ-C30 = European Organization for Research and Treatment of Cancer Quality of Life Questionnaire; ESAS = Edmonton Symptom Assessment System; ESS = Epworth Sleepiness Scale; FAACT = Functional Assessment of Cancer Therapy-Anorexia-Cachexia; FACIT-F = Functional Assessment for Chronic Illness Therapy - Fatigue; FACT-An = Functional Assessment of Cancer Therapy - Anaemia; FIS = Fatigue Impact Scale; FSC = Fatigue Symptom Checklist; FSS = Fatigue Severity Scale; HADS = Hospital Anxiety and Depression Scale; HDRS = Hamilton Depression Rating Scale; IM = intramuscular; ITT = intention-to-treat; IV = intravenous; KDQ = Kidney Disease Questionnaire; KEDSS = Kurtzke Expanded Disability Status Scale; KPS = Karnofsky Performance Status; LASA = Linear Analogue Scale Assessments; MAF = Multidimensional Assessment of Fatigue; mBDI = Modified Beck Depression Inventory; MFI = Multidimensional Fatigue Inventory; MMSE = Mini Mental State Examination; MOS-30 = Medical Outcomes Study-Short Form 30; MPMM = Monopolar Profile of Mood States; MS = multiple sclerosis; MS SFS = Multiple Sclerosis Specific Fatigue Scale; NHPT = Nine Hole Peg Test; NOSIE = Nurses' Observational Scale for Inpatient Evaluation; NRS = numeric

rating scale; PC = placebo-controlled; PFS = Piper Fatigue Scale; PO = per oral; POMS-DD = Depression-Dejection subscale of the Profile of Mood States; prn = as needed; PSQI = Pittsburgh Sleep Quality Index; pts = participants; QOL = RAYS quality of life; QOL-LAS = quality of life linear analogue scale; RCT = randomised controlled trial; RFS = Role Function Scale; RT = radiation therapy; SAFTEE = Systematic Assessment for Treatment Emergent Events; sc = subcutaneous; SCID = depression according to Structured Clinical Interview for DSM-IV; SD = standard deviation; SE = standard error; SEC = Social Experience Checklist; SED = Symptom Experience Diary; SF-36 = Short Form-36; SGIC = Subject Global Impression of Change; TMS = transcranial magnetic Social Experience Checklist; VAS = visual analogue scale; VAS-F = visual analogue scale - fatigue

Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
Agteresch 2000	Open-label study
Bruera 1987	Focus of this study was sedation related to narcotics, not fatigue
Bruera 1998	Focus of study was cancer cachexia
Bruera 2003	Focus of this study was sedation related to narcotics, not fatigue
Capuron 2002	Main focus of this study was not fatigue
Carter 2005	Open-label study, case series
Cerchietti 2004	Not blinded, not a RCT
Cruciani 2004	Open-label study
Cruciani 2006	Phase I/II study
Cueva 2012	Phase II study
Cullum 2004	Open-label study, case series
De Conno 1998	Main focus of this study was depression and appetite
Diel 2004	Part of the study was open-label
Downer 1993	Main focus of this study was cachexia
Dunlop 2007	Focus of this study was depression
Gehring 2012	Open-label study
Graziano 2002	The drug was intended to treat fatigue, which was induced by a treatment not a disease
Hannestad 2011	The participants were not at an advanced stage of illness or in a palliative situation
Hovey 2014	The drug was intended to treat fatigue, which was induced by a treatment not a disease
Inoue 2003	The drug was intended to treat fatigue, which was induced by a treatment not a disease
Lauretti 2013	Focus of this study was pain
Laval 2008	This article was published only as a protocol; no original results
Lower 2009	The drug was intended to treat fatigue, which was induced by a treatment not a disease

Study	Reason for exclusion
Mar Fan 2008	The drug was intended to treat fatigue, which was induced by a treatment not a disease
McElhiney 2010	Primary focus of this study was not fatigue
McElhiney 2013	Main focus of this study was cognitive function
Mercadante 2001	Consecutive study, not randomised
Metz 2004	No randomisation
Moertel 1974	Single-blind study
Mohr 2003	Fatigue was not the primary outcome
Monk 2006	Open-label study
Moss 2006	Open-label study, case series
Popiela 1989	Single-blind study
Rabkin 2000a	Investigators used pre-selection ("mood responders were maintained" after an open-label trial)
Rabkin 2011a	This study was a subset of another study
Rammohan 2002	Phase II study
Romani 2004	Authors state that "Due to the design of the study, [the similarly decreased fatigue scores in both groups] cannot be disjoined from a placebo effect."
Roscoe 2005	Patients cannot be considered to be in a palliative care situation
Sailer 2000	24 patients were assigned to 4 groups and patients were very heterogenous concerning disease duration
Shaw 2006	Phase II study
Shaw 2013	Phase II study
Spathis 2009	Open-label trial; aim of study was to determine feasibility of conducting a RCT for fatigue in lung cancer
Stockler 2007	Fatigue was studied as secondary outcome
Torta 2007	Main focus was depression. Fatigue was secondary outcome
Wade 2002	Investigators focus on disability, not fatigue
Wagner 1997	Open-label study, case series
Weitzner 2002	Fatigue was not the primary outcome
Wilwerding 1995	Focus was not fatigue, but narcotic-induced sedation
Zifko 2002	Open-label study

RCT: randomised controlled trial

DATA AND ANALYSES

Comparison 1. Methylphenidate versus placebo in cancer

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.1 FACIT-F score change	2		Std. Mean Difference (IV, Fixed, 95% CI)	Totals not selected

Analysis 1.1. Comparison 1: Methylphenidate versus placebo in cancer, Outcome 1: FACIT-F score change

Study or Subgroup	Methylphenidate			Placebo			Std. Mean Difference IV, Fixed, 95% CI	Std. Mean Difference IV, Fixed, 95% CI
	Mean	SD	Total	Mean	SD	Total		
Bruera 2006	9.6	9.8	52	7.5	11.3	53	0.20 [-0.19 , 0.58]	
Butler 2007	6.5	2.28	20	2.8	2.66	21	1.46 [0.76 , 2.16]	

Comparison 2. Modafinil versus placebo in multiple sclerosis

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.1 Fatigue score change	2		Std. Mean Difference (IV, Fixed, 95% CI)	Totals not selected

Analysis 2.1. Comparison 2: Modafinil versus placebo in multiple sclerosis, Outcome 1: Fatigue score change

Study or Subgroup	Modafinil			Placebo			Std. Mean Difference IV, Fixed, 95% CI	Std. Mean Difference IV, Fixed, 95% CI
	Mean	SD	Total	Mean	SD	Total		
Lange 2009	13	7.7	11	-0.4	8.5	10	1.59 [0.58 , 2.60]	
Stankoff 2005	8.7	13.64	56	13.7	13.49	59	-0.37 [-0.74 , 0.00]	

Comparison 3. Pemoline versus placebo in multiple sclerosis

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
3.1 Fatigue score change	2		Mean Difference (IV, Fixed, 95% CI)	Totals not selected

Analysis 3.1. Comparison 3: Pemoline versus placebo in multiple sclerosis, Outcome 1: Fatigue score change

ADDITIONAL TABLES

Table 1. Drug treatment of fatigue: Populations, substances and outcome measures

Patient pop- ulation	Substance	Outcome measures	Study
Fatigue in advanced cancer	Amphetamine	BFI, McGill Quality of Life Questionnaire, ECOG	Auret 2009
	L-carnitine	FACT-An, LASA, MMSE, KPS	Cruciani 2009
	L-carnitine	BFI, FACIT-F, depression and pain instrument	Cruciani 2012
	Paroxetine	FSC, MFI, Monopolar Profile of Mood States	Morrow 2003
	Methylphenidate	FACIT-F, ESAS, daily diary	Bruera 2006
	Methylphenidate	Cognitive function, FACIT-F, CESD, MMSE, KPS	Butler 2007
	Methylphenidate	BFI, SED, SF-36 vitality subscale, LASA, PSQI, SGIC	Moraska 2010
	Methylphenidate	BFI	Roth 2010
	Methylphenidate	BFI	Escalante 2014
	Donepezil	FACIT-F, ESAS	Bruera 2007
	Methylprednisolone	NOSIE, LASA, Physician's Global	Della Cuna 1989
	Modafinil	BFI, ESS, POMS-DD	Jean-Pierre 2010
	Modafinil	FACIT, ESS, HADS, QOL-LAS	Spathis 2014
	Fluoxetine	FACT-G	Fisch 2003
	Standardised Mistle- toe extract	FACT-G	Semiglazov 2006
	Medroxyprogesterone acetate (MPA)	FACT-G	Simons 1996
	Megestrol acetate	EORTC-QLQ-C30 instrument	Westman 1999

Table 1. Drug treatment of fatigue: Populations, substances and outcome measures *(Continued)*

Fatigue in amyotrophic lateral sclerosis (ALS)	Modafinil	CGI-E, FSS, ESS, BDI, RFS, VAS	Rabkin 2009
Fatigue in HIV/AIDS	Methylphenidate	PFS	Breitbart 2001
	Pemoline	PFS	Breitbart 2001
	Armodafinil	CGI-I, FSS, HDRS, BDI	Rabkin 2011
	Modafinil	CGI-I, FSS, HDRS, BDI	
	Amphetamine	SCID; HDRS, BFI, BHS, VAS for mood, CFS, VAS for energy level	Wagner 2000
	Testosterone	CGIS "was expanded to include ratings of energy, as well as mood and a global rating. Significant improvement was defined as score of 1 or 2 on CGIS"; HDRS, CFS	Rabkin 2004
	Testosterone	CGIS ratings for libido, mood, energy and erectile function; CFS	Rabkin 2000
	Testosterone	Body composition, muscle strength and physical function, MOS-30	Knapp 2008
	Fluoxetine	Body composition, muscle strength and physical function, MOS-30	Rabkin 2004
Fatigue in multiple sclerosis	Pemoline	VAS-F, KEDSS, mBDI; tolerance of adverse effects and checklist to identify their nature	Weinshenker 1992
	Pemoline	FSS, MS Specific Fatigue Scale, subjective response (verbal rating)	Krupp 1995
	L-carnitine	FSS, Fatigue Impact Scale, Beck Depression Inventory, Social Experience Checklist	Tomassini 2004
	Amantadine	FSS, Fatigue Impact Scale, Beck Depression Inventory, Social Experience Checklist	Tomassini 2004
	Amantadine	KEDSS; mBDI, "Stroop Interference Test" (attentional measure of freedom from distracting information)	Cohen 1989
	Amantadine	KEDSS; mBDI, "Stroop Interference Test" (attentional measure of freedom from distracting information)	Krupp 1995
	Amantadine	KEDSS	Murray 1985
	Amantadine	KEDSS, patient preference	Rosenberg 1988
	Amantadine	KEDSS; BDI; VAS-F	Canadian MSRG 1987
	Amantadine	FSS	Ashtari 2009
	Amantadine	FSS	Shaygannejad 2012

Table 1. Drug treatment of fatigue: Populations, substances and outcome measures (Continued)

	Acetylsalicylic acid	MFIS, KEDSS, CESD, SGIC, FSS	Wingerchuk 2005
	Acetylsalicylic acid	FSS	Shaygannejad 2012
	Modafinil	FSS, alertness test, NHPT, TMS	Lange 2009
	Modafinil	MFIS, VAS-F, ESS	Stankoff 2005
	Modafinil	FSS, VAS-F, FIS, SF-36 QOL	Vasconcelos 2007
	Dexamethasone	FIS, QOL	Yennurajalingam 2013
	Alfacalcidol (Vit. D)	FIS, QOL	Barak 2014
End-stage renal disease (ESRD)	L-carnitine	KDQ	Brass 2001
Parkinson's disease	Modafinil	MFI, ESS, CESD	Lou 2009
Multi-type advanced disease (hospice patients)	Methylphenidate	PFS, VAS-F, ESAS	Kerr 2012
End-stage COPD	Paroxetine	SF-36, chronic respiratory questionnaire: 1 of 4 domains was fatigue, geriatric depression scale	Lacasse 2004
Postpolio syndrome	Amantadine	FSS	Stein 1995

BDI = Beck Depression Inventory; BFI = Brief Fatigue Inventory; BHS = Beck Hopelessness Scale; CESD = Center for Epidemiologic Studies Depression; CFS = Chalder Fatigue Scale; DB = double-blind; ECOG = performance status according to the Eastern Cooperative Oncology Group Scale; EORTC-QLQ-C30 = European Organization for Research and Treatment of Cancer Quality of Life Questionnaire; ESAS = Edmonton Symptom Assessment System; ESS = Epworth Sleepiness Scale; FACIT-F = Functional Assessment for Chronic Illness Therapy - Fatigue; FACT-An = Functional Assessment of Cancer Therapy - Anaemia; FIS = Fatigue Impact Scale; FSC = Fatigue Symptom Checklist; FSS = Fatigue Severity Scale; HADS = Hospital Anxiety and Depression Scale; HDRS = Hamilton Depression Rating Scale; IM = intramuscular; ITT = intention-to-treat; IV = intravenous; KDQ = Kidney Disease Questionnaire; KEDSS = Kurtzke Expanded Disability Status Scale; KPS = Karnofsky Performance Status; LASA = Linear Analogue Scale Assessments; MAF = Multidimensional Assessment of Fatigue; mBDI = Modified Beck Depression Inventory; MFI = Multidimensional Fatigue Inventory; MMSE = Mini Mental State Examination; MOS-30 = Medical Outcomes Study-Short Form 30; MPMM = Monopolar Profile of Mood States; MS = multiple sclerosis; MS SFS = Multiple Sclerosis Specific Fatigue Scale; NHPT = Nine Hole Peg Test; NOSIE = Nurses' Observational Scale for Inpatient Evaluation; NRS = numeric rating scale; PC = placebo-controlled; PFS = Piper Fatigue Scale; PO = per oral; POMS-DD = Depression-Dejection subscale of the Profile of Mood States; prn = as needed; PSQI = Pittsburgh Sleep Quality Index; pts = participants; QOL = RAYS quality of life; QOL-LAS = quality of life linear analogue scale; RCT = randomised controlled trial; RFS = Role Function Scale; RT = radiation therapy; sc = subcutaneous; SCID = depression according to Structured Clinical Interview for DSM-IV; SD = standard deviation; SE = standard error; SEC = Social Experience Checklist; SED = Symptom Experience Diary; SF-36 = Short Form-36; SGIC = Subject Global Impression of Change; TMS = transcranial magnetic Social Experience Checklist; VAS = visual analogue scale; VAS-F = visual analogue scale - fatigue

Table 2. Overview of adverse reactions associated with fatigue treatment

Drug and indication	Impact of adverse reaction	Symptoms	Study
Acetyl-L-carnitine and amantadine in multiple sclerosis	1 (carnitine) and 5 (amantadine) of 36 patients withdrew	Carnitine: insomnia, nervousness Amantadine: nausea, dizziness	Tomassini 2004
Amantadine in multiple sclerosis	7 of 32 reported adverse reactions (versus 6 of 32 on placebo)	Hallucinations, nausea, gastric irritation, early morning wakening, hyperactivity, flu-like illness	Murray 1985
Amantadine in multiple sclerosis	—	Constipation, nausea, anxiety, influenza-like illness	Cohen 1989
Amantadine, pemoline in multiple sclerosis	No severe adverse reaction (did not lead to withdrawal)	Pemoline: palpitations, nausea, mood change, sleep disturbance, Amantadine: sleep disturbance, palpitations	Krupp 1995
Dextroamphetamine in HIV-positive men with fatigue	None of 23 patients discontinued during the 2 weeks of DB, PC RCT	Overstimulation, heart palpitation, sleep deprivation, loss of appetite and/or weight, headache. The authors reported that adverse reactions generally were "transient, reversible, and well managed with dose reduction; no serious medical side effects were reported"	Wagner 2000
Dexamphetamine in advanced cancer	Drop-out rate in verum group compared to placebo group was statistically not significant	Significant increase in pulse rate. Transient increase of dry mouth, insomnia, tremor, anorexia on days 6 and 8 (of 8)	Auret 2009
Pemoline in multiple sclerosis	3 of 46 dropped out because of adverse reaction; 25% of patients did not tolerate pemoline well according to the authors	Most common side effects were anorexia, irritability, "jitteriness" and insomnia	Weinshenker 1992
Acetylsalicylic acid in multiple sclerosis	No serious adverse reaction	Nausea, transient epigastric pain (nausea, headache and diarrhoea occurred with placebo)	Wingerchuk 2005
Methylphenidate and pemoline in HIV	Severe side effects relatively uncommon; 5 withdrew (2 on methylphenidate, 2 on pemoline, 1 on placebo)	Hyperactivity, jitteriness, dry mouth, rapid heart beat, difficulty sleeping, constipation, neuropathic pain (1 patient)	Breitbart 2001
Methylphenidate in advanced cancer	No severe adverse effects	Most pts reported insomnia, anorexia, restlessness, behavioural change and vertigo. Numbers in placebo groups were similar	Bruera 2006
Donepezil in advanced cancer	No severe adverse effects	Most pts reported anorexia, nausea, restlessness, dizziness, behaviour change, vertigo	Bruera 2007
Amantadine in multiple sclerosis	No statistically significant difference in prevalence of adverse events reporting	Most pts reported headache, insomnia, nausea, anxiety, dizziness, ataxia	Canadian MSRG 1987
Paroxetine in end-stage COPD	Only 1 of 12 pts dropped out because of adverse effects related to the drug	Somnolence was reported by 5 pts, constipation, nausea, headache tremor were rated by 2 pts each	Lacasse 2004
Testosterone for HIV-positive pts	3 of 46 pts discontinued because of adverse reactions	Acne, irritability, insomnia, headache, nasal congestion	Rabkin 2000

Table 2. Overview of adverse reactions associated with fatigue treatment (Continued)

Testosterone and fluoxetine in HIV	—	Fluoxetine: diarrhoea, nausea, nonspecific discomfort Testosterone: sleepiness; all treatments: dry mouth, sleepiness, loose bowels	Rabkin 2004
Modafinil for HIV patients	—	Treatment-emergent side effects were relatively uncommon CD4 cell count did not show either statistically or clinically significant changes	Rabkin 2010
Armodafinil in HIV	Armodafinil was well tolerated, with few and transient adverse events	The most common was headache	Rabkin 2011
Amantadine in post-polio syndrome	—	Insomnia was the commonest reported adverse reaction; dry mouth was noted by 1 patient	Stein 1995
L-carnitine in end-stage renal disease (ESRD)	No serious adverse event was believed by the investigators to be certainly or probably drug-related	Flu syndrome, injection-site reaction, pain, pharyngitis, headache and hypertension showed no difference in frequency between L-carnitine and placebo	Brass 2001
Methylphenidate in cancer	No serious adverse effect	Some pts experienced nausea, vomiting, facial rash	Escalante 2014
Fluoxetine in cancer	A total of 16 adverse events were judged to be possibly due to the treatment and 6 to taking the placebo	Allergic reaction, cardiac arrhythmia, dyspnoea, headaches, dizziness, mood change, myalgia, fever, diarrhoea, abdominal pain	Jean-Pierre 2010
Methylphenidate in cancer	There was a significant difference in self reported toxicities (SED)	Increased levels of nervousness and appetite loss in the methylphenidate group	Moraska 2010
Methylphenidate in cancer	No severe adverse reaction	4 (31%) men who started receiving methylphenidate had to be discontinued from the study due to increased blood pressure	Roth 2010
Medroxyprogesterone acetate (MPA) in cancer	—	Nausea and vomiting	Simons 1996
Acetylsalicylic acid and amantadine in multiple sclerosis	Acetylsalicylic acid and amantadine are both well tolerated	The most common side effects of acetylsalicylic acid were nausea (n = 53) and transient epigastric pain (n = 51). The most common side effect of amantadine was also nausea (n = 51)	Shaygannejad 2012
Modafinil in cancer	Modafinil seemed to be well tolerated	Most symptoms are nausea, vomiting, anxiety and headache	Spathis 2014
Dexamethasone in multiple sclerosis	2 adverse events were probably related to study treatment	Pain grade 3 and vomiting grade 2	Yennurajalingam 2013

COPD = chronic obstructive pulmonary disease; DB = double-blind; PC = placebo-controlled; pts = participants RCT = randomised controlled trial

APPENDICES

Appendix 1. CENTRAL search strategy

Subject search was run on 28 April 2014

#	1.	NEOPLASMS
#	2.	(neoplasm* or cancer* or carcinoma* or tumour* or tumor* or adenocarcinoma* or leukeni* or lymphoma* or malignan*):ti,ab,kw
#	3.	MULTIPLE SCLEROSIS
#	4.	"multiple sclerosis":ti,ab,kw
#	5.	AMYOTROPHIC LATERAL SCLEROSIS
#	6.	"amyotrophic lateral sclerosis":ti,ab,kw
#	7.	ACQUIRED IMMUNODEFICIENCY SYNDROME
#	8.	AIDS-RELATED COMPLEX
#	9.	HIV
#	10.	HIV WASTING SYNDROME
#	11.	"acquired immunodeficiency syndrome":ti,ab,kw
#	12.	"AIDS related complex":ti,ab,kw
#	13.	"HIV":ti,ab,kw
#	14.	"human immunodeficiency virus":ti,ab,kw
#	15.	LUNG DISEASES
#	16.	PULMONARY DISEASE CHRONIC OBSTRUCTIVE
#	17.	HEART DISEASES
#	18.	"lung disease*" or "heart disease*" or "pulmonary disease*"
#	19.	HEART FAILURE
#	20.	"cardiac failure" or "heart failure"
#	21.	"incurable disease*" or "incurable illness*"
#	22.	((terminal or advanced or progressive or "end stage" or end-stage) and (illness* or disease*))
#	23.	BONE MARROW TRANSPLANTATION
#	24.	(neutropeni* or neutropaeni*):ti,ab,kw

(Continued)

#	25.	RADIOTHERAPY
#	26.	(radioth* or radiat* or irradiat* or radiochemo* or chemotherapy*):ti,ab,kw
#	27.	“bone marrow” NEAR transplant*
#	28.	“bone-marrow” NEAR transplant*
#	29.	(fatigue near/4 treatment) or (treated near/4 treated) or (fatigue near/4 therapy) or (fatigue near/4 intervention)
#	30.	(asthenia near/4 treatment) or (treated near/4 treated) or (fatigue near/4 therapy) or (fatigue near/4 intervention)
#	31.	((tired* near/4 treatment) or (tired* near/4 treated) or (tired* near/4 therapy) or (treatment near/4 exhausted) or (exhausted near/4 therapy) or (exhaustion near/4 therapy))
#	32.	apathy or apathetic or lassitude or lethargy* or “feeling drained” or “feeling sleepy” or “feeling sluggish” or “feeling weak”
#	33.	(FATIGUE or cancer-related fatigue or cancer related fatigue):ti,ab,kw
#	34.	tired* or weary or weariness or exhaustion or exhausted or lackluster or astheni* or asthenia*
#	35.	lack* NEAR/2 energy
#	36.	lack* NEAR/2 vigour
#	37.	lack* NEAR/2 vigor
#	38.	loss NEAR/2 energy
#	39.	loss NEAR/2 vigour
#	40.	loss NEAR/2 vigor
#	41.	lost NEAR/2 energy
#	42.	lost NEAR/2 vigour
#	43.	lost NEAR/2 vigor
#	44.	palliati* or hospice or “end of life”
#	45.	#1-#28/OR
#	46.	#29-#44
#	47.	#45 AND #46
#	48.	dexamphetamine or dextroamphetamine
#	49.	ANTIDEPRESSIVE AGENTS
#	50.	ADRENERGIC UPTAKE INHIBITORS

(Continued)

#	51.	glucocorticoid or steroid or steroids or dexamethason or dexamethasone or methylprednisolone or corticosteroid or corticosteroids
#	52	Carnitine or L-carnitine
#	53.	modafinil
#	54.	amantadine or donepezil or pemoline
#	55.	Methylphenidate or d-threo-methylphenidate
#	56.	paroxetine
#	57.	Aspirin or acetylsalicylic acid
#	58.	citalopram
#	59.	bupropion
#	60.	testosterone cypionate
#	61.	#48 - #60/OR
#	62.	#47 AND #61

Appendix 2. MEDLINE search strategy

Subject search combined with the recommended study design filter was run on 28 April 2014 (using OVID)

#	1.	exp Neoplasms/
#	2.	(cancer* or malignan*).mp.
#	3.	multiple sclerosis.af.
#	4.	amyotrophic lateral sclerosis.af.
#	5.	(sclerosis adj4 (amyotroph* or multiple)).mp.
#	6.	exp acquired immune deficiency syndrome/
#	7.	aids related complex.mp.
#	8.	exp Human immunodeficiency virus/
#	9.	(HIV or AIDS).mp.
#	10.	bone marrow transplantation.mp.
#	11.	radiotherapy.mp.
#	12.	(carcinoma* or tumour* or adenocarcinoma* or leukeni* or leukaemi* or lymphoma* or tumor*).mp

(Continued)

#	13.	(neutropeni\$ or neutropaeni\$).mp.
#	14.	(radioth* or radiat* or irradiat* or radiochemo* or chemotherap*).mp.
#	15.	Lung Disorders.mp.
#	16.	heart disorders.mp.
#	17.	congestive heart failure.mp.
#	18.	cor pulmonale.mp.
#	19.	cancer fatigue.mp.
#	20.	(cancer-related fatigue or cancer related fatigue).mp.
#	21.	exp FATIGUE/
#	22.	(tired\$ or weary or weariness or exhaustion or exhausted or lacklustred or ((astheni\$ or asthenia\$) and syndrome)).mp.
#	23.	((lack\$ or loss or lost) adj2 (energy or vigour or vigor)).mp.
#	24.	(apathy or apathetic or lassitude or letharg\$ or (feeling adj3 (drained or sleepy or sluggish or weak\$))).mp.
#	25.	((advanced or terminal* or progressi* or end-stage or endstage or "endstage") adj6 (disease or illness)).mp.
#	26.	palliati*.mp.
#	27.	(dexamphetamine or dextroamphetamine).mp.
#	28.	(methylphenidate or threo\$methylphenidate or d-threo-methylphenidate).mp.
#	29.	donepezil.mp.
#	30.	amantadine.mp.
#	31.	(Carnitine or L-carntine).mp.
#	32.	pemoline.mp.
#	33.	modafinil.mp.
#	34.	paroxetine.mp.
#	35.	(testosteronecypionate or androgen).mp.
#	36.	(acetylsalicylic acid or aspirin).mp.
#	37.	fluoxetine.mp
#	38.	citalopram.mp

(Continued)

#	39.	(glucocorticoid or steroid or steroids or dexamethason or dexamethasone or methylprednisolone or corticosteroid or corticosteroids).mp.
#	40.	bupropion.mp.
#	41.	(anti depressant agents or anti depressive agents or Selective serotonin reuptake inhibitors).mp.
#	42.	random*.ti,ab.
#	43.	factorial*.ti,ab.
#	44.	(crossover* or cross over* or cross-over*).ti,ab.
#	45.	placebo*.ti,ab.
#	46.	(doubl* adj blind*).ti,ab.
#	47.	(singl* adj blind*).ti,ab.
#	48.	randomized controlled trial.mp.
#	49.	assign*.ti,ab.
#	50.	allocat*.ti,ab.
#	51.	evaluation study*.ti,ab.
#	52.	prospective study*.ti,ab.
#	53.	comparative study*.ti,ab.
#	54.	(animal* or nonhuman* or animal experiment*).ti,ab.
#	55.	#1-#18/OR
#	56.	#19-26/OR
#	57.	#27-41/OR
#	58.	#42-53/OR
#	59.	#55 AND #56 AND #57 AND #58
#	60.	#59 NOT #54

Appendix 3. EMBASE search strategy

Subject search combined with the recommended study design filter was run on 28 April 2014

#	1.	neoplasm/exp
#	2.	(cancer* or malignan*)

(Continued)

#	3.	'multiple sclerosis'/exp
#	4.	'amyotrophic lateral sclerosis'/exp
#	5.	sclerosis (amyotroph*4 or multiple)
#	6.	((('acquired immune deficiency syndrome'/exp) OR ('aids related complex'/exp))
#	7.	'Human immunodeficiency virus'/exp
#	8.	(HIV or AIDS)
#	9.	'lung disease'/exp
#	10.	'heart disease'/exp
#	11.	'cor pulmonale'/exp
#	12.	'congestive heart failure'/exp
#	13.	(neoplasm* or cancer* or carcinoma* or tumour* or adenocarcinoma* or leukemia* or leukaemia* or lymphoma* or tumor* or tumour* or malignan*):ti,ab
#	14.	(neoplasm* or cancer* or carcinoma* or tumour* or adenocarcinoma* or leukemia* or leukaemia* or lymphoma* or tumor* or tumour* or malignan*)/mj
#	15.	(neutropeni* or neutropaeni*):ti,ab
#	16.	(neutropeni* or neutropaeni*)/mj
#	17.	RADIOTHERAPY/exp
#	18.	(radioth* or radiat* or irradiat* or radiochemo* or chemotherapy*):ti,ab
#	19.	(radioth* or radiat* or irradiat* or radiochemo* or chemotherapy*)/mj
#	20.	((('bone marrow' transplant*4) or ('bone-marrow' NEAR transplant*))
#	21.	advanced or terminal*6 or progressi*6 or end-stage or endstage or 'end stage' (disease or illness)
#	22.	FATIGUE/exp
#	23.	(Fatigue or 'cancer-related fatigue' or 'cancer related fatigue'):ab,py
#	24.	'FATIGUE':de
#	25.	tired* or weary or weariness or exhaustion or exhausted or lackluster or (asteni* or asthenia*) and syndrome
#	26.	loss OR lost OR lack*2 AND (energy OR vigour OR vigor)
#	27.	apathy or apathetic or lassitude or lethargy* or (feeling (drained or sleepy or sluggish or weak*3))
#	28.	palliati* or hospice or 'end of life'

(Continued)

#	29.	#1 - #12/OR
#	30.	#13 - #21 /OR
#	31.	#22- #28 /OR
#	32.	#29 and #30 and #31
#	33.	(dexamphetamine or dextroamphetamine):ti,ab
#	34.	(methylphenidate or threomethylphenidate or d-threo-methylphenidate):ti,ab
#	35.	Donepezil:ti,ab
#	36.	amantadine:ti,ab
#	37.	Carnitine or l-carnitine:ti,ab
#	38.	Pemoline:ti,ab
#	39.	Modafinil:ti,ab
#	40.	Paroxetine:ti,ab
#	41.	(testosterone cypionate or androgen):ti,ab
#	42.	(acetylsalicylic acid or aspirin):ti,ab
#	43.	fluoxetine:ti,ab
#	44.	Citalopram:ti,ab
#	45.	(glucocorticoid or steroid or steroids or dexamethasone or dexamethasone or methylprednisolone or corticosteroid or corticosteroids):ti,ab
#	46.	Bupropion:ti,ab
#	47.	(‘anti depressant agents’ or ‘anti depressive agents’ or ‘Selective serotonin reuptake inhibitors’):ti,ab
#	48.	#33 - #47 / OR
#	49.	#32 and #48
#	50.	control*:ti,ab
#	51.	factorial:ti,ab
#	52.	(crossover or ‘cross over’ or ‘cross-over’):ti,ab
#	53.	placebo:ti,ab
#	54.	‘double blind’:ti,ab
#	55.	‘single blind’:ti,ab

(Continued)

#	56.	assign*:ti,ab
#	57.	allocate*:ti,ab
#	58.	'crossover procedure'
#	59.	'DOUBLE-BLIND PROCEDURE'
#	60.	'RANDOMIZED CONTROLLED TRIAL'
#	61.	'SINGLE BLIND PROCEDURE'
#	62.	'evaluation study'
#	63.	'prospective study'
#	64.	'comparative study'
#	65.	'animal experiment' OR 'nonhuman experiment'
#	66.	#50-#64/OR
#	67.	#49 AND #66 AND ([adult]/lim OR [aged]/lim) AND [humans]/lim AND [embase]/lim
#	68.	#67 NOT #65

Appendix 4. PsycINFO search strategy

Subject search combined with the recommended study design filter was run on 28 April 2014 (using OVID).

#	1.	exp Neoplasms/
#	2.	(cancer* or malignan*).mp.
#	3.	multiple sclerosis.af.
#	4.	amyotrophic lateral sclerosis.af.
#	5.	(sclerosis adj4 (amyotroph* or multiple)).mp.
#	6.	exp acquired immune deficiency syndrome/
#	7.	aids related complex.mp.
#	8.	exp Human immunodeficiency virus/
#	9.	(HIV or AIDS).mp.
#	10.	bone marrow transplantation.mp.
#	11.	radiotherapy.mp.

(Continued)

#	12.	(cancinoma* or tumour* or adenocarcinoma* or leukeni* or leukaemi* or lymphoma* or tumor*).mp
#	13.	(neutropeni\$ or neutropaeni\$).mp.
#	14.	(radioth* or radiat* or irradiat* or radiochemo* or chemotherap*).mp.
#	15.	Lung Disorders.mp.
#	16.	heart disorders.mp.
#	17.	congestive heart failure.mp.
#	18.	cor pulmonale.mp.
#	19.	cancer fatigue.mp.
#	20.	(cancer-related fatigue or cancer related fatigue).mp.
#	21.	exp FATIGUE/
#	22.	(tired\$ or weary or weariness or exhaustion or exhausted or lacklustred or ((astheni\$ or asthenia\$) and syndrome)).mp.
#	23.	((lack\$ or loss or lost) adj2 (energy or vigour or vigor)).mp.
#	24.	(apathy or apathetic or lassitude or letharg\$ or (feeling adj3 (drained or sleepy or sluggish or weak\$))).mp.
#	25.	((advanced or terminal* or progressi* or end-stage or endstage or "endstage") adj6 (disease or illness)).mp.
#	26.	palliati*.mp.
#	27.	(dexamphetamine or dextroamphetamine).mp.
#	28.	(methylphenidate or threo\$methylphenidate or d-threo-methylphenidate).mp.
#	29.	donepezil.mp.
#	30.	amantadine.mp.
#	31.	(Carnitine or L-carntine).mp.
#	32.	pemoline.mp.
#	33.	modafinil.mp.
#	34.	paroxetine.mp.
#	35.	(testosterone cypionate or androgen).mp.
#	36.	(acetylsalicylic acid or aspirin).mp.
#	37.	fluoxetine.mp

(Continued)

#	38.	citalopram.mp
#	39.	(glucocorticoid or steroid or steroids or dexamethason or dexamethasone or methylprednisolone or corticosteroid or corticosteroids).mp.
#	40.	bupropion.mp.
#	41.	(anti depressant agents or anti depressive agents or Selective serotonin reuptake inhibitors).mp.
#	42.	random*.ti,ab.
#	43.	factorial*.ti,ab.
#	44.	(crossover* or cross over* or cross-over*).ti,ab.
#	45.	placebo*.ti,ab.
#	46.	(doubl* adj blind*).ti,ab.
#	47.	(singl* adj blind*).ti,ab.
#	48.	randomized controlled trial.mp.
#	49.	assign*.ti,ab.
#	50.	allocat*.ti,ab.
#	51.	evaluation study*.ti,ab.
#	52.	prospective study*.ti,ab.
#	53.	comparative study*.ti,ab.
#	54.	(animal* or nonhuman* or animal experiment*).ti,ab.
#	55.	#1-#18/OR
#	56.	#19-26/OR
#	57.	#27-41/OR
#	58.	#42-53/OR
#	59.	#55 AND #56 AND #57 AND #58
#	60.	#59 NOT #54

FEEDBACK

Feedback submitted to Cochrane.org, June 2016

Summary

Comments submitted via the [Cochrane.org](https://www.cochrane.org) online contact form, and forwarded to the PaPaS Review Group by Nancy Owens, Cochrane Communications & External Affairs Department

Name: Vasiliy Vlassov

Which category applies to this message? Editorial

Subject: erroneous explanation

Message:

Here

<http://www.cochrane.org/news/featured-review-pharmacological-treatments-fatigue-associated-palliative-care>

"drugs that improve anaemia (lack of iron)," damn shaming wrong

Reply

Editorial response

The contributor highlights the definition of anaemia used by the authors in the Plain Language Summary and the summary version of the review on Cochrane.org. We attempted to contact Vasiliy Vlassov on more than one occasion in order to clarify the intended meaning of the comment, but we did not receive a response. Our interpretation of the comment is that the definition of anaemia in the PLS is misleading, and so this has now been amended to 'anaemia (a shortage of red cells or haemoglobin in the blood)'.

Contributors

Mrs Anna Erskine, Managing Editor, and Professor Christopher Eccleston, Co-ordinating Editor, at the Pain, Palliative and Supportive Care Review Group; Martin Mücke, Contact Person.

WHAT'S NEW

Date	Event	Description
15 April 2020	Review declared as stable	See Published notes .

HISTORY

Protocol first published: Issue 4, 2007

Review first published: Issue 11, 2010

Date	Event	Description
25 July 2017	Review declared as stable	See Published notes .
16 June 2016	Feedback has been incorporated	See Feedback .
27 May 2015	Amended	Contact details amended.
19 January 2015	New citation required and conclusions have changed	This review updates the original review, 'Pharmacological treatments for fatigue associated with palliative care' (Peuckmann-Post 2011), and also incorporates 'Drug therapy for the management of cancer-related fatigue' (Minton 2010). The updated search strategy identified 20 additional studies suitable for inclusion.
6 June 2014	New search has been performed	We used an updated search strategy and included 'Risk of bias' tables in this update.
5 November 2010	Amended	Minor amendment to title - text was italicised on publication and update of contact details.

Date	Event	Description
7 November 2008	Amended	Further RevMan 5 conversion changes.
13 August 2008	Amended	Converted to new review format.

CONTRIBUTIONS OF AUTHORS

MM, MC, LR: guarantors of the review, conceived the review, designed the search strategy, responsible for write up of the protocol and review.

MM and MC: responsible for the update of the review.

MM, MC: extracted data from papers and entered data into RevMan.

MM, MC, HC, LR: organised retrieval of papers, undertook searches and screened retrieved papers against inclusion criteria.

OM, PS: were involved in the original development of the search strategy (first published in Issue 3, 2007) and the appraisal of papers relating to pharmacological management of cancer-related fatigue, and approved the final manuscript.

LR, VP: were involved in the original development of the search strategy (first published in Issue 4, 2007) and the appraisal of papers relating to pharmacological treatments for fatigue associated with palliative care, and approved the final manuscript.

DECLARATIONS OF INTEREST

Martin Mücke has no relevant conflicts of interest to declare.

Mochamat has no relevant conflicts of interest to declare.

Henning Cuhls has no relevant conflicts of interest to declare.

Vera Peuckmann-Post has no relevant conflicts of interest to declare.

Ollie Minton has no relevant conflicts of interest to declare.

Patrick Stone has no relevant conflicts of interest to declare.

Lukas Radbruch has no relevant conflicts of interest to declare.

SOURCES OF SUPPORT

Internal sources

- Department of Palliative Medicine, University Hospital of Bonn, Bonn, Germany
- Department of General Medicine and Family Medicine, University Hospital of Bonn, Bonn, Germany

External sources

- Department of Anesthesiology and Intensive Care, University of Diponegoro/Kariadi Hospital Semarang, Indonesia

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

We modified the search strategy to match the inclusion criteria better and to prevent technical error. NNTB and NNTH calculations were not possible, as the information required to calculate these indicators was not included in the papers.

This updated review also incorporates another Cochrane review, 'Drug therapy for the management of cancer-related fatigue' ([Minton 2010](#)).

NOTES

2017

A restricted search in 2017 did not identify any potentially relevant studies likely to change the conclusions. Therefore, this review has now been stabilised following discussion with the authors and editors. If appropriate, we will update the review if new evidence likely to change the conclusions is published, or if standards change substantially which necessitate major revisions.

2020

A restricted search in August 2019 did not identify any potentially relevant studies likely to change the conclusions. Therefore, this review has now been stabilised following discussion with the authors and editors. The review will be re-assessed for updating in three years. If appropriate, we will update the review before this date if new evidence likely to change the conclusions is published, or if standards change substantially which necessitate major revisions.

INDEX TERMS

Medical Subject Headings (MeSH)

Amantadine [therapeutic use]; Benzhydryl Compounds [therapeutic use]; Carnitine [therapeutic use]; Central Nervous System Stimulants [therapeutic use]; Chronic Disease; Fatigue [*drug therapy] [etiology]; Kidney Failure, Chronic [complications]; Methylphenidate [therapeutic use]; Modafinil; Multiple Sclerosis [complications]; Neoplasms [complications]; *Palliative Care; Pemoline [therapeutic use]; Randomized Controlled Trials as Topic

MeSH check words

Adult; Humans