Cochrane
Library

Cochrane Database of Systematic Reviews

Anticholinergic agents for chronic asthma in adults (Review)

Westby MJ, Benson MK, Gibson PG

Westby MJ, Benson MK, Gibson PG.
Anticholinergic agents for chronic asthma in adults.
Cochrane Database of Systematic Reviews 2004, Issue 3. Art. No.: CD003269.
DOI: [10.1002/14651858.CD003269.pub2](https://doi.org/10.1002/14651858.CD003269.pub2).

www.cochranelibrary.com

TABLE OF CONTENTS

ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
BACKGROUND	3
OBJECTIVES	4
METHODS	4
RESULTS	6
Figure 1.	7
DISCUSSION	12
AUTHORS' CONCLUSIONS	13
ACKNOWLEDGEMENTS	13
REFERENCES	14
CHARACTERISTICS OF STUDIES	27
DATA AND ANALYSES	51
Analysis 1.1. Comparison 1 Longer term AC+BA vs BA, Outcome 1 Withdrawals due to treatment failure/side effects.	52
Analysis 1.2. Comparison 1 Longer term AC+BA vs BA, Outcome 2 No of patients with side effects.	52
Analysis 2.1. Comparison 2 Longer Term AC vs placebo, Outcome 1 Symptom score paired mean difference (Scale -1 to +1). ...	53
Analysis 2.2. Comparison 2 Longer Term AC vs placebo, Outcome 2 PEF paired analysis (scale -100 to +100 l/min).	54
Analysis 2.3. Comparison 2 Longer Term AC vs placebo, Outcome 3 Patient preference.	55
Analysis 2.4. Comparison 2 Longer Term AC vs placebo, Outcome 4 Number of patients dropping out.	55
Analysis 2.5. Comparison 2 Longer Term AC vs placebo, Outcome 5 Adverse effects - number of patients.	55
Analysis 3.1. Comparison 3 IB 40 vs IB 80, Outcome 1 Symptom score (scale -4 to +4).	56
Analysis 3.2. Comparison 3 IB 40 vs IB 80, Outcome 2 PEF (scale -100 to +100 l/min).	57
ADDITIONAL TABLES	57
WHAT'S NEW	67
HISTORY	67
CONTRIBUTIONS OF AUTHORS	68
DECLARATIONS OF INTEREST	68
SOURCES OF SUPPORT	68
INDEX TERMS	68

[Intervention Review]

Anticholinergic agents for chronic asthma in adults

Maggie J Westby¹, Malcolm K Benson², Peter G Gibson³

¹Division of Nursing, Midwifery & Social Work, School of Health Sciences, Faculty of Biology, Medicine & Health, University of Manchester, Manchester Academic Health Science Centre, Manchester, UK. ²Osler Chest Unit, Churchill Hospital, Oxford, UK. ³Department of Respiratory and Sleep Medicine, John Hunter Hospital, New Lambton, Australia

Contact: Maggie J Westby, Division of Nursing, Midwifery & Social Work, School of Health Sciences, Faculty of Biology, Medicine & Health, University of Manchester, Manchester Academic Health Science Centre, Jean McFarlane Building, Oxford Road, Manchester, M13 9PL, UK. maggie.westby@manchester.ac.uk.

Editorial group: Cochrane Airways Group.

Publication status and date: Edited (no change to conclusions), published in Issue 7, 2017.

Citation: Westby MJ, Benson MK, Gibson PG. Anticholinergic agents for chronic asthma in adults. *Cochrane Database of Systematic Reviews* 2004, Issue 3. Art. No.: CD003269. DOI: [10.1002/14651858.CD003269.pub2](https://doi.org/10.1002/14651858.CD003269.pub2).

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Anticholinergic agents such as ipratropium bromide are sometimes used in the treatment of chronic asthma. They effect bronchodilation and have also been used in combination with β_2 -agonists in the management of chronic asthma.

Objectives

To examine the effectiveness of anticholinergic agents versus placebo and in comparison with β_2 -agonists or as adjunctive therapy to β_2 -agonists.

Search methods

The Cochrane Airways Group asthma and wheeze database was searched with a pre-defined search strategy. Searches were current as of August 2008. Reference lists of articles were also examined.

Selection criteria

Randomised trials or quasi-randomised trials were considered for inclusion. Studies assessing an anticholinergic agent versus placebo or in combination/comparison with β_2 -agonists were included. In practice, all β_2 -agonists were short acting. Short-term (less than 24 hours duration) were not considered for this review.

Data collection and analysis

Two reviewers independently assessed abstracts for retrieval of full text articles. Papers were then assessed for suitability for inclusion in the review. Data from included studies were extracted by two reviewers and entered into the software package (RevMan 4.2). We contacted authors for missing data and some responded. Adverse effect data were analysed if reported in the included studies.

Main results

The studies analysed were in two groups: those comparing anticholinergics with placebo and those comparing the combination of anticholinergics with short acting β_2 -agonists versus short acting β_2 -agonists alone. The former group had 13 studies involving 205 participants included in this review, and the latter 9 studies involving 440 patients. Generally methodological quality was poorly reported, and there were some reservations with respect to the quality of the studies. Despite the limited number of studies that could be combined, anticholinergic agents in comparison with placebo resulted in more favourable symptom scores particularly in respect of daytime dyspnoea (WMD -0.09 (95%CI -0.14, -0.04, 3 studies, 59 patients). Daily peak flow measurements also showed a statistically significant improvement for the anticholinergic (e.g. morning PEF: WMD =14.38 litres/min (95%CI 7.69, 21.08; 3 studies, 59 patients). However the

clinical significance is small and in terms of peak flow measurements equates to approximately a 7% increase over placebo. The more clinically relevant comparison of a combination of anticholinergic plus short acting β_2 -agonist versus short acting β_2 -agonist alone gave no evidence in respect of symptom scores or peak flow rates of any significant differences between the two regimes. Again there are reservations with respect to the quality of the information from which these conclusions are drawn. An update search in August 2004 did not identify any new studies.

Authors' conclusions

Overall this review provides no justification for routinely introducing anticholinergics as part of add-on treatment for patients whose asthma is not well controlled on standard therapies. This does not exclude the possibility that there may be a sub-group of patients who derive some benefit and a trial of treatment in individual patients may still be justified. The role of long term anticholinergics such as tiotropium bromide has yet to be established in patients with asthma and any future trials might draw on the messages derived from this review.

PLAIN LANGUAGE SUMMARY

Anticholinergic agents for chronic asthma in adults

Anticholinergic agents such as Atrovent are sometimes used to treat people with asthma as a bronchodilator that opens up the airways in the lungs. This review found that although this treatment was better than placebo, the size of the effect was rather small. When the drug was used in combination with more widely used bronchodilators (beta-agonists such as fenoterol), it did not appear to add much benefit. However, there are concerns about the quality of the studies that have been analysed. It could be that there are some adults with chronic asthma who respond to treatment with anticholinergic drugs, but the review has not been able to identify their common characteristics.

BACKGROUND

Asthma is the commonest chronic disease affecting all age groups in developed countries (Harrison 1998), and has been recognised and treated in the East for centuries (Gross 1984). The anticholinergic agent, atropine was probably the first treatment used for asthma and was introduced in the West in the early 1800s (Gross 1984). Atropine acts as a bronchodilator, but has adverse effects, even at doses close to those used for bronchodilation (Gross 1988), and, as a result, fell from favour following the commercial development of ephedrine and adrenaline - beta-agonists - in the 1920s and, later, the methylxanthines. However, in the 1970s a quaternary ammonium synthetic analogue of atropine, ipratropium bromide, was developed, which gave few systemic side effects and revived interest in the anticholinergics (Gross 1988). A recent study in the USA (Taylor 1999) showed that anticholinergics were prescribed for asthma in 7 of 85 cases (8%) - age range 18-54 years, with a range of disease severities. Data from a UK general practice database on new use bronchodilators in severe asthma showed that, of 14657 patients, 20% received ipratropium bromide (Meier 1997).

Anticholinergic agents act as bronchodilators. The rationale for using them in asthma is based on the assumption that activation of the parasympathetic (cholinergic) nerve pathway is an important mechanism for producing airway obstruction. Anticholinergics work by competing with acetylcholine for receptor sites at the vagus nerve-nerve or nerve-muscle junctions. This prevents transmission of reflexes induced by asthma stimuli (Gross 1997, Beakes 1997).

Until the early 1980s, the treatment for asthma was regular bronchodilator agents (especially short-acting selective β_2 - agonists - the successors to adrenaline and ephedrine) and other agents were added only when symptoms reached an unacceptable level of frequency or severity (O'Connor 1998). Treatment was based around relieving the symptoms (airway constriction) and patients were often encouraged to tolerate continuing symptoms, treating them when necessary with repeated doses of β_2 - agonists. However, in the mid 1980s, opinions on asthma changed rather strikingly, because asthma was recognised as a chronic inflammatory disease (Barnes 1996). The therapeutic emphasis in chronic asthma in adults changed to embrace agents that suppress the underlying inflammation, notably inhaled corticosteroids (BTS 1993, Bousquet 2000).

Asthma guidelines have been available since the late 1980s and are updated regularly (Harrison 1998, Kemp 2000, Creer 1999). Initially they were arrived at by consensus, based on discussion and state-of-the-art literature reviews. More recent guidelines such as the BTS/SIGN guidelines (BTS/SIGN 2004) have adhered to systematic electronic searching and standardised guideline methodology. The grades of recommendation are derived from levels of evidence. According to current guidelines, the early introduction of anti-inflammatory treatment in the form of inhaled steroids is indicated for all but the mildest forms of asthma. Bronchodilators are adjuvant treatments used in one of two ways. Firstly, short-acting bronchodilators in the form of β_2 - agonists are used as-needed to relieve episodic symptoms. Secondly, in patients whose asthma is inadequately controlled despite regular inhaled steroids, the addition of long-acting β_2 - agonists has been shown to improve lung function and symptoms, and to decrease exacerbations (Ni

Chroinin 2005). A small proportion of patients have asthma which is not adequately controlled despite moderate doses of inhaled steroid and a long-acting β_2 - agonist. There are few clinical trials in this specific patient group and possible options include an increase in the dose of inhaled steroid or additional treatment, which might include leukotriene receptor antagonists, theophyllines or anticholinergics.

It is a widely held view that anticholinergics are less effective than β_2 - agonists in the symptomatic treatment of chronic asthma (Cazzola 1998), although there is considerable variation in treatment effect amongst patients. Attempts to identify subgroups that respond better to anticholinergics have not been very successful. In general, though, anticholinergics may be better in the following: older patients - the β_2 -agonist responsiveness apparently declines with age (Cazzola 1998, Connolly 1993); patients intolerant to β_2 -agonists (Boulet 1999); patients with nocturnal asthma (Beakes 1997); patients with chronic asthma and concurrent fixed airway obstruction (Cazzola 1998), patients with intrinsic asthma and those with longer duration of asthma (Partridge 1981). Patients with high serum immunoglobulin-E levels may be better treated with anticholinergic agents (Cazzola 1998), although earlier studies show that atopic patients respond less well to anticholinergics than those with non-atopic disease (Jolobe 1984).

The possible differences in responsiveness to anticholinergics between different patient groups illustrates the heterogeneous nature of asthma. This is further compounded when taking into account the overlap of asthma and COPD. At one end of the spectrum are patients whose airflow limitation shows marked spontaneous fluctuations and improves considerably with treatment (asthma). At the other end of the spectrum are patients whose disease fluctuates to a very limited extent and is "irreversible". Whilst an improvement of greater than 15% after inhaling a short-acting β_2 -agonist indicates a degree of reversibility, any such figure is inevitably arbitrary when there is a spectrum of airway disease ranging from "reversible" to "irreversible". Clinical features such as age, nature of symptoms, atopic status and smoking history are important factors in terms of diagnosis. This blurring of the boundaries becomes important when considering anticholinergics since it is generally regarded that they may have a small but proportionately greater effect than β_2 -agonists in patients with COPD.

The other factor that needs to be taken into consideration is the relative safety of β_2 -agonists and anticholinergics - the former can cause an increased heart rate and have also been associated with increased bronchial activity and increased mortality (Emilien 1998, Spitzer 1992). In comparison, the anticholinergics have relatively few side effects. Finally, anticholinergics are also used in combination with β_2 -agonists (Gross 1984). In severe asthma, the main treatment is based on inhaled steroids, but the benefit : harm ratio of inhaled steroids appears to be reduced once the total daily dose exceeds 1000 mg (O'Connor 1998), with adverse effects leading to increased risk of osteoporosis, skin thinning and adrenal problems (van Schayck 1995). As an alternative to high doses of inhaled steroid, moderate levels of inhaled steroid have been used in conjunction with oral steroids, leukotriene-receptor antagonists, steroid sparing alternatives, theophylline, anticholinergics or long-acting β_2 -agonists (O'Connor 1998). Oral steroids and some of the alternatives, including long-acting beta-agonists, have adverse

effects too (Barnes 1997, Hougardy 2000), which might indicate a place for the relatively safe anticholinergics.

OBJECTIVES

The objectives of this review, as defined in the original protocol ("Anticholinergic agents for chronic asthma in adults"), were to investigate the efficacy and safety of the various anticholinergic agents in chronic asthma in adults in two modes of treatment:

1. Anticholinergics for symptom relief (as-needed use), either alone or in combination with β_2 -agonists, in comparison with β_2 -agonists or placebo. This included both acute (short-term) bronchodilating effects and the longer-term effectiveness of bronchodilators for symptom relief.
2. Anticholinergics as regular, longer-term treatment, including their use as adjuncts to inhaled steroids, in comparison with β_2 -agonists (long- or short- acting) or placebo.

Protocol splitting

It was decided to split the original review in line with the two objectives because there was so much data, in an attempt to make the reviews more comprehensible. In practice there were no prn (as-needed) studies in the longer term, so that the two reviews focussed on (A) short-term bronchodilator effects (up to 24 hours) and (B) longer-term regular treatment (for periods from 2 days to 3 years). The latter focus is reported in this review and the former in the review, "Anticholinergic agents for chronic asthma in adults short term".

METHODS

Criteria for considering studies for this review

Types of studies

Randomised trials and quasi-randomised trials (e.g. allocation to treatment by alternation or date of birth) were analysed in the review. If a study did not report allocation details, but the allocation could have been randomised or quasi-randomised, further information was sought from the authors and the study was included if randomisation could be confirmed. Included studies reported the measurement of at least one of the outcome variables.

Types of participants

Adult patients with stable asthma: stability was defined as an absence of exacerbations, and stable medication prior to the study. Asthma therapy used during the study was also required to be stable.

The studies were required to report that the patients had asthma - preferably this was defined according to reliable criteria, such as those of the ATS, but less exactly, the patients had to display a reversibility in their lung function of at least 15% following administration of a short term β_2 -agonist (reversibility shown using anticholinergics was considered confounded and such studies were excluded).

Studies of patients with serious, non-respiratory diseases (e.g. cancer, heart disease) were excluded, as were those of patients with cystic fibrosis or irreversible airways disease. Studies involving a mixed group of patients having either asthma or COPD were

included if separate results were given (or could be obtained from trialists) for the patients with asthma.

Types of interventions

Studies were included if they used anticholinergic agents (including ipratropium bromide, oxitropium bromide, tiotropium bromide, atropine methonitrate and glycopyrrolate) used as a bronchodilator, in comparison with a β_2 -agonist or placebo. Delivery systems included nebulisers, metered dose inhalers and powder inhalers, with and without a spacer device.

Studies were included that compared:

1. Anticholinergics versus β_2 -agonist
2. Anticholinergics versus placebo
3. The combination of β_2 -agonist and anticholinergic versus β_2 -agonist alone
4. Different doses of anticholinergics
5. One anticholinergic versus another

Analysis was carried out only for comparisons 2 and 3 for the longer-term studies.

It was decided not to analyse trials that had atropine sulphate as the sole anticholinergic agent, because this has a different mode of action to the quaternary salts of atropine, has adverse effects and is therefore not routinely used as a bronchodilator in chronic asthma. Similarly, studies using non-selective β -agonists such as isoprenaline were included, but not analysed.

Types of outcome measures

All possible outcome measures were considered, but the main focus was as follows:

1. Daytime and night-time asthma symptom scores (primary outcome)
2. Bronchodilator use for symptom relief (i.e. 'rescue medication')
3. Daily peak flow (PEF; morning and evening)
4. Patient preference
5. Quality of life score
6. Asthma exacerbation rates/ hospital admissions - number of patients, not exacerbations
7. Adverse events and effects, e.g. tachycardia

Search methods for identification of studies

Electronic searches

Trials were identified using the Cochrane Airways Group Specialised Register of trials, which is derived from systematic searches of bibliographic databases including the Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE, EMBASE, CINAHL, AMED and PsycINFO, and handsearching of respiratory journals and meeting abstracts. All records in the Specialised Register coded as 'asthma' were searched using the following terms:

Anticholinergic* OR cholinergic* OR ipratropium OR Muscarinic* OR ipratropium OR N-isopropylatropine OR oxitropium OR tiotropium OR atropine OR glycopyrrolate OR Atrovent OR Sch1000 OR Oxivent

The most recent search was conducted in August 2008.

Other sources

Bibliographies from retrieved trials, meta-analyses and narrative reviews were checked to identify relevant cross-references. Attempts were made to obtain additional data from the authors and from the drug company, Boehringer Ingelheim (Bracknell UK); largely these were fruitless because many of the studies were about 20 years old and the data not available or we were unable to contact the authors.

Data collection and analysis

Step I. Trials that appeared potentially relevant were identified independently by two reviewers (MW and MB) from the search citation abstracts.

Step II. Using the full text of each study, two reviewers (MB and MW) independently selected trials for inclusion in the review. Agreement was measured using simple agreement and differences were resolved by discussion.

Step III. After a preliminary review of all studies to confirm the basic requirements, two reviewers independently assessed the methodological quality of the included trials with particular emphasis on the concealment of allocation, which was ranked using Cochrane criteria (grade A: adequate concealment; grade B: uncertain; grade C: clearly inadequate concealment). Any differences were resolved by discussion. The Jadad scale ([Jadad 1996](#)), described in the protocol, was not used for additional analysis because it and other scales are no longer thought to be the best approach to quality assessment ([Clarke 2003](#)).

Step IV. Two reviewers (MB and MW) independently extracted data from the included trials and one reviewer (MW) entered results into the Cochrane Collaboration software program (Review Manager, version 4.2). These data were checked by MB. A pilot for the data extraction form was carried out on a small sample of studies and was revised appropriately. In practice this led to the use of two forms, the first stage involving details of participants, the interventions, the type of outcomes and whether variability measures (e.g. standard errors, p-values) were reported. The second form contained results if available (e.g. mean and standard deviations for continuous outcomes). In some cases, information regarding outcomes was estimated from graphs; this was performed independently by the two reviewers.

Data extraction included the following items:

- Participants: age, gender, smoking status, type of asthma (intrinsic, extrinsic), allergic status (atopic, non-atopic), presence of co-existing COPD, severity of asthma (rated by MB based on baseline lung function, maintenance therapy and descriptions of the authors), history of asthma, symptom status at trial entry (symptomatic/asymptomatic)
- Intervention: anticholinergic agent, dose, schedule, spacer device, concurrent β_2 -agonist
- Control: β_2 -agonist, placebo
- Outcomes: self-rated daily symptom score/symptoms - daytime and night-time, pulmonary function measures (baseline and daily peak flow rates), bronchodilator use for symptom relief, hospital admissions/rate of acute exacerbations, quality-of-life instruments, adverse events and effects.

- Design: method of randomisation, allocation concealment, blinding, presence and type of run-in period, study design (parallel, cross-over), concurrent background therapy, duration of study and, for crossover studies, the duration of any washout period and/or the time between successive crossover arms
- Funding and multi-centre trials

STATISTICAL CONSIDERATIONS

- The results for particular outcomes for each trial were combined using RevMan (Version 4.2). An intention-to-treat analysis was completed where possible. Most of the studies in this review had a crossover design, that is, the patients were given each intervention in a random order. This means that there are within-patient correlations, which, ideally, should be treated using a paired analysis ([Elbourne 2002](#)). Where possible, this was carried out for continuous outcomes according to the guidance given by [Elbourne 2002](#). The standard error for the difference of means was calculated, either from individual patient data or from p-values, and the between-period correlation parameter, ρ , was calculated. For some studies the standard error was imputed using the lowest value of ρ from the other studies. For parallel studies the standard error was calculated as the square root of the sum of the variances for each arm. All similar parallel and crossover studies with sufficient data were pooled using the fixed effects generic inverse variance method in RevMan Version 4.2 to give a weighted mean difference and 95% confidence intervals.
- In the absence of paired data the trials were analysed by the conventional approach of treating the two arms of the crossover as if they were from a parallel trial with separate groups (i.e. setting $\rho=0$). If this was the case, parallel and crossover trials were analysed separately.
- For non-paired data and continuous variables, a fixed effects weighted mean difference (WMD) or standardised mean difference (SMD) and 95% confidence interval (CI) were calculated for each study. All similar studies were pooled using fixed effects WMD/SMD and 95% CIs.
- For dichotomous variables, a fixed effects odds ratio (OR) with 95% confidence intervals (95% CI) was calculated for individual studies. All similar studies were pooled using fixed effects OR and 95% CIs.
- For pooled effects, heterogeneity was tested using the Breslow-Day test; $p < 0.1$ was considered statistically significant. The statistical term I^2 (which is a measure of the inconsistency among results) was also used to monitor the variability among studies.
- Funnel plots were not constructed because of the paucity of included studies.

SUB-GROUP/SENSITIVITY ANALYSES:

If there had been sufficient studies sub-group analyses would have been carried out. *A-priori* defined subgroups were:

1. Concurrent therapy with inhaled corticosteroid or not
2. Asthma severity (mild, moderate or severe)
3. Concurrent COPD or not
4. Smoking status
5. Age (<40; 40-69; 70+)
6. Symptom status at trial entry (symptomatic/asymptomatic)

7. Asthma type (intrinsic, extrinsic)
8. Allergic status (atopic, non-atopic)
9. Duration of asthma
10. Dose, duration and delivery method of therapy
11. Type of anticholinergic agent

Sensitivity analyses had been proposed using the following domains:

1. Methodological quality
2. Random effects versus fixed effects modelling

However, there was insufficient data to examine these.

Rationale for the presentation of studies in the review

The original review had a very large number of included studies, separated into two modes of treatment (short-term bronchodilation and longer-term regular treatment). To manage the data we used a Microsoft Access database and prioritised the comparisons to be studied. This prioritisation took place before data extraction. For the longer-term studies, the clinical question of particular interest was whether a combination of anticholinergic with β_2 -agonist would give additional symptom relief compared to that obtained with β_2 -agonist alone, so this was the main focus. It was decided to set aside the comparison anticholinergic versus β_2 -agonist because it was of little practical interest, and to look at the comparison anticholinergic versus placebo only if no difference was found between the combination and the β_2 -agonist. This would clarify whether the anticholinergic had any effect (relative to placebo) alone. In reality, two types of comparison were analysed for the longer term studies:

1. Anticholinergic versus placebo
2. Anticholinergic plus β_2 -agonist versus β_2 -agonist alone.

It was felt appropriate to separate the studies according to duration and the following definitions were provided by MB prior to data extraction: short (2 days to 2 weeks), medium (more than 2 weeks to 2 months) and long (more than 2 months).

RESULTS

Description of studies

Results of the search

From electronic searches conducted for the original review, a total of 623 references were identified. From the abstracts of the study reports, 210 references were marked for further scrutiny, and the

remaining 413 were excluded because they clearly did not meet the inclusion criteria (for example, the patients were children or were treated for acute asthma). A further 103 possible studies were identified from other sources (e.g. references in included studies and reviews). This gave 302 possible references for which the full papers were obtained where possible. 26 reports were in non-English languages, of which 19 were translated. 239 studies were included in the original review. Of these, 45 studies had only longer-term data, 187 had only short-term data and 7 had both. The longer-term studies were all concerned with the regular addition of anticholinergics and there were no studies of longer-term as-needed use. For details of the studies that were excluded and the studies included in the short term review see the Table of Excluded Studies. Two longer-term studies were later excluded and deserve particular mention: [Tarlo 1982](#), which is a crossover comparison of anticholinergic and placebo over 90 days was excluded because the patients were instructed to reduce gradually their dose of bronchodilators - thus their concurrent medication was not stable, which we considered to be confounding. Secondly, [Brand 1992](#) (and its 7 accompanying papers) was a large parallel 3 year comparison of the combination of ipratropium bromide and terbutaline versus terbutaline alone. Unfortunately the study did not report separate results for asthma, although these were recorded, and contact with the authors has failed to produce any data. Four other studies did not state if the patients were randomised to treatments, and no further information was obtained from the authors. This left 39 reports of randomised or quasi randomised trials. Of these 39 longer-term study reports, 9 were additional reports of trials already reported, leaving 30 trials.

The 30 longer-term trials sometimes included more than two randomised interventions, giving rise to more than one comparison per trial; thus 23 trials had one comparison, 6 two comparisons and 1 trial had 3 comparisons. The number of trials reporting the different comparisons was:

1. Anticholinergic versus placebo in 13 trials (analysed in this review)
2. Anticholinergic plus β_2 -agonist versus β_2 -agonist alone in 9 trials (analysed in this review)
3. Anticholinergic versus β_2 -agonist compared in 12 trials (not analysed)
4. Anticholinergic dose comparison in 4 trials (not analysed)
5. Different anticholinergics were compared in 0 trials (not analysed).

Thus, 22 trials were included for the two types of comparison prioritised here. The flow of studies is shown in figure 1 ([Figure 1](#)).

Figure 1. Fig 1 Study flow.

Update searches to August 2008 identified two studies which we excluded as they were in children (Dutt 1990; Li 2003).

Included studies

Study size

In the placebo comparison the study size of the crossover studies ranged from 6 to 39 patients and all but two studies (Vaughan 1988, Wheeler 1987) had fewer than 20 patients with asthma. In the combination comparison there were two parallel trials each with over 100 patients and the remaining crossover trials had a study size ranging from 12 to 28, with only two studies having more than 20 participants (Rebuck 1983, Tammivaara 1993).

Design

In the placebo comparison all the trials had a crossover design; for the combination versus β_2 -agonist comparison there were 2 parallel trials and 7 with a crossover design. For the placebo comparison, 1 trial had a quasi-randomised design (Light 1977), but all other included trials reported randomisation.

Duration

In the placebo comparison the duration ranged from 4 days to 1 month per phase; for the combination versus β_2 -agonist comparison the parallel trials had durations ranging from 8-12 weeks, and the crossover trials' phases ranged from 2 weeks to 3 months.

Participants

In the anticholinergic-placebo comparison, one study (Taytard 1984) had patients with perennial asthma that became worse in spring and summer and improved over the course of the study. The authors found an order effect and for this reason only first period results were used for this study, and the individual patient data analysed as a parallel trial.

One study did not give any details about concurrent medication (Ruffin 1990), one was unclear but had no oral steroids (Coe 1986), two had no concurrent medication (Ahonen 1978, Taytard 1984), one had sodium cromoglycate only (Bellia 1988), three had inhaled steroids only (Pavia 1989, Tormey 1995, Wheeler 1987), one had unspecified steroids (Kreisman 1981) and the others had oral and inhaled steroids (Light 1977, Salorinne 1988, Taylor 1986, Vaughan 1988).

Four studies were assessed to have patients with mild asthma (Ahonen 1978, Bellia 1988, Ruffin 1990, Taytard 1984), 5 with moderate asthma (Kreisman 1981, Pavia 1989, Salorinne 1988, Tormey 1995, Vaughan 1988) and 4 with heterogeneous patient populations (Coe 1986, Light 1977, Taylor 1986, Wheeler 1987). No study claimed to have patients with concurrent COPD.

In the combination comparison, one study did not give any details about concurrent medication (Waite 1987), one had aminophylline but did not say if steroids were used (Macaluso 1986), one had no steroids (Tammivaara 1993), one had only inhaled steroids and sodium cromoglycate (Haahtela 1991), two had unspecified steroids (Mazzei 1985, Philip-Joet 1990) and the other three had oral and inhaled steroids (Pierce 1982, Rebuck 1983, Wolstenholme 1989). Most of the studies also used methylxanthines as concurrent medication. Four studies were assessed to have patients with moderate asthma (Haahtela 1991, Pierce 1982, Tammivaara

1993, Wolstenholme 1989), two gave no information on severity (Macaluso 1986, Waite 1987) and three had heterogeneous patient populations (Mazzei 1985, Rebuck 1983, Philip-Joet 1990). One study (Pierce 1982) had patients with concurrent bronchitis.

Interventions

For the anticholinergic-placebo comparisons, there were three main variables with respect to the anticholinergic. The anticholinergic drugs included ipratropium bromide (7 studies), oxitropium bromide (8), nebulised atropine methonitrate (1) and inhaled atropine sulphate (Light 1977). The latter study was not analysed. The dose of ipratropium bromide ranged from 40 μ g to 200 μ g delivered either by metered dose inhaler or powder inhaler. The frequency of administration ranged from twice daily to five times daily. For further details see table of included studies.

All of the studies comparing a combination of β_2 -agonist plus anticholinergic versus β_2 -agonist alone used ipratropium bromide as the anticholinergic agent at either 40 or 80 μ g, three or four times per day. In all studies the β_2 -agonist was short acting. In eight studies fenoterol was combined with ipratropium as Duovent (40 μ g ipratropium, 100 μ g fenoterol), Berodual (20 μ g ipratropium, 50 μ g fenoterol) or using two separate inhalers. In the β_2 comparison arm, three studies used fenoterol and the remaining studies used salbutamol. Three further studies used terbutaline as the β_2 -agonist, both in combination and as the comparator. However one of these studies (Tammivaara 1993) used oral controlled-release tablets and this study was analysed separately because of the different pharmacokinetics of the oral formulation. For further details see table of included studies.

Outcome measures

Outcome measures chosen were those generally accepted to reflect the primary outcome of asthma control. Daytime and night time asthma symptom scores were derived from daily diaries. Objective measurements of lung function were daily peak flow measurements, recorded morning and evening, before and after use of inhaler. Additional outcomes were treatment related withdrawal from the study, patient preference, bronchodilator use as rescue medication (number of patients requiring this), number of patients with exacerbations and adverse effects. Clinical and physiological assessments at the beginning and end of each trial period were regarded as being less valuable, given the recognised spontaneous fluctuations which can occur in asthma.

Risk of bias in included studies

The methodological quality was poorly reported in all 22 studies, but the authors provided further details for five studies (Philip-Joet 1990, Pierce 1982, Vaughan 1988, Wheeler 1987, Wolstenholme 1989). All but one of these (Philip-Joet 1990) still did not give an adequate description of allocation concealment, with only partial details provided (for example using sealed envelopes). One additional study (Light 1977) reported an inadequate method of allocation concealment. Generally the reporting of blinding was vague, with all but two being described as double blind or blind (1 study). The Philip-Joet study stated that it did not have a double blind design and the comparison we used for the Salorinne study (inhaled Ipratropium bromide versus powder placebo) could not have been blinded. The Macaluso study, despite stating it was double blinded, was not blinded because the combination arm used three inhalations and the β_2 -agonist four. The Pavia study

reported that some patients could distinguish by taste oxitropium bromide from placebo, and it is likely that the blinding was broken in some of the other placebo studies too. This may also be true of the combination comparison, because ipratropium bromide and salbutamol, for instance, taste quite different. Nine of the 22 studies had intention to treat analyses carried out and for two other studies it was unclear what was done. Six studies (Haahtela, Philip-Joet, Tammivaara, Wolstenholme, Ruffin and Wheeler) had at least 20% of the patients' results missing for some outcomes, and nearly 40% of the patients were not analysed for the 8-week PEF outcome in the Philip-Joet study.

Effects of interventions

Duration

Although the original intention was to separate the study durations as described in the methods section, in practice there were too few studies to make this feasible, so all durations were considered together.

A) Anticholinergic versus placebo

There were 13 studies comparing anticholinergic with placebo. Of these, one was not included in our analysis because it used atropine sulphate (Light 1977).

Seven studies were also not included in the initial analysis: 6 studies did not record means for the continuous outcomes and 1 other study did not report standard deviations or p-values. The authors of these studies were contacted and Boehringer Ingelheim provided individual patient data for Wheeler 1987; some other authors offered data, but to date these had not arrived. Thus, there were only 4 studies that provided sufficient data for analysis of the continuous outcomes, although 7 studies gave data for the dichotomous outcomes.

Ahonen 1978 compared two doses of anticholinergic and, in order to clarify the main analysis, we decided to choose one dose of ipratropium bromide, based on the head-to-head comparison. It was found that for both symptom scores and peak flow measurements the summary statistics were in favour of the lower dose (40µg), although none was statistically significant (see additional Table 1 and forest plots). Therefore for this study the lower dose was used in comparison with placebo in the main analyses.

The results for various outcomes for the comparison of anticholinergic with placebo are summarised in additional tables 2-5. Where there was more than one similar study, the results were combined and the summary statistics reported in the additional table and shown in a forest plot. Otherwise the results in the table are for single studies.

For some outcomes, the percentage increase or decrease over placebo was calculated. This is an approximation used to give an estimate of the magnitude of the effect in clinical terms. It was done using as denominator the simple average, over the individual studies, of the placebo means; the numerator was the weighted mean difference from the meta-analysis (or single study). It should be noted that the studies involved had a range of asthma severities, so that patients with more severe disease might derive greater benefit than represented by the mean percentage change, but there

is insufficient information to look at subgroups, either for peak flows or symptom scores.

1. Symptom scores (Table 2)

The data were analysed by calculating the difference in means and the paired difference standard error and pooling the data, where appropriate, using the generic inverse variance method. The Wheeler study gave individual patient data (IPD) and the Ahonen study reported p-values for the comparison. The scales for the studies were different (Ahonen 1978: 1-4; Wheeler 1987; Salorinne 1988: 0-3 and Taytard 1984 0-13), but the first two scales were linearly equivalent and so could be combined in the analysis. Taytard 1984 was analysed separately because the trialists reported a combined 'clinical score', which was a composite of day-time and night-time symptom scores and drug consumption for symptomatic relief. For this study, only first period results were used (because the authors reported an order effect); the study reported individual patient data and because of large baseline differences, the change scores were used.

The four studies had treatment periods ranging from 4 days to 4 weeks and it was decided to combine these different durations, by calculating the mean symptom score per day.

It should be noted that Wheeler 1987 (which provided individual patient data) had a number of patients who failed to report their scores every day, so we calculated symptom scores based only on the patients who completed their diary cards for more than 20 days (28/30 patients). The study reported symptom scores over the whole period for each patient, so we calculated their daily symptom score (for the number of days they reported), and then took the mean for all patients included. Ahonen 1978 reported placebo values averaged over two placebo arms.

The night-time symptom scores could not be combined because the Ahonen study measured only dyspnoea and the Wheeler study gave an overall measure of cough, breathlessness and wheeze; Salorinne 1988 gave no standard deviations, although it measured cough and dyspnoea separately. Thus there was only one symptom score outcome for which data could be pooled - daytime dyspnoea. In order to use the generic inverse variance method, we calculated the interaction parameter, ρ , for the Wheeler study ($\rho=0.817$) and used this in the Ahonen study to impute the standard error. Salorinne 1988 gave no standard deviations so for each arm we used the mean of the standard deviations from Ahonen 1978 and Wheeler 1987 and then calculated the standard error for the difference using the Wheeler value of ρ . Although not ideal, we feel that this gives the best representation of the data. This approximation for the Salorinne 1988 was examined by a sensitivity analysis and it was found that the addition of the Salorinne 1988 made little difference to the summary statistics.

The results are summarised in additional Table 2; Table 3; Table 4 Daytime dyspnoea: the summary statistics were: WMD -0.09 (95%CI -0.14, -0.04, 3 studies, 59 patients), i.e. the anticholinergic agent gave more favourable symptom scores than the placebo, and the result was statistically significant. There was little heterogeneity ($I^2=0\%$, $p=0.45$). This corresponds to a 15% decrease in symptom score over placebo. The other symptom scores could not be pooled and only the Taytard study gave a statistically significant result on its own - which was in favour of the anticholinergic agent. Otherwise the results of single studies indicated that, relative to placebo, the anticholinergic showed some improvement in

daytime cough and night-time dyspnoea, but some worsening in night-time cough.

In addition, [Kreisman 1981](#) reported that patients coughed more on the ipratropium regimes. [Tormey 1995](#) reported that there were no significant differences in symptom scores (unspecified) between placebo and oxitropium bromide over 28 days; and [Vaughan 1988](#) reported that 'total symptom scores' (cough and wheeze) were no different when using nebulised atropine methonitrate compared with placebo over 2 weeks.

2. PEF measurements (Table 3)

Four studies reported daily peak flow measurements, but one of these gave no p-values or standard deviations; the Ahonen and Salorinne studies reported some p-values, and IPD were given for the [Wheeler 1987](#). p values for this study (although not used) were respectively 0.919 and 0.911 for the am and pm peak flows. The outcomes were daily peak flow rates measured before and after the bronchodilator dose.

Summary statistics were calculated using the generic inverse variance method. The Ahonen study gave an upper bound for the p-value for the ipratropium-placebo comparison, which allowed calculation of the standard error for the mean of the difference (AC - placebo). The Salorinne study gave an upper bound for the p-value for the evening PEF, but for the morning value the standard deviation for the Wheeler study was taken and the standard error calculated from this ([Ahonen 1978](#) was not used for this purpose because of a discrepancy between the graph and table values).

The following results were found:

a) Morning PEF

The PEF daily average weighted mean difference (WMD) was: WMD = 14.38 litres/min (95%CI 7.69, 21.08; 3 studies, 59 patients) i.e., statistically significantly higher for the anticholinergic; this showed some heterogeneity ($p = 0.22$, $I^2 = 34\%$) and corresponds to an increase of about 7% over placebo. We tested the assumptions for [Salorinne 1988](#) using a sensitivity analysis, and found that the heterogeneity was removed ($p = 0.65$, $I^2 = 0\%$), but there was little effect on the WMD (16.26 L/min; 95%CI 9.21, 23.31).

Post-bronchodilator morning PEF measurements were available for 2 studies, and were taken 2 and 3h after the dose. The WMD was 34.91 litres/minute (95%CI 20.13, 49.70, 2 studies, 31 patients), a statistically significant difference, corresponding to an increase of 9% over placebo.

b) Evening PEF

The evening (about 10 pm) PEF daily average WMD was: 23.48 L/m (95%CI 12.32, 34.65, 3 studies, 60 patients) i.e., statistically significantly larger peak flow rate for the anticholinergic, but there was heterogeneity ($p = 0.01$, $I^2 = 78\%$). The WMD represents an 7% increase over placebo.

Post-bronchodilator evening PEF measurements were available only for [Salorinne 1988](#): WMD 23.94 L/m (95%CI 5.20, 42.68, 1 study, 18 patients), giving a statistically significant increase of 7%.

[Vaughan 1988](#) used nebulised atropine methonitrate over 2 weeks, and reported that the peak flow measurements were not different between placebo and Atropine methonitrate either in the morning or at 5pm.

3. Use of rescue medication (Table 4)

Three studies reported that they recorded the use of rescue medication, but relative numbers of patients were given only in the [Wheeler 1987](#). [Taytard 1984](#) (3 weeks) reported that the 'mean consumption of salbutamol was essentially identical for placebo and oxitropium bromide' and [Salorinne 1988](#) (1 week) reported that 'the number of additional treatments did not differ between the various treatments'. For the [Wheeler 1987](#) (4 weeks) 23 of 30 patients taking oxitropium and 25/30 taking placebo required rescue medication during the day (OR 0.66 ; 95%CI 0.18, 2.36 - not statistically significant) and the corresponding night time ratios were 15/30 and 19/30 (OR 0.58 ; 95%CI 0.21, 1.62).

4. Patient preference (Table 4)

The number of patients preferring anticholinergic was compared with the number preferring placebo in two studies ([Salorinne 1988](#) and [Wheeler 1987](#)) - in the latter case, some patients did not have a preference. In the former study patients chose between 40 µg, 200 µg (inhaled) and 200 µg powder ipratropium bromide as well as placebo (which was a powder); the odds ratio for patient preference was: OR 1.64 (95%CI 0.68, 3.94) i.e. non-significantly in favour of the anticholinergic, with little heterogeneity.

5. Drug-related withdrawal (Table 4)

Two studies gave data on the number of asthma patients withdrawing while taking one of the study drugs. The pooled odds ratio was 0.70 (95%CI 0.13, 3.72) - non significantly in favour of anticholinergic, with some heterogeneity.

6. Adverse effects (Table 5)

Six studies gave some information on adverse effects, but of these [Kreisman 1981](#) and [Bellia 1988](#) only gave a descriptive account, and [Ahonen 1978](#) and [Salorinne 1988](#) reported the number of effects or symptom scores, rather than the number of patients with those effects. [Kreisman 1981](#) stated that the side effects were equally reported, and [Bellia 1988](#) reported that no significant side effects were recorded for oxitropium bromide. For the studies that gave numerical data, individual side effects (e.g. mouth dryness) gave similar results for anticholinergic and placebo, with no statistically significant differences observed at the lower doses used, and for overall side effects, the pooled odds ratio was OR 1.87 (95%CI 0.51, 6.81) - not statistically significant.

B) Combination of anticholinergic and β_2 -agonist versus β_2 -agonist alone

There were 9 trials comparing the combination of anticholinergic and β_2 -agonist with β_2 -agonist alone. One of the studies ([Tammivaara 1993](#)) used oral CR terbutaline as the β_2 -agonist and we analysed this study separately because of the different pharmacokinetics. One study could not be analysed because it did not give any results and 2 studies did not report standard deviations or P-values. The authors of these studies were contacted and some offered data, but to date these had not arrived. Additional information, was however provided by [Wolstenholme 1989](#) and [Philip-Joet 1990](#). This left only six studies that provided sufficient data for analysis of the continuous outcomes, and six studies gave data for the dichotomous outcomes. For all studies the final values of the outcome measures were used, rather than adopting change scores.

In [Hahtela 1991](#), the peak flows and symptom scores were reported as a daily average for weeks 1-4 and, separately, 9-12 weeks; we used the mean of 1-4 and 9-12 weeks as a measure of the

response. Philip-Joet 1990 reported the weekly mean peak flows and standard deviations for weeks 1 to 8. However, many patients dropped out (48/196 overall) and the PEF results showed such a large number of missing patients (e.g., at 8 weeks, 37 of 101 from the combination arm and 40 of 95 from the β_2 -agonist arm) that we decided to include only the results for the first week as being the least confounded (still 11/101 and 12/95 missing).

Unlike the anticholinergic-placebo comparison, these studies differed from one another in various important ways, making pooling questionable. Firstly, there were differences in study duration, but we decided it was reasonable to combine all durations.

Secondly, there was a mixture of study designs, 2 parallel and 7 crossover trials. It was not possible to analyse the crossover studies using the generic inverse variance method because none reported IPD or P-values, and consequently we had to analyse the crossover studies as if they were parallel studies, which meant that the crossover and (true) parallel studies could not be combined in an analysis.

Thirdly, the β_2 -agonist in the two arms differed in both composition and dose. We decided to treat all the β_2 -agonists as having the same equivalence and recorded the difference in doses in a single variable, the ratio of β_2 -agonist in the combination arm to that in the single arm. Two ratios were analysed 1:1 and 1:2 - the latter can be considered as the *substitution* of anticholinergic for some of the β_2 -agonist in the single arm and the former as the *addition* of anticholinergic to the same level of β_2 -agonist. Thus the breakdown of studies is:

- Substitution of anticholinergic for β_2 -agonist (ratio 1:2 combination β_2 -agonist:single β_2 -agonist): Haahtela 1991, two comparisons (Berodual versus 200 μ g salbutamol and Duovent versus 400 μ g salbutamol) and Philip-Joet 1990
- Addition of anticholinergic to β_2 -agonist (ratio 1:1): Haahtela 1991 (Duovent versus 200 μ g salbutamol), all other studies

The differing types of design and interventions meant that, in general, the study data could not be pooled, and the results are shown in additional tables 6-9. Where there was more than one similar study, the results were combined and the summary statistics reported in the additional table and shown in a forest plot. Otherwise the results in the table are for single studies.

1. Symptom scores (Table 6)

Daily symptom scores were measured in six studies and all used a scale of 0-3. Mazzei 1985 only reported symptom scores at a clinic visit, not daily (thus not meeting our criteria). No results were reported for 2 studies and no standard deviations were given for one. Macaluso 1986 reported graphically a distribution of symptom scores, but the exact nature of the representation was unclear; despite these uncertainties, we calculated a mean symptom score and standard deviation from these data. Of the remaining studies, one reported the weekly symptom scores and two reported daily scores: the former values were adjusted to daily scores.

The only possible studies that were sufficiently similar to be combined were Macaluso 1986 and Wolstenholme 1989, but in view of the uncertainties about the Macaluso 1986 symptom score data, we decided to analyse these as single studies.

Generally there is no evidence, from the single studies analysed, for a significant difference in any symptom score between the combination and the single arm, regardless of the trial design, time duration or ratio of β_2 -agonist.

The authors of Haahtela 1991 also stated that, for all symptom score measurements, there were no significant differences among any of the 4 regimens, even when the subgroup of more severe patients was analysed. The authors of Macaluso 1986 stated that there was a statistically significant difference in favour of Duovent for daytime dyspnoea, but our analysis of their available data did not confirm this. The Pierce study stated that 'symptom score cards (measuring daytime and nocturnal dyspnoea, wheeze and cough) failed to reveal any significant differences between the combination and either drug alone'. Waite 1987 stated that 'symptom scores recorded in the morning and evening for wheeze, cough and restrictions in activity showed no significant differences between the two treatments'.

2. Lung function measurements (Table 7)

Daily peak flow measurements were measured in 5 studies. An additional study, Pierce 1982 measured daily mean flow using an airflowmeter so this study was analysed separately. Rebuck 1983 reported no data (but stated that there was a similar improvement for each arm) and Haahtela 1991 gave no standard deviations for the daily peak flow measurement.

The different characteristics of the studies (parallel/crossover, PEF/flow and oral/inhaled beta-agonist) meant that no studies were sufficiently similar to be combined. However, the general trend is that there is very little difference between interventions with no single study achieving statistical significance. Haahtela 1991 also reported that 'analysis did not reveal any significant differences, even when the subgroup of more severe patients was analysed'. Post-bronchodilator measurements were made for three studies and were taken after 20m, 30m and 1h. There was little difference between the interventions.

3. Patient preference (Table 8)

Only two crossover studies reported the preference of the patients (Tammivaara 1993; Wolstenholme 1989), but the former used oral beta-agonist. In Tammivaara 1993 the patients preferred the combination, but in Wolstenholme 1989 the reverse was the case; neither study showed statistical significance. Rebuck 1983 reported no significant difference in patient preference. Macaluso 1986 reported that the 'patient's opinion was more favourable with Duovent than salbutamol', but this may have been because the patients received respectively 3 and 4 administrations of the test drug.

4. Drug-related withdrawal (Table 8)

Four studies reported the number of patients withdrawing while taking one of the study drugs, but only 2 studies could be combined. The pooled odds ratio for the parallel study substitutional-anticholinergic was: 1.40 (95%CI 0.69, 2.84, 2, studies, using the Berodual-salbutamol comparison in Haahtela 1991, 161 patients), i.e. in favour of the β_2 -agonist, but not statistically significant and with some heterogeneity ($p=0.22$, $I^2=32\%$).

Overall, there were fewer withdrawals from the β_2 -agonist arm, regardless of the time duration, ratio of β_2 -agonist levels and trial design, but no result was statistically significant. The exception was the oral beta-agonist crossover study, Tammivaara 1993, which

showed fewer withdrawals for the combination (not statistically significant).

5. Number of patients with exacerbations (Table 8)

Two studies reported the number of patients with exacerbations (or worsening), [Philip-Joet 1990](#) and [Tammivaara 1993](#), the former shows fewer patients with exacerbations for the β_2 -agonist arm and the latter shows the reverse, neither is statistically significant, and the results have not been combined because of the use of the oral β_2 -agonist in [Tammivaara 1993](#).

6. Adverse effects (Table 9)

Six studies gave some information on adverse effects, but of these [Hahtela 1991](#) and [Mazzei 1985](#) reported the number of effects or symptom scores, rather than the number of patients with those effects and [Philip-Joet 1990](#) had too much missing data to be included without confounding. Two studies gave a descriptive account only: [Wolstenholme 1989](#) stated that there was no difference in side effects reported for the two periods and [Macaluso 1986](#) found that side effects (tremor, palpitations and tachycardia) occurred in negligible percentages for both treatments. The remaining studies were not pooled because they had different ratios of β_2 -agonist and different designs (see additional table 9). Generally, there is little evidence for a difference in adverse effects between the combination and single arm β_2 -agonist, and no result is statistically significant.

DISCUSSION

As a chronic disease, with no known cure, the accepted goals of management in chronic asthma are to minimise the adverse impact of the disease on the patient's physical and mental well-being, to prevent long term damage with fixed airflow obstruction and to reduce mortality. In the majority of patients treatment is therefore directed at improving physiological end-points and patient-perceived physical and mental health. Although anticholinergics are frequently used in patients with more severe forms of asthma, the evidence justifying its use remains limited. Both short-term and long-term anticholinergics are of some benefit in patients with COPD, their role in asthma is less clearly defined. The recent BTS guidelines include anticholinergics as one of the options for short term bronchodilator relief at Step 1, and in the management of acute severe asthma ([BTS/SIGN 2004](#)). They receive no mention as regular add-on treatment for use in combination with inhaled steroids and long-acting β_2 -agonists.

The acute bronchodilator effect of an anticholinergic is usually demonstrated by making physiological measurements before and after inhaling the drug. The most commonly used measurements include the FEV1 and vital capacity, although other end-points such as the ability to modify bronchial challenge tests have also been widely studied. It is on this basis that anticholinergics are sometimes used to relieve symptoms although the majority of acute studies do not include symptom scores as part of their evaluation.

This review was specifically directed at examining the role of anticholinergics used on a regular basis, especially when used as an adjunct to a β_2 -agonist. The comparison between a regular anticholinergic and placebo is of limited clinical relevance but was undertaken to determine whether there was a detectable subjective or objective improvement. Despite the limited number of studies, anticholinergic agents resulted

in more favourable symptom scores particularly in respect of daytime dyspnoea. Daily peak flow measurements also showed a statistically significant improvement for the anticholinergic. However, the clinical significance is small and in terms of peak flow measurements equates to approximately a 7% increase over placebo. Secondary outcome measures such as use of rescue medication, patient preference and drug-related withdrawals showed no significant differences although these conclusions were based on very limited objective data. The same reservations applied to adverse effects where again there was no evidence for a systematic difference between anticholinergic and placebo.

The second part of the review is more clinically relevant and addresses the question as to whether a combination of anticholinergic plus β_2 -agonist confers significant advantage when compared with β_2 -agonist alone. There was no evidence in respect of primary outcome measures, symptom scores or peak flow rates of any significant differences between the two regimes. Similarly, there were no significant benefits shown in respect of secondary outcome measures. There are however major reservations with respect to the quality of the information from which these conclusions are drawn. The total number of studies suitable for analysis was small and involved limited numbers of patients. A number of studies were excluded from analysis because of methodological flaws, incomplete data or because they included patients with COPD. Even for those studies included, differences in methodology meant that they could not easily be combined. The main differences which might affect analysis have already been identified but merit re-iteration.

Patient characteristics

All the included studies define the patients as having asthma or reversible airflow obstruction. A 15% improvement of FEV1 after an inhaled β_2 -agonist was the most frequently used physiological test although some studies included spontaneous variability. Criteria as defined by the American Thoracic Society were used in a significant number of studies. Patient characteristics varied considerably both between and within studies. There was a wide range of patient age, variable numbers of atopic patients and variable numbers of smokers with symptoms of chronic bronchitis. Despite this heterogeneity, it seems likely that most studies focused on patients at the more reversible end of the spectrum of airway disease.

Severity of the disease

Again there was considerable heterogeneity both between and within studies with patients ranging from having mild to severe asthma. Defining criteria were often not stated. Factors taken into consideration in trying to assess the severity of the disease included patient symptoms, baseline lung function and maintenance treatment. In addition, studies were only included if the asthma was regarded as being "stable". It is theoretically possible that patients with more severe disease might derive greater symptomatic benefit from small improvements in lung function. There were insufficient data to provide sub-group analysis.

Maintenance treatment

Once again there were significant variations both between and within studies. These ranged from low maintenance treatment to patients who were on regular treatment with theophylline, inhaled or oral steroids. Additional β_2 -agonists used as rescue treatment

were included in a number of studies and used as a secondary outcome measure.

Methodological differences: different drug combinations

There were significant differences between studies with respect to the different drug regimes studied. These included a variety of anticholinergics using different doses. The β_2 -agonist used also varied and in some studies the comparator β_2 -agonist was different from that used in the combination.

Methodological differences: outcome measures

Although diary cards were used in a number of studies, data was often incomplete and the scoring system varied. Studies using regular peak flow measurements were similar although the timing of the measurements with respect to the preceding dose may have influenced the outcome measurement.

The failure to demonstrate any subjective or objective improvement by the addition of anticholinergic to a regular β_2 -agonist may have various explanations which merit further consideration. As indicated, it may simply be that the studies are insufficiently powered to detect what may be small but significant differences. Patient selection is also likely to be critically important. Patients with mild disease will have little potential for further improvement over and above that achieved by inhaled steroids and β_2 -agonists. Conversely, patients with more intractable disease may well have airway obstruction with only limited reversibility. There were insufficient data to permit subgroup analyses, looking specifically at defined patient groups.

None of the studies specifically identified those patients whose asthma was inadequately controlled despite being on regular β_2 -agonists and inhaled steroids. In part this reflects the fact that the studies analysed have been undertaken over the past 20 years and guidelines have evolved during this period.

Overall this review provides no justification for routinely introducing anticholinergics as part of add-on treatment for patients whose asthma is not well controlled on standard therapies. This does not exclude the possibility that there may be a sub-group of patients who derive some benefit and a trial of treatment in individual patients may still be justified. The review does highlight the need for studies of sufficient power using standardised

criteria and methodology in order to answer specific questions. It is unlikely that such studies will be undertaken using the currently available short term anticholinergics. The role of long term anticholinergics such as tiotropium bromide has yet to be established in patients with asthma and any future trials might draw on the messages derived from this review.

AUTHORS' CONCLUSIONS

Implications for practice

Inhaled anticholinergics have a bronchodilator effect in patients with asthma. In clinical and physiological terms this effect is small. The addition of anticholinergics to regular inhaled bronchodilators has not been shown to confer significant benefits, so the current evidence does not justify their routine use. The study does not exclude the fact that there may be sub-groups of patients who derive some symptomatic benefit.

Implications for research

Future studies need to be sufficiently powered to answer specific questions, which need to be clearly formulated. The role of long term anticholinergics in the management of asthma is a potential case in point. Patient characteristics need to be clearly defined and particular focus directed at those patients with more severe disease that is inadequately controlled on a combination of inhaled steroids and long acting β_2 -agonists. In addition, outcome measures need to be standardised and to include patient-derived symptom scores and quality of life data in addition to physiological measurements.

ACKNOWLEDGEMENTS

Margaret Westby acknowledges the award of a Daphne Jackson research Fellowship and financial support from the Leverhulme Trust.

We would also like to thank the Airways Group for support with literature searches and retrieval of studies (namely Karen Blackhall, Liz Arnold, Sarah Tracy and Bettina Reuben) and the people who translated articles from non-English languages: Jane Dennis, Toby Lasserson, Johannes van der Wouden, Derek Scoins and Makiko Meguro.

REFERENCES

References to studies included in this review

Ahonen 1978 {published data only}

- * Ahonen A, Alanko K, Mattson K. Bronchodilator action of two different doses of ipratropium bromide (SCH 1000) compared with fenoterol and placebo. *Current therapeutic research, clinical and experimental*. 1978;**24**:65.

Ashutosh 1980 {published data only}

- * Ashutosh K, Mead G, Dickey JC Jr, Berman P, Kuppinger M. Dependency of expiratory flow and bronchodilator response in asthma. *Chest* 1980;**77**:68.

Bellia 1988 {published data only}

- * Bellia V, Ferrara G, Cibella F, Cuttitta G, Visconti A, Insalaco G, et al. Comparison of the effect of oxitropium bromide and of slow-release theophylline on nocturnal asthma. *Postgraduate Medical Journal* 1988;**64**:583.

Chapman 1983 {published data only}

- * Chapman KR, Rebuck AS. Anticholinergic and sympathomimetic combination therapy of asthma. *Annals of allergy* 1983;**50**:350.

Coe 1986 {published data only}

- * Coe CI, Barnes PJ. Reduction of nocturnal asthma by an inhaled anticholinergic drug. *Chest* 1986;**90**:485.

Dompeling 1992 {published data only}

- * Dompeling E, van Schayck CP, Molema J, Folgering H, van Grunsven PM, van Weel C. Inhaled beclomethasone improves the course of asthma and COPD. *European Respiratory Journal* 1992;**5**:945.

Dompeling 1993 {published data only}

- Dompeling E, van Schayck C, van Grunsven P, et al. Slowing the deterioration of asthma and chronic obstructive pulmonary disease observed during bronchodilator therapy by adding inhaled. *Annals of Internal Medicine* 1993;**118**:770.

Haahtela 1991 {published data only}

- * Haahtela T, Ahokas I, Ahonen A, van Assendelft A, Havu M, Havu K, et al. Inhaled bronchodilator's in asthma: low or high dose?. *Annals of Allergy* 1991;**66**:175.

Kreisman 1981 {published data only}

- Kreisman H, Frank H, Wolkove N, Gent M. Synergism between ipratropium and theophylline in asthma. *Thorax* 1981;**36**:387.

Laitinen 1979 {published data only}

- Laitinen LA, Poppius H, Haahtela T. Comparison of ipratropium bromide and salbutamol in a long-term trial in asthmatic and bronchitis patients in a cold climate. *Scandinavian Journal of Respiratory Diseases-Supplementum* 1979;**103**:163-9.

Light 1977 {published data only}

- Light RW, George RB. Oral atropine in the treatment of chronic asthma. *Annals of allergy* 1977;**38**(1):58-61.

Lightbody 1977 {published data only}

- Lightbody I, Ingram C, Legge JS, Johnston RN. Ipratropium bromide, salbutamol, and prednisolone in bronchial asthma and chronic bronchitis. *Thorax* 1977;**32**:645.

Lightbody 1978 {published data only}

- Lightbody IM, Ingram CG, Legge JS, Johnston RN. Ipratropium bromide, salbutamol and prednisolone in bronchial asthma and chronic bronchitis. *British Journal of Diseases of the Chest* 1978;**72**:181.

Macaluso 1986 {published data only}

- * Macaluso S, del Torre L. Efficacy and compliance to prolonged treatment of bronchospasm. *Respiration* 1986;**50**(Suppl 2):222-5.

Mazzei 1985 {published data only}

- Mazzei JA, Torres J. Blind randomized cross-over comparative study of salbutamol and the combination fenoterol-ipratropium IK6 in patients with bronchial asthma. *Respiration* 1986;**50**(Suppl 2):313.

- * Mazzei JA, Torres J. Comparative randomized blind cross over study between salbutamol and the fenoterol-ipratropium association (IK 6) in patients with bronchial asthma. *Prensa medica Argentina* 1985;**72**:329.

O'Driscoll 1990 {published data only}

- * O'Driscoll BR, Kay EA, Taylor RJ, Bernstein A. Home nebulizers: can optimal therapy be predicted by laboratory studies?. *Respiratory Medicine* 1990;**84**:471.

O'Driscoll 1992 {published data only}

- O'Driscoll BR, Kay EA, Taylor RJ, Weatherby H, Chetty MC, Bernstein A. A long-term prospective assessment of home nebulizer treatment. *Respiratory Medicine* 1992;**86**:317.

Pavia 1987 {published data only}

- * Pavia D, Lopez-Vidriero MT, Agnew JE, Taylor RG, Eyre-Brook A, Lawton WA, et al. Effect of four weeks' oxitropium bromide treatment on lung mucociliary clearance in patients with chronic bronchitis or asthma. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):15a.

Pavia 1989 {published data only}

- Pavia D, Lopez-Vidriero MT, Agnew JE, Taylor RG, Eyre-Brook A, Lawton WA, et al. Effect of four-week treatment with oxitropium bromide on lung mucociliary clearance in patients with chronic bronchitis or asthma. *Respiration* 1989;**55**:33.

Petersen 1979 {published data only}

- Petersen BN, Weeke E. A comparative study of the bronchodilating effects of ipratropium and terbutalin inhaled with Monaghan IPPB-M 515 by 19 patients with chronic asthma. *Scandinavian Journal of Respiratory Diseases* 1979;**103**:178.

Philip-Joet 1990 {published and unpublished data}

- * Philip-Joet F, Reynaud-Gaubert M, Jirou-Najou JL, Arnaud A. Comparison of Berodual and salbutamol in asthma: a multicenter evaluation. *Respiration* 1990;**57**:379.

Pierce 1982 {published and unpublished data}

- * Pierce RJ, Holmes PW, Campbell AH. Use of ipratropium bromide in patients with severe airways obstruction. *Australian and New Zealand journal of medicine* 1982;**12**:38.

Rebuck 1983 {published data only}

- * Rebuck AS, Gent M, Chapman KR. Anticholinergic and sympathomimetic combination therapy of asthma. *The Journal of Allergy and Clinical Immunology* 1983;**71**(3):317-23.

Ruffin 1990 {published data only}

- * Ruffin R, Blight M. Effect of inhaled oxitropium bromide in asthmatics. *Australian and New Zealand Journal of Medicine* 1990;**20**:525.

Salorinne 1988 {published data only}

- Salorinne Y, Ahonen A. Comparison of the bronchodilator effect of ipratropium bromide powder and two doses of ipratropium bromide aerosol in the treatment of asthma. *Current therapeutic research, clinical and experimental*. 1988;**43**(6):1073-80.

Storms 1986 {published data only}

- Storms WW, Bodman SF, Nathan RA, Busse WW, Bush RK, Falliers CJ, O'Hollaren JD, et al. Use of ipratropium bromide in asthma Results of a multi-clinic study. *The American Journal of Medicine* 1986;**81**:61.

Tammivaara 1993 {published data only}

- * Tammivaara R, Elo J, Mansury L. Terbutaline controlled-release tablets and ipratropium aerosol in nocturnal asthma. *Allergy* 1993;**48**:45.

Taylor 1986 {published data only}

- Taylor RG, Pavia D, Agnew J, Lopez-Vidriero M, Newman S, Lennard-Jones T, Clarke S. Effect of four weeks' high dose ipratropium bromide treatment on lung mucociliary clearance. *Thorax* 1986;**41**:295.

Taytard 1984 {published data only}

- * Taytard A, Auzeir J, Vergeret J, Tozon N, Freour P. Treatment of adult asthma: controlled double-blind clinical trial of oxitropium bromide. *European Journal of Clinical Pharmacology* 1984;**26**:429.

Tormey 1995 {published data only}

- Tormey VJ, Pathmakanthan S, Faul J, Leonard C, Poulter LW, Burke CM. A comparison of the effects of inhaled salbutamol or oxitropium bromide on bronchial hyperresponsiveness in asthmatic patients treated with. *Irish Journal of Medical Science* 1995;**164**:172.

van Schayck 1991a {published data only}

- van Schayck CP, Dompeling E, van Herwaarden CL, Folgering H, Verbeek AL, van der Hoogen HJ, et al. Bronchodilator treatment

in moderate asthma or chronic bronchitis: continuous or on demand? A randomised controlled study. *BMJ* 1991;**303**:1426.

van Schayck 1991b {published data only}

- van Schayck CP, Folgering H, Harbers H, Maas KL, van Weel C. Effects of allergy and age on responses to salbutamol and ipratropium bromide in moderate asthma and chronic bronchitis. *Thorax* 1991;**46**:355.

van Schayck 1994 {published data only}

- van Schayck CP, Folgering H, den Otter JJ, Tirimanna P, van Weel C. Does continuous use of bronchodilators disguise the progression of chronic nonspecific lung diseases?. *Tijdschrift voor geneeskunde* 1994;**50**:787.

van Schayck 1995 {published data only}

- van Schayck CP, Dompeling E, van Herwaarden CL, Folgering H, Akkermans RP, van den Broek PJ, et al. Continuous and on demand use of bronchodilators in patients with non-steroid dependent asthma and chronic bronchitis: four-year follow-up randomized controlled study. *British Journal of General Practice* 1995;**45**:239.

Vaughan 1988 {published and unpublished data}

- * Vaughan TR, Bowen RE, Goodman DL, Weber RW, Nelson HS. The development of subsensitivity to atropine methylnitrate A double-blind, placebo-controlled crossover study. *The American Review of Respiratory Disease* 1988;**138**:771.

Waite 1987 {published data only}

- Waite DA. Comparison of Duovent with salbutamol in chronic asthmatics - a long term study in community practice. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):4a.

Wheeler 1987 {published and unpublished data}

- * Wheeler AW, Guest PG, Britton MG. A three month study of oxitropium bromide in patients with bronchial asthma. *Postgraduate Medical Journal* 1987;**63**:8.

Wolstenholme 1989 {published and unpublished data}

- * Wolstenholme RJ, Shettar SP. Comparison of a combination of fenoterol with ipratropium bromide (Duovent) and salbutamol in young adults with nocturnal asthma. *Respiration* 1989;**55**:152.

References to studies excluded from this review

Abboud 1987 {published data only}

- Abboud R, Gibson N, Vickerson F, Mukherjee J. Bronchodilating effect of increasing doses of fenoterol (Berotec) given in combination with a standard dose of ipratropium (Atrovent) in asthma. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):5.

Adinoff 1998 {published data only}

- Adinoff AD, Schwartz HJ, Rickard KA, Yancey SW, Swearingen BE. Salmeterol compared with current therapies in chronic asthma. *Journal of Family Practice* 1998;**47**:278.

Advenier 1984 {published data only}

Advenier C, Cerrina J, Duroux P. Bronchodilators. *Semaine Des Hopitaux* 1984;**60**:179.

Alanko 1983 {published data only}

Alanko K, Sahlstrom K, Harkonen R. Combination of fenoterol and ipratropium bromide in the treatment of bronchial asthma. *European Journal of Respiratory Diseases-Supplement* 1983;**128**(Pt 2):546.

Allen 1980a {published data only}

Allen CJ, Campbell AH. Comparison of inhaled atropine sulphate and atropine methonitrate. *Thorax* 1980;**35**:932.

Allen 1980b {published data only}

Allen CJ, Campbell AH. Dose response of ipratropium bromide assessed by two methods. *Thorax* 1980;**35**:137.

Alliott 1972 {published data only}

Alliott RJ, Lang BD, Rawson DR, Leckie WJ. Effects of salbutamol and isoprenaline-phenylephrine in reversible airways obstruction. *BMJ* 1972;**1**:539.

Altounyan 1964 {published data only}

Altounyan RE. Variation of drug action on airway obstruction in man. *Thorax* 1964;**19**:406.

Andersen 1986 {published data only}

Andersen L. Pulmonary function and acid application in the esophagus. *Chest* 1986;**90**:358.

Anderson 1980 {published data only}

Anderson G, Jariwalla AG, Turnbull P. The effects of fenoterol, ipratropium bromide and their combination in asthma and chronic bronchitis. *British Journal of Diseases of the Chest* 1980;**74**:416.

Anderson 1981 {published data only}

Anderson G, Jariwalla AG, Turnbull P, Veale J. The effects of fenoterol, ipratropium bromide and their combination in asthma and chronic bronchitis. *IRCS (International Research Communications System) Medical Science* 1981;**9**:251.

Anderson 1983 {published data only}

Anderson G. Anticholinergic drugs in airways obstruction. *Postgraduate Medical Journal* 1983;**59**(Suppl 3):64.

Aquilina 1986 {published data only}

Aquilina R, Bergero F, Noceti P, Mirabelli S, Luciani G. Protective effect of Duovent versus salbutamol in long-term treatment. *Respiration* 1986;**50**(Suppl 2):240.

Ariano 1981 {published data only}

Ariano R, Giacca S. Variations of individual susceptibility to beta-adrenergic and anticholinergic bronchodilator drugs. *Minerva Pneumologica* 1981;**20**:141.

Baigelman 1977 {published data only}

Baigelman W, Chodosh S. Bronchodilator action of the anticholinergic drug, ipratropium bromide (Sch 1000), as an aerosol in chronic bronchitis and asthma. *Chest* 1977;**71**:324.

Banos 1983 {published data only}

Banos Hidalgo P, Ramos Martos A, Cabrera Moreno R, Luque Leal J, Palacios Giner A. Comparative double blind study of the aerosol association of ipratropium bromide and salbutamol with its individual effect in chronic bronchitis. *Revista Clinica Espanola* 1983;**171**:265.

Barber 1975 {published data only}

Barber P, Chatterjee SS, Scott R. The effect of two anticholinergic bronchodilators on ventilatory function in patients with chronic bronchitis or asthma as measured by spirometry. *Postgraduate Medical Journal* 1975;**51**(Suppl):112.

Barber 1977 {published data only}

Barber PV, Chatterjee SS, Scott R. A comparison of ipratropium bromide, deproline citrate and placebo in asthma and chronic bronchitis. *British Journal of Diseases of the Chest* 1977;**71**:101.

Bauer 1973 {published data only}

Bauer P, Kummer F. Double-blind study for the comparison of the broncholytic effect of a new tropic acid derivative with isoprenaline. *International Journal of Clinical Pharmacology and Therapeutics* 1973;**8**:135.

Beasley 1987 {published data only}

Beasley CR, Rafferty P, Holgate ST. Bronchoconstrictor properties of preservatives in ipratropium bromide (Atrovent) nebuliser solution. *BMJ* 1987;**294**:1197.

Bell 1982 {published data only}

Bell R, Sahay JN, Barber PV, Chatterjee SS, Cox G. A therapeutic comparison of ipratropium bromide and salbutamol in asthmatic patients. *Current Medical Research & Opinion* 1982;**8**:242.

Bellia 1987 {published data only}

* Bellia V, Cibella F, Insalaco G, Mirto M, Peralta G, Visconti A, Schiassi M. Comparison of the effectiveness of oxitropium bromide and of a slow-release theophylline in the prevention of 'morning dip' in asthmatics. *Postgraduate Medical Journal* 1987;**63**:9.

Bernabeu 1987 {published data only}

Bernabeu L, Bourcereau J, Delacenserie R, Prosper M, Salen JM, Vincent J, Jirou-Najou JL. Ipratropium bromide -fenoterol combination: optimum dosage ratio in asthma. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):6.

Bezel 1987 {published data only}

Bezel R, Brandli O, Braun P, Haegi V, Villiger B. Additive effect of sustained-release theophylline combined with anticholinergic and sympathomimetic therapy in COPD. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):22.

Bish 1997 {published data only}

Bish R, Thien F, Whitehead F, Walters J, Walters EH. The use of regular ipratropium bromide to reduce requirements for beta-2 agonists in symptomatic asthmatic subjects [abstract]. *Australian and New Zealand Journal of Medicine* 1997;**27**:246.

Bohadana 1980 {published data only}

Bohadana AB, Peslin R, Teculescu D, Polu JM, et al. The bronchodilator action of theophylline aerosol in subjects with chronic airflow obstruction. *Clinical Respiratory Physiology* 1980;**16**:13.

Bonsignore 1986 {published data only}

Bonsignore G, Bellia V, Peralta G, Alessi N, Migliara G. The combination of fenoterol and ipratropium bromide in bronchial asthma: comparison of the acute effects of two different dosages. *Respiration* 1986;**50**(Suppl 2):148.

Botts 1985 {published data only}

Botts LD, Pingleton SK, Schroeder CE, Robinson RG, Hurwitz A. Prolongation of gastric emptying by aerosolized atropine. *The American Review of Respiratory Disease* 1985;**131**:725.

Brady 1979 {published data only}

Brady RE, Easton JG. The value of atropine in the documentation of reversible airways obstruction. *Annals of Allergy* 1979;**42**:211.

Brand 1992 {published data only}

* Brand PL, Kerstjens HA, Postma DS, Sterk PJ, Quanjer PH, Sluiter HJ, et al. Long-term multicentre trial in chronic nonspecific lung disease: methodology and baseline assessment in adult patients Dutch CNSLD Study Group. *European Respiratory Journal* 1992;**5**:21.

Bremner 1992 {published data only}

Bremner P, Burgess C, Beasley R, Woodman K, Marshall S, Crane J, Pearce N. Nebulized fenoterol causes greater cardiovascular and hypokalaemic effects than equivalent bronchodilator doses of salbutamol in asthmatics. *Respiratory Medicine* 1992;**86**:419.

Britton 1988 {published data only}

Britton J, Hanley SP, Garrett HV, Hadfield JW, Tattersfield AE. Dose related effects of salbutamol and ipratropium bromide on airway calibre and reactivity in subjects with asthma. *Thorax* 1988;**43**:300.

Brom 1985 {published data only}

Brom AL. Prophylaxis of bronchial asthma in adults with the use of ketotifen : Prolongation of gastric emptying by aerosolized atropine. *The American review of respiratory disease* 1985;**8**:725.

Bruderman 1983 {published data only}

Bruderman I, Cohen-Aronovski R, Smorzik J. A comparative study of various combinations of ipratropium bromide and metaproterenol in allergic asthmatic patients. *Chest* 1983;**83**:208.

Brumby 1980 {published data only}

* Brumby KH. Comparative clinical study of the efficacy of Atrovent and Berotec, singly and combined, with comparison of placebo. *Therapiewoche* 1980;**30**:2796.

Bryant 1990a {published data only}

Bryant DH, Rogers P. Oxitropium bromide: an acute dose response study of a new anticholinergic drug in combination

with fenoterol in asthma and chronic bronchitis. *Pulmonary Pharmacology* 1990;**3**:55.

Bryant 1990b {published data only}

Bryant D, Rogers P. Comparison of preservative-free ipratropium bromide and atrovent nebuliser solution in treatment of asthma [abstract]. *Australian and New Zealand Journal of Medicine* 1990;**20**:525.

Bryant 1992 {published data only}

Bryant DH, Rogers P. Effects of ipratropium bromide nebulizer solution with and without preservatives in the treatment of acute and stable asthma. *Chest* 1992;**102**:742.

Burge 1980 {published data only}

Burge PS, Harries MG, l'Anson E. Comparison of atropine with ipratropium bromide in patients with reversible airways obstruction unresponsive to salbutamol. *British Journal of Diseases of the Chest* 1980;**74**:259.

Burki 1997 {published data only}

Burki NK. The effects of the combination of inhaled ipratropium and oral theophylline in asthma. *Chest* 1997;**111**:1509.

Cañete 1991 {published data only}

Cañete CE. Treatment with anticholinergic drugs in asthmatic crisis. *Archivos De Bronconeumología Suppl* 1991;**27**:32.

Capel 1964 {published data only}

Capel LH, Fletcher EC. Spirometric integrals after isoprenaline and atropine in asthma. *British Journal of Diseases of the Chest* 1964;**58**:174.

Carpentiere 1986 {published data only}

Carpentiere G, Marino S, Castello F, Bonanno CT. Comparison between morning and afternoon effects of inhaling a combined preparation of fenoterol with ipratropium bromide. *Current Therapeutic Research* 1986;**40**:265.

Carratu 1983 {published data only}

Carratu L, Sofia M. Comparison of the effects of inhalation of a single dose of a ipratropium bromide-fenoterol combination with a single dose of fenoterol. *Archivio Monaldi Per La Tisiologia e Le Malattie Dell Apparato Respiratorio* 1983;**38**:263.

Casali 1979 {published data only}

Casali L, Grassi C, Rampulla C, Rossi A. Clinical pharmacology of a combination of bronchodilators. *International Journal of Clinical Pharmacology and Biopharmacy* 1979;**17**:277.

Catterall 1988 {published data only}

Catterall JR, Rhind GB, Whyte KF, Shapiro CM, Douglas NJ. Is nocturnal asthma caused by changes in airway cholinergic activity?. *Thorax* 1988;**43**:720.

Chakravarti 1987 {published data only}

Chakravarti A, Pratley M, Barcharb J, Musiosky M, McDonald F, Vickerson F, et al. Ipratropium-fenoterol interaction study in asthmatic adults. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):5.

Chervinsky 1977 {published data only}

Chervinsky P. Double-blind study of ipratropium bromide, a new anticholinergic bronchodilator. *Journal of Allergy and Clinical Immunology* 1977;**59**:22.

Chhabra 2002 {published data only}

Chhabra SK, Pandey KK. Comparison of acute bronchodilator effects of inhaled ipratropium bromide and salbutamol in bronchial asthma. *Journal of Asthma* 2002;**39**:375.

Chick 1977 {published data only}

Chick TW, Jenne JW. Comparative bronchodilator responses to atropine and terbutaline in asthma and chronic bronchitis. *Chest* 1977;**72**:719.

Chuchalin 1985 {published data only}

Chuchalin AG, Konganova NA, Pashkova TL, Treskunov VK, Apul'tsina ID. Berodual in the treatment of bronchial asthma. *Sovetskaya Meditsina* 1985;**11**:81.

Ciappi 1986 {published data only}

Ciappi G, Magnini P, Valente S, Toscano L, Corbo GM, Culla G, Patalano F. Dose-response relationship: fenoterol, ipratropium bromide and their combination. *Respiration* 1986;**50**(Suppl 2):140.

Clauzel 1987 {published data only}

Clauzel A, Michel F, Jirou-Najou J. Study of dose response curves of fenoterol, ipratropium bromide and Berodual. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):13.

Cockcroft 1977 {published data only}

Cockcroft DW, Killian DN, Mellon JJ, Hargreave FE. Protective effect of drugs on histamine-induced asthma. *Thorax* 1977;**32**:429.

Cox 1984 {published data only}

Cox ID, Hughes DT, McDonnell KA. Ipratropium bromide in patients with nocturnal asthma. *Postgraduate Medical Journal* 1984;**60**:526.

Crane 1986a {published data only}

Crane J, Gamble S. Single dose comparison of salbutamol and a fenoterol/ipratropium combination from metered aerosols in patients with asthma. *New Zealand Medical Journal* 1986;**99**:420.

Crane 1986b {published data only}

Crane J. Single-dose comparison of salbutamol and Duovent-Berodual in asthma. *Respiration* 1986;**50**(Suppl 2):285.

Crane 1987 {published data only}

Crane J, Town G, O'donnell T. Duovent in obstructive airways disease: comparison with salbutamol by metered aerosol in respect of degree and duration of bronchodilation in. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):3.

del Buffalo 1982 {published data only}

del Bufalo DC, Fabbri M, Gunella G, Longi R. Effetti sulla funzione ventilatoria dell'associazione fenoterolo-bromuro di

ipratropium. *Rivista di Patologia e Clinica della Tuberculosis e di Pneumologia* 1982;**53**:637.

Donkers 1998 {published data only}

Donkers JM, van Schayck CP, Akkermans RP. The difference in bronchodilation effect of a beta two-mimetic and an anticholinergic agent in patients with asthma or chronic obstructive lung. *Ned Tijdschr Geneeskde* 1998;**142**:1419.

Donner 1980 {published data only}

Donner CF. [IS there a placebo ie beclo plus (IB vs placebo?)] Bronchospasm management by means of ipratropium, beclomethasone and their combination. *Giornale Italiano della Tuberculosis e Delle Malattie del Torace* 1980;**34**:307.

Dutt 1990 {published data only}

Dutt NS, Reddy TC, Raghuvver TS, Yeshwanth M, Sunil Dutt N, Chinnappa Reddy T, et al. A comparative study of atropine sulphate, terbutaline and their combination in asthmatic children. *Lung India* 1990;**8**(2):73-5.

Elwood 1982 {published data only}

Elwood RK, Abboud RT. The short-term bronchodilator effects of fenoterol and ipratropium in asthma. *Journal of Allergy and Clinical Immunology* 1982;**69**:467.

Emirgil 1975 {published data only}

Emirgil C, Dwyer K, Baskette P, Sobol BJ. A new parasympatholytic bronchodilator: a study of its onset of effect after inhalation. *Current Therapeutic Research* 1975;**17**:215.

Endoh 1998 {published data only}

Endoh N, Ichinose M, Takahashi T, Miura M, Kageyama N, Mashito Y, et al. Relationship between cholinergic airway tone and serum immunoglobulin E in human subjects. *European Respiratory Journal* 1998;**12**:71.

Filiz 1994 {published data only}

Filiz A, Ekinci E, Dikensoy O, Bulgur D, Oz M. Comparison of a single dose of aerosol salbutamol and fenoterol/ipratropium on bronchial asthmatic patients. *Delaware Medical Journal* 1994;**66**:549.

Firstater 1981 {published data only}

Firstater E, Mizrahi E, Topilsky B. The effect of vagolytic drugs on airway obstruction in patients with bronchial asthma. *Annals of Allergy* 1981;**46**:332.

Fischbacher 1984 {published data only}

Fischbacher CM, Milroy R, Giannini D, Monie RD. Comparison of Duovent and salbutamol inhalers in chronic stable asthma. *Postgraduate Medical Journal* 1984;**60**(Suppl 1):28.

Flint 1983a {published data only}

Flint KC, Hockley B, Johnson NM. A comparison between a combination of ipratropium bromide plus fenoterol in a single metered dose inhaler (Duovent) and salbutamol in asthma. *Postgraduate Medical Journal* 1983;**59**:724.

Flint 1983b {published data only}

Flint K, Hockley B, Johnson NM. Salbutamol versus Duovent (a combination of fenoterol and ipratropium bromide) in asthma. *European Journal of Respiratory Diseases-Suppl* 1983;**128**(Pt 2):548.

Flint 1984 {published data only}

Flint K, Hockley B, Johnson NM. A comparison between a combination of ipratropium bromide plus fenoterol in a single metered dose inhaler (Duovent) and salbutamol in asthma. *Postgraduate Medical Journal* 1984;**60**(Suppl 1):23.

Foster 1983 {published data only}

Foster WM, Langenback EG, Glaser ML, Bergofsky EH. Acute effect of ipratropium bromide at therapeutic dose on mucus transport of adult asthmatics. *European Journal of Respiratory Diseases-Suppl* 1983;**128**(Pt 2):554.

Freeman 1992 {published data only}

Freeman W, Javaid A, Cayton RM. The effect of ipratropium bromide on maximal exercise capacity in asthmatic and non-asthmatic men. *Respiratory Medicine* 1992;**86**:151.

Frith 1987 {published data only}

Frith PA, Atkinson J. Airway anticholinergic effects of oxitropium. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):5.

Gaba 1987 {published data only}

Gaba S, Trigui F, Michel FB, Jirou-Najou JL. Circadian rhythms in asthmatics: comparison of Berotec, Atrovent and Berodual bronchodilating action. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):13.

Gillies 1987 {published data only}

Gillies AJ, Stewart D. A dose response study comparing Duovent with salbutamol. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):19.

Ginesu 1983 {published data only}

Ginesu F, Ortu AR, Schena GP, Deiola G, Polo F. Controlled clinical study of the effects of a single aerosol administration of a fenoterol-ipratropium bromide combination in comparison with. *Archivio Monaldi Per La Tisiologia e Le Malattie Dell Apparato Respiratorio* 1983;**38**:241.

Gotz 1983 {published data only}

Gotz RS. Effect of the inhalation of an atropine derivative (ipratropium bromide) in powder form on respiratory mechanics in obstructive airway. *Praxis und Klinik der Pneumologie* 1983;**37**(Suppl 1):927.

Grandordy 1988 {published data only}

Grandordy BM, Thomas V, de Lauture D, Marsac J. Cumulative dose-response curves for assessing combined effects of salbutamol and ipratropium bromide in chronic asthma. *European Respiratory Journal* 1988;**1**:531.

Gross 1975a {published data only}

Gross NJ. Sch 1000: a new anticholinergic bronchodilator. *The American Review of Respiratory Disease* 1975;**112**:823.

Gross 1975b {published data only}

Gross HJ. Dose-response study of the effect of Sch 1000 MDI on forced vital capacity (FVC), FEV1.0, maximal mid-expiratory flow (MMEF) and heart rate, ECG. *Postgraduate Medical Journal* 1975;**51**(Suppl 7):95.

Gross 1984 {published data only}

Gross NJ, Skorodin MS. Anticholinergic, antimuscarinic bronchodilators[Review] [243 refs]. *The American review of respiratory disease* 1984;**129**:856.

Gross 1987 {published data only}

Gross NJ, Bankwala Z. Effects of an anticholinergic bronchodilator on arterial blood gases of hypoxemic patients with chronic obstructive pulmonary disease;. *The American review of respiratory disease* 1987;**136**:1091.

Gutersohn 1975 {published data only}

Gutersohn. The effect of Raw on Sch1000 MDI or fenoterol MDI and the combined administration of subthreshold dosages of both compounds. *Postgraduate Medical Journal* 1975;**51**(Suppl 7):113.

Guyatt 1986 {published data only}

Guyatt G, Sackett D, Taylor DW, Chong J, Roberts R, Pugsley S. Determining optimal therapy Randomized trials in individual patients. *New England Journal of Medicine* 1986;**314**:889.

Higgins 1991 {published data only}

Higgins BG, Powell RM, Cooper S, Tattersfield AE. Effect of salbutamol and ipratropium bromide on airway calibre and bronchial reactivity in asthma and chronic bronchitis. *European Respiratory Journal* 1991;**4**:415.

Hockley 1985 {published data only}

Hockley B, Johnson NM. A comparison of three high doses of ipratropium bromide in chronic asthma. *British Journal of Diseases of the Chest* 1985;**79**:379.

Hoffstein 1994 {published data only}

Hoffstein V, Zamel N, McClean P, Chapman KR. Changes in pulmonary function and cross-sectional area of trachea and bronchi in asthmatics following inhalation of procaterol hydrochloride and. *American Journal of Respiratory and Critical Care Medicine* 1994;**149**:81.

Holst 1976 {published data only}

* Holst PE, Gebbie T, Martin PD, O'Donnell TV. A long term trial of ipratropium in asthmatic patients. *Australian and New Zealand Journal of Medicine* 1976;**11**:367.

Huhti 1986 {published data only}

Huhti E, Poukkula A. Comparison of fenoterol, ipratropium bromide, and their combination in patients with asthma or chronic airflow obstruction. *Respiration* 1986;**50**(Suppl 2):298.

Hunt 1983 {published data only}

Hunt D, MacDonald GF, Reilly P, Plumley D, Kazim F. Bronchodilator response to several doses of ipratropium bromide in asthmatics and bronchitics. *Current Therapeutic Research, Clinical and Experimental*. 1983;**33**:651.

Irsigler 1975 {published data only}

Irsigler G. Flow volume curves and changes in FEV1.0, forced vital capacity (FVC), mid and slow vital capacity (Mid VC and SVC) in patients with bronchial. *Postgraduate medical journal Suppl* 1975;**51**:96.

Jenkins 1981 {published data only}

Jenkins CR, Chow CM, Fisher BL, Marlin GE. Comparison of ipratropium bromide and salbutamol by aerosolized solution. *Australian and New Zealand Journal of Medicine* 1981;**11**:513.

Jenkins 1982 {published data only}

Jenkins CR, Chow CM, Fisher BL, Marlin GE. Ipratropium bromide and fenoterol by aerosolized solution. *British Journal of Clinical Pharmacology* 1982;**14**:113.

Jindal 1989 {published data only}

Jindal SK, Kaur SJ. Relative bronchodilatory responsiveness attributable to sympathetic and parasympathetic activity in bronchial asthma. *Respiration* 1989;**56**:16.

Johnson 1989 {published data only}

Johnson MA, Newman SP, Bloom R, Talaei N, Clarke SW. Delivery of albuterol and ipratropium bromide from two nebulizer systems in chronic stable asthma- Efficacy and pulmonary deposition. *Chest* 1989;**96**:6.

Jolobe 1981 {published data only}

Jolobe OM, Lane DJ. Atropine responsiveness in asthma in relation to steroid aerosol therapy. *British Journal of Diseases of the Chest* 1981;**75**:413.

Kaik 1975 {published data only}

Kaik G. Untersuchungen mit dem neuen Bronchospamolytikum Ipratropiumbromid (Sch 1000, Atrovent). *Therapiewoche* 1975;**25**:2428.

Kaik 1976 {published data only}

Kaik G. [Whole-body plethysmographic studies on the bronchospasmolytic effect of 2 antiasthmatic drug combinations and ipratropium bromide]. *Wiener Medizinische Wochenschrift* 1976;**126**:127.

Kaik 1979 {published data only}

Kaik G. [Lung function studies with the bronchodilator terbutaline]. *Acta medica Austriaca* 1979;**6**:1.

Kamburoff 1975 {published data only}

Kamburoff PL, Gunthner W. Dose-dependent effects of Sch 1000 MDI on the airways resistance indices in patients with chronic airways obstruction. *Postgraduate Medical Journal* 1975;**51**(Suppl 7):97.

Kaptein 1993 {published data only}

Kaptein AA, Brand PL, Dekker FW, Kerstjens HA, Postma DS, Sluiter HJ. Quality-of-life in a long-term multicentre trial in chronic nonspecific lung disease: Assessment at baseline. *European Respiratory Journal* 1993;**6**:1479.

Kennedy 1964 {published data only}

Kennedy MC, Thursby-Pelham DC. Some adrenergic drugs and atropine methonitrate given by inhalation for asthma: a comparative study. *BMJ* 1964;**1**:1018.

Kerstjens 1992 {published data only}

Kerstjens HA, Brand PL, Hughes MD, Robinson NJ, Postma DS, Sluiter HJ, et al. A comparison of bronchodilator therapy with or without inhaled corticosteroid therapy for obstructive airways disease. *New England Journal of Medicine* 1992;**327**:1413.

Kerstjens 1993a {published data only}

Kerstjens HA, Brand PL, Quanjer PH, van der Bruggen-Bogaarts Koeter GH, Postma DS. Variability of bronchodilator response and effects of inhaled corticosteroid treatment in obstructive airways disease. Dutch CNSLD Study Group. *Thorax* 1993;**48**:722.

Kerstjens 1993b {published data only}

Kerstjens HA. Airways hyperresponsiveness, bronchodilator response, allergy and smoking predict improvement in FEV1 during long-term inhaled corticosteroid. *European respiratory journal* 1993;**6**:868.

Kerstjens 1994 {published data only}

Kerstjens HA, Brand PL, de Jong PM, Koeter GH, Postma DS. Influence of treatment on peak expiratory flow and its relation to airway hyperresponsiveness and symptoms. *Thorax* 1994;**49**:1109.

Kerstjens 1995 {published data only}

Kerstjens HA, Schouten JP, Brand PLP, Schoonbrood DF, Sterk PJ, Postma DS, et al. Importance of total serum IgE for improvement in airways hyperresponsiveness with inhaled corticosteroids in asthma and chronic obstructive. *American Journal of Respiratory & Critical Care Medicine* 1995;**151**:360.

Kok-Jensen 1979 {published data only}

Kok-Jensen AA. Combined inhalation of salbutamol and ipratropin in chronic bronchial obstruction [Kombineret inhalation af salbutamol og ipratropin ved kronisk bronkial obstruktion]. *Ugeskrift for Laeger* 1979;**141**(30):2039-41.

Kradjan 1992 {published data only}

Kradjan WA, Driesner NK, Abuan TH, Emmick G, Schoene RB. Effect of age on bronchodilator response. *Chest* 1992;**101**:1545.

Kreisman 1984 {published data only}

Kreisman H, Cohen C, Ghezzi H, Vickerson F, Frank H, Wolkove N. Combined therapy with ipratropium and theophylline in asthma. *Annals of Allergy* 1984;**52**:90.

Krivoy 1987 {published data only}

Krivoy N, Jubarin A, Gaitini L, Rubin A, Alroy G. Combination therapy of ipratropium bromide, terbutaline and slow release theophylline in asthma and chronic airways obstruction. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):21.

Kunkel 1976 {published data only}

Kunkel GS. The importance of nervus vagus for bronchial obstruction of allergic and non-allergic origin and the therapeutic consequences. *Praxis der Pneumologie* 1976;**30**:459.

Kunkel 2000 {published data only}

Kunkel G, Magnussen H, Bergmann KC, Juergens UR, de Mey C, Freund E, Hinzmann R, Beckers B. Respimat (a new soft mist inhaler) delivering fenoterol plus ipratropium bromide provides equivalent bronchodilation at half the cumulative dose. *Respiration* 2000;**67**:306.

Lahdensuo 1975 {published data only}

Lahdensuo A, Viljanen A, Muittari A. A comparative study on the effect of Sch1000 MDI, salbutamol and placebo MDI in bronchial asthma. *Postgraduate Medical Journal* 1975;**51**(Suppl 7):116.

Laitinen 1986 {published data only}

Laitinen LA, Poppius H. Combination of oxitropium bromide and salbutamol in the treatment of asthma with pressurized aerosols. *British Journal of Diseases of the Chest* 1986;**80**:179.

Lanser 1975 {published data only}

Lanser KG, Foldi M, Sill V. The effect of Tropaacidester Sch 1000 on respiratory resistance [Therapie der obstruktiven Atemwegkrankung mit einem Atropinderivat (Sch 1000)]. *Medizinische Klinik* 1975;**70**(9):380-4.

Larsson 1987 {published data only}

Larsson K. Ipratropium bromide: bronchodilator action and effect on methacholine-induced bronchoconstriction. *Journal of Asthma* 1987;**24**:29.

Lavandier 1987 {published data only}

Lavandier M, Bourcereau J, Jirou-Najou J. Comparison of bronchodilating onset of action of Berotec, Atrovent and Berodual. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):13.

Lawford 1983 {published data only}

Lawford P, Palmer KN. A single dose comparison of a combination of fenoterol and ipratropium aerosols in bronchial asthma. *Postgraduate Medical Journal* 1983;**59**:28.

Lefcoe 1982 {published data only}

Lefcoe NM, Toogood JH, Blennerhassett G, Baskerville J, Paterson NA. The addition of an aerosol anticholinergic to an oral beta agonist plus theophylline in asthma and bronchitis A double-blind single dose study. *Chest* 1982;**82**:300.

Lehrer 1993 {published data only}

Lehrer PM, Hochron SM, Isenberg S, Rausch L, Carr R. The Asthma Symptom Profile: a psychophysically based scale for assessment of asthma symptoms. *Journal of Psychosomatic Research* 1993;**37**:515.

Lehrer 1994 {published data only}

Lehrer PM, Hochron SM, Rausch L, Carr R. Effects of aerosol ipratropium bromide on cardiac vagal tone. *Chest* 1994;**105**:1701.

Li 2003 {published data only}

Li SF, Yao WJ. Combined application of salbutamol and ipratropium bromide in the treatment of bronchial asthma in children. *Herald of Medicine* 2003;**22**(2):96-7.

Lichterfeld 1979 {published data only}

Lichterfeld A. Safety of Atrovent. *Scandinavian journal of respiratory diseases Suppl* 1979;**103**:143.

Linehan 1975 {published data only}

Linehan WD. The effect of Sch1000 and terbutaline given as MDI on ventilatory function in patients with asthma as measured by spirometry. *Postgraduate medical journal Suppl* 7 1975;**51**:116.

Maesen 1975a {published data only}

Maesen F, Buytendijk HJ. Dose- and time-response curves of Sch 1000 MDI and placebo. Differential response in asthmatics and bronchitics as measured by FEV1.0, vital. *Postgraduate Medical Journal* 1975;**51**(Suppl 7):97.

Maesen 1975b {published data only}

Maesen FP. Comparison of the effects of Sch 1000 MDI and fenoterol MDI in patients with bronchial asthma and chronic bronchitis. *Postgraduate Medical Journal* 1975;**51**(Suppl 7):116.

Maesen 1982 {published data only}

Maesen FP, Smeets JJ, Rubingh G. Vergelijking van het bronchospasmolytisch effect van druk-aerosolen met fenoterol (Berotec) en van een combinatie van fenoterol en een. *Tijdschrift voor geneeskunde* 1982;**7**:1184.

Maesen 1985 {published data only}

Maesen FP, Smeets JJ, Pover GM. Comparison of the effects of a combination inhaler containing salbutamol and atropine methonitrate By comparison with each component given. *Tijdschrift voor Therapie Geneesmiddel en Onderzoek* 1985;**10**:1000.

Mann 1984 {published data only}

Mann JS, Howarth PH, Holgate ST. Bronchoconstriction induced by ipratropium bromide in asthma: relation to hypotonicity. *BMJ* 1984;**289**:469.

Marcatili 1984 {published data only}

Marcatili S, Maselli R. Effects of administration of a single dose of iprafen (I puff) compared with those of a single dose of ipratropium bromide (40 mcg). *Archivio Monaldi Per La Tisiologia e Le Malattie Dell Apparato Respiratorio* 1984;**39**:37.

Marin 1987 {published data only}

Marin J, Servera E, Verara P. Efficacy devoid of side effects in the combination of an atropine-like bronchodilator and a low dose of a beta2-mimetic. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):18.

Marlin 1978 {published data only}

Marlin GE, Bush DE, Berend N. Comparison of ipratropium bromide and fenoterol in asthma and chronic bronchitis. *British Journal of Clinical Pharmacology* 1978;**6**:547.

Marlin 1979 {published data only}

Marlin GE, Berend N, Harrison AC. Combined cholinergic antagonist and beta 2-adrenoceptor agonist bronchodilator therapy by inhalation. *Australian and New Zealand Journal of Medicine* 1979;**9**:511.

Mattila 1966 {published data only}

Mattila M, Muittari A. The effect of bronchodilator aerosols on the peak expiratory flow rate in asthmatic patients. *Acta Medica Scandinavica* 1966;**180**:421.

McIntosh 1986a {published data only}

McIntosh D. A comparison between fenoterol/ipratropium bromide combinations and salbutamol administered as nebulizer solutions assessing dose-response. *Pharmatherapeutica* 1986;**4**:416.

McIntosh 1986b {published data only}

McIntosh D. Comparison between fenoterol/ipratropium bromide combinations and salbutamol administered as nebulizer solution assessing dose response and duration of action. *Respiration* 1986;**50**(Suppl 2):310-2.

Millar 1990 {published data only}

Millar AB, Bush A, Al-Hillawi H, Goldman J, Denison DM. Ipratropium bromide: are patients treated with optimal therapy?.. *Postgraduate Medical Journal* 1990;**66**:1040.

Minette 1979 {published data only}

Minette A, Marcq M. Oxitropium bromide (Ba 253), an advance in the field of anticholinergic bronchodilating treatments Preliminary results. *Revue De l Institut d Hygiene Des Mines* 1979;**34**:115.

Minette 1985 {published data only}

Minette P, Dubois P, Delwiche JP. Validity of air-helium DVmax measurements in trials of bronchodilators. *Bulletin Europeen de Physiopathologie Respiratoire* 1985;**21**:357.

Morrison 1988 {published data only}

Morrison JF, Pearson SB, Dean HG. Parasympathetic nervous system in nocturnal asthma. *BMJ* 1988;**296**:1427.

Morrison 1989 {published data only}

Morrison JF, Pearson SB. The effect of the circadian rhythm of vagal activity on bronchomotor tone in asthma. *British Journal of Clinical Pharmacology* 1989;**28**:545.

Morrison 1991 {published data only}

Morrison JF, Pearson S. The parasympathetic nervous system and the diurnal variation of lung mechanics in asthma. *Respiratory Medicine* 1991;**85**:285.

Morton 1984 {published data only}

Morton O. Response to Duovent of chronic reversible airways obstruction--a controlled trial in general practice. *Postgraduate Medical Journal* 1984;**60**(Suppl 1):32.

Muittari 1968 {published data only}

Muittari A. Atropine methonitrate in bronchial asthma. *Annales Medicinæ Internæ Fenniae* 1968;**57**:193.

Nishi 1993 {published data only}

Nishi K, Fujimura M, Myou S, Ooka T, Sakamoto S, Saitou M, et al. Comparison of the bronchodilator activities of oxitropium bromide, fenoterol, and their combination in patients with chronic obstructive. *Clinical Autonomic Research* 1993;**3**:41.

Nishimura 1999 {published data only}

Nishimura K, Koyama H, Ishihara K, Hasegawa T, Katayama S, Nakashima A, Izumi T. Additive effect of oxitropium bromide in combination with inhaled corticosteroids in the treatment of elderly patients with chronic asthma. *Allergology International* 1999;**48**:85.

Nolte 1978 {published data only}

Nolte D. Double-blind crossover comparison between a beta-adrenergic agent and a new anticholinergic agent by metered dose inhaler. *Respiration* 1978;**36**:32.

Osterwalder 1991 {published data only}

Osterwalder JJ. Randomized and controlled single case-study - A scientific approach to difficult therapeutic decisions A practical example for use of. *Schweizerische Rundschau Fuer Medizin Praxis* 1991;**80**:1168.

Pavia 1979 {published data only}

Pavia D, Bateman JR, Sheahan NF, Clarke SW. Effect of ipratropium bromide on mucociliary clearance and pulmonary function in reversible airways obstruction. *Thorax* 1979;**34**:501.

Peel 1984 {published data only}

Peel ET, Anderson G, Cheong B, Broderick N. A comparison of oxitropium bromide and ipratropium bromide in asthma. *European Journal of Respiratory Diseases* 1984;**65**:106.

Peerless 1983 {published data only}

Peerless AG, Rachelefsky GS, Mickey MR, Katz RM, Siegel SC. Bronchodilator characteristics of nebulized metaproterenol sulfate, isoetharine, and atropine in chronic asthma. *Journal of Allergy and Clinical Immunology* 1983;**72**:702.

Petrie 1975a {published data only}

Petrie GR, Palmer KN. Comparison of aerosol ipratropium bromide and salbutamol in chronic bronchitis and asthma. *BMJ* 1975;**1**:430.

Petrie 1975b {published data only}

Petrie GR, Palmer K. Comparative trial of Sch1000 MDI and salbutamol MDI in chronic bronchitis and bronchial asthma. *Postgraduate Medical Journal* 1975;**51**(Suppl 7):117.

Petro 1981 {published data only}

Petro W. Bronchodilator effect of a combination of a beta-adrenergic agonist and a cholinergic antagonist [Broncholyse unter Kombiniertes anwendung]. *Praxis und Klinik der Pneumologie* 1981;**35**(1 Spec):779.

Philip-Joet 1986 {published data only}

Philip-Joet F, Herkert A, Vervloet D, Arnaud A. [Comparative study of anticholinergic and betamimetic agents in the treatment of asthma attacks]. *Revue De Pneumologie Clinique* 1986;**42**:292.

Pierce 1979 {published data only}

Pierce RJ, Allen CJ, Campbell AH. A comparative study of atropine methonitrate, salbutamol, and their combination in airways obstruction. *Thorax* 1979;**34**:45.

Pounsford 1983 {published data only}

Pounsford JC, Fuller RW, Saunders KB. A cumulative dose response study of oxitropium bromide (Ba 253) in asthmatics. *European Journal of Respiratory Diseases-Supplement* 1983;**128**(Pt 2):536-9.

Pounsford 2001 {published data only}

Pounsford JC. What are the relative places of ipratropium, salmeterol and montelukast in maintenance therapy asthma in the elderly? [ongoing trial]. National Research Register (UK) 2001. [N0234093764]

Rafferty 1987 {published data only}

Rafferty P, Howarth P, Mann JS, Holgate S. Bronchoconstriction induced by nebulized ipratropium bromide: relation to the bromide ion. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):12.

Rebuck 1979 {published data only}

Rebuck AS, Marcus HI. SCH 1000 in psychogenic asthma. *Scandinavian Journal of Respiratory Diseases* 1979;**103**(Suppl):186.

Riding 1970 {published data only}

Riding WD, Dinda P, Chatterjee SS. The bronchodilator and cardiac effects of five pressure-packed aerosols in asthma. *British Journal of Diseases of the Chest* 1970;**64**:37.

Ruffin 1977 {published data only}

Ruffin RE, Fitzgerald JD, Rebuck AS. A comparison of the bronchodilator activity of Sch 1000 and salbutamol. *Journal of Allergy and Clinical Immunology* 1977;**59**:136.

Ruffin 1978 {published data only}

Ruffin RE, Newhouse MT. Ipratropium bromide (Sch 1000) monohydrate aerosol: bronchodilator effect of three dose levels in asthmatics. *Lung* 1978;**155**:141.

Ruffin 1982 {published data only}

Ruffin RE, McIntyre E, Crockett AJ, Zielonka K, Alpers JH. Combination bronchodilator therapy in asthma. *Journal of Allergy and Clinical Immunology* 1982;**69**:60.

Ruffin 1987b {published data only}

Ruffin RE, Meki M, Alpers JH. Combined bronchodilator protection against histamine-induced bronchoconstriction. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):10.

Rutkowski 1994 {published data only}

Rutkowski R, Kazberuk M, Siergiejko Z, Chyrek-Borowska S. Effect of fenoterol, ipratropium bromide and combination drug-berodual-on selected clinical parameters and lung function. *Pneumonologia i Alergologia Polska* 1994;**62**:358.

Rutten-van-Molken 19 {published data only}

Rutten-van-Molken MP, van-Doorslaer EK, Jansen MC, Kerstjens HA, Rutten FF. Costs and effects of inhaled

corticosteroids and bronchodilators in asthma and chronic obstructive pulmonary disease. *American Journal of Respiratory and Critical Care Medicine* 1995;**151**:975.

Sahay 1980 {published data only}

Sahay JN, Bell R, Barber PV, Chatterjee SS, Cox GA. Comparison of ipratropium bromide and salbutamol therapy by inhalation in asthma. *British Journal of Diseases of the Chest* 1980;**75**:323.

Sahlstrom 1986 {published data only}

Sahlstrom K, Alanko K, Harkonen R. Treatment of asthma bronchiale with a combination of ipratropium bromide and fenoterol. *Respiration* 1986;**50**(Suppl 2):302.

Salome 1988 {published data only}

Salome CM, Wright W, Sedgwick CJ, Woolcock AJ. Acute effects of fenoterol (Berotec) and ipratropium bromide (Atrovent) alone and in combination on bronchial hyperresponsiveness in asthmatic. *Progress in Clinical & Biological Research* 1988;**263**:405.

Schindl 1973 {published data only}

Schindl R. [Atropine derivatives as dose aerosol in inhalation therapy of bronchial asthma]. *Medizinische Welt* 1973;**24**:603.

Schlueter 1978 {published data only}

Schlueter DP, Neumann JL. Double blind comparison of acute bronchial and ventilation-perfusion changes to atrovent and isoproterenol. *Chest* 1978;**73**(Suppl 6):982.

Schroeckenstein 1988 {published data only}

Schroeckenstein DC, Bush RK, Chervinsky P, Busse WW. Twelve-hour bronchodilation in asthma with a single aerosol dose of the anticholinergic compound glycopyrrolate. *Journal of Allergy and Clinical Immunology* 1988;**82**:115.

Schultze-Werninghaus {published data only}

Schultze-Werninghaus G. Anticholinergic versus beta 2-adrenergic therapy in allergic airways obstruction: double-blind trials on bronchodilator effect and antiallergic. *Respiration* 1981;**41**:239.

Shneerson 1986 {published data only}

Shneerson JM. Comparison of the bronchodilation of ipratropium bromide and fenoterol nebuliser solutions administered alone and in combination. *Respiration* 1986;**50**(Suppl 2):274.

Simonsson 1979 {published data only}

Simonsson BG. Acute effects of ipratropium bromide (ITBR, Sch 1000, Atrovent): a review of previous studies. *Scandinavian Journal of Respiratory Diseases* 1979;**103**(Suppl):130.

Smith 1988 {published data only}

Smith CM, Anderson SD, Seale JP. The duration of action of the combination of fenoterol hydrobromide and ipratropium bromide in protecting against asthma provoked by hyperpnea. *Chest* 1988;**94**:709.

Snow 1979 {published data only}

Snow RM, Miller WC, Blair HT, Rice DL. Inhaled atropine in asthma. *Annals of Allergy* 1979;**42**:286.

Spector 1975 {published data only}

Spector S, Ball RE. Bronchodilating effects of aerosolized Sch 1000 and atropine sulfate in asthmatics. *Chest* 1975;**68**:426.

Speirs 1987 {published data only}

Speirs D, Hill L, Kirkbride P. The dose response study to compare Atrovent metered dose inhalers with and without an extension tube. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):6.

Stewart 1987 {published data only}

Stewart DE, Gillies AJ. A dose response study comparing Duovent vs salbutamol. *New Zealand Medical Journal* 1987;**100**:742.

Storms 1975a {published data only}

Storms WW, DoPico G, Reed C. Aerosol Sch 1000. *The American Review of Respiratory Disease* 1975;**111**:419.

Storms 1975b {published data only}

Storms WW, Reed CE. Dose-titration study of Sch 1000 MDI in comparison with isoprenaline. *Postgraduate Medical Journal* 1975;**51**(Suppl 7):99.

Stresemann 1975 {published data only}

Stresemann E. Time-response curves of Sch 1000 and isoprenaline as metered dose inhalers (MDI) as assessed by measurements of total airways resistance (Rt). *Postgraduate Medical Journal* 1975;**51**(Suppl 7):119.

Stresemann 1976a {published data only}

Stresemann E. Time response determinations from analysis of pulmonary function following ipratropiumbromide and isoprenaline. *Arzneimittel Forschung* 1976;**26**(5A):1015.

Stresemann 1976b {published data only}

Stresemann-E. Short Communication: Rotation-viscosimetric determinations of the effect of ipratropiumbromide on bronchial excretion/a double blind study. *Arzneimittel Forschung* 1976;**26**(5A):1013.

Sur 1990 {published data only}

Sur S, Mohiuddin AA, Vichyanond P, Nelson HS. A random double-blind trial of the combination of nebulized atropine methylnitrate and albuterol in nocturnal asthma. *Annals of Allergy* 1990;**65**:384.

Tang 1984 {published data only}

Tang OT, Flatley M. A comparison of effects of inhaling a combined preparation of fenoterol with ipratropium bromide (Duovent) with those of fenoterol and. *Postgraduate Medical Journal* 1984;**60**(Suppl 1):24.

Tarlo 1982 {published data only}

Tarlo SM, Broder I, Corey P. Six-month double-blind cross-over trial comparing an inhaled anticholinergic agent and placebo in

the treatment of asthma. *Current Therapeutic Research, Clinical and Experimental*. 1982;**32**:265.

Taube 2002 {published data only}

Taube C, Kannies F, Gronke L, Richter K, Mucke M, Paasch K, et al. Reproducibility of forced inspiratory and expiratory volumes after bronchodilation in patients with COPD or asthma. *Respiratory Medicine* 2002;**97**:568.

Thiessen 1979 {published data only}

Thiessen B, Pedersen OF. A double blind cross-over study of maximal expiratory flows and arterial blood gas tensions in normals, asthmatics and bronchitics after. *Scandinavian Journal of Respiratory Diseases* 1979;**103**(Suppl):170.

Trinquet 1975 {published data only}

Trinquet G, Duchier J, Jacquot C. The bronchodilating effect of Sch 1000 MDI and inhalations of atropine sulphate in patients suffering from bronchial asthma or from chronic. *Postgraduate Medical Journal* 1975;**51**(Suppl):119.

Tukiainen 1985 {published data only}

Tukiainen P, Salorinne Y. Oxitropium, salbutamol and their combination in asthma. *European Journal of Respiratory Diseases* 1985;**67**:31.

Uffholtz 1975 {published data only}

Uffholtz H. The effects of Sch 1000 MDI on specific airways resistance in comparison with salbutamol in patients with asthma and chronic bronchitis. *Postgraduate Medical Journal* 1975;**51**(Suppl 7):120.

Ullah 1980 {published data only}

Ullah M, Saunders KB. Influence of age on response to ipratropium bromide and salbutamol in asthmatic patients. *Progress in Respiratory Research* 1980;**14**:150.

Ullah 1981 {published data only}

Ullah MI, Newman GB, Saunders KB. Influence of age on response to ipratropium and salbutamol in asthma. *Thorax* 1981;**36**:523.

Ulmer 1979 {published data only}

Ulmer WT. Treatment of obstructive disease with fenoterol-ipratropiumbromid dosage aerosol (IK 6). Short and long-terms results [Die Behandlung obstruktiver Atemwegserkrankungen mit fenoterol-ipratropiumbromidgemisch-Dosieraerosol (IK6)]. *Medizinische Klinik* 1979;**74**(42):1548-52.

Vakil 1985 {published data only}

Vakil DV, Ayiomamitis A, Nizami RM. Use of ipratropium aerosol in the long-term management of asthma. *Journal of Asthma* 1985;**22**:165.

Verstraeten 1974a {published data only}

Verstraeten JM. [Can the association of an atropinic (SCH 1000) and a sympathicomimetic improve the bronchodilatation obtained by a sympathicomimetic alone]. *Acta tuberculosea et pneumologica Belgica* 1974;**65**:395.

Verstraeten 1974b {published data only}

Verstraeten JM. [Comparative study of the bronchodilating effect of SCH 1000, berotec and of the association of the two drugs]. *Acta tuberculosa et pneumologica Belgica* 1974;**65**:534.

Verstraeten 1975 {published data only}

Verstraeten JM. Bronchial response of asthmatic and bronchitic patients to fenoterol MI with or without subsequent Sch 1000 MDI. *Postgraduate Medical Journal* 1975;**51**(Suppl 7):120.

Villate 1980 {published data only}

Villate Navarro JI, Sobradillo Pena V, Atxotequi Iaraoligoitia V, Salaverri Nalda A, Orive Martinez C. [Comparative effects of terbutaline sulphate and ipratropium bromide on the respiratory system]. *Medicine Clinique et Experimentale* 1980;**74**:275.

Vlagopoulos 1975 {published data only}

Vlagopoulos T, Townley R, Ghazanshahi S, Bewtra A, Burke K. Comparison of the onset of action and bronchodilation effects of the anticholinergic agent SCH 1000 with isoproterenol. *Journal of Allergy and Clinical Immunology* 1975;**55**:99.

Vlagopoulos 1976 {published data only}

Vlagopoulos T, Townley R, Ghazanshahi S, Bewtra A, Burke K. Comparison of the bronchodilating effects of Sch 1000 with isoproterenol in patients with bronchial asthma. *Annals of Allergy* 1976;**36**:223.

Vogt 1974 {published data only}

Vogt R, Rychel F. Acute test comparing ipratropium bromide (Sch 1000), fenoterol and placebo, by measured dose aerosol, in obstructive lung disease. *Therapiewoche* 1974;**24**:4228.

Walker 1987 {published data only}

Walker FB, Kaiser DL, Kowal MB, Suratt PM. Prolonged effect of inhaled glycopyrrolate in asthma. *Chest* 1987;**91**:49.

Ward 1987 {published data only}

Ward MJ, MacFarlane JT, Davies D. Anticholinergics in severe asthma. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):2.

Watson 1992 {published data only}

Watson A, Lim TK, Joyce H, Pride NB. Failure of inhaled corticosteroids to modify bronchoconstrictor or bronchodilator responsiveness in middle-aged smokers with mild airflow. *Chest* 1992;**101**:350.

Wegener 1987a {published data only}

Wegener T, Hedenstrom H. Effect of nebulized ipratropium bromide on lung function in nonallergic bronchial asthma. *Respiration* 1987;**52**:1.

Wegener 1987b {published data only}

Wegener T, Hedenstrom R. Effect of nebulized ipratropium bromide on lung function in non-allergic bronchial asthma. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):6.

Wempe 1992 {published data only}

Wempe JB, Postma DS, Breederveld N, Alting-Hebing D, van der Mark TW, Koeter GH. Separate and combined effects of

corticosteroids and bronchodilators on airflow obstruction and airway hyperresponsiveness in asthma. *Journal of Allergy and Clinical Immunology* 1992;**89**:679.

Wesolowski 1985 {published data only}

Wesolowski S. [The role of anticholinergic drugs in the treatment of bronchospastic conditions]. *Pneumonologia Polska* 1985;**53**:49.

Wildbolz 1977 {published data only}

Wildbolz U, Kyd K, Scherrer M. Ipratropium in addition to theophylline-sustained betastimulation. *Lung* 1977;**154**:141.

Wilson 1978 {published data only}

Wilson RH, Battaglia PJ, Wilson NL. Crossover study with nebulized bronchodilators and atropine. *Chest* 1978;**73**(Suppl 6):998.

Windom 1990 {published data only}

Windom HH, Burgess CD, Siebers RW, Purdie G, Pearce N, Crane J, Beasley R. The pulmonary and extrapulmonary effects of inhaled beta-agonists in patients with asthma. *Clinical Pharmacology and Therapeutics* 1990;**48**:296.

Yanai 1991 {published data only}

Yanai M, Ohru T, Sekizawa K, Shimizu Y, Sasaki H, Takishima T. Effective site of bronchodilation by antiasthma drugs in subjects with asthma. *Journal of Allergy and Clinical Immunology* 1991;**87**:1080.

Yang 1989 {published data only}

Yang SC, Wu MC. The response pattern of patients with chronic airway obstruction to bronchodilators. *Journal of the Formosan Medical Association of Taiwan* 1989;**88**:700.

Yeager 1976 {published data only}

Yeager H, Weinberg RM, Kaufman LV, Katz S. Asthma: comparative bronchodilator effects of ipratropium bromide and isoproterenol. *Journal of Clinical Pharmacology* 1976;**16**:198.

Zanen 1995 {published data only}

Zanen P, Go LT, Lammers JW. The optimal particle size for parasympatholytic aerosols in mild asthmatics. *International Journal of Clinical Pharmacology and Therapeutics* 1995;**114**:111.

Zaritsky 1999 {published data only}

Zaritsky A, Qureshi F. Ipratropium does indeed reduce admissions to hospital with severe asthma. *BMJ* 1999;**318**:738.

Zuck 1987 {published data only}

Zuck P, Peiffert G, Jirou-Najou JL. Study of fenoterol and ipratropium bromide additive action in asthmatic patients. *Postgraduate Medical Journal* 1987;**63**(Suppl 1):14.

Additional references
Barnes 1996

Barnes PJ. Pathophysiology of asthma. *British Journal of Clinical Pharmacology* 1996;**42**:3-10.

Barnes 1997

Barnes PJ. Current therapies for asthma. *Chest* 1997;**111**(Suppl.):17S-26S.

Beakes 1997

Beakes DE. The use of anticholinergics in asthma. *Journal of Asthma* 1997;**34**:357-68.

Boulet 1999

Boulet L-P, Becker A, Berube D, Beveridge R, Ernst P. Summary of recommendations from the Canadian Asthma Consensus Report 1999. *Canadian Medical Association Journal* 1999;**161**(11 Suppl):S1-S12.

Bousquet 2000

Bousquet J. Global initiative for asthma (GINA) and its objectives. *Clinical & Experimental Allergy* 2000;**30**(Suppl.1):2-5.

BTS 1993

British Thoracic Society. Guidelines on the management of asthma. *Thorax* 1993;**48**(Suppl.):s1-s24.

BTS/SIGN 2004

Scottish Intercollegiate Guidelines Network. Clinical Guidelines. <http://www.sign.ac.uk> (accessed February 2004).

Cazzola 1998

Cazzola M, Centanni S, Donner CF. Anticholinergic agents. *Pulmonary Pharmacology & Therapeutics* 1998;**11**:381-92.

Clarke 2003

Clarke M, Oxman AD (editors). *Cochrane Reviewers' Handbook* 4.2.0 [updated March 2003]. Cochrane Collaboration, 2003.

Connolly 1993

Connolly MJ. Ageing, late-onset asthma and the beta-adrenoceptor. *Pharmacology & Therapeutics* 1993;**60**:389-404.

Creer 1999

Creer TL, Winder JA, Tinkelman D. Guidelines for the diagnosis and management of asthma: accepting the challenge. *Journal of Asthma* 1999;**36**:391-407.

Elbourne 2002

Elbourne DR, Altman DG, Higgins JPT, Curtin F, Worthington HV, Vail A. Meta-analyses involving cross-over trials: methodological issues. *International Journal of Epidemiology* 2002;**31**:140-9.

Emilien 1998

Emilien G, Maloteaux J-M. Current therapeutic uses and potential of beta-adrenoceptor antagonists. *European Journal of Clinical Pharmacology* 1998;**53**:389-404.

Gross 1988

Gross NJ. Ipratropium bromide. *New England Journal of Medicine* 1988;**319**:486-94.

Gross 1997

Gross NJ. Anticholinergic drugs. In: Barnes PJ, Grunstein MM editor(s). *Asthma*. Philadelphia: Lippincott-Raven, 1997:1555-68.

Harrison 1998

Harrison B. Towards evidence-based guidelines for asthma: should we? can we? how will it affect our practice?. *Clinical Asthma Reviews* 1998;**2**:139-46.

Hougardy 2000

Hougardy DM, Peterson GM, Bleasel MD, Randall CT. Is enough attention being given to the adverse effects of corticosteroid therapy. *Journal of Clinical Pharmacology & Therapeutics* 2000;**25**:227-34.

Jadad 1996

Jadad AR, Moore RA, Carroll D, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary?. *Controlled Clinical Trials* 1996;**17**:1-12.

Jolobe 1984

Jolobe OM. Asthma vs. non-specific reversible airflow obstruction: clinical features and responsiveness to anticholinergic drugs. *Respiration* 1984;**45**:237-42.

Kemp 2000

Kemp JP. Guidelines Update. *Drugs* 2000;**59**(Suppl.1):23-28.

Meier 1997

Meier CR, Hershel J. Drug use and pulmonary death rates in increasingly symptomatic asthma patients in the UK. *Thorax* 1997;**52**:612-7.

Ni Chroinin 2005

Ni Chroinin M, Greenstone IR, Danish A, Magdolinos H, Masse V, Zhang X, Ducharme FM. Long-acting beta2-agonists versus placebo in addition to inhaled corticosteroids in children and adults with chronic asthma. *Cochrane Database of Systematic Reviews* 2005, Issue 4. [DOI: [10.1002/14651858.CD005535](https://doi.org/10.1002/14651858.CD005535)]

O'Connor 1998

O'Connor BJ. Combination therapy. *Pulmonary Pharmacology & Therapeutics* 1998;**11**:397-9.

Partridge 1981

Partridge MR, Saunders KB. Site of action of ipratropium bromide and clinical and physiological determinants of response in patients with asthma. *Thorax* 1981;**36**:530-3.

Spitzer 1992

Spitzer WO, Suissa S, Ernst P, Horwitz R, Habbick B, Cockcroft D, et al. The use of beta-agonists and the risk of death and near death from asthma. *New England Journal of Medicine* 1992;**326**:501-6.

Taylor 1999

Taylor DM, Auble TE, Calhoun WJ, Mosesso VN. Current outpatient management of asthma shows poor compliance with international consensus guidelines. *Chest* 1999;**116**:1638-45.

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Ahonen 1978

Methods	Crossover design; each phase 4 days. All patients completed the study correctly. Said to be double blind, but no details. No other concurrent asthma medication.
Participants	Inclusion criteria: patients had asthma - diagnosis according to Ciba Guest symposium criteria; stable asthma medication. Patient details: 13 participants (M/F 7/6); mean age 37; mild asthma (severity homogeneous); 5/8 atopy; no information on concurrent COPD.
Interventions	5 interventions: ipratropium bromide 40 and 80µg, fenoterol 400µg, placebo (twice) by pressurised inhaler; all interventions 5 times per day (01.00, 07.00, 13.00, 19.00, 22.00h).
Outcomes	Daily PEF (measured before dose and at 10am; 7am and 10pm (pre-bronchodilator) and 10am (3h post-bronchodilator) data used in analysis. Daily symptom scores (day and night). Adverse effects.
Notes	Placebo is the mean of 2 periods. The graph recording PEFR did not agree with the table for the SEM - likely that table incorrect (see text).

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Ashutosh 1980

Methods	Crossover design, each phase 7 days.
Participants	
Interventions	atropine sulphate and isoprenaline.
Outcomes	
Notes	This comparison not analysed, so details not extracted.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Bellia 1988

Methods	Crossover design; each phase 7 days. All patients completed the study correctly. Said to be double blind, with placebos used, but no other details. Not double blind for dose comparison. Concurrent asthma medication - sodium cromoglycate.
---------	---

Bellia 1988 (Continued)

Participants	Inclusion criteria: patients had asthma - diagnosis according to ACCP-ATS criteria; stable asthma medication; recent nocturnal asthma. Exclusions: CV, ocular & genito-urinary diseases; history of intolerance to theophylline or anti-muscarinic agents. Patient details: 12 participants; age 18-60; mild asthma; no information on concurrent COPD or atopy.
Interventions	2 interventions: oxitropium bromide 400/600µg, placebo by MDI; all interventions t.i.d. (unclear when).
Outcomes	% change daily PEF(morning/ max PEF; not analysed), symptom scores (night or morning - no numbers); % predicted FEV1 at end of period (not analysed).
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Chapman 1983

Methods	See Rebuck 1983
Participants	
Interventions	
Outcomes	
Notes	Same trial as Rebuck 1983

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Coe 1986

Methods	Crossover design; each phase 14 days. All patients completed the study correctly. Said to be double blind (vs placebo), but no details. Concurrent asthma medication unclear, but none had oral steroids or anticholinergics; 200µg salbutamol given each morning and evening as part of test procedure.
Participants	Inclusion criteria: patients had documented nocturnal asthma - difference of >20% in PEFR between evening and morning readings; nocturnal asthma. Patient details: 18 participants (M/F 7/11); age 18-76; mild-severe asthma (heterogeneous severity); 10:8 atopy; no information on concurrent COPD.
Interventions	3 interventions: oxitropium bromide 200 and 400µg, placebo ; all interventions taken once at night.

Coe 1986 (Continued)

Outcomes	Daily PEF, morning and evening (no results or SDs); difference in PEFR (evening-morning) - not analysed; rescue medication - no results; daily symptom scores - no data.	
Notes	200µg salbutamol given each morning and evening as part of test procedure.	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Dompeling 1992

Methods	See van Schayck 1991a	
Participants		
Interventions		
Outcomes		
Notes	This comparison not analysed, so details not extracted	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Dompeling 1993

Methods	See van Schayck 1991a	
Participants		
Interventions		
Outcomes		
Notes	This comparison not analysed, so details not extracted.	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Hahtela 1991

Methods	Parallel design; duration 12 weeks. Overall, 24/120 patients withdrew from study. Said to be double blind, but no details. Concurrent asthma medication: inhaled steroids, cromoglycate, methylxanthines; none had oral steroids; inhaled bronchodilators withdrawn.
Participants	Inclusion criteria: patients had asthma - reversible (>15% increase in FEV1) after beta-agonist. Exclusions: patients taking oral steroids. Patient details: 120 participants, but 96 completed - patient details for 96: M/F 35/61; age 46 (mean); moderate asthma (homogeneous severity); 56:40 atopy; no information on concurrent COPD (but all non-smokers).
Interventions	4 interventions: Berodual, Duovent, Fenoterol 200 and 400µg; all interventions q.i.d. (07.00 and 19.00h and two other unspecified times).
Outcomes	PEF (daily) for 1-4weeks and 8-12w; symptom scores; extra inhaler use (not number of patients); heart rate, side effects score; number of patient withdrawals; no standard deviations/p values for continuous outcomes.
Notes	Large baseline differences for symptom scores.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Kreisman 1981

Methods	Crossover design; each phase 14 days. 3/15 patients withdrew from study. Said to be double blind (vs placebo), but no other details. Concurrent asthma medication: theophylline (study drug), maintenance beta-agonists and steroids kept constant.
Participants	Inclusion criteria: patients had asthma - diagnosis according to ATS criteria; stable asthma medication. Exclusions: no history of chronic bronchitis, cardiac, renal or hepatic disease. Patient details: 15 participants (M/F 7/5); age 21-54 (mean 34); mild-moderate asthma; atopy not stated. No concurrent COPD.
Interventions	4 interventions: ipratropium bromide 40µg and placebo, each with/without theophylline; all interventions q.i.d. (unclear when).
Outcomes	FEV1 at 1 and 2 weeks; symptom scores (wheeze breathlessness, chest tightness, - no numbers); side effects only in words.
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Laitinen 1979

Methods	Crossover design; 1st phase 3 weeks, 2nd phase 2 weeks.
---------	---

Laitinen 1979 (Continued)

Participants

Interventions Ipratropium bromide 80µg and salbutamol 400µg; both interventions q.i.d.

Outcomes

Notes This comparison not analysed, so details not extracted.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Light 1977

Methods Crossover design; each phase 2 weeks. All patients completed the study correctly. Said to be double blind (vs placebo), but no other details. Concurrent asthma medication various, some had oral steroids.

Participants Inclusion criteria: patients had asthma - reversible after isoprenaline and atropine; stable asthma medication; continuous symptoms for 3 months. Patient details: 6 participants; age 28-42 (mean 35); moderate asthma (heterogeneous severity).

Interventions 2 interventions: oral atropine sulphate 5mg and placebo; both interventions q.i.d.

Outcomes FEV1, side effects.

Notes Oral atropine - not included in analysis.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	High risk	Investigators not blinded to randomisation sequence (Cochrane Grade C)

Lightbody 1977

Methods See [Lightbody 1978](#)

Participants

Interventions

Outcomes

Notes This comparison not analysed, so details not extracted.

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Anticholinergic agents for chronic asthma in adults (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Lightbody 1977 (Continued)

Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)
--	--------------	--

Lightbody 1978

Methods	Crossover design; each phase 3 days.
Participants	
Interventions	2 interventions: Ipratropium bromide 40µg and Salbutamol 200µg, q.i.d.
Outcomes	
Notes	This comparison not analysed, so details not extracted.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Macaluso 1986

Methods	Crossover design; each phase 2 weeks. All patients completed the study. Said to be double blind, but could not be because of different number of doses (Duovent 3; Salbutamol 4 per day). Concurrent asthma medication: all patients received aminophylline (long acting), no further details; no information on withdrawal of bronchodilators.
Participants	Inclusion criteria: patients had "partially reversible chronic bronchospasm". Patient details: 15 participants (M/F 7/8); age 32-65; no information on severity of asthma, atopy or concurrent COPD.
Interventions	2 interventions: Duovent t.i.d. (times not stated), Salbutamol 200µg q.i.d (times not stated).
Outcomes	Daily symptom scores (but unclear what is reported; standard deviation calculated from graph), use of extra aerosol.
Notes	Unblinded study - different number of inhalations; unclear reporting of results.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Mazzei 1985

Methods	Crossover design; each phase 2 weeks. All patients completed the study. Said to be 'blind', but no details. Concurrent asthma medication: steroids (unspecified), methylxanthines; usual bronchodilators withdrawn.
Participants	Inclusion criteria: patients had asthma - diagnosis according to ATS criteria; exclusions: serious cardiac pathology, arrhythmias, hepatic/renal failure, COPD, other pulmonary diseases, neoplasias, pregnancy. Patient details: 20 participants (M/F 7/13); age 16-72 (mean 46); moderate asthma (heterogeneous severity); atopy not stated; no concurrent COPD.
Interventions	2 interventions: Berodual, Salbutamol 200µg; all interventions t.i.d. (every 8h, unclear when).
Outcomes	FEV1, %FEV1/FVC, FVC, symptom scores at 1 and 2 weeks; side effects; patient & physician assessment.
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

O'Driscoll 1990

Methods	Crossover design; each phase 1 month.
Participants	
Interventions	2 interventions: Ipratropium bromide 500µg (nebuliser) and salbutamol 5mg (neb); q.i.d.
Outcomes	
Notes	This comparison not analysed, so details not extracted. Different trial to O'Driscoll 1992.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

O'Driscoll 1992

Methods	Crossover design; each phase 1 month.
Participants	
Interventions	Ipratropium bromide 500µg (nebuliser) and salbutamol 5mg (neb); q.i.d.
Outcomes	

O'Driscoll 1992 (Continued)

Notes This comparison not analysed, so details not extracted. Different trial to O'Driscoll 1990.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Pavia 1987

Methods As [Pavia 1989](#)

Participants

Interventions

Outcomes

Notes Same trial as [Pavia 1989](#)

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Pavia 1989

Methods Crossover design; each phase 28 days. 1/11 asthma patients withdrew from study. Said to be double blind (vs placebo), but some patients could distinguish the OxBr by its taste, no other details. Concurrent asthma medication: no oral steroids; inhaled steroids and oral bronchodilators.

Participants Inclusion criteria: patients had asthma - reversible (>15% increase in FEV1) after beta-agonist; stable asthma medication. Patient details: 22 participants (11 with asthma(M/F for 10: 4/6), 11 with bronchitis - not separated for most outcomes); age (asthmatics) mean 59; moderate asthma; atopy not stated.

Interventions 2 interventions: oxitropium bromide 200µg and placebo; both interventions t.i.d. (on rising, early pm and on retiring).

Outcomes PEF (am, early pm, retiring - paired difference +SEM for am); FEV1 at 4 weeks; rescue medication (not for placebo) - all these for mixed asthma/COPD; number of dropouts (asthma only).

Notes Most outcomes for asthma and COPD combined; 1 author worked for Boehringer Ingelheim.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Anticholinergic agents for chronic asthma in adults (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Petersen 1979

Methods	Crossover design; each phase 3 days
Participants	
Interventions	2 interventions: ipratropium bromide 125µg (nebuliser) and terbutaline 5mg (neb); q.i.d.
Outcomes	
Notes	This comparison not analysed, so details not extracted.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Philip-Joet 1990

Methods	Parallel design; duration 8 weeks. Overall, 48/196 patients withdrew from study, but 77/196 were not analysed at 8w. "A double blind design was not used"; not stated if outcome assessor blinded. Concurrent asthma medication: steroids (unspecified), methylxanthines; inhaled bronchodilators withdrawn.
Participants	Inclusion criteria: patients had asthma - diagnosis according to ATS criteria and >15% reversibility after beta-agonist; exclusions: pregnancy, respiratory failure, diabetes, contraindications to study drugs. Patient details: 196 participants (M/F group B 56% M; group S 57% M); age: Group B 45 (mean), Group S 48; moderate asthma (fairly heterogeneous severity); atopy not stated; no information on concurrent COPD.
Interventions	2 interventions: Berodual, Salbutamol 200µg; both interventions q.i.d. (times not stated).
Outcomes	PEF (daily) over 7 days (1,2,3,4,5,6,7,8w), symptom scores (not daily), FEV1, PEFR, FVC at 30 and 60 days; adverse effects (but too much data missing); standard deviations.
Notes	Many withdrawals and missing patient data, therefore only used week 1 data for PEFR; concealment: numbered boxes, serially allocated.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	Investigators blinded to randomisation sequence (Cochrane Grade A)

Pierce 1982

Methods	Crossover design; each phase 4 weeks. 1/10 patients not analysed because of changes in medication. Said to be double blind, with placebo inhalers, but no other details. Concurrent asthma medication: theophylline & steroids (oral/inhaled); withdrawal of inhaled bronchodilators unclear.
---------	---

Pierce 1982 (Continued)

Participants	Inclusion criteria: patients had asthma - reversibility >25% after beta-agonist. Patient details: 10 participants (all male); age 52-67 (mean 59); moderate asthma; 7/10 atopy; 8/10 concurrent COPD.
Interventions	3 interventions: ipratropium bromide (40µg)+terbutaline (500µg), terbutaline (500µg), ipratropium bromide (40µg); all interventions q.i.d. (times not clear).
Outcomes	Flow - not PEF - (daily) over 4weeks; symptom scores (no results) over 4 weeks; FEV1 at 2 & 4 weeks (mean); adverse effects.
Notes	Partial allocation concealment - opaque sealed envelopes opened after patient enrollment (not serially numbered).

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Rebuck 1983

Methods	Crossover design; each phase 30 days. 4/22 patients withdrawn and 17 patients' results were analysed. Said to be double blind, with placebos and the code was not broken until the study was complete. Concurrent asthma medication: steroids (oral and inhaled), cromoglycate, theophylline (study drug in all arms); withdrawal of bronchodilators unclear.
Participants	Inclusion criteria: patients had asthma - diagnosis according to ATS criteria and reversibility >15% after beta-agonist; all patients clinically stable for >1 month. Exclusions: cough and sputum during clinically stable periods, recent respiratory tract infections, cardiac, hepatic, renal, metabolic disease, and others; hypersensitivity to atropinic compounds. Patient details: 22 participants (M/F 9/13); age 17-75 (mean for completers 47); moderate asthma (heterogeneous severity); 13/22 atopy; unclear if concurrent COPD.
Interventions	3 interventions: ipratropium bromide (40µg)+fenoterol (200µg) - may be separate inhalers, fenoterol (200µg), ipratropium bromide (40µg); all interventions q.i.d. (times not clear) and all arms had oral theophylline (200mg) q.i.d.
Outcomes	PEF (daily) over 30d (no results); FEV1 at 30days; patient preference and number of exacerbations (descriptive); adverse effects.
Notes	Authors unable to provide any more data.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Ruffin 1990

Methods	Crossover design; phase length unclear - could be 8 weeks. 5/21 patients withdrew from study. Said to be double blind (vs placebo), but no other details. Concurrent medication: not stated.
---------	--

Anticholinergic agents for chronic asthma in adults (Review)

Ruffin 1990 (Continued)

Participants	Inclusion criteria: patients had asthma - diagnosis reasons not stated. Patient details: 16 remaining participants (M/F 9/7); mean age 56; mild asthma; no information about atopy or concurrent COPD.
Interventions	2 interventions: oxitropium bromide 200µg and placebo; both interventions t.i.d. (not stated when).
Outcomes	PEF (am and pm, probably at end of period); FEV1 at end of period.
Notes	Abstract; funded by Boehringer Ingelheim.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Salorinne 1988

Methods	Crossover design; each phase 7 days. All patients completed the study correctly. Not blinded for the comparison IB (40) vs placebo (powder); no details re outcome assessor. Concurrent medication: 5/18 had oral steroids; 17 had theophylline.
Participants	Inclusion criteria: patients had asthma - improvement in FEV1 of 16-100% (not stated how). Patient details: 18 participants (M/F 10/8); age 22-70 (mean 48); moderate-severe asthma; atopy 5/18, no information on concurrent COPD.
Interventions	4 interventions: ipratropium bromide 40 and 200µg (aerosol) and 200µg powder, and placebo powder ; all interventions q.i.d (08.00, 12.00, 18.00, 22.00h).
Outcomes	PEF (8, 10am, midday, 2, 6, 8, 10pm & midnight); Symptom scores (dyspnoea day and night, cough); rescue medication (words); patient preference.
Notes	Placebo was powder.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Storms 1986

Methods	Parallel design; 90 days.
Participants	
Interventions	2 interventions: ipratropium bromide 40µg and metaproterenol 1500µg; q i.d
Outcomes	
Notes	This comparison not analysed, so details not extracted.

Anticholinergic agents for chronic asthma in adults (Review)

Storms 1986 (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Tammivaara 1993

Methods	Crossover design; each phase 3 weeks. 7/28 patients withdrawn. Said to be double blind and double dummy, but no other details. Concurrent medication: theophylline, none had steroids; patients allowed to use inhaled bronchodilators when needed (NB study drug oral).	
Participants	Inclusion criteria: patients had asthma - nocturnal decline in PEF of at least 20%. Patient details: 28 participants (M/F 10/18); age 18-59 (mean 39); moderate asthma (homogeneous severity); no information on atopy or concurrent COPD.	
Interventions	3 interventions: ipratropium bromide 40µg+oral controlled-release terbutaline (10mg), oral CR terbutaline (10mg)+aerosol placebo and 200µg powder, and ipratropium bromide 40µg+oral placebo; aerosols q.i.d (08.00, 12.00, 16.00, 20.00h); oral tablets b.i.d. (08.00 and 20.00h).	
Outcomes	PEF (daily), symptom scores over last 2 weeks of treatment period (standard deviations); number of patients with exacerbations, number needing rescue medications (words), patient preference, adverse effects.	
Notes	Oral CR terbutaline.	

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Taylor 1986

Methods	Crossover design; each phase 4 weeks. 8/23 (of all patients) failed to complete the study. Double blind (identical inhalers), but no other details. Concurrent asthma medication: mixed asthma/COPD study - medication not distinguished for asthma (3/all patients had oral steroids, 7 had aminophylline).	
Participants	Inclusion criteria: patients had asthma - reversibility of >20% with conventional bronchodilators; stable asthma medication implied. Patient details: 23 participants (asthma 7, COPD 8, no details 8) (overall M/F 12/3); asthma & COPD age 27-71 (median 60); mild-severe asthma (severity heterogeneous); no information on concurrent COPD.	
Interventions	2 interventions: ipratropium bromide 200µg and placebo; all interventions t.i.d. (not stated when).	
Outcomes	Daily symptom scores (breathlessness, cough, wheeze), PEF and FEV1 at end of period, number of patients with exacerbations (all mixed asthma/COPD); treatment related withdrawals (mixed asthma/COPD)	
Notes	Mixed asthma/COPD mainly; Boehringer Ingelheim financial support.	

Taylor 1986 (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Taytard 1984

Methods	Crossover design, but analysed as 1st period only; each phase 21 days. All patients completed the study (although there is no patient number 2). Said to be double blind (with identical treatments), but no details about the assessor. Concurrent asthma medication: no oral or inhaled steroids, no regular treatment but 8 had theophylline for relief.	
Participants	Inclusion criteria: patients had asthma - diagnosis according to ATS criteria; presenting an allergic diathesis and consulting an allergology unit; perennial asthma. Patient details: 16 participants (M/F 9/7); age 14-48 (mean 31); mild asthma; all atopic, no information on concurrent COPD.	
Interventions	2 interventions: oxitropium bromide 200µg and placebo; all interventions t.i.d. (08.00, 14.00, 20.00h).	
Outcomes	Clinical scores (IPD); rescue medication; patient rating.	
Notes	Significant order effect, so used 1st period data only from IPD.	

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Tormey 1995

Methods	Crossover design; each phase 28 days. Not clear if intention to treat. Said to be double blind (vs placebo), but no other details. Concurrent asthma medication: inhaled steroids.	
Participants	Inclusion criteria: patients had asthma - no diagnostic evidence; stable asthma medication implied. Patient details: 12 participants; age not stated; moderate asthma; no information on atopy or concurrent COPD.	
Interventions	3 interventions: oxitropium bromide 200µg, salbutamol 200µg and placebo; all interventions t.i.d. (not stated when).	
Outcomes	Daily symptom scores (unspecified, words only); FEV1 after 4 weeks.	
Notes	Abstract	

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Tormey 1995 (Continued)

Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)
--	--------------	--

van Schayck 1991a

Methods	Crossover design; each phase 1 year
Participants	
Interventions	4 interventions: ipratropium bromide 40µg q.i.d., salbutamol 400µg q.i.d., ipratropium bromide on demand, salbutamol on demand
Outcomes	
Notes	This comparison not analysed, so details not extracted.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

van Schayck 1991b

Methods	See van Schayck 1991a
Participants	
Interventions	
Outcomes	
Notes	This comparison not analysed, so details not extracted.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

van Schayck 1994

Methods	See van Schayck 1991a
Participants	
Interventions	

van Schayck 1994 (Continued)

Outcomes

Notes This comparison not analysed, so details not extracted.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

van Schayck 1995

Methods See [van Schayck 1991a](#)

Participants

Interventions

Outcomes

Notes This comparison not analysed, so details not extracted.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Vaughan 1988

Methods Crossover design; each phase 2 weeks. All patients assumed to have completed the study. Double blind study (code in sealed envelope). Concurrent asthma medication: 2 groups of patients - 12 had inhaled & oral steroids; 10 had no steroids.

Participants Inclusion criteria: patients had asthma - diagnosis ATS criteria and >15% reversibility. Patient details: 22 participants; age 44-77; moderate-severe asthma (2 homogeneous groups); no information on atopy or concurrent COPD.

Interventions 2 interventions: atropine methonitrate 2mg and placebo, nebulised; both interventions q.i.d. (unclear when).

Outcomes FEV at 2 weeks; daily symptom scores and PEF not reported in results (but 'no difference').

Notes Authors stated that the allocation schedule was sealed in envelopes until the study was completed.

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Vaughan 1988 (Continued)

Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)
--	--------------	--

Waite 1987

Methods	Crossover design; each phase 12 weeks. No information on patient withdrawal nor on concurrent asthma medication; no information on withdrawal of bronchodilators. Said to be double blind, but no details.
Participants	Inclusion criteria: patients had asthma - reversibility >15% following beta-agonist. Patient details: 12 participants; no further details.
Interventions	2 interventions: Duovent and salbutamol; no further details.
Outcomes	Daily symptom scores; use of inhaled bronchodilator; PEF am (only descriptive).
Notes	Abstract

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Wheeler 1987

Methods	Crossover design; each phase 4 weeks. 8/39 failed to complete the study and 9/39 patients' results were not analysed. Double blind (identical, masked containers). Concurrent asthma medication: none had oral steroids; 26/31 had inhaled steroids; 22 had methylxanthines.
Participants	Inclusion criteria: patients had asthma - diagnosis >15% reversibility (28 had asthma, 2 had COAD). Patient details: 39 participants; age 17-76; mild-severe asthma (severity heterogeneous); no information on concurrent COPD.
Interventions	2 interventions: oxitropium bromide 200µg, and placebo; all interventions b.i.d. (on rising and on retiring).
Outcomes	Daily PEF (am, pm); Symptom scores (day, breathlessness; night, overall); rescue medication (day & night); patient preference; side effects.
Notes	Boehringer Ingelheim study; IPD given but some patients' diary notes were incomplete; partial concealment (randomisation carried out by BI, but list supplied).

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Wolstenholme 1989

Methods	Crossover design; each phase 4 weeks. All patients completed the study; 13/17 patients' results analysed for PEF. Said to be double blind and double dummy, but no other details. Concurrent asthma medication: steroids (oral and inhaled), cromoglycate; all bronchodilators withdrawn.
Participants	Inclusion criteria: patients had asthma - >15% reversibility following beta-agonist; morning dip of >15% for half of the 2 week run-in. Patient details: 17 participants; age 19-39 (median 25); moderate asthma (severity homogeneous); 100% atopy; no information on concurrent COPD (but all were non-smokers).
Interventions	2 interventions: Duovent+placebo and salbutamol 200µg+placebo separate inhalers; both interventions q.i.d. (times not stated).
Outcomes	PEF (daily), daily symptom scores, FEV1, FVC, PEF (at 2 and 4 weeks), nocturnal waking, patient preference, adverse effects (words); standard deviations given.
Notes	Partial allocation concealment - extra communication from the author stated that sealed envelopes were used.

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	Information not available (Cochrane Grade B)

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Abboud 1987	Wrong comparison
Adinoff 1998	Wrong comparison
Advenier 1984	Review
Alanko 1983	Short term study <24h - included in short term review
Allen 1980a	Short term study <24h - included in short term review
Allen 1980b	Short term study <24h - included in short term review
Alliott 1972	Short term study <24h - included in short term review
Altounyan 1964	Short term study <24h - included in short term review
Andersen 1986	Short term study <24h - included in short term review
Anderson 1980	Short term study <24h - included in short term review
Anderson 1981	Short term study <24h - included in short term review
Anderson 1983	Short term study <24h - included in short term review
Aquilina 1986	Not randomised

Anticholinergic agents for chronic asthma in adults (Review)

Study	Reason for exclusion
Ariano 1981	COPD
Baigelman 1977	Short term study <24h - included in short term review
Banos 1983	Short term study <24h - included in short term review
Barber 1975	Short term study <24h - included in short term review
Barber 1977	Short term study <24h - included in short term review
Bauer 1973	Short term study <24h - included in short term review
Beasley 1987	Confounded - patients selected who responded to Ipratropium
Bell 1982	Confounded - additional salbutamol given to both arms on a regular basis
Bellia 1987	Not said to be randomised; no further data from authors
Bernabeu 1987	Wrong comparison
Bezel 1987	Short term study <24h - included in short term review
Bish 1997	Confounded - patients selected who responded to Ipratropium
Bohadana 1980	Short term study <24h - included in short term review
Bonsignore 1986	wrong comparison; patients had ongoing attack
Botts 1985	Short term study <24h - included in short term review
Brady 1979	Short term study <24h - included in short term review
Brand 1992	Long term study - mixed asthma and COPD patients; separate data not available for asthma despite request to authors
Bremner 1992	Short term study <24h - included in short term review
Britton 1988	Short term study <24h - included in short term review
Brom 1985	Not anticholinergic trial
Bruderman 1983	Short term study <24h - included in short term review
Brumby 1980	Long term study - not stated if patients in long term study were randomised; no response from authors
Bryant 1990a	Short term study <24h - included in short term review
Bryant 1990b	Short term study <24h - included in short term review
Bryant 1992	Short term study <24h - included in short term review
Burge 1980	Short term study <24h - included in short term review
Burki 1997	Short term study <24h - included in short term review

Study	Reason for exclusion
Capel 1964	Short term study <24h - included in short term review
Carpentiere 1986	Wrong comparison Duovent vs placebo
Carratu 1983	Short term study <24h - included in short term review
Casali 1979	Short term study <24h - included in short term review
Catterall 1988	Short term study <24h - included in short term review
Cañete 1991	Asthmatic crisis
Chakravarti 1987	Confounded - patients selected who responded to Ipratropium
Chervinsky 1977	Short term study <24h - included in short term review
Chhabra 2002	Short term study <24h - included in short term review
Chick 1977	Short term study <24h - included in short term review
Chuchalin 1985	Short term study <24h - included in short term review
Ciappi 1986	Short term study <24h - included in short term review
Clauzel 1987	Short term study <24h - included in short term review
Cockcroft 1977	Challenge study
Cox 1984	Not randomised and wrong comparison
Crane 1986a	Short term study <24h - included in short term review
Crane 1986b	Short term study <24h - included in short term review
Crane 1987	Short term study <24h - included in short term review
del Buffalo 1982	Short term study <24h - included in short term review
Donkers 1998	Short term study <24h - included in short term review
Donner 1980	Wrong comparator
Dutt 1990	Study conducted in children
Elwood 1982	Short term study <24h - included in short term review
Emirgil 1975	Short term study <24h - included in short term review
Endoh 1998	Not asthma
Filiz 1994	Not randomised
Firstater 1981	Short term study <24h - included in short term review
Fischbacher 1984	Short term study <24h - included in short term review

Study	Reason for exclusion
Flint 1983a	Short term study <24h - included in short term review
Flint 1983b	Short term study <24h - included in short term review
Flint 1984	Short term study <24h - included in short term review
Foster 1983	Short term study <24h - included in short term review
Freeman 1992	Challenge study
Frith 1987	Short term study <24h - included in short term review
Gaba 1987	Short term study <24h - included in short term review
Gillies 1987	Challenge study
Ginesu 1983	Short term study <24h - included in short term review
Gotz 1983	Short term study <24h - included in short term review
Grandordy 1988	Short term study <24h - included in short term review
Gross 1975a	Short term study <24h - included in short term review
Gross 1975b	Short term study <24h - included in short term review
Gross 1984	Review
Gross 1987	Not asthma
Gutersohn 1975	Confounded - patients selected who responded to Ipratropium
Guyatt 1986	Asthma not stable
Higgins 1991	Short term study <24h - included in short term review
Hockley 1985	Short term study <24h - included in short term review
Hoffstein 1994	Short term study <24h - included in short term review
Holst 1976	Not stated to be randomised (abstract) - too long ago to write to authors
Huhti 1986	Short term study <24h - included in short term review
Hunt 1983	Short term study <24h - included in short term review
Irsigler 1975	Short term study <24h - included in short term review
Jenkins 1981	Short term study <24h - included in short term review
Jenkins 1982	Short term study <24h - included in short term review
Jindal 1989	Not randomised
Johnson 1989	Short term study <24h - included in short term review

Study	Reason for exclusion
Jolobe 1981	Not randomised
Kaik 1975	COPD
Kaik 1976	Wrong comparator
Kaik 1979	COPD
Kamburoff 1975	Short term study <24h - included in short term review
Kaptein 1993	As Brand 1992
Kennedy 1964	Short term study <24h - included in short term review
Kerstjens 1992	As Brand 1992
Kerstjens 1993a	As Brand 1992
Kerstjens 1993b	As Brand 1992
Kerstjens 1994	As Brand 1992
Kerstjens 1995	As Brand 1992
Kok-Jensen 1979	Not randomised
Kradjan 1992	Short term study <24h - included in short term review
Kreisman 1984	Short term study <24h - included in short term review
Krivoy 1987	Short term study <24h - included in short term review
Kunkel 1976	COPD
Kunkel 2000	Wrong comparison
Lahdensuo 1975	Short term study <24h - included in short term review
Laitinen 1986	Short term study <24h - included in short term review
Lanser 1975	COPD
Larsson 1987	Challenge study
Lavandier 1987	Short term study <24h - included in short term review
Lawford 1983	Short term study <24h - included in short term review
Lefcoe 1982	Short term study <24h - included in short term review
Lehrer 1993	Short term study <24h - included in short term review
Lehrer 1994	Short term study <24h - included in short term review
Li 2003	

Study	Reason for exclusion
Lichterfeld 1979	Review
Linehan 1975	Short term study <24h - included in short term review
Maesen 1975a	Short term study <24h - included in short term review
Maesen 1975b	Short term study <24h - included in short term review
Maesen 1982	Short term study <24h - included in short term review
Maesen 1985	Short term study <24h - included in short term review
Mann 1984	Confounded - patients selected who responded to Ipratropium
Marcatili 1984	Short term study <24h - included in short term review
Marin 1987	Short term study <24h - included in short term review
Marlin 1978	Short term study <24h - included in short term review
Marlin 1979	Short term study <24h - included in short term review
Mattila 1966	Short term study <24h - included in short term review
McIntosh 1986a	Short term study <24h - included in short term review
McIntosh 1986b	Short term study <24h - included in short term review
Millar 1990	Short term study <24h - included in short term review
Minette 1979	Short term study <24h - included in short term review
Minette 1985	Short term study <24h - included in short term review
Morrison 1988	Short term study <24h - included in short term review
Morrison 1989	Short term study <24h - included in short term review
Morrison 1991	Challenge study
Morton 1984	Not randomised
Muittari 1968	Short term study <24h - included in short term review
Nishi 1993	Short term study <24h - included in short term review
Nishimura 1999	Wrong comparison oxitropium vs nothing (long term study)
Nolte 1978	Short term study <24h - included in short term review
Osterwalder 1991	Short term study <24h - included in short term review
Pavia 1979	COPD; only 2 patients with asthma
Peel 1984	Short term study <24h - included in short term review

Study	Reason for exclusion
Peerless 1983	Short term study <24h - included in short term review
Petrie 1975a	Short term study <24h - included in short term review
Petrie 1975b	Short term study <24h - included in short term review
Petro 1981	COPD
Philip-Joet 1986	Acute asthma
Pierce 1979	Short term study <24h - included in short term review
Pounsford 1983	Short term study <24h - included in short term review
Pounsford 2001	Short term study <24h - included in short term review
Rafferty 1987	Short term study <24h - included in short term review
Rebuck 1979	Short term study <24h - included in short term review
Riding 1970	Short term study <24h - included in short term review
Ruffin 1977	Short term study <24h - included in short term review
Ruffin 1978	Short term study <24h - included in short term review
Ruffin 1982	Short term study <24h - included in short term review
Ruffin 1987b	Short term study <24h - included in short term review
Rutkowski 1994	Not reported to be randomised; no response from authors
Rutten-van-Molken 19	As Brand 1992
Sahay 1980	As Bell 1982
Sahlstrom 1986	Short term study <24h - included in short term review
Salome 1988	Challenge study
Schindl 1973	COPD
Schlueter 1978	Short term study <24h - included in short term review
Schroeckenstein 1988	Short term study <24h - included in short term review
Schultze-Werninghaus	Challenge study
Shneerson 1986	Short term study <24h - included in short term review
Simonsson 1979	Review
Smith 1988	Challenge study
Snow 1979	Short term study <24h - included in short term review

Study	Reason for exclusion
Spector 1975	Short term study <24h - included in short term review
Speirs 1987	Short term study <24h - included in short term review
Stewart 1987	Short term study <24h - included in short term review
Storms 1975a	Short term study <24h - included in short term review
Storms 1975b	Short term study <24h - included in short term review
Stresemann 1975	Short term study <24h - included in short term review
Stresemann 1976a	Short term study <24h - included in short term review
Stresemann1976b	Short term study <24h - included in short term review
Sur 1990	Short term study <24h - included in short term review
Tang 1984	Short term study <24h - included in short term review
Tarlo 1982	Long term study - asthma maintenance medication not stable
Taube 2002	Short term study <24h - included in short term review
Thiessen 1979	Short term study <24h - included in short term review
Trinquet 1975	Short term study <24h - included in short term review
Tukiainen 1985	Short term study <24h - included in short term review
Uffholtz 1975	Short term study <24h - included in short term review
Ullah 1980	Short term study <24h - included in short term review
Ullah 1981	Short term study <24h - included in short term review
Ulmer 1979	One study COPD; other study not randomised (before and after)
Vakil 1985	No comparison
Verstraeten 1974a	Short term study <24h - included in short term review
Verstraeten 1974b	COPD
Verstraeten 1975	Short term study <24h - included in short term review
Villate 1980	Short term study <24h - included in short term review
Vlagopoulos 1975	Short term study <24h - included in short term review
Vlagopoulos 1976	Short term study <24h - included in short term review
Vogt 1974	Short term study <24h - included in short term review
Walker 1987	Short term study <24h - included in short term review

Study	Reason for exclusion
Ward 1987	Acute asthma
Watson 1992	Not asthma
Wegener 1987a	Short term study <24h - included in short term review
Wegener 1987b	Short term study <24h - included in short term review
Wempe 1992	Short term study <24h - included in short term review
Wesolowski 1985	Children
Wildbolz 1977	Short term study <24h - included in short term review
Wilson 1978	Not randomised
Windom 1990	Short term study <24h - included in short term review
Yanai 1991	Short term study <24h - included in short term review
Yang 1989	Short term study <24h - included in short term review
Yeager 1976	Short term study <24h - included in short term review
Zanen 1995	Not randomised and wrong comparator
Zaritsky 1999	Children; acute asthma
Zuck 1987	Confounded - patients selected who responded to Ipratropium

DATA AND ANALYSES

Comparison 1. Longer term AC+BA vs BA

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Withdrawals due to treatment failure/side effects	2	256	Odds Ratio (M-H, Fixed, 95% CI)	1.40 [0.69, 2.84]
1.1 Medium/long term, parallel, BA alone dose = 2 x comb dose	2	256	Odds Ratio (M-H, Fixed, 95% CI)	1.40 [0.69, 2.84]
2 No of patients with side effects	2	256	Odds Ratio (M-H, Fixed, 95% CI)	4.51 [0.96, 21.27]
2.1 Parallel studies, overall side effects, BA alone dose = 2 x comb dose	2	256	Odds Ratio (M-H, Fixed, 95% CI)	4.51 [0.96, 21.27]

Analysis 1.1. Comparison 1 Longer term AC+BA vs BA, Outcome 1 Withdrawals due to treatment failure/side effects.

Analysis 1.2. Comparison 1 Longer term AC+BA vs BA, Outcome 2 No of patients with side effects.

Comparison 2. Longer Term AC vs placebo

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Symptom score paired mean difference (Scale -1 to +1)	3		Symptom score diff (Fixed, 95% CI)	Subtotals only
1.1 Symptom score (daily average) up to 28 days dyspnoea, daytime, IB 40	3		Symptom score diff (Fixed, 95% CI)	-0.09 [-0.14, -0.04]
1.2 Symptom score (daily) <28 days dyspnoea, daytime, IB 40; no Salorinne	2		Symptom score diff (Fixed, 95% CI)	-0.07 [-0.13, -0.02]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.3 Symptom score (daily average) medium term overall score, night	1		Symptom score diff (Fixed, 95% CI)	0.09 [-0.09, 0.26]
2 PEF paired analysis (scale -100 to +100 l/min)	3		Mean Difference L/mn (Fixed, 95% CI)	Subtotals only
2.1 PEF am	3		Mean Difference L/mn (Fixed, 95% CI)	14.38 [7.69, 21.08]
2.2 PEF am (without Salorinne)	2		Mean Difference L/mn (Fixed, 95% CI)	16.26 [9.21, 23.31]
2.3 PEF pm, short term	3		Mean Difference L/mn (Fixed, 95% CI)	23.48 [12.32, 34.65]
2.4 PEF am, short term, post bronchodilator	2		Mean Difference L/mn (Fixed, 95% CI)	34.91 [20.13, 49.70]
2.5 PEF pm, short term, post bronchodilator	1		Mean Difference L/mn (Fixed, 95% CI)	23.94 [5.20, 42.68]
3 Patient preference	2	96	Odds Ratio (M-H, Fixed, 95% CI)	1.64 [0.68, 3.94]
4 Number of patients dropping out	2	90	Odds Ratio (M-H, Fixed, 95% CI)	0.70 [0.13, 3.72]
5 Adverse effects - number of patients	2	104	Odds Ratio (M-H, Fixed, 95% CI)	1.87 [0.51, 6.81]
5.1 overall	2	104	Odds Ratio (M-H, Fixed, 95% CI)	1.87 [0.51, 6.81]

Analysis 2.1. Comparison 2 Longer Term AC vs placebo, Outcome 1 Symptom score paired mean difference (Scale -1 to +1).

Analysis 2.2. Comparison 2 Longer Term AC vs placebo, Outcome 2 PEF paired analysis (scale -100 to +100 l/min).

Analysis 2.3. Comparison 2 Longer Term AC vs placebo, Outcome 3 Patient preference.

Analysis 2.4. Comparison 2 Longer Term AC vs placebo, Outcome 4 Number of patients dropping out.

Analysis 2.5. Comparison 2 Longer Term AC vs placebo, Outcome 5 Adverse effects - number of patients.

Comparison 3. IB 40 vs IB 80

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Symptom score (scale -4 to +4)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
1.1 Dyspnoea daytime	1	26	Mean Difference (IV, Fixed, 95% CI)	-0.30 [-0.85, 0.25]
1.2 Cough daytime	1	26	Mean Difference (IV, Fixed, 95% CI)	-0.50 [-1.12, 0.12]
1.3 Dyspnoea night	1	26	Mean Difference (IV, Fixed, 95% CI)	-0.20 [-0.91, 0.51]
2 PEF (scale -100 to +100 l/min)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
2.1 PEF am	1	26	Mean Difference (IV, Fixed, 95% CI)	34.0 [-4.06, 72.06]
2.2 PEF pm	1	26	Mean Difference (IV, Fixed, 95% CI)	24.0 [-19.16, 67.16]
2.3 PEF post dose, am	1	26	Mean Difference (IV, Fixed, 95% CI)	29.00 [-11.21, 69.21]

Analysis 3.1. Comparison 3 IB 40 vs IB 80, Outcome 1 Symptom score (scale -4 to +4).

Analysis 3.2. Comparison 3 IB 40 vs IB 80, Outcome 2 PEF (scale -100 to +100 l/min).

ADDITIONAL TABLES

Table 1. Comparison of Ahonen study (IB 40 vs IB 80)

Outcome	Mean difference	% MD of IB 80	95% CI
Daily Symptom score - daytime dyspnoea	-0.30	17% decrease	-0.85, 0.25; not statistically significant
Daily Symptom score - daytime cough	-0.50	22% decrease	-1.12, 0.12; not statistically significant
Daily Symptom score - night dyspnoea	-0.20	11% decrease	-0.91, 0.51; not statistically significant
Daily PEFR - am pre-dose	34 litres/min	11% increase	-4.06, 74.06; not statistically significant

Table 1. Comparison of Ahonen study (IB 40 vs IB 80) *(Continued)*

Daily PEFR - pm pre-dose	24 litres/min	6% increase	-19.16, 67.16; not statistically significant
Daily PEFR - am post-dose	29 litres/min	7% increase	-11.21, 69.21; not statistically significant

Table 2. Anticholinergic versus placebo - symptom scores

Outcome	Scale	Studies	Pooled summary stats	Heterogeneity	Single study results	Stat sig?	% change on placebo	Comments
Symptom score - dyspnoea daytime		Ahonen 1978 Salorinne 1988 Wheeler 1987	Paired mean difference -0.09 (95%CI -0.14, -0.04)	p = 0.45, I-squared = 0%		Y	15% decrease (i.e., favours anticholinergic)	Wheeler: rho = 0.817
Symptom score - dyspnoea daytime (no Salorinne)	1-4 0-3	Ahonen 1978 Wheeler 1987	Paired mean difference -0.07 (95%CI -0.13, -0.02)	p = 0.97, I-squared = 0%		Y	12% decrease	Wheeler: rho = 0.817
Symptom score - cough daytime	1-4 0-3	Ahonen 1978 Salorinne 1988			Mean difference Aho -0.05 (95%CI -0.16, 0.06) Sal -0.05 (no SDs)	N	10% decrease 11% decrease	
Symptom score - dyspnoea night-time	1-4 0-3	Ahonen 1978 Salorinne 1988			Mean difference Aho -0.13 (95%CI -0.27, 0.01) Sal -0.09 (no SDs)	N	24% decrease 26% decrease	
Symptom score - cough night-time	0-3	Salorinne 1988			Mean difference +0.04 (no SDs)	N	14% Increase (i.e. anticholinergic worse than placebo)	
Symptom score - night-time overall score	0-3	Wheeler 1987			Paired mean difference +0.09 (95%CI -0.09, 0.26)	N	28% Increase	Wheeler: rho = 0.611
Clinical score - day and night (change score)	0-13	Taytard 1984			Mean difference (parallel trial) -0.72 (95%CI -1.37, -0.07)	Y	71% decrease	

Table 3. Anticholinergic versus placebo - daily peak flows

Outcome	Time (post dose)	Studies	Pooled summary stats	Heterogeneity	Single study re-sults	Stat sig?	% on placebo mean	Comments
Daily PEFR am pre-bronchodilator	07.00 (6h) 08.00 (10h) "on rising" (12h)	Ahonen 1978 Salorinne 1988 Wheeler 1987	Paired mean difference 14.38 L/min (95%CI 7.69, 21.08)	p = 0.22 I-squared = 34%		Y	7% increase (i.e. favours Antichol)	Wheeler: rho = 0.919
Daily PEFR am pre-bronchodilator (no Salorinne)	07.00 (6h) "on rising" (12h)	Ahonen 1978 Wheeler 1987	Paired mean difference 16.26 l/min (95%CI 9.21, 23.31)	p = 0.65 I-squared = 0%		Y	5% increase	
Daily PEFR pm pre-bronchodilator	22.00 (3h) 22.00 (4h) "on retiring" (12h)	Ahonen 1978 Salorinne 1988 Wheeler 1987	Paired mean difference 23.48 l/min (95%CI 12.32, 34.65)	p = 0.01 I-squared = 78%		Y	7% increase	Wheeler: rho = 0.911
Daily PEFR am post-bronchodilator	10.00 (3h) 10.00 (2h)	Ahonen 1978 Salorinne 1988	Paired mean difference 34.91 l/min (95%CI 20.13, 49.70)	p = 0.06 I-squared = 71%		Y	9% increase	
Daily PEFR pm post-bronchodilator	00.00 (2h)	Salorinne 1988			Paired mean diff. 23.94 L/min (95%CI 5.20, 42.68)	Y	7% increase	

Table 4. Anticholinergic versus placebo - other outcomes

Outcome	Studies	Pooled summary stats	Heterogeneity	Single study results	Stat sig?	Interpretation
Number of patients withdrawing	Pavia 1987 Wheeler 1987	OR 0.70 (95%CI 0.13, 3.72)	p = 0.25 I-squared = 23%		N	About two thirds the odds for withdrawal with anticholinergic as placebo
Patient preference (crossover studies)	Salorinne 1988 Wheeler 1987	OR 1.64 (95%CI 0.68, 3.94)	p = 0.82 I-squared = 0%		N	About 1.5 times the odds for patients preferring anticholinergic as placebo
No. patients using rescue medication	Wheeler			Daytime OR 0.66 (95%CI 0.18, 2.36) Night OR 0.58 (95%CI 0.21, 1.62)	N N	About half the odds for using rescue medication with anticholinergic as placebo

Table 5. Anticholinergic vs placebo - Adverse effects

Effect	Outcome	Studies	Pooled summary stats	Single study results	Stat sig?	Comments
Overall adverse effects	Number of patients	Vaughan 1988 Wheeler 1987	OR 1.87 (95%CI 0.51, 6.81) p = 0.81, I-squared = 0%		N	About twice the odds for patients having adverse effects with AC as with placebo
Gastric irritation	Number of patients	Wheeler 1987		OR 3.10 (95%CI 0.12, 79.23)	N	Authors: "no major differences"
Irritation	Number of days/pt	Ahonen 1978		anticholinergic = 1; placebo = 2.5		
	Symptom scores	Salorinne 1988		anticholinergic = 8; placebo = 4	"N"	
Headache	Symptom scores	Salorinne 1988		anticholinergic = 7; placebo = 14	N	
Unpleasant taste	Number of patients	Wheeler 1987		OR 2.07 (95%CI 0.18, 24.15)		
Mouth dryness	Number of patients	Vaughan 1988		OR 1.00 (95%CI 0.13, 7.81)	N	Authors; "not statistically significantly different"
	Symptom scores	Salorinne		anticholinerg = 44; placebo = 19	"N"	
Weight gain	Number of patients	Wheeler 1987		OR 3.10 (95%CI 0.12, 79.23)	N	Authors: "no major differences"
Tremor	Number of days/pt	Ahonen 1978		anticholinergic = 0; placebo = 6		
Tremor	Symptom scores	Salorinne 1988		anticholinergic = 0; placebo = 6	"N"	
Anxiety / fear Palpitations	Number of days/pt	Ahonen 1978		anticholinergic = 2; placebo = 0	"N"	Authors: "no major differences"
	Symptom scores	Salorinne 1988		anticholinerg = 10; placebo = 11		
Urinary hesitancy	Number of patients	Vaughan 1988		OR 3.14 (95%CI 0.12, 81.35)	N	

Table 6. Anticholinergic-Beta agonist combination versus BA alone - Symptom scores

Studies	Study design	BA Ratio (comb:s-ing)	Outcome	Single study results	Stat sig?	% change on BA
Haahtela 1991	Parallel 12 weeks	1:2 (Berodual/Sal)	Daytime Dyspnoea Daytime Cough	Mean difference -0.04 (minus = favours AC/BA) +0.03 (NB all symptom scores include large baseline differences)	"N" (authors)	6% decrease 5% increase
		1:2 (Duovent/Sal)	Daytime Dyspnoea Daytime Cough	+0.26 +0.40	"N" (authors)	26% increase 80% increase
Haahtela 1991	Parallel 12 weeks	1:1 (Duovent/Sal)	Daytime Dyspnoea	+0.26 (NB all symptom scores include large baseline differences)	"N" (authors)	27% increase
			Daytime Cough	+0.30	"N" (authors)	50% increase
Macaluso 1986	Crossover 2 weeks	1:1 (Duovent/Sal)	Daytime Dyspnoea	WMD -0.13 (95%CI -0.55, +0.29)	N	37% decrease relative to BA
				WMD -0.06 (95%CI -0.54, +0.42)	N	12% decrease
			Daytime Cough	WMD -0.10 (95%CI -0.54, +0.34)	N	27% decrease
			Night Dyspnoea Night Cough	WMD -0.03 (95%CI -0.46, +0.40)	N	7% decrease
Tammi-vaara	Crossover 3 weeks	1:1 (IB+oral CR terbutaline/ oral Ter)	Daytime Dyspnoea Night Dyspnoea	WMD +0.11 (95%CI -0.24, +0.46)	N	19% increase
				WMD +0.01 (95%CI -0.28, +0.30)	N	2% increase
Wolstenholme	Crossover 4 weeks	1:1 (Duovent/Sal)	Night Dyspnoea	WMD +0.16 (95%CI -0.30, +0.62)	N	22% increase
			Night Cough	WMD +0.12 (95%CI -0.24, +0.48)	N	28% increase
			Night Wheeze	WMD +0.01 (95%CI -0.54, +0.56)	N	2% increase

Table 7. AC/BA Combination versus BA alone - daily peak flows

Studies	Design&Duration	Ratio BA	Outcome	Time (post dose)	Single study results	Stat sig?	%change on BA
Philip-Joet 1990	Parallel design 1 week measurements only used (see text)	1:2 Comb : single	PEFR am	NS (NS but qid, so guess 6h)	Mean difference 4.65 l/m (95%CI -26.24, 35.54)	N	2% increase (i.e. in favour of combination)
			PEFR pm	Not Stated (NS)	MD 8.17 l/m (95%CI -24.08, 40.42)	N	3% increase
			PEFR am post dose	NS (20 mins)	MD 3.31 l/m (95%CI -29.78, 36.40)	N	1% increase
			PEFR pm post dose	NS (20 mins)	MD -4.61 l/m (95%CI -37.93, 28.71)	N	1% decrease
Haahtela 1991	Parallel design 12 weeks	1:2 Comb : Single	PEFR am	07.00h (NS but qid so guess 6h)	Mean difference +2 (Duovent/Sal) Mean difference -4 l/m (Berodual/Sal)	"N" authors	1% increase 1% decrease
			PEFR pm	19.00h (NS)	Mean difference 6 l/m (Duovent/Sal) Mean difference 2 l/m (Berodual/Sal)	"N" authors "N" authors	1% increase 1% increase
			PEFR am post dose	07.00h (1h post dose)	Mean difference 10.5 l/m (Duo/Sal) Mean difference -2 l/m (Bero/Sal)	"N" authors	2% increase 0.5% decrease
			PEFR pm post dose	19.00h (1h post dose)	Mean difference 10 l/m (Duo/Sal) Mean difference -2.5 l/m (Bero/Sal)		2% increase 1% decrease
Haahtela 1991	Parallel design 12 weeks	1:1 Comb : single	PEFR am	07.00h (NS but qid so guess 6h)	Mean difference +1.5 litres/min	"N" authors	0.5% increase
			PEFR pm	19.00h (NS)	Mean difference 9.5 l/m	"N" authors "N" authors	2% increase
			PEFR am post dose	07.00h (1h post dose)	Mean difference 15 l/m	"N" authors	3% increase
			PEFR pm post dose	19.00h (1h post dose)	Mean difference 14 l/m		3% increase
Wolstenholme 1989	Crossover design 4 weeks	1:1 Comb : single	PEFR am	NS (NS but qid so guess 6h)	Mean difference -8.70 l/m (95%CI -83.66, 66.26)	N	3% decrease
			PEFR pm	NS (NS)	Mean difference -1.70 l/m (95%CI -89.35, 85.95)	N	0.5% decrease
			PEFR am	NS (30mins)	Mean difference	N	1% decrease

Table 7. AC/BA Combination versus BA alone - daily peak flows (Continued)

			post dose		-3.00 l/m (95%CI -78.77, 72.77)		
			PEFR pm post dose	NS (30 mins)	Mean difference -4.20 l/m (95%CI -76.47, 68.07)	N	1% decrease
Tammivaara 1993	Crossover design 3 weeks beta-agonist was oral CR terbutaline	1:1 Comb : single	PEFR am PE-FR pm	08.00h (12h post dose) 20.00h (4h post ipratropium, 12h after oral terbutaline)	Mean difference -5.00 l/m (95%CI -50.72, 40.72) Mean difference 0.00 l/m (95%CI -47.48, 47.48)	N N	1% decrease no difference
Pierce 1982	Crossover design 4 weeks	1:1 Comb : single	Flow am	08.00h (6h post dose)	Mean difference 3.40 litres (95%CI -4.18, 10.98)	N	48% increase
FLOW not peak flow			Flow pm	19.00h (NS; probably 5-6h post dose)	Mean difference 6.86 litres (95%CI -5.03, 18.75)	N	48% increase
			Flow pm post dose	20.00h (1h post dose)	Mean difference 11.59 litres (95%CI -4.61, 27.79)	N	57% increase

Table 8. AC/BA combination vs BA - other outcomes

Study	Design&Duration	Comb:BA ratio	Outcomes	Pooled summary stats	Single study results	Stat sig?	Comment
Wolstenholme 1989	Crossover trial 4 weeks	1:1	Patient preference		OR 0.79 95%CI 0.20, 3.06	N	1.3 times worse with AC/BA
Tammivaara 1993	Crossover trial 3 weeks	1:1 ORAL CR terbutaline			OR 1.70 95%CI 0.40, 7.20	N	1.7 times better with AC/BA
Philip Joet 1990	Parallel design 8 weeks	1:2	Number of patients with exacerbations		OR 1.70 95%CI 0.64, 4.51	N	1.7 times worse with AC/BA
Tammivaara 1993	Crossover trial 3 weeks	1:1 CR ORAL terbutaline	Number of patients with exacerbations		OR 0.32 95%CI 0.01, 8.24	N	3 times better with AC/BA
			No. withdrawing due to treatment failure		OR 0.32 95%CI 0.01, 8.24	N	3 times better with AC/BA

Table 8. AC/BA combination vs BA - other outcomes (Continued)

Rebuck 1983	Crossover trial 4 weeks	1:1	No. withdrawing due to treatment failure	OR 2.10 95%CI 0.18, 25.01	N	2 times worse with AC/BA
Haahtela 1991	Parallel trial 12 weeks	1:1 Duovent / Sal	No. withdrawing due to treatment failure	OR 5.09 95%CI 0.98, 26.43	N	5 times worse with AC/BA
Haahtela 1991 Philip-Joet 1990	Parallel trial 12 weeks Parallel trial 8 weeks	1:2 Berod/Sal 1:2 Berod/Sal	Number of patients withdrawing due to treatment failure	OR 1.40 95%CI 0.69, 2.84 2 studies, 256 patients; Hetero- gen. p=0.22, I-squared =32.2%	N	1.4 times worse with AC/BA

Table 9. AC/BA combination vs BA - adverse effects

Effect	Outcome	Study	Details	Single study stats	Stat sig?	Comments
Overall side effects	Number of patients	Rebuck 1983	Crossover 1:1	OR 1.35 (95%CI 0.30, 6.13)	N	Combination worse
	Number of patients	Tammi-vaara 1993	Crossover 1:1 Oral terbutaline	OR 0.81 (95%CI 0.23, 2.88)	N	Combination better
Overall side effects	No. of pts withdrawn for side effects	Haahtela 1991	Parallel 1:1	OR 3.10 (95%CI 0.12, 79.23)	N	Combination worse
	Number of side effects			Combination 39; BA 25		Combination worse
Overall side effects	Number of patients withdrawn for side effects	Haahtela 1991	Parallel 1:2 Duoven-t:Salbutamol Berod-ual:Salbutamol	OR 1.00 (95%CI 0.06, 16.76) OR 5.35 (95%CI 0.25, 116.31)	N	No difference Combination worse
	No. of side effects			Duovent 39; Salbutamol 34 Berodual 28; Salbutamol 25		Similar Similar
Tremor Palpitation Bad taste Drying Paresthesia Somnolence	Incidence	Mazzei 1985	Crossover; 1:1	Combination 4; BA 1 Combination 4; BA 8 Combination 3; BA 3 Combination 3; BA 1 Combination 1; BA 0 Combination 0; BA 2 cont.		Mixed effects
Insomnia Excitation				Combination 2; BA 0 Combination 2; BA 3		

WHAT'S NEW

Date	Event	Description
11 July 2017	Amended	The previous what's new event was deleted. This review is no longer being updated.

HISTORY

Protocol first published: Issue 4, 2001
Review first published: Issue 3, 2004

Date	Event	Description
19 September 2008	New search has been performed	Literature search re-run between August 2005 and August 2008; no new studies found.
30 June 2008	Amended	Converted to new review format.
14 May 2004	New citation required and conclusions have changed	Substantive amendment

CONTRIBUTIONS OF AUTHORS

Airways group conceived the idea for the review and MW and MB formulated the question and wrote the protocol, with additional input from PG. Decisions about prioritisation in the review and splitting of the original protocol into two reviews were made by MW and MB in consultation with PG. MW devised the MS Access database, managed the studies and wrote to the authors for missing data. MW and MB carried out the inclusion of studies, quality assessment and data extraction procedures. MW entered the data into RevMan and conducted the analysis. MW and MB wrote the review, with additional input from PG.

DECLARATIONS OF INTEREST

None known.

SOURCES OF SUPPORT

Internal sources

- No sources of support supplied

External sources

- Daphne Jackson Trust, UK.

INDEX TERMS

Medical Subject Headings (MeSH)

Adrenergic beta-Agonists [therapeutic use]; Asthma [*drug therapy]; Cholinergic Antagonists [*therapeutic use]; Drug Therapy, Combination; Randomized Controlled Trials as Topic

MeSH check words

Adult; Humans