Cochrane
Library

Cochrane Database of Systematic Reviews

Drug-eluting stents versus bare-metal stents for acute coronary syndrome (Review)

Feinberg J, Nielsen EE, Greenhalgh J, Hounscome J, Sethi NJ, Safi S, Gluud C, Jakobsen JC

Feinberg J, Nielsen EE, Greenhalgh J, Hounscome J, Sethi NJ, Safi S, Gluud C, Jakobsen JC.
Drug-eluting stents versus bare-metal stents for acute coronary syndrome.
Cochrane Database of Systematic Reviews 2017, Issue 8. Art. No.: CD012481.
DOI: [10.1002/14651858.CD012481.pub2](https://doi.org/10.1002/14651858.CD012481.pub2).

www.cochranelibrary.com

TABLE OF CONTENTS

ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	4
BACKGROUND	6
OBJECTIVES	8
METHODS	9
RESULTS	13
Figure 1.	14
Figure 2.	16
Figure 3.	17
Figure 4.	19
Figure 5.	20
Figure 6.	23
DISCUSSION	26
AUTHORS' CONCLUSIONS	29
ACKNOWLEDGEMENTS	29
REFERENCES	30
CHARACTERISTICS OF STUDIES	42
DATA AND ANALYSES	85
Analysis 1.1. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 1: All-cause mortality	93
Analysis 1.2. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 2: All-cause mortality best-worst	94
Analysis 1.3. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 3: All-cause mortality worst-best	95
Analysis 1.4. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 4: All-cause mortality according to type of drug-eluting stent	96
Analysis 1.5. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 5: All-cause mortality according to type of ACS	98
Analysis 1.6. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 6: All-cause mortality according to length of maximum follow-up	99
Analysis 1.7. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 7: All-cause mortality according to registration status	100
Analysis 1.8. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 8: Serious adverse events	101
Analysis 1.9. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 9: Serious adverse events best-worst	102
Analysis 1.10. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 10: Serious adverse events worst-best	103
Analysis 1.11. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 11: Serious adverse events according to type of drug-eluting stent	104
Analysis 1.12. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 12: Serious adverse events according to type of ACS	106
Analysis 1.13. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 13: Serious adverse events according to length of maximum follow-up	107
Analysis 1.14. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 14: Serious adverse events according to registration status	109
Analysis 1.15. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 15: Major cardiovascular events	110
Analysis 1.16. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 16: Major cardiovascular events best-worst	111
Analysis 1.17. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 17: Major cardiovascular events worst-best	112

Analysis 1.18. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 18: Major cardiovascular events according to type of drug-eluting stent	113
Analysis 1.19. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 19: Major cardiovascular events according to type of ACS	114
Analysis 1.20. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 20: Major cardiovascular events according to length of maximum follow-up	115
Analysis 1.21. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 21: Major cardiovascular events according to registration status	116
Analysis 1.22. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 22: Cardiovascular mortality	117
Analysis 1.23. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 23: Cardiovascular mortality best-worst	117
Analysis 1.24. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 24: Cardiovascular mortality worst-best	118
Analysis 1.25. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 25: Cardiovascular mortality according to type of drug-eluting stent	119
Analysis 1.26. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 26: Cardiovascular mortality according to type of ACS	120
Analysis 1.27. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 27: Cardiovascular mortality according to length of maximum follow-up	121
Analysis 1.28. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 28: Cardiovascular mortality according to registration status	122
Analysis 1.29. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 29: Myocardial infarction	123
Analysis 1.30. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 30: Myocardial infarction best-worst	124
Analysis 1.31. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 31: Myocardial infarction worst-best	125
Analysis 1.32. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 32: Myocardial infarction according to type of drug-eluting stent	126
Analysis 1.33. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 33: Myocardial infarction according to type of ACS	127
Analysis 1.34. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 34: Myocardial infarction according to length of maximum follow-up	128
Analysis 1.35. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 35: Myocardial infarction according to registration status	129
Analysis 1.36. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 36: Stent thrombosis	130
Analysis 1.37. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 37: Stent thrombosis best-worst	131
Analysis 1.38. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 38: Stent thrombosis worst-best	132
Analysis 1.39. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 39: Stent thrombosis according to type of drug-eluting stent	133
Analysis 1.40. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 40: Stent thrombosis according to type of ACS	135
Analysis 1.41. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 41: Stent thrombosis according to length of maximum follow-up	136
Analysis 1.42. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 42: Stent thrombosis according to registration status	137
Analysis 1.43. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 43: Target vessel revascularisation	138
Analysis 1.44. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 44: Target vessel revascularisation best-worst	139
Analysis 1.45. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 45: Target vessel revascularisation worst-best	140

Analysis 1.46. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 46: Target vessel revascularisation according to type of drug-eluting stent	141
Analysis 1.47. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 47: Target vessel revascularisation according to type of ACS	143
Analysis 1.48. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 48: Target vessel revascularisation according to length of maximum follow-up	144
Analysis 1.49. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 49: Target vessel revascularisation according to registration status	146
Analysis 2.1. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 1: All-cause mortality	152
Analysis 2.2. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 2: All-cause mortality best-worst	152
Analysis 2.3. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 3: All-cause mortality worst-best	152
Analysis 2.4. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 4: All-cause mortality according to type of drug-eluting stent	153
Analysis 2.5. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 5: All-cause mortality according to type of ACS	153
Analysis 2.6. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 6: All-cause mortality according to registration status	154
Analysis 2.7. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 7: Serious adverse events	154
Analysis 2.8. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 8: Serious adverse events best-worst	155
Analysis 2.9. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 9: Serious adverse events worst-best	155
Analysis 2.10. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 10: Serious adverse events according to type of drug-eluting stent	156
Analysis 2.11. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 11: Serious adverse events according to type of ACS	157
Analysis 2.12. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 12: Serious adverse events according to registration status	158
Analysis 2.13. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 13: Major cardiovascular events	158
Analysis 2.14. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 14: Major cardiovascular events best-worst	159
Analysis 2.15. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 15: Major cardiovascular events worst-best	159
Analysis 2.16. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 16: Major cardiovascular events according to type of drug-eluting stent	160
Analysis 2.17. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 17: Major cardiovascular events according to type of ACS	160
Analysis 2.18. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 18: Major cardiovascular events according to registration status	161
Analysis 2.19. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 19: Cardiovascular mortality	161
Analysis 2.20. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 20: Cardiovascular mortality best-worst	162
Analysis 2.21. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 21: Cardiovascular mortality worst-best	162
Analysis 2.22. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 22: Cardiovascular mortality according to type of drug-eluting stent	163
Analysis 2.23. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 23: Cardiovascular mortality according to type of ACS	163
Analysis 2.24. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 24: Cardiovascular mortality according to registration status	164

Analysis 2.25. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 25: Myocardial infarction	164
Analysis 2.26. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 26: Myocardial infarction best-worst	165
Analysis 2.27. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 27: Myocardial infarction worst-best	165
Analysis 2.28. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 28: Myocardial infarction according to type of drug-eluting stent	166
Analysis 2.29. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 29: Myocardial infarction according to type of ACS	166
Analysis 2.30. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 30: Myocardial infarction according to registration status	167
Analysis 2.31. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 31: Stent thrombosis	167
Analysis 2.32. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 32: Stent thrombosis best-worst	168
Analysis 2.33. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 33: Stent thrombosis worst-best	168
Analysis 2.34. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 34: Stent thrombosis according to type of drug-eluting stent	169
Analysis 2.35. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 35: Stent thrombosis according to type of ACS	169
Analysis 2.36. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 36: Stent thrombosis according to registration status	170
Analysis 2.37. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 37: Target vessel revascularisation	170
Analysis 2.38. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 38: Target vessel revascularisation best-worst	171
Analysis 2.39. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 39: Target vessel revascularisation worst-best	171
Analysis 2.40. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 40: Target vessel revascularisation according to type of drug-eluting stent	172
Analysis 2.41. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 41: Target vessel revascularisation according to type of ACS	172
Analysis 2.42. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 42: Target vessel revascularisation according to registration status	173
APPENDICES	173
WHAT'S NEW	189
HISTORY	189
CONTRIBUTIONS OF AUTHORS	190
DECLARATIONS OF INTEREST	190
SOURCES OF SUPPORT	190
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	190
INDEX TERMS	191

[Intervention Review]

Drug-eluting stents versus bare-metal stents for acute coronary syndrome

Joshua Feinberg¹, Emil Eik Nielsen¹, Janette Greenhalgh², Juliet Hounscome², Naqash J Sethi¹, Sanam Safi¹, Christian Gluud³, Janus C Jakobsen^{4,5}

¹Copenhagen Trial Unit, Centre for Clinical Intervention Research, The Capital Region of Denmark, Rigshospitalet, Copenhagen University Hospital, Copenhagen, Denmark. ²Liverpool Reviews and Implementation Group, University of Liverpool, Liverpool, UK.

³Cochrane Hepato-Biliary Group, Copenhagen Trial Unit, Centre for Clinical Intervention Research, The Capital Region, Rigshospitalet, Copenhagen University Hospital, Copenhagen, Denmark. ⁴Cochrane Hepato-Biliary Group, Copenhagen Trial Unit, Centre for Clinical Intervention Research, The Capital Region of Denmark, Rigshospitalet, Copenhagen University Hospital, Copenhagen, Denmark.

⁵Department of Cardiology, Holbaek Hospital, Holbaek, Denmark

Contact: Joshua Feinberg, wtv945@alumni.ku.dk.

Editorial group: Cochrane Heart Group.

Publication status and date: Stable (no update expected for reasons given in 'What's new'), published in Issue 6, 2021.

Citation: Feinberg J, Nielsen EE, Greenhalgh J, Hounscome J, Sethi NJ, Safi S, Gluud C, Jakobsen JC. Drug-eluting stents versus bare-metal stents for acute coronary syndrome. *Cochrane Database of Systematic Reviews* 2017, Issue 8. Art. No.: CD012481. DOI: [10.1002/14651858.CD012481.pub2](https://doi.org/10.1002/14651858.CD012481.pub2).

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Approximately 3.7 million people died from acute coronary syndrome worldwide in 2012. Acute coronary syndrome, also known as myocardial infarction or unstable angina pectoris, is caused by a sudden blockage of the blood supplied to the heart muscle. Percutaneous coronary intervention is often used for acute coronary syndrome, but previous systematic reviews on the effects of drug-eluting stents compared with bare-metal stents have shown conflicting results with regard to myocardial infarction; have not fully taken account of the risk of random and systematic errors; and have not included all relevant randomised clinical trials.

Objectives

To assess the benefits and harms of drug-eluting stents versus bare-metal stents in people with acute coronary syndrome.

Search methods

We searched the Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE, Embase, LILACS, SCI-EXPANDED, and BIOSIS from their inception to January 2017. We also searched two clinical trials registers, the European Medicines Agency and the US Food and Drug Administration databases, and pharmaceutical company websites. In addition, we searched the reference lists of review articles and relevant trials.

Selection criteria

Randomised clinical trials assessing the effects of drug-eluting stents versus bare-metal stents for acute coronary syndrome. We included trials irrespective of publication type, status, date, or language.

Data collection and analysis

We followed our published protocol and the methodological recommendations of Cochrane. Two review authors independently extracted data. We assessed the risks of systematic error by bias domains. We conducted Trial Sequential Analyses to control the risks of random errors. Our primary outcomes were all-cause mortality, major cardiovascular events, serious adverse events, and quality of life. Our

secondary outcomes were angina, cardiovascular mortality, and myocardial infarction. Our primary assessment time point was at maximum follow-up. We assessed the quality of the evidence by the GRADE approach.

Main results

We included 25 trials randomising a total of 12,503 participants. All trials were at high risk of bias, and the quality of evidence according to GRADE was low to very low. We included 22 trials where the participants presented with ST-elevation myocardial infarction, 1 trial where participants presented with non-ST-elevation myocardial infarction, and 2 trials where participants presented with a mix of acute coronary syndromes.

Meta-analyses at maximum follow-up showed no evidence of a difference when comparing drug-eluting stents with bare-metal stents on the risk of all-cause mortality or major cardiovascular events. The absolute risk of death was 6.97% in the drug-eluting stents group compared with 7.74% in the bare-metal stents group based on the risk ratio (RR) of 0.90 (95% confidence interval (CI) 0.78 to 1.03, 11,250 participants, 21 trials/22 comparisons, low-quality evidence). The absolute risk of a major cardiovascular event was 6.36% in the drug-eluting stents group compared with 6.63% in the bare-metal stents group based on the RR of 0.96 (95% CI 0.83 to 1.11, 10,939 participants, 19 trials/20 comparisons, very low-quality evidence). The results of Trial Sequential Analysis showed that we did not have sufficient information to confirm or reject our anticipated risk ratio reduction of 10% on either all-cause mortality or major cardiovascular events at maximum follow-up.

Meta-analyses at maximum follow-up showed evidence of a benefit when comparing drug-eluting stents with bare-metal stents on the risk of a serious adverse event. The absolute risk of a serious adverse event was 18.04% in the drug-eluting stents group compared with 23.01% in the bare-metal stents group based on the RR of 0.80 (95% CI 0.74 to 0.86, 11,724 participants, 22 trials/23 comparisons, low-quality evidence), and Trial Sequential Analysis confirmed this result. When assessing each specific type of adverse event included in the serious adverse event outcome separately, the majority of the events were target vessel revascularisation. When target vessel revascularisation was analysed separately, meta-analysis showed evidence of a benefit of drug-eluting stents, and Trial Sequential Analysis confirmed this result.

Meta-analyses at maximum follow-up showed no evidence of a difference when comparing drug-eluting stents with bare-metal stents on the risk of cardiovascular mortality (RR 0.91, 95% CI 0.76 to 1.09, 9248 participants, 14 trials/15 comparisons, very low-quality evidence) or myocardial infarction (RR 0.98, 95% CI 0.82 to 1.18, 10,217 participants, 18 trials/19 comparisons, very low-quality evidence). The results of the Trial Sequential Analysis showed that we had insufficient information to confirm or reject our anticipated risk ratio reduction of 10% on cardiovascular mortality and myocardial infarction.

No trials reported results on quality of life or angina.

Authors' conclusions

The current evidence suggests that drug-eluting stents may lead to fewer serious adverse events compared with bare-metal stents without increasing the risk of all-cause mortality or major cardiovascular events. However, our Trial Sequential Analysis showed that there currently was not enough information to assess a risk ratio reduction of 10% for all-cause mortality, major cardiovascular events, cardiovascular mortality, or myocardial infarction, and there were no data on quality of life or angina. The evidence in this review was of low to very low quality, and the true result may depart substantially from the results presented in this review.

More randomised clinical trials with low risk of bias and low risks of random errors are needed if the benefits and harms of drug-eluting stents for acute coronary syndrome are to be assessed properly. More data are needed on the outcomes all-cause mortality, major cardiovascular events, quality of life, and angina to reduce the risk of random error.

PLAIN LANGUAGE SUMMARY

Benefits and harms of drug-eluting versus bare-metal stents in people with acute coronary syndrome

Background

Acute coronary syndrome is a term applied to heart attacks as well as chest pain during rest. The syndrome is caused by decreased blood supply to the heart due to either a narrowing of the vessels supplying the heart or a blood clot. According to the World Health Organization, 3.7 million people died from acute coronary syndrome in 2012. Percutaneous coronary intervention, commonly known as coronary angioplasty, is used to treat the narrowed or blocked vessel of the heart by inflating a balloon to allow blood to flow through the vessel again. Coronary stents are then inserted into the vessel to ensure it remains open. Two types of stents are available for use, a stent that is coated with a drug intended to reduce the risk of future blockages (drug-eluting stent) and an uncoated stent (bare-metal stent). In this review, we aimed to assess the benefits and harms of drug-eluting stents compared with bare-metal stents.

Study characteristics

We searched scientific databases from their inception to January 2017 and found 25 trials where people were randomly allocated to either a group receiving drug-eluting stents or a group receiving bare-metal stents. The 25 trials (26 comparisons) included 12,503 adults with a mean age of 60.8 years.

Key results

The people receiving drug-eluting stents did seem to have a reduced risk of experiencing a serious adverse event compared with those receiving bare-metal stents, without affecting the risk of dying from any cause or specifically heart disease, or of having a heart attack. The serious adverse event likely to be avoided by those receiving drug-eluting stents seems to be repeat balloon angioplasty. No data were reported on quality of life or chest pain (angina) after receiving the stent.

Quality of the evidence

The evidence should be interpreted with caution, as we judged the quality to be low to very low due to limitations of the included randomised clinical trials. The reason for the very low-quality judgement was mainly due to all included trials being at high risk of bias. Consequently, the results may show more benefit of drug-eluting stents than the 'real life' effect would show. Furthermore, most of the outcomes in our review lacked sufficient statistical power. Future well-designed randomised clinical trials may therefore change the above-mentioned results.

SUMMARY OF FINDINGS

Summary of findings 1. Drug-eluting stents compared to bare-metal stents for acute coronary syndrome

Drug-eluting stents compared to bare-metal stents for acute coronary syndrome

Patient or population: People with acute coronary syndrome

Setting: Hospital

Intervention: Drug-eluting stents

Comparison: Bare-metal stents

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	Nº of participants (studies)	Quality of the evidence (GRADE)	Comments
	Risk with bare-metal stents	Risk with drug-eluting stents				
All-cause mortality at maximum follow-up Follow-up: median 12 months	Study population		RR 0.90 (0.78 to 1.03)	11,250 (21 RCTs/22 comparisons)	⊕⊕⊕⊕ LOW ^{1,2}	Trial Sequential Analysis for a RRR of 10% showed that neither the boundary for futility, benefit or harm was breached, hence the risk of imprecision of the outcome result is high. Multiple eligible treatments were used in 1 trial, generating a further comparison (21 trials reporting on 22 experimental groups).
	78 per 1000	70 per 1000 (60 to 80)				
Serious adverse events at maximum follow-up Follow-up: median 12 months	Study population		RR 0.80 (0.74 to 0.86)	11,724 (22 RCTs/23 comparisons)	⊕⊕⊕⊕ LOW ³	Trial Sequential Analysis for a RRR of 10% showed that the boundary for benefit was breached, hence the risk of imprecision of the outcome result is low. Multiple eligible treatments were used in 1 trial, generating a further comparison (22 trials reporting on 23 experimental groups).
	230 per 1000	184 per 1000 (170 to 197)				
Major cardiovascular events at maximum follow-up Follow-up: median 12 months	Study population		RR 0.96 (0.83 to 1.11)	10,939 (19 RCTs/20 comparisons)	⊕⊕⊕⊕ VERY LOW ^{1,3}	Trial Sequential Analysis for a RRR of 10% showed that neither the boundary for futility, benefit or harm was breached, hence the risk of imprecision of the outcome result is high. Multiple eligible treatments were used in 1 trial, generating a further comparison (19 trials reporting on 20 experimental groups).
	66 per 1000	63 per 1000 (55 to 73)				
Quality of life at maximum follow-up - not reported	-	-	-	-	-	

Cardiovascular mortality at maximum follow-up Follow-up: median 12 months	Study population		RR 0.91 (0.76 to 1.09)	9248 (14 RCTs/15 comparisons)	⊕⊕⊕⊕ VERY LOW ^{1,3}	Trial Sequential Analysis for a RRR of 10% showed that neither the boundary for futility, benefit or harm was breached, hence the risk of imprecision of the outcome result is high. Multiple eligible treatments were used in 1 trial, generating a further comparison (14 trials reporting on 15 experimental groups).
	58 per 1000	53 per 1000 (44 to 63)				
Myocardial infarction at maximum follow-up Follow-up: median 12 months	Study population		RR 0.98 (0.82 to 1.18)	10,217 (18 RCTs/19 comparisons)	⊕⊕⊕⊕ VERY LOW ^{1,3}	Trial Sequential Analysis for a RRR of 10% showed that neither the boundary for futility, benefit or harm was breached, hence the risk of imprecision of the outcome result is high. Multiple eligible treatments were used in 1 trial, generating a further comparison (18 trials reporting on 19 experimental groups).
	48 per 1000	47 per 1000 (39 to 56)				
Angina at maximum follow-up - not reported	-	-	-	-	-	

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **OR:** odds ratio; **RCT:** randomised controlled trial; **RR:** risk ratio; **RRR:** risk ratio reduction

GRADE Working Group grades of evidence

High quality: We are very confident that the true effect lies close to that of the estimate of the effect.

Moderate quality: We are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low quality: Our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low quality: We have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

¹Downgraded one level for serious risk of imprecision due to our required information size not being met.

²Downgraded one level for serious risk of bias due to all trials being at high risk of bias, but as mortality is a more objective outcome, lack of blinding of participants, personnel and outcome assessors may not bias the outcome as much.

³Downgraded two levels for very serious risk of bias due to all trials being at high risk of bias.

BACKGROUND

Description of the condition

Cardiovascular disease, of which ischaemic heart disease is by far the largest component, is considered to be the number one cause of death globally (Lloyd-Jones 2010; Nichols 2014; Rosamond 2008). Ischaemic heart disease is generally divided into acute coronary syndrome and stable ischaemic heart disease (Roffi 2016). The disease remains prevalent, with more than 7 million people worldwide expected to develop acute coronary syndrome each year (White 2008). According to the World Health Organization, 7.4 million people died from ischaemic heart disease in 2012, representing 15% of all global deaths, with acute coronary syndrome accounting for approximately half of the deaths (Turpie 2006; WHO 2015). The in-hospital mortality proportions across different countries range from 5% to 10% (Gupta 2003). Ischaemic heart disease remains increasingly prevalent and costly to treat due to an increase in life expectancy and a decrease in death rates (Cooper 2000; Schmidt 2012).

Acute coronary syndrome is a collective term for the following:

1. unstable angina pectoris (myocardial ischaemia at rest or minimal exertion in the absence of cardiomyocyte necrosis) (Roffi 2016);
2. non-ST-segment elevation myocardial infarction (NSTEMI); and
3. ST-segment elevation myocardial infarction (STEMI) (Amsterdam 2014). Myocardial infarction can be recognised by clinical features, electrocardiographic (ECG) findings, elevated values of biochemical markers (biomarkers) of myocardial necrosis, imaging, or pathology (Thygesen 2012). The diagnosis of myocardial infarction is dependent on an elevation of the serum levels of cardiac-specific troponin I, troponin T, or the myocardial band isoenzyme of creatine kinase (CK-MB), among others (Roffi 2016). Cardiac troponin levels will usually be positive within one hour in people with myocardial infarction when using high-sensitivity assays (Roffi 2016). If the initial test is negative, a repeat test is advisable after three hours to rule out a myocardial infarction, since in some cases there is a delay in the elevation of cardiac enzymes (Roffi 2016). The changes in the ST-segment reflected in an ECG provide the basis for the distinction between NSTEMI and STEMI (Roffi 2016). Compared to NSTEMI and STEMI, unstable angina pectoris presents without the characteristic rise in cardiac-specific biomarkers (Roffi 2016).

For research purposes, the causes of myocardial infarction are generally divided into five main classes (Thygesen 2012).

- Type 1: spontaneous myocardial infarction related to atherosclerotic plaque rupture, ulceration, fissuring, erosion, or dissection with resulting intraluminal thrombus in one or more of the coronary arteries often caused by coronary artery disease.
- Type 2: myocardial infarction secondary to an ischaemic imbalance, such as coronary artery spasm, coronary embolism, anaemia, arrhythmias, hypertension, or hypotension.
- Type 3: myocardial infarction with symptoms suggestive of myocardial ischaemia and resulting in sudden unexpected cardiac death when biomarker values are unavailable or unobtainable before death.
- Type 4a: myocardial infarction associated with percutaneous coronary intervention.

- Type 4b: myocardial infarction associated with stent thrombosis, as documented by angiography or at autopsy.
- Type 5: myocardial infarction associated with coronary artery bypass graft.

Major complications associated with myocardial infarction

- Life-threatening ventricular arrhythmias caused by changes in the electrophysiologic characteristics of the myocyte, electrolyte imbalance, continuous ischaemia, and variations in heart rate, which are all due to obstruction and hence reduced flow to the myocardium and myocardial necrosis (Briege 2009; Stevenson 1989).
- Mechanical complications caused by necrosis of the myocardium, such as ventricular wall rupture, septum rupture, and papillary muscle rupture (Briege 2009; Pohjola-Sintonen 1989; Stevenson 1989).
- Cardiogenic shock caused by failure of the ventricle to pump an adequate amount of blood leading to systemic hypotension (Briege 2009; Stevenson 1989).
- Acute decompensated heart failure caused by impairment in systolic and diastolic function due to myocardial ischaemia (Briege 2009).
- Depression (Thombs 2006).

A narrowing of a coronary vessel causes unstable angina due to one of five reasons:

1. non-occlusive thrombus on pre-existing plaques;
2. dynamic obstruction, i.e. coronary vasoconstriction;
3. progressive mechanical obstruction, such as restenosis after percutaneous coronary intervention;
4. inflammation or infection; and
5. secondary unstable angina due to conditions increasing the oxygen demand, such as hypertension, thyrotoxicosis, and tachycardia (Braunwald 1998; Roffi 2016). Unstable angina is associated with lower mortality compared to myocardial infarction but similar rates of rehospitalisations; however, the patient may have a worse quality of life (Dudas 2013; Maddox 2007; Roffi 2016). People with unstable angina are also at risk of cardiogenic shock (Ruiz-Bailén 2008).

Historically, the degree of luminal stenosis and the number of coronary arteries involved (single-, double-, or triple-vessel disease) have defined the severity of ischaemic heart disease (Ringqvist 1983). Researchers have since developed more comprehensive scoring systems (Gensini 1983; Seizer 1982). Coronary angiography score and two additional scores, that is vascular scoring and stenosis scoring, determine the Gensini score (Gensini 1983). The results of the coronary angiography determine the SYNTAX score, which takes into account lesion complexity, lesion location, and number of lesions (Sianos 2005). The SYNTAX score II is used to improve the decision-making in choosing between percutaneous coronary intervention (PCI) and coronary artery bypass grafting (CABG) for a long-term, individualised risk assessment in people with complex ischaemic heart disease. The SYNTAX score II combines the anatomical-based SYNTAX score (Sianos 2005), as well as seven clinical variables (creatinine clearance, peripheral vascular disease, unprotected left main coronary disease, sex, chronic obstructive pulmonary

disease, age, and left ventricular ejection fraction) (Farooq 2013; Sianos 2005).

A glossary of medical terms is in [Appendix 1](#).

Description of the intervention

Percutaneous coronary intervention is performed by inserting an access sheath into a peripheral artery (most often the femoral artery or the radial artery); a catheter is advanced, under X-ray screening, through the blood vessels to the aortic root, at the origin of the coronary arteries. Other tools such as balloons and stents can then be advanced down the artery, over a guide wire, to the location of the narrowing or blockage (Cantor 2005; Hamon 2009). Andreas Roland Grüntzig performed the first percutaneous coronary intervention in 1977 (Grüntzig 1978). Percutaneous coronary intervention has since then evolved to become one of the cornerstones in the treatment of ischaemic heart disease.

Percutaneous coronary intervention in people with acute coronary syndrome can be performed both as primary percutaneous coronary intervention (primary PCI) and secondary percutaneous coronary intervention (secondary PCI) (Meyer 1981; Meyer 1982; Meyer 1982a; Meyer 1983). Primary PCI may be performed in the acute phase of myocardial infarction if patients present with STEMI. Patients presenting with NSTEMI or unstable angina pectoris that is characterised as high-risk acute coronary syndrome may also benefit from undergoing angiography and percutaneous coronary intervention either immediately or as part of an early (within 24 hours) invasive strategy (Hoenig 2010; Roffi 2016). The high-risk factors include haemodynamic instability, severe left ventricular dysfunction, recurrent or persistent rest angina, mechanical complications, sustained ventricular tachycardia, and dynamic ST-T wave electrocardiographic changes (Hoenig 2010).

Secondary PCI includes both subacute PCI and elective PCI. Subacute PCI is generally performed within the first 72 hours after symptom debut in people with unstable angina pectoris or NSTEMI who are not candidates for primary PCI because of being haemodynamically stable with medical therapy and not at high risk (Breall 2016; Roffi 2016). Elective PCI is performed in patients where coronary artery bypass grafting is not indicated, as well as in patients who are dissatisfied with their quality of life because of symptoms related to ischaemic heart disease or with adverse events due to their medical treatment (Levin 2016).

The first percutaneous coronary intervention was performed by inflating a balloon at the blockage of the coronary artery to dilate the artery ('balloon angioplasty') (Grüntzig 1978; Grüntzig 1979). The healing properties of the treatment seem to be related to the percutaneous coronary intervention expanding the outer diameter of the blocked coronary artery and not by decompression of the arterial plaque (Düber 1986). Balloon angioplasty, however, generally did not seem to achieve a long-lasting result, with restenosis occurring over time (Dangas 2002; Puel 1988). In an attempt to keep the lumen open longer, the next development was a small metallic scaffold called a 'stent'. These tubular devices are expanded over a balloon and press against the walls of the artery to keep it open (Puel 1988). These first devices were subsequently termed 'bare-metal stents' (after the later introduction of 'drug-eluting stents') and had improved outcomes over balloon angioplasty, but were still associated with high rates of restenosis over time (Cutlip 2002; Erbel 1998; Fischman 1994;

Macaya 1996; Puel 1988; Serruys 1994; Serruys 1998). A Cochrane review from 2005 found that primary stenting versus balloon angioplasty alone seemed to reduce the rate of reinfarction and target vessel revascularisation (Nordmann 2005).

The next stents were drug-eluting stents consisting of three main components:

1. a metal mesh;
2. an antiproliferative drug (e.g. sirolimus, paclitaxel, zotarolimus, or everolimus); and
3. a polymer used to coat the metal mesh.

Antiproliferative drugs to limit the excessive growth of neointima using cytotoxic or cytostatic agents, as well as a polymer to control the release of the antiproliferative drug, supplemented the metal mesh (Degertekin 2002; Fajadet 2006; Holmes 2004; Lee 2005; Morice 2002; Moses 2003; Stone 2004; Stone 2004a).

Newer drug-eluting stents have decreased strut thickness and are meant to have improved flexibility/deliverability, enhanced polymer biocompatibility/drug-eluting profiles, and superior re-endothelialisation kinetics (Serruys 2010; Stone 2010). They typically use everolimus or zotarolimus as their antiproliferative drug (Serruys 2010; Stone 2010).

The polymer-coating of the drug-eluting stents has, however, been linked with adverse events, such as stent thrombosis (Chen 2015). Both drug-eluting stents with a biodegradable polymer as well as polymer-free drug-eluting stents have therefore been developed. Polymer-free drug-eluting stents use the same antiproliferative drugs (such as paclitaxel or sirolimus) as the polymer drug-eluting stents (Abizaid 2010; Chen 2015).

In an attempt to further reduce the risk of restenosis, bioresorbable (also called biodegradable) stents were developed. The principal components of the bioresorbable stent are the same as the drug-eluting stents; in most cases, a polylactic acid mesh replaces the metal mesh (Haude 2013; Puricel 2015). The polylactic acid mesh is broken down and removed over time (Tamai 2000). The same types of drugs (everolimus, paclitaxel, sirolimus) used in drug-eluting stents along with Biolimus are used in bioresorbable stents (Haude 2013; Haude 2016; Puricel 2015).

Bare-metal stents, drug-eluting stents, and bioresorbable stents are used in modern percutaneous coronary interventions, with drug-eluting stents generally being the first choice (Windecker 2014). Guidelines recommend that acute coronary syndrome patients receive 12 months of antiplatelet therapy (aspirin plus a G protein-coupled purinergic (P2Y₁₂ receptor) blocker) regardless of whether percutaneous coronary intervention is performed (Windecker 2014). The recommended minimum length of duration of antiplatelet therapy for the implant of the bare-metal stents and drug-eluting stents is one month and six months, respectively (Windecker 2014).

Adverse events associated with percutaneous coronary intervention include death, coronary artery complications (such as perforation of the artery, distal embolisation (passage of an intravascular mass, which is capable of clogging capillaries), or stent thrombosis), myocardial infarction (type 4 myocardial infarction) (Thygesen 2012), vascular complications (such as bleeding or infection at the access site, retroperitoneal bleeding,

or atheroembolism), stroke, and acute kidney failure (Baim 1996; Cantor 1998; Stankovic 2004).

How the intervention might work

Percutaneous coronary intervention aims to decrease the stenosis of the coronary artery resulting in increased blood flow to the myocardium of the heart, which is thought to limit ischaemia and potentially reinfarction. Drug-eluting stents may be more beneficial than bare-metal stents because they release antiproliferative drugs, which may limit neointimal growth (Degertekin 2002; Fajadet 2006; Holmes 2004; Lee 2005; Morice 2002; Moses 2003; Stone 2004; Stone 2004a). The new bioresorbable stents as well as the polymer-free drug-eluting stents may be even more beneficial since they remove material that has been associated with adverse events (Abizaid 2010; Chen 2015; Haude 2013; Puricel 2015).

Why it is important to do this review

The prevalence of ischaemic heart disease is considerable and causes one-third of all deaths in people over 35 years of age (Lloyd-Jones 2010; Nichols 2014; Rosamond 2008). Beneficial treatments can therefore alleviate a considerable disease burden and healthcare cost. Former reviews have found conflicting evidence of the efficacy and safety of drug-eluting stents versus bare-metal stents.

A 2010 Cochrane review compared drug-eluting stents versus bare-metal stents in participants with both acute coronary syndrome and stable ischaemic heart disease (Greenhalgh 2010). The authors found no significant difference on all-cause mortality, incidence of acute myocardial infarction, or thrombosis. However, the review showed indications of beneficial effects of drug-eluting stents compared with bare-metal stents on target lesion revascularisation, target vessel revascularisation, and a composite outcome of cardiac events.

Six non-Cochrane reviews compared drug-eluting stents with bare-metal stents in participants with ischaemic heart disease (De Luca 2012; Kastrati 2007; Roukoz 2009; Stettler 2007; Suh 2011; Zheng 2014). Three reviews included only STEMI patients (De Luca 2012; Suh 2011; Zheng 2014). Two of these reviews found no difference between drug-eluting stents versus bare-metal stents on all-cause mortality and stent thrombosis, but a beneficial effect on target vessel revascularisation (De Luca 2012; Suh 2011). Suh 2011 observed improvement of recurrent myocardial infarction with drug-eluting stents, while De Luca 2012 found no effect. Zheng 2014, which included only four trials, compared the effects of bare-metal stents with drug-eluting stents in STEMI participants at five-year follow-up. The review found that there was no difference in the interventions with regard to all-cause mortality and acute myocardial infarction, as well as no effect on thrombosis, except for its occurrence later than one year after percutaneous coronary intervention. Three reviews assessed the effects of drug-eluting stents versus bare-metal stents in participants with ischaemic heart disease (including STEMI, NSTEMI, unstable angina pectoris, and stable angina pectoris) (Kastrati 2007; Roukoz 2009; Stettler 2007). Of the three reviews, two compared sirolimus stents and paclitaxel stents with bare-metal stents (Roukoz 2009; Stettler 2007), while one compared only sirolimus stents with bare-metal stents (Kastrati 2007). All reviews found no statistically significant difference between drug-eluting stents and bare-metal stents on all-cause mortality. While Stettler 2007 found a beneficial effect in favour of

the drug-eluting stents using sirolimus on myocardial infarction, they found no effect for drug-eluting stents using paclitaxel. Kastrati 2007 and Roukoz 2009 found no difference between drug-eluting stents versus bare-metal stents on myocardial infarction. All three reviews found a beneficial effect favouring drug-eluting stents on target vessel revascularisation (Kastrati 2007; Roukoz 2009; Stettler 2007).

Current guidelines on drug-eluting stents versus bare-metal stents for acute coronary syndrome

The 2011 American College of Cardiology Foundation/American Heart Association/Society for Cardiovascular Angiography and Interventions guideline recommends drug-eluting stents as a useful alternative to bare-metal stents to prevent restenosis in cases where there is an increased risk of restenosis, and the patient is likely to be able to tolerate and comply with prolonged dual antiplatelet therapy (so-called evidence A for STEMI, evidence C for NSTEMI/unstable angina pectoris; evidence levels are explained in the glossary in Appendix 1) (Levine 2011). The clinical situations associated with increased risk of restenosis are left main disease, small vessels, in-stent restenosis, bifurcations, diabetes, long lesions, multiple lesions, and saphenous vein grafts (Levine 2011; Levine 2016). The guideline also states that bare-metal stents should be used in people with a high risk of bleeding; inability to comply with one year of dual antiplatelet therapy; or anticipated invasive or surgical procedures in the next year, which the 2013 American College of Cardiology Foundation/American Heart Association Guideline for the Management of ST-Elevation Myocardial Infarction also recommends (ACEP 2013; Levine 2011).

The National Institute for Health and Care Excellence (NICE) has assessed the effects and cost-effectiveness of drug-eluting stents compared with bare-metal stents (NICE 2008). NICE recommends drug-eluting stents in people where the target artery for treatment has less than a 3 mm calibre or the lesion is longer than 15 mm, and the price difference between drug-eluting stents and bare-metal stents is no more than GBP 300.

This review is an update of the 2010 Cochrane review mentioned above (Greenhalgh 2010), which has now been divided into a review including acute coronary syndrome participants and a review including stable ischaemic heart disease participants. With the present review we planned to do the following:

1. take full account of the risk of systematic errors ('bias'), design errors, and risks of random errors ('play of chance') (Higgins 2011; Jakobsen 2014; Keus 2010; Thorlund 2011; Wetterslev 2008);
2. include trials irrespective of outcome, follow-up duration, and number of participants;
3. assess outcomes at several time points and take into account the variability of the follow-up period; and
4. include all types of drug-eluting stents, including polymer-free stents and bioresorbable stents.

OBJECTIVES

To assess the benefits and harms of drug-eluting stents versus bare-metal stents in people with acute coronary syndrome.

METHODS

Criteria for considering studies for this review

Types of studies

We searched for randomised clinical trials (both individual and cluster-randomised trials) irrespective of publication type, publication status, publication date, and language.

Types of participants

We included participants of any age with a diagnosis of acute coronary syndrome (according to the definition of the trialists).

Types of interventions

We included any type of drug-eluting stents, including bioresorbable stents and polymer-free drug-eluting stents.

We accepted any type of medical therapy as a co-intervention to percutaneous coronary intervention.

We included any type of bare-metal stent as a control intervention.

Types of outcome measures

Primary outcomes

1. All-cause mortality.
2. Serious adverse events, defined as any untoward medical occurrence that resulted in death, was life-threatening, was persistent, or led to significant disability, prolonged hospitalisation, or any medical event that had jeopardised the participant or required intervention to prevent it (ICH-GCP 1997).
3. Major cardiovascular event, defined as a composite outcome consisting of cardiovascular mortality and myocardial infarction.
4. Quality of life measured on any valid scale, such as the Seattle Angina Questionnaire or 36-Item Short Form Health Survey (SF-36) (Ware 1992; Wyrwich 2004).

Secondary outcomes

1. Cardiovascular mortality (defined by the trialists).
2. Myocardial infarction (defined by the trialists).
3. Angina on a continuous scale, such as 'angina stability' and 'angina frequency' used in the Seattle Angina Questionnaire (Wyrwich 2004).

Exploratory outcomes

1. Stent thrombosis.
2. Target vessel revascularisation (defined by the trialists). In general, target vessel revascularisation is any repeat percutaneous intervention or surgical bypass of any segment of the target vessel (Hicks 2010).

We conducted meta-analyses when possible of all dichotomous and continuous outcomes at the following two time points:

- outcomes assessed at maximal follow-up (the time point of primary interest); and
- outcomes assessed at one month.

Search methods for identification of studies

Electronic searches

We identified trials through systematic searches of the following bibliographic databases:

- the Cochrane Central Register of Controlled Trials (CENTRAL) in the Cochrane Library (2017, Issue 1 of 12), 18 January 2017;
- Epub Ahead of Print, In-Process & Other Non-Indexed Citations, MEDLINE Daily, and MEDLINE (Ovid, 1946 to 18 January 2017);
- Embase (Ovid, 1980 to 2017 week 3), 18 January 2017;
- LILACS (Latin American and Caribbean Health Sciences Literature) (Bireme, 1982 to 19 January 2017);
- SCI-EXPANDED (Thomson Reuters, 1900 to 19 January 2017);
- BIOSIS Web of Science (Thomson Reuters, 1926 to 19 January 2017).

We adapted the search strategy for MEDLINE Ovid to the other databases (Appendix 2). We applied the Cochrane sensitivity-maximising randomised controlled trial filter to MEDLINE Ovid, adaptations of which were applied to the other databases, except CENTRAL and LILACS (Lefebvre 2011). We searched all databases from their inception and imposed no restriction on language of publication.

We also conducted a search of ClinicalTrials.gov (www.clinicaltrials.gov, searched on 30 March 2017), the World Health Organization International Clinical Trials Registry Platform (WHO ICTRP) (www.apps.who.int/trialsearch, searched on 31 March 2017), European Medicines Agency (www.ema.europa.eu/ema/, searched on 3 April 2017), and the US Food and Drug Administration (www.fda.gov, searched on 4 April 2017). We also searched Google Scholar manually for trials not found in the preliminary search (Lefebvre 2011). We adapted the preliminary search strategy for MEDLINE (Ovid) for use in these databases.

We also searched the websites of the following pharmaceutical companies for ongoing or unpublished trials: Johnson & Johnson (www.jnj.com/about-jnj, searched on 30 March 2017), Medtronic (www.medtronic.com/us-en/index.html, searched on 30 March 2017), Boston Scientific (www.bostonscientific.com/en-US/Home.html, searched on 30 March 2017), and Abbott (www.abott.com/, searched 30 March 2017).

Searching other resources

We identified additional trials from the reference lists of review articles and included trials.

Data collection and analysis

We performed the review following the recommendations of the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). We conducted the analyses using Review Manager 5 (RevMan 2014), Stata 14 (Stata 2015), and Trial Sequential Analysis (CTU 2011).

Selection of studies

Two review authors (JF and EEN) independently screened titles and abstracts of all of the potentially eligible trials for inclusion. We coded all of these studies as 'retrieve' (eligible or potentially eligible/unclear) or 'do not retrieve'. In case of disagreement, we

asked a third review author to arbitrate (JCJ). We retrieved the full-text trial reports/publications, and two review authors (JF and EEN) independently screened the full texts and identified trials for inclusion. We reported reasons for exclusion of the ineligible studies. Any disagreements were resolved through discussion. We identified and excluded duplicates and collated multiple reports of the same trial so that each trial, rather than each report, was the unit of interest in this review. We recorded the selection process in sufficient detail to complete a PRISMA flow diagram and 'Characteristics of excluded studies' tables.

Data extraction and management

We used a data collection form that we had piloted on three trials in the review to collect trial characteristics and outcome data. Two review authors (JF and EEN) independently extracted all trial characteristics from the included trials. In case of disagreement, we asked a third review author to arbitrate (JCJ). We extracted the following trial characteristics.

1. Methods: duration of the trial, details of any 'run-in' period, and date of publication.
2. Participants: number randomised, number analysed, number lost to follow up/withdrawn, mean age, sex, inclusion criteria, and exclusion criteria.
3. Interventions: intervention, comparison, concomitant medications, and excluded medications.
4. Outcomes: primary and secondary outcomes specified and collected and time points reported.
5. Notes: trial funding and notable conflicts of interest of the trial authors.

Two review authors (JF and EEN) independently extracted outcome data from the included studies. Any disagreements were resolved by consensus or by involving a third review author (JCJ). One review author (JF) transferred the data into the Review Manager 5 file (RevMan 2014). We double-checked that data were entered correctly by comparing the data presented in the review with the study reports. A second review author (EEN) spot-checked study characteristics for accuracy against the trial report.

Assessment of risk of bias in included studies

Two review authors (JF and EEN) independently assessed risk of bias for each study using the criteria outlined in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). Any disagreements were resolved by discussion or by involving a third review author (JCJ). We assessed the risk of bias according to the random sequence generation, allocation sequence concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, selective outcome reporting, and other potential sources of bias. These 'risk of bias' domains enable the classification of randomised trials with an overall bias assessment at low or high risk of bias. Trials with one domain at either high or unclear risk of bias are considered at overall high risk of bias as previous meta-epidemiological studies demonstrate that these two categories do not differ in their overestimation of benefits and underestimation of harms (Gluud 2006; Kjaergard 2001; Lundh 2012; Lundh 2017; Moher 1998; Savović 2012; Savović 2012a; Schulz 1995; Wood 2008). For additional details on how we assessed the risk of bias, see Appendix 3. We assessed the risk of bias at both trial level and outcome level.

Assessment of bias in conducting the review

We conducted the review according to the published protocol and have reported any deviations from it in the [Differences between protocol and review](#) section of this review.

Measures of treatment effect

Dichotomous outcomes

We calculated risk ratios (RR) with 95% confidence intervals (CIs) for dichotomous outcomes.

Continuous outcomes

We had planned to use end scores in our analyses if both end scores and change scores were reported. If only change scores were reported, we would use these, but no trials reported any continuous outcomes. We planned to calculate mean differences (MD) and standardised mean differences (SMD) with 95% CIs for continuous outcomes, as well as Trial Sequential Analysis-adjusted CIs. We planned to use the SMD when all trials assessed the same outcome but measured it in a variety of ways, for example with different scales (Higgins 2011).

Unit of analysis issues

Where multiple trial arms were reported in a single trial, we only included the relevant arms. In the case of a trial with two relevant experimental arms (PASEO 2009a; PASEO 2009b), we included the two arms as different comparisons and halved the control group to avoid double-counting.

Dealing with missing data

We contacted investigators or study sponsors to obtain any missing data.

Dichotomous outcomes

We did not impute missing values for any outcomes in our primary analysis. We imputed data in two of our sensitivity analyses (see [Sensitivity analysis](#)).

Continuous outcomes

We did not impute missing values for any outcomes in our primary analysis. We planned to impute data in our sensitivity analysis for dichotomous and continuous outcomes (see [Sensitivity analysis](#)).

If studies did not report standard deviations (SD), we planned to calculate them using data from the trial where possible.

Assessment of heterogeneity

We primarily investigated forest plots to visually assess for any sign of heterogeneity. We additionally assessed the presence of statistical heterogeneity by Chi² test (threshold $P < 0.10$) and measured the quantities of heterogeneity using the I² statistic (Higgins 2002; Higgins 2003). We followed the recommendations for thresholds in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011):

- 0% to 40%: might not be important;
- 30% to 60%: may represent moderate heterogeneity;
- 50% to 90%: may represent substantial heterogeneity; and
- 75% to 100%: considerable heterogeneity.

We investigated possible heterogeneity through subgroup analyses. Ultimately, we performed meta-analyses of our outcomes (Higgins 2011).

Assessment of reporting biases

We planned to assess publication bias and other reporting biases by visual inspection of funnel plots for our primary outcomes if we included at least 10 trials (Higgins 2011). Using the asymmetry of the funnel plot, we assessed the risk of bias. To attempt to correct for the impact of any reporting bias, we used the 'trim and fill' method (Duval 2000; Taylor 1998).

For dichotomous outcomes, we tested asymmetry with the Harbord test, Harbord 2006, if τ^2 was less than 0.1 and with the Rücker test, Rücker 2008, if τ^2 was more than 0.1.

Data synthesis

Meta-analysis

We undertook this systematic review according to the recommendations provided in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011), according to Keus 2010, and according to the eight-step assessment suggested by Jakobsen 2014. We used the statistical software Review Manager 5, provided by Cochrane, to meta-analyse data (RevMan 2014).

We used Stata, Stata 2015, in case of zero-event trials where Review Manager 5's zero-event handling (replacing zero with a constant of 0.5) was not sufficient, for example in cases with a skewed number of participants between groups, we planned to handle this with reciprocal zero-event handling according to Sweeting 2004. Additionally, we planned on using Stata 2015 in case we needed to undertake meta-regression (post hoc).

We used Trial Sequential Analysis to assess and control the risk of random error (see below). If a meta-analysis did not reach the required information size, we presented Trial Sequential Analysis-adjusted confidence intervals to account for the lack of information.

We included all studies in our initial analyses and planned to conduct a sensitivity analysis of trials at low risk of bias, however no such trials were found. If the results were similar, we planned to base our primary conclusions on the overall analysis. If results differed, we planned to base our primary conclusions on trials at low risk of bias.

Trial Sequential Analysis

Cumulative meta-analyses are at risk of producing random errors due to sparse data and multiple testing of accumulating data (Brok 2008; Brok 2009; Higgins 2011a; Pogue 1997; Thorlund 2009; Wetterslev 2008); Trial Sequential Analysis can be applied to control this risk (CTU 2011; Jackson 2017; Thorlund 2011; Wetterslev 2017). The required information size (i.e. the number of participants needed in a meta-analysis to detect or reject a certain intervention effect) can be calculated in order to minimise random errors (Wetterslev 2008; Wetterslev 2009). The required information size takes into account the event proportion in the control group, the assumption of a plausible risk ratio reduction, and the heterogeneity of the meta-analysis (Turner 2013; Wetterslev 2008; Wetterslev 2009; Wetterslev 2017). Trial Sequential Analysis enables testing for significance each time a meta-analysis includes a new trial. On the basis of the required information size, trial

sequential monitoring boundaries can be constructed, enabling the determination of the statistical inference concerning cumulative meta-analysis that has not yet reached the required information size (Wetterslev 2008; Wetterslev 2017).

Firm evidence for benefit or harms may be established if the trial sequential monitoring boundary is crossed before reaching the required information size, in which case further trials may turn out to be superfluous. In contrast, if the boundaries for benefit or harm are not surpassed, it may be established that it is necessary to continue with further trials before a certain intervention effect can be detected or rejected. Firm evidence for lack of the postulated intervention effect, which occurs when the cumulative Z-score crosses the trial sequential monitoring boundaries for futility, can also be assessed with Trial Sequential Analysis.

To control the risks of random error, we used relatively conservative estimations of the anticipated intervention effect estimates (Jakobsen 2014). Large anticipated intervention effects lead to small required information sizes, and the thresholds for significance will be too lenient (Jakobsen 2014).

We analysed all primary and secondary outcomes with Trial Sequential Analysis. We used the following assumptions.

Primary outcomes

We estimated the diversity-adjusted required information size based on the proportion of participants with an outcome in the control group (Wetterslev 2009). We used a risk of a type I error of 2% (Jakobsen 2014), a risk of a type II error of 20%, and the diversity suggested by the trials in the meta-analysis (Jakobsen 2014). We used a risk of a type I error of 2% due to us including four primary outcomes, hence adjusting our type I error accordingly (Jakobsen 2014). We chose a risk of a type II error of 20% based primarily on tradition; it could be argued this value should be lower (Castellini 2016).

As anticipated intervention effects we used the following risk ratio reductions (RRR) because they seemed to be the maximum realistic intervention effect estimates based on former studies, trials, and meta-analyses.

- All-cause mortality: RRR of 10% (De Luca 2012; Suh 2011).
- Serious adverse events: RRR of 10%.
- Major cardiovascular event defined as a composite outcome consisting of cardiovascular mortality and myocardial infarction: RRR of 10% (De Luca 2012; Suh 2011).
- Quality of life measured on any valid scale, such as the Seattle Angina Questionnaire or SF-36: we planned to use the observed SD, a clinically relevant mean difference equal to SD/2.

Secondary outcomes

We estimated the diversity-adjusted required information size based on the proportion of participants with an outcome in the control group (Wetterslev 2009). We used a risk of a type I error of 2.5% (Jakobsen 2014), a risk of a type II error of 20%, and the diversity suggested by the trials in the meta-analysis (Jakobsen 2014). We used a risk of a type I error of 2.5% due to us including three secondary outcomes, hence adjusting our type I error accordingly (Jakobsen 2016). We chose a risk of a type II error of 20% based primarily on tradition; it could be argued this value should be lower (Castellini 2016).

As anticipated intervention effects for the secondary outcomes in the Trial Sequential Analysis, we used the following RRRs because we assumed that these intervention effects were realistic based on former studies, trials, and meta-analyses as cited below.

- Myocardial infarction: RRR of 10% (De Luca 2012; Suh 2011).
- Cardiovascular mortality: RRR of 10% (De Luca 2012; Suh 2011).
- Angina (continuous outcome): the observed SD, a clinically relevant mean difference equal to SD/2.

Exploratory outcomes

We estimated the diversity-adjusted required information size based on the proportion of participants with an outcome in the control group (Wetterslev 2009). We used a risk of a type I error of 3.33% (Jakobsen 2014), a risk of a type II error of 20%, and the diversity suggested by the trials in the meta-analysis (Jakobsen 2014). We used a risk of a type I error of 3.33% due to us including two exploratory outcomes, hence adjusting our type I error accordingly (Jakobsen 2016). We chose a risk of a type II error of 20% based primarily on tradition; it could be argued this value should be lower (Castellini 2016).

As anticipated intervention effects for the exploratory outcomes in the Trial Sequential Analysis, we used the following RRRs because we assumed that these intervention effects were realistic based on former studies, trials, and meta-analyses as cited below.

- Stent thrombosis: RRR of 10% (De Luca 2012; Suh 2011).
- Target vessel revascularisation: RRR of 30% (De Luca 2012; Suh 2011).

As supplementary Trial Sequential Analysis, we planned to use the limit of the confidence interval closest to zero effect as the anticipated intervention effect for all trial sequential analyses (Jakobsen 2014).

Assessment of significance

We assessed our intervention effects with both random-effects meta-analyses, DerSimonian 1986, and fixed-effect meta-analyses, Demets 1987, using the more conservative point estimate, that is the estimate closest to zero effect, of the two (Jakobsen 2014). If the two estimates were equal, we used the estimate with the widest confidence interval. We had four primary outcomes and three secondary outcomes, and therefore considered a P value less than 2% as significant for the primary outcomes and a P value less than 2.5% as significant for the secondary outcomes (Jakobsen 2014; Jakobsen 2016). We used the eight-step procedure to assess if the thresholds for significance were crossed or not (Jakobsen 2014).

'Summary of findings' tables

We used the GRADE approach to assess the quality of the body of evidence associated with each of the primary outcomes (all-cause mortality, serious adverse events, major cardiovascular events, and quality of life) and secondary outcomes (cardiovascular mortality, myocardial infarction, and angina) in our review (Guyatt 2008), constructing 'Summary of findings' tables using GRADEpro software (www.grade-pro.org). The GRADE approach appraises the quality of a body of evidence based on the extent to which one can be confident that an estimate of effect or association reflects the item being assessed. The quality measure of a body of evidence considers within-study risk of bias, directness of the evidence,

heterogeneity of the data, precision of effect estimates, and risk of publication bias (Guyatt 2008). We assessed the precision of the effect estimates according to Jakobsen and colleagues (Jakobsen 2014). We included all studies in our primary analyses, as we found no trials at low risk of bias or at low risk of bias in all domains other than 'blinding of participants and personnel' (Gluud 2006; Kjaergard 2001; Lundh 2012; Moher 1998; Savović 2012; Schulz 1995; Wood 2008).

Subgroup analysis and investigation of heterogeneity

We performed the following subgroup analyses.

- Type of drug-eluting stents used:
 - paclitaxel-eluting stents;
 - sirolimus-eluting stents;
 - zotarolimus-eluting stents;
 - everolimus-eluting stents;
 - bioresorbable stents;
 - polymer-free drug-eluting stents; and
 - mixed drug-eluting stents.
- Unstable angina pectoris/non-ST-segment elevation myocardial infarction participants compared with ST-segment elevation myocardial infarction participants.
- Length of maximum follow-up:
 - less than or equal to six months;
 - between six months and 12 months;
 - between one year and three years; and
 - more than or equal to three years.
- Comparison of the effect of drug-eluting stents versus bare-metal stents between trials with different clinical trial registration status:
 - pre-registration;
 - postregistration; and
 - no registration.

We only used the primary outcomes in our subgroup analyses.

We used the formal test for subgroup differences in Review Manager 5 (RevMan 2014).

We did not perform the following subgroup analyses due to lack of data.

- Participants with diabetes compared with participants without diabetes.
- Participants with high risk of bleeding (as defined by trialists) compared with participants without high risk of bleeding.
- Age of participants:
 - age 0 to 18 years;
 - age 19 to 75 years; and
 - age 76 years or above.

Sensitivity analysis

We planned to assess the potential impact of bias by performing a sensitivity analysis where we excluded trials at high risk of bias. However, we found no trials at low risk of bias.

As a secondary sensitivity analysis, we planned to only include trials with low risk of bias in all domains except 'blinding of participants

and personnel', as we did not expect to find any trials at low risk of bias in this domain. However, we found no such trial. We thoroughly discussed the limitations of trials at high risk of bias in the [Quality of the evidence](#).

To assess the potential impact of the missing data for dichotomous outcomes, we performed the two following analyses.

1. 'Best-worst-case' scenario: we assumed that all participants lost to follow-up in the experimental group survived, had no serious adverse event, had no major cardiovascular event, had no stent thrombosis, and had no target vessel revascularisation. We assumed that they also had a beneficial event with regard to quality of life and angina, defined as the group mean plus both one and two SDs of the group mean ([Jakobsen 2014](#)). We assumed that all of those with missing outcomes in the control group died, had a serious adverse event, had a major cardiovascular event, had stent thrombosis, and had target vessel revascularisation. We assumed that they also had a harmful event with regard to quality of life and angina, defined as the group mean plus both one and two SDs of the group mean ([Jakobsen 2014](#)).
2. 'Worst-best-case' scenario: we assumed that all participants lost to follow-up in the experimental group died, had a serious adverse event, had a major cardiovascular event, had stent thrombosis, and had target vessel revascularisation. We assumed that they also had a harmful event with regard to quality of life and angina, defined as the group mean plus both one and two SDs of the group mean ([Jakobsen 2014](#)). We assumed that all of those with missing outcomes in the control group survived, had no serious adverse event, had no major cardiovascular event, had no stent thrombosis, and had no target vessel revascularisation. We assumed that they also had a beneficial event with regard to quality of life and angina, defined as the group mean plus both one and two SDs of the group mean ([Jakobsen 2014](#)).

We presented results from both scenarios.

To assess the potential impact of missing SDs for continuous outcomes, we planned to perform the following sensitivity analyses.

- Where SDs were missing and not possible to calculate, we would impute SDs from trials with similar populations and low risk of bias.
- If we found no such trials, we would impute SDs from trials with a similar population. As the final option, we would impute SDs from all trials.

However, as we included no trials reporting data on continuous outcomes, we did not perform the above sensitivity analyses.

RESULTS

Description of studies

Results of the search

We identified a total of 76,576 potentially relevant references through searching the CENTRAL (Cochrane Library) (n = 8778), MEDLINE (n = 21,652), Embase (n = 14,916), SCI-EXPANDED (n = 19,920), BIOSIS (n = 10,657), and LILACS (n = 653) from inception to January 2017. The search strategy is presented in [Appendix 2](#). Additionally, we found 18 trials by searching Google Scholar, ClinicalTrials.gov, and references identified through previous meta-analyses and randomised clinical trials. We excluded 34,514 reference duplicates. Accordingly, we screened 42,080 records and excluded 41,985 references based on titles and abstracts. We assessed 95 full-text articles for eligibility. Of these, we excluded 27 references reporting on 25 trials based on our inclusion and exclusion criteria. We have provided reasons for exclusion the [Characteristics of excluded studies](#) table. Fourteen trials/15 comparisons reported on a mix of acute coronary syndrome participants and stable angina participants and are awaiting classification ([Characteristics of studies awaiting classification](#)) ([BASKET 2005a](#); [BASKET 2005b](#); [BASKET-PROVE I 2010](#); [BASKET-PROVE II 2015](#); [EAGLE 2006](#); [Erglis 2007](#); [FUTURE I 2004](#); [FUTURE II 2004](#); [Han 2007](#); [Li 2004](#); [NORSTENT 2016](#); [PRODIGY 2014](#); [SES-SMART 2004](#); [XIMA 2014](#); [ZEUS 2015](#)). We tried but could not establish contact with the trial authors to obtain individual participant data. We therefore did not include these trials in the primary analyses, but only in our exploratory analyses.

We included 51 publications reporting results of 25 trials (26 comparisons). The study flow chart is presented in [Figure 1](#).

Figure 1. study flow diagram.

Figure 1. (Continued)

Included studies

We included 51 references reporting on 25 trials (26 comparisons) in our primary analyses comparing drug-eluting stents versus bare-metal stents. The trials were published between 2004 and 2016. The trials (which often took place in more than one country) were conducted at sites in 21 different countries: eight from Italy; eight from the Netherlands; five from Spain; three from Germany; three from the UK; three from Denmark; two each from Switzerland, Argentina, Austria, and Israel; and one each from Brazil, China, Croatia, Finland, France, Iran, Norway, Poland, Serbia, Sweden, and the USA. For further details on included studies please see [Characteristics of included studies](#).

At one-month follow-up (median one-month follow-up), 6/25 trials reported all-cause mortality, 7/25 trials reported serious adverse events, 6/25 trials reported major cardiovascular events, 5/25 trials reported cardiovascular mortality, 6/25 trials reported myocardial infarction, 5/25 trials reported stent thrombosis, and 6/25 trials reported target vessel revascularisation. No trials reported quality of life or angina.

At maximum follow-up (median 12 months, range 1 to 72 months' follow-up), 21/25 trials reported all-cause mortality, 22/25 trials reported serious adverse events, 19/25 trials reported major cardiovascular events, 14/25 trials reported cardiovascular mortality, 18/25 trials reported myocardial infarction, 20/25 trials reported stent thrombosis, and 22/25 trials reported target vessel revascularisation. No trials reported quality of life or angina. Serious adverse events was not reported comprehensively and were limited to all-cause mortality, cardiovascular mortality, myocardial infarction, stent thrombosis, and target vessel revascularisation.

Participants

The 25 included trials (26 comparisons) randomised a total of 12,503 participants with acute coronary syndrome. The number of participants in each trial ranged from 16 to 3006. The mean age was 60.4 years (range 55.9 years to 65.9 years) in the drug-eluting stents group and 61.2 years (range 57.1 years to 64.6 years) in the bare-metal stents group. The mean proportion of women was 21.1% in the drug-eluting stents group and 31.3%

in the bare-metal stents group. In the drug-eluting stents group 15.4% of participants had diabetes, whereas in the bare-metal stents group 14.2% had diabetes. In both the drug-eluting stents group and the bare-metal stents group, 45% of participants had hypertension. In the drug-eluting stents group 40.5% had hyperlipidaemia, whereas in the bare-metal stents group 39.5% had hyperlipidaemia. The percentage of participants who had a previous myocardial infarction was 7.7% in the drug-eluting stents group and 7.8% in the bare-metal stents group. In the drug-eluting stents group 6.6% had undergone a previous PCI, whereas in the bare-metal stents group 5.7% had undergone a previous PCI. In the drug-eluting stents group 2.3% had had a coronary bypass graft inserted, whereas in the bare-metal stents group 1.9% had had a coronary bypass graft inserted.

In 22 trials (23 comparisons), the participants presented with STEMI. In two trials, the participants presented with different types of acute coronary syndrome. In one trial ([ELISA-3 2016](#)), participants presented with NSTEMI.

In the drug-eluting stents group, 50.1% had a lesion in the left anterior descending artery, 48.6% a lesion in the right circumflex artery, and 17.4% in the left circumflex artery. Under 1% had a lesion in the left main artery, intermediate branch, or in a graft. In the bare-metal stents group, 49.8% had a lesion in the left anterior descending artery, 49.3% a lesion in the right circumflex artery, and 18.6% in the left circumflex artery. Under 1% had a lesion in the left main artery, intermediate branch, or in a graft.

In the drug-eluting stents group, 62% of participants had one vessel disease and 38% had multivessel disease. In the bare-metal stents group, 59% had one vessel disease and 41% had multivessel disease. In the drug-eluting stents group, 62% of participants had a lesion with TIMI flow 0/1, 13.9% with TIMI 2, and 26.2% with TIMI 3. In the bare-metal stents group, 61.6% had a lesion with TIMI flow 0/1, 13.9% with TIMI 2, and 26.3% with TIMI 3. Participants may have had a lesion in more than one vessel, hence the total of the above percentages is equal to more than 100%.

Experimental intervention

We included 11 trials where the drug-eluting stent was a sirolimus-eluting stent (with seven of these being a Cypher sirolimus-eluting

stent) and eight trials (nine comparisons) where it was a paclitaxel-eluting stent (with six trials using a Taxus stent, either Express, Liberte, or unspecified). Two trials used an everolimus-eluting stent, one trial a Biolimus-eluting stent, one trial a dexamethasone-eluting stent, one trial a zotarolimus-eluting stent, and one trial a mix of drug-eluting stents (sirolimus-, paclitaxel- and zotarolimus-eluting stents). In one trial it was unclear what type of drug-eluting stent was used.

Successful PCI was performed in 95.2% of participants in the drug-eluting stents group (reported by 16 trials). The average length of the drug-eluting stent was 23.87 mm. The mean duration from onset of chest pain to PCI was 235.5 minutes (range 144 to 438 minutes) in the drug-eluting stents group

Control intervention

The bare-metal stent used in three trials was an Express bare-metal stent, two trials a Driver stent, one trial an Express2 stent, one trial a Gazelle stent, one trial a Genius MAGIC stent, one trial a Multi-Link Vision cobalt chromium stent, one trial a Vision stent, one trial a BiodivYsio stent, one trial a BX stent, and one trial a Liberte stent. In 10 trials the bare-metal stent used was not specified. One trial used a mix of different bare-metal stents.

Successful PCI was performed in 93.0% of participants in the bare-metal stents group (reported by 16 trials). The average length of the bare-metal stent was 22.29 mm. The mean duration from onset of chest pain to PCI was 242.2 minutes (range 164 to 426 minutes) in the bare-metal stents group

Co-interventions

All 25 trials (26 comparisons) employed co-interventions. In 22 of these trials the study was designed to deliver the co-interventions similarly in the drug-eluting stents group and the bare-metal stents group. In two trials the co-interventions were not specified, and in one trial the drug-eluting stents group and the bare-metal stents group received different co-interventions (different length of antithrombotic therapy). Before or at the time of PCI, aspirin was given to 99.8% of participants in both groups in the 10 trials that reported this. Clopidogrel was given to 85.4% of participants in the drug-eluting stents group and 84.6% of participants in the bare-metal stents group in the 10 trials that reported this. Heparin was given to 88.9% of participants in the drug-eluting stents group and

89.8% of participants in the bare-metal stents group in the eight trials that reported this. Glycoprotein IIb/IIIa receptor blocker was given to 63.5% of participants in the drug-eluting stents group and 61.4% of participants in the bare-metal stents group in the nine trials that reported this. For further details see [Characteristics of included studies](#).

The post-stent implantation therapy in 14 of the 25 trials was 80 mg to 100 mg of aspirin every day indefinitely. One trial administered 80 mg to 325 mg indefinitely; a second trial administered 250 mg for at least a year; and a third trial stated that participants received aspirin, but the dose was unclear. In eight trials it was unclear if the participants received aspirin in the months after stent implantation.

In three trials clopidogrel was administered for a year at an unspecified dose. In two trials 75 mg clopidogrel was administered for three months; in nine trials 75 mg was administered for nine to 12 months; and in four trials 75 mg was administered for six months. In five trials it was unclear if participants received clopidogrel in the months after stent implantation. One trial administered 500 mg ticlopidine for six weeks. Another trial administered 60 mg of prasugrel or 75 mg of clopidogrel for one year.

Excluded studies

Based on our inclusion and exclusion criteria, we excluded 27 publications on 25 trials after full-text assessment. Seven trials did not assess drug-eluting stents; four trials did not assess bare-metal stents; one trial used a different medical intervention in the drug-eluting stents group compared with the bare-metal stents group; and 13 trials were not randomised clinical trials. For further details see [Characteristics of excluded studies](#).

Risk of bias in included studies

Based on the information we collected from the published reports and received from authors, we judged all 25 trials (26 comparisons) to be at high risk of bias. We judged many trials to be at unclear risk of bias in several domains, and we were unable to obtain additional information from the authors when contacted. Additional information is provided in the 'Risk of bias' summary (Figure 2), 'Risk of bias' graph (Figure 3), and in the [Characteristics of included studies](#) table.

Figure 2. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.

Figure 3. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias): All outcomes	Blinding of outcome assessment (detection bias): All outcomes	Incomplete outcome data (attrition bias): All outcomes	Selective reporting (reporting bias)	Other bias
COMFORTABLE 2012	+	+	+	+	+	+	+
Darkahian 2014	?	?	?	?	?	?	?
DEB-AMI 2012	+	+	?	?	?	+	+
DEBATER 2012	+	?	+	+	?	+	+
DEDICATION 2008	+	+	+	+	?	+	+
De Ribamar Costa 2012	?	?	?	?	?	?	?
DEVINE 2007	?	?	?	?	?	?	?
Diaz 2007	+	?	?	?	?	?	?
ELISA-3 2016	?	+	+	+	+	+	+
EXAMINATION 2012	+	+	?	+	+	+	+
FIBISTEMI 2007	?	?	?	?	?	?	?
GRACIA-3 2010	?	+	?	+	?	?	+
HAAMU-STENT 2006	?	?	?	?	+	?	?
HORIZONS-AMI 2009	+	+	?	+	?	?	+
König 2007	?	?	?	?	?	?	?
Mission 2008	?	?	+	+	?	+	+
MULTISTRATEGY 2008	+	+	+	+	+	+	+
OCTAMI 2010	?	?	?	?	+	+	+
PASEO 2009a	+	+	+	+	?	+	+
PASEO 2009b	+	+	+	+	?	+	+
PASSION 2006	?	+	?	+	+	?	+
SELECTION 2007	?	?	?	?	+	?	?
SESAMI 2007	+	+	?	+	+	?	+

Figure 3. (Continued)

SELECTION 2007	?	?	?	?	?	?	?
SESAMI 2007	+	+	?	+	+	?	+
Steinwender 2008	?	?	?	?	+	?	?
Strozzi 2007	+	?	?	?	?	-	?
Typhoon 2006	?	+	-	?	+	?	-

Allocation

The generation of the allocation sequence was at low risk of bias in 11 trials. The remaining 14 trials were described as being randomised, but the method used for sequence generation was not mentioned.

The method used to conceal allocation was at low risk of bias in 12 trials. In the remaining 13 trials the method used for allocation concealment was insufficiently described.

Blinding

The blinding of participants was at low risk of bias in one trial ([ELISA-3 2016](#)). Nine trials stated that participants and personnel were not blinded. In the remaining 15 trials the method used to blind participants and personnel was insufficiently described.

The blinding of outcome assessors was performed and adequately described in 11 trials. One trial stated that outcome assessors were not blinded. In the remaining 13 trials the method used to blind outcome assessors was insufficiently described.

Incomplete outcome data

Thirteen trials adequately addressed incomplete outcome data. In the remaining 12 trials the method used to handle incomplete outcome data was insufficiently described.

Selective reporting

Four trials reported the results of the outcomes as specified in their respective protocols, or reported our primary outcomes, resulting in a judgement of low risk of bias according to our predefined 'Risk of bias' assessment. Five trials did not report the same outcomes as stated in the pre-published protocol. For the remaining 16 trials, we could not obtain a protocol and the trial did not report our primary outcomes sufficiently.

Other potential sources of bias

Four trials had no other biases, resulting in a judgement of low risk of bias. Eleven trials were industry funded, resulting in an

assessment of high risk of bias. The remaining 10 trials did not report how they were funded and were therefore judged to be at unclear risk of bias.

Effects of interventions

See: [Summary of findings 1 Drug-eluting stents compared to bare-metal stents for acute coronary syndrome](#)

Results at maximum follow-up

Primary outcomes

All-cause mortality

Twenty-one out of 25 trials (22 comparisons) with a total of 11,250 participants and a median follow-up of 12 months (range 1 to 72 months) reported all-cause mortality at maximum follow-up. A total of 431/6250 (6.87%) participants receiving drug-eluting stents died versus 387/5000 (7.74%) participants receiving bare-metal stents. Meta-analysis showed no evidence of a difference (risk ratio (RR) 0.90, 95% confidence interval (CI) 0.78 to 1.03, 11,250 participants, 21 trials/22 comparisons, low-quality evidence, [Analysis 1.1](#)). Hence, the absolute risk for mortality at maximum follow-up was 6.97% in the drug-eluting stents group compared with 7.74% in the bare-metal stents group.

Heterogeneity

Visual inspection of the forest plots and the tests for statistical heterogeneity ($I^2 = 0\%$; $P = 0.68$) indicated that heterogeneity might not be important.

Trial Sequential Analysis

Trial Sequential Analysis showed that there was not enough information to confirm or reject that drug-eluting stents versus bare-metal stents results in a risk ratio reduction (RRR) of 10% on all-cause mortality at maximum follow-up ([Figure 4](#)). The Trial Sequential Analysis-adjusted CI was 0.70 to 1.14.

Figure 4. Trial Sequential Analysis of drug-eluting stents versus bare-metal stents on all-cause mortality at maximum follow-up in 21 trials. The diversity-adjusted required information size (RIS) was calculated based on mortality in the control group of 7.76%; risk ratio reduction (RRR) of 10% in the experimental group; type I error of 2.0%; and type II error of 20% (80% power). No diversity was noted. The diversity-adjusted required information size was 45,046 participants. The cumulative Z-curve (blue line) did not cross the trial sequential monitoring boundaries for benefit or harm (red inward-sloping lines). The cumulative Z-curve did not cross the inner-wedge futility line (the inner-wedge futility could not be calculated due to too little information). Additionally, the cumulative Z-score did not cross the RIS. The green dotted line shows conventional boundaries (2.0%).

Bayes factor

Bayes factor was calculated based on a RR of 0.90 and the meta-analysis result (RR 0.90). Bayes factor of 0.79 was above the Bayes factor threshold for significance of 0.1, supporting that there seems to be no evidence of a beneficial effect of drug-eluting stent on all-cause mortality at maximum follow-up.

Risk of bias and sensitivity analyses

We assessed the risk of bias of the outcome result in all trials as high, therefore we could not conduct sensitivity analysis for bias.

The best-worst and worst-best case meta-analyses showed that incomplete outcome data bias had the potential to influence the results (best-worst meta-analysis (RR 0.60, 95% CI 0.44 to 0.80, 11,775 participants, 21 trials/22 comparisons, [Analysis 1.2](#)); worst-best meta-analysis (RR 1.27, 95% CI 0.99 to 1.64, 11,775 participants, 21 trials/22 comparisons, [Analysis 1.3](#))). Data were imputed for 11 trials.

Visual inspection of the funnel plots showed some signs of asymmetry. Harbord's test for small-study effects was not significant ($P = 0.19$). Using the trim and fill method, we found our meta-analysis results were not affected. Hence, we assessed the risk of publication bias as low.

Subgroup analyses

The formal test for subgroup difference in Review Manager 5 found no evidence of a difference in subgroup analyses according to type of drug-eluting stent, type of participants (STEMI compared with mixed acute coronary syndrome (ACS) participants), length of maximum follow-up, and protocol registration status ([Analysis 1.4](#); [Analysis 1.5](#); [Analysis 1.6](#); [Analysis 1.7](#)).

Serious adverse events

Twenty-two out of 25 (23 comparisons) trials with a total of 11,724 participants and a median follow-up of 12 months (range 1 to 72) reported serious adverse events at maximum follow-up. A total of 1292/6484 (19.92%) participants receiving drug-eluting stents had

20

influence the results (best-worst case meta-analysis (RR 0.67, 95% CI 0.62 to 0.71, 12,249 participants, 22 trials/23 comparisons, [Analysis 1.9](#)); worst-best case meta-analysis (RR 0.98, 95% CI 0.91 to 1.05, 12,249 participants, 22 trials/23 comparisons, [Analysis 1.10](#))). Data were imputed for 11 trials.

Visual inspection of the funnel plots showed no signs of asymmetry. Harbord's test for small-study effects was not significant ($P = 0.08$). Hence, we assessed the risk of publication bias as low.

Subgroup analyses

The formal test for subgroup difference in Review Manager 5 found no evidence of a difference in subgroup analyses according to type of drug-eluting stent, type of participants (STEMI compared with mixed ACS participants), length of maximum follow-up, and protocol registration status ([Analysis 1.11](#); [Analysis 1.12](#); [Analysis 1.13](#); [Analysis 1.14](#)).

Major cardiovascular events

Nineteen out of 25 trials (20 comparisons) with a total of 10,939 participants and a median follow-up of 12 months (range 1 to 72) reported major cardiovascular events (defined as cardiovascular death or myocardial infarction) at maximum follow-up. A total of 400/6096 (6.56%) participants receiving drug-eluting stents had a major cardiovascular event versus 321/4843 (6.63%) participants receiving bare-metal stents. Meta-analysis showed no evidence of a difference (RR 0.96, 95% CI 0.83 to 1.11, 10,939 participants, 19 trials/20 comparisons, very low-quality evidence, [Analysis 1.15](#)). Hence, the absolute risk for major cardiovascular event at maximum follow-up was 6.36% in the drug-eluting stents group compared with 6.63% in the bare-metal stents group.

Heterogeneity

Visual inspection of the forest plots and the tests for statistical heterogeneity ($I^2 = 0\%$; $P = 0.87$) indicated that heterogeneity might not be important.

Trial Sequential Analysis

Trial Sequential Analysis showed that there was not enough information to confirm or reject that drug-eluting stents versus bare-metal stents results in a RRR of 10% on major cardiovascular events at maximum follow-up. The Trial Sequential Analysis-adjusted CI was 0.53 to 1.76.

Bayes factor

Bayes factor was calculated based on a RR of 0.90 and the meta-analysis result (RR 0.96). Bayes factor of 0.57 was above the Bayes factor threshold for significance of 0.1, supporting that there seems to be no evidence of an effect of drug-eluting stents on major cardiovascular events at maximum follow-up.

Risk of bias and sensitivity analyses

We assessed the risk of bias of the outcome result in all trials as high, therefore we could not conduct sensitivity analysis for bias.

The best-worst case and worst-best case meta-analyses showed that incomplete outcome data bias alone had the potential to influence the results (best-worst case meta-analysis (RR 0.53, 95% CI 0.39 to 0.72, 11,596 participants, 19 trials/20 comparisons, [Analysis 1.16](#)); worst-best case meta-analysis (RR 1.40, 95% CI 0.97

to 2.00, 11,596 participants, 19 trials/20 comparisons, [Analysis 1.17](#))). Data were imputed for nine trials.

Visual inspection of the funnel plots showed no signs of asymmetry. Harbord's test for small-study effects was not significant ($P = 0.21$). Hence, we assessed the risk of publication bias as low.

Subgroup analyses

The formal test for subgroup difference in Review Manager 5 found no evidence of a difference in subgroup analyses according to type of drug-eluting stent, type of participants (STEMI compared with mixed ACS participants), length of maximum follow-up, and protocol registration status ([Analysis 1.18](#); [Analysis 1.19](#); [Analysis 1.20](#); [Analysis 1.21](#)).

Quality of life

No trials reported quality of life at any time point.

Secondary outcomes

Cardiovascular mortality

Fourteen out of 25 trials (15 comparisons) with a total of 9248 participants and a mean follow-up of 44.5 months (range 3 to 137) reported cardiovascular mortality at maximum follow-up. A total of 260/5402 (4.81%) participants receiving drug-eluting stents died by a cardiovascular cause versus 223/3846 (5.80%) participants receiving bare-metal stents. Meta-analysis showed no evidence of a difference (RR 0.91, 95% CI 0.76 to 1.09, 9248 participants, 14 trials/15 comparisons, very low-quality evidence, [Analysis 1.22](#)).

Heterogeneity

Visual inspection of the forest plots and the tests for statistical heterogeneity ($I^2 = 0\%$; $P = 0.44$) indicated that heterogeneity might not be important.

Trial Sequential Analysis

Trial Sequential Analysis showed that there was not enough information to confirm or reject that drug-eluting stents versus bare-metal stents results in a RRR of 10% on cardiovascular mortality at maximum follow-up. The Trial Sequential Analysis-adjusted CI was 0.44 to 1.89.

Bayes factor

Bayes factor was calculated based on a RR of 0.90 and the meta-analysis result (RR 0.91). Bayes factor of 0.9 was above the Bayes factor threshold for significance of 0.1, supporting that there seems to be no evidence of an effect of drug-eluting stents on cardiovascular mortality at maximum follow-up.

Risk of bias and sensitivity analyses

We assessed the risk of bias of the outcome result in all trials as high, therefore we could not conduct sensitivity analysis for bias.

The best-worst case and worst-best case meta-analyses showed that incomplete outcome data bias alone had the potential to influence the results (best-worst case meta-analysis (RR 0.50, 95% CI 0.32 to 0.77, 9742 participants, 14 trials, [Analysis 1.23](#)); worst-best case meta-analysis (RR 1.49, 95% CI 1.06 to 2.11, 9742 participants, 14 trials, [Analysis 1.24](#))). Data were imputed for eight trials.

Visual inspection of the funnel plots showed no signs of asymmetry. Harbord's test for small-study effects was not significant ($P = 0.39$). Hence, we assessed the risk of publication bias as low.

Subgroup analyses

The formal test for subgroup difference in Review Manager 5 found no evidence of a difference in subgroup analyses according to type of drug-eluting stent, type of participants (STEMI compared with mixed ACS participants), length of maximum follow-up, and protocol registration status ([Analysis 1.25](#); [Analysis 1.26](#); [Analysis 1.27](#); [Analysis 1.28](#)).

Myocardial infarction

Eighteen out of 25 trials (19 comparisons) with a total of 10,217 participants and a mean follow-up of 44.5 months (range 3 to 137) reported myocardial infarction at maximum follow-up. A total of 296/5732 (5.15%) participants receiving drug-eluting stents had a myocardial infarction versus 214/4485 (4.75%) participants receiving bare-metal stents. Meta-analysis showed no evidence of a difference (RR 0.98, 95% CI 0.82 to 1.18, 10,217 participants, 18 trials/19 comparisons, very low-quality evidence, [Analysis 1.29](#)).

Heterogeneity

Visual inspection of the forest plots and the tests for statistical heterogeneity ($I^2 = 0\%$; $P = 0.56$) indicated that heterogeneity might not be important.

Trial Sequential Analysis

Trial Sequential Analysis showed that there was not enough information to confirm or reject that drug-eluting stents versus bare-metal stents results in a RRR of 10% on myocardial infarction at maximum follow-up. The Trial Sequential Analysis-adjusted CI was 0.46 to 2.08.

Bayes factor

Bayes factor was calculated based on a RR of 0.90 and the meta-analysis result (RR 0.98). Bayes factor of 1.09 was above the Bayes factor threshold for significance of 0.1, supporting that there seems to be no evidence of an effect of drug-eluting stents on myocardial infarction at maximum follow-up.

Risk of bias and sensitivity analyses

We assessed the risk of bias of the outcome result in all trials as high, therefore we could not conduct sensitivity analysis for bias.

The best-worst case and worst-best case meta-analyses showed that incomplete outcome data bias alone had the potential to influence the results (best-worst case meta-analysis (RR 0.46, 95% CI 0.33 to 0.66, 10,851 participants, 18 trials/19 comparisons, [Analysis 1.30](#)); worst-best case meta-analysis (RR 1.56, 95% CI 1.00 to 2.45, 10,851 participants, 18 trials/19 comparisons, [Analysis 1.31](#))). Data were imputed for 10 trials.

Visual inspection of the funnel plots showed no signs of asymmetry. Harbord's test for small-study effects was not significant ($P = 0.39$). Hence, we assessed the risk of publication bias as low.

Subgroup analyses

The formal test for subgroup difference in Review Manager 5 found no evidence of a difference in subgroup analyses according to type of drug-eluting stent, type of participants (STEMI compared

with mixed ACS participants), length of maximum follow-up, and protocol registration status ([Analysis 1.32](#); [Analysis 1.33](#); [Analysis 1.34](#); [Analysis 1.35](#)).

Angina

No trials reported angina at any time point.

Exploratory outcomes

Stent thrombosis

Twenty out of 25 trials (21 comparisons) with a total of 11,350 participants and a mean follow-up of 44.5 months (range 3 to 137) reported stent thrombosis at maximum follow-up. A total of 268/6299 (4.23%) participants receiving drug-eluting stents had a stent thrombosis versus 223/5051 (4.41%) of participants receiving bare-metal stents. Meta-analysis showed no evidence of a difference (RR 0.96, 95% CI 0.80 to 1.16, 11,350 participants, 20 trials/21 comparisons, [Analysis 1.36](#)).

Heterogeneity

Visual inspection of the forest plots and the tests for statistical heterogeneity ($I^2 = 0\%$; $P = 0.53$) indicated that heterogeneity might not be important.

Trial Sequential Analysis

Trial Sequential Analysis showed that there was not enough information to confirm or reject that drug-eluting stents versus bare-metal stents results in a RRR of 10% on stent thrombosis at maximum follow-up. The Trial Sequential Analysis-adjusted CI was 0.45 to 2.05.

Bayes factor

Bayes factor was calculated based on a RR of 0.90 and the meta-analysis result (RR 0.96). Bayes factor of 1.03 was above the Bayes factor threshold for significance of 0.1, supporting that there seems to be no evidence of an effect of drug-eluting stents on stent thrombosis at maximum follow-up.

Risk of bias and sensitivity analyses

We assessed the risk of bias of the outcome result in all trials as high, therefore we could not conduct sensitivity analysis for bias.

The best-worst case and worst-best case meta-analyses showed that incomplete outcome data bias alone had the potential to influence the results (best-worst case meta-analysis (RR 0.47, 95% CI 0.32 to 0.67, 12,007 participants, 20 trials/21 comparisons, [Analysis 1.37](#)); worst-best case meta-analysis (RR 1.55, 95% CI 1.12 to 2.15, 12,007 participants, 20 trials/21 comparisons, [Analysis 1.38](#))). Data were imputed for 12 trials.

Visual inspection of the funnel plots showed signs of asymmetry. Harbord's test for small-study effects was not significant ($P = 0.05$). The trim and fill method did not change our results. Hence, we assessed the risk of publication bias as low.

Subgroup analyses

The formal test for subgroup difference in Review Manager 5 found no evidence of a difference in subgroup analyses according to type of drug-eluting stent, type of participants (STEMI compared with mixed ACS participants), length of maximum follow-up, and

protocol registration status ([Analysis 1.39](#); [Analysis 1.40](#); [Analysis 1.41](#); [Analysis 1.42](#)).

Target vessel revascularisation

Twenty-two out of 25 trials (23 comparisons) with a total of 11,770 participants and a median follow-up of 12 months (range 1 to 72) reported target vessel revascularisation at maximum follow-up. A total of 576/6504 (8.85%) participants receiving drug-eluting stents had a target vessel revascularisation versus 702/5266 (13.33%) participants receiving bare-metal stents. Meta-analysis showed evidence of a beneficial effect of drug-eluting stents compared with bare-metal stents on the risk of target vessel revascularisation at maximum follow-up (RR 0.58, 95% CI 0.52 to 0.65, 11,770 participants, 22 trials/23 comparisons, [Analysis 1.43](#)).

Heterogeneity

Visual inspection of the forest plots indicated no statistical heterogeneity. The test for statistical heterogeneity ($I^2 = 37\%$; $P = 0.04$) indicated moderate heterogeneity.

Trial Sequential Analysis

Trial Sequential Analysis showed that the Z-curve crossed the boundary for benefit. Hence, there seems to be firm evidence that drug-eluting stents versus bare-metal stents results in a RRR of 30% or more on the risk of target vessel revascularisation at maximum follow-up ([Figure 6](#)). In accordance, the Trial Sequential Analysis-adjusted CI was 0.50 to 0.67.

Figure 6. Trial Sequential Analysis of drug-eluting stents versus bare-metal stents on target vessel revascularisation at maximum follow-up in 22 trials. The diversity-adjusted required information size (RIS) was calculated based on a rate of target vessel revascularisations in the control group of 13.28%; risk ratio reduction (RRR) of 30% in the experimental group; type I error of 3.33%; and type II error of 20% (80% power). No diversity was noted. The diversity-adjusted required information size was 5361 participants. The cumulative Z-curve (blue line) crossed the trial sequential monitoring boundaries for benefit. The green dotted line shows conventional boundaries (3.33%).

Bayes factor

Bayes factor was calculated based on a RR of 0.70 and the meta-analysis result (RR 0.58). Bayes factor of 0.0002 was below the Bayes factor threshold for significance of 0.1, supporting that there seems to be evidence of an effect of drug-eluting stents on target vessel revascularisation at maximum follow-up.

Risk of bias and sensitivity analyses

We assessed the risk of bias of the outcome result in all trials as high, therefore we could not conduct sensitivity analysis for bias.

The best-worst case and worst-best case meta-analyses showed that incomplete outcome data bias alone had the potential

to influence the results (best-worst case random-effects meta-analysis (RR 0.41, 95% CI 0.37 to 0.45, 12,368 participants, 22 trials/23 comparisons, [Analysis 1.44](#)); worst-best case meta-analysis (RR 0.98, 95% CI 0.89 to 1.07, 12,368 participants, 22 trials/23 comparisons, [Analysis 1.45](#))). Data were imputed for 14 trials.

Visual inspection of the funnel plots showed no clear signs of asymmetry. Harbord's test for small-study effects was not significant ($P = 0.05$). Hence, we assessed the risk of publication bias as low.

Subgroup analyses

The formal test for subgroup difference in Review Manager 5 found no evidence of a difference in subgroup analyses according to type of drug-eluting stent, type of participants (STEMI compared with mixed ACS participants), length of maximum follow-up, and protocol registration status ([Analysis 1.46](#); [Analysis 1.47](#); [Analysis 1.48](#); [Analysis 1.49](#)).

Results at one-month follow-up

Primary outcomes

All-cause mortality

Six out of 25 trials with a total of 3233 participants reported target vessel revascularisation at one-month follow-up. A total of 25/1621 (1.54%) participants receiving drug-eluting stents had a target vessel revascularisation versus 51/1612 (3.16%) participants receiving bare-metal stents. Meta-analysis showed evidence of a beneficial effect of drug-eluting stents compared with bare-metal stents on the risk of target vessel revascularisation at one-month follow-up (RR 0.50, 95% CI 0.31 to 0.82, 3233 participants, 6 trials, [Analysis 2.1](#)).

Heterogeneity

Visual inspection of the forest plots and the tests for statistical heterogeneity ($I^2 = 0\%$; $P = 0.55$) indicated that heterogeneity might not be important.

Bayes factor

Bayes factor was calculated based on a RR 0.70 and the meta-analysis result (RR 0.50). Bayes factor of 0.26 was above the Bayes factor threshold for significance of 0.1, supporting that there seems to be no evidence of an effect of drug-eluting stents on target vessel revascularisation at one-month follow-up.

Risk of bias and sensitivity analyses

We assessed the risk of bias of the outcome result in all trials as high, therefore we could not conduct sensitivity analysis for bias.

The best-worst case and worst-best case meta-analyses showed that incomplete outcome data bias alone did not have the potential to influence the results (best-worst case meta-analysis (RR 0.41, 95% CI 0.26 to 0.66, 3251 participants, 6 trials, [Analysis 2.2](#)); worst-best case meta-analysis (RR 0.62, 95% CI 0.40 to 0.98, 3251 participants, 6 trials, [Analysis 2.3](#))). Data were imputed for two trials.

We did not visually inspect the funnel plots or perform Harbord's test as fewer than 10 trials were included for this outcome. Hence, we assessed the risk of publication bias as unclear.

Subgroup analyses

The formal test for subgroup difference in Review Manager 5 found no evidence of a difference in subgroup analyses according to type of drug-eluting stent, type of participants (STEMI compared with mixed ACS participants), and protocol registration status ([Analysis 2.4](#); [Analysis 2.5](#); [Analysis 2.6](#)).

Serious adverse events

Seven out of 25 trials with a total of 3313 participants reported serious adverse events at one-month follow-up. A total of 48/1661 (2.89%) participants receiving drug-eluting stents had a serious adverse event versus 74/1652 (4.48%) participants receiving bare-metal stents. Meta-analysis showed evidence of a beneficial effect of drug-eluting stents compared with bare-metal stents on the risk of serious adverse events at one-month follow-up (RR 0.65, 95% CI 0.45 to 0.93, 3313 participants, 7 trials, [Analysis 2.7](#)).

Heterogeneity

Visual inspection of the forest plots and the tests for statistical heterogeneity ($I^2 = 0\%$; $P = 0.94$) indicated that heterogeneity might not be important.

Bayes factor

Bayes factor was calculated based on a RR of 0.90 and the meta-analysis result (RR 0.65). Bayes factor of 0.68 was above the Bayes factor threshold for significance of 0.1, supporting that there seems to be no beneficial significant effect of drug-eluting stents on serious adverse events at one-month follow-up.

Risk of bias and sensitivity analyses

We assessed the risk of bias of the outcome result in all trials as high, therefore we could not conduct sensitivity analysis for bias.

The best-worst case and worst-best case meta-analyses showed that incomplete outcome data bias alone had the potential to influence the results (best-worst case meta-analysis (RR 0.56, 95% CI 0.40 to 0.80, 3331 participants, 7 trials, [Analysis 2.8](#)); worst-best case meta-analysis (RR 0.73, 95% CI 0.52 to 1.03, 3331 participants, 7 trials, [Analysis 2.9](#))). Data were imputed for three trials.

We did not visually inspect the funnel plots or perform Harbord's test as fewer than 10 trials were included for this outcome. Hence, we assessed the risk of publication bias as unclear.

Subgroup analyses

The formal test for subgroup difference in Review Manager 5 found no evidence of a difference in subgroup analyses according to type of drug-eluting stent, type of participants (STEMI compared with mixed ACS participants), and protocol registration status ([Analysis 2.10](#); [Analysis 2.11](#); [Analysis 2.12](#)).

Major cardiovascular events

Six out of 25 trials with a total of 3150 participants reported major cardiovascular events (defined as cardiovascular mortality or myocardial infarction) at one-month follow-up. A total of 30/1580 (1.90%) participants receiving drug-eluting stents had a major cardiovascular event versus 44/1570 (2.80%) participants receiving bare-metal stents. Meta-analysis showed no evidence of a difference (RR 0.68, 95% CI 0.43 to 1.08, 3150 participants, 6 trials, [Analysis 2.13](#)).

Heterogeneity

Visual inspection of the forest plots and the tests for statistical heterogeneity ($I^2 = 0\%$; $P = 0.98$) indicated that heterogeneity might not be important.

Bayes factor

Bayes factor was calculated based on a RR 0.90 and the meta-analysis result (RR 0.68). Bayes factor of 0.93 was above the Bayes factor threshold for significance of 0.1, supporting that there seems to be no evidence of an effect of drug-eluting stents on major cardiovascular events at one-month follow-up.

Risk of bias and sensitivity analyses

We assessed the risk of bias of the outcome result in all trials as high, therefore we could not conduct sensitivity analysis for bias.

The best-worst case and worst-best case meta-analyses showed that incomplete outcome data bias alone had the potential to influence the results (best-worst case meta-analysis (RR 0.54, 95% CI 0.35 to 0.84, 3168 participants, 6 trials, [Analysis 2.14](#)); worst-best case meta-analysis (RR 0.82, 95% CI 0.53 to 1.27, 3168 participants, 6 trials [Analysis 2.15](#))). Data were imputed for three trials.

We did not visually inspect the funnel plots or perform Harbord's test as fewer than 10 trials were included for this outcome. Hence, we assessed the risk of publication bias as unclear.

Subgroup analyses

The formal test for subgroup difference in Review Manager 5 found no evidence of a difference in subgroup analyses according to type of drug-eluting stent, type of participants (STEMI compared with mixed ACS participants), and protocol registration status ([Analysis 2.16](#); [Analysis 2.17](#); [Analysis 2.18](#)).

Secondary outcomes

Cardiovascular mortality

Five out of 25 trials with a total of 2406 participants reported cardiovascular mortality at one-month follow-up. A total of 22/1208 (1.82%) participants receiving drug-eluting stents died by a cardiovascular cause versus 30/1198 (2.50%) of participants receiving bare-metal stents. Meta-analysis showed no evidence of a difference (RR 0.73, 95% CI 0.43 to 1.25, 2406 participants, 5 trials, [Analysis 2.19](#)).

Heterogeneity

Visual inspection of the forest plots and the test for statistical heterogeneity ($I^2 = 0\%$; $P = 0.88$) indicated that heterogeneity might not be important.

Bayes factor

Bayes factor was calculated based on a RR 0.90 and the meta-analysis result (RR 0.73). Bayes factor of 1.02 was above the Bayes factor threshold for significance of 0.1, supporting that there seems to be no evidence of an effect of drug-eluting stents on cardiovascular mortality at one-month follow-up.

Risk of bias and sensitivity analyses

We assessed the risk of bias of the outcome result in all trials as high, therefore we could not conduct sensitivity analysis for bias.

The best-worst case and worst-best case meta-analyses showed that incomplete outcome data bias alone had the potential to influence the results (best-worst case meta-analysis (RR 0.53, 95% CI 0.32 to 0.88, 2423 participants, 5 trials, [Analysis 2.20](#)); worst-best case meta-analysis (RR 0.90, 95% CI 0.54 to 1.50, 2423 participants, 5 trials, ; [Analysis 2.21](#))). Data were imputed for two trials.

We did not visually inspect the funnel plots or perform Harbord's test as fewer than 10 trials were included for this outcome. Hence, we assessed the risk of publication bias as unclear.

Subgroup analyses

The formal test for subgroup difference in Review Manager 5 found no evidence of a difference in subgroup analyses according to type of drug-eluting stent, type of participants (STEMI compared with mixed ACS participants), and protocol registration status ([Analysis 2.22](#); [Analysis 2.23](#); [Analysis 2.24](#)).

Myocardial infarction

Six out of 25 trials with a total of 3150 participants reported myocardial infarction at one-month follow-up. A total of 13/1580 (0.82%) participants receiving drug-eluting stents had a myocardial infarction versus 26/1570 (1.66%) of participants receiving bare-metal stents. Meta-analysis showed no evidence of a difference (RR 0.51, 95% CI 0.26 to 0.98, 3150 participants, 6 trials, [Analysis 2.25](#)).

Heterogeneity

Visual inspection of the forest plots and the tests for statistical heterogeneity ($I^2 = 0\%$; $P = 0.99$) indicated that heterogeneity might not be important.

Bayes factor

Bayes factor was calculated based on a RR 0.90 and the meta-analysis result (RR 0.51). Bayes factor of 0.88 was above the Bayes factor threshold for significance of 0.1, supporting that there seems to be no evidence of an effect of drug-eluting stents on myocardial infarction at one-month follow-up.

Risk of bias and sensitivity analyses

We assessed the risk of bias of the outcome result in all trials as high, therefore we could not conduct sensitivity analysis for bias.

The best-worst case and worst-best case meta-analyses showed that incomplete outcome data bias alone had the potential to influence the results (best-worst case meta-analysis (RR 0.37, 95% CI 0.20 to 0.68, 3168 participants, 6 trials, [Analysis 2.26](#)); worst-best case meta-analysis: (RR 0.74, 95% CI 0.41 to 1.32, 3168 participants, 6 trials, [Analysis 2.27](#))). Data were imputed for three trials.

We did not visually inspect the funnel plots or perform Harbord's test as fewer than 10 trials were included for this outcome. Hence, we assessed the risk of publication bias as unclear.

Subgroup analyses

The formal test for subgroup difference in Review Manager 5 found no evidence of a difference in subgroup analyses according to type of drug-eluting stent, type of participants (STEMI compared with mixed ACS participants), and protocol registration status ([Analysis 2.28](#); [Analysis 2.29](#); [Analysis 2.30](#)).

Exploratory outcomes

Stent thrombosis

Five out of 25 trials with a total of 3070 participants reported stent thrombosis at one-month follow-up. A total of 15/1540 (0.97%) participants receiving drug-eluting stents had a stent thrombosis versus 31/1530 (2.03%) participants receiving bare-metal stents. Meta-analysis showed evidence of a beneficial effect of drug-eluting stents compared with bare-metal stents on the risk of stent thrombosis (RR 0.48, 95% CI 0.26 to 0.90, 3070 participants, 5 trials, [Analysis 2.31](#)).

Heterogeneity

Visual inspection of the forest plots and the tests for statistical heterogeneity ($I^2 = 0\%$; $P = 0.88$) indicated that heterogeneity might not be important.

Bayes factor

Bayes factor was calculated based on a RR 0.90 and the meta-analysis result (RR 0.48). Bayes factor of 0.87 was above the Bayes factor threshold for significance of 0.1, supporting that there seems to be no evidence of an effect of drug-eluting stents on stent thrombosis at one-month follow-up.

Risk of bias and sensitivity analyses

We assessed the risk of bias of the outcome result in all trials as high, therefore we could not conduct sensitivity analysis for bias.

The best-worst case and worst-best case meta-analyses showed that incomplete outcome data bias alone had the potential to influence the results (best-worst case meta-analysis (RR 0.36, 95% CI 0.20 to 0.65, 3088 participants, 5 trials, [Analysis 2.32](#)); worst-best case meta-analysis (RR 0.68, 95% CI 0.39 to 1.18, 3088 participants, 5 trials, [Analysis 2.33](#))). Data were imputed for three trials.

We did not visually inspect the funnel plots or perform Harbord's test as fewer than 10 trials were included for this outcome. Hence, we assessed the risk of publication bias as unclear.

Subgroup analyses

The formal test for subgroup difference in Review Manager 5 found no evidence of a difference in subgroup analyses according to type of drug-eluting stent, type of participants (STEMI compared with mixed ACS participants), length of maximum follow-up, and protocol registration status ([Analysis 2.34](#); [Analysis 2.35](#); [Analysis 2.36](#)).

Target vessel revascularisation

Six out of 25 trials with a total of 3233 participants reported target vessel revascularisation at one-month follow-up. A total of 25/1621 (1.54%) participants receiving drug-eluting stents had a target vessel revascularisation versus 51/1612 (3.16%) participants receiving bare-metal stents. Meta-analysis showed evidence of a beneficial effect of drug-eluting stents compared with bare-metal stents on the risk of target vessel revascularisation at one-month follow-up (RR 0.50, 95% CI 0.31 to 0.82, 3233 participants, 6 trials, [Analysis 2.37](#)).

Heterogeneity

Visual inspection of the forest plots and the tests for statistical heterogeneity ($I^2 = 0\%$; $P = 0.55$) indicated that heterogeneity might not be important.

Bayes factor

Bayes factor was calculated based on a RR 0.70 and the meta-analysis result (RR 0.50). Bayes factor of 0.26 was above the Bayes factor threshold for significance of 0.1, supporting that there seems to be no evidence of an effect of drug-eluting stents on target vessel revascularisation at one-month follow-up.

Risk of bias and sensitivity analyses

We assessed the risk of bias of the outcome result in all trials as high, therefore we could not conduct sensitivity analysis for bias.

The best-worst case and worst-best case meta-analyses showed that incomplete outcome data bias alone did not have the potential to influence the results (best-worst case meta-analysis (RR 0.41, 95% CI 0.26 to 0.66, 3251 participants, 6 trials, [Analysis 2.38](#)); worst-best case meta-analysis (RR 0.62, 95% CI 0.40 to 0.98, 3251 participants, 6 trials, [Analysis 2.39](#))). Data were imputed for two trials.

We did not visually inspect the funnel plots or perform Harbord's test as fewer than 10 trials were included for this outcome. Hence, we assessed the risk of publication bias as unclear.

Subgroup analyses

The formal test for subgroup difference in Review Manager 5 found no evidence of a difference in subgroup analyses according to type of drug-eluting stent, type of participants (STEMI compared with mixed ACS participants), and protocol registration status ([Analysis 2.40](#); [Analysis 2.41](#); [Analysis 2.42](#)).

GRADE assessments

Our main results are summarised in the [Summary of findings 1](#).

DISCUSSION

Summary of main results

We included 25 trials (26 comparisons) randomising a total of 12,503 participants. All trials and outcomes were at high risk of bias, and the quality of the evidence according to the GRADE assessment was low to very low for all outcomes.

Meta-analyses showed no evidence of a difference when comparing drug-eluting stents with bare-metal stents on the risk of all-cause mortality and on major cardiovascular events. The results of our Trial Sequential Analysis showed that we did not have sufficient information to confirm or reject our anticipated RRR of 10% on both risk of all-cause mortality and on major cardiovascular events.

Meta-analyses showed evidence of a beneficial effect of drug-eluting stents compared with bare-metal stents on the risk of serious adverse events. The Trial Sequential Analysis for serious adverse events showed that the Z-curve crossed the boundary for benefit for drug-eluting stents. Hence, there seems to be firm evidence that drug-eluting stents versus bare-metal stents results in a RRR of 10% or more on the risk of a serious adverse event at maximum follow-up. When assessing specific

types of events included in the serious adverse event outcome separately, we found the majority of the events were target vessel revascularisation.

Formal tests for subgroup differences showed no evidence of a difference according to drug-eluting stent type (including bioresorbable stents), type of participants (STEMI, NSTEMI, ACS), protocol registration status, and length of follow-up for any of our primary outcomes.

Meta-analyses showed no evidence of a difference when comparing drug-eluting stents with bare-metal stents on the risk of cardiovascular mortality and myocardial infarction. The results of our Trial Sequential Analysis showed that we did not have sufficient information to confirm or reject our anticipated RRR of 10% on both risk of cardiovascular mortality and on myocardial infarction.

No trials reported results on angina or quality of life.

Overall completeness and applicability of evidence

This review provides the most comprehensive and contemporary appraisal of the evidence on drug-eluting stents versus bare-metal stents to date. We searched for published and unpublished trials irrespective of publication type, publication status, publication date, and language. We also searched bibliographies of both Cochrane and non-Cochrane reviews for any trials we might have missed. The high risk of bias in our trials suggests that our results may show an overestimation of the benefit of drug-eluting stents and an underestimation of the harm of drug-eluting stents.

No trials reported quality of life or the number of participants with angina.

A number of trials did not report data separately on ACS participants and participants with other ischaemic heart disease, therefore we could not include these participants in our analyses. We have included these trials as 'studies awaiting classification'. The results from these trials deviate from our analyses in showing a difference between interventions for major cardiovascular events and myocardial infarction. We have attempted to contact the trial authors for specific information on ACS participants, and hope to receive these data in the future.

Quality of the evidence

Heterogeneity

We assessed the statistical heterogeneity in the planned analyses of our primary and secondary outcomes as low.

Risk of systematic error ('bias')

Our 'Risk of bias' assessment showed that all trials were at high risk of bias. It is, therefore, highly probable that our review results are also biased, that is that there is a great risk that our results overestimate benefits and underestimate harm (Hrobjartsson 2012; Hrobjartsson 2013; Hrobjartsson 2014; Hrobjartsson 2014a; Jakobsen 2014; Lundh 2012; Savović 2012; Schulz 1995; Wood 2008). We have downgraded all of our outcomes by two levels for a very serious risk of bias.

Only one trial had low risk of bias in blinding of participants and personnel, due to the nature of the PCI procedure. Trials without blinding of participants and personnel tend to overestimate

benefits and underestimate harm (Hrobjartsson 2014; Pocock 2015; Savović 2012; Wood 2008). This may be due to a lack of expectations in a control group; differential behaviours across intervention groups (e.g. differential dropout or differential cross-over to an alternative intervention); or different perception and reporting of an outcome such as pain or feeling more obliged to consider participants for target vessel revascularisation when knowing that a participant had received a bare-metal stent (Higgins 2011a).

We assessed 12 out of 25 trials as at unclear or high risk of bias for the incomplete outcome data bias domain. Our best-worst case and worst-best case analyses showed that incomplete outcome data alone had the potential to change our results at maximum follow-up. Where insufficient data were reported by the trialists, we attempted to contact the authors, but frequently did not receive a reply. Hence, the extent of the incomplete outcome data was often unclear. Our best-worst case and worst-best case analyses might underestimate the potential impact of missing data because only the data on the reported population were used if no other information was available. Incomplete outcome data bias might potentially have an even greater bias impact than our best-worst case/worst-best case scenarios show, that is the 'true' difference between the actually observed cases and the intention-to-treat population might be larger than our data suggest.

Publication bias

Most of our funnel plots and Harbord's tests did not show clear signs of asymmetry (only all-cause mortality and stent thrombosis showed some signs). We assessed those that did show signs with a trim and fill method, which showed that the results were not changed. Hence, there was no strong suspicion of significant small-study bias or publication bias for any of our results.

Indirectness

In our review the composite outcome 'serious adverse events' was largely driven by target vessel revascularisation. The reason for performing target vessel revascularisation is typically based on clinical characteristics such as unstable angina or an angiographic indication. The angiographic indication could be problematic: it may be that there is an angiographic difference at follow-up between participants receiving drug-eluting stents and participants receiving bare-metal stents, but that this angiographic difference does not necessarily translate into different clinical outcomes. Despite the above, we have not chosen to downgrade for indirectness, but the reader should be aware of this limitation.

Imprecision (risk of random error)

The Trial Sequential Analysis of serious adverse events confirmed that we had enough information to assess a 10% RRR at maximum follow-up. Trial Sequential Analysis of all-cause mortality, major cardiovascular events, cardiovascular mortality, and myocardial infarction showed that we did not have enough information to assess a 10% RRR at maximum follow-up, resulting in a GRADE downgrade for imprecision for these outcomes.

GRADE

We have assessed the quality of the evidence for the results of each outcome using GRADE (Summary of findings 1). The GRADE assessment generally showed the evidence to be of low to very low quality. Reasons for the GRADE assessment are given in the footnotes of the Summary of findings 1.

Potential biases in the review process

Strengths

Our review has several strengths. We included trials regardless of language of publication and whether they reported data for the outcomes we had planned to assess. Where additional information was needed, we contacted all relevant authors. We included more participants than any previous systematic review, which gave us increased power and precision to detect any significant differences between the intervention and the control group. We followed our peer-reviewed protocol, which was published before the literature search began ([Feinberg 2016](#)), and we conducted the review using the methods recommended by Cochrane and findings from additional methodological studies ([Higgins 2011a](#)). We also performed Trial Sequential Analyses and used an eight-step procedure to assess if the thresholds for statistical and clinical significance were crossed ([Jakobsen 2014](#)), which adds further robustness to our results and conclusions. We also tested the robustness of our results with sensitivity analyses (best-worst case and worst-best case). Our meta-analyses had little statistical heterogeneity, strengthening the validity of our results.

Limitations

Our systematic review has several limitations. Our findings, interpretations, and conclusions are affected by the quality and quantity of the included trials. The relatively short follow-up might also make it harder to detect a difference; it may be that longer follow-up would reveal such a difference.

Furthermore, by only including randomised clinical trials and ignoring harms assessed in quasi-randomised studies and observational studies, we run the risk of focusing overly on potential benefits at the cost of overlooking late and rare harms.

Another, and likely often ignored, limitation is the fact that we have compared drug-eluting stents versus bare-metal stents without having solid proof that the latter intervention is significantly more beneficial from a statistical and clinical point of view than just balloon dilatation or no intervention at all (see [Description of the intervention](#)). Evidence points in this direction, but solid evidence based on systematic reviews with meta-analysis and Trial Sequential Analysis taking risks of bias into consideration is lacking ([Garattini 2016](#)).

In our primary analyses, we included only one trial using a biodegradable drug-eluting stent in the experimental group ([COMFORTABLE 2012](#)). Our subgroup assessing different types of drug-eluting stents compared with bare-metal stents found no differences between the experimental stents, but this may be due to a low number of included trials evaluating these types of stents. In the future, network meta-analyses and individual participant data meta-analysis may assess these types of stents ([Garattini 2016](#); [Skoog 2015](#)).

We used observed data in the estimation of a required information size, which might lead to lack of consistency in stopping boundaries across analyses. The alternative would be to predefine diversity for each analysis the incidence in all of the control groups before data were known. This also has several limitations, that is how these were values estimated, and if they are erroneous then this will lead to erroneous results. We chose from a pragmatic perspective the observed data ([Jakobsen 2014](#)).

Composite outcome

To increase the statistical power, we chose to use two composite outcomes: major cardiovascular events and serious adverse events. A potential limitation when using composite outcomes is that each component of a composite outcome will not necessarily have similar degrees of severity, especially with serious adverse events, and this might bias the results of the outcome ([Garattini 2016](#)). For example, if certain more severe serious adverse events occur in one intervention group and other less severe serious adverse events occur in the other intervention group, then there is a risk of overlooking actual severity differences between the compared groups on this composite outcome ([Garattini 2016](#)).

In our review, the composite outcome serious adverse events was largely driven by target vessel revascularisation. The reason for performing target vessel revascularisation is typically based on clinical characteristics such as unstable angina or an angiographic indication. The angiographic indication could be problematic: it may be that there is an angiographic difference at follow-up between participants receiving drug-eluting stents and participants receiving bare-metal stents, but that this angiographic difference does not necessarily translate into different clinical outcomes. For this reason we did not use the traditional outcome MACE (major adverse cardiac events). MACE usually consists of a composite of cardiac death, myocardial infarction, and either target vessel revascularisation or target lesion revascularisation, but may include other outcomes ([Kip 2008](#)). In recent years, two major reports have described and analysed several issues regarding the cardiac composite outcome MACE ([Cutlip 2007](#); [Kip 2008](#)). The main issues regarding MACE concern the variability and lack of consistency in which outcomes are included in the composite outcome ([Cutlip 2007](#); [Kip 2008](#)), which can lead to misleading conclusions. The main issue with MACE in this particular review was using target vessel revascularisation or target lesion revascularisation as a component of MACE. This was problematic for several reasons. Firstly, it is important to note that the decision of whether or not target vessel revascularisation will be performed is based on subjective opinion. As both treatment providers and participants in the included trials were not blinded to treatment allocation, target vessel revascularisation and target lesion revascularisation may introduce bias. Secondly, using a composite outcome consisting of safety outcomes (death and myocardial infarction) and outcomes presumed to be a measure of procedural effectiveness (target vessel revascularisation and target lesion revascularisation) could lead to erroneous conclusions ([Kip 2008](#)). We therefore decided not to use MACE as a composite outcome, and instead used a composite cardiovascular outcome consisting of cardiovascular mortality and myocardial infarction ('major cardiovascular events').

It must be noted that we found no evidence of a difference when assessing all-cause mortality, cardiovascular mortality, and myocardial infarction, but the Trial Sequential Analysis showed that we did not have enough information to confirm or reject our anticipated intervention effects, so the lack of a shown difference on these clinical outcomes might be caused by lack of statistical power.

Another limitation when assessing serious adverse events is that the results on serious adverse events were not comprehensively reported in any of the included trials ([Ioannidis 2009](#)). The only serious adverse events reported were usually all-cause mortality,

cardiovascular mortality, myocardial infarction, and target vessel revascularisation.

Patient group

Our primary analyses assessed the effects of drug-eluting stents versus bare-metal stents for acute coronary syndrome. During our literature search we found 14 trials pooling participants with both ACS and stable angina (BASKET 2005a; BASKET 2005b; BASKET-PROVE I 2010; BASKET-PROVE II 2015; EAGLE 2006; Erglis 2007; FUTURE I 2004; FUTURE II 2004; Han 2007; Li 2004; NORSTENT 2016; PRODIGY 2014; SES-SMART 2004; XIMA 2014; ZEUS 2015). The results were similar to our primary results except for the outcome major cardiovascular events (RR 0.72, 95% CI 0.60 to 0.85, $I^2 = 26\%$, 17,628 participants, 12 trials) and myocardial infarction (RR 0.57, 95% CI 0.41 to 0.79, $I^2 = 58\%$, 17,628 participants, 12 trials).

Agreements and disagreements with other studies or reviews

A 2010 Cochrane review compared drug-eluting stents versus bare-metal stents in participants with both ACS and stable ischaemic heart disease (Greenhalgh 2010). They found no significant difference on all-cause mortality, incidence of acute myocardial infarction, or thrombosis. However, the review showed indications of beneficial effects of drug-eluting stents compared with bare-metal stents on target lesion revascularisation, target vessel revascularisation, and a composite outcome of cardiac events. They did not perform Trial Sequential Analysis or another method to assess the risk of random error, neither did they assess the strength of evidence for the alternative hypothesis (Jakobsen 2014).

We identified three previous reviews assessing the effects of drug-eluting stents versus bare-metal stents in STEMI participants. All three reviews confirmed our results showing that there is no difference between drug-eluting stents and bare-metal stents on all-cause mortality.

De Luca 2012 found no difference in terms of mortality at 440 days of follow-up (hazard ratio (HR) 0.85, 95% CI 0.70 to 1.04, $P = 0.11$). No difference in stent thrombosis was also observed. A significant reduction in target vessel revascularisation was found (HR 0.57, 95% CI 0.50 to 0.66, $P = 0.001$). They did not perform Trial Sequential Analysis or other method to assess the risk of random error, neither did they assess the strength of evidence for the alternative hypothesis (Jakobsen 2014).

Suh 2011 included 14 trials and found no evidence of an effect on mortality (RR 0.88, 95% CI 0.70 to 1.10, $P = 0.26$). Suh 2011 found evidence of an effect on myocardial infarction (RR 0.77, 95% CI 0.77 to 0.97, $P = 0.03$), but did not perform Trial Sequential Analysis or other method to assess the risk of random error, nor did they assess the strength of evidence for the alternative hypothesis (Jakobsen 2014).

Zheng 2014 found no difference in terms of mortality in a meta-analysis of four trials all with a minimum follow-up of four years

(odds ratio (OR) 0.85, 95% CI 0.59 to 1.20, $P = 0.35$) or stent thrombosis (OR 3.03, 95% CI 1.28 to 7.18, $P = 0.01$). Additionally, Zheng 2014 also found a reduction in target vessel revascularisation (OR 0.55, 95% CI 0.55 to 0.77). They did not perform Trial Sequential Analysis or other method to assess the risk of random error, neither did they assess the strength of evidence for the alternative hypothesis (Jakobsen 2014).

AUTHORS' CONCLUSIONS

Implications for practice

The current evidence suggests that drug-eluting stents may lead to fewer serious adverse events compared with bare-metal stents. When assessing each specific type of event included in the serious adverse event outcome separately, we found that the majority of events were target vessel revascularisation. The reduction in serious adverse events came without increasing the risk of all-cause mortality or major cardiovascular events. Our Trial Sequential Analysis showed that there was currently not enough information to assess a risk ratio reduction of 10% for all-cause mortality, major cardiovascular events, cardiovascular mortality, or myocardial infarction, and there were no data on quality of life or angina. The evidence for the above outcomes was of low to very low quality, and the true result may depart substantially from the results presented in this systematic review.

Implications for research

Our Trial Sequential Analysis showed that there was currently not enough information to assess a risk ratio reduction of 10% for all-cause mortality, major cardiovascular events, cardiovascular mortality, and myocardial infarction. We therefore recommend that future randomised clinical trials assessing these outcomes be conducted, as currently the superiority of drug-eluting stents relies on a possible reduced risk of target vessel revascularisation. In addition, we found no trials reporting quality of life or angina, therefore future trials should also assess these outcomes. Furthermore, trials conducted up to now have not comprehensively reported serious adverse events, and future trials should avoid continuing this practice. Future trials should be conducted with low risk of systematic error and low risk of random errors. Such trials ought to be designed and reported according to the SPIRIT (Standard Protocol Items: Recommendations for Interventional Trials) and CONSORT guidelines (Chan 2013; Schulz 2010).

There is a need for systematic reviews assessing serious adverse events in quasi-randomised and observational studies.

ACKNOWLEDGEMENTS

We thank Professor Rumona Dickson for her work on the protocol and the previous version of the review.

We thank Jørn Wetterslev for his work on the protocol.

We thank Cochrane Heart for the provision of a template for this protocol.

REFERENCES

References to studies included in this review

COMFORTABLE 2012 {published data only}

Magro M, Raber L, Heg D, Taniwaki M, Kelbaek H, Ostojic M, et al. The MI SYNTAX score for risk stratification in patients undergoing primary percutaneous coronary intervention for treatment of acute myocardial infarction: a substudy of the COMFORTABLE AMI trial. *International Journal of Cardiology* 2014;**175**(2):314-22.

Raber L, Kelbaek H, Baumbach A, Tuller D, Ostojic M, Juni P, et al. Long-term clinical outcomes of biolimus-eluting stents with biodegradable versus bare-metal stents in patients with acute STEMI: 5 year results of the randomized COMFORTABLE AMI trial. *European Heart Journal* 2016;**37**:35.

Raber L, Kelbaek H, Ostojic M, Baumbach A, Tuller D, von Birgelen C, et al. Comparison of biolimus eluted from an erodible stent coating with bare metal stents in acute ST-elevation myocardial infarction (COMFORTABLE AMI trial): rationale and design. *EuroIntervention* 2012;**7**(12):1435-43. [PMID: 22301368]

Raber L, Kelbaek H, Taniwaki M, Ostojic M, Heg D, Baumbach A, et al. Biolimus-eluting stents with biodegradable polymer versus bare-metal stents in acute myocardial infarction: two-year clinical results of the COMFORTABLE AMI trial. *Circulation: Cardiovascular Interventions* 2014;**7**(3):355-64.

* Raber L, Ostojic M. Effect of biolimus-eluting stents with biodegradable polymer vs bare-metal stents on cardiovascular events among patients with acute myocardial infarction: The COMFORTABLE AMI randomized trial. *JAMA* 2012;**308**(8):777-87.

Darkahian 2014 {published data only}

Darkahian M, Peighambari M. Comparison of the mid-term outcome between drug-eluting stent and bare metal stent implantation in patients undergoing primary PCI in Rajaie Heart Center January 2012 - April 2013. *Iranian Heart Journal* 2014;**15**(2):12-9.

DEB-AMI 2012 {published data only}

Belkacemi A, Agostoni P, Nathoe H, Shao C, Sangiorgi G, Voskuil M, et al. Results of the drug-eluting balloon in acute STEMI (DEB-AMI) trial: a multicenter randomised comparison of drug-eluting balloon plus bare-metal stent versus bare-metal stent versus DES in primary PCI with 6-month angiographic, intravascular, functional and 12-month clinical outcomes. *EuroIntervention* 2012;**8**(Suppl N):N25.

* Belkacemi A, Agostoni P, Nathoe HM, Voskuil M, Shao C, Van Belle E, et al. First results of the DEB-AMI (drug eluting balloon in acute ST-segment elevation myocardial infarction) trial: a multicenter randomized comparison of drug-eluting balloon plus bare-metal stent versus bare-metal stent versus drug-eluting stent in primary percutaneous coronary intervention with 6-month angiographic, intravascular, functional, and clinical outcomes. *Journal of the American College of Cardiology* 2012;**59**(25):2327-37. [PMID: 22503057]

DEBATER 2012 {published data only}

* Wijnbergen I, Helmes H, Tijssen J, Brueren G, Peels K, van Dantzig JM, et al. Comparison of drug-eluting and bare-metal stents for primary percutaneous coronary intervention with or without abciximab in ST-segment elevation myocardial infarction: DEBATER: the Eindhoven reperfusion study. *JACC: Cardiovascular Interventions* 2012;**5**(3):313-22.

Wijnbergen I, Tijssen J, Brueren G, Peels K, van Dantzig JM, Veer MV, et al. Long-term comparison of sirolimus-eluting and bare-metal stents in ST-segment elevation myocardial infarction. *Coronary Artery Disease* 2014;**25**(5):378-83.

DEDICATION 2008 {published data only}

Holmvang L, Kelbaek H, Kaltoft A, Thuesen L, Lassen JF, Clemmensen P, et al. Long-term outcome after drug-eluting versus bare-metal stent implantation in patients with ST-segment elevation myocardial infarction: 5 years follow-up from the randomized DEDICATION trial (Drug Elution and Distal Protection in Acute Myocardial Infarction). *JACC: Cardiovascular Interventions* 2013;**6**(6):548-53.

Kaltoft A, Kelbaek H, Clemmensen P, Helqvist S, Lassen JF, Klovgaard L, et al. Long-term outcome after drug-eluting versus bare metal stent implantation in patients with ST-segment elevation myocardial infarction: 3-year follow-up of the DEDICATION trial. *EuroIntervention* 2010;**6**(Suppl H):H132.

Kaltoft A, Kelbaek H, Thuesen L, Lassen JF, Clemmensen P, Klovgaard L, et al. Long-term outcome after drug-eluting versus bare-metal stent implantation in patients with ST-segment elevation myocardial infarction: 3-year follow-up of the randomized DEDICATION (Drug Elution and Distal Protection in Acute Myocardial Infarction) trial. *Journal of the American College of Cardiology* 2010;**56**(8):641-5.

* Kelbaek H, Thuesen L, Helqvist S, Clemmensen P, Klovgaard L, Kaltoft A, et al. Drug-eluting versus bare metal stents in patients with ST-segment-elevation myocardial infarction - eight-month follow-up in the drug elution and distal protection in acute myocardial infarction (DEDICATION) trial. *Circulation* 2008;**118**(11):1155-62.

De Ribamar Costa 2012 {published data only}

De Ribamar Costa J, Siqueira D, Abizaid A, Chamie D, Costa R, Viana R, et al. Serial grey-scale and radiofrequency intravascular assessment of plaque modification and vessel geometry at proximal and distal edges of bare-metal and drug-eluting stents. *Journal of the American College of Cardiology* 2012;**1**:E141.

DEVINE 2007 {published data only}

Pitt JE, Reeve R, Watkin H, Whitlam G, Pulikal J, Ment N, et al. Drug eluting versus bare metal stents in acute ST elevation myocardial infarction (DEVINE) - a randomised control trial. *European Heart Journal* 2007;**28**(1 Suppl):206.

Díaz 2007 {published data only}

Díaz de la Llera LS, Ballesteros S, Nevado J, Fernández M, Villa M, Sánchez A, et al. Sirolimus-eluting stents compared

with standard stents in the treatment of patients with primary angioplasty. *American Heart Journal* 2007;**154**(1):164.e1-6.

ELISA-3 2016 {published data only}

Remkes WS, Badings EA, Hermanides RS, Rasoul S, Dambrink JHE, Koopmans PC, et al. Randomised comparison of drug-eluting versus bare-metal stenting in patients with non-ST elevation myocardial infarction. *Open Heart* 2016;**3**(2):e000455.

EXAMINATION 2012 {published data only}

Brugaletta S, Sabate M, Cequier A, Iniguez A, Serra A, Hernandez-Antolin R, et al. Impact of everolimus-eluting stents on stent thrombosis as compared to conventional bare metal stents in patients with ST-segment elevation myocardial infarction. Insights from the EXAMINATION trial. *European Heart Journal* 2012;**33**:175.

* Sabate M, Cequier A, Iniguez A, Serra A, Hernandez-Antolin R, Mainar V, et al. Everolimus-eluting stent versus bare-metal stent in ST-segment elevation myocardial infarction (EXAMINATION): 1 year results of a randomised controlled trial. *Lancet* 2012;**380**(9852):1482-90. [PMID: 22951305]

Sabate M, Cequier A, Iniguez A, Serra A, Hernandez-Antolin R, Mainar V, et al. Rationale and design of the EXAMINATION trial: a randomised comparison between everolimus-eluting stents and cobalt-chromium bare-metal stents in ST-elevation myocardial infarction. *EuroIntervention* 2011;**7**(8):977-84.

FIBISTEMI 2007 {published data only}

Gao H, Yan HB, Zhu XL, Li N, Ai H, Wang J, et al. Firebird sirolimus eluting stent versus bare metal stent in patients with ST-segment elevation myocardial infarction. *Chinese Medical Journal* 2007;**120**(10):863-7.

GRACIA-3 2010 {published data only}

Sanchez PL, Gimeno F, Ancillo P, Sanz JJ, Alonso-Briaes JH, Bosa F, et al. Role of the paclitaxel-eluting stent and tirofiban in patients with ST-elevation myocardial infarction undergoing postfibrinolysis angioplasty: the GRACIA-3 randomized clinical trial. *Circulation* 2010;**3**(4):297-307.

HAAMU-STENT 2006 {published data only}

Tierala I, Syvanne M, Kupari M. Randomised comparison of a paclitaxel-eluting and a bare metal stent in STEMI-PCI. *American Journal of Cardiology - TCT abstracts* 2006;**98**(Meeting abstract):78M.

HORIZONS-AMI 2009 {published data only}

Mehran R, Brodie B, Cox DA, Grines CL, Rutherford B, Bhatt DL, et al. The Harmonizing Outcomes with Revascularization and Stents in Acute Myocardial Infarction (HORIZONS-AMI) Trial: study design and rationale. *American Heart Journal* 2008;**156**(1):44-56. [PMID: 18585496]

* Stone GW, Lansky AJ, Pocock SJ, Gersh BJ, Dangas G, Wong SC, et al. Paclitaxel-eluting stents versus bare-metal stents in acute myocardial infarction. *New England Journal of Medicine* 2009;**360**(19):1946-59. [DOI: 10.1056/NEJMoa0810116] [PMID: 19420364]

Stone GW, Parise H, Witzenbichler B, Kirtane A, Guagliumi G, Peruga JZ, et al. Selection criteria for drug-eluting versus bare-metal stents and the impact of routine angiographic follow-up: 2-year insights from the HORIZONS-AMI (Harmonizing Outcomes With Revascularization and Stents in Acute Myocardial Infarction) trial. *Journal of the American College of Cardiology* 2010;**56**(19):1597-604.

Stone GW, Witzenbichler B, Guagliumi G, Peruga JZ, Brodie BR, Dudek D, et al. Heparin plus a glycoprotein IIb/IIIa inhibitor versus bivalirudin monotherapy and paclitaxel-eluting stents versus bare-metal stents in acute myocardial infarction (HORIZONS-AMI): final 3-year results from a multicentre, randomised controlled trial. *Lancet* 2011;**377**(9784):2193-204. [PMID: 21665265]

König 2007 {published data only}

König A, Leibig M, Rieber J, Schiele TM, Theisen K, Siebert U, et al. Randomized comparison of dexamethasone-eluting stents with bare metal stent implantation in patients with acute coronary syndrome: serial angiographic and sonographic analysis. *American Heart Journal* 2007;**153**(6):979.e1-8.

Mission 2008 {published data only}

Atary JZ, van der Hoeven BL, Liem SS, Jukema JW, van der Bom JG, Atsma DE, et al. Three-year outcome of sirolimus-eluting versus bare-metal stents for the treatment of ST-segment elevation myocardial infarction (from the MISSION! Intervention Study). *American Journal of Cardiology* 2010;**106**(1):4-12.

Boden H, Van Der Hoeven BL, Atary JZ, Liem SS, Van Der Wall EE, Schalij MJ. 5-year outcomes of the MISSION intervention study: a randomized comparison of sirolimus-eluting to bare-metal stent implantation in patients with STEMI. *European Heart Journal* 2011;**32**:654.

Boden H, van der Hoeven BL, Liem SS, Atary JZ, Cannegieter SC, Atsma DE, et al. Five-year clinical follow-up from the MISSION! Intervention Study: sirolimus-eluting stent versus bare metal stent implantation in patients with ST-segment elevation myocardial infarction, a randomised controlled trial. *EuroIntervention* 2012;**7**(9):1021-9.

* van der Hoeven BL, Liem S, Jukema JW, Suraphakdee N, Putter H, Dijkstra J, et al. Sirolimus-eluting stents versus bare-metal stents in patients with ST-segment elevation myocardial infarction: 9-month angiographic and intravascular ultrasound results and 12-month clinical outcome results from the MISSION! intervention study. *Journal of the American College of Cardiology* 2008;**51**(6):618-26.

MULTISTRATEGY 2008 {published data only}

Valgimigli M, Campo G, Gambetti S, Bolognese L, Ribichini F, Colangelo S, et al. Three-year follow-up of the MULTICentre evaluation of Single high-dose Bolus Tirofiban versus Abciximab with Sirolimus-eluting STent or Bare-Metal Stent in Acute Myocardial Infarction Study (MULTISTRATEGY). *International Journal of Cardiology* 2013;**165**(1):134-41. [PMID: 21864917]

* Valgimigli M, Campo G, Percoco G, Bolognese L, Vassanelli C, Colangelo S, et al. Comparison of angioplasty with infusion

of tirofiban or abciximab and with implantation of sirolimus-eluting or uncoated stents. The MULTISTRATEGY Randomized Trial. *JAMA* 2008;**299**(15):1788-99. [PMID: 18375998]

OCTAMI 2010 {published data only}

Guagliumi G, Sirbu V, Bezerra H, Biondi-Zoccai G, Fiocca L, Musumeci G, et al. Strut coverage and vessel wall response to zotarolimus-eluting and bare-metal stents implanted in patients with ST-segment elevation myocardial infarction: the OCTAMI (Optical Coherence Tomography in Acute Myocardial Infarction) Study. *JACC: Cardiovascular Interventions* 2010;**3**(6):680-7.

PASEO 2009a {published data only}

* Di Lorenzo E, De Luca G, Sauro R, Varricchio A, Capasso M, Lanzillo T, et al. The PASEO (paclitaxel- or sirolimus-eluting stent versus bare metal stent in primary angioplasty) randomized trial. *JACC: Cardiovascular Interventions* 2009;**2**(6):515-23.

Di Lorenzo E, Sauro R, Varricchio A, Capasso M, Lanzillo T, Manganelli F, et al. Benefits of drug-eluting stents as compared to bare metal stent in ST-segment elevation myocardial infarction: four year results of the paclitaxel or sirolimus-eluting stent vs bare metal stent in primary angioplasty (PASEO) randomized trial. *American Heart Journal* 2009;**158**(4):e43-50.

PASEO 2009b {published data only}

PASSION 2006 {published data only}

Dirksen M, Vink M, Suttorp M, Tijssen J, Patterson M, Slagboom T, et al. Two year follow up after primary PCI with a paclitaxel-eluting stent versus a bare-metal stent for acute ST-elevation myocardial infarction (the PASSION trial): a follow-up study. *EuroIntervention* 2009;**4**:64-70.

Laarman GJ, Suttorp MJ, Dirksen MT, van Heerebeek L, Kiemeneij F, Slagboom T, et al. Paclitaxel-eluting versus uncoated stents in primary percutaneous coronary intervention. *New England Journal of Medicine* 2006;**355**(11):1105-13.

* Vink MA, Dirksen MT, Suttorp MJ, Tijssen JG, van Etten J, Patterson MS, et al. 5-year follow-up after primary percutaneous coronary intervention with a paclitaxel-eluting stent versus a bare-metal stent in acute ST-segment elevation myocardial infarction: a follow-up study of the PASSION (Paclitaxel-Eluting Versus Conventional Stent in Myocardial Infarction with ST-Segment Elevation) trial. *JACC: Cardiovascular Interventions* 2011;**4**(1):24-9.

SELECTION 2007 {published data only}

Cechi T, Vittori G, Biondi Zoccai GGL, Vecchio S, Falchetti E, Spaziani G. Single-center randomized evaluation of paclitaxel-eluting versus conventional stent in acute myocardial infarction (SELECTION). *Journal of Interventional Cardiology* 2007;**20**(4):282-91.

SESAMI 2007 {published data only}

Menichelli M, Parma A, Pucci E, Fiorilli R, De Felice F, Nazzaro M, et al. Randomized trial of sirolimus-eluting stent versus bare-

metal stent in acute myocardial infarction (SESAMI). *Journal of the American College of Cardiology* 2007;**49**(19):1924-30.

Musto C, Fiorilli R, De Felice F, Patti G, Nazzaro M, Scappaticci M, et al. Long-term outcome of sirolimus-eluting versus bare-metal stent in the setting of acute myocardial infarction: 5-year results of the SESAMI trial. *Giornale Italiano di Cardiologia* 2011;**12**(Suppl 1):4S.

* Violini R, Musto C, De Felice F, Nazzaro MS, Cifarelli A, Petitti T, et al. Maintenance of long-term clinical benefit with sirolimus-eluting stents in patients with ST-segment elevation myocardial infarction: 3-year results of the SESAMI (sirolimus-eluting stent versus bare-metal stent in acute myocardial infarction) trial. *Journal of the American College of Cardiology* 2010;**55**(8):810-4.

Steinwender 2008 {published data only}

Steinwender C, Hofmann R, Kypta A, Kammler J, Kerschner K, Grund M, et al. In-stent restenosis in bare metal stents versus sirolimus-eluting stents after primary coronary intervention for acute myocardial infarction and subsequent transcatheter transplantation of autologous stem cells. *Clinical Cardiology* 2008;**31**(8):356-9.

Strozzi 2007 {published data only}

Strozzi M, Anic D. Comparison of stent graft, sirolimus stent, and bare metal stent implanted in patients with acute coronary syndrome: clinical and angiographic follow up. *Croatian Medical Journal* 2007;**48**(3):348-52.

Typhoon 2006 {published data only}

Spaulding C, Henry P, Teiger E, Beatt K, Bramucci E, Carrie D, et al. Sirolimus-eluting versus uncoated stents in acute myocardial infarction. *New England Journal of Medicine* 2006;**355**(11):1093-104.

* Spaulding C, Teiger E, Commeau P, Varenne O, Bramucci E, Slama M, et al. Four-year follow-up of TYPHOON (trial to assess the use of the CYPHer sirolimus-eluting coronary stent in acute myocardial infarction treated with Balloon angioplasty). *JACC: Cardiovascular Interventions* 2011;**4**(1):14-23.

References to studies excluded from this review

ACUITY 2006 {published data only}

Stone GW, McLaurin BT, Cox DA, Bertrand ME, Lincoff AM, Moses JW, et al. Bivalirudin for patients with acute coronary syndromes. *New England Journal of Medicine* 2006;**355**:2203-16.

BARRICADE 2011 {published data only}

Stone GW, Goldberg S, O'Shaughnessy C, Midei M, Siegel RM, Cristea E, et al. 5-year follow-up of polytetrafluoroethylene-covered stents compared with bare-metal stents in aortocoronary saphenous vein grafts: the randomized BARRICADE (barrier approach to restenosis: restrict intima to curtail adverse events) trial. *JACC: Cardiovascular Interventions* 2012;**4**(3):300-9.

DESSOLVE-1 2011 {published data only}

Sakamoto K, Waseda K, Yock PG, Honda Y, Wijns W, Ormiston J, et al. Arterial response to sirolimus eluting stents with

bioabsorbable polymer: First IVUS Report from the DESSOLVE-I FIM Trial. *Journal of the American College of Cardiology* 2011;**58**(20, Suppl B):B32.

Gioia 2006 {published data only}

Gioia G, Matthai W, Benassi A, Rana H, Levite HA, Ewing LG. Improved survival with drug-eluting stent implantation in comparison with bare metal stent in patients with severe left ventricular dysfunction. *Catheterization & Cardiovascular Interventions* 2006;**68**:392-8.

Halkin 2006 {published data only}

Halkin A, Selzer F, Marroquin O, Laskey W, Detre K, Cohen H. Clinical outcomes following percutaneous coronary intervention with drug-eluting vs. bare-metal stents in dialysis patients. *Journal of Invasive Cardiology* 2006;**18**:577-83.

Han 2006 {published data only}

* Han SH, Ahn TH, Kang WC, Oh KJ, Chung WJ, Shin MS, et al. The favorable clinical and angiographic outcomes of a high-dose dexamethasone-eluting stent: randomized controlled prospective study. *American Heart Journal* 2006;**152**(5):887.e1-7.

Park YM, Han SH, Shin KC, Kim MG, Lee KH, Kang WC, et al. The favorable long term clinical outcomes of a high-dose dexamethasone-eluting stent: randomized controlled prospective study. *Journal of the American College of Cardiology* 2011;**1**:E1699.

Hausleiter 2005 {published data only}

Hausleiter J, Kastrati A, Wessely R, Dibra A, Mehilli J, Schratzenstaller T, et al. Prevention of restenosis by a novel drug-eluting stent system with a dose-adjustable, polymer-free, on-site stent coating. *European Heart Journal* 2005;**26**:1475-81.

Hoffmann 2007 {published data only}

Hoffmann R, Stellbrink E, Schroder J, Grawe A, Vogel G, Blindt R, et al. Impact of the metabolic syndrome on angiographic and clinical events after coronary intervention using bare-metal or sirolimus-eluting stents. *American Journal of Cardiology* 2007;**100**:1347-52.

Hokimoto 2015 {published data only}

Hokimoto S, Mizuno Y, Sueta D, Morita S, Akasaka T, Tabata N, et al. High incidence of coronary spasm after percutaneous coronary interventions: comparison between new generation drug-eluting stent and bare-metal stent. *International Journal of Cardiology* 2015;**182**:171-3.

Ishii 2012 {published data only}

Ishii H, Toriyama T, Aoyama T, Takahashi H, Tanaka M, Yoshikawa D, et al. Percutaneous coronary intervention with bare metal stent vs. drug-eluting stent in hemodialysis patients. *Circulation Journal* 2012;**76**(7):1609-15.

JACK-EPC 2013 {published data only}

Wojakowski W, Pyrlík A, Król M, Buszman P, Ochala A, Milewski K, et al. Circulating endothelial progenitor cells are inversely correlated with in-stent restenosis in patients with non-ST-segment elevation acute coronary syndromes

treated with EPC-capture stents (JACK-EPC trial). *Minerva Cardioangiologica* 2013;**61**:301-11.

Kim 2006 {published data only}

Kim Y-H, Park S-W, Lee CW, Hong M-K, Gwon H-C, Jang Y, et al. Comparison of sirolimus-eluting stent, paclitaxel-eluting stent, and bare metal stent in the treatment of long coronary lesions. *Catheterization & Cardiovascular Interventions* 2006;**67**:181-7.

Lansky 2000 {published data only}

Lansky AJ, Roubin GS, O'Shaughnessy CD, Moore PB, Dean LS, Raizner AE, et al. Randomized comparison of GR-II stent and Palmaz-Schatz stent for elective treatment of coronary stenoses. *Circulation* 2000;**102**(12):1364-8.

Lasave 2007 {published data only}

Lasave LI, de Ribamar Costa J Jr, Abizaid AA, Feres F, Tanajura LF, Staico R, et al. A three-dimensional intravascular ultrasound comparison between the new zotarolimus-eluting stent (ZotMaxx) and the non-drug-eluting TriMaxx stent. *Journal of Invasive Cardiology* 2007;**19**:303-8.

MASTER 2012 {published data only}

Costa JR Jr, Abizaid A, Dudek D, Silber S, Leon MB, Stone GW. Rationale and design of the MGuard for acute ST elevation reperfusion MASTER trial. *Catheterization & Cardiovascular Interventions* 2013;**82**(2):184-90.

Stone GW, Abizaid A, Silber S, Dizon JM, Merkely B, Costa RA, et al. Prospective, randomized, multicenter evaluation of a polyethylene terephthalate micronet mesh-covered stent (MGuard) in ST-segment elevation myocardial infarction: The MASTER Trial. *Journal of the American College of Cardiology* 2012;**60**(19):1975-84.

Meredith 2007 {published data only}

Meredith IT, Ormiston J, Whitbourn R, Kay IP, Muller D, Popma JJ, et al. Four-year clinical follow-up after implantation of the Endeavor zotarolimus-eluting stent: ENDEAVOR I, the first-in-human study. *American Journal of Cardiology* 2007;**100**:56M-61M.

Oyabu 2006 {published data only}

Oyabu J, Ueda Y, Ogasawara N, Okada K, Hirayama A, Kodama K. Angioscopic evaluation of neointima coverage: sirolimus drug-eluting stent versus bare metal stent. *American Heart Journal* 2006;**152**:1168-74.

PATENCY 2007 {published data only}

Heldman A, Farhat N, Fry E, Cummins F, Roubin G, McGarvey J, et al. Paclitaxel-eluting stents for cytostatic prevention of restenosis: patency study follow-up. *American Journal of Cardiology* 2002;**1**(6 (S1)):H1-4.

PISCES 2005 {published data only}

Serruys PW, Sianos G, Abizaid A, Aoki J, den Heijer P, Bonnier H, et al. The effect of variable dose and release kinetics on neointimal hyperplasia using a novel paclitaxel-eluting stent platform: the Paclitaxel In-Stent Controlled Elution Study (PISCES). *Journal of the American College of Cardiology* 2005;**46**(2):253-60.

Strategy 2005 {published data only}

Valgimigli M, Percoco G, Malagutti P, Campo G, Ferrari F, Barbieri D, et al. Tirofiban and sirolimus-eluting stent vs abciximab and bare-metal stent for acute myocardial infarction: a randomized trial. *JAMA* 2005;**293**(17):2109-17. [PMID: 15870414]

Suzuki 2013 {published data only}

Suzuki S, Ishii H, Matsudaira K, Okumura N, Yoshikawa D, Hayashi M, et al. Long-term outcome of drug-eluting vs. bare-metal stents in patients with acute myocardial infarction. Subgroup analysis of the Nagoya Acute Myocardial Infarction Study (NAMIS). *Circulation Journal* 2013;**77**(8):2024-31.

TRUST 2003 {published data only}

Hamm CW, Hugenholtz PG, Trust Investigators. Silicon carbide-coated stents in patients with acute coronary syndrome. *Catheterization & Cardiovascular Interventions* 2003;**60**(3):375-81.

Turan 1998 {published data only}

Turan F, Degertekin M, Gencbay M, Basaran Y, Yilmaz H, Dindar I, et al. Heparin-coated stent implantation in patients with unstable angina pectoris: a randomized trial. *European Heart Journal* 1998;**19**:284.

VESTASYNC 2014 {published data only}

Costa JR Jr, Oliveira BA, Abizaid A, Costa R, Perin M, Abizaid A, et al. Clinical, angiographic, and intravascular ultrasound results of the VestSync II trial. *Catheterization & Cardiovascular Interventions* 2014;**84**(7):1073-9. [PMID: 23460415]

Windecker 2005 {published data only}

Windecker S, Simon R, Lins M, Klauss V, Eberli FR, Roffi M, et al. Randomized comparison of a titanium-nitride-oxide-coated stent with a stainless steel stent for coronary revascularization - The TiNOX trial. *Circulation* 2005;**111**:2617-22.

XAMI 2013 {published data only}

Hofma SH, Smits PC, Van't Hof A, Velders MA, Van Boven AJ. Low event rates at long-term follow-up in the randomized myocardial infarction XAMI trial comparing first and second generation drug eluting stents. *Journal of the American College of Cardiology* 2013;**1**:B78.

References to studies awaiting assessment

BASKET 2005a {published data only}

Kaiser C, Brunner-La Rocca H, Buser P, Bonetti P, Osswald S, Linka A, et al. Incremental cost-effectiveness of drug-eluting stents compared with a third-generation bare-metal stent in a real world setting: Randomised Basel Stent Kosten Effektivitäts Trial (BASKET). *Lancet* 2005;**366**:921-9.

BASKET 2005b {published data only}

BASKET-PROVE I 2010 {published data only}

* Kaiser C, Galatius S, Erne P, Eberli F, Alber H, Rickli H, et al. Drug-eluting versus bare-metal stents in large coronary arteries.

New England Journal of Medicine 2010;**363**(24):2310-9. [DOI: 10.1056/NEJMoa1009406] [PMID: 21080780]

BASKET-PROVE II 2015 {published data only}

* Kaiser C, Galatius S, Jeger R, Gilgen N, Skov Jensen J, Naber CK, et al. Long-term efficacy and safety of biodegradable-polymer biolimus-eluting stents: main results of the Basel Stent Kosten-Effektivitäts Trial-PROspective Validation Examination II (BASKET-PROVE II), a randomized, controlled noninferiority 2-year outcome trial. *Circulation* 2015;**131**(1):74-81. [DOI: 10.1161/CIRCULATIONAHA.114.013520] [PMID: 25411159]

EAGLE 2006 {published data only}

Ischinger T. European study of Axxion and Glycocalix long-term evaluation. *Herz* 2006;**31**:596.

Erglis 2007 {published data only}

Erglis A, Narbutė I, Kumsars I, Jegere S, Mintale I, Zakke I, et al. A randomized comparison of paclitaxel-eluting stents versus bare-metal stents for treatment of unprotected left main coronary artery stenosis. *Journal of the American College of Cardiology* 2007;**50**(6):491-7.

FUTURE I 2004 {published data only}

Costa RA, Lansky AJ, Mintz GS, Mehran R, Tsuchiya Y, Negoita M, et al. Angiographic results of the first human experience with everolimus-eluting stents for the treatment of coronary lesions (the FUTURE I trial). *American Journal of Cardiology* 2005;**95**(1):113-6.

* Grube E, Sonoda S, Ikeno F, Honda Y, Kar S, Chan C, et al. Six-and twelve-month results from first human experience using everolimus-eluting stents with bioabsorbable polymer. *Circulation* 2004;**109**:2168-71.

FUTURE II 2004 {published data only}

Storger H, Grube E, Hofmann M, Schwartz F, Haase J. Clinical experiences using everolimus-eluting stents in patients with coronary artery disease. *Journal of Interventional Cardiology* 2004;**17**(6):387-90.

Han 2007 {published data only}

Han YL, Wang SL, Jing QM, Yu HB, Wang B, Ma YY, et al. Midterm outcomes of prospective, randomized, single-center study of the Janus tacrolimus-eluting stent for treatment of native coronary artery lesions. *Chinese Medical Journal* 2007;**120**(7):552-6.

Li 2004 {published data only}

Li S, Fu X, Liu J, Wu W, Gu X, Ma N, et al. Randomized study to evaluate sirolimus-eluting stents implanted at coronary small vessel lesions. *American Journal of Cardiology* 2004;**94**(6A):218E.

NORSTENT 2016 {published data only}

Bønnaa KH, Mannsverk J, Wiseth R, Aaberge L, Myreng Y, Nygård O, et al. Drug-eluting or bare-metal stents for coronary artery disease. *New England Journal of Medicine* 2016;**375**:1242-52.

PRODIGY 2014 {published data only}

Valgimigli M, Campo G, Percoco G, Monti M, Ferrari F, Tumscitz C, et al. Randomized comparison of 6- versus 24-month clopidogrel therapy after balancing anti-intimal hyperplasia stent potency in all-comer patients undergoing percutaneous coronary intervention: Design and rationale for the PROlonging Dual-antiplatelet treatment after Grading stent-induced Intimal hyperplasia study (PRODIGY). *American Heart Journal* 2010;**160**(5):804-11. [PMID: 21095265]

* Valgimigli M, Tebaldi M, Borghesi M, Branckx P, Campo G, Tumscitz C, et al. Two-year outcomes after first- or second-generation drug-eluting or bare-metal stent implantation in all-comer patients undergoing percutaneous coronary intervention. *JACC: Cardiovascular Interventions* 2014;**7**(1):20-8. [PMID: 24332420]

SES-SMART 2004 {published data only}

Ardisino D, Cavallini C, Bramucci E, Indolfi C, Marzocchi A, Manari A, et al. Sirolimus-eluting vs uncoated stents for prevention of restenosis in small coronary arteries: a randomized trial. *JAMA* 2004;**292**(22):2727-34. [PMID: 15585732]

XIMA 2014 {published data only}

de Belder A, de la Torre Hernandez JM, Lopez-Palop R, O'Kane P, Hernandez FH, Strange J, et al. A prospective randomized trial of everolimus-eluting stents versus bare-metal stents in octogenarians. *Journal of the American College of Cardiology* 2014;**63**(14):1371-5. [DOI: [10.1016/j.jacc.2013.10.053](https://doi.org/10.1016/j.jacc.2013.10.053)] [PMID: 24216285]

ZEUS 2015 {published data only}

Valgimigli M, Patialiakas A, Thury A, Colangelo S, Campo G, Tebaldi M, et al. Randomized comparison of zotarolimus-eluting Endeavor Sprint versus bare-metal stent implantation in uncertain drug-eluting stent candidates: rationale, design, and characterization of the patient population for the Zotarolimus-Eluting Endeavor Sprint Stent in Uncertain DES Candidates study. *American Heart Journal* 2013;**166**(5):831-8. [PMID: 24176438]

* Valgimigli M, Patialiakas A, Thury A, McFadden E, Colangelo S, Campo G, et al. Zotarolimus-eluting versus bare-metal stents in uncertain drug-eluting stent candidates. *Journal of the American College of Cardiology* 2015;**65**(8):805-15. [PMID: 25720624]

Additional references

Abizaid 2010

Abizaid A, Costa JR Jr. New drug-eluting stents: an overview on biodegradable and polymer-free next-generation stent systems. *Circulation: Cardiovascular Interventions* 2010;**3**(4):384-93.

ACEP 2013

American College of Emergency Physicians (ACEP), Society for Cardiovascular Angiography and Interventions, O'Gara PT, Kushner FG, Ascheim DD, Casey DE Jr, Chung MK, de Lemos JA, et al. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction: a report of the American College of Cardiology Foundation/American Heart Association

Task Force on Practice Guidelines. *Journal of the American College of Cardiology* 2013;**61**(4):e78-140.

Amsterdam 2014

Amsterdam EA, Wenger NK, Brindis RG, Casey DE Jr, Ganiats TG, Holmes DR Jr, et al. 2014 AHA/ACC guideline for the management of patients with non-ST-elevation acute coronary syndromes: executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* 2014;**130**(25):2354-94.

Baim 1996

Baim DS, Grossman W. Complications of cardiac catheterization. In: Baim DS, Grossman W, editors(s). *Cardiac Catheterization, Angiography and Intervention*. Williams & Wilkins, 1996:17.

Braunwald 1998

Braunwald E. Unstable angina: an etiologic approach to management. *Circulation* 1998;**98**(21):2219-22.

Breall 2016

Breall JA, Aroesty JM, Simons M. Overview of the non-acute management of unstable angina and non-ST elevation myocardial infarction. www.uptodate.com/contents/overview-of-the-non-acute-management-of-unstable-angina-and-non-st-elevation-myocardial-infarction (accessed 19 November 2015).

Brieger 2009

Brieger D, Fox KA, Fitzgerald G, Eagle KA, Budaj A, Avezum A, et al. Predicting freedom from clinical events in non-ST-elevation acute coronary syndromes: the Global Registry of Acute Coronary Events. *Heart* 2009;**95**(11):888-94.

Brok 2008

Brok J, Thorlund K, Gluud C, Wetterslev J. Trial sequential analysis reveals insufficient information size and potentially false positive results in many meta-analyses. *Journal of Clinical Epidemiology* 2008;**61**(8):763-9.

Brok 2009

Brok J, Thorlund K, Wetterslev J, Gluud C. Apparently conclusive meta-analyses may be inconclusive. Trial sequential analysis adjustment of random error risk due to repetitive testing of accumulating data in apparently conclusive neonatal meta-analyses. *International Journal of Epidemiology* 2009;**38**(1):287-98.

Cantor 1998

Cantor WJ, Lazzam C, Cohen EA, Bowman KA, Dolman S, Mackie K, et al. Failed coronary stent deployment. *American Heart Journal* 1998;**136**(6):1088-95.

Cantor 2005

Cantor WJ, Puley G, Natarajan MK, Dzavik V, Madan M, Fry A, et al. Radial versus femoral access for emergent percutaneous coronary intervention with adjunct glycoprotein IIb/IIIa inhibition in acute myocardial infarction - the RADIAL-AMI pilot randomized trial. *American Heart Journal* 2005;**150**(3):543-9. [PMID: 16169338]

Castellini 2016

Castellini G, Nielsen EE, Gluud C. Comment on: "Cell therapy for heart disease: Trial sequential analyses of two Cochrane reviews". *Clinical Pharmacology and Therapeutics* 2016 Oct 11 [Epub ahead of print]. [DOI: [10.1002/cpt.536](https://doi.org/10.1002/cpt.536)]

Chan 2013

Chan A-W, Tetzlaff JM, Altman DG, Laupacis A, Gøtzsche PC, Krleža-Jerić K, et al. SPIRIT 2013 Statement: Defining standard protocol items for clinical trials. *Annals of Internal Medicine* 2013;**158**(3):200-7.

Chen 2015

Chen W, Habraken TC, Hennink WE, Kok RJ. Polymer-free drug-eluting stents: an overview of coating strategies and comparison with polymer-coated drug-eluting stents. *Bioconjugate Chemistry* 2015;**26**(7):1277-88. [PMID: 26041505]

Cooper 2000

Cooper R, Cutler J, Desvigne-Nickens P, Fortmann SP, Friedman L, Havlik R, et al. Trends and disparities in coronary heart disease, stroke, and other cardiovascular diseases in the United States: findings of the national conference on cardiovascular disease prevention. *Circulation* 2000;**102**(25):3137-47. [PMID: 11120707]

CTU 2011 [Computer program]

Trial Sequential Analysis (TSA) (www.ctu.dk/tsa). Copenhagen Trial Unit (CTU), Centre for Clinical Intervention Research. Copenhagen, Denmark, 2011.

Cutlip 2002

Cutlip DE, Chauhan MS, Baim DS, Ho KK, Popma JJ, Carrozza JP, et al. Clinical restenosis after coronary stenting: perspectives from multicenter clinical trials. *Journal of the American College of Cardiology* 2002;**40**(12):2082-9.

Cutlip 2007

Cutlip DE, Windecker S, Mehran R, Boam A, Cohen DJ, van Es GA, et al. Clinical end points in coronary stent trials: a case for standardized definitions. *Circulation* 2007;**115**(17):2344-51. [PMID: 17470709]

Dangas 2002

Dangas G, Kuepper F. Restenosis: repeat narrowing of a coronary artery. *Circulation* 2002;**105**(22):2586.

Degertekin 2002

Degertekin M, Regar E, Tanabe K, Lee CH, Serruys PW. Sirolimus eluting stent in the treatment of atherosclerosis coronary artery disease. *Minerva Cardioangiologica* 2002;**50**(5):405-18. [PMID: 12384623]

De Luca 2012

De Luca G, Dirksen MT, Spaulding C, Kelbaek H, Schalić M, Thuesen L, et al. Drug-eluting vs bare-metal stents in primary angioplasty: a pooled patient-level meta-analysis of randomized trials. *Archives of Internal Medicine* 2012;**172**(8):611-21.

Demets 1987

Demets DL. Methods for combining randomized clinical trials: strengths and limitations. *Statistics in Medicine* 1987;**6**(3):341-50. [PMID: 3616287]

DerSimonian 1986

DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled Clinical Trials* 1986;**7**(3):177-88. [PMID: 3802833]

Düber 1986

Düber C, Jungbluth A, Rumpelt HJ, Erbel R, Meyer J, Thoenes W. Morphology of the coronary arteries after combined thrombolysis and percutaneous transluminal coronary angioplasty for acute myocardial infarction. *American Journal of Cardiology* 1986;**58**(9):698-703.

Dudas 2013

Dudas K, Björck L, Jernberg T, Lappas G, Wallentin L, Rosengren A. Differences between acute myocardial infarction and unstable angina: a longitudinal cohort study reporting findings from the Register of Information and Knowledge about Swedish Heart Intensive Care Admissions (RIKS-HIA). *BMJ Open* 2013;**3**(1):e002155.

Duval 2000

Duval S, Tweedie R. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* 2000;**56**:455-63.

Erbel 1998

Erbel R, Haude M, Höpp HW, Franzen D, Rupprecht HJ, Heublein B, et al. Coronary-artery stenting compared with balloon angioplasty for restenosis after initial balloon angioplasty. Restenosis Stent Study Group. *New England Journal of Medicine* 1998;**339**(23):1672-8.

Fajadet 2006

Fajadet J, Wijns W, Laarman GJ, Kuck KH, Ormiston J, Münzel T, et al. Randomized, double-blind, multicenter study of the Endeavor zotarolimus-eluting phosphorylcholine-encapsulated stent for treatment of native coronary artery lesions: clinical and angiographic results of the ENDEAVOR II trial. *Circulation* 2006;**114**(8):798-806. [PMID: 16908773]

Farooq 2013

Farooq V, van Klaveren D, Steyerberg EW, Meliga E, Vergouwe Y, Chieffo A, et al. Anatomical and clinical characteristics to guide decision making between coronary artery bypass surgery and percutaneous coronary intervention for individual patients: development and validation of SYNTAX score II. *Lancet* 2013;**381**(9867):639-50.

Fischman 1994

Fischman DL, Leon MB, Baim DS, Schatz RA, Savage MP, Penn I, et al. A randomized comparison of coronary-stent placement and balloon angioplasty in the treatment of coronary artery disease. Stent Restenosis Study Investigators. *New England Journal of Medicine* 1994;**331**(8):496-501. [PMID: 8041414]

Garattini 2016

Garattini S, Jakobsen JC, Wetterslev J, Bertele V, Banzi R, Rath A, et al. Evidence-based clinical practice: overview of threats to the validity of evidence and how to minimise them. *European Journal of Internal Medicine* 2016;**32**:13-21.

Gensini 1983

Gensini GG. A more meaningful scoring system for determining the severity of coronary heart disease. *American Journal of Cardiology* 1983;**51**(3):606. [PMID: 6823874]

Gluud 2006

Gluud LL. Bias in clinical intervention research. *American Journal of Epidemiology* 2006;**163**(6):493-501.

Greenhalgh 2010

Greenhalgh J, Hockenhull J, Rao N, Dundar Y, Dickson RC, Bagust A. Drug-eluting stents versus bare metal stents for angina or acute coronary syndromes. *Cochrane Database of Systematic Reviews* 2010, Issue 5. Art. No: CD004587. [DOI: [10.1002/14651858.CD004587.pub3](https://doi.org/10.1002/14651858.CD004587.pub3)]

Grüntzig 1978

Grüntzig A. Transluminal dilatation of coronary-artery stenosis. *Lancet* 1978;**1**(8058):263. [PMID: 74678]

Grüntzig 1979

Grüntzig AR, Senning A, Siegenthaler WE. Nonoperative dilatation of coronary-artery stenosis: percutaneous transluminal coronary angioplasty. *New England Journal of Medicine* 1979;**301**(2):61-8. [PMID: 449946]

Gupta 2003

Gupta M, Chang WC, Van de Werf F, Granger CB, Midodzi W, Barbash G, et al. International differences in in-hospital revascularization and outcomes following acute myocardial infarction. *European Heart Journal* 2003;**24**(18):1640-50.

Guyatt 2008

Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;**336**(7650):924-6. [PMID: 18436948]

Hamon 2009

Hamon M, Rasmussen LH, Manoukian SV, Cequier A, Lincoff MA, Rupprecht HJ, et al. Choice of arterial access site and outcomes in patients with acute coronary syndromes managed with an early invasive strategy: the ACUITY trial. *EuroIntervention* 2009;**5**(1):115-20. [PMID: 19577992]

Harbord 2006

Harbord RM, Egger M, Sterne JA. A modified test for small-study effects in meta-analyses of controlled trials with binary endpoints. *Statistics in Medicine* 2006;**25**(20):3443-57. [PMID: 16345038]

Haude 2013

Haude M, Erbel R, Erne P, Verheye S, Degen H, Böse D, et al. Safety and performance of the drug-eluting absorbable metal scaffold (DREAMS) in patients with de-novo coronary lesions:

12 month results of the prospective, multicentre, first-in-man BIOSOLVE-I trial. *Lancet* 2013;**381**(9869):836-44.

Haude 2016

Haude M, Ince H, Abizaid A, Toelg R, Lemos PA, von Birgelen C, et al. Safety and performance of the second-generation drug-eluting absorbable metal scaffold in patients with de-novo coronary artery lesions (BIOSOLVE-II): 6 month results of a prospective, multicentre, non-randomised, first-in-man trial. *Lancet* 2016;**387**(10013):31-9.

Hicks 2010

Hicks KA, Hung JHM, Mahaffey KW, Mehran R, Nissen SE, Stockbridge NL, et al. Standardized definitions for end point events in cardiovascular trials. www.clinpage.com/images/uploads/endpoint-defs_11-16-2010.pdf (accessed 24 June 2016).

Higgins 2002

Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine* 2002;**21**(11):1539-58.

Higgins 2003

Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327**(7414):557-60.

Higgins 2011

Higgins JP, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Higgins 2011a

Higgins JP, Whitehead A, Simmonds M. Sequential methods for random-effects meta-analysis. *Statistics in Medicine* 2011;**30**(9):903-21. [PMID: 21472757]

Hoenig 2010

Hoenig MR, Aroney CN, Scott IA. Early invasive versus conservative strategies for unstable angina and non-ST elevation myocardial infarction in the stent era. *Cochrane Database of Systematic Reviews* 2010, Issue 3. Art. No: CD004815. [DOI: [10.1002/14651858.CD004815.pub4](https://doi.org/10.1002/14651858.CD004815.pub4)]

Holmes 2004

Holmes DR Jr, Leon MB, Moses JW, Popma JJ, Cutlip D, Fitzgerald PJ, et al. Analysis of 1-year clinical outcomes in the SIRIUS trial: a randomized trial of a sirolimus-eluting stent versus a standard stent in patients at high risk for coronary restenosis. *Circulation* 2004;**109**(5):634-40. [PMID: 14769686]

Hrobjartsson 2012

Hrobjartsson A, Thomsen AS, Emanuelsson F, Tendal B, Hilden J, Boutron I, et al. Observer bias in randomised clinical trials with binary outcomes: systematic review of trials with both blinded and non-blinded outcome assessors. *BMJ (Clinical Research Ed)* 2012;**344**:e1119.

Hrobjartsson 2013

Hrobjartsson A, Thomsen ASS, Emanuelsson F, Tendal B, Hilden J, Boutron I, et al. Observer bias in randomized clinical

trials with measurement scale outcomes: a systematic review of trials with both blinded and nonblinded assessors. *CMAJ: Canadian Medical Association Journal* 2013;**185**(4):E201-11.

Hrobjartsson 2014

Hrobjartsson A, Thomsen AS, Emanuelsson F, Tendal B, Rasmussen JV, Hilden J, et al. Observer bias in randomized clinical trials with time-to-event outcomes: systematic review of trials with both blinded and non-blinded outcome assessors. *International Journal of Epidemiology* 2014;**43**(3):937-48.

Hrobjartsson 2014a

Hrobjartsson A, Emanuelsson F, Skou Thomsen AS, Hilden J, Brorson S. Bias due to lack of patient blinding in clinical trials. A systematic review of trials randomizing patients to blind and nonblind sub-studies. *International Journal of Epidemiology* 2014;**43**(4):1272-83.

ICH-GCP 1997

International Conference on Harmonisation Expert Working Group (ICH-EWG). International Conference on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use. ICH Harmonised Tripartite Guideline. Guideline for Good Clinical Practice in CFR & ICH Guidelines. Vol. 1. Barnett International/PAREXEL, 1997.

Ioannidis 2009

Ioannidis JP. Adverse events in randomized trials: neglected, restricted, distorted, and silenced. *Archives of Internal Medicine* 2009;**169**(19):1737-9.

Jackson 2017

Jackson D, Turner R. Power analysis for random-effects meta-analysis. *Research Synthesis Methods* 2017 Apr 4 [Epub ahead of print]. [DOI: [10.1002/jrsm.1240](https://doi.org/10.1002/jrsm.1240)] [PMID: 28378395]

Jakobsen 2014

Jakobsen JC, Wetterslev J, Winkel P, Lange T, Gluud C. Thresholds for statistical and clinical significance in systematic reviews with meta-analytic methods. *BMC Medical Research Methodology* 2014;**14**:120. [PMID: 25416419]

Jakobsen 2016

Jakobsen JC, Wetterslev J, Lange T, Gluud C. Viewpoint: taking into account risks of random errors when analysing multiple outcomes in systematic reviews. *The Cochrane Database of Systematic Reviews* 2016;**3**:ED000111. [DOI: [10.1002/14651858.ED000111](https://doi.org/10.1002/14651858.ED000111)]

Kastrati 2007

Kastrati A, Mehilli J, Pache J, Kaiser C, Valgimigli M, Kelbaek H, et al. Analysis of 14 trials comparing sirolimus-eluting stents with bare-metal stents. *New England Journal of Medicine* 2007;**356**(10):1030-9. [PMID: 17296823]

Keus 2010

Keus F, Wetterslev J, Gluud C, van Laarhoven CJ. Evidence at a glance: error matrix approach for over-viewing available evidence. *BMC Medical Research Methodology* 2010;**10**:90.

Kip 2008

Kip KE, Hollabaugh K, Marroquin OC, Williams DO. The problem with composite end points in cardiovascular studies: the story of major adverse cardiac events and percutaneous coronary intervention. *Journal of the American College of Cardiology* 2008;**51**(7):701-7. [PMID: 18279733]

Kjaergard 2001

Kjaergard LL, Villumsen J, Gluud C. Reported methodologic quality and discrepancies between large and small randomized trials in meta-analyses. *Annals of Internal Medicine* 2001;**135**(11):982-9.

Lee 2005

Lee MS, Shah AP, Aragon J, Jamali A, Dohad S, Kar S, et al. Drug-eluting stenting is superior to bare metal stenting in saphenous vein grafts. *Catheterization & Cardiovascular Interventions* 2005;**66**(4):507-11. [PMID: 16270361]

Lefebvre 2011

Lefebvre C, Manheimer E, Glanville J. Chapter 6: Searching for studies. In: Higgins JP, Green S, editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Levin 2016

Levin T, Aroesty JM. Stable ischemic heart disease: indications for revascularization. www.uptodate.com/contents/stable-ischemic-heart-disease-indications-for-revascularization (accessed 10 September 2015).

Levine 2011

Levine GN, Bates ER, Blankenship JC, Bailey SR, Bittl JA, Cercek B, et al. 2011 ACCF/AHA/SCAI guideline for percutaneous coronary intervention: executive summary: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Society for Cardiovascular Angiography and Interventions. *Circulation* 2011;**124**(23):2574-609. [PMID: 22064598]

Levine 2016

Levine GN, Bates ER, Blankenship JC, Bailey SR, Bittl JA, Cercek B, et al. 2015 ACC/AHA/SCAI focused update on primary percutaneous coronary intervention for patients with ST-elevation myocardial infarction: an update of the 2011 ACCF/AHA/SCAI guideline for percutaneous coronary intervention and the 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction. *Journal of the American College of Cardiology* 2016;**67**(10):1235-50.

Lloyd-Jones 2010

Lloyd-Jones D, Adams RJ, Brown TM, Carnethon M, Dai S, De Simone G, et al. Executive summary: heart disease and stroke statistics - 2010 update: a report from the American Heart Association. *Circulation* 2010;**121**(7):948-54. [PMID: 20177011]

Lundh 2012

Lundh A, Sismondo S, Lexchin J, Busuioac OA, Bero L. Industry sponsorship and research outcome. *Cochrane Database of*

Systematic Reviews 2012, Issue 12. Art. No: MR000033. [DOI: [10.1002/14651858.MR000033.pub2](https://doi.org/10.1002/14651858.MR000033.pub2)]

Lundh 2017

Lundh A, Bero L. The ties that bind. *BMJ* 2017;**356**:j176.

Macaya 1996

Macaya C, Serruys PW, Ruygrok P, Suryapranata H, Mast G, Klugmann S, et al. Continued benefit of coronary stenting versus balloon angioplasty: one-year clinical follow-up of Benestent trial. Benestent Study Group. *Journal of the American College of Cardiology* 1996;**27**(2):255-61. [PMID: 8557891]

Maddox 2007

Maddox TM, Reid KJ, Rumsfeld JS, Spertus JA. One-year health status outcomes of unstable angina versus myocardial infarction: a prospective, observational cohort study of ACS survivors. *BMC Cardiovascular Disorders* 2007;**7**:28.

Meyer 1981

Meyer J, Schmitz H, Erbel R, Kiesslich T, Böcker-Josephs B, Krebs W, et al. Treatment of unstable angina pectoris with percutaneous transluminal coronary angioplasty (PTCA). *Catheterization and Cardiovascular Diagnosis* 1981;**7**(4):361-71.

Meyer 1982

Meyer J, Merx W, Dörr R, Lambertz H, Bethge C, Effert S. Successful treatment of acute myocardial infarction shock by combined percutaneous transluminal coronary recanalization (PTCR) and percutaneous transluminal coronary angioplasty (PTCA). *American Heart Journal* 1982;**103**(1):132-4.

Meyer 1982a

Meyer J, Merx W, Schmitz H, Erbel R, Kiesslich T, Dörr R, et al. Percutaneous transluminal coronary angioplasty immediately after intracoronary streptolysis of transmural myocardial infarction. *Circulation* 1982;**66**(5):905-13.

Meyer 1983

Meyer J, Schmitz HJ, Kiesslich T, Erbel R, Krebs W, Schulz W, et al. Percutaneous transluminal coronary angioplasty in patients with stable and unstable angina pectoris: analysis of early and late results. *American Heart Journal* 1983;**106**(5 Pt 1):973-80.

Moher 1998

Moher D, Pham B, Jones A, Cook DJ, Jadad AR, Moher M, et al. Does quality of reports of randomised trials affect estimates of intervention efficacy reported in meta-analyses? *Lancet* 1998;**352**(9128):609-13.

Morice 2002

Morice MC, Serruys PW, Sousa JE, Fajadet J, Ban Hayashi E, Perin M, et al. A randomized comparison of a sirolimus-eluting stent with a standard stent for coronary revascularization. *New England Journal of Medicine* 2002;**346**(23):1773-80. [PMID: 12050336]

Moses 2003

Moses JW, Leon MB, Popma JJ, Fitzgerald PJ, Holmes DR, O'Shaughnessy C, et al. Sirolimus-eluting stents versus standard stents in patients with stenosis in a native coronary artery.

New England Journal of Medicine 2003;**349**(14):1315-23. [PMID: 14523139]

NICE 2008

NICE (National Institute for Health and Care Excellence). Drug-eluting stents for the treatment of coronary artery disease. www.nice.org.uk/guidance/ta152?unlid=1066690929201610722145 (accessed prior to 12 February 2016).

Nichols 2014

Nichols M, Townsend N, Scarborough P, Rayner M. Cardiovascular disease in Europe 2014: epidemiological update. *European Heart Journal* 2014;**35**(42):2929. [PMID: 25139896]

Nordmann 2005

Nordmann AJ, Bucher HC, Hengstler P, Harr T, Young J. Primary stenting versus primary balloon angioplasty for treating acute myocardial infarction. *Cochrane Database of Systematic Reviews* 2005, Issue 2. Art. No: CD005313. [DOI: [10.1002/14651858.CD005313](https://doi.org/10.1002/14651858.CD005313)]

Pocock 2015

Pocock SJ, Clayton TC, Stone GW. Design of major randomized trials: part 3 of a 4-part series on statistics for clinical trials. *Journal of the American College of Cardiology* 2015;**66**(24):2757-66.

Pogue 1997

Pogue JM, Yusuf S. Cumulating evidence from randomized trials: utilizing sequential monitoring boundaries for cumulative meta-analysis. *Controlled Clinical Trials* 1997;**18**(6):580-93; discussion 661-6. [PMID: 9408720]

Pohjola-Sintonen 1989

Pohjola-Sintonen S, Muller JE, Stone PH, Willich SN, Antman EM, Davis VG, et al. Ventricular septal and free wall rupture complicating acute myocardial infarction: experience in the Multicenter Investigation of Limitation of Infarct Size. *American Heart Journal* 1989;**117**(4):809-18.

Puel 1988

Puel J, Juilliere Y, Bertrand ME, Rickards AF, Sigwart U, Serruys PW. Early and late assessment of stenosis geometry after coronary arterial stenting. *American Journal of Cardiology* 1988;**61**(8):546-53.

Puricel 2015

Puricel S, Arroyo D, Corpataux N, Baeriswyl G, Lehmann S, Kallinikou Z, et al. Comparison of everolimus- and biolimus-eluting coronary stents with everolimus-eluting bioresorbable vascular scaffolds. *Journal of the American College of Cardiology* 2015;**65**(8):791-801.

RevMan 2014 [Computer program]

Nordic Cochrane Centre, The Cochrane Collaboration Review Manager 5 (RevMan 5). Version 5.3. Copenhagen: Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Ringqvist 1983

Ringqvist I, Fisher LD, Mock M, Davis KB, Wedel H, Chaitman BR, et al. Prognostic value of angiographic indices of coronary artery disease from the Coronary Artery Surgery Study (CASS). *Journal of Clinical Investigation* 1983;**71**(6):1854-66. [PMID: 6863543]

Roffi 2016

Roffi M, Patrono C, Collet JP, Mueller C, Valgimigli M, Andreotti F, et al. 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: Task Force for the Management of Acute Coronary Syndromes in Patients Presenting Without Persistent ST-Segment Elevation of the European Society of Cardiology (ESC). *European Heart Journal* 2016;**37**(3):267-315.

Rosamond 2008

Rosamond W, Flegal K, Furie K, Go A, Greenlund K, Haase N, et al. Heart disease and stroke statistics - 2008 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation* 2008;**117**(4):e25-146. [PMID: 18086926]

Roukoz 2009

Roukoz H, Bavry AA, Sarkees ML, Mood GR, Kumbhani DJ, Rabbat MG, et al. Comprehensive meta-analysis on drug-eluting stents versus bare-metal stents during extended follow-up. *American Journal of Medicine* 2009;**122**(6):581. [PMID: 19486720]

Rücker 2008

Rücker G, Schwarzer G, Carpenter J. Arcsine test for publication bias in meta-analyses with binary outcomes. *Statistics in Medicine* 2008;**27**(5):746-63. [PMID: 17592831]

Ruiz-Bailén 2008

Ruiz-Bailén M, Rucabado-Aguilar L, Castillo-Rivera AM, Expósito-Ruiz M, Morante-Valle A, Rodríguez-García JJ, et al. Cardiogenic shock in acute coronary syndrome in the Spanish population. *Medical Science Monitor* 2008;**14**(11):PH46-57.

Savović 2012

Savović J, Jones HE, Altman DG, Harris RJ, Jüni P, Pildal J, et al. Influence of reported study design characteristics on intervention effect estimates from randomized, controlled trials. *Annals of Internal Medicine* 2012;**157**(6):429-38.

Savović 2012a

Savović J, Jones H, Altman D, Harris R, Jüni P, Pildal J, et al. Influence of reported study design characteristics on intervention effect estimates from randomised controlled trials: combined analysis of meta-epidemiological studies. *Health Technology Assessment* 2012;**16**(35):1-82.

Schmidt 2012

Schmidt M, Jacobsen JB, Lash TL, Bøtker HE, Sørensen HT. 25 year trends in first time hospitalisation for acute myocardial infarction, subsequent short and long term mortality, and the prognostic impact of sex and comorbidity: a Danish nationwide cohort study. *BMJ* 2012;**344**:e356. [PMID: 22279115]

Schulz 1995

Schulz KF, Chalmers I, Hayes RJ, Altman DG. Empirical evidence of bias. Dimensions of methodological quality associated with estimates of treatment effects in controlled trials. *JAMA* 1995;**273**(5):408-12.

Schulz 2010

Schulz KF, Altman DG, Moher D, CONSORT Group. CONSORT 2010 Statement: Updated guidelines for reporting parallel group randomised trials. *Annals of Internal Medicine* 2010;**152**(11):726-732.

Seizer 1982

Seizer A. On the limitation of therapeutic intervention trials in ischemic heart disease: a clinician's viewpoint. *American Journal of Cardiology* 1982;**49**(1):252-5.

Serruys 1994

Serruys PW, de Jaegere P, Kiemeneij F, Macaya C, Rutsch W, Heyndrickx G, et al. A comparison of balloon-expandable-stent implantation with balloon angioplasty in patients with coronary artery disease. Benestent Study Group. *New England Journal of Medicine* 1994;**331**(8):489-95. [PMID: 8041413]

Serruys 1998

Serruys PW, van Hout B, Bonnier H, Legrand V, Garcia E, Macaya C, et al. Randomised comparison of implantation of heparin-coated stents with balloon angioplasty in selected patients with coronary artery disease (Benestent II). *Lancet* 1998;**352**(9129):673-81. [PMID: 9728982]

Serruys 2010

Serruys PW, Silber S, Garg S, van Geuns RJ, Richardt G, Buszman PE, et al. Comparison of zotarolimus-eluting and everolimus-eluting coronary stents. *New England Journal of Medicine* 2010;**363**(2):136-46. [PMID: 20554978]

Sianos 2005

Sianos G, Morel MA, Kappetein AP, Morice MC, Colombo A, Dawkins K, et al. The SYNTAX score: an angiographic tool grading the complexity of coronary artery disease. *EuroIntervention* 2005;**1**(2):219-27. [PMID: 19758907]

Skoog 2015

Skoog M, Saarimäki JM, Gluud C, Sheinin M, Erlendsson K, Aamdal S, et al. Transparency and registration in clinical research in the Nordic countries. The Nordic Trial Alliance Working Group on Transparency and Registration under NordForsk 2015.

Stankovic 2004

Stankovic G, Orlic D, Corvaja N, Airolidi F, Chieffo A, Spanos V, et al. Incidence, predictors, in-hospital, and late outcomes of coronary artery perforations. *American Journal of Cardiology* 2004;**93**(2):213-6.

Stata 2015 [Computer program]

Stata Statistical Software: Release 14. College Station, TX: StataCorp LP, 2015.

Sterne 2009

Sterne JA, White IR, Carlin JB, Spratt M, Royston P, Kenward MG, et al. Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls. *BMJ* 2009;**338**:b2393. [PMID: 19564179]

Stettler 2007

Stettler C, Wandel S, Allemann S, Kastrati A, Morice MC, Schömig A, et al. Outcomes associated with drug-eluting and bare-metal stents: a collaborative network meta-analysis. *Lancet* 2007;**370**(9591):937-48. [PMID: 17869634]

Stevenson 1989

Stevenson WG, Linssen GC, Havenith MG, Brugada P, Wellens HJ. The spectrum of death after myocardial infarction: a necropsy study. *American Heart Journal* 1989;**118**(6):1182-8.

Stone 2004

Stone GW, Ellis SG, Cox DA, Hermiller J, O'Shaughnessy C, Mann JT, et al. A polymer-based, paclitaxel-eluting stent in patients with coronary artery disease. *New England Journal of Medicine* 2004;**350**(3):221-31. [PMID: 14724301]

Stone 2004a

Stone GW, Ellis SG, Cox DA, Hermiller J, O'Shaughnessy C, Mann JT, et al. One-year clinical results with the slow-release, polymer-based, paclitaxel-eluting TAXUS stent: the TAXUS-IV trial. *Circulation* 2004;**109**(16):1942-7. [PMID: 15078803]

Stone 2010

Stone GW, Rizvi A, Newman W, Mastali K, Wang JC, Caputo R, et al. Everolimus-eluting versus paclitaxel-eluting stents in coronary artery disease. *New England Journal of Medicine* 2010;**362**(18):1663-74.

Suh 2011

Suh HS, Song HJ, Choi JE, Jang EJ, Son HJ, Lee SM, et al. Drug-eluting stents versus bare-metal stents in acute myocardial infarction: a systematic review and meta-analysis. *International Journal of Technology Assessment in Health Care* 2011;**27**(1):11-22. [PMID: 21262083]

Sweeting 2004

Sweeting MJ, Sutton AJ, Lambert PC. What to add to nothing? Use and avoidance of continuity corrections in meta-analysis of sparse data. *Statistics in Medicine* 2004;**23**(9):1351-75. [PMID: 15116347]

Tamai 2000

Tamai H, Igaki K, Kyo E, Kosuga K, Kawashima A, Matsui S, et al. Initial and 6-month results of biodegradable poly-L-lactic acid coronary stents in humans. *Circulation* 2000;**102**(4):399-404.

Taylor 1998

Taylor SJ, Tweedie RL. Practical estimates of the effect of publication bias in meta-analysis. *Australian Epidemiologist* 1998;**5**:14-7.

Thombs 2006

Thombs BD, Bass EB, Ford DE, Stewart KJ, Tsilidis KK, Patel U, et al. Prevalence of depression in survivors of acute myocardial infarction. *Journal of General Internal Medicine* 2006;**21**(1):30-8.

Thorlund 2009

Thorlund K, Devereaux PJ, Wetterslev J, Guyatt G, Ioannidis JP, Thabane L, et al. Can trial sequential monitoring boundaries reduce spurious inferences from meta-analyses? *International Journal of Epidemiology* 2009;**38**(1):276-86.

Thorlund 2011

Thorlund K, Engstrøm J, Wetterslev J, Brok J, Imberger G, Gluud C. User manual for Trial Sequential Analysis (TSA). www.ctu.dk/tsa/files/tsa_manual.pdf (accessed 15 May 2015).

Thygesen 2012

Thygesen K, Alpert JS, Jaffe AS, Simoons ML, Chaitman BR, White HD, Writing Group on behalf of the Joint ESC/ACCF/AHA/WHF Task Force for the Universal Definition of Myocardial Infarction. Third universal definition of myocardial infarction. *Global Heart* 2012;**7**(4):275-95. [PMID: 25689940]

Turner 2013

Turner RM, Bird SM, Higgins JP. The impact of study size on meta-analyses: examination of underpowered studies in Cochrane reviews. *PLoS ONE* 2013;**8**(3):e59202. [PMID: 23544056]

Turpie 2006

Turpie AG. Burden of disease: medical and economic impact of acute coronary syndromes. *American Journal of Managed Care* 2006;**12**(16 Suppl):S430-4.

Ware 1992

Ware JE Jr, Sherbourne CD. The MOS 36-item Short-Form Health Survey (SF-36). I. Conceptual framework and item selection. *Medical Care* 1992;**30**(6):473-83.

Wetterslev 2008

Wetterslev J, Thorlund K, Brok J, Gluud C. Trial sequential analysis may establish when firm evidence is reached in cumulative meta-analysis. *Journal of Clinical Epidemiology* 2008;**61**(1):64-75.

Wetterslev 2009

Wetterslev J, Thorlund K, Brok J, Gluud C. Estimating required information size by quantifying diversity in random-effects model meta-analyses. *BMC Medical Research Methodology* 2009;**9**:86.

Wetterslev 2017

Wetterslev J, Jakobsen JC, Gluud C. Trial Sequential Analysis in systematic reviews with meta-analysis. *BMC Medical Research Methodology* 2017;**17**(1):39.

White 2008

White HD, Chew DP. Acute myocardial infarction. *Lancet* 2008;**372**(9638):570-84.

WHO 2015

World Health Organization (WHO). Cardiovascular diseases. www.who.int/cardiovascular_diseases/en/ (accessed 27 January 2015).

Windecker 2014

Windecker S, Kolh P, Alfonso F, Collet JP, Cremer J, Falk V, et al. 2014 ESC/EACTS Guidelines on myocardial revascularization: The Task Force on Myocardial Revascularization of the European Society of Cardiology (ESC) and the European Association for Cardio-Thoracic Surgery (EACTS). Developed with the special contribution of the European Association of Percutaneous Cardiovascular Interventions (EAPCI). *European Heart Journal* 2014;**35**(37):2541-619.

Wood 2008

Wood L, Egger M, Gluud LL, Schulz KF, Jüni P, Altman DG, et al. Empirical evidence of bias in treatment effect estimates in controlled trials with different interventions and outcomes: meta-epidemiological study. *BMJ* 2008;**336**(7644):601-5.

Wyrwich 2004

Wyrwich KW, Spertus JA, Kroenke K, Tierney WM, Babu AN, Wolinsky FD, Heart Disease Expert Panel. Clinically important

differences in health status for patients with heart disease: an expert consensus panel report. *American Heart Journal* 2004;**147**(4):615-22.

Zheng 2014

Zheng F, Xing S, Gong Z, Xing Q. Five-year outcomes for first generation drug-eluting stents versus bare-metal stents in patients with ST-segment elevation myocardial infarction: a meta-analysis of randomised controlled trials. *Heart, Lung & Circulation* 2014;**23**(6):542-8. [PMID: 24529648]

References to other published versions of this review

Feinberg 2016

Feinberg J, Nielsen EE, Greenhalgh J, Hounscome J, Sethi NJ, Safi S, et al. Drug-eluting stents versus bare-metal stents for acute coronary syndrome. *Cochrane Database of Systematic Reviews* 2016, Issue 12. Art. No: CD012481. [DOI: [10.1002/14651858.CD012481](https://doi.org/10.1002/14651858.CD012481)]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

COMFORTABLE 2012

Study characteristics

Methods	Parallel-design randomised clinical trial in Switzerland, Denmark, Serbia, UK, the Netherlands, and Israel
Participants	<p>1161 participants with symptom onset within 24 hours and ST-segment elevation of at least 1 mm in 2 or more contiguous leads, true posterior MI, or new left bundle branch block were eligible for randomisation in the presence of at least 1 culprit lesion within the infarct vessel. There was no limit regarding the number of treated lesions, vessels, or complexity.</p> <p>Male:female = 918:243</p> <p>Mean age = 60.5 years</p> <p>Exclusion criteria: presence of mechanical complications of acute MI, known allergy to any study medication, use of vitamin K antagonists, planned surgery unless dual antiplatelet therapy could be maintained throughout the perisurgical period, history of bleeding diathesis or known coagulopathy, pregnancy, participation in another trial before reaching the primary endpoint, inability to provide informed consent, and non-cardiac comorbid conditions with life expectancy of less than 1 year.</p>
Interventions	<p>Experimental group: stents eluting Biolimus from a biodegradable polylactic acid polymer (BioMatrix, Biosensors Europe SA).</p> <p>Control group 1: bare-metal stents of otherwise identical design (Gazelle, Biosensors Europe SA).</p> <p>Co-intervention: acetylsalicylic acid (250 mg) was administered before the procedure. In centres where prasugrel was available, an initial dose of 60 mg (including participants preloaded with clopidogrel) was administered followed up with a 10 mg daily dose. If prasugrel was not available or contraindicated, clopidogrel was administered at a loading dose of 600 mg, followed up with a dose of 75 mg twice daily for 7 days, followed up with a maintenance dose of 75 mg once daily. Dual antiplatelet therapy was prescribed for a duration of at least 1 year in all participants. Unfractionated heparin was routinely</p>

Drug-eluting stents versus bare-metal stents for acute coronary syndrome (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

COMFORTABLE 2012 (Continued)

administered with a minimal dose of 5000 IU or a dose of 70 to 100 IU/kg to maintain an activated clotting time of 250 seconds. Bivalirudin was administered at a dose of 0.75 mg/kg intravenously followed up with an infusion of 1.75 mg/kg per hour during the duration of the procedure. The use of glycoprotein IIb/IIIa inhibitors was left to the discretion of the operator.

Outcomes	Death, myocardial infarction, biomarkers, TVR, TLR, device success, lesion success, procedural success, stent thrombosis, transient ischaemic attack, ischaemic stroke
Notes	NCT00962416 (prospectively)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated web-based system
Allocation concealment (selection bias)	Low risk	Computer-generated web-based system
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants and personnel were not blinded to the allocation.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	An independent data and safety monitoring board blinded to treatment groups periodically reviewed all event information and compared safety outcomes between groups.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Less than 5% dropped out, and dropout was judged to be equal in both groups.
Selective reporting (reporting bias)	Low risk	All outcomes stated in the pre-published protocol were reported.
Other bias	High risk	The trial was funded by the Swiss National Science Foundation (grant 33CM30-124112) and an unrestricted research grant from Biosensors Europe SA.

Darkahian 2014
Study characteristics

Methods	Parallel-design randomised clinical trial in Iran
Participants	384 participants with ST-segment elevation myocardial infarction. Male:female = not reported Mean age = not reported Exclusion criteria: none stated.
Interventions	Experimental group: drug-eluting stent Control group: bare-metal stent

Drug-eluting stents versus bare-metal stents for acute coronary syndrome (Review)

Darkahian 2014 (Continued)

Co-intervention: none stated.

Outcomes	Death, MI, TVR
Notes	Abstract only. We contacted the authors at mehdipei@gmail.com on 23 April 2017 regarding all bias domains, additional data, and request for full publication, but have received no reply.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	The number of dropouts was unclear.
Selective reporting (reporting bias)	Unclear risk	There was no pre-published protocol, and the trial did not report on our primary outcomes.
Other bias	Unclear risk	It was unclear how the trial was funded.

DEB-AMI 2012

Study characteristics

Methods	Parallel-design randomised clinical trial in the Netherlands and Italy
Participants	<p>100 participants presenting in the first 12 h after the onset of STEMI (diagnosed by the presence of anginal symptoms associated with electrocardiographic ST-segment elevation of 1 mm in 2 contiguous leads or new left bundle branch block) and undergoing primary PCI, with angiographic evidence of a single culprit lesion in the target vessel, after successful thrombus aspiration (defined by no angiographically evident flow-limiting residual thrombus at the site of [sub]occlusion, and Thrombolysis In Myocardial Infarction (TIMI) flow grade 1) were deemed eligible for inclusion.</p> <p>Male:female = 83:17</p> <p>Mean age = 57.9 years</p> <p>Exclusion criteria: contraindications to study medications (acetylsalicylic acid, clopidogrel, paclitaxel), life expectancy 12 months, lesion length 25 mm, reference vessel diameter 2.5 mm and 4.0 mm, severe triple vessel disease, left main stenosis 50%, and a combination of type C coronary lesion and diabetes mellitus (in which DES was favoured).</p>

DEB-AMI 2012 (Continued)

Interventions	<p>Experimental group: DES (Taxus Liberté, Boston Scientific, Natick, MA) as a stainless steel stent platform coated with a permanent polymer that allows the release of paclitaxel (1 g/mm²). The total strut thickness including stent and polymer is 132 µm.</p> <p>Control group: BMS (Genius MAGIC stent, Eurocor) cobalt chromium stent platform with a strut thickness of 60 µm.</p> <p>Co-intervention: pre-dilation with a standard balloon (balloon-to-artery ratio 0.8:1).</p> <p>All participants received routinely in the ambulance or at the first medical contact a loading dose of acetylsalicylic acid (325 to 500 mg) and of clopidogrel (600 mg). Heparin was administered before and during the procedure to maintain an activated clotting time 250 s. Additional administration of glycoprotein IIb/IIIa inhibitors was recommended, but was left to the physician's discretion.</p> <p>After fulfilling angiographic inclusion criteria, patients underwent mandated thrombus aspiration of the culprit lesion with a manual thrombus aspiration device.</p>	
Outcomes	Angiographic in-segment late luminal loss (expressed in millimetres) as determined by quantitative coronary analysis (QCA), binary restenosis using QCA, stent malapposition and re-endothelialisation assessed by optical coherence tomography, coronary endothelial dysfunction after acetylcholine infusion, and clinical outcomes (death, MI, repeated revascularisation procedures).	
Notes	<p>The trial had a third arm receiving paclitaxel drug-eluting balloon dilatation before BMS implantation. We were unable to include this arm due to multi-intervention.</p> <p>NCT00856765 - prospectively</p> <p>We contacted the authors at P.Stella@umcutrecht.nl on 21 March 2017 regarding blinding, dropouts, and additional data, but have received no reply.</p>	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Automatically generated by a computer
Allocation concealment (selection bias)	Low risk	Opaque, sealed envelopes
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Outcome assessors were only described as blinded for images (QCA analyses) and not for our primary outcomes.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	The number of dropouts was unclear.
Selective reporting (reporting bias)	Low risk	The authors reported on all outcomes described on ClinicalTrials.gov, and the outcomes were submitted before the trial was started.
Other bias	High risk	The trial was funded by Eurocor GmbH.

DEBATER 2012

Study characteristics

Methods	Parallel-design randomised clinical trial in the Netherlands
Participants	<p>907 patients 18 years of age or older with STEMI were eligible if they presented within 12 h of onset of symptoms.</p> <p>Male:female = 668:202</p> <p>Mean age = 60.5 years</p> <p>Exclusion criteria: patients who were on oral anticoagulation and patients who had received thrombolytic therapy or treatment with a glycoprotein IIb/IIIa inhibitor in the previous 24 h were not eligible. Other exclusion criteria were contraindications for DES, contraindications for clopidogrel or glycoprotein IIb/IIIa inhibitors, comorbid conditions with a predictable fatal outcome in the short run, cardiogenic shock, and inability to give informed consent.</p>
Interventions	<p>Experimental group: sirolimus-eluting stents (Cypher, Cordis Corporation, Bridgewater, NJ).</p> <p>Control group: BMS (type was left to the discretion of the operator).</p> <p>Co-intervention: all participants received aspirin (300 mg chewed or 500 mg intravenously), clopidogrel (600 mg), and a fixed bolus of intravenous unfractionated heparin (5000 IU) in the ambulance. Before angiography, all participants received an additional intravenous bolus of heparin (5000 IU). After primary PCI, aspirin 80 mg per day was given indefinitely, and clopidogrel was prescribed (75 mg/day) for at least 1 month after BMS and 6 to 12 months after sirolimus-eluting stents. Participants allocated to abciximab received a 0.25 mg/kg bolus after stent placement followed by an infusion of 0.125 g/kg/min for 12 h.</p>
Outcomes	Death, MI, revascularisation, stroke, bleeding
Notes	Participants included in this trial were randomised to another trial comparing abciximab with no abciximab (clinical trial number NCT00986050). We contacted the authors at nico.pijls@cze.nl on 23 April 2017 regarding allocation concealment, specific data on cardiovascular mortality, and additional data, but have received no reply.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated allocation
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants and personnel were not blinded.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	All events were adjudicated by a clinical endpoints committee under blinded conditions.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	More than 5% dropped out at 5 years, however not at 1-year follow-up (4.1%).

DEBATER 2012 (Continued)

Selective reporting (reporting bias)	High risk	The protocol was published retrospectively.
Other bias	High risk	The trial was funded by Johnson & Johnson (Cordis), Guidant, Abbott, and the Friends of the Heart Foundation.

DEDICATION 2008
Study characteristics

Methods	Randomised clinical trial with a factorial (2x2) design in Denmark
Participants	<p>626 participants with chest pain of 30-minute duration who had a cumulated ST-segment elevation of 4 mm in at least 2 contiguous leads of the ECG, provided that they were 18 years of age and had a high-grade stenosis or occlusion of a coronary artery without excessive tortuosity or calcification prohibiting advancement of a filter wire to the distal vascular bed of the vessel. We did not distinguish between patients who were admitted directly to the tertiary unit with laboratory facilities or via a referring hospital as long as they presented to the catheterisation laboratory within 12 hours from symptom onset.</p> <p>Male:female = 458:168</p> <p>Mean age = 62.2 years</p> <p>Exclusion criteria: previous MI in the target vessel area, development of cardiogenic shock before enrolment, culprit lesions in an unprotected left main coronary artery, gastrointestinal bleeding within 1 month, pregnancy, known renal failure, life expectancy 1 year, and linguistic problems.</p>
Interventions	<p>Experimental group: drug-eluting stents (47% were sirolimus-eluting (Cordis, Bridgewater, NJ), 40% were paclitaxel-eluting (Boston Scientific, Natick, MA), and 13% were zotarolimus-eluting stents (Medtronic, USA).</p> <p>Control group: BMS, 38% were made of cobalt alloy (Vision, Abbott and Driver, Medtronic); 39% were stainless steel stents from Boston Scientific; and 23% were miscellaneous stainless steel stents from Biotronik (Seoul, South Korea), Cordis, Guidant (Diegem, Belgium), Jomed (Helsingborg, Sweden), and Terumo (Tokyo, Japan).</p> <p>Co-intervention: participants were examined during and after the index procedure with ST-segment monitoring, cardiac markers, and echocardiography. At discharge, participants received a daily dose of a statin, clopidogrel (for 12 months), and aspirin (indefinitely). A beta-blocker was administered in the absence of contraindications, and an angiotensin-converting enzyme inhibitor was given in case of reduced (45%) left ventricular ejection fraction.</p>
Outcomes	ST-segment resolution, late loss (difference between minimal lumen diameter in stented segment immediately after and at 8 months' follow-up), restenosis, maximal elevations, wall motion index, minimal lumen diameter, frequency of binary restenosis, occurrence of stent thrombosis, and MACE
Notes	Clinical trial number NCT00192868. The participants in the trial were additionally randomised to distal protection or no distal protection, however this was done at random, and did not break the protocol for drug-eluting stents versus bare-metal stents.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computerised assignment

DEDICATION 2008 (Continued)

Allocation concealment (selection bias)	Low risk	Central telephone system
Blinding of participants and personnel (performance bias) All outcomes	High risk	Unblinded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The angiographic analyses were performed by blinded investigators, and the Clinical Events Committee adjudicated the serious events without knowledge of the treatment sequence.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	More than 5% dropped out on re-angiography.
Selective reporting (reporting bias)	High risk	The primary outcome were changed.
Other bias	High risk	The trial was funded by Cordis/Johnson & Johnson, Medtronic, Abbott, and Boston Scientific companies.

De Ribamar Costa 2012
Study characteristics

Methods	Parallel-design randomised clinical trial in Brazil
Participants	40 people with acute coronary syndrome. Male:female = unclear Mean age = unclear Exclusion criteria: not described.
Interventions	Experimental group: drug-eluting stent (Cypher). Control group 1: bare-metal stent (Driver). Co-intervention: not described.
Outcomes	Modification in vessel, lumen and plaque area and in the composition of the plaque in the mean time between the baseline and follow-up procedure
Notes	Abstract only. No contact information could be obtained.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described

De Ribamar Costa 2012 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	The number of dropouts was unclear.
Selective reporting (reporting bias)	Unclear risk	There was no pre-published protocol, and the trial did not report on our primary outcomes.
Other bias	Unclear risk	It was unclear how the trial was funded.

DEVINE 2007
Study characteristics

Methods	Parallel-design randomised clinical trial in the UK
Participants	167 participants presented with STEMI Male:female = not reported Mean age = 57.5 years Exclusion criteria: not reported.
Interventions	Experimental group: paclitaxel-eluting stents. Control group: bare-metal stent. Co-intervention: abciximab was used unless contraindicated.
Outcomes	TLR, stent thrombosis, mortality
Notes	Abstract only. Trial is ongoing. We were unable to locate data from the completed trial. Estimated recruitment was 250 participants. We were unable to locate contact information for the authors.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding of participants and personnel (performance bias)	Unclear risk	Not described

DEVINE 2007 (Continued)

All outcomes

Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	The number of dropouts was unclear.
Selective reporting (reporting bias)	Unclear risk	There was no pre-published protocol, and the trial did not report on our primary outcomes.
Other bias	Unclear risk	It was unclear how the trial was funded.

Diaz 2007
Study characteristics

Methods	Parallel-design randomised clinical trial in Spain
Participants	<p>120 participants with STEMI over the age of 18 years who were candidates for primary angioplasty and who met the following criteria were included: (1) chest pain lasting for > 30 minutes with elevation of the ST-segment by 1 mm or more on 2 or more contiguous electrocardiographic leads or recent-onset left branch blocking, and (2) admitted to the hospital centre within the first 12 hours after the onset of symptoms.</p> <p>Male:female = 95:19</p> <p>Mean age = 64.5 years</p> <p>Exclusion criteria: patients in a state of cardiogenic shock and/or Killip IV before randomisation; the partial or total administration of prior fibrinolytic treatment or administration of any glycoprotein IIb/IIIa inhibitors during the previous 30 days; chronic kidney failure requiring dialysis; pregnant women; history of haemorrhagic diathesis or allergy to aspirin, clopidogrel, and/or abciximab; major surgery in the last 15 days; active bleeding or previous stroke in the last 6 months; and a life expectancy of less than 6 months. Patients with a reference diameter < 2.25 mm and > 4.0 mm by visual estimation in the infarction-related artery were excluded from the study.</p>
Interventions	<p>Experimental group: drug-eluting stent and clopidogrel for at least 9 months.</p> <p>Control group: bare-metal stent and clopidogrel for at least 1 month.</p> <p>Co-intervention: before the procedure, all participants received aspirin (300 to 500 mg by mouth as the loading dose and then 100 mg a day indefinitely) plus clopidogrel (300 or 600 mg as the loading dose). Abciximab was administered to all participants as a bolus at a dose of 0.25 Ag/kg, followed by an infusion at a dose of 0.125 Ag/kg per minute for 12 hours. Heparin was administered as a bolus in relation to the participant's body weight at a dose of 70 U/kg (maximum 7000 U), with additional doses to maintain an activated clotting time of between 200 and 250 seconds.</p>
Outcomes	<p>Clinical follow-up was performed at 1, 3, 6, 9, and 12 months through visits to the clinic by the participant; the participant was also contacted afterward by phone.</p> <p>The primary endpoint is a composite of death, nonfatal MI, and recurrent myocardial ischaemia within a follow-up of 360 days after initial procedure. Participants with > 1 event will be assigned the highest-ranked event.</p>

Diaz 2007 (Continued)

Secondary endpoints include freedom, at day 30 and month 12, from death, recurrent MI, and clinically driven TVR, defined as any coronary artery bypass graft surgery or a second PCI of the original target vessel performed for recurrent myocardial ischaemia.

Notes

Under the protocol, those participants randomised to sirolimus-eluting stent plus abciximab could only receive BMS in the infarct-related artery if there had previously been a failure in the implanting of a sirolimus-eluting stent that prevented its repeated use due to the impossibility of reaching the implant area and if the use of BMS resolved this situation.

Contacted about allocation concealment, blinding, pre-published protocol, funding, complete number of stent thrombosis, cardiovascular death at 1 year, serious adverse events at 1 months and 1 year, data for participants not receiving PCI/stent by email: luissalvadordiaz@hotmail.com on 13 March 2017.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Randomization was performed with the aid of a basic computer program."
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	The trial was described as "double-blind", but it was unclear who was blinded and how blinding was maintained.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	The trial was described as "double-blind", but it was unclear who was blinded and how blinding was maintained.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	There was incomplete outcome data for above 5% of participants in the BMS group (6/60), and there was a difference in dropout rate (6/60 vs 0/60, BMS and DES respectively).
Selective reporting (reporting bias)	Unclear risk	There was no pre-published protocol, and the trial did not report on serious adverse events.
Other bias	Unclear risk	It was unclear how the trial was funded.

ELISA-3 2016

Study characteristics

Methods	Parallel-design randomised clinical trial in the Netherlands
Participants	<p>474 participants with NSTEMI hospitalised with ischaemic chest pain or dyspnoea at rest, with the last episode occurring 24 hours or less before randomisation, and had at least 2 of 3 of the following high-risk characteristics: (1) evidence of extensive myocardial ischaemia on ECG (shown by new cumulative ST depression > 5 mm or temporary ST-segment elevation in 2 contiguous leads < 30 min), (2) elevated biomarkers (troponin T > 0.10 µg/L or myoglobin > 150 µg/L) or elevated CK-MB fraction (> 6% of total CK), (3) age above 65 years from the ELISA-3 trial (178 participants). Additionally, NSTEMI participants who did not meet the inclusion criteria for high-risk NSTEMI from the ELISA-3 registry were included (296 participants).</p> <p>Male:female = 351:123</p>

ELISA-3 2016 (Continued)

Mean age = 65.2 years

Exclusion criteria: persistent ST-segment elevation, symptoms of ongoing myocardial ischaemia despite optimal medical therapy, contraindication for diagnostic angiography, active bleeding, cardiogenic shock, acute posterior infarction, and life expectancy < 1 year.

Interventions	Experimental group: everolimus-eluting stent. Control group: bare-metal stent. Co-intervention: dual antiplatelet therapy (acetylsalicylic acid and clopidogrel) for the duration of 1 year.
Outcomes	Angiographic for the participants randomised from the ELISA-3 study, MACE and stent thrombosis for all participants
Notes	ISRCTN Register (ISRCTN39230163) retrospectively registered. Study participants were taken from both the ELISA-3 study and the ELISA-3 registry of participants not included due to being high-risk NSTEMI patients. We contacted the authors at v.r.c.derks@isala.nl on the 23 March 2017 regarding random sequence generation and specific data on death, cardiac death, or myocardial infarction.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Low risk	Closed-envelope system
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Participants and operators were blinded.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Clinical endpoints were adjudicated by investigators blinded to participants' treatment allocation.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Less than 5% dropped out.
Selective reporting (reporting bias)	High risk	The protocol was retrospectively published.
Other bias	High risk	The trial was funded by Abbott.

EXAMINATION 2012

Study characteristics

Methods	Parallel-design randomised clinical trial in Spain, Italy, and the Netherlands
Participants	1504 participants presented with STEMI with the following electrocardiogram criteria: at least 1 mm in 2 or more standard leads or at least 2 mm in 2 or more contiguous precordial leads or left bundle

Drug-eluting stents versus bare-metal stents for acute coronary syndrome (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

EXAMINATION 2012 (Continued)

branch block that was not known to be old, within the first 48 h after the symptoms onset requiring emergent PCI with a vessel size ranging between 2.25 mm and 4.0 mm without other anatomical restrictions could be included.

Male:female = 1244:260

Mean age = 61.2 years

Exclusion criteria: age younger than 18 years; pregnancy; patients with known intolerance to aspirin, clopidogrel, heparin, stainless steel, everolimus or contrast material; patients on chronic treatment with anti-vitamin K agents, and STEMI secondary to stent thrombosis.

Interventions	<p>Experimental group: Xience V Multi-Link Vision everolimus-eluting stent.</p> <p>Control group 1: cobalt chromium Multi-Link Vision BMS.</p> <p>Co-intervention: at the index procedure, participants received appropriate anticoagulation and other therapy according to standard hospital practice. Either unfractionated heparin or bivalirudin might be used for procedural anticoagulation. The use of glycoprotein IIb/IIIa inhibitors was left to the discretion of the investigator. Aspirin (loading dose 250 to 500 mg) and clopidogrel (loading dose of at least 300 mg) had to be given before percutaneous coronary intervention for those participants not on chronic antiplatelet treatment. Neither prasugrel nor ticagrelor was approved during the recruitment period. Clopidogrel was prescribed for at least 1 year (75 mg per day) and aspirin (100 mg) indefinitely. Compliance to dual antiplatelet therapy refers to the concomitant use of both drugs during the prescription time of 1 year. Manual thrombectomy followed by direct stenting was the recommended technique in this setting, although other devices could also be used if considered necessary. Operators were instructed to use only the assigned stent type at the index procedure.</p>
Outcomes	Death, MI, TVR, TLR, stent thrombosis, procedural success, bleeding
Notes	NCT00828087 prospectively published.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated random numbers
Allocation concealment (selection bias)	Low risk	Centralised by telephone
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	The participants were blinded to the allocation, however the personnel were not.
Blinding of outcome assessment (detection bias) All outcomes	High risk	There was no description of blinding of outcome assessors.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Less than 5% dropped out.
Selective reporting (reporting bias)	Low risk	The protocol was pre-published, and all outcomes were reported.
Other bias	High risk	The trial was funded by Abbott Vascular to the Spanish Heart Foundation.

FIBISTEMI 2007

Study characteristics

Methods	Parallel-design randomised clinical trial in China
Participants	<p>156 participants with (1) onset of chest discomfort within the prior 12 hours and ST-segment elevation (at least 1 mm in 2 standard leads or at least 2 mm in 2 contiguous precordial leads) or new or presumably new left bundle branch block; (2) no upper limitation for age; (3) willing to comply with the follow-up schedule and procedures; (4) candidate for primary PCI; (5) no contraindications of anticoagulants and antiplatelets; (6) suitable angiographic anatomy for primary PCI.</p> <p>Male:female = 126:30</p> <p>Mean age = 58.7 years</p> <p>Exclusion criteria: (1) fever and cough during the past 48 hours; (2) severely calcified lesion unsuitable for balloon pre-dilation; (3) low left ventricular ejection fraction ($\leq 40\%$); (4) renal dysfunction (serum creatinine $> 103 \mu\text{mol/L}$); (5) life expectancy < 1 year; (6) intolerant or hypersensitive to sirolimus, aspirin, clopidogrel, heparin, stainless steel, or iodine-containing contrast medium; (7) participated in another research protocol; and (8) unsuitable for stent implantation.</p>
Interventions	<p>Experimental group: Firebird stent (Firebird stent group).</p> <p>Control group 1: Driver stent (cobalt-chromium alloy stent, Medtronic Ltd., USA) (BMS group).</p> <p>Co-intervention: all procedures were performed via a femoral or radial access using standard techniques. Oral loading dose of aspirin (300 mg) and clopidogrel (300 mg) were given immediately after establishment of STEMI diagnosis. Weight-adjusted heparin (8000 to 10,000 IU) was administered at the same time. Postprocedural antiplatelet regimen included 300 mg aspirin per day for 4 weeks and lifelong 100 mg per day, 75 mg clopidogrel per day for 9 to 12 months, and use of low molecular weight heparin for 1 week. Successful stent implantation was defined as $< 20\%$ residual stenosis in the stented segment with TIMI flow 3.</p>
Outcomes	In-stent and in-segment late luminal loss, binary angiographic restenosis, MACE, stent thrombosis
Notes	It was unclear how many participants were actually randomised since all participants received Firebird stent the last 4 months, therefore no data could be used in our meta-analyses. We contacted the authors on 21 March 2017 at yan591204@yahoo.com.cn for data on the participants randomised, sequence generation, allocation concealment, blinding, incomplete outcome data, whether they pre-published a protocol, and funding information.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not described
Blinding of outcome assessment (detection bias)	Unclear risk	Not described

FIBISTEMI 2007 (Continued)

All outcomes

Incomplete outcome data (attrition bias) All outcomes	Unclear risk	The number of dropouts was unclear.
Selective reporting (reporting bias)	Unclear risk	There was no pre-published protocol, and the trial did not report on our primary outcomes.
Other bias	Unclear risk	It was unclear how the trial was funded.

GRACIA-3 2010
Study characteristics

Methods	Parallel-design randomised clinical trial in Spain
Participants	<p>436 participants age 18 years or older; symptom onset within 12 hours before random assignment; chest pain lasting more than 30 minutes; ST-segment elevation of at least 0.1 mV in at least 2 limb leads, ST-segment elevation of at least 0.2 mV in 2 or more contiguous precordial leads, or left bundle branch block or paced rhythm; and no severe heart failure (Killip class 3).</p> <p>Male:female = 358:78</p> <p>Mean age = 61 years</p> <p>Exclusion criteria: cardiogenic shock, defined as a systolic blood pressure 90 mmHg with no response to fluid administration or 100 mmHg in patients with supportive treatment and no bradycardia; suspected mechanical complications of acute myocardial infarction; previous coronary artery bypass graft; non-cardiac disease that is likely to jeopardise the planned termination of the study; women of childbearing potential unless they had a negative pregnancy test result; active bleeding and recent surgery (within 2 weeks) contraindicating the use of heparin, tirofiban, or platelet aggregation inhibitors; contraindications for thrombolysis (previous haemorrhagic stroke at any time, history of non-haemorrhagic cerebrovascular accident within the previous 12 months, intracerebral neoplasm, active internal bleeding, suspected aortic dissection, uncontrolled hypertension 180/110 in several measurements, any other known intracerebral condition not covered in contraindications, current use of anti-coagulants or heparin use within 8 hours, known bleeding diathesis, recent trauma (4 weeks) including head trauma or traumatic or prolonged (10 minutes) cardiopulmonary resuscitation or recent major surgery or biopsy (8 weeks), non-compressible vascular punctures, recent (4 weeks) internal bleeding, pregnancy, and active peptic ulcer); history of hypersensitivity to aspirin, ticlopidine, clopidogrel, heparin, tirofiban, or stainless steel; known renal failure, creatinine 2.5 mg/dL; known impaired hepatic function that contraindicates the use of clopidogrel; known thrombocytopenia (100,000); participation in other trials; known multivessel disease identified as not suitable for revascularisation; and known peripheral vascular disease that makes cardiac catheterisation difficult.</p>
Interventions	<p>Experimental group: paclitaxel-eluting stents (Taxus stents; Boston Scientific, Natick, MA).</p> <p>Control group: bare-metal stents (Express stents; Boston Scientific).</p> <p>Co-intervention: participants underwent catheterisation within 3 to 12 hours of random assignment. The infarct-related artery was dilated if there was total occlusion, the stenosis was greater than 50%, or the TIMI flow grade was less than 3. Stenting of culprit lesions was performed when they were morphologically suitable for stenting and when the procedure was expected to achieve an adequate result. To avoid thrombus compression and embolisation, direct stenting was attempted when possible. Immediately after stenting, participants received a loading dose of 300 mg of clopidogrel and 75 mg daily for 1 year. Maintenance aspirin therapy was administered indefinitely at 80 to 325 mg once daily (coated or uncoated), unless contraindicated.</p>

GRACIA-3 2010 (Continued)

The trial was divided in 4 groups; 2 groups received tirofiban as co-intervention. Participants randomly assigned to tirofiban received 150 to 325 mg of non-enteric-coated aspirin orally (chewed) or 500 mg IV as soon as STEMI was identified. Immediate thrombolysis with tenecteplase was administered according to the instructions in the Summary of Product Characteristics for the treatment of STEMI (30 mg if body weight was less than 60 kg, 35 mg if it was 60 to 69 kg, 40 mg if it was 70 to 79 kg, 45 mg if it was 80 to 89 kg, and 50 mg if it was 90 kg or more). No enoxaparin bolus was given to these participants; enoxaparin was administered as a subcutaneous injection at 0.75 mg/kg. This subcutaneous dose was the only dose administered if the coronary intervention was successfully performed within 8 hours of randomisation. Tirofiban was started 2 hours after tenecteplase with a fixed 25 g/kg IV bolus followed by 0.15 g/kg/min in IV infusion for 24 hours. The infusion was reduced by 50% if severe renal failure (creatinine clearance 30 mL/h or creatinine 2 mg/dL) was observed. The rationale for withholding tirofiban for 2 hours after fibrinolytic treatment was based on the pharmacokinetics and pharmacodynamics of tenecteplase. In a previous study with 103 participants, 14 participants receiving tenecteplase exhibited biphasic elimination from the plasma with a mean initial half-life of 22 minutes and a mean terminal half-life of 115 minutes. This mean terminal half-life of 115 minutes motivated the rationale for withholding tirofiban for 2 hours in order to avoid bleeding complications.

Participants randomly assigned to no tirofiban received 150 to 325 mg of non-enteric-coated aspirin orally (chewed) or 500 mg IV as soon as STEMI was identified. Immediate thrombolysis with tenecteplase was administered, with the doses and mode of administration described above. An enoxaparin bolus of 30 mg followed by a subcutaneous dose of 1 mg/kg was administered in participants 75 years old. Participants over 75 years did not receive the enoxaparin bolus, and the subcutaneous injection was reduced to 0.75 mg/kg. This subcutaneous dose was also the only dose administered if coronary intervention was successfully performed within 8 hours of random assignment.

Outcomes	Rate of binary stenosis, epicardial and myocardial flow, stent thrombosis, complicated myocardial infarction, major bleeding
Notes	Cross-overs, NCT00306228 retrospectively. We contacted the authors on 21.03.17 at faviles@secardiologia.es regarding random sequence generation, blinding, and additional data.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Low risk	Central telephone system
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Clinical outcome was adjudicated by an independent clinical events committee blinded to study group assignment.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	More than 5% dropped out for angiographic follow-up, but not for clinical follow-up.
Selective reporting (reporting bias)	Unclear risk	The protocol was produced after enrolment.

GRACIA-3 2010 (Continued)

Other bias	Low risk	The trial was funded by Red Temática de Enfermedades Cardiovasculares (RECAVA) of the Instituto de Salud Carlos III (Spanish Ministry of Science and Innovation), from the Fondo de Investigación Sanitaria of the Instituto de Salud Carlos III (Spanish Ministry of Science and Innovation), and from the Junta de Castilla y León.
------------	----------	---

HAAMU-STENT 2006
Study characteristics

Methods	Randomised clinical trial with a factorial (2x2) design in Finland
Participants	164 participants with STEMI. Male:female = not available Mean age = 63 years (from meta-analysis from 2010) Exclusion criteria: none stated.
Interventions	Experimental group: paclitaxel-eluting Taxus Express stent. Control group: BMS (Liberte). Co-intervention: in addition to abciximab or a thrombolytic agent, all participants received aspirin, enoxaparin, and clopidogrel in the acute phase. Those treated with prehospital thrombolysis also received a glycoprotein IIb/IIIa blocker during and after PCI. Clopidogrel was prescribed for 12 months if tolerated.
Outcomes	Target vessel revascularisation, stent thrombosis, quantitative angiographic analysis, mortality (not reported in a manner that could be used)
Notes	The trial was part of a larger trial randomising participants to abciximab-facilitated percutaneous coronary intervention (afPCI, n= 90) with prehospital thrombolysis (PHT, n= 74). 0 cross-overs. 12 participants died, but from which groups was not described in the publication. It should be noted that it seems more participants are missing from the DES group at angiographic follow-up (13) than the BMS group (7). The publication states that 12 of these participants died, 5 declined to have follow-up, 1 was lost to follow-up, and 1 had renal failure. 1 participant was unaccounted for. Data for mortality were taken from previous meta-analysis. We contacted the authors at if.suh@alareit.akkli on 23 April 2017, but received no reply.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not described

HAAMU-STENT 2006 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Just 1 participant was lost to follow-up.
Selective reporting (reporting bias)	Unclear risk	There was no pre-published protocol, and the trial did not report on mortality, serious adverse events, or MACE.
Other bias	Unclear risk	It was unclear how the trial was funded.

HORIZONS-AMI 2009
Study characteristics

Methods	Randomised clinical trial with parallel/factorial design in USA, Netherlands, Italy, Spain, Germany, UK, Israel, Argentina, Poland, Austria, Norway, Sweden, and Denmark
Participants	<p>3006 participants 18 years of age or older presenting within 12 hours after the onset of symptoms and who had ST-segment elevation of 1 mm or more in 2 or more contiguous leads, new left bundle branch block, or true posterior myocardial infarction were considered for enrolment. Patients were considered eligible for random assignment to paclitaxel-eluting stents or bare-metal stents if an acute infarct-related artery was present in which all lesions requiring PCI had a visually estimated reference-vessel diameter between 2.25 mm and 4.0 mm, without excessive tortuosity or severe calcification.</p> <p>Male:female = 2307:699</p> <p>Mean age = 59.6 years</p> <p>Exclusion criteria: hypersensitivity or contraindication to heparin, both abciximab and eptifibatide, aspirin, both clopidogrel and ticlopidine, bivalirudin, paclitaxel of Taxol, the polymer components of the Taxus stent, stainless steel, or contrast media (refractory to medications or history of anaphylaxis). Prior administration of thrombolytic therapy, bivalirudin, glycoprotein IIb/IIIa inhibitors, low molecular weight heparin, or fondaparinux for this admission. Current use of warfarin. Systemic (IV) paclitaxel or Taxol use within 12 months. Female of childbearing potential, unless a recent pregnancy test is negative, who possibly plans to become pregnant any time after enrolment into this study. History of bleeding diathesis or known coagulopathy (including heparin-induced thrombocytopenia) or refusal of blood transfusions. History of intracerebral mass, aneurysm, arteriovenous malformation, or haemorrhagic stroke. Stroke or transient ischaemic attack within the past 6 months or any permanent residual neurologic defect. Gastrointestinal or genitourinary bleeding within the last 2 months or major surgery within 6 wk. Recent history or known current platelet count < 100,000 cells/mm³ or haemoglobin < 10 g/dL. Extensive peripheral vascular disease, such that emergent angiography and intervention in the opinion of the investigator is likely to be difficult or complicated. An elective surgical procedure is planned that would necessitate interruption of thienopyridines during the first 6 months' postenrolment. Non-cardiac comorbid conditions are present with life expectancy < 1 year or that could result in protocol non-compliance. Patients who are actively participating in another drug or device investigational study, who have not completed the primary endpoint follow-up period. Previous enrolment in this trial. Patients who underwent coronary stent implantation within the past 30 days. 1 or more haemodynamically significant lesion(s) present in the infarct vessel (or side branches) that can only undergo balloon angioplasty or cannot be stented with a study stent (i.e. do not meet the angiographic inclusion criteria for a study stent). The presence of a bifurcation lesion in the infarct vessel that will definitely require the implantation of 2 stents for treatment. Anticipated need for greater than 100 mm of study stent length. The infarct-related artery is an unprotected left main segment. Patients with significant multivessel disease or anatomical features otherwise unfavourable for angioplasty such that the patient will have a high likelihood of requiring bypass surgery before 30 days. The culprit vessel or lesion cannot be identified. Patient presenting with possible/probable stent thrombosis. Any patient</p>

Drug-eluting stents versus bare-metal stents for acute coronary syndrome (Review)

HORIZONS-AMI 2009 (Continued)

in whom angiography demonstrates the infarct lesion to be at the site of a previously implanted stent (bare metal or drug eluting).

Interventions	<p>Experimental group: paclitaxel-eluting stents (Taxus Express, Boston Scientific).</p> <p>Control group: identical bare-metal stents (Express, Boston Scientific).</p> <p>Co-intervention: aspirin (324 mg administered in chewable form or 500 mg administered intravenously) was given in the emergency room, after which 300 to 325 mg was given orally every day during the hospitalisation and 75 to 81 mg every day thereafter indefinitely. A loading dose of clopidogrel (either 300 mg or 600 mg, at the discretion of the investigator) was administered before catheterisation, followed by 75 mg orally every day for at least 6 months (with a recommendation of 1 year or longer).</p>
Outcomes	Major adverse cardiovascular events consisting of death, reinfarction, stroke, and stent thrombosis, TLR, TVR
Notes	Before randomisation for stents, participants were randomised either to heparin + glycoprotein IIa/IIIa inhibitors or bivalirudin alone without breaking the protocol for randomisation for stents. We contacted the authors on 23 April 2017 at gs2184@columbia.edu regarding blinding and unpublished data, but have received no reply.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computerised interactive voice response system
Allocation concealment (selection bias)	Low risk	Computerised interactive voice response system
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Independent clinical events committee, unaware of treatment assignment
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	More than 5% dropped out at 3 years' follow-up, but not at 1 year.
Selective reporting (reporting bias)	Unclear risk	Enrolment begun before the protocol were published.
Other bias	High risk	The trial was funded by Merck, Nycomed, OrbusNeich, Pfizer, Possis Medical, Promed, Sanofi-Aventis, Siemens, Solvay, Terumo, and Tyco.

König 2007

Study characteristics

Methods	Parallel-design randomised clinical trial in Germany
---------	--

König 2007 (Continued)

Participants	<p>135 participants referred from the emergency department for stent implantation due to ACS, diagnosed by ST-elevation myocardial infarction or cardiac enzyme elevation.</p> <p>Male:female = 83:37 (analysed only)</p> <p>Mean age = 62.5 years</p> <p>Exclusion criteria: not described.</p>
Interventions	<p>Experimental group: dexamethasone-eluting stent (Dexamet, Abbott Vascular Devices).</p> <p>Control group 1: BMS (BiodivYsio, Biocompatibles Cardiovascular Ltd.) consisting of the same stent design without the drug.</p> <p>Co-intervention: the target lesion was a de novo lesion in a native coronary artery. The balloon size was required to reach a balloon-to-artery ratio of 1.0 to 1.1. Predilation was performed before stent implantation. All participants received aspirin (100 mg/d, indefinitely) and clopidogrel as a 300 mg loading dose before stent implantation, followed by a dose of 75 mg/d for 12 months. Stents of various dimensions were used (nominal diameter, 2.50 to 4.00 mm; length, 8.00 to 18.00 mm). Postinterventional medical therapy was similar in both groups.</p>
Outcomes	Quantitative coronary angiographic analysis, intracoronary ultrasound analysis and quantitative measurements
Notes	<p>We were unable to find a protocol.</p> <p>MACE was not defined.</p> <p>We contacted the authors at andreas.koenig@med.uni-muenchen.de on 23 April 2017, but received no reply.</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Only the quantitative coronary angiographic analysis was described as blinded.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	The number of dropouts was unclear.
Selective reporting (reporting bias)	Unclear risk	There was no pre-published protocol, and the trial did not report on serious adverse events.
Other bias	Unclear risk	It was unclear how the trial was funded.

Mission 2008

Study characteristics

Methods	Parallel-design randomised clinical trial in 1 centre in the Netherlands
Participants	<p>310 participants with STEMI symptoms started 9 h before the procedure and the ECG demonstrated STEMI (ST-segment elevation 0.2 mV in 2 contiguous leads in V 1 through V 3 or 0.1 mV in other leads, or (presumed) new left bundle branch block). The target lesion length should be equal to or less than 24 mm.</p> <p>Male:female = 241:69</p> <p>Mean age = 59.15 years</p> <p>Exclusion criteria: 1) age 18 years or 80 years; 2) left main stenosis of 50%; 3) triple-vessel disease, defined as 50% stenosis in 3 major epicardial branches; 4) previous PCI or coronary artery by-pass grafting of the infarct-related artery; 5) thrombolytic therapy for the index infarction; 5) target vessel reference diameter 2.25 mm or 3.75 mm; 6) need for mechanical ventilation; 7) contraindication to the use of aspirin, clopidogrel, heparin, or abciximab; 8) known renal failure; or 9) a life expectancy 12 months.</p>
Interventions	<p>Experimental group: sirolimus-eluting stent (Cypher, Cordis Corp., Miami Lakes, FL).</p> <p>Control group: BMS (Vision, Guidant Corp., Indianapolis, IN).</p> <p>Co-intervention: during the study period, all participants were treated according to the institutional STEMI protocol, which included standardised outpatient follow-up. Before the procedure all participants received 300 mg of aspirin, 300 to 600 mg of clopidogrel, and an intravenous bolus of abciximab (25 g/kg), followed by a continuous infusion of 10 g/kg/min for 12 h.</p> <p>Aspirin (80 to 100 mg/day) was prescribed indefinitely and clopidogrel (75 mg/day) for 12 months.</p> <p>Participants were treated with beta-blocking agents, statins, and angiotensin-converting enzyme inhibitors or angiotensin II blockers.</p>
Outcomes	<p>The primary endpoint of the study was in-segment late luminal loss at 9-month follow-up angiography.</p> <p>Secondary endpoints were angiographic restenosis and late stent malapposition at 9 months. Additional secondary endpoints were death, MI, TVR, TLR, target vessel failure, stent thrombosis, procedural success, and clinical success.</p>
Notes	<p>1 participant crossed over from the DES group to the BMS group.</p> <p>ISRCTN62825862 (registered retrospectively).</p> <p>We contacted the authors at m.j.schalij@lumc.nl on 23 April 2017, but have received no reply.</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding of participants and personnel (performance bias)	High risk	Single-blind

Mission 2008 (Continued)

All outcomes

Blinding of outcome assessment (detection bias) All outcomes	Low risk	"Patients and operators performing the follow-up angiography were blinded to the treatment assignment."
Incomplete outcome data (attrition bias) All outcomes	Low risk	6 participants were lost from the BMS group and none from the DES group, but due to the low number of participants with missing data, we have assessed the trial as low risk.
Selective reporting (reporting bias)	Unclear risk	There was no pre-published protocol, and the trial did not report on serious adverse events.
Other bias	High risk	This study was supported by the Netherlands Heart Foundation and by an unrestricted research grant from Guidant Inc., Nieuwegein, the Netherlands. Dr Jukema is an established clinical investigator of the Netherlands Heart Foundation (grant 2001D032).

MULTISTRATEGY 2008
Study characteristics

Methods	Parallel-design randomised clinical trial in Italy, Argentina, and Spain	
Participants	<p>745 participants with (1) chest pain for longer than 30 minutes with an electrocardiographic ST-segment elevation of 1 mm or greater in 2 or more contiguous electrocardiogram leads, or with a new left bundle branch block, and (2) admission either within 12 hours of symptom onset or between 12 and 24 hours after onset with evidence of continuing ischaemia.</p> <p>Male:female = 565:180</p> <p>Mean age = 63.8 years</p> <p>Exclusion criteria: administration of fibrinolytics in the previous 30 days, major surgery within 15 days, and active bleeding or previous stroke in the last 6 months.</p>	
Interventions	<p>Experimental group: sirolimus-eluting stent.</p> <p>Control group 1: uncoated stent type approved by the regulatory agency.</p> <p>Co-intervention: the participants were additionally randomised to either abciximab or tirofiban. Heparin was given at 40 to 70 U/kg, targeting an activated clotting time of at least 200 seconds. Participants received aspirin (160 to 325 mg orally or 250 mg intravenously, followed by 80 to 125 mg/d orally indefinitely) and clopidogrel (300 mg orally and then 75 mg/d for at least 3 months).</p>	
Outcomes	Death, reinfarction, TVR, stent thrombosis, bleeding, thrombocytopenia	
Notes	12 cross-overs from DES to BMS, NCT00229515 (retrospectively). We contacted the authors on 23 April 2017 at vlgmrc@unife.it regarding which stents were used and additional data, but have received no reply.	

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated sequence

Drug-eluting stents versus bare-metal stents for acute coronary syndrome (Review)

MULTISTRATEGY 2008 (Continued)

Allocation concealment (selection bias)	Low risk	Sealed envelopes
Blinding of participants and personnel (performance bias) All outcomes	High risk	Open-label study
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Clinical outcomes were assessed by an independent adjudication committee whose members were blinded to treatment assignments.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Less than 5% dropped out.
Selective reporting (reporting bias)	High risk	The protocol was published after enrolment of participants.
Other bias	High risk	The trial was funded by University of Ferrara and a grant from Merck.

OCTAMI 2010
Study characteristics

Methods	Parallel-design, single-centre, randomised clinical trial in Italy
Participants	<p>44 participants with STEMI 12 hours after symptom onset (prolonged chest pain for more than 20 min, unresponsive to nitroglycerin, and ST-segment elevation of at least 1 mm in 2 or more contiguous leads, or true posterior myocardial infarction), an infarct artery in a native coronary vessel with 70% diameter stenosis, a reference vessel diameter of 2.5 to 3.75 mm, and underwent primary PCI with stent implantation.</p> <p>Male:female = 34:10</p> <p>Mean age = 61.1 years</p> <p>Exclusion criteria: patients with left main disease, infarct lesions in bypass grafts, cardiogenic shock, renal failure, recent major bleeding, allergy to aspirin or clopidogrel, on anticoagulant therapy, or with no suitable anatomy for optical coherence tomography (ostial lesions, extreme tortuosity, and large vessels 3.75 mm in diameter) were excluded.</p>
Interventions	<p>Experimental group: Endeavor zotarolimus-eluting stent.</p> <p>Control group: Driver bare-metal stent.</p> <p>Co-intervention: percutaneous coronary intervention was performed according to standard techniques. Direct stenting, thrombus aspiration, and use of glycoprotein IIb/IIIa inhibitors were allowed and left to the operator's discretion. All participants were pre-treated with aspirin 250 mg intravenously and clopidogrel 300 mg orally before PCI, followed by daily administration of clopidogrel 75 mg for at least 6 months after discharge and aspirin indefinitely. Participants received unfractionated heparin during PCI to maintain an activated clotting time of 300 s or more.</p>
Outcomes	Uncovered struts, maximum length of uncovered segments, percentage of malapposed struts and maximum length of malapposed segments, neointimal response, death, MI, and stent thromboses

OCTAMI 2010 (Continued)

Notes No cross-overs, ClinicalTrials.gov number NCT00704561. We contacted the authors at guagliumig@gmail.com on 23 April 2017, but have received no reply.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding of participants and personnel (performance bias) All outcomes	High risk	"Open label"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Core laboratory personnel were blinded to the treatment assignment.
Incomplete outcome data (attrition bias) All outcomes	Low risk	"Clinical follow-up was complete in all patients at 1 year."
Selective reporting (reporting bias)	Low risk	The outcomes were stated on ClinicalTrials.gov 2 months after trial start.
Other bias	High risk	"Supported by Ospedali Riuniti di Bergamo (Bergamo, Italy) with grant support from Medtronic CardioVascular, Santa Rosa, California. Dr. Guagliumi has received grant/research support from Boston Scientific Corporation, Medtronic Vascular, LightLab Imaging, and Labcoat and is a consultant for Boston Scientific Corporation and Volcano Corporation. Dr. Costa is on the Speakers' Bureau and is a consultant for Boston Scientific Corporation, Sanofi-Aventis, and Eli Lilly and is on the Speakers' Bureau and is a member of the Scientific Advisory Board for Abbott, Cordis, LightLab Imaging, and Scitec. Dr. Sirbu has received grant/research support from LightLab Imaging."

PASEO 2009a

Study characteristics

Methods	Parallel-design randomised clinical trial with 2 experimental groups in Italy
Participants	<p>270 participants with STEMI who fulfilled all of the following criteria: 1) chest pain for more than 30 min; 2) ST-segment elevation of 1 mm or more in 2 or more contiguous electrocardiograph leads or with presumably new left bundle branch block; 3) hospital admission within 12 h from symptoms onset.</p> <p>Male:female = 190:80</p> <p>Mean age = 62 years</p> <p>Exclusion criteria: "Exclusion criteria included: 1) active internal bleeding or a history of bleeding diathesis within the previous 30 days; 2) history of intracranial hemorrhage, intracranial neoplasm, arteriovenous malformation or aneurysm; 3) known allergy to sirolimus, paclitaxel, heparin, aspirin, or clopidogrel; 4) history of stroke within 30 days or any history of hemorrhagic stroke; 5) major surgical</p>

Drug-eluting stents versus bare-metal stents for acute coronary syndrome (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

PASEO 2009a (Continued)

procedure or severe physical trauma within the previous month; 6) history, symptoms, or findings suggestive of aortic dissection; 7) thrombolytic/fibrinolytic therapy within 24 h; 8) history of thrombocytopenia; 9) hemorrhagic retinopathy; 10) patients on warfarin or acenocoumarol with international normalized ratio 2; and 11) pregnancy. A vessel size 2.25 mm was the only angiographic exclusion criteria."

Interventions	<p>Experimental group: Taxus paclitaxel-eluting stent.</p> <p>Control group: bare-metal stent.</p> <p>Co-intervention: in the coronary care unit, all participants received 70 U/kg IV bolus of unfractionated heparin plus 1000 U/h infusion (to maintain an activated clotting time of at least 200 s), aspirin intravenously (500 mg), and clopidogrel (300 mg loading dose). All participants received up-stream glycoprotein IIb/IIIa inhibitors as a routine adjunctive therapy before primary PCI. Postinterventional antiplatelet therapy for all participants included in the 3 study groups consisted of aspirin (100 mg) indefinitely and clopidogrel (75 mg for 6 months).</p>
Outcomes	<p>The primary endpoint was TLR at 1-year follow-up.</p> <p>Secondary endpoints were: 1) cumulative combined incidence of death and/or recurrent MI at 2-year follow-up; 2) cumulative incidence of in-stent thrombosis (assessed according to Academic Research Consortium's definition) at 2-year follow-up; and 3) major adverse cardiac events (combined death and/or recurrent MI and/or TLR) at 2-year follow-up.</p>
Notes	<p>We halved the control group to account for the 2 experimental arms and to avoid double-counting. Clinical trial number NCT00759850, registered retrospectively. We contacted the authors at emidilorenzo@tin.it on 23 April 2017, but have received no reply.</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer generated
Allocation concealment (selection bias)	Low risk	Sealed envelopes
Blinding of participants and personnel (performance bias) All outcomes	High risk	"Open Label"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	<p>"All events were reviewed by 2 cardiologists blinded to treatment assignment."</p> <p>"Offline quantitative coronary angiography (Integris Allura, Philips, Best, the Netherlands) were performed by 2 experienced technicians who were unaware of treatment assignment with the averaging scores if they were not in agreement."</p>
Incomplete outcome data (attrition bias) All outcomes	Low risk	"No patient was lost to follow-up."
Selective reporting (reporting bias)	Unclear risk	There was no pre-published protocol (registered retrospectively at ClinicalTrials.gov), and the trial did not report on serious adverse events.
Other bias	Low risk	San Giuseppe Moscati Hospital funded the study (from ClinicalTrials.gov).

PASEO 2009b

Study characteristics

Methods	Parallel-design, single-centre, randomised clinical trial with 2 experimental groups in Italy
Participants	<p>270 participants with STEMI who fulfilled all the following criteria: 1) chest pain for more than 30 min; 2) ST-segment elevation of 1 mm or more in 2 or more contiguous electrocardiograph leads or with presumably new left bundle branch block; 3) hospital admission within 12 h from symptoms onset.</p> <p>Male:female = 190:80</p> <p>Mean age = 62 years</p> <p>Exclusion criteria: "Exclusion criteria included: 1) active internal bleeding or a history of bleeding diathesis within the previous 30 days; 2) history of intracranial hemorrhage, intracranial neoplasm, arteriovenous malformation or aneurysm; 3) known allergy to sirolimus, paclitaxel, heparin, aspirin, or clopidogrel; 4) history of stroke within 30 days or any history of hemorrhagic stroke; 5) major surgical procedure or severe physical trauma within the previous month; 6) history, symptoms, or findings suggestive of aortic dissection; 7) thrombolytic/fibrinolytic therapy within 24 h; 8) history of thrombocytopenia; 9) hemorrhagic retinopathy; 10) patients on warfarin or acenocoumarol with international normalized ratio 2; and 11) pregnancy. A vessel size 2.25 mm was the only angiographic exclusion criteria."</p>
Interventions	<p>Experimental group: Cypher sirolimus-eluting stent.</p> <p>Control group: bare-metal stent.</p> <p>Co-intervention: in the coronary care unit, all participants received 70 U/kg IV bolus of unfractionated heparin plus 1000 U/h infusion (to maintain an activated clotting time of at least 200 s), aspirin intravenously (500 mg), and clopidogrel (300 mg loading dose). All participants received up-stream glycoprotein IIb/IIIa inhibitors as a routine adjunctive therapy before primary PCI. Postinterventional antiplatelet therapy for all participants included in the 3 study groups consisted of aspirin (100 mg) indefinitely and clopidogrel (75 mg for 6 months).</p>
Outcomes	<p>The primary endpoint was TLR at 1-year follow-up.</p> <p>Secondary endpoints were: 1) cumulative combined incidence of death and/or recurrent MI at 2-year follow-up; 2) cumulative incidence of in-stent thrombosis (assessed according to Academic Research Consortium's definition) at 2-year follow-up; and 3) major adverse cardiac events (combined death and/or recurrent MI and/or TLR) at 2-year follow-up.</p>
Notes	<p>We halved the control group to account for the 2 experimental arms and to avoid double-counting.</p> <p>Clinical trial number NCT00759850, registered retrospectively.</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer generated
Allocation concealment (selection bias)	Low risk	Sealed envelopes
Blinding of participants and personnel (performance bias) All outcomes	High risk	"Open Label"

PASEO 2009b (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	"All events were reviewed by 2 cardiologists blinded to treatment assignment." "Offline quantitative coronary angiography (Integris Allura, Philips, Best, the Netherlands) were performed by 2 experienced technicians who were unaware of treatment assignment with the averaging scores if they were not in agreement."
Incomplete outcome data (attrition bias) All outcomes	Low risk	"No patient was lost to follow-up."
Selective reporting (reporting bias)	Unclear risk	There was no pre-published protocol (registered retrospectively at ClinicalTrials.gov), and the trial did not report on serious adverse events.
Other bias	Low risk	San Giuseppe Moscati Hospital funded the study (from ClinicalTrials.gov).

PASSION 2006

Study characteristics

Methods	Parallel-design randomised clinical trial in the Netherlands
Participants	<p>619 participants with an acute myocardial infarction with ST-segment elevation (> 20 minutes of chest pain and at least 1 mm of ST-segment elevation in at least 2 contiguous leads or a new left bundle branch block), reperfusion was expected to be achieved within 6 hours after the onset of symptoms, and the native coronary artery was considered to be suitable for primary PCI with stent implantation.</p> <p>Male:female = 470:149</p> <p>Mean age = 61 years</p> <p>Exclusion criteria: patients were excluded if they had received thrombolytic therapy; the infarction was caused by in-stent thrombosis or restenosis; there was a contraindication to aspirin, clopidogrel, or both; patients were participating in another clinical trial; cardiogenic shock was evident before randomisation; the neurologic outcome after resuscitation was uncertain; they had undergone intubation, ventilation, or both; there was known intracranial disease; or the estimated life expectancy was less than 6 months.</p>
Interventions	<p>Experimental group: paclitaxel-eluting stent (Taxus Express2, Boston Scientific).</p> <p>Control group: uncoated stent (Express2 or Liberté, Boston Scientific).</p> <p>Co-intervention: aspirin (at a dose of 100 to 500 mg) and clopidogrel (300 mg) were administered upon arrival at the hospital. A glycoprotein IIb/IIIa receptor blocker was administered at the discretion of the operator. A bolus of 10,000 IU of unfractionated heparin was administered before the procedure. Coronary angiography was performed through either the radial or the femoral artery. The target segment was filmed in at least 2 orthogonal planes after the intracoronary administration of 100 to 200 µg of nitroglycerin; quantitative coronary angiography was then performed. The use of thrombectomy devices and predilatation balloons was at the operator's discretion. Participants were prescribed 80 to 100 mg of aspirin daily for life and 75 mg of clopidogrel daily for at least 6 months.</p>
Outcomes	The primary endpoint was the first occurrence of a serious adverse cardiac event at 12 months, including death from cardiac causes, recurrent myocardial infarction requiring hospitalisation, and ischaemia-driven revascularisation of a target lesion. The secondary endpoints of the study were revas-

PASSION 2006 (Continued)

cularisation of a target lesion and a composite of death from cardiac causes or recurrent myocardial infarction.

Notes
ISRCTN65027270 (registered retrospectively). We contacted the authors at m.a.vink@olvg.nl on 7 March 2017, but received no reply.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Low risk	"Assignment to study groups was performed with the use of sealed envelopes"
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	"Patients, referring physicians, investigators responsible for obtaining follow-up information, and interventionalists performing repeated procedures were all unaware of treatment assignments" "Drs. Laarman and Suttorp adjudicated all end points of the study in a blinded fashion."; however the trial was described as single-blinded.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"Patients, referring physicians, investigators responsible for obtaining follow-up information, and interventionalists performing repeated procedures were all unaware of treatment assignments" "Drs. Laarman and Suttorp adjudicated all end points of the study in a blinded fashion."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Less than 5% with incomplete outcome data in both groups.
Selective reporting (reporting bias)	Unclear risk	There was no pre-published protocol, and the trial did not report on quality of life.
Other bias	Low risk	"The trial was entirely funded by the Department of Interventional Cardiology at Onze Lieve Vrouwe Gasthuis"

SELECTION 2007

Study characteristics

Methods	Parallel-design, single-centre, randomised clinical trial in Italy
Participants	80 participants younger than 75 years with chest pain persisting ≥ 30 minutes that was associated with ST-segment elevation ≥ 0.1 mV in ≥ 2 continuous electrocardiographic leads. Male:female = 66:14 Mean age = 60.7 years Exclusion criteria: clinical exclusion criteria were cardiogenic shock, thrombolytic therapy, oral anticoagulant therapy, prolonged cardiopulmonary resuscitation, previous coronary artery bypass graft, PCI, or ischaemic stroke within 6 months, and haemorrhagic diathesis. Angiographic exclusion criteria were

SELECTION 2007 (Continued)

represented by previous stenting of the infarct-related artery and a reference vessel diameter of the culprit lesion of < 2.25 mm by visual estimate.

Interventions	<p>Experimental group: Taxus paclitaxel-eluting stent.</p> <p>Control group: Express bare-metal stents.</p> <p>Co-intervention: primary PCI was performed according to standard care. After stent implantation, intravascular ultrasound was performed on the stented segment and further postdilation was left to operator discretion. All participants received abciximab (ReoPro, Centocor, Malvern, PA) as a bolus of 0.25 mg/kg of body weight followed by a 12-hour infusion at a rate of 0.125 µg/kg/min. Unfractionated heparin was given as an initial bolus of 70 IU/kg, and additional boluses were administered during the procedure to achieve an activated clotting time of 250 to 300 seconds. Participants were routinely treated with aspirin (250 to 500 mg IV bolus followed by 100 mg/day orally indefinitely) and clopidogrel (300 mg loading dose and 75 mg/day for 9 months).</p>
Outcomes	<p>The primary endpoint was a percentage of the stent volume obstructed by neointimal hyperplasia measured by intravascular ultrasound at 7 ± 1 months. Secondary endpoints were 7 ± 1 months angiographic binary restenosis rate and MACE, i.e. any death, non-fatal myocardial infarction, and TLR at discharge, at 1 and 7 ± 1 months' follow-up.</p>
Notes	<p>We contacted the authors at vittoriguido@gmail.com regarding random sequence generation, allocation concealment, blinding of outcome assessors, pre-published protocol, and trial funding, but have received no reply.</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding of participants and personnel (performance bias) All outcomes	High risk	Operators not blinded.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts
Selective reporting (reporting bias)	Unclear risk	There was no pre-published protocol, and the trial did not report on serious adverse events.
Other bias	Unclear risk	It was unclear how the trial was funded.

SESAMI 2007

Study characteristics

Drug-eluting stents versus bare-metal stents for acute coronary syndrome (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

SESAMI 2007 (Continued)

Methods	Parallel-design randomised clinical trial in Italy
Participants	<p>320 participants who were 18 years of age, had symptoms of acute MI for 30 min but 12 h, and had 1 mm ST-segment elevation in at least 2 contiguous leads or left bundle branch block.</p> <p>Male:female = 256:64</p> <p>Mean age = 62.5 years</p> <p>Exclusion criteria: cardiogenic shock (systolic blood pressure 80 mmHg for 30 min or need for intra-venous pressors or intra-aortic balloon counterpulsation); a history of bleeding diathesis, leukopenia, thrombocytopenia, or severe hepatic or renal dysfunction; non-cardiac illness associated with a life expectancy of 1 year; left main coronary artery or graft disease; participation in another study; or inability to give informed consent owing to prolonged cardiopulmonary resuscitation.</p>
Interventions	<p>Experimental group: Cypher sirolimus-eluting stent.</p> <p>Control group: BX stent (bare-metal stent).</p> <p>Co-intervention: the study protocol recommended that aspirin (500 mg intravenously) and beta-blockers (in the absence of contraindications) be administered in the emergency room. Clopidogrel, an inhibitor of adenosine diphosphate-induced platelet aggregation, was given as a bolus of 4 tablets immediately after the procedure and was continued for 1 year in both groups. The glycoprotein IIb/IIIa receptor inhibitor abciximab (ReoPro, Eli Lilly, Indianapolis, IN; Centocor, Horsham, PA) was administered as a 0.25 mg/kg bolus followed by a 12-hour infusion (0.125 g/kg/min; maximum 10 g/min). If it was not started in the emergency room, abciximab therapy was initiated in the catheterisation laboratory before coronary angiography. The heparin dose was calculated to achieve an activated clotting time of 200 to 250 s.</p>
Outcomes	<p>The primary endpoint for the trial was binary restenosis at the 1-year angiographic follow-up.</p> <p>The secondary endpoints were TLR, TVR, MACE, and target vessel failure at 1 year.</p>
Notes	No mention of cross-overs. Clinical trial number NCT00288210 (registered retrospectively). We contacted the authors at cmusto@hotmail.it on 12 April 2017 regarding a pre-published protocol, whether they had recorded serious adverse events comprehensively, and whether they had data for cardiovascular mortality.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	The allocation schedule was based on computer-generated random numbers.
Allocation concealment (selection bias)	Low risk	Sealed, sequentially numbered, opaque allocation envelopes
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	All serious clinical events, including stent thrombosis, were reviewed by 2 authors (AG and DA), who were unaware of stent assignment.

SESAMI 2007 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	Less than 5% with incomplete outcome data for clinical outcomes in both groups. More than 5% missing angiographic control at 1 year (50% in both groups).
Selective reporting (reporting bias)	Unclear risk	There was no pre-published protocol, and the trial did not report on serious adverse events or cardiovascular mortality.
Other bias	Low risk	San Camillo Hospital, Rome, funded the study (taken from ClinicalTrials.gov).

Steinwender 2008

Study characteristics

Methods	Parallel-design randomised clinical trial in Austria
Participants	16 participants with a first ST-elevation anterior myocardial infarction who were eligible for primary percutaneous coronary intervention. Male:female = 12:4 Mean age = 55.5 years Exclusion criteria: not described.
Interventions	Experimental group: sirolimus-eluting stent (Cypher, Cordis Corp., Miami, FL). Control group: bare-metal stent (Driver, Medtronic, Minneapolis, MN). Co-intervention: postinterventional therapy consisted of aspirin (100 mg/d), clopidogrel (75 mg/d for 3 months), beta-blockers, angiotensin-converting enzyme or -receptor blockers, and a statin, if indicated. The second day after stent implantation, granulocyte-colony stimulating factor mobilised autologous stem cells therapy at a daily dose of 10 µg/kg body weight divided into 2 subcutaneous injections was initiated.
Outcomes	Quantitative coronary angiography, death
Notes	We contacted the authors at clemens.steinwender@akh.linz.at on 23 March 2017 regarding randomisation, blinding, funding, protocol, and additional clinical outcomes, but have received no reply.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not described
Blinding of outcome assessment (detection bias)	Unclear risk	Not described

Steinwender 2008 (Continued)

All outcomes

Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up.
Selective reporting (reporting bias)	Unclear risk	No protocol could be obtained.
Other bias	Unclear risk	It was unclear how the trial was funded.

Strozzi 2007

Study characteristics

Methods	Parallel-design randomised clinical trial in Croatia
Participants	<p>79 participants who underwent stent implantation in acute coronary syndrome from January 2003 to May 2004. A diagnosis of ACS included acute myocardial infarction with ST elevation, prolonged angina for more than 20 minutes, or recurrent episodes at rest with indicators of cardiac ischaemia or injury (cardiac enzyme elevation and ST-segment denivelation).</p> <p>Male:female = 62:17</p> <p>Mean age = 57.8 years</p> <p>Exclusion criteria: patients with previous PCI or coronary artery bypass graft surgery, multivessel, diffuse disease, tortuous vessel, arteries less than 3 mm in diameter, distal stenosis location, and left main and bifurcation lesions.</p>
Interventions	<p>Experimental group: sirolimus-eluting stent.</p> <p>Control group 1: bare-metal stent.</p> <p>Co-intervention: all procedures were performed using standard transfemoral approach (6) with 7 French guiding catheters (6). All participants received aspirin (300 mg), heparin (10,000 IU or more in longer procedures), and eptifibatide. Standard PCI was performed with balloon predilation, stent placement, and postdilation if needed. Postprocedural medications included aspirin 100 mg/d and ticlopidine 500 mg/d (clopidogrel was not available). Ticlopidine was stopped 6 weeks after the procedure.</p>
Outcomes	Quantitative coronary arteriography analysis, death, MI, coronary artery bypass grafting, TLR, MACE
Notes	<p>The trial had a third arm randomising participants to stent grafts. NCT00452517 (retrospectively registered).</p> <p>We contacted the authors at maja.strozzi@zg.t-com.hr on 12 April 2017 regarding random sequence generation, blinding of outcome assessment, pre-published protocol, the number of participants with complete data, trial funding, and serious adverse events. The email appeared to be no longer functioning, and we were unable to find additional contact information.</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomisation codes

Strozzi 2007 (Continued)

Allocation concealment (selection bias)	Unclear risk	Not described
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not described
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	The number of dropouts was unclear.
Selective reporting (reporting bias)	High risk	Protocol was published retrospectively.
Other bias	Unclear risk	Not described

Typhoon 2006
Study characteristics

Methods	Parallel-design randomised clinical trial conducted in 48 centres in France and Germany
Participants	<p>715 participants with symptoms that began less than 12 hours before catheterisation and if the electrocardiogram showed ST-segment elevation (at least 1 mm in 2 or more standard leads or at least 2 mm in 2 or more contiguous precordial leads).</p> <p>Male:female = 558:154</p> <p>Mean age = 59.25 years</p> <p>Exclusion criteria: clinical criteria for exclusion included the administration of fibrinolytic agents for the index infarction, overt acute heart failure, a previously documented left ventricular ejection fraction of less than 30%, previous myocardial infarction, and an estimated life expectancy of less than 12 months.</p>
Interventions	<p>Experimental group: sirolimus-eluting stent (Cypher or Cypher Select, Cordis, Johnson & Johnson).</p> <p>Control group: any commercially available uncoated stent.</p> <p>Co-intervention: participants were premedicated with aspirin (at least 100 mg) and unfractionated heparin (5000 to 10,000 IU). A loading dose of 300 mg of clopidogrel was administered either before or immediately after PCI. Heparin was administered throughout the procedure in order to maintain an activated clotting time of 250 seconds or longer. Administration of platelet glycoprotein IIb/IIIa inhibitors was left to the investigator's discretion.</p> <p>Combined antiplatelet therapy included daily administration of aspirin (100 mg) and either clopidogrel (75 mg) or ticlopidine (250 mg). Dual antiplatelet therapy was recommended for at least 6 months, and aspirin therapy was recommended indefinitely.</p>
Outcomes	The primary endpoint was target vessel failure, defined as the composite of TVR, recurrent infarction, or target vessel-related death at 1 year.

Typhoon 2006 (Continued)

The secondary endpoints included the rate of successful treatment of the lesion, defined by residual stenosis of less than 50% of the reference luminal diameter on quantitative coronary analysis and a TIMI flow grade of 3, and in-stent late luminal loss on quantitative coronary analysis at 8 months.

Notes No cross-overs noted. ClinicalTrials.gov: NCT00232830 (registered retrospectively). We contacted the authors at cspauldi@its.jnj.com on 12 April 2017 regarding random sequence generation, blinding of outcome assessment, pre-published protocol, and serious adverse events, but it seemed that our emails were rejected by the author's email service.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Low risk	Sealed envelopes
Blinding of participants and personnel (performance bias) All outcomes	High risk	"Patients, but not investigators, were unaware of the treatment assignment"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not described
Incomplete outcome data (attrition bias) All outcomes	Low risk	Only 3 participants had missing data at 1-year follow-up. At 4 years' follow-up 81% had follow-up data. The trial used multiple imputation to deal with missing data.
Selective reporting (reporting bias)	Unclear risk	There was no pre-published protocol, and the trial did not report on serious adverse events.
Other bias	High risk	"The TYPHOON trial was an investigator-initiated trial sponsored by Cordis Corporation, a Johnson & Johnson company. Dr. Spaulding was a member of a Cordis scientific advisory board; has received honorarium from Cordis for speaker's fees; and has been a full-time employee of the Cordis Corporation since June 1, 2010. Dr. Varenne is a speaker for Cordis, Boston Scientific, and Abbott. Drs. Cebrian, Wang, and Stoll are full-time employees of the Cordis Corporation and have Johnson & Johnson stock options. Dr. Fajadet is a consultant for Cordis and Abbott. Dr. Erglis is on the advisory board for Cordis. Dr. Hosten is employed by the Cordis Corporation as a consultant. Dr. Henry is part of the Cordis Exchange program."

ACS: acute coronary syndrome

BMS: bare-metal stent

CK-MB: myocardial band isoenzyme of creatine kinase

DES: drug-eluting stent

ECG: electrocardiogram

IV: intravenous

MACE: major adverse cardiac events

MI: myocardial infarction

NSTEMI: non-ST-segment elevation myocardial infarction

PCI: percutaneous coronary intervention

STEMI: ST-segment elevation myocardial infarction

TLR: target lesion revascularisation

TVR: target vessel revascularisation

Drug-eluting stents versus bare-metal stents for acute coronary syndrome (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
ACUITY 2006	Wrong intervention: the participants were not randomised to drug-eluting stents versus bare-metal stents but to bivalirudin versus no bivalirudin
BARRICADE 2011	Wrong intervention: not drug-eluting stent
DESSOLVE-1 2011	Not a randomised clinical trial
Gioia 2006	Not a randomised clinical trial
Halkin 2006	Not a randomised clinical trial
Han 2006	Wrong control: not bare-metal stent
Hausleiter 2005	Not a randomised clinical trial
Hoffmann 2007	Not a randomised clinical trial
Hokimoto 2015	Not a randomised clinical trial
Ishii 2012	Not a randomised clinical trial
JACK-EPC 2013	Wrong intervention: the intervention group did not receive a drug-eluting stent, but a bio-engineered stent with struts covered by murine monoclonal antibodies against human CD34 antigen (Genous)
Kim 2006	Not a randomised clinical trial
Lansky 2000	Wrong intervention: bare-metal stent versus bare-metal stent
Lasave 2007	Not a randomised clinical trial
MASTER 2012	Wrong control: the control group received either drug-eluting or bare-metal stent
Meredith 2007	Not a randomised clinical trial
Oyabu 2006	Not a randomised clinical trial
PATENCY 2007	Excludes myocardial infarction participants
PISCES 2005	Not a randomised clinical trial
Strategy 2005	Multi-intervention: the experimental group received tirofiban in addition to a drug-eluting stent, whereas the control group received abciximab in addition to a bare-metal stent
Suzuki 2013	Not a randomised clinical trial
TRUST 2003	Wrong intervention: not drug-eluting stent
Turan 1998	Wrong intervention: not drug-eluting stent
VESTASYNC 2014	Wrong control: not bare-metal stent

Study	Reason for exclusion
Windecker 2005	Wrong intervention: not drug-eluting stent
XAMI 2013	Wrong control: not bare-metal stent

Characteristics of studies awaiting classification *[ordered by study ID]*

BASKET 2005a

Methods	Parallel design, single-centre, randomised clinical trial with 2 different experimental arms in Switzerland
Participants	<p>826 participants treated with PCI and stenting at the University Hospital of Basel, Switzerland between 5 May 2003 and 31 May 2004 were included in BASKET, irrespective of indication for PCI.</p> <p>Male:female = 320:84</p> <p>Mean age = 64 years</p> <p>Exclusion criteria: a target vessel diameter of 4 mm or greater, as the largest available DES size was 3.5 mm (n = 23); presence of restenotic lesions, given the different causes and outcome of restenotic lesions (n = 49); and no consent, primarily due to patients' or referring physicians' preference for DES before angiography and inability of patients to give informed consent (n = 90).</p>
Interventions	<p>Experimental group 1: sirolimus-coated Cypher stent.</p> <p>Control group: cobalt chromium-based Vision/Pixel stent.</p> <p>Co-intervention: in participants with ST-elevation myocardial infarction, primary PCI was the treatment of choice. Participants with non-ST-elevation myocardial infarction or unstable angina were treated urgently within 24 h of chest pain if possible, mostly with glycoprotein IIb/IIIa blocker therapy.</p> <p>All participants were treated with clopidogrel for 6 months (300 mg periprocedurally, maintenance dose 75 mg per day) irrespective of stents used. Otherwise, participants received usual standard of care, i.e. aspirin 100 mg daily and a statin; other drugs, including glycoprotein IIb/IIIa blockers, were used as clinically indicated.</p>
Outcomes	The primary endpoint was cost-effectiveness after 6 months, with effectiveness defined as reduction of major adverse cardiac events, i.e. cardiac death, non-fatal myocardial infarction, and target vessel revascularisation. Mortality from other causes was also analysed, but these events were not included as primary endpoints. Non-fatal myocardial infarction apart from the index intervention was diagnosed on the basis of a typical rise (and fall) of cardiac enzyme concentration, typical chest pain, and new pathological Q-waves or ischaemic ST-T wave changes, or both, in the electrocardiogram, according to current guidelines. Quality of life was assessed with the EQ-5D.
Notes	<p>No cross-overs reported. No ClinicalTrials.gov number or other pre-published protocol found.</p> <p>We contacted the author regarding random sequence generation, pre-published protocol, blinding of participants and personnel, composite outcome of cardiac death and myocardial infarction, data after 6 months, and data for STEMI, NSTEMI, unstable and stable angina separately by email: pfisterer@email.ch on 19 March 2017.</p>

BASKET 2005b

Methods	Parallel design, single-centre, randomised clinical trial with 2 different experimental arms in Switzerland
Participants	<p>826 participants treated with PCI and stenting at the University Hospital of Basel, Switzerland between 5 May 2003 and 31 May 2004 were included in BASKET, irrespective of indication for PCI.</p> <p>Male:female = 320:84</p> <p>Mean age = 64 years</p> <p>Exclusion criteria: a target vessel diameter of 4 mm or greater, as the largest available DES size was 3.5 mm (n = 23); presence of restenotic lesions, given the different causes and outcome of restenotic lesions (n = 49); and no consent, primarily due to patients' or referring physicians' preference for DES before angiography and inability of patients to give informed consent (n = 90).</p>
Interventions	<p>Experimental group 2: paclitaxel-coated Taxus stent.</p> <p>Control group: cobalt chromium-based Vision/Pixel stent.</p> <p>Co-intervention: in participants with ST-elevation myocardial infarction, primary PCI was the treatment of choice. Participants with non-ST-elevation myocardial infarction or unstable angina were treated urgently within 24 h of chest pain if possible, mostly with glycoprotein IIb/IIIa blocker therapy.</p> <p>All participants were treated with clopidogrel for 6 months (300 mg periprocedurally, maintenance dose 75 mg per day) irrespective of stents used. Otherwise, participants received usual standard of care, i.e. aspirin 100 mg daily and a statin; other drugs, including glycoprotein IIb/IIIa blockers, were used as clinically indicated.</p>
Outcomes	The primary endpoint was cost-effectiveness after 6 months, with effectiveness defined as reduction of major adverse cardiac events, i.e. cardiac death, non-fatal myocardial infarction, and target vessel revascularisation. Mortality from other causes was also analysed, but these events were not included as primary endpoints. Non-fatal myocardial infarction apart from the index intervention was diagnosed on the basis of a typical rise (and fall) of cardiac enzyme concentration, typical chest pain, and new pathological Q-waves or ischaemic ST-T wave changes, or both, in the electrocardiogram, according to current guidelines. Quality of life was assessed with the EQ-5D.
Notes	<p>No cross-overs reported. No ClinicalTrials.gov number or other pre-published protocol found.</p> <p>We contacted the author regarding random sequence generation, pre-published protocol, blinding of participants and personnel, composite outcome of cardiac death and myocardial infarction, data after 6 months, and data for STEMI, NSTEMI, unstable and stable angina separately by email: pfisterer@email.ch on 19 March 2017.</p>

BASKET-PROVE I 2010

Methods	Parallel-design randomised clinical trial in Switzerland, Denmark, Austria, and Italy
Participants	<p>2314 participants who presented with chronic or acute coronary disease, who underwent angioplasty with stenting, and who required only stents that were 3.0 mm or more in diameter. No restrictions were placed on the number of treated lesions or vessels, the length of treated lesions, or the number of stents placed.</p> <p>Male:female = 1749:565</p> <p>Mean age = 66.3 years</p> <p>Exclusion criteria: cardiogenic shock; in-stent restenosis or thrombosis of stents placed before the study; unprotected left main coronary artery (i.e. with no functioning bypass graft) or substantial</p>

BASKET-PROVE I 2010 (Continued)

	stenosis in a bypass graft; plans for any surgery within 12 months; a need for oral anticoagulation, an increased risk of bleeding, or known intolerance to or suspected noncompliance with long-term antiplatelet therapy; or circumstances that would have made follow-up impossible. In addition, patients requiring stents larger than 4.0 mm in diameter were excluded because no sirolimus-eluting stents of this size were available.
Interventions	<p>Experimental group 1: first-generation sirolimus-eluting stent (Cypher Select, Cordis).</p> <p>Experimental group 2: second-generation everolimus-eluting stent (Xience V, Abbott Vascular).</p> <p>Control group: bare-metal (cobalt–chromium) stent (Vision, Abbott Vascular).</p> <p>Co-intervention: all participants were prescribed aspirin at a daily dose of 75 to 100 mg indefinitely and clopidogrel at a daily dose of 75 mg for 1 year, after a loading dose of 300 mg or 600 mg, regardless of stent type. Therapeutic agents for secondary prevention, such as statins, were prescribed according to current guidelines.</p>
Outcomes	<p>Primary outcomes: death from cardiac causes or non-fatal myocardial infarction at 2 years.</p> <p>Secondary outcomes: efficacy endpoint of target vessel revascularisation and the safety endpoint of late death from cardiac causes or non-fatal myocardial infarction.</p>
Notes	Current Controlled Trials number ISRCTN72444640. We contacted the authors at Christoph.kaiser@usb.ch on 13 April 2017, but have received no reply.

BASKET-PROVE II 2015

Methods	Randomised clinical trial with a parallel/factorial design in Switzerland, Denmark, Germany, and Austria
Participants	<p>2299 participants with chronic or acute coronary artery disease requiring angioplasty and stenting with stents ≥ 3.0 mm in diameter by visual assessment. No restrictions applied to the number of treated lesions or vessels, length of treated lesions, or number of stents implanted.</p> <p>Male:female = 1780:519</p> <p>Mean age = 62.3 years</p> <p>Exclusion criteria: cardiogenic shock; in-stent restenosis or thrombosis; unprotected left main coronary artery or bypass graft disease; planned surgery within 12 months; need for oral anticoagulation, increased bleeding risk, known intolerance to or suspected noncompliance with long-term antiplatelet drug therapy; history of transient ischaemic attack or stroke; or circumstances that would have made follow-up impossible.</p>
Interventions	<p>Experimental group 1: second-generation Biolimus A9–eluting biodegradable-polymer stainless-steel DES (Nobori, Terumo).</p> <p>Experimental group 2: second-generation everolimus-eluting durable-polymer cobalt-chromium DES (Xience Prime, Abbott Vascular).</p> <p>Control group: generation thin-strut BMS coated with a biocompatible silicone-carbide layer (PRO-Kinetic, Biotronik).</p> <p>Co-intervention: all participants received a loading dose of 60 mg prasugrel with a maintenance dose of 10 mg daily, risk-adjusted to 5 mg in participants aged > 75 years or body weight < 60 kg. Prasugrel was prescribed for 12 months after stenting with DES and for participants with acute coronary syndrome and for 4 weeks after elective stenting with BMS. To ensure adherence to antiplatelet therapy, participants were given the drug in monthly packages at regular intervals from</p>

BASKET-PROVE II 2015 (Continued)

	the study centres. No further prasugrel was provided after 12 months. All other concomitant treatments were prescribed according to current guidelines.
Outcomes	<p>Primary outcomes: major adverse cardiac events (i.e. a combination of cardiac death, non-fatal myocardial infarction, or clinically driven non-myocardial infarction-related target vessel revascularisation).</p> <p>Secondary outcomes: TVR, stent thrombosis, myocardial infarction, cardiac death.</p>
Notes	ClinicalTrials.gov NCT01166685. We contacted the authors at Christoph.kaiser@usb.ch on 13 April 2017, but have received no reply.

EAGLE 2006

Methods	Parallel-design randomised clinical trial in Germany
Participants	<p>129 participants with a single, de-novo, coronary stenosis (diameter 2.5 to 4.0 mm, length of stenosis $\leq$ 25 mm).</p> <p>Male:female = 65:64 (50% male)</p> <p>Mean age = not stated</p> <p>Exclusion criteria: none stated.</p>
Interventions	<p>Experimental group: Axxion (paclitaxel) polymer-free stent.</p> <p>Control group: bare-metal stent.</p> <p>Co-intervention: none stated.</p>
Outcomes	<p>Primary: MACE after 30 days.</p> <p>Secondary: MACE and angina status after 6 months, binary restenosis and late luminal loss after 6 to 9 months.</p>
Notes	No contact information could be obtained.

Erglis 2007

Methods	Parallel-design randomised clinical trial in Latvia and Australia
Participants	<p>103 participants with clinically symptomatic left main coronary artery disease with angiographic evidence of 50% diameter stenosis of left main coronary artery suitable for stent implantation. The left main coronary artery was considered unprotected if there was no CABG to left anterior descending artery branches or left circumflex branches. All patients were good candidates for CABG.</p> <p>Male:female = 86:17</p> <p>Mean age = 61.7 years</p> <p>Exclusion criteria: none stated.</p>
Interventions	<p>Experimental group: Taxus Express paclitaxel-eluting stent.</p> <p>Control group: Express or Liberte BMS.</p>

Erglis 2007 (Continued)

Co-intervention: all participants received cutting balloon intervention before stenting. A maximum of 24 hours before the PCI procedure, all participants received 100 mg aspirin and a loading dose (300 mg) of clopidogrel. Participants were required to take clopidogrel 75 mg daily for at least 6 months after the PCI procedure. During the procedure, all participants were administered 10,000 IU of heparin or low-molecular-weight heparin, depending on the participant's weight. Glycoprotein IIb/IIIa inhibitors were given at the operator's discretion.

Outcomes	Intravascular ultrasound characteristics, death, cardiac death, MI, stent thrombosis, TLR, MACE
Notes	We contacted the authors at a.a.erglis@stradini.lv regarding separate data on ACS participants.

FUTURE I 2004

Methods	Parallel-design randomised clinical trial in Germany
Participants	<p>42 participants with de novo coronary lesions with a reference diameter between 2.75 and 4.0 mm with lesion length 18 mm.</p> <p>Male:female = 36:6</p> <p>Mean age = 65 years</p> <p>Exclusion criteria: none stated.</p>
Interventions	<p>Experimental group: S-Stent (Biosensors International) everolimus eluting with a bioresorbable polymer.</p> <p>Control group: bare-metal stent.</p> <p>Co-intervention: all participants received aspirin (325 mg/d) and clopidogrel (300 mg loading dose immediately and 75 mg/d for 6 months).</p>
Outcomes	<p>Primary outcomes: the primary endpoint was MACE, including death, CABG to the target vessel, Q-wave and non-Q-wave myocardial infarction, and TLR at 30 days.</p> <p>Secondary outcomes: the secondary endpoint was to compare quantitative angiographic and intravascular ultrasound observation within the vessel as well as the clinical performance of the everolimus-eluting stent with regard to device success, MACE, and restenosis rate at 6-month follow-up.</p>
Notes	No registration with ClinicalTrials.gov. We contacted the authors at ivus@crci.stanford.edu on 15 April 2017, but have received no reply.

FUTURE II 2004

Methods	Parallel-design randomised clinical trial at 3 sites in Germany
Participants	<p>64 participants with de novo coronary lesions with a reference diameter between 2.75 and 4.0 mm with lesion length 18 mm.</p> <p>Male:female = not stated</p> <p>Mean age = not stated</p> <p>Exclusion criteria: none stated.</p>

FUTURE II 2004 (Continued)

Interventions	<p>Experimental group: S-Stent (Biosensors International) everolimus eluting with a bioresorbable polymer.</p> <p>Control group: bare-metal stent.</p> <p>Co-intervention: aspirin and clopidogrel for 6 months.</p>
Outcomes	MACE, including death, CABG to the target vessel, Q-wave and non-Q-wave myocardial infarction, and TLR at 1 and 6 months. Stent thrombosis and restenosis.
Notes	No registration with ClinicalTrials.gov. We contacted the authors at ivus@crci.stanford.edu on 15 April 2017, but have received no reply.

Han 2007

Methods	Parallel-design randomised clinical trial in China
Participants	<p>200 participants with coronary heart disease (36.5% acute myocardial infarction).</p> <p>Male:female = 144:56</p> <p>Mean age = not stated</p> <p>Exclusion criteria: none stated.</p>
Interventions	<p>Experimental group: Janus tacrolimus-eluting stent.</p> <p>Control group: Tecnic Carbstent (bare-metal stent).</p> <p>Co-intervention: clopidogrel for 4 months and aspirin lifelong.</p>
Outcomes	MACE, cardiovascular mortality, myocardial infarction, TLR
Notes	Abstract only. No contact information could be obtained.

Li 2004

Methods	Parallel-design randomised clinical trial in China
Participants	<p>152 participants with coronary small vessel lesions (inside diameter 3.0 mm on quantitative coronary angiography).</p> <p>Male:female = not stated</p> <p>Mean age = not stated</p> <p>Exclusion criteria: none stated.</p>
Interventions	<p>Experimental group: sirolimus-eluting Cypher stents.</p> <p>Control group: uncoated stents.</p> <p>Co-intervention: standard antiplatelet and anticoagulation agents and individual modifications as needed.</p>
Outcomes	Mortality, stent thrombosis, restenosis

Li 2004 (Continued)

Notes	Abstract only. No contact information could be found.
-------	---

NORSTENT 2016

Methods	Parallel-design randomised clinical trial at 8 sites in Norway
Participants	<p>9013 men and women who were at least 18 years of age and who presented with stable angina or an acute coronary syndrome, had lesions in native coronary arteries or coronary artery grafts (all of which were amenable for implantation of either drug-eluting stents or bare-metal stents), had a Norwegian national identification number and were able to communicate in Norwegian, and provided informed consent.</p> <p>Male:female = 6657:2356</p> <p>Mean age = 62.6 years</p> <p>Exclusion criteria: "Patients were excluded if they had previously been treated with a coronary stent, had a bifurcation lesion requiring treatment with a two-stent technique, had a serious medical condition other than coronary artery disease with a life expectancy of less than 5 years, were participating in another randomized trial, had intolerable side effects to any drug in use during PCI or contraindications to long-term dual-antiplatelet therapy or had been prescribed warfarin, or were not able to follow the trial protocol, as judged by the investigator."</p>
Interventions	<p>Experimental group: any commercially available stent in Norway (Promus, Xience, Endeavor, Endeavor Resolute, Cypher, Taxus).</p> <p>Control group 1: any commercially available stent in Norway (Driver, Integrity, Liberte, Multi-Link Vision, Multi-Link 8, Chrono Carbostent, Multi-Link Flexmaster, Graftmaster, Coroflex Blue, Multi-Link Zeta).</p> <p>Co-intervention: "All the patients in the two groups were prescribed aspirin at a daily dose of 75 mg indefinitely and clopidogrel at a daily dose of 75 mg for 9 months after the procedure regardless of the randomized assignment or the indication for PCI."</p>
Outcomes	<p>The primary outcome was a composite of death from any cause and non-fatal spontaneous myocardial infarction at a median follow-up of 5 years, as specified in an amendment to the protocol made by the steering committee in May 2012.</p> <p>Secondary outcomes were subcategories of death; fatal and non-fatal spontaneous and periprocedural myocardial infarction and stroke; hospitalisation for unstable angina pectoris; revascularisation of a target lesion, target vessel, or non-target vessel with PCI or CABG; definite stent thrombosis; major bleeding episodes; and health-related quality of life.</p>
Notes	ClinicalTrials.gov NCT00811772 (prospectively). We contacted the authors at kaare.harald.bonaa@ntnu.no on 12 April 2017. We received an initial reply and were informed we would receive data for ACS and stable angina participants separately. We have not yet received these data.

PRODIGY 2014

Methods	Randomised clinical trial with a parallel design in Italy
Participants	2013 participants ≥ 18 years old with chronic stable coronary artery disease or acute coronary syndrome, including non-ST-elevation and ST-elevation MI. Patients were eligible if they had ≥ 1 lesion with a diameter stenosis of $\geq 50\%$ that was suitable for coronary stent implantation in a vessel with a reference vessel diameter of ≥ 2.25 mm and were undergoing elective, urgent, or emergent coronary angioplasty with intended stent implantation.

Drug-eluting stents versus bare-metal stents for acute coronary syndrome (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

PRODIGY 2014 (Continued)

Male:female = 1538:475

Mean age = 68.3 years

Exclusion criteria: known allergy to acetylsalicylic acid or clopidogrel, planned surgery within 24 months of PCI unless the dual antiplatelet therapy could be maintained throughout the perisurgical period, history of bleeding diathesis, major surgery within 15 days, active bleeding or previous stroke in the last 6 months, concomitant or foreseeable need for oral anticoagulation therapy, pregnancy, life expectancy less than 24 months, participation in another trial, and inability to provide informed consent.

Interventions	<p>Experimental group 1: Endeavor Sprint zotarolimus-eluting stent (Medtronic, Santa Rosa, CA).</p> <p>Experimental group 2: Taxus paclitaxel-eluting stent (Boston Scientific, Natick, MA).</p> <p>Experimental group 3: Xience V everolimus-eluting stent (Abbott Vascular, Santa Clara, CA).</p> <p>Control group: third-generation thin-strut BMS.</p> <p>Co-intervention: all participants received aspirin (160 to 325 mg orally or 500 mg intravenously as a loading dose and then 80 to 160 mg orally indefinitely) and clopidogrel as follows: 300 or 600 mg orally as a loading dose to be given preferably before PCI but not later than at the time the participant leaves the catheterisation laboratory and then 75 mg/d for the treatment duration provided by the randomisation scheme. At 30 days, participants in each stent group were randomised in a balanced fashion to 6 months of dual antiplatelet treatment (SHORT arm) versus prolonging aspirin and clopidogrel for 24 months (LONG arm).</p>
Outcomes	Myocardial infarction, death, stroke, cardiac death, stent thrombosis, TVR
Notes	<p>We contacted the authors at m.valgimigli@erasmusmc.nl regarding separate data.</p> <p>All participants were re-randomised after 30 days to 6 or 24 months clopidogrel.</p> <p>ClinicalTrials.gov NCT00611286.</p>

SES-SMART 2004

Methods	Randomised clinical trial with a parallel/factorial design in Italy
Participants	<p>257 participants aged 18 years or older, with a documented diagnosis of acute coronary syndrome (without persistent ST-segment elevation), stable angina pectoris, or silent myocardial ischaemia as shown by exercise stress test. Additional eligibility criteria were the presence of a single, previously untreated 50% to 99% target lesion in a native coronary artery 2.75 mm in diameter or less, which could be completely covered by a single stent (maximum length 33 mm). The patients could have had single-vessel or multivessel disease, but in the latter case had to have the non-randomised lesion located in other coronary vessels.</p> <p>Male:female = 184:73</p> <p>Mean age = 63.6 years</p> <p>Exclusion criteria: recent ST-segment elevation acute coronary syndrome (within the previous 15 days), severe calcifications or thrombus-containing lesions, a left ventricular ejection fraction less than 30%, and known allergies to aspirin, clopidogrel, ticlopidine, heparin, stainless steel, contrast agents, or sirolimus.</p>
Interventions	Experimental group: sirolimus-eluting stent (Cypher balloon-expandable stent; Cordis, Miami Lakes, FL).

SES-SMART 2004 (Continued)

	<p>Control group 1: uncoated stent of identical architecture and radiographic appearance (BX Sonic balloon-expandable stent; Cordis)</p> <p>Co-intervention: all participants received oral aspirin once daily and clopidogrel (a loading dose of 300 mg at least 2 hours before the procedure). Participants who had been pre-treated with ticlopidine (250 mg twice a day) or clopidogrel (75 mg once daily) for at least 72 hours did not receive a clopidogrel loading dose. During the procedure, heparin was given as a bolus at 70 U/kg, with additional boluses to maintain an activated clotting time of more than 250 seconds. The use of glycoprotein IIb/IIIa inhibitors was encouraged but left to the discretion of the attending physician. Heparin administration was discontinued immediately after the procedure.</p>
Outcomes	<p>Primary outcome: 8-month angiographic binary in-segment restenosis rate.</p> <p>Secondary outcomes: procedural success, 8-month in-segment minimal luminal diameter, late luminal loss, late loss index, and major adverse cardiac and cerebrovascular events.</p>
Notes	<p>We contacted the authors at cardiologia.parma@ao.pr.it regarding separate data for NSTEMI participants and stable participants, but have received no reply.</p>

XIMA 2014

Methods	<p>Randomised clinical trial with a parallel/factorial design in the UK and Spain</p>
Participants	<p>800 participants > 80 years of age. Coronary disease warranting use of DES (15 mm long or < 3 mm wide) was a requirement. Patients with non-ST-segment elevation myocardial infarction, unstable angina, and stable angina were eligible to participate.</p> <p>Male:female = 481:319</p> <p>Mean age = 83.5</p> <p>Exclusion criteria: acute ST-segment elevation myocardial infarction, cardiogenic shock, thrombocytopenia ($< 50 \times 10^9/\text{mm}^3$), poor life expectancy, gastrointestinal haemorrhage < 3 months, or previous intracerebral bleeding.</p>
Interventions	<p>Experimental group: second-generation everolimus-eluting stents (Xience, Abbott Vascular, Santa Clara, CA).</p> <p>Control group: bare-metal Vision stents (Abbott Vascular).</p> <p>Co-intervention: before PCI, loading doses of aspirin 300 mg and clopidogrel 600 mg were given unless the participants were established on these drugs. Long-term treatment with warfarin was not a contraindication, but caution was emphasised. The use of glycoprotein IIb/IIIa inhibitors was at the discretion of the operator, but caution was urged.</p>
Outcomes	<p>Death, haemorrhage, MI, TVR</p>
Notes	<p>ISRCTN92243650. We contacted the authors at adam.debelder@bsuh.nhs.uk regarding separate data on participants with acute coronary syndrome and participants with stable angina pectoris as well as regarding blinding.</p>

ZEUS 2015

Methods	<p>Randomised clinical trial with a parallel/factorial design in Italy, Switzerland, Portugal, and Hungary</p>
---------	--

Drug-eluting stents versus bare-metal stents for acute coronary syndrome (Review)

ZEUS 2015 (Continued)

Participants	<p>1606 participants with high bleeding risk and/or the presence of relative or absolute contraindications to long-term dual antiplatelet treatment and/or high thrombosis risk due to systemic disorders or planned non-cardiac surgery and/or low restenosis risk based on angiographic findings.</p> <p>Male:female = 1133:473</p> <p>Mean age = 73.7 years</p> <p>Exclusion criteria: pregnancy; women of childbearing potential must have had a negative pregnancy test (urine or serum human chorionic gonadotropin), preferably < 24 hours, but at a minimum within 7 days prior to randomisation. Patients who were unable to give informed consent or who were unwilling to undergo planned follow-up through 12 months, or both.</p>
Interventions	<p>Experimental group: Zotarolimus-Eluting Endeavor Sprint stent (E-ZES) (Santa Rosa, CA).</p> <p>Control group: all types of third-generation thin-strut BMS.</p> <p>Co-intervention: all eligible participants received aspirin (160 to 325 mg orally or 500 mg intravenously as a loading dose and then 80 to 160 mg orally per day) and clopidogrel (300 or 600 mg orally as a loading dose followed by 75 mg/day) or prasugrel (60 mg loading dose followed by 10 or 5 mg/day) or ticagrelor (180 mg loading dose followed by 90 mg twice daily). Participants who were not eligible for dual antiplatelet treatment were treated with either aspirin or clopidogrel (or prasugrel or ticagrelor) monotherapy.</p>
Outcomes	All-cause death, non-fatal MI, or any target vessel revascularisation, stent thrombosis, bleeding
Notes	ClinicalTrials.gov NCT01385319 (prospectively). We contacted the authors at marco.valgimigli@insel.ch; vlgmrc@unife.it on 13 April 2017, but have received no reply.

ACS: acute coronary syndrome
BMS: bare-metal stent
CABG: coronary artery bypass grafting
DES: drug-eluting stent
MACE: major adverse cardiac events
MI: myocardial infarction
NSTEMI: non-ST-segment elevation myocardial infarction
PCI: percutaneous coronary intervention
STEMI: ST-segment elevation myocardial infarction
TLR: target lesion revascularisation
TVR: target vessel revascularisation

DATA AND ANALYSES

Comparison 1. Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.1 All-cause mortality	22	11250	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.78, 1.03]
1.2 All-cause mortality best-worst	22	11775	Risk Ratio (M-H, Random, 95% CI)	0.60 [0.44, 0.80]
1.3 All-cause mortality worst-best	22	11775	Risk Ratio (M-H, Random, 95% CI)	1.27 [0.99, 1.64]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.4 All-cause mortality according to type of drug-eluting stent	22	11250	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.78, 1.03]
1.4.1 Biodegradable (Biolimus)	1	1157	Risk Ratio (M-H, Random, 95% CI)	0.89 [0.54, 1.45]
1.4.2 Everolimus	1	1458	Risk Ratio (M-H, Random, 95% CI)	0.73 [0.54, 1.00]
1.4.3 Paclitaxel	8	4468	Risk Ratio (M-H, Random, 95% CI)	0.89 [0.71, 1.12]
1.4.4 Sirolimus	9	3113	Risk Ratio (M-H, Random, 95% CI)	0.89 [0.68, 1.16]
1.4.5 Zotarolimus	1	44	Risk Ratio (M-H, Random, 95% CI)	Not estimable
1.4.6 Mixed	1	626	Risk Ratio (M-H, Random, 95% CI)	1.34 [0.91, 1.98]
1.4.7 Unclear	1	384	Risk Ratio (M-H, Random, 95% CI)	0.43 [0.02, 7.39]
1.5 All-cause mortality according to type of ACS	22	11250	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.78, 1.03]
1.5.1 STEMI	21	11171	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.78, 1.03]
1.5.2 Acute coronary syndrome	1	79	Risk Ratio (M-H, Random, 95% CI)	Not estimable
1.6 All-cause mortality according to length of maximum follow-up	22	11250	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.78, 1.03]
1.6.1 Less or equal to 6 months	4	726	Risk Ratio (M-H, Random, 95% CI)	0.85 [0.29, 2.53]
1.6.2 Between 6 and 12 months	6	1315	Risk Ratio (M-H, Random, 95% CI)	0.81 [0.54, 1.21]
1.6.3 Between 1 and 3 years	3	703	Risk Ratio (M-H, Random, 95% CI)	0.99 [0.55, 1.76]
1.6.4 More or equal to 3 years	8	8490	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.77, 1.05]
1.6.5 Unclear	1	16	Risk Ratio (M-H, Random, 95% CI)	Not estimable
1.7 All-cause mortality according to registration status	22	11250	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.78, 1.03]
1.7.1 Pre-registration	3	1301	Risk Ratio (M-H, Random, 95% CI)	0.85 [0.52, 1.39]
1.7.2 Post-registration	13	9028	Risk Ratio (M-H, Random, 95% CI)	0.89 [0.77, 1.03]
1.7.3 No registration	6	921	Risk Ratio (M-H, Random, 95% CI)	1.09 [0.54, 2.20]
1.8 Serious adverse events	23	11724	Risk Ratio (M-H, Fixed, 95% CI)	0.80 [0.74, 0.86]
1.9 Serious adverse events best-worst	23	12249	Risk Ratio (M-H, Fixed, 95% CI)	0.67 [0.62, 0.71]
1.10 Serious adverse events worst-best	23	12249	Risk Ratio (M-H, Fixed, 95% CI)	0.98 [0.91, 1.05]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.11 Serious adverse events according to type of drug-eluting stent	23	11724	Risk Ratio (M-H, Fixed, 95% CI)	0.80 [0.74, 0.86]
1.11.1 Biodegradable (Biolimus)	1	1157	Risk Ratio (M-H, Fixed, 95% CI)	0.75 [0.58, 0.98]
1.11.2 Everolimus	2	1932	Risk Ratio (M-H, Fixed, 95% CI)	0.82 [0.69, 0.97]
1.11.3 Paclitaxel	8	4468	Risk Ratio (M-H, Fixed, 95% CI)	0.86 [0.76, 0.96]
1.11.4 Sirolimus	9	3113	Risk Ratio (M-H, Fixed, 95% CI)	0.74 [0.64, 0.84]
1.11.5 Zotarolimus	1	44	Risk Ratio (M-H, Fixed, 95% CI)	1.33 [0.17, 10.70]
1.11.6 Mixed	1	626	Risk Ratio (M-H, Fixed, 95% CI)	0.74 [0.54, 1.01]
1.11.7 Unclear	1	384	Risk Ratio (M-H, Fixed, 95% CI)	0.93 [0.52, 1.69]
1.12 Serious adverse events according to type of ACS	23	11724	Risk Ratio (M-H, Fixed, 95% CI)	0.80 [0.74, 0.86]
1.12.1 STEMI	21	11171	Risk Ratio (M-H, Fixed, 95% CI)	0.80 [0.75, 0.86]
1.12.2 Acute coronary syndrome	1	79	Risk Ratio (M-H, Fixed, 95% CI)	0.58 [0.29, 1.15]
1.12.3 NSTEMI	1	474	Risk Ratio (M-H, Fixed, 95% CI)	0.78 [0.50, 1.23]
1.13 Serious adverse events according to length of maximum follow-up	23	11724	Risk Ratio (M-H, Fixed, 95% CI)	0.80 [0.74, 0.86]
1.13.1 Less or equal to 6 months	4	726	Risk Ratio (M-H, Fixed, 95% CI)	0.67 [0.45, 0.99]
1.13.2 Between 6 and 12 months	7	1748	Risk Ratio (M-H, Fixed, 95% CI)	0.85 [0.69, 1.05]
1.13.3 Between 1 and 3 years	3	744	Risk Ratio (M-H, Fixed, 95% CI)	0.64 [0.47, 0.88]
1.13.4 More or equal to 3 years	8	8490	Risk Ratio (M-H, Fixed, 95% CI)	0.81 [0.75, 0.88]
1.13.5 Unclear	1	16	Risk Ratio (M-H, Fixed, 95% CI)	Not estimable
1.14 Serious adverse events according to registration status	23	11724	Risk Ratio (M-H, Fixed, 95% CI)	0.80 [0.74, 0.86]
1.14.1 Pre-registration	3	1301	Risk Ratio (M-H, Fixed, 95% CI)	0.71 [0.55, 0.91]
1.14.2 Post-registration	14	9502	Risk Ratio (M-H, Fixed, 95% CI)	0.81 [0.75, 0.87]
1.14.3 No registration	6	921	Risk Ratio (M-H, Fixed, 95% CI)	0.83 [0.57, 1.20]
1.15 Major cardiovascular events	20	10939	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.83, 1.11]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.16 Major cardiovascular events best-worst	20	11596	Risk Ratio (M-H, Random, 95% CI)	0.53 [0.39, 0.72]
1.17 Major cardiovascular events worst-best	20	11596	Risk Ratio (M-H, Random, 95% CI)	1.40 [0.97, 2.00]
1.18 Major cardiovascular events according to type of drug-eluting stent	20	10939	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.83, 1.11]
1.18.1 Biodegradable (Biolimus)	1	1104	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.54, 1.49]
1.18.2 Everolimus	1	1458	Risk Ratio (M-H, Random, 95% CI)	0.85 [0.58, 1.24]
1.18.3 Paclitaxel	7	4305	Risk Ratio (M-H, Random, 95% CI)	1.01 [0.81, 1.26]
1.18.4 Sirolimus	8	3018	Risk Ratio (M-H, Random, 95% CI)	0.89 [0.64, 1.24]
1.18.5 Zotarolimus	1	44	Risk Ratio (M-H, Random, 95% CI)	Not estimable
1.18.6 Mixed	1	626	Risk Ratio (M-H, Random, 95% CI)	1.04 [0.60, 1.81]
1.18.7 Unclear	1	384	Risk Ratio (M-H, Random, 95% CI)	1.36 [0.54, 3.40]
1.19 Major cardiovascular events according to type of ACS	20	10939	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.83, 1.11]
1.19.1 STEMI	19	10860	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.83, 1.12]
1.19.2 Acute coronary syndrome	1	79	Risk Ratio (M-H, Random, 95% CI)	0.68 [0.12, 3.87]
1.20 Major cardiovascular events according to length of maximum follow-up	20	10939	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.83, 1.11]
1.20.1 Less or equal to 6 months	3	563	Risk Ratio (M-H, Random, 95% CI)	1.04 [0.48, 2.28]
1.20.2 Between 1 and 3 years	9	2018	Risk Ratio (M-H, Random, 95% CI)	0.89 [0.69, 1.16]
1.20.3 More or equal to 3 years	8	8358	Risk Ratio (M-H, Random, 95% CI)	0.99 [0.82, 1.19]
1.21 Major cardiovascular events according to registration status	20	10939	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.83, 1.11]
1.21.1 Pre-registration	3	1248	Risk Ratio (M-H, Random, 95% CI)	0.86 [0.52, 1.42]
1.21.2 Post-registration	13	8949	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.83, 1.14]
1.21.3 No registration	4	742	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.47, 1.93]
1.22 Cardiovascular mortality	15	9248	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.76, 1.09]
1.23 Cardiovascular mortality best-worst	15	9742	Risk Ratio (M-H, Random, 95% CI)	0.50 [0.32, 0.77]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.24 Cardiovascular mortality worst-best	15	9742	Risk Ratio (M-H, Random, 95% CI)	1.49 [1.06, 2.11]
1.25 Cardiovascular mortality according to type of drug-eluting stent	15	9248	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.76, 1.09]
1.25.1 Biodegradable (Biolimus)	1	1104	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.54, 1.49]
1.25.2 Everolimus	1	1458	Risk Ratio (M-H, Random, 95% CI)	0.85 [0.58, 1.24]
1.25.3 Paclitaxel	6	4141	Risk Ratio (M-H, Random, 95% CI)	0.86 [0.65, 1.13]
1.25.4 Sirolimus	5	1875	Risk Ratio (M-H, Random, 95% CI)	0.81 [0.52, 1.25]
1.25.5 Zotarolimus	1	44	Risk Ratio (M-H, Random, 95% CI)	Not estimable
1.25.6 Mixed	1	626	Risk Ratio (M-H, Random, 95% CI)	2.40 [1.17, 4.93]
1.26 Cardiovascular mortality according to type of ACS	15	9248	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.76, 1.09]
1.26.1 STEMI	15	9248	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.76, 1.09]
1.26.2 Acute coronary syndrome	0	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
1.27 Cardiovascular mortality according to length of maximum follow-up	15	9248	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.76, 1.09]
1.27.1 Less or equal to 6 months	1	100	Risk Ratio (M-H, Random, 95% CI)	0.21 [0.01, 4.23]
1.27.2 Between 6 and 12 months	6	1584	Risk Ratio (M-H, Random, 95% CI)	0.94 [0.65, 1.35]
1.27.3 Between 1 and 3 years	2	270	Risk Ratio (M-H, Random, 95% CI)	0.56 [0.23, 1.32]
1.27.4 More or equal to 3 years	6	7294	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.72, 1.27]
1.28 Cardiovascular mortality according to registration status	15	9248	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.76, 1.09]
1.28.1 Pre-registration	3	1248	Risk Ratio (M-H, Random, 95% CI)	0.86 [0.52, 1.42]
1.28.2 Post-registration	10	7806	Risk Ratio (M-H, Random, 95% CI)	0.93 [0.73, 1.18]
1.28.3 No registration	2	194	Risk Ratio (M-H, Random, 95% CI)	0.71 [0.16, 3.16]
1.29 Myocardial infarction	19	10217	Risk Ratio (M-H, Random, 95% CI)	0.98 [0.82, 1.18]
1.30 Myocardial infarction best-worst	19	10851	Risk Ratio (M-H, Random, 95% CI)	0.46 [0.33, 0.66]
1.31 Myocardial infarction worst-best	19	10851	Risk Ratio (M-H, Random, 95% CI)	1.56 [1.00, 2.45]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.32 Myocardial infarction according to type of drug-eluting stent	19	10217	Risk Ratio (M-H, Random, 95% CI)	0.98 [0.82, 1.18]
1.32.1 Biodegradable (Biolimus)	1	1118	Risk Ratio (M-H, Random, 95% CI)	0.64 [0.36, 1.15]
1.32.2 Everolimus	1	1458	Risk Ratio (M-H, Random, 95% CI)	1.29 [0.79, 2.11]
1.32.3 Paclitaxel	7	4305	Risk Ratio (M-H, Random, 95% CI)	1.05 [0.75, 1.46]
1.32.4 Sirolimus	7	2282	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.58, 1.39]
1.32.5 Zotarolimus	1	44	Risk Ratio (M-H, Random, 95% CI)	Not estimable
1.32.6 Mixed	1	626	Risk Ratio (M-H, Random, 95% CI)	0.74 [0.40, 1.36]
1.32.7 Unclear	1	384	Risk Ratio (M-H, Random, 95% CI)	1.36 [0.54, 3.40]
1.33 Myocardial infarction according to type of ACS	19	10217	Risk Ratio (M-H, Random, 95% CI)	0.98 [0.82, 1.18]
1.33.1 STEMI	18	10138	Risk Ratio (M-H, Random, 95% CI)	0.98 [0.82, 1.18]
1.33.2 Acute coronary syndrome	1	79	Risk Ratio (M-H, Random, 95% CI)	0.68 [0.12, 3.87]
1.34 Myocardial infarction according to length of maximum follow-up	19	10217	Risk Ratio (M-H, Random, 95% CI)	0.98 [0.82, 1.18]
1.34.1 Less or equal to 6 months	3	563	Risk Ratio (M-H, Random, 95% CI)	1.17 [0.52, 2.63]
1.34.2 Between 6 and 12 months	7	1748	Risk Ratio (M-H, Random, 95% CI)	0.98 [0.58, 1.66]
1.34.3 Between 1 and 3 years	2	270	Risk Ratio (M-H, Random, 95% CI)	0.55 [0.24, 1.25]
1.34.4 More or equal to 3 years	7	7636	Risk Ratio (M-H, Random, 95% CI)	1.00 [0.80, 1.25]
1.35 Myocardial infarction according to registration status	19	10217	Risk Ratio (M-H, Random, 95% CI)	0.98 [0.82, 1.18]
1.35.1 Pre-registration	3	1262	Risk Ratio (M-H, Random, 95% CI)	0.64 [0.36, 1.15]
1.35.2 Post-registration	12	8213	Risk Ratio (M-H, Random, 95% CI)	1.03 [0.85, 1.26]
1.35.3 No registration	4	742	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.44, 2.12]
1.36 Stent thrombosis	21	11350	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.80, 1.16]
1.37 Stent thrombosis best-worst	21	12007	Risk Ratio (M-H, Random, 95% CI)	0.47 [0.32, 0.67]
1.38 Stent thrombosis worst-best	21	12007	Risk Ratio (M-H, Random, 95% CI)	1.55 [1.12, 2.15]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.39 Stent thrombosis according to type of drug-eluting stent	21	11350	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.80, 1.16]
1.39.1 Biodegradable (Biolimus)	1	1104	Risk Ratio (M-H, Random, 95% CI)	0.72 [0.40, 1.30]
1.39.2 Everolimus	2	1932	Risk Ratio (M-H, Random, 95% CI)	0.60 [0.34, 1.08]
1.39.3 Paclitaxel	7	4305	Risk Ratio (M-H, Random, 95% CI)	1.01 [0.73, 1.39]
1.39.4 Sirolimus	8	2957	Risk Ratio (M-H, Random, 95% CI)	1.04 [0.76, 1.41]
1.39.5 Zotarolimus	1	44	Risk Ratio (M-H, Random, 95% CI)	Not estimable
1.39.6 Mixed	1	626	Risk Ratio (M-H, Random, 95% CI)	1.00 [0.52, 1.92]
1.39.7 Unclear	1	382	Risk Ratio (M-H, Random, 95% CI)	2.16 [0.82, 5.70]
1.40 Stent thrombosis according to type of ACS	21	11350	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.80, 1.16]
1.40.1 STEMI	20	10876	Risk Ratio (M-H, Random, 95% CI)	0.98 [0.81, 1.18]
1.40.2 NSTEMI	1	474	Risk Ratio (M-H, Random, 95% CI)	0.44 [0.12, 1.68]
1.41 Stent thrombosis according to length of maximum follow-up	21	11350	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.80, 1.16]
1.41.1 Less or equal to 6 months	2	482	Risk Ratio (M-H, Random, 95% CI)	2.16 [0.82, 5.70]
1.41.2 Between 6 and 12 months	7	1748	Risk Ratio (M-H, Random, 95% CI)	0.83 [0.46, 1.49]
1.41.3 Between 1 and 3 years	3	744	Risk Ratio (M-H, Random, 95% CI)	0.57 [0.25, 1.29]
1.41.4 More or equal to 3 years	8	8360	Risk Ratio (M-H, Random, 95% CI)	0.98 [0.78, 1.22]
1.41.5 Unclear	1	16	Risk Ratio (M-H, Random, 95% CI)	0.20 [0.01, 3.61]
1.42 Stent thrombosis according to registration status	21	11350	Risk Ratio (M-H, Random, 95% CI)	0.96 [0.80, 1.16]
1.42.1 Pre-registration	3	1248	Risk Ratio (M-H, Random, 95% CI)	0.72 [0.40, 1.30]
1.42.2 Post-registration	13	9346	Risk Ratio (M-H, Random, 95% CI)	0.99 [0.81, 1.21]
1.42.3 No registration	5	756	Risk Ratio (M-H, Random, 95% CI)	0.70 [0.22, 2.20]
1.43 Target vessel revascularisation	23	11770	Risk Ratio (M-H, Fixed, 95% CI)	0.58 [0.52, 0.65]
1.44 Target vessel revascularisation best-worst	23	12368	Risk Ratio (M-H, Fixed, 95% CI)	0.41 [0.37, 0.45]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.45 Target vessel revascularisation worst-best	23	12368	Risk Ratio (M-H, Fixed, 95% CI)	0.98 [0.89, 1.07]
1.46 Target vessel revascularisation according to type of drug-eluting stent	23	11770	Risk Ratio (M-H, Fixed, 95% CI)	0.58 [0.52, 0.65]
1.46.1 Biodegradable (Biolimus)	1	1118	Risk Ratio (M-H, Fixed, 95% CI)	0.45 [0.29, 0.70]
1.46.2 Everolimus	2	1932	Risk Ratio (M-H, Fixed, 95% CI)	0.57 [0.42, 0.78]
1.46.3 Sirolimus	9	3062	Risk Ratio (M-H, Fixed, 95% CI)	0.54 [0.44, 0.66]
1.46.4 Paclitaxel	8	4530	Risk Ratio (M-H, Fixed, 95% CI)	0.63 [0.53, 0.75]
1.46.5 Mixed	1	626	Risk Ratio (M-H, Fixed, 95% CI)	0.52 [0.35, 0.76]
1.46.6 Unclear	1	382	Risk Ratio (M-H, Fixed, 95% CI)	2.16 [0.82, 5.70]
1.46.7 Dexamethasone	1	120	Risk Ratio (M-H, Fixed, 95% CI)	0.60 [0.32, 1.12]
1.47 Target vessel revascularisation according to type of ACS	23	11770	Risk Ratio (M-H, Fixed, 95% CI)	0.58 [0.52, 0.65]
1.47.1 STEMI	20	11097	Risk Ratio (M-H, Fixed, 95% CI)	0.59 [0.53, 0.66]
1.47.2 Acute coronary syndrome	2	199	Risk Ratio (M-H, Fixed, 95% CI)	0.49 [0.27, 0.86]
1.47.3 NSTEMI	1	474	Risk Ratio (M-H, Fixed, 95% CI)	0.37 [0.18, 0.77]
1.48 Target vessel revascularisation according to length of maximum follow-up	23	11770	Risk Ratio (M-H, Fixed, 95% CI)	0.58 [0.52, 0.65]
1.48.1 Less or equal to 6 months	4	724	Risk Ratio (M-H, Fixed, 95% CI)	0.46 [0.25, 0.84]
1.48.2 Between 6 and 12 months	8	1930	Risk Ratio (M-H, Fixed, 95% CI)	0.58 [0.43, 0.78]
1.48.3 Between 1 and 3 years	3	744	Risk Ratio (M-H, Fixed, 95% CI)	0.33 [0.19, 0.56]
1.48.4 More or equal to 3 years	8	8372	Risk Ratio (M-H, Fixed, 95% CI)	0.61 [0.54, 0.69]
1.48.5 Unclear	0	0	Risk Ratio (M-H, Fixed, 95% CI)	Not estimable
1.49 Target vessel revascularisation according to registration status	23	11770	Risk Ratio (M-H, Fixed, 95% CI)	0.58 [0.52, 0.65]
1.49.1 Pre-registration	3	1262	Risk Ratio (M-H, Fixed, 95% CI)	0.42 [0.28, 0.65]
1.49.2 Post-registration	14	9425	Risk Ratio (M-H, Fixed, 95% CI)	0.60 [0.53, 0.67]
1.49.3 No registration	6	1083	Risk Ratio (M-H, Fixed, 95% CI)	0.56 [0.38, 0.82]

Analysis 1.1. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 1: All-cause mortality

Analysis 1.2. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 2: All-cause mortality best-worst

Analysis 1.3. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 3: All-cause mortality worst-best

Analysis 1.4. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 4: All-cause mortality according to type of drug-eluting stent

Study or Subgroup	Drug-eluting stents		Bare-metal stents		Weight	Risk Ratio	Risk Ratio
	Events	Total	Events	Total		M-H, Random, 95% CI	M-H, Random, 95% CI
1.4.1 Biodegradable (Biolimus)							
COMFORTABLE 2012	28	575	32	582	7.5%	0.89 [0.54 , 1.45]	
Subtotal (95% CI)		575		582	7.5%	0.89 [0.54 , 1.45]	
Total events:	28		32				
Heterogeneity: Not applicable							
Test for overall effect: Z = 0.48 (P = 0.63)							
1.4.2 Everolimus							
EXAMINATION 2012	65	731	88	727	19.8%	0.73 [0.54 , 1.00]	
Subtotal (95% CI)		731		727	19.8%	0.73 [0.54 , 1.00]	
Total events:	65		88				
Heterogeneity: Not applicable							
Test for overall effect: Z = 1.99 (P = 0.05)							
1.4.3 Paclitaxel							
DEB-AMI 2012	0	49	2	51	0.2%	0.21 [0.01 , 4.23]	
DEVINE 2007	5	81	4	82	1.1%	1.27 [0.35 , 4.54]	
GRACIA-3 2010	21	217	15	216	4.5%	1.39 [0.74 , 2.63]	
HAAMU-STENT 2006	8	82	4	82	1.4%	2.00 [0.63 , 6.38]	
HORIZONS-AMI 2009	123	2103	48	687	17.6%	0.84 [0.61 , 1.16]	
PASEO 2009a	6	90	5	45	1.4%	0.60 [0.19 , 1.86]	
PASSION 2006	27	302	35	301	8.1%	0.77 [0.48 , 1.24]	
SELECTION 2007	1	40	3	40	0.4%	0.33 [0.04 , 3.07]	
Subtotal (95% CI)		2964		1504	34.7%	0.89 [0.71 , 1.12]	
Total events:	191		116				
Heterogeneity: Tau ² = 0.00; Chi ² = 6.69, df = 7 (P = 0.46); I ² = 0%							
Test for overall effect: Z = 0.98 (P = 0.33)							
1.4.4 Sirolimus							
DEBATER 2012	39	408	39	422	10.2%	1.03 [0.68 , 1.58]	
Diaz 2007	2	60	3	54	0.6%	0.60 [0.10 , 3.46]	
Mission 2008	2	158	4	152	0.6%	0.48 [0.09 , 2.59]	
MULTISTRATEGY 2008	26	369	28	367	6.9%	0.92 [0.55 , 1.54]	
PASEO 2009b	5	90	4	45	1.1%	0.63 [0.18 , 2.21]	
SESAMI 2007	5	157	8	156	1.5%	0.62 [0.21 , 1.86]	
Steinwender 2008	0	8	0	8		Not estimable	
Strozzi 2007	0	39	0	40		Not estimable	
Typhoon 2006	17	294	20	286	4.7%	0.83 [0.44 , 1.55]	
Subtotal (95% CI)		1583		1530	25.7%	0.89 [0.68 , 1.16]	
Total events:	96		106				
Heterogeneity: Tau ² = 0.00; Chi ² = 1.98, df = 6 (P = 0.92); I ² = 0%							
Test for overall effect: Z = 0.90 (P = 0.37)							
1.4.5 Zotarolimus							
OCTAMI 2010	0	33	0	11		Not estimable	
Subtotal (95% CI)		33		11		Not estimable	
Total events:	0		0				
Heterogeneity: Not applicable							
Test for overall effect: Not applicable							
1.4.6 Mixed							
DEDICATION 2008	51	313	38	313	12.0%	1.34 [0.91 , 1.98]	
Subtotal (95% CI)		313		313	12.0%	1.34 [0.91 , 1.98]	
Total events:	51		38				
Heterogeneity: Not applicable							
Test for overall effect: Z = 1.48 (P = 0.14)							

Analysis 1.4. (Continued)

Heterogeneity: Not applicable

Test for overall effect: $Z = 1.48$ ($P = 0.14$)

1.4.7 Unclear

Darkahian 2014	0	51	7	333	0.2%	0.43 [0.02 , 7.39]
----------------	---	----	---	-----	------	--------------------

Subtotal (95% CI)		51		333	0.2%	0.43 [0.02 , 7.39]
--------------------------	--	-----------	--	------------	-------------	---------------------------

Total events:	0		7			
---------------	---	--	---	--	--	--

Heterogeneity: Not applicable

Test for overall effect: $Z = 0.58$ ($P = 0.56$)

Total (95% CI)		6250		5000	100.0%	0.90 [0.78 , 1.03]
-----------------------	--	-------------	--	-------------	---------------	---------------------------

Total events:	431		387			
---------------	-----	--	-----	--	--	--

Heterogeneity: $\text{Tau}^2 = 0.00$; $\text{Chi}^2 = 14.72$, $\text{df} = 18$ ($P = 0.68$); $I^2 = 0\%$

Test for overall effect: $Z = 1.57$ ($P = 0.12$)

Test for subgroup differences: $\text{Chi}^2 = 6.04$, $\text{df} = 5$ ($P = 0.30$), $I^2 = 17.3\%$

Analysis 1.5. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 5: All-cause mortality according to type of ACS

Analysis 1.6. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 6: All-cause mortality according to length of maximum follow-up

Analysis 1.6. (Continued)

Heterogeneity: $\tau^2 = 0.00$; $\chi^2 = 14.72$, $df = 18$ ($P = 0.68$); $I^2 = 0\%$

Test for overall effect: $Z = 1.57$ ($P = 0.12$)

Test for subgroup differences: $\chi^2 = 0.37$, $df = 3$ ($P = 0.95$), $I^2 = 0\%$

0.01 0.1 1 10 100
Favours DES Favours BMS

Analysis 1.7. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 7: All-cause mortality according to registration status

Analysis 1.8. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 8: Serious adverse events

Analysis 1.9. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 9: Serious adverse events best-worst

Analysis 1.10. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 10: Serious adverse events worst-best

Analysis 1.11. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 11: Serious adverse events according to type of drug-eluting stent

Analysis 1.11. (Continued)

Subtotal (95% CI) 313 313 5.7% **0.74 [0.54 , 1.01]**

Total events: 53 72

Heterogeneity: Not applicable

Test for overall effect: $Z = 1.89$ ($P = 0.06$)

1.11.7 Unclear

Darkahian 2014 10 51 70 333 1.5% 0.93 [0.52 , 1.69]

Subtotal (95% CI) 51 333 **1.5%** **0.93 [0.52 , 1.69]**

Total events: 10 70

Heterogeneity: Not applicable

Test for overall effect: $Z = 0.23$ ($P = 0.82$)

Total (95% CI) 6484 5240 **100.0%** **0.80 [0.74 , 0.86]**

Total events: 1292 1206

Heterogeneity: $\text{Chi}^2 = 26.95$, $\text{df} = 21$ ($P = 0.17$); $I^2 = 22\%$

Test for overall effect: $Z = 6.09$ ($P < 0.00001$)

Test for subgroup differences: $\text{Chi}^2 = 3.85$, $\text{df} = 6$ ($P = 0.70$), $I^2 = 0\%$

Analysis 1.12. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 12: Serious adverse events according to type of ACS

Analysis 1.13. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 13: Serious adverse events according to length of maximum follow-up

Analysis 1.13. (Continued)

Total (95% CI) 6484 5240 100.0% 0.80 [0.74, 0.86]

Total events: 1292 1206

Heterogeneity: $\text{Chi}^2 = 26.95$, $\text{df} = 21$ ($P = 0.17$); $I^2 = 22\%$

Test for overall effect: $Z = 6.09$ ($P < 0.00001$)

Test for subgroup differences: $\text{Chi}^2 = 3.20$, $\text{df} = 3$ ($P = 0.36$), $I^2 = 6.3\%$

Analysis 1.14. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 14: Serious adverse events according to registration status

Analysis 1.15. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 15: Major cardiovascular events

Analysis 1.16. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 16: Major cardiovascular events best-worst

Analysis 1.17. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 17: Major cardiovascular events worst-best

Analysis 1.18. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 18: Major cardiovascular events according to type of drug-eluting stent

Analysis 1.18. (Continued)

1.18.7 Unclear

Darkahian 2014	5	51	24	333	2.6%	1.36 [0.54 , 3.40]
Subtotal (95% CI)		51		333	2.6%	1.36 [0.54 , 3.40]

Total events:

Heterogeneity: Not applicable

Test for overall effect: $Z = 0.66$ ($P = 0.51$)

Total (95% CI)		6096		4843	100.0%	0.96 [0.83 , 1.11]
-----------------------	--	-------------	--	-------------	---------------	---------------------------

Total events:

Heterogeneity: $\text{Tau}^2 = 0.00$; $\text{Chi}^2 = 11.50$, $df = 18$ ($P = 0.87$); $I^2 = 0\%$

Test for overall effect: $Z = 0.55$ ($P = 0.58$)

Test for subgroup differences: $\text{Chi}^2 = 1.54$, $df = 5$ ($P = 0.91$), $I^2 = 0\%$

Analysis 1.19. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 19: Major cardiovascular events according to type of ACS

Study or Subgroup	Drug-eluting stents		Bare-metal stents		Weight	Risk Ratio M-H, Random, 95% CI	Risk Ratio M-H, Random, 95% CI
	Events	Total	Events	Total			
1.19.1 STEMI							
COMFORTABLE 2012	27	553	30	551	8.5%	0.90 [0.54 , 1.49]	
Darkahian 2014	5	51	24	333	2.6%	1.36 [0.54 , 3.40]	
DEB-AMI 2012	0	49	2	51	0.2%	0.21 [0.01 , 4.23]	
DEBATER 2012	6	408	2	422	0.9%	3.10 [0.63 , 15.28]	
DEDICATION 2008	24	313	23	313	7.2%	1.04 [0.60 , 1.81]	
Diaz 2007	3	60	3	54	0.9%	0.90 [0.19 , 4.27]	
EXAMINATION 2012	47	731	55	727	15.5%	0.85 [0.58 , 1.24]	
GRACIA-3 2010	21	217	15	216	5.4%	1.39 [0.74 , 2.63]	
HAAMU-STENT 2006	1	82	4	82	0.5%	0.25 [0.03 , 2.19]	
HORIZONS-AMI 2009	150	2103	47	687	21.8%	1.04 [0.76 , 1.43]	
Mission 2008	9	158	14	152	3.3%	0.62 [0.28 , 1.39]	
MULTISTRATEGY 2008	21	369	21	367	6.3%	0.99 [0.55 , 1.79]	
OCTAMI 2010	0	33	0	11		Not estimable	
PASEO 2009a	11	90	8	45	3.1%	0.69 [0.30 , 1.59]	
PASEO 2009b	10	90	9	45	3.2%	0.56 [0.24 , 1.27]	
PASSION 2006	45	302	44	301	14.8%	1.02 [0.69 , 1.50]	
SELECTION 2007	1	40	2	40	0.4%	0.50 [0.05 , 5.30]	
SESAMI 2007	4	157	4	156	1.2%	0.99 [0.25 , 3.90]	
Typhoon 2006	13	251	11	250	3.5%	1.18 [0.54 , 2.58]	
Subtotal (95% CI)		6057		4803	99.3%	0.96 [0.83 , 1.12]	
Total events:	398		318				
Heterogeneity: Tau² = 0.00; Chi² = 11.35, df = 17 (P = 0.84); I² = 0%							
Test for overall effect: Z = 0.52 (P = 0.61)							
1.19.2 Acute coronary syndrome							
Strozzi 2007	2	39	3	40	0.7%	0.68 [0.12 , 3.87]	
Subtotal (95% CI)		39		40	0.7%	0.68 [0.12 , 3.87]	
Total events:	2		3				
Heterogeneity: Not applicable							
Test for overall effect: Z = 0.43 (P = 0.67)							
Total (95% CI)		6096		4843	100.0%	0.96 [0.83 , 1.11]	
Total events:	400		321				
Heterogeneity: Tau² = 0.00; Chi² = 11.50, df = 18 (P = 0.87); I² = 0%							
Test for overall effect: Z = 0.55 (P = 0.58)							
Test for subgroup differences: Chi² = 0.15, df = 1 (P = 0.70), I² = 0%

Analysis 1.20. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 20: Major cardiovascular events according to length of maximum follow-up

Analysis 1.21. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 21: Major cardiovascular events according to registration status

Analysis 1.22. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 22: Cardiovascular mortality

Analysis 1.23. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 23: Cardiovascular mortality best-worst

Analysis 1.24. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 24: Cardiovascular mortality worst-best

Analysis 1.25. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 25: Cardiovascular mortality according to type of drug-eluting stent

Analysis 1.25. (Continued)

Test for overall effect: $Z = 1.06$ ($P = 0.29$)

Favours DES

Favours BMS

Test for subgroup differences: $\text{Chi}^2 = 7.55$, $\text{df} = 4$ ($P = 0.11$), $I^2 = 47.0\%$

Analysis 1.26. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 26: Cardiovascular mortality according to type of ACS

Analysis 1.27. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 27: Cardiovascular mortality according to length of maximum follow-up

Analysis 1.28. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 28: Cardiovascular mortality according to registration status

Analysis 1.29. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 29: Myocardial infarction

Analysis 1.30. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 30: Myocardial infarction best-worst

Analysis 1.31. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 31: Myocardial infarction worst-best

Analysis 1.32. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 32: Myocardial infarction according to type of drug-eluting stent

Analysis 1.32. (Continued)

1.32.7 Unclear

Darkahian 2014	5	51	24	333	4.0%	1.36 [0.54 , 3.40]
Subtotal (95% CI)		51		333	4.0%	1.36 [0.54 , 3.40]

Total events: 5 24

Heterogeneity: Not applicable

Test for overall effect: $Z = 0.66$ ($P = 0.51$)

Total (95% CI) 5732 4485 100.0% **0.98 [0.82 , 1.18]**

Total events: 296 214

Heterogeneity: $\tau^2 = 0.00$; $\chi^2 = 14.55$, $df = 16$ ($P = 0.56$); $I^2 = 0\%$

Test for overall effect: $Z = 0.21$ ($P = 0.83$)

Test for subgroup differences: $\chi^2 = 4.84$, $df = 5$ ($P = 0.44$), $I^2 = 0\%$

Analysis 1.33. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 33: Myocardial infarction according to type of ACS

Study or Subgroup	Drug-eluting stents Events	Drug-eluting stents Total	Bare-metal stents Events	Bare-metal stents Total	Weight	Risk Ratio M-H, Random, 95% CI	Risk Ratio M-H, Random, 95% CI
1.33.1 STEMI							
COMFORTABLE 2012	18	559	28	559	9.9%	0.64 [0.36 , 1.15]	
Darkahian 2014	5	51	24	333	4.0%	1.36 [0.54 , 3.40]	
DEB-AMI 2012	0	49	0	51		Not estimable	
DEBATER 2012	6	408	2	422	1.3%	3.10 [0.63 , 15.28]	
DEDICATION 2008	17	313	23	313	9.1%	0.74 [0.40 , 1.36]	
Diaz 2007	1	60	1	54	0.4%	0.90 [0.06 , 14.04]	
EXAMINATION 2012	35	731	27	727	13.9%	1.29 [0.79 , 2.11]	
GRACIA-3 2010	5	216	3	217	1.7%	1.67 [0.41 , 6.92]	
HAAMU-STENT 2006	1	82	4	82	0.7%	0.25 [0.03 , 2.19]	
HORIZONS-AMI 2009	150	2103	47	687	33.5%	1.04 [0.76 , 1.43]	
Mission 2008	9	158	14	152	5.1%	0.62 [0.28 , 1.39]	
OCTAMI 2010	0	33	0	11		Not estimable	
PASEO 2009a	5	90	5	45	2.4%	0.50 [0.15 , 1.64]	
PASEO 2009b	6	90	5	45	2.6%	0.60 [0.19 , 1.86]	
PASSION 2006	19	302	12	301	6.7%	1.58 [0.78 , 3.19]	
SELECTION 2007	0	40	1	40	0.3%	0.33 [0.01 , 7.95]	
SESAMI 2007	4	157	4	156	1.8%	0.99 [0.25 , 3.90]	
Typhoon 2006	13	251	11	250	5.4%	1.18 [0.54 , 2.58]	
Subtotal (95% CI)		5693		4445	98.9%	0.98 [0.82 , 1.18]	
Total events:	294		211				
Heterogeneity: $\tau^2 = 0.00$; $\chi^2 = 14.39$, $df = 15$ ($P = 0.50$); $I^2 = 0\%$							
Test for overall effect: $Z = 0.17$ ($P = 0.87$)							
1.33.2 Acute coronary syndrome							
Strozzi 2007	2	39	3	40	1.1%	0.68 [0.12 , 3.87]	
Subtotal (95% CI)		39		40	1.1%	0.68 [0.12 , 3.87]	
Total events:	2		3				
Heterogeneity: Not applicable							
Test for overall effect: $Z = 0.43$ ($P = 0.67$)							
Total (95% CI)		5732		4485	100.0%	0.98 [0.82 , 1.18]	
Total events:	296		214				
Heterogeneity: $\tau^2 = 0.00$; $\chi^2 = 14.55$, $df = 16$ ($P = 0.56$); $I^2 = 0\%$							
Test for overall effect: $Z = 0.21$ ($P = 0.83$)							
Test for subgroup differences: $\chi^2 = 0.17$, $df = 1$ ($P = 0.68$), $I^2 = 0\%$

Analysis 1.34. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 34: Myocardial infarction according to length of maximum follow-up

Analysis 1.35. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 35: Myocardial infarction according to registration status

Analysis 1.36. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 36: Stent thrombosis

Analysis 1.37. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 37: Stent thrombosis best-worst

Analysis 1.38. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 38: Stent thrombosis worst-best

Analysis 1.39. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 39: Stent thrombosis according to type of drug-eluting stent

Analysis 1.39. (Continued)

Test for overall effect: $Z = 0.00$ ($P = 1.00$)

1.39.7 Unclear

Darkahian 2014	5	51	15	331	3.6%	2.16 [0.82 , 5.70]
----------------	---	----	----	-----	------	--------------------

Subtotal (95% CI)		51		331	3.6%	2.16 [0.82 , 5.70]
--------------------------	--	-----------	--	------------	-------------	---------------------------

Total events:	5		15			
---------------	---	--	----	--	--	--

Heterogeneity: Not applicable

Test for overall effect: $Z = 1.56$ ($P = 0.12$)

Total (95% CI)		6299		5051	100.0%	0.96 [0.80 , 1.16]
-----------------------	--	-------------	--	-------------	---------------	---------------------------

Total events:	268		223			
---------------	-----	--	-----	--	--	--

Heterogeneity: $\tau^2 = 0.00$; $\chi^2 = 16.88$, $df = 18$ ($P = 0.53$); $I^2 = 0\%$

Test for overall effect: $Z = 0.41$ ($P = 0.68$)

Test for subgroup differences: $\chi^2 = 6.44$, $df = 5$ ($P = 0.27$), $I^2 = 22.3\%$

Analysis 1.40. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 40: Stent thrombosis according to type of ACS

Analysis 1.41. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 41: Stent thrombosis according to length of maximum follow-up

Analysis 1.41. (Continued)

Test for overall effect: $Z = 0.41$ ($P = 0.68$)

Favours DES

Favours BMS

Test for subgroup differences: $\text{Chi}^2 = 5.64$, $df = 4$ ($P = 0.23$), $I^2 = 29.1\%$

Analysis 1.42. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 42: Stent thrombosis according to registration status

Analysis 1.43. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 43: Target vessel revascularisation

Analysis 1.44. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 44: Target vessel revascularisation best-worst

Analysis 1.45. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 45: Target vessel revascularisation worst-best

Analysis 1.46. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 46: Target vessel revascularisation according to type of drug-eluting stent

Analysis 1.46. (Continued)

Subtotal (95% CI) 51 331 0.5% 2.16 [0.82, 5.70]

Total events: 5 15

Heterogeneity: Not applicable

Test for overall effect: $Z = 1.56$ ($P = 0.12$)

1.46.7 Dexamethasone

König 2007 12 60 20 60 2.6% 0.60 [0.32, 1.12]

Subtotal (95% CI) 60 60 2.6% 0.60 [0.32, 1.12]

Total events: 12 20

Heterogeneity: Not applicable

Test for overall effect: $Z = 1.62$ ($P = 0.11$)

Total (95% CI) 6504 5266 100.0% 0.58 [0.52, 0.65]

Total events: 576 702

Heterogeneity: $\text{Chi}^2 = 35.01$, $\text{df} = 22$ ($P = 0.04$); $I^2 = 37\%$

Test for overall effect: $Z = 9.82$ ($P < 0.00001$)

Test for subgroup differences: $\text{Chi}^2 = 10.16$, $\text{df} = 6$ ($P = 0.12$), $I^2 = 41.0\%$

Analysis 1.47. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 47: Target vessel revascularisation according to type of ACS

Analysis 1.48. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 48: Target vessel revascularisation according to length of maximum follow-up

Study or Subgroup	Drug-eluting stents		Bare-metal stents		Weight	Risk Ratio M-H, Fixed, 95% CI	Risk Ratio M-H, Fixed, 95% CI
Events	Total	Events	Total				
1.48.1 Less or equal to 6 months							
Darkahian 2014	5	51	15	331	0.5%	2.16 [0.82 , 5.70]
DEB-AMI 2012	1	49	9	51	1.2%	0.12 [0.02 , 0.88]
DEVINE 2007	2	81	8	82	1.0%	0.25 [0.06 , 1.16]
Strozzi 2007	2	39	9	40	1.2%	0.23 [0.05 , 0.99]
Subtotal (95% CI)		220		504	3.9%	0.46 [0.25 , 0.84]
Total events:	10		41				
Heterogeneity: Chi ² = 13.04, df = 3 (P = 0.005); I ² = 77%							
Test for overall effect: Z = 2.53 (P = 0.01)							
1.48.2 Between 6 and 12 months							
Diaz 2007	1	60	3	54	0.4%	0.30 [0.03 , 2.80]
GRACIA-3 2010	9	217	10	216	1.3%	0.90 [0.37 , 2.16]
HAAMU-STENT 2006	2	69	7	75	0.9%	0.31 [0.07 , 1.44]
König 2007	12	60	20	60	2.6%	0.60 [0.32 , 1.12]
Mission 2008	8	158	20	152	2.7%	0.38 [0.17 , 0.85]
OCTAMI 2010	4	33	1	11	0.2%	1.33 [0.17 , 10.70]
PASSION 2006	22	302	30	303	3.9%	0.74 [0.43 , 1.25]
SELECTION 2007	7	80	17	80	2.2%	0.41 [0.18 , 0.94]
Subtotal (95% CI)		979		951	14.3%	0.58 [0.43 , 0.78]
Total events:	65		108				
Heterogeneity: Chi ² = 5.02, df = 7 (P = 0.66); I ² = 0%							
Test for overall effect: Z = 3.66 (P = 0.0002)							
1.48.3 Between 1 and 3 years							
ELISA-3 2016	9	234	25	240	3.3%	0.37 [0.18 , 0.77]
PASEO 2009a	5	90	8	45	1.4%	0.31 [0.11 , 0.90]
PASEO 2009b	4	90	8	45	1.4%	0.25 [0.08 , 0.79]
Subtotal (95% CI)		414		330	6.1%	0.33 [0.19 , 0.56]
Total events:	18		41				
Heterogeneity: Chi ² = 0.32, df = 2 (P = 0.85); I ² = 0%							
Test for overall effect: Z = 4.06 (P < 0.0001)							
1.48.4 More or equal to 3 years							
COMFORTABLE 2012	26	559	58	559	7.6%	0.45 [0.29 , 0.70]
DEBATER 2012	54	408	66	422	8.5%	0.85 [0.61 , 1.18]
DEDICATION 2008	33	313	64	313	8.4%	0.52 [0.35 , 0.76]
EXAMINATION 2012	49	731	76	727	10.0%	0.64 [0.45 , 0.90]
HORIZONS-AMI 2009	265	2103	125	687	24.8%	0.69 [0.57 , 0.84]
MULTISTRATEGY 2008	23	369	50	367	6.6%	0.46 [0.29 , 0.73]
SESAMI 2007	13	157	25	156	3.3%	0.52 [0.27 , 0.97]
Typhoon 2006	20	251	48	250	6.3%	0.42 [0.25 , 0.68]
Subtotal (95% CI)		4891		3481	75.7%	0.61 [0.54 , 0.69]
Total events:	483		512				
Heterogeneity: Chi ² = 12.05, df = 7 (P = 0.10); I ² = 42%							
Test for overall effect: Z = 7.98 (P < 0.00001)							
1.48.5 Unclear							
Subtotal (95% CI)		0		0		Not estimable	
Total events:	0		0				
Heterogeneity: Not applicable							
Test for overall effect: Not applicable							
Total (95% CI)		6504		5266	100.0%	0.58 [0.52 , 0.65]
Total events:	576		702				

Analysis 1.48. (Continued)

Total (95% CI) 6504 5266 100.0% 0.58 [0.52, 0.65]

Total events: 576 702

Heterogeneity: $\chi^2 = 35.01$, $df = 22$ ($P = 0.04$); $I^2 = 37\%$

Test for overall effect: $Z = 9.82$ ($P < 0.00001$)

Test for subgroup differences: $\chi^2 = 5.40$, $df = 3$ ($P = 0.14$), $I^2 = 44.4\%$

Analysis 1.49. Comparison 1: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at maximum follow-up, Outcome 49: Target vessel revascularisation according to registration status

Comparison 2. Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.1 All-cause mortality	6	3213	Risk Ratio (M-H, Fixed, 95% CI)	0.78 [0.49, 1.22]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.2 All-cause mortality best-worst	6	3231	Risk Ratio (M-H, Random, 95% CI)	0.61 [0.40, 0.95]
2.3 All-cause mortality worst-best	6	3231	Risk Ratio (M-H, Random, 95% CI)	0.92 [0.60, 1.43]
2.4 All-cause mortality according to type of drug-eluting stent	6	3213	Risk Ratio (M-H, Fixed, 95% CI)	0.78 [0.49, 1.22]
2.4.1 Everolimus	1	1498	Risk Ratio (M-H, Fixed, 95% CI)	0.78 [0.36, 1.71]
2.4.2 Paclitaxel	3	857	Risk Ratio (M-H, Fixed, 95% CI)	0.83 [0.42, 1.64]
2.4.3 Sirolimus	2	858	Risk Ratio (M-H, Fixed, 95% CI)	0.68 [0.26, 1.78]
2.5 All-cause mortality according to type of ACS	6	3213	Risk Ratio (M-H, Fixed, 95% CI)	0.78 [0.49, 1.22]
2.5.1 STEMI	6	3213	Risk Ratio (M-H, Fixed, 95% CI)	0.78 [0.49, 1.22]
2.6 All-cause mortality according to registration status	6	3213	Risk Ratio (M-H, Fixed, 95% CI)	0.78 [0.49, 1.22]
2.6.1 Post-registration	3	2856	Risk Ratio (M-H, Fixed, 95% CI)	0.68 [0.41, 1.14]
2.6.2 No registration	3	357	Risk Ratio (M-H, Fixed, 95% CI)	1.28 [0.47, 3.47]
2.7 Serious adverse events	7	3313	Risk Ratio (M-H, Random, 95% CI)	0.65 [0.45, 0.93]
2.8 Serious adverse events best-worst	7	3331	Risk Ratio (M-H, Random, 95% CI)	0.56 [0.40, 0.80]
2.9 Serious adverse events worst-best	7	3331	Risk Ratio (M-H, Random, 95% CI)	0.73 [0.52, 1.03]
2.10 Serious adverse events according to type of drug-eluting stent	7	3313	Risk Ratio (M-H, Random, 95% CI)	0.65 [0.45, 0.93]
2.10.1 Everolimus	1	1498	Risk Ratio (M-H, Random, 95% CI)	0.55 [0.30, 0.98]
2.10.2 Paclitaxel	4	957	Risk Ratio (M-H, Random, 95% CI)	0.77 [0.41, 1.46]
2.10.3 Sirolimus	2	858	Risk Ratio (M-H, Random, 95% CI)	0.67 [0.35, 1.28]
2.11 Serious adverse events according to type of ACS	7	3313	Risk Ratio (M-H, Random, 95% CI)	0.65 [0.45, 0.93]
2.11.1 STEMI	7	3313	Risk Ratio (M-H, Random, 95% CI)	0.65 [0.45, 0.93]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.12 Serious adverse events according to registration status	7	3313	Risk Ratio (M-H, Random, 95% CI)	0.65 [0.45, 0.93]
2.12.1 Pre-registration	1	100	Risk Ratio (M-H, Random, 95% CI)	0.35 [0.01, 8.31]
2.12.2 Post-registration	3	2856	Risk Ratio (M-H, Random, 95% CI)	0.60 [0.41, 0.88]
2.12.3 No registration	3	357	Risk Ratio (M-H, Random, 95% CI)	1.09 [0.43, 2.76]
2.13 Major cardiovascular events	6	3150	Risk Ratio (M-H, Random, 95% CI)	0.68 [0.43, 1.08]
2.14 Major cardiovascular events best-worst	6	3168	Risk Ratio (M-H, Random, 95% CI)	0.54 [0.35, 0.84]
2.15 Major cardiovascular events worst-best	6	3168	Risk Ratio (M-H, Random, 95% CI)	0.82 [0.53, 1.27]
2.16 Major cardiovascular events according to type of drug-eluting stent	6	3150	Risk Ratio (M-H, Random, 95% CI)	0.68 [0.43, 1.08]
2.16.1 Everolimus	1	1498	Risk Ratio (M-H, Random, 95% CI)	0.78 [0.36, 1.71]
2.16.2 Paclitaxel	3	794	Risk Ratio (M-H, Random, 95% CI)	0.66 [0.32, 1.37]
2.16.3 Sirolimus	2	858	Risk Ratio (M-H, Random, 95% CI)	0.60 [0.25, 1.45]
2.17 Major cardiovascular events according to type of ACS	6	3150	Risk Ratio (M-H, Random, 95% CI)	0.68 [0.43, 1.08]
2.17.1 STEMI	6	3150	Risk Ratio (M-H, Random, 95% CI)	0.68 [0.43, 1.08]
2.18 Major cardiovascular events according to registration status	6	3150	Risk Ratio (M-H, Random, 95% CI)	0.68 [0.43, 1.08]
2.18.1 Pre-registration	1	100	Risk Ratio (M-H, Random, 95% CI)	0.35 [0.01, 8.31]
2.18.2 Post-registration	3	2856	Risk Ratio (M-H, Random, 95% CI)	0.67 [0.41, 1.09]
2.18.3 No registration	2	194	Risk Ratio (M-H, Random, 95% CI)	0.92 [0.24, 3.58]
2.19 Cardiovascular mortality	5	2406	Risk Ratio (M-H, Fixed, 95% CI)	0.73 [0.43, 1.25]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.20 Cardiovascular mortality best-worst	5	2423	Risk Ratio (M-H, Fixed, 95% CI)	0.53 [0.32, 0.88]
2.21 Cardiovascular mortality worst-best	5	2423	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.54, 1.50]
2.22 Cardiovascular mortality according to type of drug-eluting stent	5	2406	Risk Ratio (M-H, Fixed, 95% CI)	0.73 [0.43, 1.25]
2.22.1 Everolimus	1	1498	Risk Ratio (M-H, Fixed, 95% CI)	0.78 [0.36, 1.71]
2.22.2 Paclitaxel	3	794	Risk Ratio (M-H, Fixed, 95% CI)	0.67 [0.30, 1.46]
2.22.3 Sirolimus	1	114	Risk Ratio (M-H, Fixed, 95% CI)	0.90 [0.13, 6.17]
2.23 Cardiovascular mortality according to type of ACS	5	2406	Risk Ratio (M-H, Fixed, 95% CI)	0.73 [0.43, 1.25]
2.23.1 STEMI	5	2406	Risk Ratio (M-H, Fixed, 95% CI)	0.73 [0.43, 1.25]
2.24 Cardiovascular mortality according to registration status	5	2406	Risk Ratio (M-H, Fixed, 95% CI)	0.73 [0.43, 1.25]
2.24.1 Pre-registration	1	100	Risk Ratio (M-H, Fixed, 95% CI)	0.35 [0.01, 8.31]
2.24.2 Post-registration	2	2112	Risk Ratio (M-H, Fixed, 95% CI)	0.70 [0.39, 1.25]
2.24.3 No registration	2	194	Risk Ratio (M-H, Fixed, 95% CI)	1.30 [0.27, 6.38]
2.25 Myocardial infarction	6	3150	Risk Ratio (M-H, Random, 95% CI)	0.51 [0.26, 0.98]
2.26 Myocardial infarction best-worst	6	3168	Risk Ratio (M-H, Random, 95% CI)	0.37 [0.20, 0.68]
2.27 Myocardial infarction worst-best	6	3168	Risk Ratio (M-H, Random, 95% CI)	0.74 [0.41, 1.32]
2.28 Myocardial infarction according to type of drug-eluting stent	6	3150	Risk Ratio (M-H, Random, 95% CI)	0.51 [0.26, 0.98]
2.28.1 Everolimus	1	1498	Risk Ratio (M-H, Random, 95% CI)	0.55 [0.19, 1.64]
2.28.2 Paclitaxel	3	794	Risk Ratio (M-H, Random, 95% CI)	0.38 [0.09, 1.63]
2.28.3 Sirolimus	2	858	Risk Ratio (M-H, Random, 95% CI)	0.54 [0.20, 1.46]
2.29 Myocardial infarction according to type of ACS	6	3150	Risk Ratio (M-H, Random, 95% CI)	0.51 [0.26, 0.98]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.29.1 STEMI	6	3150	Risk Ratio (M-H, Random, 95% CI)	0.51 [0.26, 0.98]
2.30 Myocardial infarction according to registration status	6	3150	Risk Ratio (M-H, Random, 95% CI)	0.51 [0.26, 0.98]
2.30.1 Pre-registration	1	100	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.30.2 Post-registration	3	2856	Risk Ratio (M-H, Random, 95% CI)	0.50 [0.25, 1.00]
2.30.3 No registration	2	194	Risk Ratio (M-H, Random, 95% CI)	0.59 [0.07, 4.69]
2.31 Stent thrombosis	5	3070	Risk Ratio (M-H, Random, 95% CI)	0.48 [0.26, 0.90]
2.32 Stent thrombosis best-worst	5	3088	Risk Ratio (M-H, Random, 95% CI)	0.36 [0.20, 0.65]
2.33 Stent thrombosis worst-best	5	3088	Risk Ratio (M-H, Random, 95% CI)	0.68 [0.39, 1.18]
2.34 Stent thrombosis according to type of drug-eluting stent	5	3070	Risk Ratio (M-H, Random, 95% CI)	0.48 [0.26, 0.90]
2.34.1 Everolimus	1	1498	Risk Ratio (M-H, Random, 95% CI)	0.37 [0.15, 0.95]
2.34.2 Paclitaxel	2	714	Risk Ratio (M-H, Random, 95% CI)	0.66 [0.11, 3.94]
2.34.3 Sirolimus	2	858	Risk Ratio (M-H, Random, 95% CI)	0.58 [0.23, 1.46]
2.35 Stent thrombosis according to type of ACS	5	3070	Risk Ratio (M-H, Random, 95% CI)	0.48 [0.26, 0.90]
2.35.1 STEMI	5	3070	Risk Ratio (M-H, Random, 95% CI)	0.48 [0.26, 0.90]
2.36 Stent thrombosis according to registration status	5	3070	Risk Ratio (M-H, Random, 95% CI)	0.48 [0.26, 0.90]
2.36.1 Pre-registration	1	100	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.36.2 Post-registration	3	2856	Risk Ratio (M-H, Random, 95% CI)	0.47 [0.25, 0.88]
2.36.3 No registration	1	114	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.06, 14.04]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.37 Target vessel revascularisation	6	3233	Risk Ratio (M-H, Random, 95% CI)	0.50 [0.31, 0.82]
2.38 Target vessel revascularisation best-worst	6	3251	Risk Ratio (M-H, Random, 95% CI)	0.41 [0.26, 0.66]
2.39 Target vessel revascularisation worst-best	6	3251	Risk Ratio (M-H, Random, 95% CI)	0.62 [0.40, 0.98]
2.40 Target vessel revascularisation according to type of drug-eluting stent	6	3233	Risk Ratio (M-H, Random, 95% CI)	0.50 [0.31, 0.82]
2.40.1 Everolimus	1	1498	Risk Ratio (M-H, Random, 95% CI)	0.36 [0.17, 0.76]
2.40.2 Paclitaxel	3	877	Risk Ratio (M-H, Random, 95% CI)	0.52 [0.18, 1.49]
2.40.3 Sirolimus	2	858	Risk Ratio (M-H, Random, 95% CI)	0.77 [0.29, 2.04]
2.41 Target vessel revascularisation according to type of ACS	6	3233	Risk Ratio (M-H, Random, 95% CI)	0.50 [0.31, 0.82]
2.41.1 STEMI	6	3233	Risk Ratio (M-H, Random, 95% CI)	0.50 [0.31, 0.82]
2.42 Target vessel revascularisation according to registration status	6	3233	Risk Ratio (M-H, Random, 95% CI)	0.50 [0.31, 0.82]
2.42.1 Pre-registration	1	100	Risk Ratio (M-H, Random, 95% CI)	Not estimable
2.42.2 Post-registration	3	2856	Risk Ratio (M-H, Random, 95% CI)	0.53 [0.32, 0.90]
2.42.3 No registration	2	277	Risk Ratio (M-H, Random, 95% CI)	0.34 [0.09, 1.29]

Analysis 2.1. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 1: All-cause mortality

Analysis 2.2. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 2: All-cause mortality best-worst

Analysis 2.3. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 3: All-cause mortality worst-best

Analysis 2.4. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 4: All-cause mortality according to type of drug-eluting stent

Analysis 2.5. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 5: All-cause mortality according to type of ACS

Analysis 2.6. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 6: All-cause mortality according to registration status

Analysis 2.7. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 7: Serious adverse events

Analysis 2.8. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 8: Serious adverse events best-worst

Analysis 2.9. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 9: Serious adverse events worst-best

Analysis 2.10. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 10: Serious adverse events according to type of drug-eluting stent

Analysis 2.11. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 11: Serious adverse events according to type of ACS

Analysis 2.12. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 12: Serious adverse events according to registration status

Analysis 2.13. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 13: Major cardiovascular events

Analysis 2.14. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 14: Major cardiovascular events best-worst

Analysis 2.15. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 15: Major cardiovascular events worst-best

Analysis 2.16. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 16: Major cardiovascular events according to type of drug-eluting stent

Analysis 2.17. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 17: Major cardiovascular events according to type of ACS

Analysis 2.18. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 18: Major cardiovascular events according to registration status

Analysis 2.19. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 19: Cardiovascular mortality

Analysis 2.20. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 20: Cardiovascular mortality best-worst

Analysis 2.21. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 21: Cardiovascular mortality worst-best

Analysis 2.22. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 22: Cardiovascular mortality according to type of drug-eluting stent

Analysis 2.23. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 23: Cardiovascular mortality according to type of ACS

Analysis 2.24. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 24: Cardiovascular mortality according to registration status

Analysis 2.25. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 25: Myocardial infarction

Analysis 2.26. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 26: Myocardial infarction best-worst

Analysis 2.27. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 27: Myocardial infarction worst-best

Analysis 2.28. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 28: Myocardial infarction according to type of drug-eluting stent

Analysis 2.29. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 29: Myocardial infarction according to type of ACS

Analysis 2.30. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 30: Myocardial infarction according to registration status

Analysis 2.31. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 31: Stent thrombosis

Analysis 2.32. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 32: Stent thrombosis best-worst

Analysis 2.33. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 33: Stent thrombosis worst-best

Analysis 2.34. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 34: Stent thrombosis according to type of drug-eluting stent

Analysis 2.35. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 35: Stent thrombosis according to type of ACS

Analysis 2.36. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 36: Stent thrombosis according to registration status

Analysis 2.37. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 37: Target vessel revascularisation

Analysis 2.38. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 38: Target vessel revascularisation best-worst

Analysis 2.39. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 39: Target vessel revascularisation worst-best

Analysis 2.40. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 40: Target vessel revascularisation according to type of drug-eluting stent

Analysis 2.41. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 41: Target vessel revascularisation according to type of ACS

Analysis 2.42. Comparison 2: Drug-eluting stents versus bare-metal stents for acute coronary syndrome at one month, Outcome 42: Target vessel revascularisation according to registration status

APPENDICES

Appendix 1. Glossary

Acute decompensated heart failure: heart failure where the heart is unable to overcome the pressure in the blood vessels, resulting in symptoms such as difficulty breathing, oedema, and fatigue.

Atheroembolism: embolism originating from an atherosclerotic plaque.

Atherosclerosis: arterial wall-thickening due to build up of plaque.

Angiography: visualisation of the blood vessels typically by injection of contrast and using X-ray.

Angina pectoris: medical term for chest pain or discomfort due to ischaemic heart disease.

Arrhythmias: deviation from the normal heart rhythm.

Balloon angioplasty: using a balloon to open a narrowed vessel.

Bifurcation: when a blood vessel splits into two different blood vessels.

Biolimus: the trade name of the drug umirolimus. The mechanism of action is believed to be antiproliferation of smooth muscle cells.

Bioresorbable stents: stents that are absorbed after initial placement with the intent of reducing restenosis.

Cardiomyocyte necrosis: undesirable death of the cells of the heart.

Continuous ischaemia: prolonged reduced blood supply (to the heart).

Coronary vasoconstriction: narrowing of the blood vessels of the heart due to muscle contraction.

Diastolic: the time period in which the ventricles of the heart relax.

Everolimus: a drug used in stents with the aim of reducing restenosis.

Hypertension: increased pressure typically inside the arteries.

Hypotension: decreased pressure typically inside the arteries.

Intraluminal thrombus: formation of a clot inside a vessel.

Ischaemic: reduced blood supply to an organ.

Level of evidence A: data derived from multiple randomised clinical trials or meta-analyses.

Level of evidence B: data derived from a single randomised clinical trial or large non-randomised studies.

Level of evidence C: consensus of opinion of the experts and/or small studies, retrospective studies, registries.

Myocardial band isoenzyme of creatine kinase (CK-MB): a type of the enzyme creatine kinase that is highly specific for the heart.

Myocardial ischaemia: reduced blood supply to the heart.

Myocardial necrosis: death of the muscle tissue of the heart.

Myocardium: the muscle tissue of the heart.

Myocyte: a heart muscle cell.

Neointimal: scar tissue formed in a vessel after an injury.

Nonocclusive thrombus: the blood clot does not completely prevent blood flow through the vessel.

Non-ST-elevation myocardial infarction: a type of heart attack that does not show ST-segment elevation on an electrocardiogram.

Paclitaxel: a drug used in stents with the aim of reducing restenosis.

Paclitaxel stent: a stent using the paclitaxel drug, a drug used in stents with the aim of reducing restenosis.

Percutaneous: through the skin.

Percutaneous coronary intervention: an intervention where a balloon is guided up to the heart through an access sheath penetrating the skin to reduce the narrowing of a blood vessel.

Re-endothelialisation: regrowth of endothelium after injury.

Restenosis: narrowing of a previously narrowed blood vessel due to a blood clot.

Retroperitoneal bleeding: bleeding behind the peritoneum, a membrane lining the abdominal cavity.

Revascularisation: removing the cause of the stenosed blood vessel, thereby allowing blood flow to resume.

Saphenous vein grafts: when performing bypass surgery, the saphenous vein (located in the leg) may be used to bypass the occluded vessel, ultimately re-establishing heart flow.

Sirolimus: a drug used in stents with the aim of reducing restenosis.

Sirolimus stent: a stent using the sirolimus drug, a drug used in stents with the aim of reducing restenosis.

ST: ST is short for the ST-segment, which is a specific segment of the printout when recording an electrocardiogram that is used to differentiate between ST and non-ST myocardial infarction.

ST-elevation myocardial infarction: a type of heart attack that shows ST-segment elevation on an electrocardiogram.

Stenosis: narrowing of a vessel.

Stent thrombosis: blockage of the stent by a blood clot.

Systolic: the time period in which the ventricles of the heart contract.

Tachycardia: a faster-than-normal heart beat, typically above 100 beats per minute.

Thyrotoxicosis: excessive levels of the hormone produced by the thyroid gland, resulting in unwanted symptoms.

Troponin I: a cardiac protein that is released during a heart attack.

Troponin T: a cardiac protein that is released during a heart attack.

Ventricular arrhythmias: deviation from the normal heart rhythm involving the ventricles of the heart.

Zotarolimus: a drug used in stents with the aim of reducing restenosis.

Appendix 2. Search strategies

Database	Date searched	Search strategy
CENTRAL (Cochrane Library)	18/01/2017	#1 MeSH descriptor: [Stents] explode all trees #2 stent* #3 #1 or #2 #4 drug elut* #5 MeSH descriptor: [Sirolimus] explode all trees #6 sirolimus #7 rapamycin #8 paclitaxel #9 taxol #10 MeSH descriptor: [Immunosuppressive Agents] explode all trees #11 coat* stent* #12 MeSH descriptor: [Taxoids] explode all trees #13 taxane* #14 qp2 #15 hexanoyltaxol #16 everolimus #17 abt-578 #18 MeSH descriptor: [Tacrolimus] explode all trees #19 MeSH descriptor: [Dactinomycin] explode all trees #20 actinomycin #21 batimastat #22 MeSH descriptor: [Dexamethasone] explode all trees #23 dexamethasone

(Continued)

#24 MeSH descriptor: [Estradiol] explode all trees

#25 estradiol

#26 praxel

#27 paxene

#28 onxol

#29 anzatax

#30 immunosuppress*

#31 prograf*

#32 meractinomycin

#33 cosmegen

#34 dactinomycin

#35 millicorten

#36 maxidex

#37 decaspray

#38 dexpak

#39 dexasone

#40 oradexon

#41 hexadecadrol

#42 decaject

#43 hexadrol

#44 decameth

#45 methylfluorprednisolone

#46 vivelle

#47 oestradiol

#48 estrace

#49 aerodiol

#50 estraderm

#51 ovocyclin

#52 estramustin*

#53 estracyt

#54 emcyt

#55 tacrolimus

#56 taxoids

#57 zotarolimus

(Continued)

#58 umirolimus

#59 biolimus

#60 pimecrolimus

#61 elidel

#62 #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or #14 or #15 or #16 or #17 or #18 or #19 or #20 or #21 or #22 or #23 or #24 or #25 or #26 or #27 or #28 or #29 or #30 or #31 or #32 or #33 or #34 or #35 or #36 or #37 or #38 or #39 or #40 or #41 or #42 or #43 or #44 or #45 or #46 or #47 or #48 or #49 or #50 or #51 or #52 or #53 or #54 or #55 or #56 or #57 or #58 or #59 or #60 or #61

#63 #3 and #62

#64 eluting stent*

#65 #63 or #64

#66 MeSH descriptor: [Angioplasty, Balloon, Coronary] explode all trees

#67 MeSH descriptor: [Percutaneous Coronary Intervention] explode all trees

#68 balloon angioplast*

#69 (percutaneous adj6 coronary intervention*)

#70 PCI

#71 (intervention* adj6 percutaneous coronary)

#72 (revascularization* adj6 percutaneous coronary)

#73 (angioplast* adj6 coronary)

#74 percutaneous coronary

#75 ((transluminal or trans-luminal) adj6 coronary)

#76 #66 or #67 or #68 or #69 or #70 or #71 or #72 or #73 or #74 or #75

#77 MeSH descriptor: [Myocardial Ischemia] explode all trees

#78 ((myocardial or myocardium or subendocardial or transmural or cardiac or cardial or coronary or heart) adj2 (infarct* or postinfarct* or hypoxi* or anoxi* or failure* or decompensation or insufficien*))

#79 (heart disease* or coronary disease* or IHD or CIHD or CHD)

#80 (myocardial dysfunction or angina or stenocardia)

#81 ((ischemi* or ischaemi*) adj2 (myocardium or myocardial or heart or coronary or cardiac or cardial or subendocardial or cardiomyopath*))

#82 ((artery occlusion* or artery disease* or arterioscleros* or atheroscleros*) adj2 coronary)

#83 #77 or #78 or #79 or #80 or #81 or #82

#84 MeSH descriptor: [Acute Coronary Syndrome] explode all trees

#85 MeSH descriptor: [Myocardial Infarction] explode all trees

#86 MeSH descriptor: [Coronary Thrombosis] explode all trees

#87 coronary thrombosis

(Continued)

#88 acute coronary

#89 MeSH descriptor: [Angina, Unstable] explode all trees

#90 myocardial infarct*

#91 heart infarct*

#92 acs

#93 ami

#94 (coronary adj3 syndrome*)

#95 acute angina

#96 (unstable adj3 angina)

#97 unstable coronary

#98 #84 or #85 or #86 or #87 or #88 or #89 or #90 or #91 or #92 or #93 or #94 or #95 or #96 or #97

#99 #65 or #76

#100 #83 or #98

#101 #99 and #100

MEDLINE (Ovid)

18/01/2017

1. Stents/
2. stent*.tw.
3. 1 or 2
4. drug elut*.tw.
5. Sirolimus/
6. sirolimus.tw.
7. rapamycin.tw.
8. paclitaxel.tw.
9. taxol.tw.
10. exp Immunosuppressive Agents/
11. coat* stent*.tw.
12. exp Taxoids/
13. taxane*.tw.
14. qp2.tw.
15. hexanoyltaxol.tw.
16. everolimus.tw.
17. abt-578.tw.
18. Tacrolimus/
19. Dactinomycin/

(Continued)

20. actinomycin.tw.
21. batimastat.tw.
22. exp Dexamethasone/
23. dexamethasone.tw.
24. exp Estradiol/
25. estradiol.tw.
26. praxel.tw.
27. paxene.tw.
28. onxol.tw.
29. anzatax.tw.
30. immunosuppress*.tw.
31. prograf*.tw.
32. meractinomycin.tw.
33. cosmegen.tw.
34. dactinomycin.tw.
35. millicorten.tw.
36. maxidex.tw.
37. decaspray.tw.
38. dexpak.tw.
39. dexasone.tw.
40. oradexon.tw.
41. hexadecadrol.tw.
42. decaject.tw.
43. hexadrol.tw.
44. decameth.tw.
45. methylfluorprednisolone.tw.
46. vivelle.tw.
47. oestradiol.tw.
48. estrace.tw.
49. aerodiol.tw.
50. estraderm.tw.
51. ovocyclin.tw.
52. estramustin*.tw.
53. estracyt.tw.

(Continued)

54. emcyt.tw.
55. tacrolimus.tw.
56. taxoids.tw.
57. zotarolimus.tw.
58. umirolimus.tw.
59. biolimus.tw.
60. pimecrolimus.tw.
61. elidel.tw.
62. or/4-61
63. 3 and 62
64. eluting stent*.tw.
65. 63 or 64
66. exp Angioplasty, Balloon, Coronary/ or exp Percutaneous Coronary Intervention/
67. balloon angioplast*.tw.
68. (percutaneous adj6 coronary intervention*).tw.
69. PCI.tw.
70. (intervention* adj6 percutaneous coronary).tw.
71. (revascularization* adj6 percutaneous coronary).tw.
72. (angioplast* adj6 coronary).tw.
73. percutaneous coronary.tw.
74. ((transluminal or trans-luminal) adj6 coronary).tw.
75. or/66-74
76. exp Myocardial Ischemia/
77. ((myocardial or myocardium or subendocardial or transmural or cardiac or cardial or coronary or heart) adj2 (infarct* or postinfarct* or hypoxi* or anoxi* or failure* or decompensation or insufficien*)).tw.
78. (heart disease* or coronary disease* or IHD or CIHD or CHD).tw.
79. (myocardial dysfunction or angina or stenocardia).tw.
80. ((ischemi* or ischaemi*) adj2 (myocardium or myocardial or heart or coronary or cardiac or cardial or subendocardial or cardiomyopath*)).tw.
81. ((artery occlusion* or artery disease* or arterioscleros* or atheroscleros*) adj2 coronary).tw.
82. or/76-81
83. Acute Coronary Syndrome/
84. exp Myocardial Infarction/

(Continued)

85. exp Coronary Thrombosis/
86. coronary thrombosis.tw.
87. acute coronary.tw.
88. exp Angina, Unstable/
89. myocardial infarct*. tw.
90. heart infarct*.tw.
91. acs.tw.
92. ami.tw.
93. (coronary adj3 syndrome*).tw.
94. acute angina.tw.
95. (unstable adj3 angina).tw.
96. unstable coronary.tw.
97. or/83-96
98. randomized controlled trial.pt.
99. controlled clinical trial.pt.
100. randomized.ab.
101. placebo.ab.
102. drug therapy.fs.
103. randomly.ab.
104. trial.ab.
105. groups.ab.
106. or/98-105
107. exp animals/ not humans.sh.
108. 106 not 107
109. 65 or 75
110. 82 or 97
111. 108 and 109 and 110

Embase (Ovid)

18/01/2017

1. exp stent/
2. stent*.tw.
3. 1 or 2
4. exp drug eluting stent/
5. drug elut*.tw.
6. sirolimus.tw.
7. exp rapamycin/

(Continued)

8. rapamycin.tw.
9. paclitaxel/
10. paclitaxel.tw.
11. taxol.tw.
12. exp immunosuppressive agent/
13. coat* stent*.tw.
14. taxoid/
15. taxane*.tw.
16. qp2.tw.
17. hexanoyltaxol.tw.
18. everolimus/
19. everolimus.tw.
20. abt-578.tw.
21. tacrolimus/
22. dactinomycin/
23. actinomycin.tw.
24. batimastat.tw.
25. dexamethasone/
26. dexamethasone.tw.
27. estradiol/
28. estradiol.tw.
29. praxel.tw.
30. paxene.tw.
31. onxol.tw.
32. anzatax.tw.
33. immunosuppress*.tw.
34. prograf*.tw.
35. meractinomycin.tw.
36. cosmegen.tw.
37. dactinomycin/
38. dactinomycin.tw.
39. millicorten.tw.
40. maxidex.tw.
41. decaspray.tw.

(Continued)

42. dexpak.tw.
43. dexasone.tw.
44. oradexon.tw.
45. hexadecadrol.tw.
46. decaject.tw.
47. hexadrol.tw.
48. decameth.tw.
49. methylfluorprednisolone.tw.
50. vivelle.tw.
51. oestradiol.tw.
52. estrace.tw.
53. aerodiol.tw.
54. estraderm.tw.
55. ovocyclin.tw.
56. estramustin*.tw.
57. estracyt.tw.
58. emcyt.tw.
59. tacrolimus/
60. tacrolimus.tw.
61. taxoids.tw.
62. zotarolimus/
63. zotarolimus.tw.
64. umirolimus/
65. umirolimus.tw.
66. biolimus.tw.
67. pimecrolimus/
68. pimecrolimus.tw.
69. elidel.tw.
70. or/4-69
71. 3 and 70
72. eluting stent*.tw.
73. 71 or 72
74. exp percutaneous coronary intervention/
75. balloon angioplast*.tw.

(Continued)

76. (percutaneous adj6 coronary intervention*).tw.
77. PCI.tw.
78. (intervention* adj6 percutaneous coronary).tw.
79. (revascularization* adj6 percutaneous coronary).tw.
80. (angioplast* adj6 coronary).tw.
81. percutaneous coronary.tw.
82. ((transluminal or trans-luminal) adj6 coronary).tw.
83. or/74-82
84. exp heart muscle ischemia/
85. ((myocardial or myocardium or subendocardial or transmural or cardiac or cardial or coronary or heart) adj2 (infarct* or postinfarct* or hypoxi* or anoxi* or failure* or decompensation or insufficien*)).tw.
86. (heart disease* or coronary disease* or IHD or CIHD or CHD).tw.
87. (myocardial dysfunction or angina or stenocardia).tw.
88. ((ischemi* or ischaemi*) adj2 (myocardium or myocardial or heart or coronary or cardiac or cardial or subendocardial or cardiomyopath*)).tw.
89. ((artery occlusion* or artery disease* or arterioscleros* or atheroscleros*) adj2 coronary).tw.
90. or/84-89
91. exp acute coronary syndrome/
92. exp heart infarction/
93. coronary artery thrombosis/
94. coronary thrombosis.tw.
95. acute coronary.tw.
96. exp unstable angina pectoris/
97. myocardial infarct*.tw.
98. heart infarct*.tw.
99. acs.tw.
100. ami.tw.
101. (coronary adj3 syndrome*).tw.
102. acute angina.tw.
103. (unstable adj3 angina).tw.
104. unstable coronary.tw.
105. or/91-104
106. random\$.tw.
107. factorial\$.tw.

(Continued)

108. crossover\$.tw.
109. cross over\$.tw.
110. cross-over\$.tw.
111. placebo\$.tw.
112. (doubl\$ adj blind\$).tw.
113. (singl\$ adj blind\$).tw.
114. assign\$.tw.
115. allocat\$.tw.
116. volunteer\$.tw.
117. crossover procedure/
118. double blind procedure/
119. randomized controlled trial/
120. single blind procedure/
121. or/106-120
122. (animal/ or nonhuman/) not human/
123. 121 not 122
124. 73 or 83
125. 90 or 105
126. 123 and 124 and 125

SCI-Expanded (Web
of Science) (Thomson
reuters)

19/01/2017

#33 #32 AND #31 AND #14

#32 TS=(random* or blind* or allocat* or assign* or trial* or placebo* or crossover* or cross-over*)

#31 #30 OR #20

#30 #29 OR #28 OR #27 OR #26 OR #25 OR #24 OR #23 OR #22 OR #21

#29 TS=unstable coronary

#28 TS=(unstable NEAR/3 angina)

#27 TS=acute angina

#26 TS=(coronary NEAR/3 syndrome*)

#25 TS=(acs or ami)

#24 TS=heart infarct*

#23 TS=myocardial infarct*

#22 TS=acute coronary

#21 TS=coronary thrombosis

#20 #19 OR #18 OR #17 OR #16 OR #15

(Continued)

#19 TS=((("artery occlusion*" or "artery disease*" or arterioscleros* or atheroscleros*) NEAR/2 coronary)

#18 TS=((("ischemi*" or ischaemi*" NEAR/2 (myocardium or myocardial or heart or coronary or cardiac or cardial or subendocardial or cardiomyopath*))

#17 TS=(myocardial dysfunction or angina or stenocardia)

#16 TS=(heart disease* or coronary disease* or IHD or CIHD or CHD)

#15 TS=((("myocardial or myocardium or subendocardial or transmural or cardiac or cardial or coronary or heart) NEAR/2 (infarct* or postinfarct* or hypoxi* or anoxi* or failure* or decompensation or insufficien*))

#14 #13 OR #5

#13 #12 OR #11 OR #10 OR #9 OR #8 OR #7 OR #6

#12 TS=(trans-luminal NEAR/6 coronary)

#11 TS=(transluminal NEAR/6 coronary)

#10 TS="percutaneous coronary"

#9 TS=(angioplast* NEAR/6 coronary)

#8 TS=PCI

#7 TS=(percutaneous NEAR/6 "coronary intervention*")

#6 TS="balloon angioplast*"

#5 #4 AND #1

#4 #3 OR #2

#3 TS=(sirolimus or rapamycin or paclitaxel or taxol or taxoids or taxane* or gp2 or hexanoyltaxol or everolimus or abt-578 or tacrolimus or dactinomycin or actinomycin or batimastat or dexamethasone or estradiol or praxel or paxene or onxol or anzatax or immunosuppress* or prograf* or meractinomycin or cosmegen or dactinomycin or millicorten or maxidex or decaspray or dexpak or dexasone or oradexon or hexadecadrol or decaject or hexadrol or decameth or methylfluorprednisolone or vivelle or oestradiol or estrace or aerodiol or estraderm or ovocyclin or estramustin* or estracyt or emcyt or tacrolimus or zotarolimus or umirolimus or biolimus or pimecrolimus or elidel)

#2 TS=(drug elut* or eluting stent* or coat* stent*)

#1 TS=stent*

BIOSIS (Web of Science) 19/01/2017

#33 #32 AND #31 AND #14

#32 TS=(random* or blind* or allocat* or assign* or trial* or placebo* or crossover* or cross-over*)

#31 #30 OR #20

#30 #29 OR #28 OR #27 OR #26 OR #25 OR #24 OR #23 OR #22 OR #21

#29 TS=unstable coronary

#28 TS=(unstable NEAR/3 angina)

#27 TS=acute angina

#26 TS=(coronary NEAR/3 syndrome*)

(Continued)

#25 TS=(acs or ami)

#24 TS=heart infarct*

#23 TS=myocardial infarct*

#22 TS=acute coronary

#21 TS=coronary thrombosis

#20 #19 OR #18 OR #17 OR #16 OR #15

#19 TS=((("artery occlusion*" or "artery disease*" or arterioscleros* or atheroscleros*) NEAR/2 coronary)

#18 TS=((("ischemi*" or ischaemi*) NEAR/2 (myocardium or myocardial or heart or coronary or cardiac or cardial or subendocardial or cardiomyopath*))

#17 TS=(myocardial dysfunction or angina or stenocardia)

#16 TS=(heart disease* or coronary disease* or IHD or CIHD or CHD)

#15 TS=((("myocardial or myocardium or subendocardial or transmural or cardiac or cardial or coronary or heart) NEAR/2 (infarct* or postinfarct* or hypoxi* or anoxi* or failure* or decompensation or insufficien*))

#14 #13 OR #5

#13 #12 OR #11 OR #10 OR #9 OR #8 OR #7 OR #6

#12 TS=(trans-luminal NEAR/6 coronary)

#11 TS=(transluminal NEAR/6 coronary)

#10 TS="percutaneous coronary"

#9 TS=(angioplast* NEAR/6 coronary)

#8 TS=PCI

#7 TS=(percutaneous NEAR/6 "coronary intervention*")

#6 TS="balloon angioplast*"

#5 #4 AND #1

#4 #3 OR #2

#3 TS=(sirolimus or rapamycin or paclitaxel or taxol or taxoids or taxane* or gp2 or hexanoyltaxol or everolimus or abt-578 or tacrolimus or dactinomycin or actinomycin or batimastat or dexamethasone or estradiol or praxel or paxene or onxol or anzatax or immunosuppress* or prograf* or meractinomycin or cosmegen or dactinomycin or millicorten or maxidex or decaspray or dexpak or dexasone or oradexon or hexadecadrol or decasject or hexadrol or decameth or methylfluorprednisolone or vivelle or oestradiol or estrace or aerodirol or estraderm or ovocyclin or estramustin* or estracyt or emcyt or tacrolimus or zotarolimus or umirolimus or biolimus or pimecrolimus or elidel)

#2 TS=(drug elut* or eluting stent* or coat* stent*)

#1 TS=stent*

LILACS (Bireme)

19/01/2017

(stent\$ or drug elut\$ or balloon angioplast\$ or percutaneous coronary intervention\$ or PCI) and (heart disease\$ or coronary disease\$ or IHD or CIHD or CHD or coronary thrombosis or acute coronary or myocardial infarct\$ or heart

(Continued)

infarct\$ or acs or ami or coronary syndrome\$ or unstable coronary or myocardial dysfunction or angina or stenocardia or myocardial ischemia)

www.clinicaltrials.gov	30/03/17	Searched using 'stent' and using either 'acute coronary syndrome', 'ischaemic heart disease', 'stemi', 'nSTEMI', or 'unstable angina' as the condition
the World Health Organization International Clinical Trials Registry Platform (ICTRP) (www.app-s.who.int/trialsearch)	31/03/17	Searched using 'stent' and 'acute coronary syndrome' or 'stemi' or 'nSTEMI' or 'unstable angina'
European Medicines Agency (EMA) (www.ema.europa.eu/ema/)	03/04/17	Searched using 'drug-eluting stent', 'drug eluting stent', 'bare-metal stent', 'bare metal stent'
the Food and Drug Administration (FDA) (www.fda.gov)	04/04/17	Searched using 'drug-eluting stent', 'drug eluting stent', 'bare-metal stent', 'bare metal stent'
Google Scholar	05/04/17	drug eluting stent bare metal stent "randomized trial" OR "randomised trial"

Appendix 3. Details on assessment of risk of bias

We will classify each trial according to the domains below for each outcome result.

Random sequence generation

- Low risk: if sequence generation is achieved using a computer random number generator or a random numbers table. We will also consider drawing lots, tossing a coin, shuffling cards, and throwing dice as adequate if an independent adjudicator performs these methods.
- Unclear risk: if there is insufficient information to permit judgement of 'low risk' or 'high risk'.
- High risk: if the allocation sequence is not randomised or only quasi-randomised.

Allocation sequence concealment

- Low risk: if the allocation of participants results from a central independent unit, on-site locked computer, identical-looking numbered, sealed, opaque envelopes, drug bottles or containers prepared by an independent investigator.
- Unclear risk: if there is insufficient information to permit judgement of 'low risk' or 'high risk'.
- High risk: if the allocation sequence is known to the investigators who assigned participants.

Blinding of participants and personnel

- Low risk: if the participants and the personnel are blinded to treatment allocation and this is described.
- Unclear risk: if there is insufficient information to permit judgement of 'low risk' or 'high risk'.
- High risk: if blinding of participants and personnel is not performed.

Blinding of outcome assessment

- Low risk: if the trial investigators performing the outcome assessments, analyses, and calculations are blinded to the intervention.
- Unclear risk: if there is insufficient information to permit judgement of 'low risk' or 'high risk'.
- High risk: if blinding of outcome assessment is not performed.

Incomplete outcome data

- Low risk: (1) there are no dropouts or withdrawals for all outcomes, or (2) the numbers and reasons for withdrawals and dropouts for all outcomes are clearly stated, can be described as being similar in both groups, and the trial handles missing data appropriately in

intention-to-treat analysis using proper methodology, e.g. multiple imputations.* As a general rule, we will judge the trial as at low risk of bias due to incomplete outcome data if the number of dropouts is less than 5%. However, the 5% cutoff is not definitive.

- Unclear risk: if there is insufficient information to permit judgement of 'low risk' or 'high risk'.
- High risk: the pattern of dropouts can be described as being different in the two intervention groups, or the trial uses improper methodology in dealing with the missing data, e.g. last observation carried forward.

*Multiple imputation is a general approach to the problem of missing data. It aims to allow for the uncertainty about the missing data by creating several different plausible imputed datasets and appropriately combining results obtained from each of them. The first stage is to create multiple copies of the dataset, with the missing values replaced by imputed values. These are sampled from their predictive distribution based on the observed data; thus, multiple imputation is based on a Bayesian approach. The imputation procedure must fully account for all uncertainty in predicting the missing values by injecting appropriate variability into the multiple imputed values. The second stage is to use standard statistical methods to fit the model of interest to each of the imputed datasets. The estimated associations from the imputed datasets will differ and are only useful when averaged together to give overall estimated associations. Valid inferences are obtained because we are averaging over the distribution of the missing data given the observed data ([Sterne 2009](#)).

Selective outcome reporting

- Low risk: a protocol is published before or at the time the trial begins, and the outcomes called for in the protocol are reported on. If there is no protocol, or the protocol is published after the trial begins, reporting of the primary outcomes will grant the trial a grade of low risk of bias.
- Unclear risk: if there is no protocol and the primary outcomes are not reported on.
- High risk: if the outcomes that are called for in the protocol are not reported on.

Other bias risk

- Low risk of bias: the trial appears to be free of other components (e.g. academic bias or for-profit bias) that could put it at risk of bias.
- Unclear risk of bias: the trial may or may not be free of other components that could put it at risk of bias.
- High risk of bias: there are other factors in the trial that could put it at risk of bias (e.g. authors have conducted trials on the same topic, for-profit bias, etc.).

Overall risk of bias

- Low risk of bias: we will classify the outcome result as overall 'low' risk of bias only if we classify all of the bias domains described in the aforementioned text as low risk of bias. Due to the nature of the percutaneous coronary intervention procedure, we do not expect to find any trials at low risk of bias. We provide a description of how we will deal with this scenario in [Data synthesis](#).
- High risk of bias: we will classify the outcome result as 'high' risk of bias if we classify any of the bias domains in the aforementioned text as 'unclear' or 'high' risk of bias.

We will grade each potential source of bias as high, low, or unclear and provide a quote from the study report together with a justification for our judgement in the 'Risk of bias' table. We will summarise the 'Risk of bias' judgements across different studies for each of the domains listed. Where information on risk of bias relates to unpublished data or correspondence with a trialist, we will note this in the 'Risk of bias' table.

When considering treatment effects, we will take into account the risk of bias for the studies that contribute to that outcome.

WHAT'S NEW

Date	Event	Description
9 June 2021	Review declared as stable	The review question is considered by the editorial team not to be of current clinical relevance and this review is therefore not being updated.

HISTORY

Protocol first published: Issue 12, 2016

Review first published: Issue 8, 2017

CONTRIBUTIONS OF AUTHORS

Joshua Feinberg (JF): drafted the protocol, extracted data, co-ordinated the review, conceived the review, designed the review, interpreted the data by providing a methodological view, and revised the review.

Emil Eik Nielsen (EEN): drafted the protocol, extracted data, drafted the review, interpreted the data by providing a methodological view, and revised the review.

Janette Greenhalgh (JG) commented on the review.

Juliet Hounscome (JH) commented on the review.

Naqash J Sethi (NJS): commented on the review.

Sanam Safi (SS): commented on the review.

Christian Gluud (CG): revised the protocol, interpreted the data by providing a methodological and clinical view, and commented on and revised the review.

Janus C Jakobsen (JCJ): revised the protocol, analysed the data, interpreted the data by providing a methodological and clinical view, and commented on and revised the review.

All authors agreed on the final version.

DECLARATIONS OF INTEREST

The performance of this review is free of any real or perceived bias introduced by receipt of any benefit in cash or kind, on any subsidy derived from any source that may have or be perceived to have an interest in the outcomes of this review.

Joshua Feinberg (JF): no conflict of interest.

Emil Eik Nielsen (EEN): no conflict of interest.

Janette Greenhalgh (JG): no conflict of interest.

Juliet Hounscome (JH): no conflict of interest.

Naqash J Sethi (NS): no conflict of interest.

Sanam Safi (SS): no conflict of interest.

Christian Gluud (CG): member of the Copenhagen Trial Unit task force for developing Trial Sequential Analysis methods, manuals, and software.

Janus C Jakobsen (JCJ): no conflict of interest.

SOURCES OF SUPPORT

Internal sources

- No sources of support provided

External sources

- The Cochrane Heart Group US Satellite is supported by intramural support from the Northwestern University Feinberg School of Medicine and the Northwestern University Clinical and Translational Science (NUCATS) Institute (UL1TR000150), USA
- This project was supported by the National Institute for Health Research (NIHR), via Cochrane Infrastructure funding to the Heart Group. The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the Systematic Reviews Programme, NIHR, National Health Service (NHS), or the Department of Health, UK

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

We did not present a table describing the serious adverse events in each trial. The only serious adverse events reported were all-cause mortality, cardiovascular mortality, myocardial infarction, and target vessel revascularisation, and these were already reported in our other outcome analyses.

We renamed the time point 'time point at three months or less' to 'one month', as all included trials for this outcome reported at one month. If in the future we find trials reporting at any time point other than one month that also report at three months or less, we will change the name back.

In analyses where our visual inspection showed signs of asymmetry, we used the 'trim and fill' method to assess the potential effect of publication bias.

We changed the diversity-adjusted required information size for secondary outcomes from 10% to 20% (it was a typo in the original protocol).

We added anticipated intervention effects for the secondary outcomes in the Trial Sequential Analysis.

We added an estimate of the diversity-adjusted required information size for the exploratory outcomes.

We added that a P value less than 2.5% would be considered as significant for the secondary outcomes.

INDEX TERMS

Medical Subject Headings (MeSH)

Acute Coronary Syndrome [mortality] [*therapy]; Cause of Death; *Drug-Eluting Stents [adverse effects]; Randomized Controlled Trials as Topic; *Stents [adverse effects]

MeSH check words

Humans