Published in final edited form as:

Immunobiology. 2019 March ; 224(2): 196–206. doi:10.1016/j.imbio.2018.11.007.

Comparative Analysis of Expression of Microbial Sensing Molecules in Mucosal Tissues with Periodontal Disease

J.L. Ebersole^{1,*}, S. Kirakodu¹, M.J. Novak¹, L. Orraca³, A. J. Stromberg⁴, J. Gonzalez-Martinez⁵, A. Burgos⁵, and O.A. Gonzalez^{1,2}

¹Center for Oral Health Research, University of Kentucky, Lexington, KY

²Division of Periodontics, College of Dentistry, University of Kentucky, Lexington, KY

³School of Dental Medicine, University of Puerto Rico, San Juan, PR

⁴Department of Statistics, College of Arts and Sciences, University of Kentucky, Lexington, KY

⁵Caribbean Primate Research Center, University of Puerto Rico, Toa Baja, PR.

Abstract

Host-derived pattern recognition receptors (PRRs) are necessary for effective innate immune engagement of pathogens that express microbial-associated molecular patterns (MAMP) ligands for these PRRs. This study used a nonhuman primate model to evaluate the expression of these sensing molecules in gingival tissues. *Macaca mulatta* aged 12–24 with a healthy periodontium (n=13) or periodontitis (n=11) provided gingival tissues for assessment of naturally-occurring periodontitis. An additional group of animals (12–23 years; n=18) was subjected to a 5 month longitudinal study examining the initiation and progression of periodontitis, RNA was isolated and microarray analysis conducted for gene expression of the sensing PRRs. The results demonstrated increased expression of various PRRs in naturally-occurring established periodontitis. Selected PRRs also correlated with both bleeding on probing (BOP) and pocket depth (PD) in the animals. The longitudinal model demonstrated multiple TLRs, as well as selected other PRRs that were significantly increased by 2 weeks during initiation of the lesion. While gene expression levels of various PRRs correlated with BOP and PD at baseline and resolution of disease, few correlated with these clinical parameters during initiation and progression of the lesion. These findings suggest that the levels of various PRRs are affected in established periodontitis lesions, and that PRR expression increased most dramatically during the initiation of the disease process, presumably in response to the juxtaposed microbial challenge to the tissues and goal of reestablishing homeostasis.

*Corresponding author: Jeffrey L. Ebersole, Complete Address: Department of Biomedical Sciences, School of Dental Medicine, University of Nevada Las Vegas, 1001 Shadow Lane, Las Vegas, NV 89106. Phone: (702)774-4152. Fax number: (702)774-2632. jeffrey.ebersole@unlv.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The authors state no conflict of interest in the experimental design or data reported.

Keywords

Pattern recognition receptors; periodontal disease; nonhuman primates; gingiva; aging

INTRODUCTION

The oral mucosa harbors a very complex microbial ecology that varies among the various niches in the oral cavity (Avila, et al., 2009, Jenkinson, 2011, Alcaraz, et al., 2012, Zarco, et al., 2012). Furthermore in these niches there is an array of commensal bacteria that colonizes the sites during health, as well as a range of potential opportunistic pathogenic bacteria that can emerge within the microbiota and trigger disease processes (Wade, 2013, Darveau, 2009, Tlaskalova-Hogenova, et al., 2004, Dixon, et al., 2004, Cugini, et al., 2013, Periasamy and Kolenbrander, 2009, Hajishengallis and Lamont, 2012).

The innate immune system has evolved to encompass an array of cellular receptors that are capable of recognizing a plethora of microbial components, enabling the cells to recognize, engage and respond to these microbial or pathogen-associated molecular patterns (MAMPs, PAMPs)(Hajishengallis, 2014, Ji and Choi, 2013, Hans and Hans, 2011). The pattern recognition receptors (PRRs) on host cells were originally described related to immune cells, such as macrophages ($M\Phi$) and dendritic cells (DCs). However, it is now recognized that the PRRs are distributed more broadly across most cell types, including epithelial cells at mucosal surfaces (Walsh, et al., 2013, Kumar, et al., 2013, Jounai, et al., 2012). The PRRs have now been delineated to exist as cell-surface associated [*e.g.* Toll-like receptors (TLRs)], intracellular [*e.g.* nucleotide-binding oligomerization domains (NODs)], and soluble PRRs (*e.g.* C-reactive protein, serum amyloid A) that comprise early warning system for innate immunity (Kumar, et al., 2011). Thus, these microbial sensing molecules interact with a range of bacterial (*e.g.* lipopolysaccharide, lipoteichoic acid, flagella, DNA), viral (*e.g.* DNA, RNA), and fungal (*e.g.* mannan) ligands.

It is well recognized that an array of PRRs are differentially expressed in periodontitis tissues (Ebersole, et al., 2013, Benakanakere and Kinane, 2012). These response relationships have been developed conceptually in describing the disease as a host-mediated disruption of microbial homeostasis (Ji and Choi, 2013, Bartold and Van Dyke, 2013). The innate responses critical to this biologic interactions clearly (Fatemi, et al., 2013) involve recognition of microbial components mediated by various PRRs, including the toll-like receptors in both resident and immune cells (Di Benedetto, et al., 2013).

Multiple reports have identified alterations in TLR2 and TLR4 expression level in inflamed tissues of periodontitis (Myneni, et al., 2013), associated with smoking (Fatemi, et al., 2013), and with diabetes as a co-morbidity (Promsudthi, et al., 2014). These specific PRRs have also been demonstrated to be significant in a rodent model of *P. gingivalis* exacerbated ligature-induced RANKL-dependent periodontitis (Lin, et al., 2014).

More recently, studies have broadened the perspective of PRRs contribution to the chronic inflammation of tissue destruction of periodontitis. MYD88 mRNA was increased in periodontitis tissues and was elevated to a greater degree in severe periodontitis. While this

study did not find much effect on various TLRs, NLRs, or signaling molecules CD14 and TRIF, changes in MYD88 support a critical TLR transduction signaling change in periodontitis (Ghaderi, et al., 2014). A recent report by Sahingur et al. (Sahingur, et al., 2013) demonstrated upregulation of TLR9 and TLR8 in chronic periodontitis tissues, as well as significant correlations among PRRs, and detection of changes in intracellular PRRs, AIM2 and DAI, in diseased gingival tissues. These types of findings have been extended in attempts to better understand potential triggers and molecular mechanisms underlying these differences. Recent work has identified that gingival tissues in periodontitis displayed a differential TLR2 promoter methylation, thus suggesting a likelihood of altered modulation in the expression of this important PRR in inflamed tissues (Benakanakere, et al., 2015). This PRR receptor changes connects with a report by Park et al. (Park, et al., 2014) that demonstrated elevated inflammasome components were expressed in chronic periodontitis, including IL-1 β . These authors also showed that *P. gingivalis* induced IL-1 β and pyroptotic cell death in a macrophage cell line through engagement of NLRP3 and AIM2 inflammasome activation. Additionally, they described the detection of priming signals from TLR2 and TLR4 activation that preceded the *P. gingivalis* induced IL-1 β release representing a proinflammatory mediator activity in the periodontitis tissues.

Thus, while numerous PRRs have been identified in gingival tissues, the relationship of changes in the expression and levels of these innate immune molecules to health and disease processes remains unclear. This report uses a nonhuman primate model of naturally-occurring periodontitis and a prospective model of ligature-induced periodontitis to document alterations in expression of these microbial sensing molecules that are related to this mucosal disease.

METHODS

Nonhuman primate model and Oral Clinical Evaluation

Rhesus monkeys (*Macaca mulatta*) (n=24; 10 females and 14 males) housed at the Caribbean Primate Research Center (CPRC) at Sabana Seca, Puerto Rico, were used in these studies. Healthy animals (n=13) and naturally-occurring periodontitis animals (n=11) were 12–23 years of age. The 18 animals (10 females and 8 males) in the longitudinal study of ligature-induced disease were 12–23 years of age. The nonhuman primates are typically fed a 20% protein, 5% fat, and 10% fiber commercial monkey diet (diet 8773, Teklad NIB primate diet modified: Harlan Teklad). The diet is supplemented with fruits and vegetables, and water is provided *ad libitum* in an enclosed corral setting.

A protocol approved by the Institutional Animal Care and Use Committee (IACUC) of the University of Puerto Rico, enabled anesthetized animals to be examined for clinical measures of periodontal including probing pocket depth (PD), and bleeding on probing (BOP) as we have described previously (Ebersole, et al., 2008).

Tissue sampling and gene expression microarray analysis

A buccal gingival sample from either healthy or periodontitis-affected tissue from the premolar/molar maxillary region of each animal was taken using a standard gingivectomy

technique, and maintained frozen in RNAlater solution. Total RNA was isolated from each gingival tissue using a standard procedure as we have described and tissue RNA samples submitted to the microarray core to assess RNA quality analyze the transcriptome using the GeneChip® Rhesus Macaque Genome Array or GeneChip® Rhesus Gene 1.0 ST Array (Affymetrix) (Affymetrix) (Meka, et al., Gonzalez, et al., 2011). Individual samples were used for gene expression analyses. Table 1 lists the microbial sensing gene set examined in this report.

Based upon the microarray outcomes we selected 7 representative genes and performed a qPCR analysis using a standard technique in our laboratory employing a Roche 480 LightCycler (Gonzalez, et al., 2014) with Tm 62°C. Primers were prepared for SAA1 (forward - CAGCGATGCCAGAGAGAATATC; reverse – CAGCGATGCCAGAGAGAATATC; amplicon 121 bp), CD14 (forward – GCCCTAAACTCCCTCAATCTG; reverse – CAGTCTGTTGCAGCTGAGAT; amplicon 99 bp), NAIP (forward - GGCTCTTGGATGCAGATGATA; reverse – ATGATTGGAGAGAACGGCAATA; amplicon 112 bp), TLR2 (forward - GATGCTGCCATTCTTGTCTTC; reverse – CAGGTAGGTCTTGGTGTTCATT; amplicon 99 bp), ZBP1/DAI (forward – TCAGCCCATCACTCGAAAC; reverse – ATGAGGCTTCATCCACATAGTC; amplicon 113 bp), CLEC7A (forward – CAAGTGTCTTCCCAACCTGATA; reverse – GAATGTTGATCCATCCTCCCA; amplicon 93 bp), NOD2 (forward – GAGGCAGTTCCATTTCATTTGT; reverse – TGCTTAGAAGGAAGGGCTTAAT; amplicon 96 bp), and GAPDH (forward – GGTGTGAACCATGAGAAGTATGA; reverse – GAGTCCTTCCACGATACCAAAG; amplicon 123 bp) genes, designed using software PrimerQuest at Integrated DNA Technologies website (www.idtdna.com) and were synthesized by Integrated DNA Technologies, Inc. (Coralville, IA). The level of message was determined and those levels compared across the RNA samples prepared from the healthy animals compared with the periodontitis group.

Data analysis

Normalization of values across the chips was accomplished through signal intensity standardization across each chip using Affymetrix PLIER algorithm. The arrays contained matched and mismatched pairs allowing the MAS 5 algorithm to be used. For each gene we first determined differences in expression across the groups using ANOVA (version 9.3, SAS Inc., Cary, NC). The healthy tissues were compared to the periodontitis tissues using a *t*-test and accepting a p-value ≤ 0.05 for significance. The choice of Least Significant Difference for multiple comparisons (ANOVA followed by *t*-tests) provided maximum power given our necessarily small sample sizes. We did determine a correlation between gene expression and clinical measures of periodontitis using a Spearman Rank correlation analysis that was fit to the gene expression by age. A p-value ≤ 0.05 was used to evaluate the significance of the correlation. A portion of the data has been uploaded into the ArrayExpress data base (www.ebi.ac.uk) under accession number: E-MTAB-1977.

RESULTS

Profiles of PRRs

Both CD14 and CLEC4M expression levels were increased in periodontitis compared to healthy gingival tissues (Figure 1A). TLR2, TLR4 and TLR9 levels were also increased significantly in the periodontitis samples, as was the intracellular sensor NAIP. NOD2 another intracellular detection receptor was significantly decreased, whereas ZBP1/DAI increased in the periodontitis samples (Figure 1A&B). An array of soluble PRRs was also examined in disease with no significant differences observed in established periodontitis lesions (Figure 1C).

Figure 2A&B depict the correlations of the PRR gene expression in the periodontitis animals for both BOP and PD. The results demonstrate multiple sensing genes are correlated with increasing BOP during periodontitis, including CLEC isoforms and TLRs. Additionally, COLEC12, PTX3 and TLR1 were all significantly negatively correlated with this measure of inflammation. In contrast, very few of these genes were correlated with mean PD in the established periodontitis animals, although LBP expression levels were positively correlated with both BOP and PD.

PRRs in Progressing Periodontitis

We also evaluated the expression of the PRRs using a longitudinal model of progressing and resolving periodontitis. Following ligature placement, a substantial accumulation of microbial plaque occurs, which initiates an acute and chronic inflammatory response in the local tissues. This leads to progressing periodontitis through approximately 3 months (Smith, et al., 1993, Moritz, et al., 1998, Branch-Mays, et al., 2008). Removal of the ligatures enables a clinical resolution within about 60 days. The clinical changes of inflammation (BOP) and tissue loss (PD) are demonstrated in Figure 3.

Of the 36 PRRs evaluated, 15 demonstrated significant differences from baseline during the longitudinal study (Figure 4). Interestingly, TLR1, TLR2, TLR4, TLR6, TLR7, TLR8, and TLR9 all demonstrated a similar pattern with significant increases within 2 weeks of initiation of the disease. TLR1 levels dropped to baseline by 1 month, while TLR2, 4, 6, 7, 8, and 9 remained elevated throughout the disease progression (3 months). All TLR message levels decreased to baseline after resolution of the disease.

Other cellular PRRs, including NAIP (intracellular), CLEC4E and CD14 (surface) demonstrated a pattern similar to the TLRs with increases during the initiation of the disease that were, in some cases, maintained during disease progression and returned to baseline at 5 months (resolution). In contrast, both CLEC7A and NOD2 showed a pattern of changes with significant decreases during initiation and progression of disease, and returned to baseline at resolution. ZBP1/DAI exhibited a unique profile of expression with increases peaking at 1–3 months of disease; again returning to baseline by 5 months. AIM2 (intracellular) also showed a somewhat unique profile of expression with peak levels attained by 2 weeks that were maintained throughout the experiment, including remaining elevated after resolution of disease. Finally, the only soluble PRR level that was affected by the disease process was

FCN1 (ficolin 1) that was significantly increased at 2 weeks during the acute inflammatory response but returned to baseline by 1 month during the progression of destructive disease.

The relationship of the gene expression profiles during the initiation, progression, and resolution of the disease was examined related to the baseline expression of these microbial sensing molecules. Table 2 shows that for a few of these sensing molecules, the changes in gene expression that occurred with disease were related to baseline gingival levels of the expression in the animals. Interestingly, the primary positive correlations with baseline levels of the expression of these genes occurred at 1 and 3 months of the disease process, suggesting some individual regulation of these responses based upon the existing interaction with the oral microbial ecology prior to disease initiation.

Table 3 summarizes the relationship of gene expression for the sensing molecules and clinical features of BOP and PD in the animals at baseline, during the disease process, and at resolution. Generally, a limited number of these host signaling genes were significantly correlated with BOP at baseline (3/36), although 9 were correlated with PD, not overlapping those related to BOP. These relationships were lost during disease initiation and progression, with no correlations detected. Of particular interest was the substantial frequency of gene expression levels that were significantly correlated with BOP (15/36) and PD (11/36) during the resolution phase of the disease.

Table 4 provides a summary of the qPCR analysis of selected genes as a validation of the microarray results. The findings demonstrate an agreement in level differences for 4/5 genes in the cross-sectional samples of established lesions and all 5 genes from the longitudinal study of periodontitis using these 2 independent techniques.

DISCUSSION

Innate immune responses are controlled by the capacity of the host cells and biomolecules to recognize bacteria, viruses, and fungi through specific ligands (PAMPs, MAMPs) that are structural motifs of the microorganisms (Walsh, et al., 2013, Kumar, et al., 2013, Kumar, et al., 2011, Qian and Cao, 2013, Mogensen, 2009, Atkinson, 2008). This is accomplished through the expression and production of an array of pattern recognition receptors (PRR) that target specific components of these bacterial motifs. These interactions result in changes in cell functions and release of various molecules that more broadly engage host innate immune, inflammatory, and even adaptive immune cells and molecules (Kumar, et al., 2013, Mogensen, 2009). Generally the PRRs appear to function in a nonspecific fashion, albeit individual PRRs have a predilection for certain structural motifs and are thus more individually effective within from a more limited range of microbes (Qian and Cao, 2013, Sukhithasri, et al., 2013).

This report describes alterations in the expression of these PRRs in gingival tissues from nonhuman primates with established periodontal lesions, as well as evaluating the dynamics of the PRR changes in gingival tissues during the initiation, progression, and resolution of ligature-induced periodontitis (Gonzalez, et al., 2012, Oz and Puleo, 2011, Struillou, et al., 2010). We observed 9/35 sensing genes were significantly different in the established

periodontitis tissues. These included cell surface (TLR2, TLR4, TLR9, CD14, and CLEC4M) and intracellular (NAIP, NOD2, ARHGEF2, and ZBP1/DAI) PRRs. TLR2 receptors are important PRRs in immune and non-immune host recognition of microbes. TLR2 primarily engages lipoteichoic acids from Gram-positive bacteria although it can form heterodimers with other TLR and non-TLR receptors [*e.g.* CD36, CD14, CLEC7A (Dectin-1)] to bind additional specific molecular ligand structures on microbes (van Bergenhenegouwen, et al., 2013). We also noted that TLR2 was positively correlated with mean pocket depth in established disease lesions. TLR4 and TLR9 expression was significantly positively correlated with the level of bleeding on probing. In contrast, TLR1 was negatively correlated with BOP and TLR5 was negatively correlated with pocket depth in established lesions. These findings supported potential roles for these PRRs in the diseased gingival tissues; however, there appeared to be some selectivity in their regulation reflecting the level of disease. This might be explained by the broader biologic processes occurring in the gingival tissues in trying to reestablish homeostasis, and/or it may reflect variations in the quality of the microbial biofilms that result in more selective regulatory influences on these cell signaling molecules. Extension of these findings to the longitudinal model of periodontitis provided additional evidence for the role of these PRRs in the process of inflammation and disease progression in the gingival tissues. TLR1, TLR2, TLR4, TLR6, TLR8, and TLR9 all demonstrated significant increases by 2 weeks following initiation of disease. All except TLR1 remained elevated through the first month of disease, with TLR2, TLR4 and TLR6 remaining elevated through the 3 month progression of disease. All of these returned to baseline levels following disease resolution at 5 months. Only TLR7 demonstrated a different pattern with increases peaking at 1 month and remaining significantly elevated within gingival tissues even following resolution. TLR7, like TLR3 and TLR8 receptors, has been suggested to primarily engage nucleic acids as ligands, related to detection of viral infections (Wu, et al., 2013, Szabo and Rajnavolgyi, 2013, Mansson Kvarnhammar, et al., 2013, Kaiko, et al., 2013). Its role as an intracellular sensing PRR may be related to increasing data supporting a role for bacterial invasion of gingival tissues related to the pathogenic potential and expression of disease by these bacteria (Darveau, 2009, Eick, et al., 2006, Colombo, et al., 2006, Vitkov, et al., 2005, Atanasova and Yilmaz, 2014, Park, et al., 2004). Additionally, correlation analyses showed that multiple TLRs at 1 and 3 months of disease were related to baseline gingival tissue levels. Additionally, TLR3, 5, and 9 were significantly positively correlated with BOP at resolution of the disease, while TLR4 and TLR8 were negatively correlated. TLR4 and TLR8 were significantly negatively correlated with pocket depth (PD) in the resolved lesions, supporting a potential role contributing to reestablishing tissue homeostasis with the microbial ecology. Interestingly, while TLR message levels were related to baseline levels in individual animals, they were not quantitatively correlated with clinical disease parameters during disease initiation and progression of disease. These results suggest individual heterogeneity within this group of subject animals related to the characteristics of response to the microbial challenge. However, the complex microbial biofilms appear to uncouple TLR signaling potential related to clinical features of disease during initiation and progression of the lesion.

The CD14 molecule, a co-receptor for bacterial LPSs, exists both anchored to the membrane of immune cells and shed into a soluble form to interact with non-immune cells (Litvack and

Palaniyar, 2010, Jin and Lee, 2008). The expression of CD14 was significantly increased in established periodontitis lesions. It demonstrated a response profile during disease initiation and progression similar to the TLRs. The level of CD14 mRNA was shown to be related to baseline levels of expression during disease initiation and progression, but was generally unrelated to the level of bleeding or pocket depth. As with the TLR transcription, regulation of CD14 appears in response to the increased microbial burden during disease initiation and progression, with levels seeming more animal specific rather than directly linked to the severity of the disease process, *e.g.* BOP, PD. Some information has been developed regarding levels of sCD14 and CD14 gene polymorphisms as related to periodontitis. The limited data available suggested that sCD14 levels in serum were elevated in periodontitis (Pussinen, et al., 2007), while elevated levels of sCD14 in gingival crevicular fluid were correlated with fewer deep pockets in untreated chronic periodontitis patients (Jin and Darveau, 2001). Additionally, it appears that a particular polymorphism was related to higher serum levels of sCD14 and periodontitis (Gong, et al., 2013). Our data implies that elevated levels of CD14 may reflect an initiating and progressing disease process.

Multiple C-type lectin family members are both endocytic and cell surface receptors that are linked to cell signaling directly through NF- κ B activation (Yan, et al., 2013). CLEC4E (Mincle) binds nuclear proteins from damaged cells, as well as carbohydrate residues of pathogens (Yamasaki, et al., 2009, Miyake, et al., 2013, Sharma, et al., 2014). CLEC4M, which is similar to CD209 (DC-SIGN) recognizes various pathogens and interacts with ICAM-3 (Khoo, et al., 2008, Rydz, et al., 2013). CLEC7A (Dectin-1) is a PRR primarily expressed on immune cells and engages various glucan moieties on microbes, and contributes to development of Th17 cells (Sukhithasri, et al., 2013, Rao, et al., 2014, Plato, et al., 2013, Gringhuis, et al., 2012). Multiple of these receptors were significantly increased in established periodontitis lesions, and positively correlated with BOP as a measure of inflammation. Prospectively, CLEC4E showed a significant increase by 2 weeks, while CLEC7A was significantly decreased through the initiation and progression of periodontitis. Rather limited information exist examining C-type lectin PRRs in the gingival tissues related to health or disease. Dectin-1 expression has been shown to be increased in gingival tissues from aged mice unrelated to disease (Liang, et al., 2010), while Tamai et al. (Tamai, et al., 2011) demonstrated the importance of Dectin-1 on oral epithelial cells related to detection of *Candida albicans*. Carvalho et al. (Carvalho, et al., 2012) have also identified that Dectin-1 appears to trigger a predominant Th17 response resulting from engagement with *C. albicans*. While this type of activity has not been identified in gingival tissues, and this type of fungal pathogen is not generally associated with periodontitis, alterations in the family of CLEC PRRs remains rather unexplored for a role in the disease process or resolution. Since Dectin-1 is primarily an immune cell PRR, the contrast between the developing lesion data and established natural lesion results provide a temporal contrast for how this type of receptor might function within the periodontium as part of the innate immune response to the microbial challenge.

The nucleotide-binding oligomerization domain-containing protein receptors (NLRs) are intracellular PRRs (NOD1, NOD2, NLRC4, NLRP3, and NAIP). The NOD PRRs bind intracellular bacterial peptidoglycan components leading to activation of NF κ B (Rasmussen, et al., 2009, Wen, et al., 2013). Various studies in host-pathogen responses in periodontitis

have identified a role for NOD1/2 in sensing periodontal pathogens, regulating molecules that enhance interaction of neutrophils with endothelial cells, and triggering innate responses in periodontal ligament cells (Benakanakere and Kinane, 2012, Zelkha, et al., 2010). NOD1 has been shown to be a critical molecule in bone loss in mice to a murine-specific oral microorganism (Jiao, et al., 2013). Also, Yuan and colleagues (Yuan, et al., 2013) have suggested a pivotal role for NOD2 in controlling inflammatory responses that may link periodontitis with chronic systemic inflammation, such as occurs in atherosclerosis. In our study, NOD2 was significantly decreased in established lesions and was decreased during initiation and progression of the experimental lesions, as well as negatively correlated with BOP at the resolution phase. In contrast, while NOD1 was not found to be significantly different or change with disease, it was highly positively correlated with both BOP and PD in the resolved lesions. Thus, the exact role for NOD involvement in periodontitis initiation and progression, and details on the components of the complex microbial ecology at disease sites remain to be evaluated, but would seem to hold some promise for understanding risk for expression of disease.

NAIP is a sensor component of the NLRC4 inflammasome that recognizes intracellular bacteria pathogens. This molecule is a part of a larger family of Inhibitors of Apoptosis (IAP) and links this biological function with the NLR aspects of the inflammasomes (Kofoed and Vance, 2012). Little information is available regarding NAIP alterations in periodontitis tissues; however, it appears to be increased during inflammatory responses (Beug, et al., 2012) and has recently been reported to interact with bacterial type III secretion proteins to engage the NLRC4 inflammasomes in response to both respiratory and gastrointestinal pathogens (Gong and Shao, 2012). In our models of periodontitis, NAIP was significantly elevated in established periodontitis tissues and was positively correlated with BOP in disease. In addition, it increased significantly during the initiation of experimental periodontitis. This is consistent with our previous findings of decreased apoptotic gene profiles in periodontitis tissues (Gonzalez, et al., 2011, Gonzalez, et al., 2013). NLRC4 (IPAF) indirectly senses specific proteins from pathogenic microbes and helps to assemble the inflammasome complex (Lupfer and Kanneganti, 2013). NLRP3, AIM2 (Absent in Melanoma 2), and NLRC4 inflammasomes have all been shown to contribute to caspase-1 activation and IL-1 β secretion by activated macrophages (Rathinam, et al., 2012, Martinon, et al., 2009). This process has been documented to contribute to mucosal immunity to both lung (Cai, et al., 2012) and intestinal pathogens (Nordlander, et al., 2014) in murine models of infection. Both the NLRP3 and AIM2 inflammasome complexes have been suggested to have a role in periodontitis with NLRP3 being activated in diseased tissues and macrophages (Bostanci, et al., 2009), and AIM2 up-regulated in gingival fibroblasts (Belibasakis, et al., 2013). However, *in vitro*, it has been shown that *P. gingivalis* can repress NLRP3 inflammasome activation that could be triggered by PAMPs and danger associated molecular pattern (DAMPs) signals that require endocytosis (Park, et al., 2014). Our results demonstrated no effects on NLRP3 in either established or progressing lesions in the nonhuman primates. However, NLRC4 was positively correlated with BOP in periodontitis. AIM2 is also a critical part of the inflammasome responses that contributes to host resistance through recognition of bacterial and viral DNA (Vilaysane and Muruve, 2009). IFN γ induces expression of AIM2 that has a role in processing of pro-inflammatory cytokines,

like IL-1 β and IL-18 (Fang, et al., 2014). Recent work by Sahingur and colleagues (Sahingur, et al., 2013) reported an up-regulation of AIM2 in chronic periodontitis tissues, coupled with increases in TLR9 and DAI mRNA and products in the tissues. Our current study showed that AIM2 levels increased by 2 weeks following initiation of disease and maintained a significantly increased level throughout the entire longitudinal study. In contrast, the expression of this gene was unrelated to established lesions, or directly to the clinical features of progressing periodontitis. These findings support alterations in intracellular detection of microbial PAMPs from both Gram-positive and Gram-negative bacteria with disease, and suggest that triggering of the inflammasome assembly relates to the gingival inflammation and bleeding that occurs in periodontitis. The results also suggest that increases in gene products related to inflammasome development and activity may be a process that takes place at earlier stages of the progression of disease, but is not necessarily sustained in established lesions. Alternatively, these findings could indicate that many clinical lesions in the nonhuman primates and humans could actually represent a biologically stable environment at the molecular level.

ARHGEF2 (Rho guanine nucleotide exchange factor 2) activates Rho-GTPases via extracellular stimuli. It has been suggested to be involved in epithelial barrier permeability, antigen presentation, and innate immune responses. It is a tight junction-associated molecule that forms an intracellular sensing system along with NOD1 for the detection of invasive bacterial pathogens that exploit tight junctions for entry/crossing epithelial barriers (Philpott, et al., 2014). Involved not only in sensing peptidoglycan through NOD1, but also in the activation of NF κ B and in innate immune signaling by promoting IL-6 and TNF α secretion by macrophages.

As with the NLRs, Z-DNA-binding protein 1 (ZBP1 or DAI/DLM-1) is an intracellular PRR that recognizes DNA in the cytoplasm, primarily considered as an antiviral mechanism. ZBP1 once activated, elevates the level of antiviral cytokines, such as type I interferons (Vilaysane and Muruve, 2009, Yanai, et al., 2009). Expression of this PRR was significantly elevated in established periodontitis lesions. It was significantly increased during the initiation and progression of disease, albeit peak levels were seen at 1–3 months, which were somewhat different kinetics than other PRRs. It was also highly significantly positively correlated with BOP during disease resolution. Based on its DNA ligand, this PRR has been related to antiviral innate immunity, as well as its potential to signal biomolecules that are immunomodulatory. However, these functions could overlap in detecting intracellular periodontopathogens (Eick, et al., 2006, Vitkov, et al., 2005, Atanasova and Yilmaz, 2014, Park, et al., 2004, Tamai, et al., 2011), or is consistent with observations of the potential role of DNA viruses (*e.g.* CMV, EBV) in the human disease process (Slots, 2010).

Soluble PRRs are antimicrobial molecules that function by binding to the cell wall of various microorganisms, enhancing phagocytosis and complement activation, that lead to microbe destruction. Collectins, ficolins and the family of pentraxins [*e.g.* Serum Amyloid A (SAA), C-Reactive Protein (CRP)] are all secreted by cells as soluble PRRs. COLEC12 promotes binding and phagocytosis of Gram-positive, Gram-negative bacteria and yeast, and also interacts with oxidized LDLs (Canton, et al., 2013, Taylor and Drickamer, 2007). COLEC12 was highly negatively correlated with BOP in the cross-sectional analysis

comparing health to periodontitis tissues. Thus, this relationship could signify a contribution of elevated COLEC12 in the tissues towards modulating the intensity of the vascular inflammatory response in the gingival tissues. The ficolins (FCN1, FCN2) are secreted PRRs released from neutrophils following triggering of fMLP receptors (Endo, et al., 2011, Thomsen, et al., 2011) and can activate the lectin pathway of complement activation upon engagement of N-acetylglucosamine (GlcNAc) residues. We observed an increased levels of ficolins in periodontitis, with FCN2 positively correlated with pocket depth measures in naturally-occurring periodontitis and FCN1 demonstrating a significant increase in levels within 2 weeks of disease initiation in the longitudinal model, but then returning to baseline levels. These finding are consistent with the acute role of the neutrophils in responding to the infectious challenge of the accumulating microbial biofilms at sites of ligature placement. The lack of sustainability of the elevated gene expression may signify the transient role of these PRRs in the disease process and engagement of more robust biomolecules to help control the infection. Pentraxins are divided into short proteins, such as CRP and serum amyloid protein, and long proteins, such as PTX3 and PTX4. PTX3 binds to various microbes to activate the classical complement pathway and enhances the clearance of apoptotic cells (Cieslik and Hrycek, 2012, Deban, et al., 2009). This PRR has been found to be increased in gingival crevicular fluid and plasma with disease progression from health to gingivitis to periodontitis (Pradeep, et al., 2011). In addition, gingival tissues from patients with generalized aggressive periodontitis had a higher mean concentration of PTX3 than tissues from patients with generalized chronic periodontitis or control subjects (Lakshmanan, et al., 2013). Finally, salivary levels of PTX3 correlated with plaque index and bleeding on probing in the chronic periodontitis patients, while serum and salivary PTX3 levels correlated with those of IL-1 β in aggressive periodontitis individuals (Gumus, et al., 2014). We observed a significant negative correlation in expression levels of PTX3 with both BOP and PD only during resolution of disease, which is consistent with a role in for the molecule in the reestablishment of homeostasis of the gingival tissues.

This study provides data that support a role for an array of PRRs during the initiation and progression of periodontal lesions. The results also indicated that the profiles of PRR gene expression during the initiation and early progression of a periodontal disease lesion are somewhat different than those detected in naturally occurring established lesions and suggest that the heterogeneity of local changes in human disease may be a reflection of fundamental differences in the “molecular stage” of the lesion that cannot be discerned from standard clinical measures of gingival index, bleeding on probing, and pocket depth.

ACKNOWLEDGEMENTS

This project was supported by National Institute of Health grants P20GM103538 and UL1TR000117. We express our gratitude to the Caribbean Primate Research Center (CPRC) supported by grant P40RR03640, and the Microarray Core of University Kentucky for their invaluable technical assistance. We thank M. Kirakodu for data management support. The authors state no conflict of interest in the experimental design or data reported.

REFERENCES

- Avila M, Ojcius DM, Yilmaz O 2009 The oral microbiota: living with a permanent guest. *DNA Cell Biol* 28, 405. [PubMed: 19485767]
- Jenkinson HF 2011 Beyond the oral microbiome. *Environ Microbiol* 13, 3077. [PubMed: 21906224]

- Alcaraz LD, Belda-Ferre P, Cabrera-Rubio R, Romero H, Simon-Soro A, Pignatelli M, Mira A 2012 Identifying a healthy oral microbiome through metagenomics. *Clin Microbiol Infect* 18 Suppl 4, 54.
- Zarco MF, Vess TJ, Ginsburg GS 2012 The oral microbiome in health and disease and the potential impact on personalized dental medicine. *Oral Dis* 18, 109. [PubMed: 21902769]
- Wade WG 2013 The oral microbiome in health and disease. *Pharmacol Res* 69, 137. [PubMed: 23201354]
- Darveau RP 2009 The oral microbial consortium's interaction with the periodontal innate defense system. *DNA Cell Biol* 28, 389. [PubMed: 19435427]
- Tlaskalova-Hogenova H, Stepankova R, Hudcovic T, Tuckova L, Cukrowska B, Lodinova-Zadnikova R, Kozakova H, Rossmann P, Bartova J, Sokol D, Funda DP, Borovska D, Rehakova Z, Sinkora J, Hofman J, Drastich P, Kokesova A 2004 Commensal bacteria (normal microflora), mucosal immunity and chronic inflammatory and autoimmune diseases. *Immunol Lett* 93, 97. [PubMed: 15158604]
- Dixon DR, Reife RA, Cebra JJ, Darveau RP 2004 Commensal bacteria influence innate status within gingival tissues: a pilot study. *J Periodontol* 75, 1486. [PubMed: 15633325]
- Cugini C, Klepac-Ceraj V, Rackaityte E, Riggs JE, Davey ME 2013 *Porphyromonas gingivalis*: keeping the pathos out of the biont. *J Oral Microbiol* 5.
- Periasamy S, Kolenbrander PE 2009 *Aggregatibacter actinomycetemcomitans* builds mutualistic biofilm communities with *Fusobacterium nucleatum* and *Veillonella* species in saliva. *Infect Immun* 77, 3542. [PubMed: 19564387]
- Hajishengallis G, Lamont RJ 2012 Beyond the red complex and into more complexity: the polymicrobial synergy and dysbiosis (PSD) model of periodontal disease etiology. *Mol Oral Microbiol* 27, 409. [PubMed: 23134607]
- Hajishengallis G 2014 The inflammophilic character of the periodontitis-associated microbiota. *Mol Oral Microbiol*
- Ji S, Choi Y 2013 Innate immune response to oral bacteria and the immune evasive characteristics of periodontal pathogens. *J Periodontal Implant Sci* 43, 3. [PubMed: 23507986]
- Hans M, Hans VM 2011 Toll-like receptors and their dual role in periodontitis: a review. *J Oral Sci* 53, 263. [PubMed: 21959652]
- Walsh D, McCarthy J, O'Driscoll C, Melgar S 2013 Pattern recognition receptors--molecular orchestrators of inflammation in inflammatory bowel disease. *Cytokine Growth Factor Rev* 24, 91. [PubMed: 23102645]
- Kumar S, Ingle H, Prasad DV, Kumar H 2013 Recognition of bacterial infection by innate immune sensors. *Crit Rev Microbiol* 39, 229. [PubMed: 22866947]
- Jounai N, Kobiyama K, Takeshita F, Ishii KJ 2012 Recognition of damage-associated molecular patterns related to nucleic acids during inflammation and vaccination. *Front Cell Infect Microbiol* 2, 168. [PubMed: 23316484]
- Kumar H, Kawai T, Akira S 2011 Pathogen recognition by the innate immune system. *Int Rev Immunol* 30, 16. [PubMed: 21235323]
- Ebersole JL, Dawson DR 3rd, Morford LA, Peyyala R, Miller CS, Gonzalez OA 2013 Periodontal disease immunology: 'double indemnity' in protecting the host. *Periodontol* 2000 62, 163. [PubMed: 23574466]
- Benakanakere M, Kinane DF 2012 Innate cellular responses to the periodontal biofilm. *Front Oral Biol* 15, 41. [PubMed: 22142956]
- Bartold PM, Van Dyke TE 2013 Periodontitis: a host-mediated disruption of microbial homeostasis. Unlearning learned concepts. *Periodontol* 2000 62, 203. [PubMed: 23574467]
- Fatemi K, Radvar M, Rezaee A, Rafatpanah H, Azangoo khiavi H., Dadpour Y, Radvar N 2013 Comparison of relative TLR-2 and TLR-4 expression level of disease and healthy gingival tissue of smoking and non-smoking patients and periodontally healthy control patients. *Aust Dent J* 58, 315. [PubMed: 23981212]
- Di Benedetto A, Gigante I, Colucci S, Grano M 2013 Periodontal disease: linking the primary inflammation to bone loss. *Clin Dev Immunol* 2013, 503754. [PubMed: 23762091]
- Myneni SR, Settem RP, Sharma A 2013 Bacteria take control of tolls and T cells to destruct jaw bone. *Immunol Invest* 42, 519. [PubMed: 24004056]

- Promsudthi A, Poomsawat S, Limsricharoen W 2014 The role of Toll-like receptor 2 and 4 in gingival tissues of chronic periodontitis subjects with type 2 diabetes. *J Periodontol* 49, 346. [PubMed: 23844856]
- Lin J, Bi L, Yu X, Kawai T, Taubman MA, Shen B, Han X 2014 Porphyromonas gingivalis exacerbates ligature-induced, RANKL-dependent alveolar bone resorption via differential regulation of Toll-like receptor 2 (TLR2) and TLR4. *Infect Immun* 82, 4127. [PubMed: 25047844]
- Ghaderi H, Kiany F, Razmkhah M, Dadras S, Chenari N, Hosseini A, Younesi V, Ghaderi A 2014 mRNA expression of pattern recognition receptors and their signaling mediators in healthy and diseased gingival tissues. *J Indian Soc Periodontol* 18, 150. [PubMed: 24872620]
- Sahingur SE, Xia XJ, Voth SC, Yeudall WA, Gunsolley JC 2013 Increased nucleic Acid receptor expression in chronic periodontitis. *J Periodontol* 84, e48. [PubMed: 23646855]
- Benakanakere M, Abdolhosseini M, Hosur K, Finoti LS, Kinane DF 2015 TLR2 promoter hypermethylation creates innate immune dysbiosis. *J Dent Res* 94, 183. [PubMed: 25389002]
- Park E, Na HS, Song YR, Shin SY, Kim YM, Chung J 2014 Activation of NLRP3 and AIM2 inflammasomes by Porphyromonas gingivalis infection. *Infect Immun* 82, 112. [PubMed: 24126516]
- Ebersole JL, Steffen MJ, Gonzalez-Martinez J, Novak MJ 2008 Effects of age and oral disease on systemic inflammatory and immune parameters in nonhuman primates. *Clin Vaccine Immunol* 15, 1067. [PubMed: 18448617]
- Meka A, Bakthavatchalu V, Sathishkumar S, Lopez MC, Verma RK, Wallet SM, Bhattacharyya I, Boyce BF, Handfield M, Lamont RJ, Baker HV, Ebersole JL, Kesavalu L Porphyromonas gingivalis infection-induced tissue and bone transcriptional profiles. *Mol Oral Microbiol* 25, 61. [PubMed: 20331794]
- Gonzalez OA, Stromberg AJ, Huggins PM, Gonzalez-Martinez J, Novak MJ, Ebersole JL 2011 Apoptotic genes are differentially expressed in aged gingival tissue. *J Dent Res* 90, 880. [PubMed: 21471327]
- Gonzalez OA, Novak MJ, Kirakodu S, Orraca L, Chen KC, Stromberg A, Gonzalez-Martinez J, Ebersole JL 2014 Comparative analysis of gingival tissue antigen presentation pathways in ageing and periodontitis. *J Clin Periodontol* 41, 327. [PubMed: 24304139]
- Smith MA, Braswell LD, Collins JG, Boyd DL, Jeffcoat MK, Reddy M, Li KL, Wilensky S, Vogel R, Alfano M, et al. 1993 Changes in inflammatory mediators in experimental periodontitis in the rhesus monkey. *Infect Immun* 61, 1453. [PubMed: 8384162]
- Moritz AJ, Cappelli D, Lantz MS, Holt SC, Ebersole JL 1998 Immunization with Porphyromonas gingivalis cysteine protease: effects on experimental gingivitis and ligature-induced periodontitis in Macaca fascicularis. *J Periodontol* 69, 686. [PubMed: 9660338]
- Branch-Mays GL, Dawson DR, Gunsolley JC, Reynolds MA, Ebersole JL, Novak KF, Mattison JA, Ingram DK, Novak MJ 2008 The effects of a calorie-reduced diet on periodontal inflammation and disease in a non-human primate model. *J Periodontol* 79, 1184. [PubMed: 18597600]
- Qian C, Cao X 2013 Regulation of Toll-like receptor signaling pathways in innate immune responses. *Ann N Y Acad Sci* 1283, 67. [PubMed: 23163321]
- Mogensen TH 2009 Pathogen recognition and inflammatory signaling in innate immune defenses. *Clin Microbiol Rev* 22, 240. [PubMed: 19366914]
- Atkinson TJ 2008 Toll-like receptors, transduction-effector pathways, and disease diversity: evidence of an immunobiological paradigm explaining all human illness? *Int Rev Immunol* 27, 255. [PubMed: 18574739]
- Sukhithasri V, Nisha N, Biswas L, Anil Kumar V., Biswas R 2013 Innate immune recognition of microbial cell wall components and microbial strategies to evade such recognitions. *Microbiol Res* 168, 396. [PubMed: 23578963]
- Gonzalez O, Tobia C, Ebersole J, Novak MJ 2012 Caloric restriction and chronic inflammatory diseases. *Oral Dis* 18, 16. [PubMed: 21749581]
- Oz HS, Puleo DA 2011 Animal models for periodontal disease. *J Biomed Biotechnol* 2011, 754857. [PubMed: 21331345]
- Struillou X, Boutigny H, Soueidan A, Layrolle P 2010 Experimental animal models in periodontology: a review. *Open Dent J* 4, 37. [PubMed: 20556202]

- van Bergenhenegouwen J, Plantinga TS, Joosten LA, Netea MG, Folkerts G, Kraneveld AD, Garssen J, Vos AP 2013 TLR2 & Co: a critical analysis of the complex interactions between TLR2 and coreceptors. *J Leukoc Biol* 94, 885. [PubMed: 23990624]
- Wu S, Jiang ZY, Sun YF, Yu B, Chen J, Dai CQ, Wu XL, Tang XL, Chen XY 2013 Microbiota regulates the TLR7 signaling pathway against respiratory tract influenza A virus infection. *Curr Microbiol* 67, 414. [PubMed: 23677145]
- Szabo A, Rajnavolgyi E 2013 Collaboration of Toll-like and RIG-I-like receptors in human dendritic cells: tRIGgering antiviral innate immune responses. *Am J Clin Exp Immunol* 2, 195. [PubMed: 24179728]
- Mansson Kvarnhammar A., Tengroth L, Adner M, Cardell LO 2013 Innate immune receptors in human airway smooth muscle cells: activation by TLR1/2, TLR3, TLR4, TLR7 and NOD1 agonists. *PLoS One* 8, e68701. [PubMed: 23861935]
- Kaiko GE, Loh Z, Spann K, Lynch JP, Lalwani A, Zheng Z, Davidson S, Uematsu S, Akira S, Hayball J, Diener KR, Baines KJ, Simpson JL, Foster PS, Phipps S 2013 Toll-like receptor 7 gene deficiency and early-life Pneumovirus infection interact to predispose toward the development of asthma-like pathology in mice. *J Allergy Clin Immunol* 131, 1331. [PubMed: 23561801]
- Eick S, Reissmann A, Rodel J, Schmidt KH, Pfister W 2006 Porphyromonas gingivalis survives within KB cells and modulates inflammatory response. *Oral Microbiol Immunol* 21, 231. [PubMed: 16842507]
- Colombo AV, Silva CM, Haffajee A, Colombo AP 2006 Identification of oral bacteria associated with crevicular epithelial cells from chronic periodontitis lesions. *J Med Microbiol* 55, 609. [PubMed: 16585650]
- Vitkov L, Krautgartner WD, Hannig M 2005 Bacterial internalization in periodontitis. *Oral Microbiol Immunol* 20, 317. [PubMed: 16101968]
- Atanasova KR, Yilmaz O 2014 Looking in the Porphyromonas gingivalis cabinet of curiosities: the microbium, the host and cancer association. *Mol Oral Microbiol* 29, 55. [PubMed: 24506890]
- Park Y, Yilmaz O, Jung IY, Lamont RJ 2004 Identification of Porphyromonas gingivalis genes specifically expressed in human gingival epithelial cells by using differential display reverse transcription-PCR. *Infect Immun* 72, 3752. [PubMed: 15213115]
- Litvack ML, Palaniyar N 2010 Review: Soluble innate immune pattern-recognition proteins for clearing dying cells and cellular components: implications on exacerbating or resolving inflammation. *Innate Immun* 16, 191. [PubMed: 20529971]
- Jin MS, Lee JO 2008 Structures of the toll-like receptor family and its ligand complexes. *Immunity* 29, 182. [PubMed: 18701082]
- Pussinen PJ, Paju S, Mantyla P, Sorsa T 2007 Serum microbial- and host-derived markers of periodontal diseases: a review. *Curr Med Chem* 14, 2402. [PubMed: 17896988]
- Jin L, Darveau RP 2001 Soluble CD14 levels in gingival crevicular fluid of subjects with untreated adult periodontitis. *J Periodontol* 72, 634.
- Gong Y, Bi W, Cao L, Yang Y, Chen J, Yu Y 2013 Association of CD14–260 polymorphisms, red-complex periodontopathogens and gingival crevicular fluid cytokine levels with cyclosporine A-induced gingival overgrowth in renal transplant patients. *J Periodontal Res* 48, 203. [PubMed: 22934794]
- Yan H, Ohno N, Tsuji NM 2013 The role of C-type lectin receptors in immune homeostasis. *Int Immunopharmacol* 16, 353. [PubMed: 23623943]
- Yamasaki S, Matsumoto M, Takeuchi O, Matsuzawa T, Ishikawa E, Sakuma M, Tateno H, Uno J, Hirabayashi J, Mikami Y, Takeda K, Akira S, Saito T 2009 C-type lectin Mincle is an activating receptor for pathogenic fungus, Malassezia. *Proc Natl Acad Sci U S A* 106, 1897. [PubMed: 19171887]
- Miyake Y, Toyonaga K, Mori D, Kakuta S, Hoshino Y, Oyamada A, Yamada H, Ono K, Suyama M, Iwakura Y, Yoshikai Y, Yamasaki S 2013 C-type lectin MCL is an FcRgamma-coupled receptor that mediates the adjuvanticity of mycobacterial cord factor. *Immunity* 38, 1050. [PubMed: 23602766]

- Sharma A, Steichen AL, Jondle CN, Mishra BB, Sharma J 2014 Protective role of Mincle in bacterial pneumonia by regulation of neutrophil mediated phagocytosis and extracellular trap formation. *J Infect Dis* 209, 1837. [PubMed: 24353272]
- Khoo US, Chan KY, Chan VS, Lin CL 2008 DC-SIGN and L-SIGN: the SIGNs for infection. *J Mol Med (Berl)* 86, 861. [PubMed: 18458800]
- Rydz N, Swystun LL, Notley C, Paterson AD, Riches JJ, Sponagle K, Boonyawat B, Montgomery RR, James PD, Lillcrap D 2013 The C-type lectin receptor CLEC4M binds, internalizes, and clears von Willebrand factor and contributes to the variation in plasma von Willebrand factor levels. *Blood* 121, 5228. [PubMed: 23529928]
- Rao R, Graffeo CS, Gulati R, Jamal M, Narayan S, Zambirinis CP, Barilla R, Deutsch M, Greco SH, Ochi A, Tomkoter L, Blobstein R, Avanzi A, Tippens DM, Gelbstein Y, Van Heerden E, Miller G 2014 Interleukin 17-Producing gammadeltaT Cells Promote Hepatic Regeneration in Mice. *Gastroenterology* 147, 473. [PubMed: 24801349]
- Plato A, Willment JA, Brown GD 2013 C-type lectin-like receptors of the dectin-1 cluster: ligands and signaling pathways. *Int Rev Immunol* 32, 134. [PubMed: 23570314]
- Gringhuis SI, Kaptein TM, Wevers BA, Theelen B, van der Vlist M, Boekhout T, Geijtenbeek TB 2012 Dectin-1 is an extracellular pathogen sensor for the induction and processing of IL-1beta via a noncanonical caspase-8 inflammasome. *Nat Immunol* 13, 246. [PubMed: 22267217]
- Liang S, Hosur KB, Domon H, Hajishengallis G 2010 Periodontal inflammation and bone loss in aged mice. *J Periodontol Res* 45, 574. [PubMed: 20337897]
- Tamai R, Sugamata M, Kiyoura Y 2011 *Candida albicans* enhances invasion of human gingival epithelial cells and gingival fibroblasts by *Porphyromonas gingivalis*. *Microb Pathog* 51, 250. [PubMed: 21742026]
- Carvalho A, Giovannini G, De Luca A, D'Angelo C, Casagrande A, Iannitti RG, Ricci G, Cunha C, Romani L 2012 Dectin-1 isoforms contribute to distinct Th1/Th17 cell activation in mucosal candidiasis. *Cell Mol Immunol* 9, 276. [PubMed: 22543832]
- Rasmussen SB, Reinert LS, Paludan SR 2009 Innate recognition of intracellular pathogens: detection and activation of the first line of defense. *APMIS* 117, 323. [PubMed: 19400860]
- Wen H, Miao EA, Ting JP 2013 Mechanisms of NOD-like receptor-associated inflammasome activation. *Immunity* 39, 432. [PubMed: 24054327]
- Zelkha SA, Freilich RW, Amar S 2010 Periodontal innate immune mechanisms relevant to atherosclerosis and obesity. *Periodontol* 2000 54, 207. [PubMed: 20712641]
- Jiao Y, Darzi Y, Tawaratsumida K, Marchesan JT, Hasegawa M, Moon H, Chen GY, Nunez G, Giannobile WV, Raes J, Inohara N 2013 Induction of bone loss by pathobiont-mediated Nod1 signaling in the oral cavity. *Cell Host Microbe* 13, 595. [PubMed: 23684310]
- Yuan H, Zelka S, Burkatovskaya M, Gupte R, Leeman SE, Amar S 2013 Pivotal role of NOD2 in inflammatory processes affecting atherosclerosis and periodontal bone loss. *Proc Natl Acad Sci U S A* 110, E5059. [PubMed: 24324141]
- Kofoed EM, Vance RE 2012 NAIPs: building an innate immune barrier against bacterial pathogens. NAIPs function as sensors that initiate innate immunity by detection of bacterial proteins in the host cell cytosol. *Bioessays* 34, 589. [PubMed: 22513803]
- Beug ST, Cheung HH, LaCasse EC, Korneluk RG 2012 Modulation of immune signalling by inhibitors of apoptosis. *Trends Immunol* 33, 535. [PubMed: 22836014]
- Gong YN, Shao F 2012 Sensing bacterial infections by NAIP receptors in NLRC4 inflammasome activation. *Protein Cell* 3, 98. [PubMed: 22426978]
- Gonzalez OA, John Novak M., Kirakodu S, Stromberg AJ, Shen S, Orraca L, Gonzalez-Martinez J, Ebersole JL 2013 Effects of aging on apoptosis gene expression in oral mucosal tissues. *Apoptosis* 18, 249. [PubMed: 23334583]
- Lupfer C, Kanneganti TD 2013 Unsolved Mysteries in NLR Biology. *Front Immunol* 4, 285. [PubMed: 24062750]
- Rathinam VA, Vanaja SK, Fitzgerald KA 2012 Regulation of inflammasome signaling. *Nature immunology* 13, 333. [PubMed: 22430786]
- Martinon F, Mayor A, Tschopp J 2009 The inflammasomes: guardians of the body. *Annu Rev Immunol* 27, 229. [PubMed: 19302040]

- Cai S, Batra S, Wakamatsu N, Pacher P, Jeyaseelan S 2012 NLRC4 inflammasome-mediated production of IL-1 β modulates mucosal immunity in the lung against gram-negative bacterial infection. *J Immunol* 188, 5623. [PubMed: 22547706]
- Nordlander S, Pott J, Maloy KJ 2014 NLRC4 expression in intestinal epithelial cells mediates protection against an enteric pathogen. *Mucosal Immunol* 7, 775. [PubMed: 24280936]
- Bostanci N, Emingil G, Saygan B, Turkoglu O, Atilla G, Curtis MA, Belibasakis GN 2009 Expression and regulation of the NALP3 inflammasome complex in periodontal diseases. *Clin Exp Immunol* 157, 415. [PubMed: 19664151]
- Belibasakis GN, Guggenheim B, Bostanci N 2013 Down-regulation of NLRP3 inflammasome in gingival fibroblasts by subgingival biofilms: involvement of *Porphyromonas gingivalis*. *Innate Immun* 19, 3. [PubMed: 22522430]
- Vilaysane A, Muruve DA 2009 The innate immune response to DNA. *Semin Immunol* 21, 208. [PubMed: 19497760]
- Fang R, Hara H, Sakai S, Hernandez-Cuellar E, Mitsuyama M, Kawamura I, Tsuchiya K 2014 Type I interferon signaling regulates activation of the absent in melanoma 2 inflammasome during *Streptococcus pneumoniae* infection. *Infect Immun* 82, 2310. [PubMed: 24643540]
- Philpott DJ, Sorbara MT, Robertson SJ, Croitoru K, Girardin SE 2014 NOD proteins: regulators of inflammation in health and disease. *Nat Rev Immunol* 14, 9. [PubMed: 24336102]
- Yanai H, Savitsky D, Tamura T, Taniguchi T 2009 Regulation of the cytosolic DNA-sensing system in innate immunity: a current view. *Curr Opin Immunol* 21, 17. [PubMed: 19362700]
- Slots J 2010 Human viruses in periodontitis. *Periodontol* 2000 53, 89. [PubMed: 20403107]
- Canton J, Neculai D, Grinstein S 2013 Scavenger receptors in homeostasis and immunity. *Nat Rev Immunol* 13, 621. [PubMed: 23928573]
- Taylor ME, Drickamer K 2007 Paradigms for glycan-binding receptors in cell adhesion. *Curr Opin Cell Biol* 19, 572. [PubMed: 17942297]
- Endo Y, Matsushita M, Fujita T 2011 The role of ficolins in the lectin pathway of innate immunity. *Int J Biochem Cell Biol* 43, 705. [PubMed: 21315829]
- Thomsen T, Schlosser A, Holmskov U, Sorensen GL 2011 Ficolins and FIBCD1: soluble and membrane bound pattern recognition molecules with acetyl group selectivity. *Mol Immunol* 48, 369. [PubMed: 21071088]
- Cieslik P, Hrycek A 2012 Long pentraxin 3 (PTX3) in the light of its structure, mechanism of action and clinical implications. *Autoimmunity* 45, 119. [PubMed: 21988562]
- Deban L, Bottazzi B, Garlanda C, de la Torre YM, Mantovani A 2009 Pentraxins: multifunctional proteins at the interface of innate immunity and inflammation. *Biofactors* 35, 138. [PubMed: 19449441]
- Pradeep AR, Kathariya R, Raghavendra NM, Sharma A 2011 Levels of pentraxin-3 in gingival crevicular fluid and plasma in periodontal health and disease. *J Periodontol* 82, 734. [PubMed: 21080790]
- Lakshmanan R, Jayakumar ND, Sankari M, Padmalatha O, Varghese S 2013 Estimation of Pentraxin-3 Levels in the Gingival Tissues of Chronic And Aggressive Periodontitis Participants-An in Vivo Study. *J Periodontol*
- Gumus P, Nizam N, Nalbantsoy A, Ozcaka O, Buduneli N 2014 Saliva and serum levels of pentraxin-3 and interleukin-1 β in generalized aggressive or chronic periodontitis. *J Periodontol* 85, e40. [PubMed: 23952075]

Figure 1:

Distribution of gene expression of PRRs in healthy and periodontitis gingival tissues. The bars denote the mean expression levels in healthy (n=13), and periodontitis (n=11) samples for Surface PRR (A), Intracellular PRR (B), and Soluble PRR (C). Significant differences are denoted by p-values.

Figure 2:
Correlation of PRR gene expression in periodontitis gingival tissues with (A) Bleeding on Probing (BOP) and (B) Pocket Depth (PD). Red dashed lines denote the r-value required for reaching p<0.01 for the 24 animals. Genes reaching statistical significance are identified.

Figure 3: Clinical features of (A) Bleeding on Probing and (B) Pocket Depth in longitudinal ligature-induced periodontitis (n=18). Points denote mean level for the group at each sampling point. Vertical lines signify 1 SD. The asterisk (*) denotes significantly different from baseline at least at $p < 0.01$ and the hashtag (#) signifies a significant difference from peak levels at 1 mo. at least at $p < 0.01$.

Figure 4:

PRRs whose gene expression changed significantly from baseline during initiation, progression and resolution of ligature-induced periodontitis. Points denote group means and vertical brackets enclose 1 SD. The asterisk (*; $p < 0.01$) and hashtag (#; $p < 0.05$) signify difference from baseline.

Table 1:

Gene expression targets for microbial sensing molecules

Gene ID	Gene Title
<i>Surface Cell Associated PRRs</i>	
CD14	Binds to monomeric lipopolysaccharide and delivers it to the MD-2/TLR4 complex; also soluble molecule
CD209 (DC-SIGN)	Surface of immature dendritic cells (DCs) and involved in initiation of primary immune response
CLEC7A	C-Type Lectin Domain Family 7, Member A, beta-1,3-linked and beta-1,6-linked glucans from fungi
CLEC4E	C-Type Lectin domain family 4, member E, mycobacteria is via trehalose 6,6'-dimycolate; SAP130, a nuclear protein that is released by dead or dying cells
CLEC4M (L-SIGN)	C-Type Lectin Domain Family 4, Member M, parasites to viruses
CLEC6A (Dectin-2)	C-Type Lectin Domain Family 6, Member A, alpha-mannans on <i>C.albicans</i> hyphae
COLEC12	Collectin Sub-Family Member 12, Scavenger receptor for phagocytosis of Gram-positive, Gram-negative bacteria and yeast
FCN2	Ficolin (Collagen/Fibrinogen Domain Containing Lectin) 2 (Hucolin), Complement-activating lectin, phagocytosis of <i>S.typhimurium</i> by neutrophils
MARCO	Macrophage Receptor With Collagenous Structure, class A scavenger receptor for both Gram-negative and Gram-positive bacteria
MRC1	Mannose Receptor, C Type 1, bind high-mannose structures on the surface of potentially pathogenic viruses, bacteria, and fungi for phagocytosis
TLR1	Toll-like receptor 1, diacylated and triacylated lipopeptides
TLR2	Toll-like receptor 2, bacterial lipoproteins and other microbial cell wall components
TLR4	Toll-like receptor 4, bacterial lipopolysaccharide (LPS)
TLR5	Toll-like receptor 5, bacterial flagellins
TLR6	Toll-like receptor 6, Gram-positive bacteria and fungi
<i>Intracellular Associated PRRs</i>	
AIM2	Cytosolic double-stranded DNA
ARHGEF2	Rho/Rac Guanine Nucleotide Exchange Factor (GEF) 2, intracellular sensing system along with NOD1 for the detection of microbial effectors during cell invasion
IFIH1 (MDA5)	Interferon induced with helicase C domain 1, RIG-1-like receptor family, viral sensor
LGP2 (DEXH58)	DEXH (Asp-Glu-X-His) Box Polypeptide 58, RIG-1-like receptor family, viral sensor
NAIP (BIRC1)	Baculoviral IAP repeat-containing protein 1, effects apoptosis
NLRC4 (IPAF)	NLR family CARD domain-containing protein 4, inflammasome
NOD1	Nucleotide-binding oligomerization domain 1, intracellular bacterial lipopolysaccharides (LPS), senses peptidoglycan (PGN)-derived muropeptides
NOD2	Nucleotide-binding oligomerization domain 2, intracellular bacterial lipopolysaccharides (LPS) by recognizing the muramyl dipeptide
RIG-1 (DDX58)	Retinoic acid-inducible gene 1, RIG-1-like receptor family, viral sensor
TLR3	Toll-like receptor 3, nucleotide-sensing for double-stranded RNA
TLR7	Toll-like receptor 7, nucleotide-sensing for single-stranded RNA
TLR8	Toll-like receptor 8, G-rich oligonucleotides
TLR9	Toll-like receptor 9, nucleotide-sensing for unmethylated cytidine-phosphate-guanosine (CpG) dinucleotides
ZBP1/DAI	Z-DNA Binding Protein 1, cytoplasmic sensor binds to foreign DNA and induces type-I interferon
<i>Soluble PRRs</i>	

Gene ID	Gene Title
CRP (PTX1)	C-reactive protein, promotes agglutination, bacterial capsular swelling, phagocytosis and complement fixation
FCN1	Ficolin (Collagen/Fibrinogen Domain Containing) 1, Complement-activating lectin, 9-O-acetylated 2–6-linked sialic acid derivatives and to various glycans
MBL2	Mannose-Binding Lectin (Protein C) 2, Soluble, soluble mannose-binding lectin or mannose-binding protein found in serum
PTX3	Pentraxin 3, Long, mediating agglutination, complement activation, and opsonization
PTX4	Pentraxin 4, Long, mediating agglutination, complement activation, and opsonization
SAA1	Serum Amyloid A1

Table 2:

Correlation of gene expression levels with level of mRNA at baseline. R-values >0.4200 or <-0.4200 were considered significant at $p < 0.05$.

	PERIODONTITIS			
Gene ID	2 wk	1 mo	3 mo	5 mo
Cell surface				
TLR2			0.5671	-0.5935
TLR3		0.6495	0.6574	
TLR4			0.4444	
CD209		0.4568	-0.4383	
CLEC4E			0.7392	
CLEC6A		0.5970		
FCN2				0.5267
MARCO	0.4879			
Intracellular				
AIM2		0.5277		
NLRC4			0.4468	-0.4258
TLR7		0.5482		
ZBP1/DAI		0.4391		
Soluble				
CD14		0.4313	0.6575	
LBP			0.4452	
TOTAL +	1	7	7	1
TOTAL -	0	0	1	2

Table 3:

Correlation of gene expression levels with clinical parameters of bleeding on probing (BOP) and pocket depth (PD) at different states of disease. Values >0.4200 or <-0.4200 for $n=18$ were considered significant at $p<0.05$ as highlighted in boldface type. Cells with no values showed no significant correlation related to BOP or PD.

Bleeding on Probing				Pocket Depth		
Gene ID	B	Disease	Resolution	B	Disease	Resolution
CD14				0.4655	-0.0258	-0.1814
CD209	-0.2233	-0.1697	-0.5570			
CLEC4E	0.2321	0.3004	-0.5606	0.1828	0.0956	-0.6819
CLEC6A				0.5948	0.1988	0.1613
COLEC12	0.2213	0.1251	-0.6052	-0.3121	-0.1615	-0.7203
CRP				-0.0399	-0.1067	-0.6471
FCN1	0.3966	0.2063	-0.6687			
FCN2	0.3174	0.0957	0.4591	-0.5645	0.2568	0.7122
IFIH1				0.4573	0.3536	0.4213
MBL2				-0.5908	-0.0192	-0.3087
NAIP	0.5023	0.3027	-0.6901	0.3658	0.1880	-0.5163
NOD1	-0.3031	0.1701	0.5815	-0.1197	0.0707	0.6773
NOD2	-0.2192	-0.1159	-0.4445	0.5057	-0.3414	0.2027
PTX3	-0.2574	0.0941	-0.6679	-0.3840	-0.0495	-0.6538
PTX4				-0.4941	-0.1414	0.6646
TLR3	-0.1057	0.1523	0.4695			
TLR4	0.5312	0.3030	-0.6913	0.2847	0.0608	-0.6144
TLR5	-0.1089	-0.0019	0.5675			
TLR6	0.5960	0.3712	-0.2864			
TLR7				0.4538	0.0949	-0.0439
TLR8	0.2647	0.3638	-0.8853	0.4059	0.1743	-0.4872
TLR9	0.2013	0.3023	0.7415			
ZBP1	0.0816	0.2028	0.8422	0.5261	-0.0679	0.2559

Table 4:

Comparison of gene expression profiles using qPCR and microarray analyses. Values represent fold-difference compared to Healthy tissue message levels assigned a value of 1.0 from cross-sectional samples.

Gene ID	X-Sectional Periodontitis (Fold vs. Health)	Longitudinal (Fold vs. Baseline)
SAA1		
qPCR	3.69 ± 0.25	
GeneChip	2.57 ± 0.81	
CD14		
qPCR	1.12 ± 0.48	2.16 ± 0.13 *
GeneChip	1.54 ± 0.30	1.76 ± 0.12
NAIP		
qPCR	1.79 ± 0.06	
GeneChip	1.37 ± 0.40	
TLR2		
qPCR	0.97 ± 0.17	2.80 ± 0.17 *
GeneChip	1.56 ± 0.40	2.52 ± 0.30
ZBP1/DAI		
qPCR	1.41 ± 0.54	2.68 ± 0.45 #
GeneChip	2.25 ± 0.84	1.58 ± 0.14
CLEC7A		
qPCR		0.90 ± 0.11 *
GeneChip		0.84 ± 0.07
NOD2		
qPCR		0.62 ± 0.04 *
GeneChip		0.78 ± 0.06

* denotes samples from 2 weeks compared with baseline

denotes samples from 3 months compared with baseline