Published in final edited form as:

J Am Chem Soc. 2019 March 20; 141(11): 4526–4530. doi:10.1021/jacs.8b13042.

Designing a Green Fluorogenic Protease Reporter by Flipping a Beta Strand of GFP for Imaging Apoptosis in Animals

Qiang Zhang^{¶,§}, Antonino Schepis^{§,‡}, Hai Huang^{§,‡}, Junjiao Yang^{¶,§,‡}, Wen Ma^{¶,‡}, Joaquim Torra^{¶,§,†}, Shao-Qing Zhang[⊥], Lina Yang[#], Haifan Wu^{¶,§}, Santi Nonell[†], Zhiqiang Dong[#], Thomas B. Kornberg[§], Shaun R. Coughlin[§], and Xiaokun Shu^{*,¶,§}

[¶]Department of Pharmaceutical Chemistry, University of California–San Francisco, San Francisco, California 94158, United States

[§]Cardiovascular Research Institute, University of California–San Francisco, San Francisco, California 94158, United States

[¶]Department of Chemistry and Biochemistry, University of California San Diego, La Jolla, California 92093, United States

[†]Institut Quimic de Sarria, Universitat Ramon Llull, Via Augusta 390, 08017 Barcelona, Spain

[⊥]Department of Neurology, Harvard Medical School and MassGeneral Institute for Neurodegenerative Disease, Massachusetts General Hospital, Charlestown, Massachusetts 02129, United States

[#]Bio-Medical Center, College of Life Science and Technology, Huazhong Agricultural University, Wuhan, Hubei 430070, China

Abstract

A family of proteases called caspases mediate apoptosis signaling in animals. We report a GFP-based fluorogenic protease reporter, dubbed “FlipGFP”, by flipping a beta strand of the GFP. Upon protease activation and cleavage, the beta strand is restored, leading to reconstitution of the GFP and fluorescence. FlipGFP-based TEV protease reporter achieves 100-fold fluorescence change. A FlipGFP-based executioner caspase reporter visualized apoptosis in live zebrafish embryos with spatiotemporal resolution. FlipGFP also visualized apoptotic cells in the midgut of *Drosophila*. Thus, the FlipGFP-based caspase reporter will be useful for monitoring apoptosis during animal development and for designing reporters of proteases beyond caspases. The design strategy can be further applied to a red fluorescent protein for engineering a red fluorogenic protease reporter.

*Corresponding Author xiaokun.shu@ucsf.edu.

[‡]These authors contributed equally.

The authors declare no competing financial interest.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.8b13042.

Movies S1–10 (ZIP)

DNA constructs, protein purification and characterization, imaging, characterization of FlipGFP, engineering a red fluorogenic protease reporter (PDF)

Proteases play fundamental roles in many signaling pathways. One family of protease, caspase, executes cell apoptosis. The apoptosis signaling pathway is composed of extrinsic and intrinsic pathways that are activated by stimuli such as DNA damage, oncogenes, and developmental cues.^{1,2} Both pathways lead to activation of the executioner caspases that cleave thousands of substrates, resulting in cell apoptosis.³ Activity reporters of executioner caspases thus visualize apoptosis signaling. Furthermore, mechanistic understanding of cell and developmental biology requires investigation of cell signaling in living animals. To image spatiotemporal dynamics of apoptosis in animals especially developing animals, it is ideal to use genetically encoded fluorescent protease reporters. However, fluorescence imaging of living animals is challenging because of tissue autofluorescence, cell heterogeneity and rapid shape and position changes. Thus, to enable in vivo imaging, protease reporters should have large dynamic range (i.e., large fluorescence increase upon protease activation) and high brightness (i.e., high quantum yield). Over the past few years, to overcome limitations of FRET-based executioner caspase reporters,^{4–7} many different types of reporters have been developed.^{8–13} For example, the bipartite GFP was previously engineered into a light-driven protease sensor.¹¹ Here we report a new type of rationally designed fluorogenic protease reporter based on GFP with both large dynamic range and high brightness. And we demonstrate its application in imaging apoptosis in zebrafish embryos.

To design a GFP-based fluorogenic protease reporter with large dynamic range, we modified the self-assembling split GFP so that its self-assembly is regulated by protease activity. GFP contains 11 beta strands (β -strands) and a central α -helix (Movie S1),^{14,15} which can be split into three parts. One part contains nine β -strands and the central α -helix (β 1–9); a second part contains 10th β strand (β 10); and the third part contains 11th β strand (β 11).¹⁶ β 1–9 contains three amino acids that form the chromophore;^{17,18} whereas β 11 contains the highly conserved Glu222 that catalyzes chromophore maturation.¹⁹ It has been shown that when β 10 and β 11 are linked together or are in close proximity, they rapidly bind to β 1–9, and green fluorescence develops within several tens of minutes.²⁰ This has been used to detect protein-protein interactions,²⁰ including interaction between G protein coupled receptor (GPCR) and beta-arrestin upon activation and phosphorylation of GPCR.²¹

GFP structure suggests a compact fit of β 1–9 and β 10–11 (Figure 1a), which presumably provides structural basis of high binding affinity between them (Movie S2). On this basis, we sought to prevent β 10–11 binding to β 1–9 until protease activation. To achieve this, we redesigned β 10–11. GFP structure shows that β 10–11 forms an antiparallel β -strands, which fit well with the rest of the structure (Figure 1a). We hypothesized that if β 10–11 was redesigned so that it forms a parallel structure, then it would no longer fit β 1–9. To flip β 11, we utilized heterodimerizing coiled coils E5 and K5: we linked β 10/ β 11 with E5; β 11 was followed by K5; the protease cleavage sequence was inserted between β 11 and K5 (Figure 1b, c).

Based on this design, E5/K5 heterodimer “flips” β 11 and prevents self-assembly of the split GFP (Figure 1b, Figure S1). Upon protease cleavage, β 11 flips back, forming an antiparallel structure with β 10, which enables self-assembly with β 1–9 and leads to fluorescence increase. We named this reporter FlipGFP. As a proof of concept, we designed a TEV

protease reporter by inserting TEV cleavage sequence into FlipGFP, which achieved 77-fold fluorescence increase upon TEV cleavage in HEK293 cells (Figure 1d, linker length = 5 amino acids).

Because efficient assembly between cleaved $\beta 10$ –E5– $\beta 11$ and $\beta 1$ –9 is likely dependent on distance between $\beta 10$ and E5 (Figure 1b), we increased length of the linker between $\beta 10$ and E5 from 5 amino acid (aa) to 10, 15, 20, 30 aa. Characterization of the FlipGFP variants in HEK293 cells indicated that the FlipGFP with 10aa linker showed the largest fluorescence increase (106-fold) upon TEV cleavage and the highest brightness (Figure 1d,e). On the other hand, when the linker was 30aa, the fluorescence increase was narrowed to 26-fold. Interestingly, in the inactive state (i.e., no TEV protease), this FlipGFP with 30aa linker was as dim as the FlipGFP with 5aa linker. But in the active state (i.e., cleaved by TEV protease), the FlipGFP with 30aa linker was much dimmer than the FlipGFP with 5aa linker. This suggests that while elongation of the linker length from 5aa to 10aa increases the assembly efficiency of the split GFP, further elongation of the linker to 30aa decreases reconstitution efficiency. We thus decided to use 10aa linker length for FlipGFP from now on. Fluorescence quantum yield of reassembled FlipGFP was determined to be 0.66 ± 0.03 .

We next measured kinetics of FlipGFP-based protease reporter to determine its temporal resolution. First, we separately purified two parts of FlipGFP in the presence of TEV protease. Then we mixed the cleaved FlipGFP and monitored green fluorescence over time. The green fluorescence increased over time with time to half-maximal fluorescence ($T_{1/2}$) of 43 ± 6 min (Figure 1f,g), which is similar to the previously reported kinetics of the split GFP self-assembly.^{16,20} The proposed working mechanism of FlipGFP was supported by circular dichroism spectroscopy and molecular dynamics simulation (Figures S1–3, Movies S3–4, SI). Lastly, to obtain large fluorescence change, the expression level of FlipGFP needs to be titrated in cells (Figure S4).

To design a FlipGFP-based caspase reporter for imaging apoptosis, we replaced the TEV cleavage sequence with the consensus cleavage sequence (DEVD) of caspase-3 in FlipGFP (Figure 2a). The well characterized signaling pathways of apoptosis are composed of extrinsic and intrinsic pathways, both of which lead to activation of the executioner caspases that include caspase-3.³ And the conserved cleavage sequence (DEVD) of caspase-3 has been used in the previous caspase reporters.^{7,9,10,12,22,23}

We expressed FlipGFP in the HeLa cells. Upon addition of staurosporine, which induces apoptosis by activating caspase 3,²⁴ FlipGFP's green fluorescence increased significantly in cells treated with staurosporine (Figure 2b, Movie S5), whereas the fluorescence signal of FlipGFP has no or little change in cells without treatment with staurosporine (Movie S6). And the onset of apoptosis in single cells as detected by FlipGFP varied around 2 to 5 h after addition of staurosporine (Figure 2c, Movies S7–8, Figure S5), consistent with previous results.²⁵ Activation of caspase-3 in FlipGFP fluorescing cells was confirmed using a commercially available small molecule active-site modifier for far-red fluorescent imaging of caspase-3 activity (Figure 2d).

To determine whether the fluorescence signal and the temporal resolution of FlipGFP are sufficient for imaging apoptosis in animals, we expressed the FlipGFP-based caspase reporter in zebrafish. Zebrafish was chosen based on optical transparency and its importance as a model organism among vertebrate animals.²⁶ To demonstrate that FlipGFP could be used to visualize spatiotemporal dynamics of apoptosis during embryo development, we injected mRNA encoding two parts of FlipGFP caspase reporter linked with a T2A site and mCherry into the embryo at the one-cell stage. T2A is a “self-cleaving” peptide used in coexpression of multiple genes.²⁷

Time-lapse imaging of the head was started at 24 h postfertilization (hpf) (Figure 2e). At this time, little green fluorescence was observed (Figure 2e, Movie S9). Ten hours later, green fluorescence was detected in a couple of cells. After 15–20 h, more cells showed strong green fluorescence. By contrast, mCherry fluorescence was weak and decreased over time, presumably because its mRNA injected at the one-cell stage was degraded over time and/or diluted as development progressed. Green, presumed apoptotic cells were distributed around the developing forebrain and the retina. In particular, time-lapse imaging first detected green fluorescence on the rostral aspect of the forebrain. The number of green cells increased over time and they appeared to form clusters. These cells appeared to have neuron-like long extensions or processes (Figure 2e). We also imaged at 15 μ m deeper layer of the brain, which revealed similar distribution of the apoptotic cells (Figure 2e, right). Furthermore, z-section imaging indicated that many of them appeared to be located on the outer layer of the forebrain and the retina (Movie S10). The activity of caspase-3 was confirmed using antibody staining against cleaved caspase-3 (Figure 2f, Figure S6). The spatial pattern of FlipGFP fluorescence we observed is consistent with a previous study based on TUNEL staining that showed apoptotic cells in the developing brain.²⁸

After demonstration of FlipGFP in zebrafish, we decided to show whether it can also be used in detecting apoptosis in another important model organism *Drosophila*. Because of the genetic tractability and plethora of available genetic tools, *Drosophila melanogaster* is an attractive model organism in developmental biology. To demonstrate FlipGFP-based caspase reporter in imaging apoptosis in this model organism, we first created a transgenic UAS-FlipGFP *Drosophila*.

In the midgut of adult *Drosophila*, enterocytes are the largest class of differentiated intestinal cells.^{29,30} At physiological condition, a small number of enterocytes are believed to undergo apoptosis that is caused by passage, digestion and absorption of food and various xenobiotics.^{30–32} However, detection of this low level of physiological apoptosis has been difficult, presumably due to lack of a sensitive and robust method. To express FlipGFP in the enterocytes of the midgut, we crossed UAS-FlipGFP with Myo1A-GAL4. To mark all of the enterocytes, we coexpressed the red fluorescent protein mCherry. Fluorescence imaging of the midgut of the adult fly revealed a couple of enterocytes that were green fluorescent while most enterocytes were not, though they were all red fluorescent from mCherry (Figure 3a). This data suggests that FlipGFP is able to visualize physiological apoptosis in the midgut. Furthermore, we confirmed that green fluorescent cells were apoptotic cells using antibody staining against cleaved caspase-3 (Figure 3b). Additionally, we also showed that FlipGFP imaged apoptosis in the eye disc of *Drosophila* (Figure S7). Our data demonstrates that

FlipGFP-based caspase reporter can visualize apoptotic cells and will thus be a valuable tool for investigating apoptosis in *Drosophila*.

Lastly, we attempted to design a red fluorogenic protease reporter based on the same approach. Initially, the protease reporter based on the super folder Cherry (sfCherry) failed to fluoresce in HEK293 cells (Figure S8),³³ presumably due to its inefficient self-assembly. On the other hand, after seven rounds of directed evolution of sfCherry (Figure 4a, b) and three rounds of optimization of the “flipped” β 10–11 (Figures S8–9, Table S1), we eventually obtained a red fluorogenic protease reporter: it was not red fluorescent in the absence of TEV protease; but it became red fluorescent (22-fold) when TEV protease was coexpressed in HEK293 cells (Figure 4c–e).

In summary, we have designed a new approach, i.e. “flipping” a beta strand of GFP, in developing GFP-based fluorogenic protease reporters. FlipGFP-based protease reporter achieves ~100-fold fluorescence increase upon proteolytic cleavage, which greatly facilitates imaging of protease activity *in vivo*.^{34,35} This approach can be applied to GFP-like other color fluorescent proteins, and multicolor protease reporters will enable simultaneous imaging of multiple protease activities in living cells and animals.³⁶

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

W.M. gratefully acknowledges the use of the computing facility at Beckman Institute, University of Illinois. S.-Q. Z. is grateful to Dr. Meng-Long Feng for inspiration.

Funding

This work was supported by National Institute of General Medical Sciences (NIGMS) R01 GM115399 (to X.S.), R35 GM122548 (to T.B.K.), R01 HL054737 and R01 HL121387 (to S.R.C.), 5T32HL007731 (to H.H.), National Natural Science Foundation of China (Grant No. 31571107, 31871481 to Z.D.), and R35 GM122603 (to William F. DeGrado for support of Haifan Wu).

REFERENCES

- (1). Tait SWG; Green DR Mitochondrial regulation of cell death. Cold Spring Harbor Perspect. Biol 2013, 5 (9), a008706.
- (2). Strasser A; Jost PJ; Nagata S The many roles of FAS receptor signaling in the immune system. Immunity 2009, 30 (2), 180. [PubMed: 19239902]
- (3). Crawford ED; Wells JA Caspase substrates and cellular remodeling. Annu. Rev. Biochem 2011, 80, 1055. [PubMed: 21456965]
- (4). Campbell DS; Okamoto H Local caspase activation interacts with Slit-Robo signaling to restrict axonal arborization. J. Cell Biol 2013, 203 (4), 657. [PubMed: 24385488]
- (5). Yamaguchi Y; Shinotsuka N; Nonomura K; Takemoto K; Kuida K; Yosida H; Miura M Live imaging of apoptosis in a novel transgenic mouse highlights its role in neural tube closure. J. Cell Biol 2011, 195 (6), 1047. [PubMed: 22162136]
- (6). Takemoto K; Kuranaga E; Tonoki A; Nagai T; Miyawaki A; Miura M Local initiation of caspase activation in *Drosophila* salivary gland programmed cell death *in vivo*. Proc. Natl. Acad. Sci. U. S. A 2007, 104 (33), 13367. [PubMed: 17679695]

- (7). Takemoto K; Nagai T; Miyawaki A; Miura M Spatiotemporal activation of caspase revealed by indicator that is insensitive to environmental effects. *J. Cell Biol* 2003, 160 (2), 235. [PubMed: 12527749]
- (8). To T-L; Schepis A; Ruiz-González R; Zhang Q; Yu D; Dong Z; Coughlin SR; Shu X Rational design of a GFP-based fluorogenic caspase reporter for imaging apoptosis *in vivo*. *Cell Chemical Biology* 2016, 23 (7), 875. [PubMed: 27447051]
- (9). To T-L; Piggott BJ; Makhijani K; Yu D; Jan Y-N; Shu X Rationally designed fluorogenic protease reporter visualizes spatiotemporal dynamics of apoptosis *in vivo*. *Proc. Natl. Acad. Sci. U. S. A* 2015, 112 (11), 3338. [PubMed: 25733847]
- (10). Ding Y; Li J; Enterina JR; Shen Y; Zhang I; Tewson PH; Mo GCH; Zhang J; Quinn AM; Hughes TE; Maysinger D; Alford SC; Zhang Y; Campbell RE Ratiometric biosensors based on dimerization-dependent fluorescent protein exchange. *Nat. Methods* 2015, 12 (3), 195. [PubMed: 25622108]
- (11). (a)Do K; Boxer SG GFP Variants with Alternative β -strands and Application as a Light-driven Protease Sensor: A Tale of Two Tails. *J. Am. Chem. Soc* 2013, 135 (28), 10226. [PubMed: 23819615] (b)Deng A; Boxer SG Structural Insight into the Photochemistry of Split Green Fluorescent Proteins: A Unique Role for a His-Tag. *J. Am. Chem. Soc* 2018, 140 (1), 375. [PubMed: 29193968] (c)Do K; Boxer SG Thermodynamics, Kinetics, and Photochemistry of β -Strand Association and Dissociation in a Split-GFP System. *J. Am. Chem. Soc* 2018, 140 (1), 375. [PubMed: 29193968]
- (12). Nicholls SB; Chu J; Abbruzzese G; Tremblay KD; Hardy JA Mechanism of a genetically encoded dark-to-bright reporter for caspase activity. *J. Biol. Chem* 2011, 286 (28), 24977. [PubMed: 21558267]
- (13). Yu D; Baird MA; Allen JR; Howe ES; Klassen MP; Reade A; Makhijani K; Song Y; Liu S; Murthy Z; Zhang S-Q; Weiner OD; Kornberg TB; Jan Y-N; Davidson MW; Shu X A naturally monomeric infrared fluorescent protein for protein labeling *in vivo*. *Nat. Methods* 2015, 12 (8), 763. [PubMed: 26098020]
- (14). Yang F; Moss LG; Phillips GN The molecular structure of green fluorescent protein. *Nat. Biotechnol* 1996, 14 (10), 1246. [PubMed: 9631087]
- (15). Ormö M; Cubitt AB; Kallio K; Gross LA; Tsien RY; Remington SJ Crystal structure of the *Aequorea victoria* green fluorescent protein. *Science* 1996, 273 (5280), 1392. [PubMed: 8703075]
- (16). Cabantous S; Terwilliger TC; Waldo GS Protein tagging and detection with engineered self-assembling fragments of green fluorescent protein. *Nat. Biotechnol* 2005, 23 (1), 102. [PubMed: 15580262]
- (17). Tsien RY Constructing and exploiting the fluorescent protein paintbox (Nobel Lecture). *Angew. Chem. Int. Ed* 2009, 48 (31), 5612.
- (18). Tsien RY The green fluorescent protein. *Annu. Rev. Biochem* 1998, 67, 509. [PubMed: 9759496]
- (19). Sniegowski JA Base Catalysis of Chromophore Formation in Arg⁹⁶ and Glu²²² Variants of Green Fluorescent Protein. *J. Biol. Chem* 2005, 280 (28), 26248. [PubMed: 15888441]
- (20). Cabantous S; Nguyen HB; Pédelacq J-D; Koraïchi F; Chaudhary A; Ganguly K; Lockard MA; Favre G; Terwilliger TC; Waldo GS A new protein-protein interaction sensor based on tripartite split-GFP association. *Sci. Rep* 2013, 3, 2854. [PubMed: 24092409]
- (21). Zhang Q; Zheng Y-W; Coughlin SR; Shu X A rapid fluorogenic GPCR- β -arrestin interaction assay. *Protein Sci.* 2018, 27 (4), 874. [PubMed: 29411438]
- (22). Sorrells S; Toruno C; Stewart RA; Jette C Analysis of apoptosis in zebrafish embryos by whole-mount immunofluorescence to detect activated Caspase 3. *J. Visualized Exp* 2013, e51060.
- (23). Zhang J; Wang X; Cui W; Wang W; Zhang H; Liu L; Zhang Z; Li Z; Ying G; Zhang N; Li B Visualization of caspase-3-like activity in cells using a genetically encoded fluorescent biosensor activated by protein cleavage. *Nat. Commun* 2013, 4, 2157. [PubMed: 23857461]
- (24). Tee AR; Proud CG Staurosporine inhibits phosphorylation of translational regulators linked to mTOR. *Cell Death Differ.* 2001, 8 (8), 841. [PubMed: 11526437]

- (25). Albeck JG; Burke JM; Aldridge BB; Zhang M; Lauffenburger DA; Sorger PK Quantitative analysis of pathways controlling extrinsic apoptosis in single cells. *Mol. Cell* 2008, 30 (1), 11. [PubMed: 18406323]
- (26). Grunwald DJ; Eisen JS Headwaters of the zebrafish – emergence of a new model vertebrate. *Nat. Rev. Genet* 2002, 3 (9), 717. [PubMed: 12209146]
- (27). Szymczak AL; Workman CJ; Wang Y; Vignali KM; Dilioglou S; Vanin EF; Vignali DAA Correction of multi-gene deficiency in vivo using a single ‘self-cleaving’ 2A peptide-based retroviral vector. *Nat. Biotechnol* 2004, 22 (5), 589. [PubMed: 15064769]
- (28). Cole L; Ross LS Apoptosis in the developing zebrafish embryo. *Dev. Biol* 2001, 240 (1), 123. [PubMed: 11784051]
- (29). Jiang H; Edgar BA Intestinal stem cell function in *Drosophila* and mice. *Curr. Opin. Genet. Dev* 2012, 22 (4), 354. [PubMed: 22608824]
- (30). Apidianakis Y; Pitsouli C; Perrimon N; Rahme L Synergy between bacterial infection and genetic predisposition in intestinal dysplasia. *Proc. Natl. Acad. Sci. U. S. A* 2009, 106 (49), 20883. [PubMed: 19934041]
- (31). Liang J; Balachandra S; Ngo S; O’Brien LE Feedback regulation of steady-state epithelial turnover and organ size. *Nature* 2017, 548 (7669), 588. [PubMed: 28847000]
- (32). Okumura T; Takeda K; Taniguchi K; Adachi-Yamada T β v integrin inhibits chronic and high level activation of JNK to repress senescence phenotypes in *Drosophila* adult midgut. *PLoS One* 2014, 9 (2), No. e89387. [PubMed: 24586740]
- (33). Nguyen HB; Hung LW; Yeates TO; Terwilliger TC; Waldo GS Split green fluorescent protein as a modular binding partner for protein crystallization. *Acta Crystallogr., Sect. D: Biol. Crystallogr* 2013, 69 (12), 2513. [PubMed: 24311592]
- (34). Fuchs Y; Steller H Programmed cell death in animal development and disease. *Cell* 2011, 147 (4), 742. [PubMed: 22078876]
- (35). Jacobson MD; Weil M; Raff MC Programmed cell death in animal development. *Cell* 1997, 88 (3), 347. [PubMed: 9039261]
- (36). Lopez-Otin C; Bond JS Proteases: multifunctional enzymes in life and disease. *J. Biol. Chem* 2008, 283 (45), 30433. [PubMed: 18650443]

Figure 1. Rational design of a green fluorogenic protease reporter FlipGFP. (a) Schematic. (b) FlipGFP. (c) Sequence. (d) Normalized fluorescence (HEK293). The error bars represent standard deviation ($n = 3$). (e) Fluorescence images. (f) Representative emission spectra of FlipGFP over time. FlipGFP (β 1–9 and the cleaved β 10–11) were purified and mixed (β 1–9:2.9 μ M; β 10–11:6.5 μ M). (g) Fluorescence intensity (integrated from 490 to 550 nm) over time after mixing β 1–9 and cleaved β 10–11. Scale bars, 20 μ m.

Figure 2. FlipGFP-based caspase reporter in imaging apoptosis in mammalian cells and in zebrafish. (a) Schematic. (b) Fluorescence images of HeLa cells upon addition of staurosporine (Movie S5). (c) Representative cells (Figure S5). (d) FLICA staining. (e) Time-lapse images of zebrafish embryo expressing FlipGFP (green), mCherry (purple). Imaging started at 24 hpf (Movie S9). (f) Whole mount immunostaining of zebrafish embryo expressing FlipGFP. Arrow points to apoptotic cells. Embryos were fixed with 4% PFA at 27 hpf. Scale bars: 20 μm (b, d); 70 μm (e); 50 μm (f).

Figure 3. FlipGFP caspase reporter in imaging apoptosis in *Drosophila*. (a) Confocal images of FlipGFP in the midgut of *Drosophila*. FlipGFP was expressed in enterocytes by crossing UAS-FlipGFP with Myo1A-GAL4. mCherry was expressed. (b) Antibody staining against cleaved caspase 3 (and thus active caspase 3). Arrows point to apoptotic cells. Scale bars: 30 μm (a, top); 10 μm (a, bottom); 10 μm (b).