Published in final edited form as:

Addict Behav. 2019 June ; 93: 46–51. doi:10.1016/j.addbeh.2019.01.014.

Culturally Relevant Risk and Protective Factors for Nonmedical Use of Prescription Opioids among Incarcerated African American Men

Paris B. Wheeler, M.S.¹, Danelle Stevens-Watkins, Ph.D.¹, Myles Moody, M.A.², Jardin Dogan, M.Ed.¹, and Dominiqueca Lewis, B.S.¹

¹University of Kentucky, College of Education, Department of Educational, School, and Counseling Psychology, Dickey Hall, 251 Scott Street, Lexington, KY 40508, Paris.wheeler@uky.edu, d.stevenswatkins@uky.edu, jndo224@g.uky.edu, dle274@g.uky.edu

²University of Kentucky, College of Liberal Arts and Sciences, Department of Sociology, 1515 Patterson Office Tower, Lexington KY 40506-0027, myles.moody@uky.edu

Abstract

Background: Recent studies have demonstrated that nonmedical use of prescription opioids (NMUPO) is a national phenomenon affecting a multitude of subpopulations, including incarcerated African American men. However, there has been little investigation of the correlates of NMUPO among this population.

Objective: Grounded in primary socialization theory, the current study aimed to examine the association between family bonds, family history of prescription drug misuse, and mental health symptoms on NMUPO among African American incarcerated men.

Method: A step-wise logistic regression was conducted to determine whether family and mental health factors affected the likelihood of lifetime NMUPO.

Results: Prescription drug misuse among immediate family members ($p < .001$) and lifetime experience of serious anxiety ($p = .003$) were significantly associated with an increased likelihood of NMUPO. Having a close personal relationship with one's father was significantly associated with a decreased likelihood of NMUPO ($p = .034$).

Conclusions/Importance: Risk and protective factors are identified that can be incorporated into interventions aimed at reducing or preventing NMUPO among African American men. Directions for future research on NMUPO among African American incarcerated men are discussed.

Keywords

Incarceration; African American; prescription drug misuse; anxiety; family influence

Introduction

Prescription drugs are among the most commonly misused illicit drugs in the United States (Substance Abuse and Mental Health Services Administration [SAMHSA], 2017). Originally depicted as a predominately “White” problem (Netherland & Hansen, 2017), recent literature suggests that nonmedical use of prescription opioids (NMUPO) among African Americans has increased over time (Pouget, Fong, & Rosenblum, 2018) and may now be similar to rates among Whites (Wang, Becker, Fiellin, 2013; Ford & Rigg, 2015; McCabe, West, & Boyd, 2013). However, few studies have begun to examine correlates of prescription opioid misuse among African Americans (Nicholson & Ford, 2018; Nicholson & Vincent, 2018; Harrell & Broman, 2009; Ford & Rigg, 2015; Vaughn, Nelson, Salas-Wright, Qian, & Schootman, 2015; Nicholson, Dawson-Edwards, Higgins, & Walton, 2016; Oberleitner et al., 2018). Further, recent evidence suggests that incarcerated African American men in particular demonstrate high rates of prescription opioid misuse (Wheeler, Stevens-Watkins, Knighton, Mahaffey, & Lewis, 2018). National datasets exclude institutionalized populations, resulting in underrepresentation of incarcerated African American men in the literature on NMUPO (Pettit, 2012). However, NMUPO among incarcerated African Americans is a critical area of study due to increased risk of overdose among ex-prisoners (Binswanger et al., 2007; Binswanger, Blatchford, Mueller, & Stern, 2013). For example, evidence suggests that pre-incarceration prescription opioid misuse may increase risk for overdose after release (Winter et al., 2015) and that prescription opioids are involved in over 40% of post-release overdose deaths (Binswanger et al., 2013). In order to understand risk and protective factors of prescription opioid misuse among this understudied and vulnerable population, it is important to examine within-group, culturally relevant correlates of NMUPO (Harrell & Broman, 2009). Therefore, the purpose of the current study is to examine family influence and mental health symptoms as correlates of lifetime NMUPO among incarcerated African American men nearing community re-entry using a primary socialization theoretical framework.

Primary Socialization Theory

Primary socialization theory posits that the family is a primary influence associated with drug misuse (Oetting & Donnermeyer, 1998). Family bonds facilitate the transmission of norms through the direct monitoring, reinforcement, and sanctioning of the expression of norms (Oetting & Donnermeyer, 1998). For example, families with drug misusing parents may transmit norms about drug misuse, resulting in an increased likelihood of their offspring becoming drug misusers (Oetting & Donnermeyer, 1998). Oetting Deffenbacher, and Donnermeyer (1998) also note mental health symptoms increase risk for drug misuse. While the primary socialization model was originally developed to explain adolescent behaviors, Oetting (1999) called for examination of adult behaviors using primary socialization theory. Further, Leukefeld and Leukefeld (1999) noted the importance of using primary socialization theory to examine drug-using behaviors in the context of the criminal justice system. Thus, a primary socialization framework is used in the current study to examine the relationship between family bonds, family prescription drug misuse, mental health, and NMUPO among incarcerated African American men.

Family Influence and Prescription Drug Misuse.

Family is a cornerstone of African American culture and has been a primary source of support since the time of slavery (McCoy, 2011). The literature demonstrates that strength of bonds with parents and family history of drug misuse relate to African American individuals' prescription drug misuse. A study with an African American sample showed that adolescents who have stronger parental bonds are less likely to report NMUPO (Nicholson et al., 2016). Similarly, nationally representative data has demonstrated presence of both parents in the household and parental involvement are associated with decreased odds of reporting NMUPO (Monnat & Rigg, 2016). Further, one study showed that parental bonds were significantly associated with NMUPO among African American adolescents but not among Whites (Ford & Rigg, 2015). These results suggest that close parental bonds are protective against prescription opioid misuse and may be a culturally relevant factor for NMUPO among African Americans.

There is limited study on the impact of prescription drug misuse among family members on African American individuals' NMUPO. However, research indicates that prescription opioid misuse is related to illicit drug use. Liebschutz and colleagues (2010) found that among a majority African American sample of adults, family history of illicit drug misuse was associated with greater odds of prescription drug misuse. Blanco and colleagues (2016) found similar results using a nationally representative adult sample. However, neither of these studies examined family history of prescription drug misuse specifically, but rather examined illicit drug use among family members. Thus, the current study contributes to the literature by investigating the relationship between family prescription drug misuse on NMUPO among African American adults.

Mental Health and Prescription Drug Misuse.

The majority of research on mental health factors and prescription drug misuse comes from studies with predominantly White samples. However, findings from these studies indicate that mental health problems are also important factors to consider in the study of African American individuals' NMUPO. For example, reporting a major depressive episode (Monnat & Rigg, 2016; Edlund et al., 2015; Ford & Rigg, 2015) or depressive symptoms (Zullig & Divin, 2012; Benotsch et al., 2013) are associated with increased odds of NMUPO. Similarly, anxiety symptoms (Benotsch et al., 2013; Jeffers et al., 2015) are associated with increased likelihood of prescription opioid misuse.

Mental health problems like anxiety and depression may be more common among African American men with a history of incarceration compared to those without (Assari et al., 2018). However, only one study to date has examined the association between mental health and prescription opioid misuse among this population, which showed that more frequent mental health symptoms were associated with increased odds of NMUPO (Knighton, Stevens-Watkins, Staton, & Pangburn, 2018). This indicates the importance of considering mental health problems when examining NMUPO among African American incarcerated men.

The Current Study

Numerous gaps in the literature on NMUPO among African Americans have been identified. The majority of the literature on the relationship between family factors and NMUPO among African Americans focuses on adolescent samples. However, African Americans may be more likely to initiate substance misuse in adulthood (Doherty, Green, Reisinger, Ensminger, 2008), indicating that these factors may be pertinent to examine among adults as well. The current study adds to the literature by examining family factors among incarcerated African American men. Further, the current study extends the literature by examining mental health influences on NMUPO among African American incarcerated men, who are at greater risk for mental health problems than their non-institutionalized counterparts (Assari et al., 2018).

Because of the significant risk of post-release overdose posed by those who misused prescriptions prior to incarceration (Binswanger et al., 2013; Winter et al., 2015), there is a need to address these gaps in the literature in order to increase our understanding of risk and protective factors of NMUPO among African American incarcerated men.. The timeliness of this topic is underscored by evidence that this understudied and vulnerable population has poorer post-release drug misuse outcomes, even with drug intervention during incarceration (Rounds-Bryant, Motivans, & Pelissier, 2004). Therefore, it is critical to investigate risk and protective factors of NMUPO relevant to incarcerated African American men to better inform drug treatment among this particular population. We aimed to address this need by examining family bonds and family prescription drug misuse, mental health problems, and their impact on NMUPO among African American incarcerated men nearing re-entry.

Methods

The current study uses data derived from a larger project examining mental health and drug misuse among incarcerated African American men nearing community re-entry. Participants were recruited from the four minimum to medium security prisons across the state of Kentucky with the highest percentages of African American male prisoners. The study had the following eligibility criteria: (1) self-identified as Black or African American; (2) at least 18 years of age; (3) willing to participate in the study; (4) eligible for community re-entry within 120 days of participating in the study; and (5) returning to Jefferson or Fayette County following release. Eligible men were recruited through letters in the mail inviting them to participate in a health study and were provided with a day and time to meet with study staff.

Procedures

This study was approved by the Kentucky Department of Corrections Research Ethics Committee and the University of Kentucky Institutional Review Board (IRB), and research staff obtained written informed consent from each participant. Study interviews were conducted by masters-level mental health clinicians using Audio Computer Assisted Self-Interviewing (ACASI) techniques in order to enhance validity of responses due to the sensitive nature of the questionnaires (Djave, Brown, Gaul, & Sutton, 2014). A Federal

Certificate of Confidentiality was obtained so that participant data could not be requested by a third-party. Participants were compensated \$25 to complete the study.

Measures

Dependent variable.

Lifetime nonmedical use of prescription opioids. A modified version of the Addiction Severity Index, fifth edition (ASI-V; McLellan et al., 1992), was administered to assess impairment among drug misusers across seven domains. Lifetime nonmedical use of 11 different prescription opioids was assessed by the question, *Have you ever used ____ that was not prescribed to you?* Participants responded either 1 = *yes* or 0 = *no*. Then, a new variable was created in SPSS totaling lifetime misuse for all 11 drugs and then dichotomized, such that 1 = *ever used any prescription opioid* and 0 = *never used*.

Independent variables.

Prescription drug misuse among family members. Participants were asked whether their immediate family members ever misused prescription drugs using questions from the modified ASI-V. Family use was assessed with the question, *Does anyone in your immediate family abuse prescription drugs?* Participants responded either 1 = *yes* or 0 = *no*.

Relationship with parents. The modified ASI-V was used to ask participants to report on their relationships with several family members. Participants were asked two separate questions about their mother and father: *During your life, have you had a close, long lasting, personal relationship with your [mother; father]?* Participants responded either 1 = *yes* or 0 = *no*.

Family social support. Social support was assessed by the Family subscale of the Multidimensional Scale of Perceived Social Support (MSPSS; Zimet, Dahlem, Zimet, & Farley, 1988). The MSPSS is a 12-item measure that assesses perceived social support from three domains: friends, family, and significant others. The Family subscale consists of four items and has a Cronbach's alpha of 0.93, indicating good reliability in this sample. Participants responded to each question on a Likert scale ranging from 1 = *very strongly disagree* to 7 = *very strongly agree*. To compute the subscale score, the averages of the responses were taken to obtain a score that ranged from 1 to 7.

Lifetime mental health symptoms. Using the modified ASI-V, participants were asked the following questions: *Have you had a significant period (that was not a direct result of drug/alcohol use) in which you have experienced serious depression? Have you had a significant period (that was not a direct result of drug/alcohol use) in which you have experienced serious anxiety or tension in your lifetime?* Participants responded either 1 = *yes* or 0 = *no* to each question.

Analytical Approach

Analyses were conducted using SPSS 24.0. Descriptive analyses were conducted for all demographic information and each of the study variables. Bivariate correlations were conducted to determine significant associations between the independent and outcome

variables. A step-wise logistic regression analysis was conducted based on the theoretical framework of primary socialization theory. The variance accounted for by each step in the model is presented using Nagelkerke's R^2 .

Results

The study sample consisted of 208 African American male prisoners. All demographic information was obtained through self-report (Table 1). Over half (56.5%) of participants reported ever misusing any of the 11 prescription opioids in their lifetime. The most commonly misused prescription opioids were codeine cough syrup (44.7%) and Lortab/Hydrocodone (31.7%). Among participants, 20.6% reported prescription drug misuse among their immediate family members. Seventy-seven percent of participants reported having a close relationship with their mother whereas 39.3% had a close relationship with their father. Similarly, about a third of participants (37%) reported having a significant period of serious depression in their lifetime compared to 42.6% experiencing serious anxiety in their lifetime. The median score on the Family subscale of the MSPSS was 6.

Bivariate correlations showed significant associations between NMUPO and the demographic variables of age, ever incarcerated before 18, and total number of children, so these variables were retained for the model (Table 2). There were significant correlations between lifetime NMUPO and all independent variables except for MSPSS Family (Table 3), so this subscale was excluded from the model. A linear regression was conducted to examine multicollinearity between predictor variables and Mahalanobis distances were calculated to identify multivariate outliers. All tolerance statistics were greater than 0.1, suggesting variables were not excessively correlated. The chi-square distribution was used to determine the critical value for Mahalanobis distance at the 0.001 level, resulting in a critical value of 26.125. All cases exceeding a Mahalanobis distance of 26.125 were identified as outliers and were excluded from further analyses, resulting in a final sample size of 201.

Results from the step-wise logistic regression are presented in Table 4. At step one, participants who had ever been incarcerated before age 18 were nearly two times more likely to report lifetime NMUPO ($OR = 1.92$, 95% $CI[1.06, 3.49]$ $p = .032$). At step two, ever incarcerated before 18 was no longer a significant predictor ($p = .52$) and ever having had a severe period of anxiety was associated with 3.70 greater odds of ever using prescription opioids nonmedically ($OR = 3.70$, 95% $CI[1.56, 8.66]$, $p = .003$). At step three, ever having had a severe period of anxiety remained significant ($OR = 3.97$, 95% $CI[1.58, 9.86]$, $p = .003$). Participants with a close relationship with their father were 51% less likely to engage in NMUPO ($OR = 0.49$, 95% $CI[0.25, 0.95]$, $p = 0.034$) and participants who had prescription drug misuse among their immediate family were nearly seven times more likely to engage in NMUPO ($OR = 6.94$, 95% $CI[2.56, 18.86]$, $p < 0.001$). The final model accounted for 31% of variance in likelihood of engaging in NMUPO.

Discussion

Correlates of nonmedical use of prescription opioids (NMUPO) are understudied among African American populations (Nicholson & Ford, 2018; Harrell & Broman, 2009; Ford &

Rigg, 2015; Vaughn et al., 2015). This is particularly true for incarcerated African Americans, who are excluded from the national survey data which comprises the majority of our knowledge on NMUPO (Pettit, 2012). The significance of this exclusion is highlighted by the increased risk for overdose associated with prescription opioid misuse among incarcerated individuals (Binswanger et al., 2013; Winter et al., 2015) and poor drug outcomes among recently released African Americans (Rounds-Bryant et al., 2004). To address this gap, the current study examined the impact of family bonds, family prescription drug misuse, and mental health problems on pre-incarceration NMUPO among African American incarcerated men nearing re-entry.

Results showed that history of severe anxiety was found to increase the likelihood of NMUPO, which is consistent with research linking mental health symptoms with prescription opioid misuse among incarcerated African American men (Knighton et al., 2018). In addition, men who had close personal relationships with their fathers were 51% less likely to engage in NMUPO compared to those who did not. This is consistent with previous findings on the influence of family bonds (Monnat & Rigg, 2015; Nicholson et al., 2016). Further, men who had immediate family members who engaged in prescription drug misuse were nearly seven times more likely to also engage in NMUPO. Results from our study extend previous findings by providing evidence that in addition to illicit substance misuse (Liebschutz et al., 2010; Blanco et al., 2016), prescription drug misuse among immediate family increases likelihood of NMUPO.

The present study contributes to the research on NMUPO among African Americans because it is one of the few to examine culturally relevant factors among this population rather than using race as a covariate in a predominately White sample. Further, we were able to examine these relationships in a criminal justice context. Previous findings in a non-institutionalized sample also showed a decrease in likelihood of NMUPO when participants had strong parental bonds (Ford, 2009); however, the magnitude of this decreased likelihood (i.e. odds ratio) was not as large as in the current study. It is possible that the influence of criminal justice involvement may exacerbate the prescription opioid misuse outcomes associated with poor parent bonds. This could also contribute to the possible discrepancy in rates of prescription drug misuse between community-dwelling and incarcerated African American samples (Wheeler et al., 2018; SAMHSA, 2014).

Depressive symptoms were not significantly associated with NMUPO as previous research has indicated (Monnat & Rigg, 2015; Edlund et al., 2015; Ford & Rigg, 2015; Zullig & Divin, 2012; Benotsch et al., 2013), while anxiety remained a significant correlate. There are two possible explanations for the link between anxiety and NMUPO. Martins and colleagues suggest there may be a genetic component linking propensity towards prescription opioid misuse and anxiety disorders, meaning that certain people may be more susceptible to both of these conditions (Martins, Keyes, Storr, Zhu, & Chilcoat, 2009). If so, people who are more prone to opioid misuse and anxiety disorders may be more susceptible to the outcomes associated with poor parent bonding and family prescription drug misuse.

Another possibility is that for our sample, prescription opioids may be used for self-medication of anxious distress. To our knowledge no studies have examined self-medication

of mental health symptoms among incarcerated African Americans. However, Young and colleagues found that among adolescents (29.5% African American), those who were identified as self-treaters (i.e., exhibit nonmedical use of opioids for the treatment of pain) were more likely to exhibit symptoms of anxiety and depression (Young, McCabe, Cranford, Ross-Durow, & Boyd, 2012). This suggests that there may be an association between anxiety symptoms and using prescription opioids to self-medicate. Further study is needed to determine whether this relationship exists among incarcerated African American men, and if so, whether they are using prescription opioids to self-medicate their anxiety symptoms.

Importantly, the men from the current study sample were nearing community re-entry. There has been little study on the post-release prescription opioid misuse of African American ex-offenders. Nevertheless, African Americans face numerous barriers post-release, such as finding work and financial stress (Baron, Draine, & Salzer, 2013; Western, Braga, Davis, Sirois, 2015), that are associated with increased likelihood of mental health problems (Mahaffey, Stevens-Watkins, & Leukefeld, 2018). Given the presence of these stressors upon release, it is likely that many African American men experience serious anxiety during community re-entry, thus exacerbating post-release NMUPO outcomes.

Implications for Practice

Our findings have implications for clinical practice and development of interventions aimed at reducing the rates and associated risks of NMUPO. The potential for post-release anxiety to exacerbate NMUPO is critical given the increased risk for overdose in the months following release (Winter et al., 2015). Pre- and post-release interventions aimed at reducing prescription drug misuse should screen for anxiety and provide individuals with skills and resources needed to reduce their anxiety symptoms and cope with re-entry stressors that these men are likely to face upon release (Mahaffey et al., 2018; Baron et al., 2013; Western et al., 2015).

Our results also indicate the importance of father involvement as a protective factor against NMUPO. Close relationships with mothers was not a significant influence on NMUPO, whereas having a close relationship with one's father was particularly impactful among our sample. These results are consistent with evidence that retrospective report of greater father involvement is associated with less frequent current drug use among African American men (Barton, Kogan, Cho, & Brown 2015). Interventions incorporating father involvement may be beneficial in preventing and reducing prescription opioid misuse among African American youth transitioning into adulthood.

Notably, incarceration also tends to weaken familial bonds between African American men and their families (McLeod & Tirmazi, 2017). Thus, the children of incarcerated African American men are at risk for weaker bonds with their fathers, which the current study has demonstrated is a risk factor for NMUPO. Further, African American men who misused prescription opioids prior to incarceration may transmit NMUPO norms to their children. The combined risk factors of family NMUPO and lack of close relationship with one's father may be perpetuated onto the offspring of the same African American men who experienced these risk factors themselves, resulting in a cycle of prescription drug misuse. Clinicians working in the criminal justice system should target African American men who

misuse prescription opioids in efforts to (1) help treat their NMUPO and (2) aid in their re-entry efforts to reconnect with their families and facilitate bonds so that the cycle of NMUPO is not perpetuated onto their children.

Limitations and Future Directions

The results of the current study should be considered in the context of some limitations. This cross-sectional study utilized a sample of African American incarcerated men, thus limiting the ability to draw causal conclusions. Further, our results may also have limited external validity and may not be generalizable to women, non-African American, and non-institutionalized populations. In addition, the current study only examined nonmedical use of prescription opioids. Further study is needed to understand the risk and protective factors associated with the nonmedical use of other classes of prescription drugs, such as sedatives. Prescription sedatives may be particularly pertinent to study given the risk associated with history of anxiety demonstrated in the current study. Evidence suggests the sedative medications prescribed for anxiety may be commonly misused among incarcerated African American men (Wheeler et al., 2018). It is possible that the need to cope with anxiety symptoms paired with increased access to prescriptions intended to treat anxiety could have a multiplicative influence on the likelihood of prescription sedative misuse.

Future research should extend the investigation of risk and protective factors of NMUPO using a primary socialization framework among African American men by examining the influence of peer relationships and community factors. Utilizing a sample of incarcerated African American men provided a way to examine primary socialization theory in the context of criminal justice involvement. However, because we did not compare offenders to community members, we were unable to examine the differential impact that criminal justice involvement has as a secondary socialization source, as suggested by Leukefeld and Leukefeld (1999). Further study is needed to determine the specific ways in which incarceration impacts one's propensity towards prescription drug misuse among African American men. Overall, there is still much to learn about the risk and protective factors of NMUPO among incarcerated African American men. Given the increased risk for poor drug outcomes and post-release stressors (Rounds-Bryant et al., 2004; Baron et al., 2013; Western et al., 2015), this population deserves much more attention in the literature.

References

- Assari S, Miller RJ, Taylor RJ, Mouzon D, Keith V, & Chatters LM (2017). Discrimination fully mediates the effects of incarceration history on depressive symptoms and psychological distress among African American men. *Journal of Racial and Ethnic Health Disparities*, 1–10. doi: 10.1007/s40615-017-0364-y
- Baron RC, Draine J, & Salzer MS (2013). "I'm not sure that I can figure out how to do that": Pursuit of work among people with mental illnesses leaving jail. *American Journal of Psychiatric Rehabilitation*, 16(2), 115–135. doi: 10.1080/15487768.2013.789696 [PubMed: 23935455]
- Barton AW, Kogan SM, Cho J, & Brown GL (2015). Father involvement and young, rural African American men's engagement in substance misuse and multiple sexual partnerships. *American Journal of Community Psychology*, 56(3–4), 241–251. doi: 10.1007/s10464-015-9748-5 [PubMed: 26362297]
- Benotsch EG, Zimmerman R, Cathers L, McNulty S, Pierce J, Heck T, ... & Snipes D (2013). Non-medical use of prescription drugs, polysubstance use, and mental health in transgender adults. *Drug*

& Alcohol Dependence, 132(1), 391–394. doi: 10.1016/j.drugalcdep.2013.02.027 [PubMed: 23510637]

Binswanger IA, Blatchford PJ, Mueller SR, & Stern MF (2013). Mortality after prison release: opioid overdose and other causes of death, risk factors, and time trends from 1999 to 2009. *Annals of Internal Medicine*, 159(9), 592–600. doi: 10.7326/0003-4819-159-9-201311050-00005 [PubMed: 24189594]

Binswanger IA, Stern MF, Deyo RA, Heagerty PJ, Cheadle A, Elmore JG, & Koepsell TD (2007). Release from prison— A high risk of death for former inmates. *New England Journal of Medicine*, 356(2), 157–165. doi: 10.1056/NEJMsa064115 [PubMed: 17215533]

Blanco C, Wall MM, Okuda M, Wang S, Iza M, & Olfson M (2016). Pain as a predictor of opioid use disorder in a nationally representative sample. *American Journal of Psychiatry*, 173(12), 1189–1195. doi: 10.1176/appi.ajp.2016.15091179 [PubMed: 27444794]

Djave K, Brown EE, Gaul Z, & Sutton M (2014). Community-based electronic data collections for HIV prevention research with black/African-American men in the rural, Southern USA. *AIDS Care*, 26(10), 1309–1317. doi: 10.1080/09540121.2014.911812 [PubMed: 24773187]

Doherty EE, Green KM, Reisinger HS, & Ensminger ME (2008). Long-term patterns of drug use among an urban African-American cohort: The role of gender and family. *Journal of Urban Health*, 85(2), 250–267. doi: 10.1007/s11524-007-9246-7 [PubMed: 18247122]

Edlund MJ, Forman-Hoffman VL, Winder CR, Heller DC, Kroutil LA, Lipari RN, & Colpe LJ (2015). Opioid misuse and depression in adolescents: results from the National Survey on Drug Use and Health. *Drug & Alcohol Dependence*, 152, 131–138. doi: 10.1016/j.drugalcdep.2015.04.010 [PubMed: 25981310]

Ford JA (2009). Nonmedical prescription drug use among adolescents. *Youth & Society*, 40, 336–352. doi: 10.1177/0044118X08316345

Ford JA, & Rigg KK (2015). Racial/ethnic differences in factors that place adolescents at risk for prescription opioid misuse. *Prevention Science*, 16(5), 633–641. doi:10.1007/s11121-014-0514-y [PubMed: 25344348]

Harrell ZA, & Broman CL (2009). Racial/ethnic differences in correlates of prescription drug misuse among young adults. *Drug & Alcohol Dependence*, 104(3), 268–271. doi: 10.1016/j.drugalcdep.2009.05.017 [PubMed: 19616903]

Jeffers AJ, Benotsch EG, Green BA, Bannerman D, Darby M, Kelley T, & Martin AM (2015). Health anxiety and the non-medical use of prescription drugs in young adults: A cross-sectional study. *Addictive Behaviors*, 50, 74–77. doi: 10.1016/j.addbeh.2015.06.012 [PubMed: 26103426]

Jordan LC, & Lewis ML (2005). Paternal relationship quality as a protective factor: Preventing alcohol use among African American adolescents. *Journal of Black Psychology*, 31(2), 152–171. doi: 10.1177/0095798405274881

Knighton J, Stevens-Watkins D, Staton M, & Pangburn K (2018). Trends and mental health correlates of nonmedical opioid use among criminal justice-involved African American men. *Addictive Behaviors*, 85, 14–20. doi: 10.1016/j.addbeh.2018.04.022 [PubMed: 29803098]

Leukefeld CG, & Leukefeld S (1999). Primary socialization theory and a bio/psycho/social/spiritual practice model for substance use. *Substance Use & Misuse*, 34(7), 983–991. doi: 10.3109/10826089909039390 [PubMed: 10359216]

Liebschutz JM, Saitz R, Weiss RD, Averbuch T, Schwartz S, Meltzer EC, ... & Samet JH. (2010). Clinical factors associated with prescription drug use disorder in urban primary care patients with chronic pain. *The Journal of Pain*, 11(11), 1047–1055. doi:10.1016/j.jpain.2009.10.012 [PubMed: 20338815]

Mahaffey CC, Stevens-Watkins D, & Leukefeld C (2018). Life After: Examining the relationship between sociobehavioral factors and mental health among African American ex-offenders. *International Journal of Offender Therapy and Comparative Criminology*, 62(12), 3873–3889. doi: 10.1177/0306624X17750327 [PubMed: 29295666]

Martins SS, Keyes KM, Storr CL, Zhu H, & Chilcoat HD (2009). Pathways between nonmedical opioid use/dependence and psychiatric disorders: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Drug & Alcohol Dependence*, 103(1), 16–24. doi: 10.1016/j.drugalcdep.2009.01.019 [PubMed: 19414225]

- McCabe SE, West BT, & Boyd CJ (2013). Medical use, medical misuse, and nonmedical use of prescription opioids: results from a longitudinal study. *Pain*, 154(5), 708–713. doi:10.1016/j.pain.2013.01.011 [PubMed: 23433943]
- McCoy R (2011). African American elders, cultural traditions, and the family reunion. *Generations*, 35(3), 16–21.
- McLellan AT Kushner H, Metzger D, Peters RH, Smith I, Grissom G, Pettinati H, & Argeriou M (1992). The fifth edition of the Addiction Severity Index. *Journal of Substance Misuse Treatment*, 10, 81–89. doi: 10.1016/0740-5472(92)90062-S
- McLeod B, & Tirmazi T (2017). Paternal dimensions and complexities: Understanding the relationships between parental dyads and fathers' involvement among Black fathers with criminal records. *Journal of Family Social Work*, 20(5), 376–398. doi: 10.1080/10522158.2017.1283653
- Monnat SM, & Rigg KK (2016). Examining rural/urban differences in prescription opioid misuse among US adolescents. *Journal of Rural Health*, 32, 204–218. doi:10.1111/jrh.12141 [PubMed: 26344571]
- Netherland J, & Hansen H (2017). White opioids: Pharmaceutical race and the war on drugs that wasn't. *Biosocieties*, 12(2), 217–238. doi:10.1057/biosoc.2015.46 [PubMed: 28690668]
- Nicholson J, Dawson-Edwards C, Higgins GE, & Walton IN (2016). The nonmedical use of pain relievers among African-Americans: A test of primary socialization theory. *Journal of Substance Misuse*, 21(6), 636–639. doi:10.3109/14659891.2015.1122101
- Nicholson HL, & Ford JA (2018). Correlates of prescription opioid misuse among Black adults: Findings from the 2015 National Survey on Drug Use and Health. *Drug and alcohol dependence*, 186, 264–267. doi: 10.1016/j.drugalcdep.2017.12.006 [PubMed: 29477575]
- Nicholson HL, & Vincent J (2018). Gender differences in prescription opioid misuse among U.S. Black adults. *Substance Use & Misuse*, 1–12. doi: 10.1080/10826084.2018.1531427
- Oberleitner LM, Lumley MA, Grekin ER, MZ Smith K, Loree AM, Carty JN, & Valentino D (2018). Problematic prescription opioid use in a chronic pain treatment facility: The role of emotional processes. *Substance Use & Misuse*, 1–11. doi: 10.1080/10826084.2018.1521426
- Oetting ER (1999). Primary socialization theory. Developmental stages, spirituality, government institutions, sensation seeking, and theoretical implications. V. *Substance Use & Misuse*, 34(7), 947–982. doi: 10.3109/10826089909039389 [PubMed: 10359215]
- Oetting ER, Deffenbacher JL, & Donnermeyer JF (1998). Primary socialization theory: The role played by personal traits in the etiology of drug use and deviance. II. *Substance Use & Misuse*, 33(6), 1337–1366. doi: 10.3109/10826089809062220 [PubMed: 9603274]
- Oetting ER, & Donnermeyer JF (1998). Primary socialization theory: The etiology of drug use and deviance. I. *Substance Use & Misuse*, 33(4), 995–1026. doi: 10.3109/10826089809056252 [PubMed: 9548633]
- Pettit. (2012). *Invisible Men: Mass Incarceration and the Myth of Black Progress*. New York, NY: The Russel Sage Foundation.
- Pouget ER, Fong C, & Rosenblum A (2018). Racial/Ethnic Differences in Prevalence Trends for heroin use and non-medical use of prescription opioids among entrants to Opioid Treatment Programs, 2005–2016. *Substance Use & Misuse*, 53(2), 290–300. doi: 10.1080/10826084.2017.1334070 [PubMed: 28854060]
- Rounds-Bryant JL, Motivans MA, & Pelissier BM (2004). Correlates of Drug Treatment Outcomes for African American and White Male Federal Prisoners: Results from the TRIAD Study. *American Journal of Drug & Alcohol Abuse*, 30(3), 495–514. doi:10.1081/ADA-200032265 [PubMed: 15540489]
- Substance Abuse and Mental Health Services Administration. (2017). Key Substance Use and Mental Health Indicators in the United States: Results from the 2016 National Survey on Drug Use and Health (NSDUH Series H-52). Retrieved from <https://www.samhsa.gov/data/>
- Substance Abuse and Mental Health Services Administration. (2014). Results from the 2013 National Survey on Drug Use and Health: Detailed Tables (NSDUH Series H-48, HHS Publication No. SMA 14–4863). Rockville, MD: Substance Abuse and Mental Health Services Administration.
- Vaughn MG, Nelson EJ, Salas-Wright CP, Qian Z, & Schootman M (2016). Racial and ethnic trends and correlates of non-medical use of prescription opioids among adolescents in the United States

- 2004–2013. *Journal of Psychiatric Research*, 73, 17–24. doi: 10.1016/j.jpsychires.2015.11.003 [PubMed: 26679761]
- Wang KH, Becker WC, & Fiellin DA (2013). Prevalence and correlates for nonmedical use of prescription opioids among urban and rural residents. *Drug and Alcohol Dependence*, 127(1–3), 156–162. doi:10.1016/j.drugalcdep.2012.06.027 [PubMed: 22819293]
- Western B, Braga AA, Davis J, & Sirois C (2015). Stress and hardship after prison. *American Journal of Sociology*, 120(5), 1512–1547. doi: 10.2307/585264
- Wheeler P, Stevens-Watkins D, Mahaffey C, Knighton J, Lewis D (2018). Non-medical use of prescription drugs among Black men: Implications for community health. *The Journal of Urban Health*. 1–10. doi: 10.1007/s11524-018-0258-2 [PubMed: 29270709]
- Winter RJ, Stoové M, Degenhardt L, Hellard ME, Spelman T, Jenkinson R, ... & Kinner SA (2015). Incidence and predictors of non-fatal drug overdose after release from prison among people who inject drugs in Queensland, Australia. *Drug and Alcohol Dependence*, 153, 43–49. doi: 10.1016/j.drugalcdep.2015.06.011 [PubMed: 26105708]
- Young A, McCabe SE, Cranford JA, Ross-Durow P, & Boyd CJ (2012). Nonmedical use of prescription opioids among adolescents: subtypes based on motivation for use. *Journal of Addictive Diseases*, 31(4), 332–341. doi: 10.1080/10550887.2012.735564 [PubMed: 23244552]
- Zinet GD, Dahlem NW, Zimet SG, & Farley GK (1988). The multidimensional scale of perceived social support. *Journal of Personality Assessment*, 42(1), 30–41. doi: 10.1207/s15327752jpa5201_2
- Zullig KJ, & Divin AL (2012). The association between non-medical prescription drug use, depressive symptoms, and suicidality among college students. *Addictive Behaviors*, 37(8), 890–899. doi: 10.1016/j.addbeh.2012.02.008 [PubMed: 22541802]

Table 1.

Demographics

	<i>M</i>	<i>SD</i>	Range	<i>n</i>	%
Ethnicity					
African American				187	
Black and bi-racial				19	
Black and Moorish American				1	
Black and Cuban				1	
Black and Puerto Rican				1	
Age (years)	35.92	10.56	19.00-69.00		
Education (years)	12.00 ^a	2.53	6.00-16.00		
Partner Status (1 = Yes; 0 = No)					18
Months of Lifetime Incarceration	72.00 ^a	84.00	1.00-420.00		
Ever incarcerated before 18 (Yes)					61.5
Number of children	2.82				

^a Median years of education and months of lifetime incarceration

Table 2.**Bivariate Correlations of Demographic Factors**

	1.	2.	3.	4.	5.	6.	7.	8.
1. Ever Used prescription opioids nonmedically	-							
2. Age	-.140 *	-						
3. Incarcerated before 18	.170 ^a *	-.141 *	-					
4. Number of months incarcerated before 18	.004	.346 **	-.019	-				
5. Homeless in past year before incarceration	-.069 ^a	.089	.080 ^a	.041	-			
6. Years of education	-.116	-.044	.009	.006	-.169 *	-		
7. Marital status	-.034 ^a	-.223 **	-.031 ^a	.058	.011 ^a	-.032	-	
8. Number of children	-.214 **	-.340 **	-.153 *	.094	-.015	.016	0.355 **	-

^a = Spearman correlation

*
 $p < .05$

**
 $p < .01$

Table 3.**Bivariate Correlations of Family and Personal Factors**

	1.	2.	3.	4.	5.	6.	7.
1. Ever Used prescription opioids nonmedically	-						
2. PDU among immediate family ^a	.317 **	-					
3. Close relationship with mother ^a	-.137 *	-.113	-				
4. Close relationship with father ^a	-.183 *	-.047	.162 *	-			
5. Significant period of serious depression ^a	.191 *	.164 *	-.164 *	-.062	-		
6. Significant period of serious anxiety ^a	.280 **	.134	-.171 *	-.034	.673 **	-	
7. MPSS—Family	.104	.020	.280 **	.104	-.211 **	-.168 **	-

^a = Spearman correlation

*
 $p < .05$

**
 $p < .01$

Table 4.

Regression Coefficients

	Step 1			Step 2			Step 3		
	B	Wald	df	Exp(B)	B	Wald	df	Exp(B)	Exp(B)
Age	-.016	1.186	1	.984	-.029	3.258	1	.971	.973
Ever Incarcerated before 18	.653	4.589	1	1.922*	.614	3.766	1	1.848	1.466
Total Number of Children	-.113	3.566	1	.893	-.090	2.029	1	.914	.895
Lifetime Serious Anxiety					1.307	9.050	1	3.696**	3.947**
Lifetime Serious Depression					-.145	.109	1	.865	.695
Close Rel with Mother								-.214	.808
Close Rel with Father								-.720	.487*
NMPDU among immediate family								1.938	6.943***
Nagelkerke's R^2				0.089				0.177	0.310

* $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$