Hemorrhagic Infarction of Cerebellum and Brainstem Due to Cervical Epidural Steroid Injection – Case Report and Review of the Literature

Connie Bao, Shipping Bao

ABSTRACT

Radiculopathy is a painful condition usually due to a compressed nerve in the spine. It is reported that cervical radiculopathy has an annual incidence of 83.2 cases per 100 000 people. Cervical epidural steroid injection is one of the main treatments, but it can result in serious adverse events, including loss of vision, stroke, paralysis, and death.

A 69-year-old female died after receiving the second of three planned C5/C6 epidural steroid injections. The autopsy revealed extensive hemorrhage and necrosis of the entire right side of the cerebellum and nearby brainstem, and extensive petechial hemorrhage of the left side of the cerebellum.

This report discusses the importance of a complete autopsy in the case of death due to cervical epidural steroid injection and reporting the case either to peer review literature or to the Food and Drug Administration Adverse Event Reporting System (FAERS) database. *Acad Forensic Pathol.* 2018 8(4): 952-956

AUTHORS

Connie Bao, Champaign County Coroner's Office

Roles: Project conception and/or design, data acquisition, analysis and/or interpretation, manuscript creation and/or revision, approved final version for publication, accountable for all aspects of the work, general administrative support, writing assistance and/or technical editing.

Shipping Bao MD, Champaign County Coroner's Office

Roles: Project conception and/or design, data acquisition, analysis and/or interpretation, manuscript creation and/or revision, approved final version for publication, accountable for all aspects of the work, principal investigator of the current study, general supervision.

CORRESPONDENCE

Connie Bao, 202 S. Art Bartell Road, Urbana IL 61802, conniewbao@gmail.com

ETHICAL APPROVAL

As per Journal Policies, ethical approval was not required for this manuscript

STATEMENT OF HUMAN AND ANIMAL RIGHTS

This article does not contain any studies conducted with animals or on living human subjects

STATEMENT OF INFORMED CONSENT

No identifiable personal data were presented in this manuscript

DISCLOSURES & DECLARATION OF CONFLICTS OF INTEREST

The authors, reviewers, editors, and publication staff do not report any relevant conflicts of interest

FINANCIAL DISCLOSURE

The authors have indicated that they do not have financial relationships to disclose that are relevant to this manuscript

KEYWORDS

Forensic pathology, Steroid injection, Death, Cerebellar infarction

INFORMATION

ACADEMIC FORENSIC PATHOLOGY: THE OFFICIAL PUBLICATION OF THE NATIONAL ASSOCIATION OF MEDICAL EXAMINERS

©2018 Academic Forensic Pathology International • (ISSN: 1925-3621) • <https://doi.org/10.1177/1925362118821494>

Submitted for consideration on 28 Jul 2018. Accepted for publication on 24 Sep 2018

INTRODUCTION

Radiculopathy is a painful condition usually due to a compressed nerve in the spine. It is characterized by pain, numbness, tingling, or weakness along the course of the nerve. Radiculopathy can occur in any part of the spine, but it is most common in the neck (cervical radiculopathy) and in the lower back. It is reported that cervical radiculopathy has an annual incidence of 83.2 cases per 100 000 people (1). The primary etiology of spinal nerve compression is intervertebral foraminal stenosis associated with cervical spondylosis (70-75%) or herniation of the nucleus pulposus (20-25%). To date, treatment consists of conservative management (e.g., analgesics, cervical collar, cervical traction, physical therapy), interlaminar and transforaminal epidural steroid injections, and surgery (1).

A literature review by Benny et al. classified complications associated with cervical epidural steroid injections into two categories – minor and major (2). Minor complications are associated with temporary deficits, including vasovagal reactions and transient neurologic deficits, such as pain or weakness, prolonged paresthesias, headaches, hypersensitivity reactions, transient global amnesia, peripheral neuropraxia, and skin rashes (2).

Major complications are considered permanent injuries. Benny et al. reported there are 105 known cases of severe complications, of which only 68 have been well documented (2). These complications include brain infarction and edema, spinal cord infarction and edema, combined spinal cord and brain infarcts, cortical blindness, high spinal anesthesia, seizures, bleeding, and death. There have been 15 reports of death due to brain and spinal cord infarcts, seizure, high spinal anesthesia, and other unspecified causes (2).

CASE REPORT

A 69-year-old female was admitted to the hospital for the second of three planned C5/C6 epidural steroid injections (each containing 1 mL of 1% bupivacaine, 1 mL of 1% lidocaine, and 40 mg of methylprednis-

olone). Shortly after the injection, she complained of slurred speech, difficulty breathing, and right-sided weakness. A few hours after the procedure, a “Code Blue” was called and the patient had cardiac arrest after being noted to be hypoxic and not apneic. A computed tomography scan of the head showed a brainstem stroke with hemorrhage. She was on life support and pronounced dead two days later.

A complete autopsy was performed. The positive findings included a needle mark on the lower right side of the neck, extensive hemorrhage and necrosis of the entire right side of the cerebellum and nearby brainstem (**Image 1**), and extensive petechial hemorrhage of the left side of the cerebellum. The right side of the cerebellum was dusky, soft, and in a near-liquid state (**Image 2**). There was no hemorrhage in the cerebral hemispheres or cervical spinal cord. She also had early cirrhosis of the liver and atherosclerosis with 70% stenosis of the left anterior descending coronary artery. The cause of death in this case was hemorrhagic infarction of the cerebellum and brainstem due to right cervical epidural steroid injection.

This case was a private autopsy performed in a coroner’s office. The policy is to return all organs to the family right after the autopsy, so the brain was not fixed prior to dissection. The entire right side of the cerebellum and nearby brainstem revealed extensive hemorrhage and necrosis; they were not suitable for microscopic study. The cervical spinal cord was dissected and revealed no abnormalities. The vertebral arteries were not evaluated.

DISCUSSION

Multiple types of spinal injections, whether epidural/translaminar, transforaminal, or facet injections, are offered to patients with or without surgical spinal lesions for pain management. Although not approved by the US Food and Drug Administration (FDA), injections are being performed with increased frequency (160%) (3). Studies suggest they are typically short-acting and ineffective over the long-term and they expose patients to major risks and complications (3). The overall complication rate is about 0-16.8% (4).

Image 1: Extensive hemorrhage and necrosis of the entire right side of the cerebellum and nearby brainstem.

Image 2: The right side of the cerebellum in a dusky, soft, and near-liquid state, consistent with hemorrhagic infarction.

The Centers for Disease Control and Prevention (CDC) specifically identified 25 deaths, 337 patients sickened, and 14 000 cases of exposure to contaminated steroids; many other patients likely develop additional complications that go unreported or underreported (3). Epstein concluded that although the benefits for epidural steroid injections may include transient pain relief for those with or without surgical disease, the multitude of risks attributed to these injections outweighs the benefits (3). Injections may relieve symptoms for those without surgical lesions, but this relief may simply coincide with the self-limited course of the disease. For those with surgical lesions, injections may significantly delay requisite surgery, resulting in increased permanent neurological deficits (3, 4).

Periprocedure infarctions due to either vasospasm or embolus from particulate matter associated with the steroid injection have been postulated as mechanisms for causing neurologic symptoms (5). The embolism may be related to inadvertent intraarterial injection as corticosteroid suspensions may contain particles that occlude arterioles (5). Other mechanisms for major complications may include arterial intimal tear and vertebral artery perforation or dissection.

CONCLUSION

The FDA has warned that injection of corticosteroids into the epidural space of the spine may result in serious adverse events, including loss of vision, stroke,

paralysis, and death (6). There are many unreported or unpublished cases either because of pending civil litigation or reluctance of some physicians to report complications (2). In any death associated with cervical epidural steroid injections, a complete autopsy may be warranted to rule out other causes of death, such as acute cardiac events. The cases could be published in peer review literature or reported to the FDA Adverse Event Reporting System (FAERS) database to provide more data for research and regulations.

REFERENCES

- 1) Scanlon GC, Moeller-Bertram T, Romanowsky SM, Wallace MS. Cervical transforaminal epidural steroid injections: more dangerous than we think? *Spine (Phila Pa 1976)*. 2007 May 15; 32(11):1249-56. PMID: 17495784. <https://doi.org/10.1097/BRS.0b013e318053ec50>.
- 2) Benny B, Azari P, Briones D. Complications of cervical transforaminal epidural steroid injections. *Am J Phys Med Rehabil*. 2010 Jul; 89(7): 601-7. PMID: 20567139. <https://doi.org/10.1097/PHM.0b013e3181e2d07c>.
- 3) Epstein N. The risks of epidural and transforaminal steroid injections in the spine: commentary and a comprehensive review of the literature. *Surg Neurol Int*. 2013 Mar 22; 4(Suppl 2):S74-93. PMID: 23646278. PMCID: PMC3642757. <https://doi.org/10.4103/2152-7806.109446>.
- 4) Epstein NE. Major risks and complications of cervical epidural steroid injections: an updated review. *Surg Neurol Int*. 2018 Apr 23; 9:86. PMID: 29740507. PMCID: PMC5926212. https://doi.org/10.4103/sni.sni_85_18.
- 5) Popescu A, Lai D, Lu A, Gardner K. Stroke following epidural injections – case report and review of literature. *J Neuroimaging*. 2013 Jan; 23(1):118-21. PMID: 21699610. <https://doi.org/10.1111/j.1552-6569.2011.00615.x>.
- 6) U.S. Food and Drug Administration [Internet]. Silver Spring (MD): U.S. Department of Health and Human Services; c2018. FDA Drug Safety Communication: FDA requires label changes to warn of rare but serious neurologic problems after epidural corticosteroid injections for pain; 2014 Apr 23 [cited 2018 Jul 28]. Available from: <https://www.fda.gov/Drugs/DrugSafety/ucm394280.htm>.