Antibiotic Prescription Patterns among Children Younger than 5 Years in Nouna District, Burkina Faso

Ali Sié,¹ Boubacar Coulibaly,¹ Sawadogo Adama,¹ Lucienne Ouermi,¹ Clarisse Dah,¹ Charlemagne Tapsoba,¹ Till Bärnighausen,^{2,3,4} John Daniel Kelly,^{5,6} Thuy Doan,^{6,7} Thomas M. Lietman,^{5,6,7} Jeremy D. Keenan,^{6,7} and Catherine E. Oldenburg^{5,6,7*}

¹Centre de Recherche en Santé de Nouna, Nouna, Burkina Faso; ²Heidelberg Institute of Global Health, Medical Faculty and University Hospital, University of Heidelberg, Heidelberg, Germany; ³Africa Health Research Institute, Somkehe, South Africa; ⁴Department of Global Health and Population, Harvard T.H. Chan School of Public Health, Boston, Massachusetts; ⁵Department of Epidemiology and Biostatistics, University of California, San Francisco, San Francisco, California; ⁶Francis I. Proctor Foundation, University of California, San Francisco, San Francisco, California; ⁷Department of Ophthalmology, University of California, San Francisco, San Francisco, California

Abstract. Understanding antibiotic use may help predict emergence of antimicrobial resistance. We evaluated antibiotic prescription trends in rural Burkina Faso, where little is known about antibiotic consumption. Antibiotic prescription data for 20 communities were extracted through record review in six primary health-care facilities serving the communities. The number of antibiotic prescriptions per child-year was calculated using population-based census data from the communities. A total of 1,444 antibiotic prescriptions were made from March to June 2017 among 3,401 children in the communities. The frequency of antibiotic prescription was 1.70 prescriptions per child-year (95% CI: 1.61–1.79). Penicillins were the most common (1.04 prescriptions per child-year, 95% CI: 1.01–1.06), followed by sulfonamides (0.69 prescriptions per child-year, 95% CI: 0.67–0.71) and macrolides (0.38 prescriptions per child-year, 95% CI: 0.37–0.40). Continued monitoring of antibiotic consumption in diverse settings will be important to understand the potential for emergence of antibiotic resistance.

INTRODUCTION

Increasing antibiotic consumption has been shown to lead to selection for resistant bacterial strains at the individual and community levels.¹ In sub-Saharan Africa, pediatric antibiotic therapy is often guided by clinical symptoms because of lack of microbiologic diagnostic facilities in many regions.² Such a strategy is thought to be sensitive but nonspecific because of viral or noninfectious illnesses being classified as bacterial infection,² with sensitivity ranging from 80% to 95% and specificity ranging from 40% to 80% in young infants.³ This results in antibiotic treatment of infections or illnesses that are non-bacterial in nature. Furthermore, lack of susceptibility data to guide treatment decisions may result in inappropriate treatment, which can result in adverse clinical outcomes and mortality.^{4,5} Although the WHO recommends tailoring empiric therapy to local resistance patterns, in practice such a strategy is difficult in the absence of laboratory facilities. Understanding pediatric antibiotic use patterns may provide some evidence of which areas are at greater risk of selection for antibiotic resistance.

Pediatric antibiotic consumption has been shown to be common in multiple settings.^{6,7} However, few estimates of childhood antibiotic use exist in rural sub-Saharan African settings. Here, we conducted a retrospective review of antibiotic prescriptions in six health-care facilities in a rural setting in Burkina Faso to better understand antibiotic prescription practices in this setting.

METHODS

Study setting. This study took place in 20 communities and six Centres de Santé et de Promotion Sociale (CSPSs) located

in the rural Health and Demographic Surveillance Site (HDSS) of Nouna in northwest Burkina Faso serving a population of 3,401 children younger than 5 years.⁸ CSPSs are government primary health-care facilities in Burkina Faso that deliver preventive and curative care to a catchment area covering several villages. Health care for children younger than 5 years has been provided free of charge since 2016, and the median distance from communities to the CSPS is 8 km.^{8,9} Childhood vaccination coverage in the Nouna HDSS has increased over time, from 72% in 2012 to 81% in 2014.¹⁰ The HDSS conducts a triannual census, in which all members of all HDSS communities are enumerated.

Data collection for the study took place in July 2017, corresponding with the malaria season in this region of Burkina Faso.¹¹ The data collection period for antibiotic prescriptions covered March through June, before the malaria season. The Institutional Review Boards at the University of California, San Francisco, and the Centre de Recherche en Santé de Nouna reviewed and approved all study procedures. Verbal informed consent was obtained from the caregivers of all children participating in the caregiver survey.

Health facility data collection. We collected antibiotic prescription data from the six participating CSPSs over a 3-month period from March 23 to June 23, 2017. At each CSPS in the country, pediatric prescription data are recorded on paper forms that are submitted to the government to cover expenses related to care for children younger than 5 years. On each form, the child's age, gender, and village of residence are recorded, as well as the treatment prescribed, the dose, and duration of treatment.

Individual caregiver questionnaire. We used the most recent census (March 2017) from the HDSS to generate a list of all children aged 6–59 months residing in each community. Caregivers were interviewed in July 2017. Fifteen children were randomly selected from each census list and a caregiver interviewer was administered for each randomly selected child. In one community (Biron Badala), only seven children

* Address correspondence to Catherine E. Oldenburg, Francis I. Proctor Foundation, University of California, San Francisco, San Francisco, CA 94143. E-mail: catherine.oldenburg@ucsf.edu

were found from the list, and, therefore, children were over-sampled from a neighboring community (Biron Bobo). Caregivers were asked if they had sought care for the child at a health-care facility in the past 30 days, and if so, the reason for the visit and treatment received (antibiotics, antimalarials, or antiparasitics). To aid caregivers in accurately reporting treatments, caregivers were shown photographs of commonly used antibiotics, antimalarials, and antiparasitics. Caregivers were also asked if their child had received any antibiotics in the past 3 months, and if so, where the child received the antibiotics, for how many days the child was treated, and how many weeks had elapsed since the antibiotic prescription.

Statistical methods. We used the most recent HDSS census to calculate the denominator for each village included in the study. We tabulated the number of prescriptions of each antibiotic over the 3-month period. We calculated antibiotic prescriptions per child-year rather than defined daily doses because pediatric dosing is weight based, and weight data were not available.¹² We did not have data on adherence to antibiotic prescriptions or the duration of prescription for the indication, and, thus, were unable to calculate days of therapy. The number of prescriptions per child-year was calculated overall and by age group by dividing the total number of prescriptions by the number of child-months in each study community, and multiplying by 12 to estimate prescriptions per child-year, with 95% CIs calculated assuming a Poisson distribution. Descriptive characteristics were calculated overall for caregiver responses. All analyses were conducted in R version 3.1.5 (The R Foundation for Statistical Computing, Vienna, Austria).

RESULTS

A total of 1,444 prescriptions of 1,509 unique antibiotics were recorded during the 3-month observation period among 3,401 children residing in the study communities, corresponding to 10,203 child-months of observation. The median age of children receiving an antibiotic prescription was 18 months (IQR 8–30 months), and 47.9% were female. The overall frequency of antibiotic prescription was 0.141 prescriptions per child-month (95% CI: 0.134–0.149), equivalent to 1.70 prescriptions per child-year (95% CI: 1.61–1.79) based on the 3-month period of observation (Table 1). The most commonly prescribed antibiotic class was penicillin, 0.76 prescriptions per child-year (95% CI: 0.70–0.82). The next most commonly prescribed antibiotics were sulfonamides (0.50 prescriptions per child-year, 95% CI: 0.46–0.55) and macrolides (0.28 prescriptions per child-year, 95% CI: 0.25–0.32). Antibiotic prescriptions were highest in the youngest children, ranging from 3.6 prescriptions per child-year in children aged 0–6 months (95% CI: 3.5–3.7) to 0.32 prescriptions per child-year (95% CI: 0.34–0.29) in children aged 49–60 months (Figure 1).

Three hundred and one caregiver questionnaires were administered: The median age of the children was 37 months (IQR 24–49 months) and 50.2% were female. Caregivers of 82 children (27.3%) reported that their child had visited a health facility for a health problem in the previous month, mostly for malaria and fever (Table 2). Most who visited the health facility reported receiving antibiotics (84.1%). More than one-third (38.6%) of the caregivers reported that their child had received

TABLE 1
Antibiotic prescriptions per child-year by antibiotic class

Drug class	Number of prescriptions	Courses per child-year (95% CI)
Any antibiotic	1,444	1.70 (1.61–1.79)
Penicillins	646	0.76 (0.70–0.82)
Amoxicillin	574	0.68 (0.62–0.73)
Penicillin	61	0.07 (0.05–0.09)
Ampicillin	11	0.01 (0.006–0.02)
Sulfonamides (cotrimoxazole)	428	0.50 (0.46–0.55)
Macrolides (erythromycin)	240	0.28 (0.25–0.32)
Cephalosporins (ceftriaxone)	8	0.009 (0.004–0.02)
Fluoroquinolones (ciprofloxacin)	7	0.008 (0.003–0.02)
Aminoglycosides (gentamicin)	6	0.007 (0.003–0.02)
Tetracyclines	4	0.005 (0.001–0.01)
Nitroimidazoles (metronidazole)	170	0.20 (0.17–0.23)

an antibiotic in the past 3 months. Only one caregiver reported that the antibiotic came from a source other than the CSPS or the dispensary associated with the CSPS.

DISCUSSION

We documented relatively high antibiotic consumption among children younger than 5 years compared with that in other geographic areas in a rural area of Burkina Faso during the hot, dry period immediately before the rainy season. Prescriptions were highest among the youngest children, and dropped steadily with increasing age. Previous estimates have shown that antibiotic consumption varies widely geographically among children below age 2 years, with the highest use in countries in South Asia (10 courses per child-year) and lowest use in Brazil and South Africa (one course per child-year).⁶ In the present study, our estimate of approximately two prescriptions per child-year was lower than all but two sites included in the previous report.⁶ Recent work has shown that although antibiotic consumption in Francophone West Africa, including Burkina Faso, is among the lowest in the world, antibiotic use is increasing.¹³

Continued evaluation of antibiotic prescription trends, including in areas where consumption is thought to be lower, will be an important component of ongoing antibiotic use surveillance. Surveillance studies that evaluate microbiological trends of drug resistance in settings with increasing antibiotic prescription trends will be important for guiding empiric treatment recommendations using syndromic algorithms.

The most commonly prescribed antibiotic class was penicillin. Amoxicillin is recommended by the WHO for treatment of pneumonia.¹⁴ Malaria was the commonest reason for seeking care in this study. Malaria often presents with similar clinical symptoms as respiratory illness, including tachypnea and fever, and children receiving care for malaria were commonly given an antibiotic. In other community-based treatment settings in sub-Saharan Africa, the most common reasons for presentation include fever, upper respiratory infection, malaria, and diarrhea, and penicillins have been reported as the most commonly used antibiotic class.^{3,15–17} Antibiotic consumption data collection was primarily conducted in the dry season, during which time pneumococcal carriage and

FIGURE 1. Antibiotic prescriptions per child-year in Nouna District, Burkina Faso, by age category in months. Error bars indicate 95% CIs. Black lines indicate all antibiotic prescriptions; red, sulfonamides; purple, macrolides; and green, penicillins. This figure appears in color at www.ajtmh.org.

pneumonia have been shown to be increased relative to the rainy season.^{18,19} Extrapolating the results of this antibiotic survey to the entire calendar year requires the assumption that antibiotic prescription trends are constant over the year, which may not be true because of the seasonality of common childhood infections. Furthermore, establishing the causative agents of childhood infection was not possible in this study, as all treatments were presumptive based on symptoms. Future studies evaluating causative agents as well as duration of

therapy and completion of antibiotic courses would be useful to better understand antibiotic use in rural communities.

We were only able to extract antibiotic prescriptions from the CSPS, and if children received antibiotics from other sources, we were not able to capture it. However, in this study caregiver report of antibiotic consumption indicated that obtaining antibiotics from sources other than the CSPS was uncommon. Adding this reported usage would have resulted in negligible differences in our effect estimates. Previous work had shown high concordance between mothers' report of antibiotic use and antibiotic prescription.⁶ Most prescriptions in this setting likely came from the CSPS. However, we were unable to link children in the HDSS to the CSPS prescription data, and thus cannot validate caregiver report with CSPS records.

In this study of children in rural Burkina Faso, we demonstrated that antibiotic prescription is common, with children receiving an average of two prescriptions per year. This estimate is lower than what has been reported in high-income settings. Continued monitoring of antibiotic use in this and similar settings will be important to understand the potential for the emergence of antibiotic resistance and to develop rational antibiotic use programs.

Received September 30, 2018. Accepted for publication December 20, 2018.

TABLE 2
Child health-care access per caregiver report (N = 301)

	Number (%)
Visited health facility, past month	82 (27.3%)
Reason for health facility visit	
Malaria	66 (22.0%)
Fever	57 (19.0%)
Cough	22 (7.3%)
Diarrhea	12 (4.0%)
Antibiotics received at health visit (N = 82)	69 (84.1%)
Antimalarials received at health visit (N = 82)	79 (96.3%)
Antiparasitics received at health visit (N = 82)	3 (3.7%)
Any antibiotic use, past 3 months	113 (38.6%)
Number of days child received antibiotic (IQR)	3 (3–7)
Number of weeks since antibiotic prescription (IQR)	3 (1–5)

Published online January 28, 2019.

Financial support: This work was supported in part by the Research to Prevent Blindness Career Development Award (to C. E. O. and T. D.) and the National Eye Institute of the National Institutes of Health under Award Number K08EY026986 (to T. D.). The funders did not play a role in study design, analysis, interpretation of results, or decision to publish.

Authors' addresses: Ali Sié, Boubacar Coulibaly, Sawadogo Adama, Lucienne Ouermi, Clarisse Dah, and Charlemagne Tapsoba, Centre de Recherche en Santé de Nouna, Nouna, Burkina Faso, E-mails: sieali@yahoo.fr, boubacar@fasonet.bf, sawadama75@gmail.com, ouermil@yahoo.fr, n.clarissedah@yahoo.fr, and charlemagnetapsoba@gmail.com. Till Bärnighausen, Universitätsklinikum Heidelberg Institut für Public Health, Heidelberg Institute of Public Health, Heidelberg, Germany, E-mail: till.baernighausen@uni-heidelberg.de. John Daniel Kelly, Department of Medicine, University of California, San Francisco, San Francisco, CA, E-mail: dan.kelly@ucsf.edu. Thuy Doan, Thomas M. Lietman, and Catherine E. Oldenburg, Francis I. Proctor Foundation, University of California, San Francisco, San Francisco, CA, E-mails: thuy.doan@ucsf.edu, tom.lietman@ucsf.edu, and catherine.oldenburg@ucsf.edu. Jeremy D. Keenan, Francis I. Proctor Foundation, University of California, San Francisco, San Francisco, CA, and Department of Ophthalmology, University of California, San Francisco, San Francisco, CA, E-mail: jeremy.keenan@ucsf.edu.

REFERENCES

1. Bell BG, Schellevis F, Stobberingh E, Goossens H, Pringle M, 2014. A systematic review and meta-analysis of the effects of antibiotic consumption on antibiotic resistance. *BMC Infect Dis* 14: 13.
2. Williams PCM, Isaacs D, Berkley JA, 2018. Antimicrobial resistance among children in sub-Saharan Africa. *Lancet Infect Dis* 18: e33–e44.
3. Lee AC, Chandran A, Herbert HK, Kozuki N, Markell P, Shah R, Campbell H, Rudan I, Baqui AH, 2014. Treatment of infections in young infants in low- and middle-income countries: a systematic review and meta-analysis of frontline health worker diagnosis and antibiotic access. *PLoS Med* 11: e1001741.
4. Blomberg B et al., 2007. Antimicrobial resistance predicts death in Tanzanian children with bloodstream infections: a prospective cohort study. *BMC Infect Dis* 7: 1114–1147.
5. Thaver D, Ali SA, Zaidi AKM, 2009. Antimicrobial resistance among neonatal pathogens in developing countries. *Pediatr Infect Dis J* 28: S19–S21.
6. Rogawski ET et al., 2016. Use of antibiotics in children younger than two years in eight countries: a prospective cohort study. *Bull World Health Organ* 95: 49–61.
7. Youngster I et al., 2017. Antibiotic use in children—a cross-national analysis of 6 countries. *J Pediatr* 182: 239–244.
8. Sie A, Louis VR, Gbangou A, Müller O, Niamba L, Stieglbauer G, Yé M, Kouyaté B, Sauerborn R, Becher H, 2010. The health and demographic surveillance system (HDSS) in Nouna, Burkina Faso, 1993–2007. *Glob Health Action* 3: 5284.
9. Koulidiati J-L, Nesbitt RC, Ouedraogo N, Hien H, Robyn PJ, Compaoré P, Souares A, Brenner S, 2018. Measuring effective coverage of curative child health services in rural Burkina Faso: a cross-sectional study. *BMJ Open* 8: e020423.
10. Kagoné M, Yé M, Nébié E, Sie A, Schoeps A, Becher H, Muller O, Fisker AB, 2017. Vaccination coverage and factors associated with adherence to the vaccination schedule in young children of a rural area in Burkina Faso. *Glob Health Action* 10: 1399749.
11. Otte im Kampe E, Muller O, Sie A, Becher H, 2015. Seasonal and temporal trends in all-cause and malaria mortality in rural Burkina Faso, 1998–2007. *Malar J* 14: 300.
12. Monnet DL, 2007. Measuring antimicrobial use: the way forward. *Clin Infect Dis* 44: 671–673.
13. Klein EY, Van Boeckel TP, Martinez EM, Pant S, Gandra S, Levin SA, Goossens H, Laxminarayan R, 2018. Global increase and geographic convergence in antibiotic consumption between 2000 and 2015. *Proc Natl Acad Sci* 115: E3463–E3470.
14. World Health Organization, 2012. *Recommendations for Management of Common Childhood Conditions*.
15. Joseph F, Oladele O, Oludare O, Olatunde O, 2015. Drug prescribing pattern for under-fives in a paediatric clinic in south-western Nigeria. *Ethiop J Health Sci* 25: 73.
16. Brembilla A, Mauny F, Garcia A, Koura KG, Deloron P, Faucher JF, 2016. Antibiotics usage in infants during the first 18 months of life in Benin: a population-based cohort study. *Eur J Clin Microbiol Infect Dis* 35: 681–689.
17. Ayele B, Belay T, Gebre T, Zerihun M, Amere A, Assefa Y, Habte D, Loh AR, Stoller NE, Keenan JD, 2011. Association of community antibiotic consumption with clinically active trachoma in rural Ethiopia. *Int Health* 3: 282–288.
18. Bojang A, Jafali J, Egere UE, Hill PC, Antonio M, Jeffries D, Greenwood BM, Roca A, 2015. Seasonality of pneumococcal nasopharyngeal carriage in rural Gambia determined within the context of a cluster randomized pneumococcal vaccine trial. *PLoS One* 10: e0129649.
19. Buchner H, Rehfuess EA, 2015. Cooking and season as risk factors for acute lower respiratory infections in African children: a cross-sectional multi-country analysis. *PLoS One* 10: e0128933.