Published in final edited form as:

Sleep Med Clin. 2019 June ; 14(2): 199–208. doi:10.1016/j.jsmc.2019.01.003.

DELIVERING CBT-I IN MILITARY PERSONNEL AND VETERANS

Monica R. Kelly, PhD^a, Ruth Robbins, PhD^b, and Jennifer L. Martin, PhD^{a,c}

^aGeriatric Research, Education and Clinical Center, VA Greater Los Angeles Healthcare System, VA Sepulveda Ambulatory Care Center, 11E, 16111 Plummer Street, North Hills, CA 91343, USA

^bVA Greater Los Angeles Healthcare System, VA Sepulveda Ambulatory Care Center, 11E, 16111 Plummer Street, North Hills, CA 91343, USA

^cDepartment of Medicine, David Geffen School of Medicine, University of California, Los Angeles, Los Angeles, 16111 Plummer Street, North Hills, CA 91343, USA

Keywords

Cognitive Behavioral Therapy for Insomnia; Military; Veteran; Posttraumatic Stress Disorder; Depression; Aging; Sleep Apnea; Chronic Pain

INTRODUCTION

Insomnia is one of the two most commonly diagnosed sleep disorders among military servicemembers and veterans, and both the Department of Defense (DOD) and Veterans Administration (VA) have embraced cognitive-behavioral therapy for insomnia (CBT-I) as a valuable first-line approach to treatment, conducting large-scale provider training initiatives to meet the needs of individuals served by these two large healthcare systems.¹

Insomnia disorder seldom occurs on its own in servicemembers and veterans; rather, it is typically seen in the context of other physical and mental health conditions. It is also more common among women servicemembers and veterans compared to men, although treatment studies with military populations have included a predominance of men (typically 80–90%) with some studies including exclusively men.^{2,3} Results of randomized controlled trials and clinical implementation of CBT-I show that it retains its effectiveness when delivered to servicemembers and veterans in “real world” clinical practice, even in the context of comorbid conditions.^{1,4}

Correspondence to: Jennifer L. Martin.

*Corresponding Author:.

DISCLOSURES: The authors have no potential conflicts of interest to disclose. This work was supported by the VA Greater Los Angeles Healthcare System Geriatric Research, Education and Clinical Center and NIH/NHLBI K24HL143055 (PI: Martin).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

MENTAL HEALTH COMORBIDITIES

In servicemembers and veterans, insomnia disorder commonly co-occurs with mental health conditions, an association which has adverse consequences for functioning. Disrupted sleep is also a risk factor for mental health disorders including posttraumatic stress disorder (PTSD) and depression. A recent study of active duty Army personnel found insomnia is highly comorbid with major depressive disorder (85%), generalized anxiety disorder (83%), and PTSD (70%).⁵ The relationship between insomnia and mental health symptoms is bidirectional and chronic in nature. Insomnia symptoms prior to deployment are predictive of a greater risk for development of depression, anxiety and PTSD,⁶ while sleep difficulties are also core diagnostic criteria of PTSD and depression.⁷ Younger veterans deployed during Operation Iraqi Freedom/Operation Enduring Freedom/Operation New Dawn (OEF/OIF/OND) with premorbid psychiatric disorders are more likely to suffer from insufficient and poor quality compared to those without premorbid psychiatric conditions further supporting a bidirectional relationship between sleep and psychiatric illness.⁸

Posttraumatic stress disorder (PTSD)

Insomnia is the most frequently endorsed symptom of PTSD in servicemembers postdeployment,⁹ and evidence suggests CBT-I is highly effective in improving sleep and reducing trauma symptoms in veterans with PTSD. Nearly half of Vietnam-era veterans with combat-related PTSD endorse regular difficulties with sleep onset and fragmentation.¹⁰ Furthermore, insomnia is the most commonly reported enduring symptom following PTSD treatment in active duty servicemembers¹¹ and in individuals exposed to trauma at night.¹² Insomnia is also a predictor of poor response to exposure therapy for PTSD.¹³ These findings suggest veterans meeting criteria for insomnia disorder and PTSD may benefit from sleep-focused interventions in multiple ways.

Some sleep symptoms present differently for individuals with military-related PTSD, particularly symptoms of hyperarousal at night. Veterans with PTSD show greater night-to-night and person-to-person variability in sleep symptoms¹⁴ and distinctive maladaptive cognitions such as fear of reexperiencing nocturnal traumatic events or losing the ability to maintain vigilance during sleep.¹⁵ These hypervigilance-related features may be related to the context of the trauma (i.e. sexual assault in bed, nighttime combat), and manifest behaviorally through actions such as trying to avoid falling sleep, checking behaviors, and keeping weapons at the bedside. A theoretical model describing factors that predispose, precipitate, and perpetuate insomnia (the “3 P’s” model of insomnia)¹⁶ can be applied in the case of PTSD (see Table 1). Utilizing this conceptualization within insomnia treatment, it may be helpful to highlight factors such as regularity in sleep scheduling, addressing unhelpful thoughts with cognitive therapy, and employing coping strategies (e.g. relaxation, grounding, distraction) for distress and arousal at night.

CBT-I confers significant benefit for veterans with PTSD in terms of reducing insomnia severity across subjective sleep measures, general functioning, nightmares and sleep-related fear.^{17,18} Similar reductions in symptoms of insomnia severity have also been demonstrated for veterans with comorbid PTSD and insomnia in residential programs,¹⁹ and when CBT-I is delivered via telehealth.²⁰ CBT-I is a robust treatment for individuals with co-occurring

insomnia and PTSD even when delivered to patient populations that may not be able to easily access outpatient mental health services.

Over 70% of patients with PTSD have chronic nightmares,²¹ and combining nightmare-focused treatments with CBT-I is supported by several studies. Imagery Rehearsal Therapy (IRT) and Exposure, Rescripting and Relaxation Therapy (ERRT) have demonstrated positive results for veterans with PTSD and insomnia, when combined with CBT-I.²² Combining CBT-I with IRT improved subjective sleep, insomnia severity, nightmare frequency and overall PTSD symptoms.^{23,24} Sleep measured objectively by actigraphy also improved in one study.²⁵ Cognitive Behavioral Social Rhythm Therapy (CBSRT) utilized techniques from Social Rhythm Therapy, CBT-I and IRT and is associated with improved sleep-related and PTSD-related symptoms.²⁴ Combining CBT-I with ERRT may also be beneficial for general sleep outcomes, nightmare frequency and nightmare-related distress.²⁶ Addressing nightmares and insomnia together therefore has significant advantages.

Depression

Depression is a common mental health condition among military personnel and veterans with insomnia. Sleep difficulties are both prodromal to and diagnostic of depression,²⁷ and insomnia is a risk factor for suboptimal depression treatment response,²⁸ chronicity²⁹ and relapse after depression treatment.³⁰ Even following successful depression treatment, insomnia symptoms frequently remain, even among individuals considered in remission.³¹ In military populations, depression is often comorbid with other mental health diagnoses such as PTSD,³² further complicating the clinical picture.^{33,34,35} Additionally, in military personnel, poor sleep is related to increased suicidality and history of suicide attempts³⁶ over and above other risk factors.³⁷

Patients with comorbid insomnia and depression may present with symptoms that complicate CBT-I delivery. For example, depressed individuals may have low motivation and may find it more difficult to adhere to treatment recommendations.³⁸ Similarly, depressed individuals may have difficulty with components of CBT-I that require them to get out of bed when not sleeping. Patients with comorbid depression may benefit from better understanding and addressing common symptoms of depression, such as low mood and anhedonia, that are not considered core insomnia symptoms but can interfere with treatment progress. Behavioral activation through working together with the patient to identify previously enjoyable or mastery-oriented activities could address depression-related difficulty rising at a prescribed time as well as using the bed as coping strategy. Other modifications to CBT-I may be helpful, such as motivational interviewing to enhance adherence, early morning bright light exposure to improve mood and stabilize sleep, scheduled worry time to address rumination, and relaxation techniques. Collaborating with prescribing physicians around pharmacotherapy for depression is also essential to ensure sleep is not adversely impacted by antidepressant therapy. Results of VA's CBT-I dissemination program shows that CBT-I reduces depression severity and suicidal ideation in the context of routine clinical care, suggesting CBT-I should be offered to individuals with comorbid depression and insomnia.³⁹

Substance Use Disorders (SUD)

Insomnia and substance use often co-occur, frequently in conjunction with other mental health difficulties. Veterans often report using substances such as alcohol and marijuana to cope with insomnia symptoms,^{40,41} and insomnia symptoms predict alcohol relapse for military personnel and veterans.⁴² While mental health conditions are risk factors for substance use disorders, the relationship between depression or PTSD symptoms and alcohol use is mediated by insomnia severity among veterans, suggesting complex relationships among psychiatric conditions, substance use, and sleep.⁴³ Importantly, sleep symptoms may not remit after PTSD treatment in dual-diagnosis substance use residential treatment programs,⁴⁴ but there is evidence CBT-I may reduce risk of substance use relapse among individuals with comorbid mental health condition. While substances such as alcohol and cannabis may induce drowsiness or relaxation, sleep fragmentation is increased as substances are metabolized, and tolerance to sedation develops with continued use.⁴⁵ CBT-I in substance users may require multi-model interventions for co-occurring diagnoses as problematic alcohol use is associated with reduced treatment adherence,⁴⁶ and treatment for insomnia typically is not initiated until the recovery phase of substance use disorders. CBT-I techniques are practical and effective interventions for sleep disturbance in individuals with a history of substance use disorders. Although alcohol use relapse rates may not be reduced by CBT-I,^{47,48} a pilot study of veterans with cannabis use disorder found the CBT-I Coach mobile app was effective in reducing cannabis use and improving sleep.⁴⁹ Despite evidence for the interrelationship between sleep disturbance and substance use, limited studies exist investigating the effectiveness of CBT-I for military populations with problematic substance use patterns.

MEDICAL ISSUES

Pain

There exists a bidirectional relationship between insomnia and chronic pain, suggesting that symptoms of both must be addressed to improve functioning. Among individuals with chronic pain, 50%–70% report significantly disrupted sleep.⁵⁰ Chronic pain may cause patients to remain in lighter stages of sleep,⁵¹ and interrupted sleep can, in turn, decrease pain thresholds.^{52,53} Insomnia is a risk factor for chronic pain after acute injury,⁵⁴ and experimentally induced pain disrupts sleep.⁵⁵ In older veterans, poor sleep is associated with worse next-day subjective pain ratings.⁵⁶ Sleep difficulties are endorsed by 90% of Operation Iraqi Freedom/Operation Enduring Freedom (OIF/OEF) veterans who present with the “polytrauma clinical triad” of chronic pain, TBI, and PTSD.^{57,58} PTSD-related hyperarousal can disrupt sleep^{59,60} and lower pain thresholds as well⁶¹ (see Figure 1). This underscores the complexity of sleep in relation to polytrauma and overall health.

Multiple factors complicate insomnia treatment in the context of chronic pain. Assessing medication use and working with prescribers is important as pain medications may lead to disruptions in sleep,^{62,61} worsen sleep apnea,⁶³ and impair cognition.⁵⁰ Dysfunctional beliefs about the pain-sleep relationship, and behaviors such as extended time in bed for “rest” may be linked to poorly managed pain and contribute to insomnia symptoms. Providing psychoeducation about sleep and pain, making pain-specific behavioral

recommendations (i.e., separate “sleep” and “rest” locations), using cognitive therapy techniques to challenge unhelpful beliefs, employing behavioral activation with activity pacing, and providing relaxation strategies may be helpful for chronic pain patients with insomnia.

There is a positive effect of CBT-I on sleep and mood outcomes in chronic pain patients;^{64–66} however, despite improvements in sleep, differences in pain ratings from pre-to post-intervention are not always achieved.⁶⁷ Among OIF/OEF veterans with polytrauma, modified CBT-I in combination with prazosin reduced headache intensity and frequency, reduced daytime sleepiness, and improved cognitive performance.⁶⁸ Overall, improvements from CBT-I in sleep indices for comorbid insomnia and pain suggests CBT-I is a viable treatment option for patients with comorbid chronic pain and may improve their overall function.

Aging

Approximately one half of older veterans endorse disturbed sleep,⁶⁹ and chronic, untreated sleep disturbances are associated with increased risk for negative physical,⁷⁰ psychological,⁷¹ cognitive,^{72–74} and social⁷⁵ outcomes. CBT-I is effective first-line treatment for insomnia in older adults,^{76–78} and hypnotic medications are not recommended due to risk for adverse effects.⁷⁹ The likelihood of taking sedative-hypnotic medications to treat sleep difficulties is higher in older adults, although they increase risk for falls,^{80,81} fractures,⁸² strokes,⁸³ and mortality.⁸⁴ Studies of older veterans show benefit of CBT-I,^{4,69} although older veterans underutilize behavioral health services, posing unique challenges in delivering CBT-I to older veterans.^{4,78,85}

Adaptations to CBT-I can be made to accommodate the physical and cognitive changes associated with advanced age.⁸⁶ Older adults may adopt unhelpful expectations regarding sleep and aging,⁸⁷ and psychoeducation is important to address these maladaptive thoughts. In addition, engagement in stimulating evening activities and eliminating activities that increase risk of evening dozing are important tools in implementing CBT-I. To accommodate physical limitations and maintain safety from falls at night, modification to stimulus control instructions may be required. A recent study showed that CBT-I, delivered by trained non-mental health providers, produced significant improvements in sleep initiation and maintenance in older veterans and that benefits were sustained over a 12-month follow-up.⁶⁹ A second study explored benefits of an adapted version of CBT-I for older veterans in and Adult Day Health Care program and found benefits for this population in which mild cognitive impairment, functional limitations and limited social milieu are all commonplace.⁸⁶

Sleep Apnea

Insomnia and obstructive sleep apnea (OSA) frequently co-occur⁸⁸ and contribute to impairments in functioning and quality of life, over and above either disorder in isolation.⁸⁹ Risk factors for OSA include obesity, use of central nervous system depressants (i.e., opioids, alcohol), older age, male gender, and smoking.⁹⁰ Data from active duty military personnel indicate increases in medical encounters for both insomnia and OSA.⁹¹ Over 65%

of active duty personnel with PTSD have comorbid OSA,⁹² and OSA is the most commonly-diagnosed sleep disorder among VA users.² OSA is typically comorbid with medical conditions such as hypertension, cardiovascular diseases, diabetes, and is associated with increased mortality,^{93,94} and psychiatric disorders including depression, anxiety and PTSD.^{90,95} While the first line treatment for OSA, positive airway pressure (PAP) therapy,⁹⁶ can improve sleep-related symptoms and health,⁹⁷ comorbid insomnia is associated with lower PAP adherence.⁹⁸ This suggests CBT-I is an appropriate adjunctive treatment for OSA patients with insomnia.

Special treatment considerations or modifications may be needed to address compliance with PAP in veterans with comorbid OSA and insomnia. Early PAP use is predictive of long-term PAP adherence;⁹⁹ therefore, to encourage long-term PAP use, early intervention for insomnia is important. A Cochrane review reported that behavioral and educational interventions effectively improve PAP adherence,¹⁰⁰ and this can easily be woven into CBT-I treatments. In fact, interventions targeting veterans with co-existing OSA and insomnia have demonstrated positive sleep and PAP adherence outcomes.^{101,102} Taken together, the literature provides support for using CBT-I in veterans with comorbid insomnia and OSA and significant sleep-related benefits can be achieved. The available literature is currently limited in terms of studies assessing the role of CBT-I in OSA for active duty populations where additional challenges to PAP use are present.

ACCESS TO CARE

Because insomnia is highly prevalent in the veteran population, and CBT-I has demonstrated effectiveness in patients with complex medical and mental health histories, a system-wide dissemination has been implemented by the VA and DOD.¹ The CBT-I dissemination program utilizes didactic and competency training components through lectures and telephone-based consultation about ongoing cases for mental health providers from a variety of disciplines. Outcomes demonstrate high success rates for provider competency¹ and self-efficacy,¹⁰³ in addition to large effects for improvement in insomnia symptoms for veterans.⁴

Despite these efforts, there are still barriers to engaging CBT-I for military populations such as the limited access to care and the stigma of mental health care. VA/DOD guidelines recommend CBT-I as a first-line treatment for insomnia disorder in PTSD;¹⁰⁴ however, veterans with insomnia and PTSD remain likely to receive sedative-hypnotic medications instead.¹⁰⁵ It may be the case that providers are unaware of CBT-I or the service may not be readily available (e.g. limited to weekday sessions during standard work hours and larger, urban clinics). In addition to provider training initiatives described above, alternative delivery modalities have been explored as a means of increasing access to CBT-I. There is evidence that small-group CBT-I is as effective as individual CBT-I for veterans.¹⁰⁶ Telehealth interventions for veterans in rural areas are also effective and may enable trained providers to more easily reach veterans living in remote areas.²⁰ While internet-based delivery may further increase access, available evidence suggests it is inferior to face-to-face treatment for active duty military personnel.¹⁰⁷

Aside from systemic barriers to accessing CBT-I, there are also individual beliefs about mental health treatment, which are more prevalent in veterans who have been diagnosed with a mental health condition such as PTSD or depression, are younger, or identify as a racial/ethnic minority.¹⁰⁸ Nonetheless, one study of women veterans reported that patients actually prefer behavioral over pharmacological insomnia treatment.¹⁰⁹ Connecting active duty military personnel and veterans with CBT-I through primary care or non-mental health settings may be a means of reducing stigma and promoting engagement.

SUMMARY

Insomnia is commonly reported by military populations, especially those with comorbid mental and physical health conditions. Co-occurring conditions may result in an altered presentation of insomnia symptoms, and complicate provision of CBT-I, requiring supplementary assessment or modifications to traditional CBT-I techniques. CBT-I has consistently demonstrated positive outcomes for active duty military personnel and veterans, even in the context of significant comorbidities such as PTSD, depression, sleep apnea, and chronic pain. Despite its promise, studies of CBT-I in some military populations, including women active duty service members and active duty and veteran patients with substance use disorders, remain relatively lacking. Access to care remains a challenge, which is being addressed via provider training initiatives and exploration of alternative delivery modalities.

REFERENCES

1. Karlin BE, Trockel M, Taylor CB, Gimeno J, Manber R. National dissemination of cognitive behavioral therapy for insomnia in veterans: therapist- and patient-level outcomes. *J Consult Clin Psychol.* 2013;81(5):912–917. [PubMed: 23586730]
2. Alexander M, Ray MA, Hebert JR, et al. The National Veteran Sleep Disorder Study: Descriptive Epidemiology and Secular Trends, 2000–2010. *Sleep.* 2016;39(7):1399–1410. [PubMed: 27091538]
3. Babson KA, Wong AC, Morabito D, Kimerling R. Insomnia Symptoms Among Female Veterans: Prevalence, Risk Factors, and the Impact on Psychosocial Functioning and Health Care Utilization. *J Clin Sleep Med.* 2018;14(6):931–939. [PubMed: 29852900]
4. Karlin BE, Trockel M, Spira AP, Taylor CB, Manber R. National evaluation of the effectiveness of cognitive behavioral therapy for insomnia among older versus younger veterans. *International journal of geriatric psychiatry.* 2015;30(3):308–315. [PubMed: 24890708]
5. Brownlow JA, Klingaman EA, Boland EM, Brewster GS, Gehrman PR. Psychiatric disorders moderate the relationship between insomnia and cognitive problems in military soldiers. *J Affect Disord.* 2017;221:25–30. [PubMed: 28628764]
6. Gehrman P, Seelig AD, Jacobson IG, et al. Predeployment Sleep Duration and Insomnia Symptoms as Risk Factors for New-Onset Mental Health Disorders Following Military Deployment. *Sleep.* 2013;36(7):1009–1018. [PubMed: 23814337]
7. Diagnostic and statistical manual of mental disorders : DSM-5. Arlington, VA: American Psychiatric Association; 2013.
8. Ulmer CS, Van Voorhees E, Germain AE, et al. A Comparison of Sleep Difficulties among Iraq/Afghanistan Theater Veterans with and without Mental Health Diagnoses. *J Clin Sleep Med.* 2015;11(9):995–1005. [PubMed: 26094928]
9. McLay RN, Klam WP, Volkert SL. Insomnia is the most commonly reported symptom and predicts other symptoms of post-traumatic stress disorder in U.S. service members returning from military deployments. *Mil Med.* 2010;175(10):759–762. [PubMed: 20968266]

10. Neylan TC, Marmar CR, Metzler TJ, et al. Sleep disturbances in the Vietnam generation: findings from a nationally representative sample of male Vietnam veterans. *Am J Psychiatry*. 1998;155(7): 929–933. [PubMed: 9659859]
11. Pruiksma KE, Taylor DJ, Wachen JS, et al. Residual sleep disturbances following PTSD treatment in active duty military personnel. *Psychol Trauma*. 2016;8(6):697–701. [PubMed: 27243567]
12. Zayfert C, DeViva JC. Residual insomnia following cognitive behavioral therapy for PTSD. *J Trauma Stress*. 2004;17(1):69–73. [PubMed: 15027796]
13. Lopez CM, Lancaster CL, Gros DF, Acierno R. Residual Sleep Problems Predict Reduced Response to Prolonged Exposure among Veterans with PTSD. *J Psychopathol Behav Assess*. 2017;39(4):755–763. [PubMed: 29225414]
14. Straus LD, Drummond SP, Nappi CM, Jenkins MM, Norman SB. Sleep variability in military-related PTSD: a comparison to primary insomnia and healthy controls. *J Trauma Stress*. 2015;28(1):8–16. [PubMed: 25630526]
15. Hull A, Holliday SB, Reinhard M, Kobayashi I, Mellman T. The Role of Fear of Loss of Vigilance and Reexperiencing in Insomnia Among Veterans. *Military Behavioral Health*. 2016;4(4):373–382.
16. Spielman AGP. The varied nature of insomnia In: PJ H, ed. *Case Studies in Insomnia* Springer; 1991.
17. Talbot LS, Maguen S, Metzler TJ, et al. Cognitive behavioral therapy for insomnia in posttraumatic stress disorder: a randomized controlled trial. *Sleep*. 2014;37(2):327–341. [PubMed: 24497661]
18. Kanady JC, Talbot LS, Maguen S, et al. Cognitive Behavioral Therapy for Insomnia Reduces Fear of Sleep in Individuals With Posttraumatic Stress Disorder. *J Clin Sleep Med*. 2018;14(7):1193–1203. [PubMed: 29991428]
19. DeViva JC, McCarthy E, Bieu RK, et al. Group cognitive-behavioral therapy for insomnia delivered to veterans with posttraumatic stress disorder receiving residential treatment is associated with improvements in sleep independent of changes in posttraumatic stress disorder. *Traumatology*. 2018.
20. Laurel Franklin C, Walton JL, Raines AM, et al. Pilot study comparing telephone to inperson delivery of cognitive-behavioural therapy for trauma-related insomnia for rural veterans. *J Telemed Telecare*. 2018;24(9):629–635. [PubMed: 28950755]
21. Leskin GA, Woodward SH, Young HE, Sheikh JI. Effects of comorbid diagnoses on sleep disturbance in PTSD. *JPsychiatr Res*. 2002;36(6):449–452. [PubMed: 12393315]
22. Ho FY, Chan CS, Tang KN. Cognitive-behavioral therapy for sleep disturbances in treating posttraumatic stress disorder symptoms: A meta-analysis of randomized controlled trials. *Clin Psychol Rev*. 2016;43:90–102. [PubMed: 26439674]
23. Ulmer CS, Edinger JD, Calhoun PS. A multi-component cognitive-behavioral intervention for sleep disturbance in veterans with PTSD: a pilot study. *J Clin Sleep Med*. 2011;7(1):57–68. [PubMed: 21344046]
24. Haynes PL, Kelly M, Warner L, Quan SF, Krakow B, Bootzin RR. Cognitive Behavioral Social Rhythm Group Therapy for Veterans with posttraumatic stress disorder, depression, and sleep disturbance: Results from an open trial. *J Affect Disord*. 2016;192:234–243. [PubMed: 26748739]
25. Margolies SO, Rybarczyk B, Vrana SR, Leszczyszyn DJ, Lynch J. Efficacy of a cognitive-behavioral treatment for insomnia and nightmares in Afghanistan and Iraq veterans with PTSD. *J Clin Psychol*. 2013;69(10):1026–1042. [PubMed: 23629959]
26. Swanson LM, Favorite TK, Horin E, Arnedt JT. A combined group treatment for nightmares and insomnia in combat veterans: a pilot study. *J Trauma Stress*. 2009;22(6):639–642. [PubMed: 19908322]
27. Baglioni C, Battagliese G, Feige B, et al. Insomnia as a predictor of depression: a meta-analytic evaluation of longitudinal epidemiological studies. *J Affect Disord*. 2011; 135(1–3): 10–19. [PubMed: 21300408]
28. Troxel WM, Kupfer DJ, Reynolds CF 3rd, et al. Insomnia and objectively measured sleep disturbances predict treatment outcome in depressed patients treated with psychotherapy or psychotherapy-pharmacotherapy combinations. *J Clin Psychiatry*. 2012;73(4):478–485. [PubMed: 22152403]

29. Pigeon WR, Hegel M, Unutzer J, et al. Is insomnia a perpetuating factor for late-life depression in the IMPACT cohort? *Sleep*. 2008;31(4):481–488. [PubMed: 18457235]
30. Dombrovski AY, Mulsant BH, Houck PR, et al. Residual symptoms and recurrence during maintenance treatment of late-life depression. *J Affect Disord*. 2007;103(1–3):77–82. [PubMed: 17321595]
31. Carney CE, Segal ZV, Edinger JD, Krystal AD. A comparison of rates of residual insomnia symptoms following pharmacotherapy or cognitive-behavioral therapy for major depressive disorder. *Journal of Clinical Psychiatry*. 2007;68(2):254–260. [PubMed: 17335324]
32. Marmar CR, Schlenger W, Henn-Haase C, et al. Course of Posttraumatic Stress Disorder 40 Years After the Vietnam War: Findings From the National Vietnam Veterans Longitudinal Study. *JAMA Psychiatry* 2015;72(9):875–881. [PubMed: 26201054]
33. Ramsawh HJ, Fullerton CS, Mash HB, et al. Risk for suicidal behaviors associated with PTSD, depression, and their comorbidity in the U.S. Army. *J Affect Disord*. 2014;161:116–122. [PubMed: 24751318]
34. Curry JF, Aubuchon-Endsley N, Brancu M, et al. Lifetime major depression and comorbid disorders among current-era women veterans. *J Affect Disord*. 2014;152–154:434–440.
35. Magruder KM, Frueh BC, Knapp RG, et al. Prevalence of posttraumatic stress disorder in Veterans Affairs primary care clinics. *Gen Hosp Psychiatry*. 2005;27(3):169–179. [PubMed: 15882763]
36. Pigeon WR, Pinquart M, Conner K. Meta-analysis of sleep disturbance and suicidal thoughts and behaviors. *J Clin Psychiatry*. 2012;73(9):e1160–1167. [PubMed: 23059158]
37. Ribeiro JD, Pease JL, Gutierrez PM, et al. Sleep problems outperform depression and hopelessness as cross-sectional and longitudinal predictors of suicidal ideation and behavior in young adults in the military. *J Affect Disord*. 2012;136(3):743–750. [PubMed: 22032872]
38. Trivedi MH, Lin EH, Katon WJ. Consensus recommendations for improving adherence, self-management, and outcomes in patients with depression. *CNS Spectr*. 2007;12(8 Suppl 13): 1–27.
39. Trockel M, Karlin BE, Taylor CB, Brown GK, Manber R. Effects of cognitive behavioral therapy for insomnia on suicidal ideation in veterans. *Sleep*. 2015;38(2):259–265. [PubMed: 25515115]
40. Cucciare MA, Darrow M, Weingardt KR. Characterizing binge drinking among U. S. military Veterans receiving a brief alcohol intervention. *AddictBehav*. 2011;36(4):362–367.
41. Metrik J, Bassett SS, Aston ER, Jackson KM, Borsari B. Medicinal versus Recreational Cannabis Use among Returning Veterans. *Transl Issues Psychol Sci*. 2018;4(1):6–20. [PubMed: 30003119]
42. Williams EC, Frasco MA, Jacobson IG, et al. Risk factors for relapse to problem drinking among current and former US military personnel: a prospective study of the Millennium Cohort. *Drug Alcohol Depend*. 2015;148:93–101. [PubMed: 25599962]
43. Miller MB, DiBello AM, Carey KB, Borsari B, Pedersen ER. Insomnia severity as a mediator of the association between mental health symptoms and alcohol use in young adult veterans. *Drug Alcohol Depend*. 2017;177:221–227. [PubMed: 28618286]
44. Colvonen PJ, Ellison J, Haller M, Norman SB. Examining Insomnia and PTSD Over Time in Veterans in Residential Treatment for Substance Use Disorders and PTSD. *BehavSleep Med*. 2018:1–12.
45. Garcia AN, Salloum IM. Polysomnographic sleep disturbances in nicotine, caffeine, alcohol, cocaine, opioid, and cannabis use: A focused review. *Am J Addict*. 2015;24(7):590–598. [PubMed: 26346395]
46. Magura S, Mateu PF, Rosenblum A, Matusow H, Fong C. Risk factors for medication non-adherence among psychiatric patients with substance misuse histories. *Ment Health Subst Use*. 2014;7(4):381–390. [PubMed: 25309623]
47. Currie SR, Clark S, Hodgins DC, El-Guebaly N. Randomized controlled trial of brief cognitive-behavioural interventions for insomnia in recovering alcoholics. *Addiction*. 2004;99(9): 1121–1132. [PubMed: 15317632]
48. Arnedt JT, Conroy DA, Armitage R, Brower KJ. Cognitive-behavioral therapy for insomnia in alcohol dependent patients: a randomized controlled pilot trial. *Behav Res Ther*. 2011;49(4):227–233. [PubMed: 21377144]

49. Babson KA, Ramo DE, Baldini L, Vandrey R, Bonn-Miller MO. Mobile App-Delivered Cognitive Behavioral Therapy for Insomnia: Feasibility and Initial Efficacy Among Veterans With Cannabis Use Disorders. *JMIR Res Protoc*. 2015;4(3):e87. [PubMed: 26187404]
50. Menefee LA, Cohen MJ, Anderson WR, Doghramji K, Frank ED, Lee H. Sleep disturbance and nonmalignant chronic pain: a comprehensive review of the literature. *Pain medicine (Malden, Mass)*. 2000;1(2):156–172.
51. Moldofsky H Sleep influences on regional and diffuse pain syndromes associated with osteoarthritis. *Semin Arthritis Rheum*. 1989;18(4 Suppl 2):18–21. [PubMed: 2658072]
52. Moldofsky H, Lue FA, Smythe HA. Alpha EEG sleep and morning symptoms in rheumatoid arthritis. *The Journal of rheumatology*. 1983;10(3):373–379. [PubMed: 6887162]
53. Older SA, Battafarano DF, Danning CL, et al. The effects of delta wave sleep interruption on pain thresholds and fibromyalgia-like symptoms in healthy subjects; correlations with insulin-like growth factor I. *The Journal of rheumatology*. 1998;25(6):1180–1186. [PubMed: 9632083]
54. Finan PH, Goodin BR, Smith MT. The association of sleep and pain: an update and a path forward. *The journal of pain : official journal of the American Pain Society*. 2013;14(12):1539–1552.
55. Sutton BC, Opp MR. Musculoskeletal sensitization and sleep: chronic muscle pain fragments sleep of mice without altering its duration. *Sleep*. 2014;37(3):505–513. [PubMed: 24587573]
56. Dzierzewski JM, Williams JM, Roditi D, et al. Daily variations in objective nighttime sleep and subjective morning pain in older adults with insomnia: evidence of covariation over time. *Journal of the American Geriatrics Society*. 2010;58(5):925–930. [PubMed: 20406316]
57. Lew HL, Otis JD, Tun C, Kerns RD, Clark ME, Cifu DX. Prevalence of chronic pain, posttraumatic stress disorder, and persistent postconcussive symptoms in OIF/OEF veterans: polytrauma clinical triad. *Journal of rehabilitation research and development*. 2009;46(6):697–702. [PubMed: 20104399]
58. Lew HL, Pogoda TK, Hsu PT, et al. Impact of the “polytrauma clinical triad” on sleep disturbance in a department of veterans affairs outpatient rehabilitation setting. *American journal of physical medicine & rehabilitation*. 2010;89(6):437–445. [PubMed: 20489391]
59. Kobayashi I, Boarts JM, Delahanty DL. Polysomnographically measured sleep abnormalities in PTSD: a meta-analytic review. *Psychophysiology*. 2007;44(4):660–669. [PubMed: 17521374]
60. Yetkin S, Aydin H, Ozgen F. Polysomnography in patients with post-traumatic stress disorder. *Psychiatry Clin Neurosci*. 2010;64(3):309–317. [PubMed: 20602730]
61. Onen SH, Alloui A, Gross A, Eschallier A, Dubray C. The effects of total sleep deprivation, selective sleep interruption and sleep recovery on pain tolerance thresholds in healthy subjects. *Journal of sleep research*. 2001;10(1):35–42. [PubMed: 11285053]
62. Morasco BJ, O’Hearn D, Turk DC, Dobscha SK. Associations between prescription opioid use and sleep impairment among veterans with chronic pain. *Pain medicine (Malden, Mass)*. 2014;15(11):1902–1910.
63. Molony RR, MacPeck DM, Schiffman PL, et al. Sleep, sleep apnea and the fibromyalgia syndrome. *The Journal of rheumatology* 1986;13(4):797–800. [PubMed: 3464759]
64. Currie SR, Wilson KG, Pontefract AJ, deLaplante L. Cognitive-behavioral treatment of insomnia secondary to chronic pain. *J Consult Clin Psychol*. 2000;68(3):407–416. [PubMed: 10883557]
65. Edinger JD, Wohlgemuth WK, Krystal AD, Rice JR. Behavioral insomnia therapy for fibromyalgia patients: a randomized clinical trial. *Archives of internal medicine*. 2005;165(21):2527–2535. [PubMed: 16314551]
66. Rybarczyk B, Stepanski E, Fogg L, Lopez M, Barry P, Davis A. A placebo-controlled test of cognitive-behavioral therapy for comorbid insomnia in older adults. *J Consult Clin Psychol*. 2005;73(6): 1164–1174. [PubMed: 16392989]
67. Jungquist CR, O’Brien C, Matteson-Rusby S, et al. The efficacy of cognitive-behavioral therapy for insomnia in patients with chronic pain. *Sleep medicine*. 2010;11(3):302–309. [PubMed: 20133188]
68. Ruff RL, Ruff SS, Wang XF. Improving sleep: initial headache treatment in OIF/OEF veterans with blast-induced mild traumatic brain injury. *Journal of rehabilitation research and development*. 2009;46(9):1071–1084. [PubMed: 20437313]

69. Alessi C, Martin JL, Fiorentino L, et al. Cognitive Behavioral Therapy for Insomnia in Older Veterans Using Nonclinician Sleep Coaches: Randomized Controlled Trial. *Journal of the American Geriatrics Society*. 2016;64(9):1830–1838. [PubMed: 27550552]
70. Song Y, Dzierzewski JM, Fung CH, et al. Association Between Sleep and Physical Function in Older Veterans in an Adult Day Healthcare Program. *Journal of the American Geriatrics Society*. 2015;63(8):1622–1627. [PubMed: 26200520]
71. Jaussent I, Bouyer J, Ancelin ML, et al. Insomnia and daytime sleepiness are risk factors for depressive symptoms in the elderly. *Sleep*. 2011; 34(8): 1103–1110. [PubMed: 21804672]
72. Hahn EA, Wang HX, Andel R, Fratiglioni L. A change in sleep pattern may predict Alzheimer disease. *The American journal of geriatric psychiatry: official journal of the American Association for Geriatric Psychiatry*. 2014;22(11):1262–1271.
73. Yaffe K, Falvey CM, Hoang T. Connections between sleep and cognition in older adults. *The Lancet Neurology*. 2014;13(10):1017–1028. [PubMed: 25231524]
74. Yaffe K, Nettiksimmons J, Yesavage J, Byers A. Sleep Quality and Risk of Dementia Among Older Male Veterans. *The American journal of geriatric psychiatry: official journal of the American Association for Geriatric Psychiatry*. 2015;23(6): 651–654. [PubMed: 25794635]
75. Spira AP, Kaufmann CN, Kasper JD, et al. Association between insomnia symptoms and functional status in U.S. older adults. *The journals of gerontology Series B, Psychological sciences and social sciences*. 2014;69 Suppl 1: S35–41.
76. Bloom HG, Ahmed I, Alessi CA, et al. Evidence-based recommendations for the assessment and management of sleep disorders in older persons. *Journal of the American Geriatrics Society*. 2009;57(5):761–789. [PubMed: 19484833]
77. Morgenthaler T, Kramer M, Alessi C, et al. Practice parameters for the psychological and behavioral treatment of insomnia: an update. *An american academy of sleep medicine report*. *Sleep*. 2006;29(11):1415–1419. [PubMed: 17162987]
78. Irwin MR, Cole JC, Nicassio PM. Comparative meta-analysis of behavioral interventions for insomnia and their efficacy in middle-aged adults and in older adults 55+ years of age. *Health psychology: official journal of the Division of Health Psychology, American Psychological Association*. 2006;25(1):3–14.
79. Bertisch SM, Herzig SJ, Winkelman JW, Buettner C. National use of prescription medications for insomnia: NHANES 1999–2010. *Sleep*. 2014;37(2):343–349. [PubMed: 24497662]
80. Stone KL, Blackwell TL, Ancoli-Israel S, et al. Sleep disturbances and risk of falls in older community-dwelling men: the outcomes of Sleep Disorders in Older Men (MrOS Sleep) Study. *Journal of the American Geriatrics Society*. 2014;62(2):299–305. [PubMed: 24428306]
81. Diem SJ, Ewing SK, Stone KL, et al. Use of non-benzodiazepine sedative hypnotics and risk of falls in older men. *Journal of gerontology & geriatric research*. 2014;3(3):158. [PubMed: 25587493]
82. Bakken MS, Engeland A, Engesaeter LB, Ranhoff AH, Hunskaar S, Ruths S. Risk of hip fracture among older people using anxiolytic and hypnotic drugs: a nationwide prospective cohort study. *European journal of clinical pharmacology*. 2014;70(7): 873–880. [PubMed: 24810612]
83. Wu MP, Lin HJ, Weng SF, Ho CH, Wang JJ, Hsu YW. Insomnia subtypes and the subsequent risks of stroke: report from a nationally representative cohort. *Stroke*. 2014;45(5):1349–1354. [PubMed: 24699057]
84. Weich S, Pearce HL, Croft P, et al. Effect of anxiolytic and hypnotic drug prescriptions on mortality hazards: retrospective cohort study. *BMJ (Clinical research ed)*. 2014;348:g1996.
85. Karlin BE, Duffy M, Gleaves DH. Patterns and predictors of mental health service use and mental illness among older and younger adults in the United States. *Psychological Services*. 2008;5(3): 275–294.
86. Martin JL, Song Y, Hughes J, et al. A Four-Session Sleep Intervention Program Improves Sleep for Older Adult Day Health Care Participants: Results of a Randomized Controlled Trial. *Sleep*. 2017;40(8).
87. Morin CM, Stone J, Trinkle D, Mercer J, Remsberg S. Dysfunctional beliefs and attitudes about sleep among older adults with and without insomnia complaints. *Psychology and aging*. 1993;8(3): 463–467. [PubMed: 8216967]

88. Hein M, Lanquart JP, Loas G, Hubain P, Linkowski P. Prevalence and risk factors of moderate to severe obstructive sleep apnea syndrome in insomnia sufferers: a study on 1311 subjects. *Respir Res.* 2017;18(1):135. [PubMed: 28683800]
89. Sweetman AM, Lack LC, Catchside PG, et al. Developing a successful treatment for co-morbid insomnia and sleep apnoea. *Sleep Med Rev.* 2017;33:28–38. [PubMed: 27401786]
90. Colvonen PJ, Masino T, Drummond SP, Myers US, Angkaw AC, Norman SB. Obstructive Sleep Apnea and Posttraumatic Stress Disorder among OEF/OIF/OND Veterans. *J Clin Sleep Med.* 2015;11(5):513–518. [PubMed: 25665698]
91. J AC, Knapik JJ, Lieberman HR. Trends and factors associated with insomnia and sleep apnea in all United States military service members from 2005 to 2014. *Journal of sleep research.* 2017;26(5):665–670. [PubMed: 28448689]
92. Williams SG, Collen J, Orr N, Holley AB, Lettieri CJ. Sleep disorders in combat-related PTSD. *Sleep & breathing = Schlaf & Atmung.* 2015; 19(1): 175–182. [PubMed: 24752303]
93. Punjabi NM, Caffo BS, Goodwin JL, et al. Sleep-disordered breathing and mortality: a prospective cohort study. *PLoSmedicine.* 2009;6(8):e1000132.
94. Solomon SD, Davidson JR. Trauma: prevalence, impairment, service use, and cost. *J Clin Psychiatry.* 1997;58 Suppl 9:5–11.
95. Sharafkhaneh A, Giray N, Richardson P, Young T, Hirshkowitz M. Association of psychiatric disorders and sleep apnea in a large cohort. *Sleep.* 2005;28(11):1405–1411. [PubMed: 16335330]
96. Qaseem A, Holty JE, Owens DK, et al. Management of obstructive sleep apnea in adults: A clinical practice guideline from the American College of Physicians. *Ann Intern Med.* 2013;159(7):471–483. [PubMed: 24061345]
97. Orr JE, Smales C, Alexander TH, et al. Treatment of OSA with CPAP Is Associated with Improvement in PTSD Symptoms among Veterans. *J Clin Sleep Med.* 2017;13(1):57–63. [PubMed: 27707436]
98. Pieh C, Bach M, Popp R, et al. Insomnia symptoms influence CPAP compliance. *Sleep & breathing = Schlaf & Atmung.* 2013;17(1):99–104. [PubMed: 22311553]
99. Budhiraja R, Parthasarathy S, Drake CL, et al. Early CPAP use identifies subsequent adherence to CPAP therapy. *Sleep.* 2007;30(3):320–324. [PubMed: 17425228]
100. Wozniak DR, Lasserson TJ, Smith I. Educational, supportive and behavioural interventions to improve usage of continuous positive airway pressure machines in adults with obstructive sleep apnoea. *Cochrane Database Syst Rev.* 2014(1):CD007736.
101. Alessi CA, Martin JL, Fung CH, et al. 0407 Randomized Controlled Trial of an Integrated Behavioral Treatment in Veterans with Obstructive Sleep Apnea and Coexisting Insomnia. *Sleep.* 2018;41(suppl_1):A155–A155.
102. Fung CH, Martin JL, Josephson K, et al. Efficacy of Cognitive Behavioral Therapy for Insomnia in Older Adults With Occult Sleep-Disordered Breathing. *Psychosomatic medicine.* 2016;78(5): 629–639. [PubMed: 27136498]
103. Manber R, Trockel M, Batdorf W, et al. Lessons learned from the national dissemination of cognitive behavioral therapy for insomnia in the Veterans Health Administration: Impact of training on therapists' self-efficacy and attitudes. *Sleep medicine clinics.* 2013;8(3):399–405.
104. VA/DoD clinical practice guideline for management of post-traumatic stress. Washington (DC): Department of Veterans Affairs, Department of Defense;2017.
105. Bramoweth AD, Renqvist JG, Hanusa BH, Walker JD, Germain A, Atwood CW Jr. Identifying the Demographic and Mental Health Factors That Influence Insomnia Treatment Recommendations Within a Veteran Population. *Behav Sleep Med.* 2017:1–12.
106. Koffel E, Farrell-Carnahan L. Feasibility and preliminary real-world promise of a manualized group-based cognitive behavioral therapy for insomnia protocol for veterans. *Mil Med.* 2014;179(5): 521–528. [PubMed: 24806497]
107. Taylor DJ, Peterson AL, Pruiksma KE, et al. Internet and In-Person Cognitive Behavioral Therapy for Insomnia in Military Personnel: A Randomized Clinical Trial. *Sleep.* 2017;40(6).
108. Pietrzak RH, Johnson DC, Goldstein MB, Malley JC, Southwick SM. Perceived stigma and barriers to mental health care utilization among OEF-OIF veterans. *Psychiatr Serv.* 2009;60(8): 1118–1122. [PubMed: 19648201]

109. Culver NC, Song Y, Kate McGowan S, et al. Acceptability of Medication and Nonmedication Treatment for Insomnia Among Female Veterans: Effects of Age, Insomnia Severity, and Psychiatric Symptoms. *Clin Ther*. 2016;38(11):2373–2385. [PubMed: 28314434]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

SYNOPSIS:

Insomnia is commonly reported by military populations, especially those with comorbid mental and physical health conditions. Co-occurring conditions result in an altered presentation of insomnia symptoms, and complicate provision of Cognitive Behavioral Therapy for Insomnia (CBT-I), requiring supplementary assessment or modifications to traditional techniques. CBT-I has consistently demonstrated positive outcomes for active duty servicemembers and veterans, even in the context of significant comorbidities such as posttraumatic stress disorder, depression, sleep apnea, and chronic pain. Despite its promise, studies of CBT-I in some populations, including women and individuals with substance use disorders remain relatively understudied in active duty and veteran populations.

KEY POINTS

- Insomnia is frequently reported by active duty military personnel and veterans
- There is a bidirectional relationship between insomnia and mental and physical health disorders that commonly occur in military populations
- Cognitive Behavioral Therapy for Insomnia (CBT-I) is efficacious to improve sleep and functioning in military populations from active duty service members to older veterans
- Military personnel and veterans with complex presentations including posttraumatic stress disorder, depression, sleep apnea, and chronic pain benefit from CBT-I
- Data evaluating CBT-I are limited in women servicemembers and veterans and among those with substance use disorders despite high prevalence and risk of sleep difficulties in these subgroups

Figure 1:
Sleep and pain interact and impact implementation of insomnia treatment.

Table 1.**3 P's Model of Insomnia and PTSD in Military Populations**

Predisposing factors	Precipitating factors	Perpetuating factors
<i>Elevate risk for developing insomnia and PTSD</i>	<i>Trigger insomnia and PTSD</i>	<i>Maintain insomnia and PTSD</i>
Family history of sleep or anxiety problems	Military trauma exposure (e.g. combat, sexual or interpersonal assault)	Behaviors (e.g. irregular sleep/wake schedule, hypervigilance - checking locks)
Early-life trauma exposure	Change in military status (e.g. deployment, discharge)	Cognitions (e.g. fears of vulnerability or nightmares leading to sleep avoidance, distress regarding daytime consequences of sleep loss)
Mental health symptoms (e.g. high stress reactivity, anxiety, depression)	Social/interpersonal life stressors (e.g. divorce, childbirth)	Maladaptive coping strategies (e.g. napping, substance use including alcohol or caffeine)
Medical conditions (e.g. apnea, pain)	Economic/occupational stressors (e.g. job loss, shift work)	Environmental factors (e.g. light, temperature)