Published in final edited form as:

AIDS. 2018 November 13; 32(17): 2497–2505. doi:10.1097/QAD.0000000000001965.

The Association Between Oral Disease and Type of Antiretroviral Therapy among Perinatally HIV-Infected Youth

Caroline H SHIBOSKI¹, Tzy-Jyun YAO², Jonathan S. RUSSELL², Mark I. RYDER¹, Russell B. VAN DYKE⁴, George R. SEAGE III³, Anna-Barbara MOSCICKI⁵, and Pediatric HIV/AIDS Cohort Study

¹Department of Orofacial Sciences, School of Dentistry, University of California, San Francisco, San Francisco, California, United States of America

²Center for Biostatistics in AIDS Research (CBAR), Harvard T.H. Chan School of Public Health, Boston, Massachusetts, United States of America

³Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, United States of America

⁴School of Medicine, Tulane University, New Orleans, Louisiana, United States of America

⁵Department of Pediatrics, Division of Adolescent and Young Adult Medicine, University of California, Los Angeles, Los Angeles, California, United States of America

Abstract

Objectives: This study explores the association between combination antiretroviral therapy (cART) and oral health outcomes (dental and periodontal) among perinatally HIV-infected (PHIV) youth.

Methods: We conducted a cross-sectional study of oral health among PHIV youth participating in the Oral Health substudy of Pediatric HIV/AIDS Cohort Study (PHACS). Dentists at research sites were trained/calibrated on how to perform a standardized oral mucosal, dental, and periodontal examination. They assessed the decayed-missing-filled-surfaces and teeth index (DMFS/T). The number of decayed surfaces and teeth (DS/DT), and the number of teeth with bleeding on probing for each participant were derived from the examination. Data for analysis included lifetime measurements of CD4 count and viral load (VL), socio-demographic information, and current/ past history of ART.

Corresponding Author: Caroline H. Shiboski, DDS, MPH, PhD, Leland A. and Gladys K. Barber Distinguished Professor in Dentistry, Chair Department of Orofacial Sciences, Box 0422, Room S612, 513 Parnassus Avenue, University of California San Francisco, San Francisco, CA 94143-0422, Phone: (415)476-5976; Fax: (415)476-4204, caroline.shiboski@ucsf.edu.

Authors contributions:

Caroline H SHIBOSKI: study design; data interpretation; manuscript writing

Tzy-Jyun YAO: statistical data analysis; data interpretation; manuscript writing

Jonathan S. RUSSELL: statistical data analysis; manuscript writing

Mark I. RYDER: study design; data interpretation; manuscript writing

Russell B. VAN DYKE: study design, manuscript writing

George R. SEAGE III: study design, manuscript writing

Anna-Barbara MOSCICKI: study design; data interpretation; manuscript writing

Note: The conclusions and opinions expressed in this article are those of the authors and do not necessarily reflect those of the National Institutes of Health or U.S. Department of Health and Human Services.

Results: Among 209 PHIV youth, 95% were on ART at the time of enrollment. Among 143 PHIV youth on the same cART for at least one year, we found that the mean DT score of those receiving cART containing an integrase inhibitor (II) was 86% higher than that of those on cART without an II after adjusting for age, lifetime proportion of unsuppressed VL and CD4 nadir. Initiating protease inhibitors (PI) before age 6 years was associated with a significantly lower DMFT score compared to participants who initiated at age 6 and older.

Conclusion: Our study revealed that PHIV youth who received cART containing II had a significantly higher number of untreated active caries than those on cART without II. This may warrant closer dental surveillance of those receiving II.

INTRODUCTION

Combination antiretroviral therapy (cART) has been very successful at preserving immune function and controlling opportunistic infections among individuals infected with the human Immunodeficiency virus (HIV).[1] Oral mucosal diseases associated with advanced immunosuppression such as candidiasis and hairy leukoplakia are significantly less common among patients on cART.[2–11] In a recent study among perinatally HIV-infected (PHIV) youth participating in the Adolescent Master Protocol (AMP) of the Pediatric HIV/AIDS Cohort Study (PHACS), oral mucosal lesions were very rare.[12] This was not surprising since the majority of youth had been on cART for many years. In contrast, this group was found to have a high burden of dental and periodontal disease with 61% having at least one tooth with untreated caries, and over 30% having some form of periodontitis.[12, 13] In comparison to perinatally HIV exposed uninfected (PHEU) youth in AMP, PHIV youth had a greater mean number of untreated decayed teeth whereas no differences were found between groups with respect to periodontal disease. Of note, there were no differences in cumulative history of caries between HIV infected and uninfected groups suggesting a more recent influence on the development of current caries. Although these analyses were comprehensive in their consideration of potential risk factors for dental and periodontal disease, the effect of type of ART exposure was not assessed.

To date, 24 Food and Drug Administration-approved drugs, organized into six distinct classes, are available for treatment of HIV-1 infections: nucleoside-analog reverse transcriptase inhibitors (NRTI), non-nucleoside reverse transcriptase inhibitors (NNRTI), protease inhibitors (PI), fusion inhibitors (FI), integrase inhibitors (II), and co-receptor antagonists.[14] Given the different mechanisms of action of these drugs, their effect on the oral microbiome might vary, which may in turn affect dental and periodontal conditions. Thus the objective of this study was to examine the association between ART and oral health outcomes (dental and periodontal diseases) among PHIV youth participating in the AMP/PHACS Oral Health Substudy, controlling for indicators of HIV disease severity. We did not include oral mucosal disease as an endpoint given its very low prevalence in this study population.[12]

METHODS

Study Design and Population

We conducted a cross-sectional study of oral health among participants in 11 AMP/PHACS sites.[12, 13]. As described previously, PHACS is a prospective cohort study designed to determine the impact of HIV infection and ART among PHIV youth.[15, 16].

The study was approved by Institutional Review Boards (IRB) at clinical sites, and at the Harvard T.H. Chan School of Public Health. Parents or legal guardians provided written informed consent for their child's participation. Youth consented or assented per local IRB guidelines.

Variables and Measures

PHACS Database, and Data Collection Overview—Variables and measures collected as part of the oral study and/or obtained from the PHACS/AMP database have been described previously.[12, 13] Sixteen dentists at 11 sites were trained and calibrated on how to perform a standardized oral mucosal, dental, and periodontal examination. The mean (SD; range) concordance level was 97.7% (2.4%; 92% - 100%) for probing depths, 98.3% (2.4%; 92% - 100%) for gingival margin, and 99.2% (1.3%; 96% - 100%) for the presence/absence of clinically detectable caries.[12]

Dental and periodontal endpoints—The decayed-missing-filled-surfaces and teeth index (DMFS/T) was used to assess the prevalence and history of dental caries, and the number of decayed surfaces and teeth (DS/T) was used to assess untreated active caries.[17] The Plaque Index developed by Silness and Le was measured on all teeth (one buccal and one lingual site per tooth) as an objective measure of oral hygiene.[18] Extensive periodontal parameters were measured on 6 sites per tooth, including the bleeding on probing (BOP). [13] The number of teeth with BOP at more than 1 site was derived. All tooth counts were corrected for missing teeth by multiplying by the ratio of 28 over the number of scored teeth.

Based on the periodontal parameters of attachment loss and probing depth, periodontitis was defined as either absent, mild, moderate, or severe in accordance with the Centers for Disease Control and Prevention and the American Academy of Periodontology (CDC-AAP case definition).[19] Among those with no periodontitis, we used a definition for gingivitis adapted from work by Offenbacher and colleagues where individuals had to have BOP on at least 10% of the sites scored to be considered as having gingivitis.[20] A 5 minute unstimulated whole saliva flow rate was performed.[21, 22]

Exposure to ART—The main variables of interest reflected exposure to ART. For this analysis, cART was defined as any regimen containing at least three drugs from at least two drug classes. Classes considered were NRTI, NNRTI, PI, FI, and II.

Any regimen that stopped before a cutoff of 6 weeks (42 days) after birth was considered prophylaxis and not considered as part of our analyses. Likewise, any regimen that lasted less than 3 days was not considered.

When exploring history of ART, we considered if a participant had ever received cART and any drug within each class of medication as binary variables. Age at first exposure to any ART, cART, and class of medication was explored both as a continuous variable (years), and as a categorized variable: <2 years, 2 to <6 years, 6 years or older.

When exploring the current ART regimen, we defined as “current stable ART regimen” participants on a specific regimen with a start date at least one year prior to the oral health exam, and who had no changes to the regimen during that time. To explore the effect of specific classes of medications included in cART, we grouped the regimens by several categorized variables for the most common classes as follows: cART inclusive of an NNRTI versus cART without NNRTI; cART inclusive of a PI versus cART without PI; and cART inclusive of II versus cART without II.

Clinical indicators of history of, and current, HIV disease severity—HIV disease severity history was considered in the characterization of the study participants and in various analyses using the following clinical Indicators (among all available lifetime measures up to the time of the oral examination): percent of HIV RNA measures detectable (>400 copies/mL), CD4 cell count nadir (dichotomized to <200 or ≥200 cells/mm³), and history of a clinical AIDS-Defining Illness (as defined in CDC Stage 3).[23]

Current HIV disease severity was considered in the characterization of the study sample and in various analyses using the following binary clinical Indicators measured at or within 90 days of the oral study visit: HIV RNA viral load (<400 or ≥400 copies/mL), and CD4 cell count (<350 or ≥350 cells/mm³).

Statistical Analyses

The study population was characterized descriptively for history of ART exposure, current ART exposure, history of HIV disease severity, and current HIV disease severity. Summary statistics were calculated for continuous measures, and frequencies were calculated for discrete measures.

Summary statistics of the dental and periodontal tooth count endpoints were calculated. Considering the excess zero and over-dispersion relative to Poisson distribution for DT or other tooth counts, zero-inflated negative binomial (ZINB) models [12, 13] were applied to assess the association between each ART exposure measure and dental or periodontal outcomes (SAS proc genmod). We used a Directed Acyclic Graph (DAG) to distinguish between variables which could play potential confounding versus mediating roles.[24, 25] Thus all models were adjusted for age, and historical HIV disease severity using two indicators selected based on their associations with the ART exposures and oral health outcomes, the input of the principal investigators, and their interpretability: the percent of detectable HIV RNA measurements over the participant’s lifetime up to the oral health visit and CD4 nadir (< versus ≥200 cells/mm³). An example of DAG drawn in exploring DT as primary outcome is shown in supplemental Figure 1. All variables were only assessed through the negative binomial component of the ZINB models. We used a similar approach, relying on a similar DAG with age and historical HIV disease severity as confounders, to

explore the association between age at first cART exposure and dental or periodontal outcomes. We explored both cART and specific drug classes.

RESULTS

Socio-demographic, HIV-disease, and oral disease characteristics

Among 335 participants enrolled from 11 sites, 209 were PHIV, 53% of the PHIV youth were female, 63% were non-Hispanic Blacks, 27% were Hispanic, 10% were White/other, 16% were 10 to 13 years, 61% were 14 to 18 years, and 23% were 19 years or older at enrollment.[12] Only 38% lived with a caregiver who was their biological parent, 71% had a caregiver who had graduated high school and for 44% the main caregiver's income was \$20,000 per year.

The median current HIV RNA viral load was 40 copies/mL (1st quartile (Q1): 20; 3rd quartile (Q3): 1760), and 66 (32%) PHIV youth had a current viral load > 400 copies/mL; the median current CD4 cell count was 626 cells/mm³ (Q1: 441; Q3: 825), and 38 (18%) had a current CD4 cell count < 350 cells/mm³ (supplemental Table 1). Over their lifetime, 70 (33%) had a CD4 cell count nadir < 200 cells/mm³, 51 (24%) had a history of an AIDS-defining illness, and the median percentage of lifetime viral load measures which were detectable was 45% (Q1: 20%; Q3: 70%).

As previously described, 61% of PHIV youth had at least one tooth with untreated dental caries, and the mean number of DT among PHIV was 2.2 (SD 3.3).[12] The mean number of DT and DMFT scores were higher in PHIV youth with a CD4 nadir < 200 cells/mm³ (3.2; SD: 4.6) than among those with a nadir ≥ 200 cells/mm³ (1.7; SD:2.4).

History of ART and current regimens received by PHIV youth

All PHIV youth had received ART at some point since birth, and 198 (95%) were on ART at the time of the oral health visit. All youth received an NRTI at some point in time, the next most common antiretroviral class ever received by this group was a PI (91%) followed by an NNRTI (67%; Table 1a). The median age when an NRTI was initiated was 0.6 year, and around 3 years for the first PI among those who ever received PI, while the median age at first II was 14.7 years. Nearly three quarter of PHIV youth had been on the same regimen (or no treatment) for the past year, including 143 (93%) on cART. Among the 143 on stable cART, 78% were receiving a regimen containing PI, 30% an NNRTI, and 20% II (Table 1b).

Association between current cART regimen and oral outcomes

Current cART, dental disease and salivary function—Among 143 PHIV youth on the same cART for at least one year, cART with PI or cART with NNRTI were not associated with either DMFS/T or DS/T. However, those who received a cART regimen containing II had a significantly higher number of DT and DS than those receiving cART without II ($p < 0.001$ and $p = 0.002$, respectively), and a higher DMFT score, although this was not statistically significant (Table 2). The mean DT score for youth who received cART with an II was 86% higher than those who received cART without II after adjusting for the lifetime proportion of VL which were suppressed and nadir CD4 (Table 3).

Because a low salivary flow rate is associated with caries, we examined its association with cART. While no difference was found with respect to unstimulated salivary flow rate between those on cART with II and those on cART without, the median flow rate among PHIV youth on the same cART with NNRTI for at least 1 year was 0.5 mL/minute (Q1: 0.3 and Q3: 1 mL/minute) compared to 0.8 mL/minute (Q1: 0.5 and Q3: 1.2 mL/minute) among youth on cART without NNRTI ($p = 0.007$). However, no difference was detected with respect to DMFS/T or DS/T between those receiving a cART with NNRTI compared to those receiving a cART without. Similarly, no difference was detected with respect to DMFS/T or DS/T among those receiving the same cART with PI for at least 1 year compared to those receiving a cART without.

Current cART and periodontal disease—With respect to periodontal disease, there was no difference between those receiving a cART with versus without II, or between those receiving a cART with versus without PI. Mild to moderate periodontitis was diagnosed in 25% of PHIV youth receiving a cART with NNRTI compared to 41% among PHIV youth receiving a cART without NNRTI, and number of teeth with BOP at 2-6 sites were also lower in youth receiving a cART with NNRTI (mean 7.9 versus 10.0, Supplemental Table 2), but this difference was not statistically significant ($p = 0.09$).

Association between age of first exposure to cART regimen and oral outcomes

PHIV youth with a later initiation of cART had significantly higher DMFS and DMFT scores, and higher number of teeth with multiple BOP sites, than youth whose exposure initiation was at a younger age (Table 4). The DMFS/T difference was particularly notable among PHIV youth with a later initiation of PI with a median DMFS of 10 [Q1: 4; Q3: 21] in youth 6 years of age at initiation compared to a median DMFS of 7 [Q1: 2; Q3: 13] and 4 [Q1: 0; Q3: 8] among those who initiated PI between age 2 to 6 years and before age 2 years, respectively (Kruskal-Wallis test, $p < 0.001$). However, a similar difference was not observed among different age groups of NNRTI initiation, and number of untreated caries (DS/T) did not differ across the various initiation age groups.

The association between age at first exposure to cART and DMFS/T did not persist after adjusting for age at evaluation, lifetime proportion of unsuppressed viral load and low CD4 nadir. However, initiating PI before the age of 2, or between the ages of 2 and <6 years, was associated with a significantly lower DMFT score compared to participants who initiated at age 6 and older (Table 5a).

With respect to periodontal disease, the most notable finding was an association between age at first exposure of cART and BOP. The median number of teeth with at least 2 sites with BOP was significantly higher among youth who started cART older (Table 4). This association was still significant after adjustment (Table 5b). Also of note, the median number of teeth with at least 2 sites exhibiting BOP was significantly higher among youth who were 6 years and older at the time NNRTI initiation (7 [Q1: 2; Q3: 15]), and among those ages 2-6 years at NNRTI initiation (9.5 [Q1: 5; Q3: 16]), compared to those who initiated NNRTI younger than 2 years (4.5 [Q1: 2.5; Q3: 9]; $p = 0.02$ using a Kruskal-Wallis test).

Discussion

In this unique study exploring the effect of various cART regimens on caries and periodontal outcomes among PHIV youth, those who received a cART regimen containing II had a significantly higher number of untreated active caries than those on cART without II, after adjusting for age at oral examination, life history of unsuppressed viral load and low CD4 nadir. Of note, II were initiated, if ever, at a much older age. Considering lifetime exposure to ART, PHIV youth with a later initiation of PI (the majority of them received PI as part of a cART) had significantly higher DMFT/S scores than those initiating PI at an earlier age, after controlling for age at evaluation. With respect to periodontal disease, there was no difference between those receiving cART with versus without II. However, those with a later initiation of cART had significantly higher number of teeth with multiple BOP sites than youth whose initial exposure was at a younger age. The associations between age at first cART exposure and BOP, persisted after adjusting for current age, lifetime proportion of unsuppressed viral load, and low CD4 nadir.

While multiple studies among adults with HIV disease have shown that the prevalence of oral mucosal lesions is significantly lower among adult patients on ART in both the developed and developing world,[2–11] scarce data addressing this topic have been published among HIV-infected children. The few studies reported among children have been conducted mainly in developing countries.[26–31] Among those that included children receiving ART, while the prevalence of oral mucosal lesions was lower among these children than among those not receiving ART, these estimates were overall much higher than those we observed among PHIV youth, who had almost no oral mucosal lesions.[12] This is likely because PHIV youth in AMP/PHACS were prescribed cART early in life, and have remained on it for most of their lives. This is consistent with our finding that the PHIV youth who initiated cART before age 2 had a lower prevalence of gingival inflammation, as suggested by a lower number of teeth with BOP, and those who initiated PI earlier in life had a significantly lower score of DMFS/T, which reflects the history of caries. Another possible explanation is that the group of youth who started cART earlier in life may have had better access to care, including dental care, although 87% of PHIV youth who initiated cART before age 2 years reported having a source of dental care at the time of this study, compared to 74% of the youth who initiated cART after age 2 years (data not shown), and adjusting for this factor did not change the estimate of the association.

With respect to our finding that among PHIV youth on the same cART for at least one year, those who received cART containing II had significantly higher numbers of DT and DS than those on cART without II: since II are often used as salvage therapy, the group receiving II was possibly sicker closer to the time of the oral examination (since they typically initiated II during adolescence), which could partly explain the higher prevalence of caries in that group, that persisted when explored with a multivariable model. The DMFT/S was not found to be significantly higher in the group receiving II, presumably because it captures long-term history of caries, rather than more recent active disease. Sicker youth may be less likely to seek dental care due to ascendance of medical concerns, decreased energy. This finding may also reflect decreased ability of caregiver to address medication adherence of youth. Older age may be another possible explanation, however, age was not found to be associated with

DT in our initial multivariable model exploring this association. It may also be that II alter the oral microbiome to select more cariogenic bacteria. We are currently exploring this hypothesis through ongoing oral microbiome studies. Salivary hypofunction is also a known risk factor for the development of sudden onset caries, and while we did not find that those on II had lower saliva flow rate than those not receiving II, a difference may not have been detected reliably given the cross sectional nature of our study. Furthermore, both groups had an unstimulated salivary flow rate that was above the 0.1 mL/min threshold and were not considered as having salivary hypofunction.[22]

Our study is unique in that our team of trained/calibrated dentists performed a comprehensive standardized oral examination (including an oral soft tissue examination, caries assessment, and full mouth periodontal evaluation) in a group of PHIV youth with extensive and complete medical history data, including ART, collected at regular time intervals since the onset of the AMP/PHACS project more than a decade ago. The limitation of our study is that it is cross-sectional, which hampers the determination of causality in the associations we uncovered. However, we are in the process of implementing a follow-up visit to our oral AMP/PHACS sub-study, which will allow us to further explore some of our findings as youth transition to adulthood.

Studies have shown that early initiation of cART is associated with better health outcomes in children,[15, 32–34] and the present study suggests that this is also true for oral health outcomes, including dental and gingival health. For instance, we found that PHIV youth who initiated PI earlier in life had significantly lower DMFS/T scores, and those who initiated cART before age 2 had a lower prevalence of gingival inflammation. Furthermore, the finding that PHIV youth who received a cART regimen containing II had a significantly higher number of DT/S than those on cART without II suggest that youth receiving II should undergo closer surveillance of their dental status through preventative visits scheduled perhaps as often as three times per year. Daily topical fluoride home applications and/or regular fluoride varnish applications as part of preventative dental visits should be promoted in this group. Meanwhile, further research into the effect of various cART regimens on the oral microbiome are warranted to better understand the pathogenesis of dental caries among PHIV youth exposed to these regimens almost since birth. Such studies are currently ongoing.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank the children and families for their participation in PHACS, and the individuals and institutions involved in the conduct of PHACS. The study was supported by the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development with co-funding from the National Institute on Drug Abuse, the National Institute of Allergy and Infectious Diseases, the Office of AIDS Research, the National Institute of Mental Health, the National Institute of Neurological Disorders and Stroke, the National Institute on Deafness and Other Communication Disorders, the National Institute of Dental and Craniofacial Research, and the National Institute on Alcohol Abuse and Alcoholism, through cooperative agreements with the Harvard T.H. Chan School of Public Health (HD052102) (Principal Investigator: George Seage; Project Director: Julie Alperen) and the Tulane University School of Medicine (HD052104) (Principal Investigator: Russell Van Dyke; Co-Principal Investigator: Ellen Chadwick;

Project Director: Patrick Davis). Data management services were provided by Frontier Science and Technology Research Foundation (PI: Suzanne Siminski), and regulatory services and logistical support were provided by Westat, Inc (PI: Julie Davidson).

The following institutions, clinical site investigators and staff participated in conducting PHACS AMP and AMP Up in 2016, in alphabetical order: **Ann & Robert H. Lurie Children's Hospital of Chicago:** Ram Yogeve, Margaret Ann Sanders, Kathleen Malee, Scott Hunter; **Baylor College of Medicine:** William Shearer, Mary Paul, Norma Cooper, Lynnette Harris; **Bronx Lebanon Hospital Center:** Murli Purswani, Mahboobullah Mirza Baig, Alma Villegas; **Children's Diagnostic & Treatment Center:** Ana Puga, Sandra Navarro, Patricia A. Garvie, James Blood; **Boston Children's Hospital:** Sandra K. Burchett, Nancy Karthas, Betsy Kammerer; **Jacobi Medical Center:** Andrew Wiznia, Marlene Burey, Ray Shaw, Raphaelle Auguste; **Rutgers - New Jersey Medical School:** Arry Dieudonne, Linda Bettica, Juliette Johnson; **St. Christopher's Hospital for Children:** Janet S. Chen, Maria Garcia Bulkley, Latreaca Ivey, Mitzie Grant; **St. Jude Children's Research Hospital:** Katherine Knapp, Kim Allison, Megan Wilkins, Jamie Russell-Bell; **San Juan Hospital/Department of Pediatrics:** Midnela Acevedo-Flores, Heida Rios, Vivian Olivera; **Tulane University School of Medicine:** Margarita Sillio, Medea Gabriel, Patricia Sirois; **University of California, San Diego:** Stephen A. Spector, Kim Norris, Sharon Nichols; **University of Colorado Denver Health Sciences Center:** Elizabeth McFarland, Eric Cagwin, Emily Barr, Alisa Katai; **University of Miami:** Gwendolyn Scott, Grace Alvarez, Gabriel Fernandez, Anai Cuadra.

References

1. Buchacz K, Baker RK, Palella FJ Jr., Chmiel JS, Lichtenstein KA, Novak RM, et al. AIDS-defining opportunistic illnesses in US patients, 1994–2007: a cohort study. *AIDS* 2010;24:1549–1559. [PubMed: 20502317]
2. Ramirez-Amador V, Esquivel-Pedraza L, Sierra-Madero J, Anaya-Saavedra G, Gonzalez-Ramirez I, Ponce-de-Leon S. The Changing Clinical Spectrum of Human Immunodeficiency Virus (HIV)-Related Oral Lesions in 1,000 Consecutive Patients: A 12-Year Study in a Referral Center in Mexico. *Medicine (Baltimore)* 2003;82:39–50. [PubMed: 12544709]
3. Arribas J, Hernandez-Albujar S, Gonzalez-Garcia J, Pena J, Gozalez A, Canedo T, et al. Impact of Protease Inhibitor Therapy on HIV-related Oropharyngeal Candidiasis. *AIDS* 2000;14:979–985. [PubMed: 10853979]
4. Greenspan D, Gange S, Phelan J, Navazesh M, Alves M, MacPhail L, et al. Incidence of Oral Lesions in HIV-1-infected Women: Reduction with HAART. *J Dent Res* 2004;83:145–150. [PubMed: 14742653]
5. Nicolatou-Galitis O, Velegraki A, Paikos S, Economopoulou P, Stefaniotis T, Papanikolaou I, et al. Effect of PI-HAART on the prevalence of oral lesions in HIV-I infected patients. A Greek study. *Oral Diseases* 2004;10:145–150. [PubMed: 15089923]
6. Patton LL, McKaig RG, Strauss RP, Rogers D, Eron JJ Jr. Changing prevalence of oral manifestations of human immuno-deficiency virus in the era of protease inhibitor therapy. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2000;89:299–304. [PubMed: 10710453]
7. Schmidt-Westhausen A, Pripke F, Bergmann F, Reichart P. Decline in the rate of oral opportunistic infections following introduction of highly active antiretroviral therapy. *J Oral Pathol Med* 2000;29:336–341. [PubMed: 10947250]
8. Ceballos-Salobrena A, Gaitan-Cepeda L, Ceballos-Garcia L, Lezama-Del Valle D. Oral Lesions in HIV/AIDS Patients Undergoing Highly Active Antiretroviral Treatment Including Protease Inhibitors: A New Face of Oral AIDS? *AIDS Patient Care and STDs* 2000;14:627–635. [PubMed: 11119429]
9. Hodgson TA, Greenspan D, Greenspan JS. Oral lesions of HIV disease and HAART in industrialized countries. *Adv Dent Res* 2006;19:57–62. [PubMed: 16672551]
10. Tappuni AR, Fleming GJ. The effect of antiretroviral therapy on the prevalence of oral manifestations in HIV-infected patients: a UK study. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2001;92:623–628. [PubMed: 11740479]
11. Umadevi KM, Ranganathan K, Pavithra S, Hemalatha R, Saraswathi TR, Kumarasamy N, et al. Oral lesions among persons with HIV disease with and without highly active antiretroviral therapy in southern India. *J Oral Pathol Med* 2007;36:136–141. [PubMed: 17305634]

12. Moscicki AB, Yao TJ, Ryder MI, Russell JS, Dominy SS, Patel K, et al. The Burden of Oral Disease among Perinatally HIV-Infected and HIV-Exposed Uninfected Youth. *PLoS One* 2016, 6 14;11(6):e0156459. doi: 10.1371/journal.pone.0156459. [PubMed: 27299992]
13. Ryder MI, Yao TJ, Russell JS, Moscicki AB, Shiboski CH, Pediatric HIVACS. Prevalence of periodontal diseases in a multicenter cohort of perinatally HIV-infected and HIV-exposed and uninfected youth. *J Clin Periodontol* 2017;44:2–12. [PubMed: 27801947]
14. Arts EJ, Hazuda DJ. HIV-1 antiretroviral drug therapy. *Cold Spring Harb Perspect Med* 2012;2:a007161. [PubMed: 22474613]
15. Van Dyke RB, Patel K, Siberry GK, Burchett SK, Spector SA, Chernoff MC, et al. Antiretroviral treatment of US children with perinatally acquired HIV infection: temporal changes in therapy between 1991 and 2009 and predictors of immunologic and virologic outcomes. *J Acquir Immune Defic Syndr* 2011;57:165–173. [PubMed: 21407086]
16. Alperen J, Brummel S, Tassiopoulos K, Mellins CA, Kacanek D, Smith R, et al. Prevalence of and risk factors for substance use among perinatally human immunodeficiency virus-infected and perinatally exposed but uninfected youth. *J Adolesc Health* 2014;54:341–349. [PubMed: 24239286]
17. Klein H PC, Knutson JW. Studies on dental caries: I. Dental status and dental needs of elementary school children. *Public Health Rep* 1938;53:751–765.
18. Silness J, Løe H. Periodontal Disease in Pregnancy. II. Correlation between Oral Hygiene and Periodontal Condition. *Acta Odontol Scand* 1964;22:121–135. [PubMed: 14158464]
19. Eke PI, Page RC, Wei L, Thornton-Evans G, Genco RJ. Update of the case definitions for population-based surveillance of periodontitis. *J Periodontol* 2012;83:1449–1454. [PubMed: 22420873]
20. Offenbacher S, Barros SP, Beck JD. Rethinking periodontal inflammation. *J Periodontol* 2008;79:1577–1584. [PubMed: 18673013]
21. Navazesh M Methods for collecting saliva. *Ann N Y Acad Sci* 1993;694:72–77. [PubMed: 8215087]
22. Navazesh M, Christensen C, Brightman V. Clinical criteria for the diagnosis of salivary gland hypofunction. *J Dent Res* 1992;71:1363–1369. [PubMed: 1629451]
23. Centers for Disease C, Prevention. Revised surveillance case definition for HIV infection--United States, 2014. *MMWR Recomm Rep* 2014;63:1–10.
24. Textor J Drawing and Analyzing Causal DAGs with DAGitty: User Manual for Version 2.2. In. <http://www.dagitty.net/history/v2.2/manual-2.x.pdf>; 2014.
25. Glymour MM, Greenland S. Causal diagrams Chapter 12 In: *Modern Epidemiology*. Edited by Rothman KJGS, Lash TL. Philadelphia, PA: Lippincott Williams & Wilkins; 2008.
26. Naidoo S, Chikte U. Oro-facial manifestations in paediatric HIV: a comparative study of institutionalized and hospital outpatients. *Oral Dis* 2004;10:13–18. [PubMed: 14996288]
27. Meless D, Ba B, Faye M, Diby JS, N'Zore S, Datte S, et al. Oral lesions among HIV-infected children on antiretroviral treatment in West Africa. *Trop Med Int Health* 2014;19:246–255. [PubMed: 24386972]
28. Okunseri C, Badner V, Wiznia A, Rosenberg M. Prevalence of oral lesions and percent CD4+ T-lymphocytes in HIV-infected children on antiretroviral therapy. *AIDS Patient Care STDS* 2003;17:5–11. [PubMed: 12614515]
29. Oyedemi OA, Gbolahan OO, Abe EO, Agelebe E. Oral and dental lesions in HIV infected Nigerian children. *Pan Afr Med J* 2015;20:287. [PubMed: 26161210]
30. Ranganathan K, Geethalakshmi E, Krishna Mohan Rao U, Vidya KM, Kumarasamy N, Solomon S. Orofacial and systemic manifestations in 212 paediatric HIV patients from Chennai, South India. *Int J Paediatr Dent* 2010;20:276–282. [PubMed: 20536589]
31. Rwenyonyi CM, Kutesa A, Muwazi L, Okullo I, Kasangaki A, Kekitinwa A. Oral Manifestations in HIV/AIDS-Infected Children. *Eur J Dent* 2011;5:291–298. [PubMed: 21769270]
32. Patel K, Hernan MA, Williams PL, Seeger JD, McIntosh K, Van Dyke RB, et al. Long-term effectiveness of highly active antiretroviral therapy on the survival of children and adolescents with HIV infection: a 10-year follow-up study. *Clin Infect Dis* 2008;46:507–515. [PubMed: 18199042]

33. Neilan AM, Karalius B, Patel K, Van Dyke RB, Abzug MJ, Agwu AL, et al. Association of Risk of Viremia, Immunosuppression, Serious Clinical Events, and Mortality With Increasing Age in Perinatally Human Immunodeficiency Virus-Infected Youth. *JAMA Pediatr* 2017,171:450–460. [PubMed: 28346597]
34. Brady MT, Oleske JM, Williams PL, Elgie C, Mofenson LM, Dankner WM, et al. Declines in mortality rates and changes in causes of death in HIV-1-infected children during the HAART era. *J Acquir Immune Defic Syndr* 2010,53:86–94. [PubMed: 20035164]

Table 1.

a) History of antiretroviral therapy (ART) received by 209 perinatally HIV-infected (PHIV) youth participating in the Oral Health study in the AMP/PHACS¹; **b)** Current stable combination antiretroviral therapy (cART)² regimen (unchanged for past year prior to study entry)³ among 153 PHIV youth participating in the Oral Health study in the AMP/PHACS

a) History of ART Regimen (N=209):		Ever received	Age at first ART exposure	Age at first ART exposure Categorized:	
	N (%)	Median ⁴ (Q1, Q3)	Age groups (years)	N (%)	
cART ²	205 (98)	2.7 (0.9, 5.2)	< 2	82 (39)	
			2 to < 6	85 (41)	
			6+	38 (18)	
			Not exposed	4 (2)	
NRTIs ⁵	209 (100)	0.59 (0.2, 2.4)	< 2	145 (69)	
			2 to < 6	52 (25)	
			6+	12 (6)	
PIs ⁵	190 (91)	2.7 (1.1, 4.8)	< 2	73 (35)	
			2 to < 6	87 (42)	
			6+	30 (14)	
			Not exposed	19 (9)	
NNRTIs ⁵	140 (67)	4.6 (2.4, 10.3)	< 2	28 (13)	
			2 to < 6	62 (30)	
			6+	50 (24)	
			Not exposed	69 (33)	
IIs ⁵	57 (27)	14.7 (12.6, 17.2)	6+	57 (27)	
			Not exposed	152 (73)	
b) Stable cART(or no cART/ART) for 1 year prior to study entry (N=153) ³				N (%)	
With NNRTIs				43 (28)	
No NNRTIs				100 (65)	
Non cART or no ART				10 (7)	
With PIs				111 (73)	
No PIs				32 (21)	
Non cART or no ART				10 (7)	
With IIs				29 (19)	
No IIs				114 (75)	
Non cART or no ART				10 (7)	

¹ Adolescent Master Protocol (AMP) of the Pediatric HIV/AIDS Cohort Study (PHACS)

² cART was defined as any regimen containing at least three drugs from at least two drug classes.

³ Among 209 participants, 153 had a stable regimen (143 on cART, 10 not on cART/ART) in the past year prior to study entry

⁴ Median and 1st and 3rd quantiles (Q1, Q3) among those ever received.

⁵ NRTIs: Nucleoside-analog reverse transcriptase inhibitors; PIs: Protease inhibitors; NNRTIs: Non-nucleoside reverse transcriptase inhibitors; IIs: Integrase inhibitors (Note: only 10 (5%) ever received a fusion inhibitor in their regimen, and 144 (69%) ever received Didanosine (DDI))

Table 2.

Exposure to integrase inhibitors (II) as part of combination antiretroviral therapy (cART)¹ regimen by oral health outcomes among 143 perinatally HIV-infected youth participating in the Oral Health study in the AMP/PHACS² and having received the same cART for the past year prior to oral health study entry

	cART no II N=114	cART with II N=29	Total	P-value
DMFS³ score				
- Mean (SD)	7.0 (6.8)	12.2 (14.6)	8.1 (9.2)	0.14 ⁴
- Median (Q1, Q3)	5.5 (1, 11)	5 (3, 19)	5 (2, 12)	
DMFT³ score				
- Mean (SD)	4.7 (4.1)	6.9 (5.5)	5.1 (4.5)	0.07 ⁴
- Median (Q1, Q3)	4 (1, 7)	5 (3, 12)	4 (1, 8)	
DS³				
- Mean (SD)	1.9 (3.3)	5.6 (10.8)	2.7 (5.8)	0.002 ⁴
- Median (Q1, Q3)	1 (0, 2)	2 (1, 5)	1 (0, 3)	
DT³				
- Mean (SD)	1.6 (2.5)	3.6 (4.2)	2.0 (3.0)	<0.001 ⁴
- Median (Q1, Q3)	1 (0, 2)	2 (1, 5)	1 (0, 3)	
Unstimulated whole saliva flow rate (mL/mn)				
- Mean (SD)	0.8 (0.5)	0.8 (0.5)	0.8 (0.5)	0.90 ⁴
- Median (Q1, Q3)	0.6 (0.4, 1.0)	0.8 (0.4, 1.0)	0.7 (0.4, 1.0)	
Periodontal disease				
N (%)				
- None	22 (19)	7(24)	29 (20)	0.66 ⁵
- Gingivitis	49 (43)	10 (34)	59 (41)	
- Mild Periodontitis	18 (16)	7 (24)	25 (17)	
- Moderate Periodontitis	22 (19)	5 (17)	27 (19)	
- Missing	3 (3)	0	3 (2)	

¹ cART was defined as any regimen containing at least three drugs from at least two drug classes.

² Adolescent Master Protocol (AMP) of the Pediatric HIV/AIDS Cohort Study (PHACS)

³ DMFS/T: decayed-missing-filled-surface/teeth score; DS: number of decayed surfaces; DT: number of teeth with caries

⁴ Wilcoxon Rank Sum Test

⁵ Chi-Square Test

Table 3.

Zero-Inflated Negative Binomial (ZINB) multivariable model exploring the association between stable combination antiretroviral therapy (cART)¹ with II and number decayed teeth (DT), controlling for age, HIV RNA viral load, and CD4 cell count nadir, among 143 perinatally HIV-infected youth participating in the Oral Health study in the AMP/PHACS² and having received the same cART regimen for the past year prior to study entry

		Adj. Mean Ratio Estimate	95%CI	P-value
cART with II versus no II		1.86	1.05 – 3.30	0.03
Age at time of oral exam (years)	14-<17 vs < 14	1.45	0.68 – 3.10	0.34
	17-<19 vs < 14	1.49	0.71 – 3.12	0.29
	19+ vs < 14	1.41	0.65 – 3.08	0.39
% unsuppressed HIV Viral load ³		1.01 ⁴	1.00 – 1.02	0.15
CD4 nadir < 200 cells/mm		1.77	1.09 – 2.86	0.02

¹ cART was defined as any regimen containing at least three drugs from at least two drug classes.

² Adolescent Master Protocol (AMP) of the Pediatric HIV/AIDS Cohort Study (PHACS)

³ > 400 copies/mL (% among lifetime number of VL assays)

⁴ Per one percent increase in HIV RNA unsuppressed viral load

Table 4.

Age of first exposure to combination antiretroviral therapy (cART)¹ by dental and periodontal outcomes² among 205 perinatally HIV-infected youth ever exposed to cART participating in the Oral Health study in the AMP/PHACS³

	< 2 yrs N = 82	2 to 5 yrs N = 85	6+ yrs N = 38	P-value ⁴
DMFS score				
- Mean (SD)	5.5 (6.2)	9.5 (11.5)	13.8 (17.2)	0.001
- Median (Q1, Q3)	4 (0; 9)	6 (2; 13)	8.5 (4; 19)	
DMFT score				
- Mean (SD)	3.9 (4.0)	5.6 (5.0)	7.5 (5.8)	< 0.001
- Median (Q1, Q3)	3 (0; 6)	4 (2; 8)	6 (4; 11)	
DS				
- Mean (SD)	2.3 (3.7)	3.3 (7.8)	4.4 (8.2)	0.27
- Median (Q1, Q3)	1 (0; 3)	1 (0; 3)	1.5 (0; 5)	
DT				
- Mean (SD)	1.9 (2.7)	2.2 (3.7)	3.0 (4.0)	0.20
- Median (Q1, Q3)	1 (0; 3)	1 (0; 2)	1.5 (0; 5)	
Teeth with 2-6 BOP sites				
- Mean (SD)	7.4 (7.2)	9.9 (7.0)	11.3 (8.6)	0.01
- Median (Q1, Q3)	5 (2; 11)	8 (4.5; 15)	11.5 (3; 18)	

¹ cART was defined as any regimen containing at least three drugs from at least two drug classes.

² DMFS/T: decayed-missing-filled-surface/teeth score; DS: number of decayed surfaces; DT: number of teeth with caries; BOP: bleeding on probing

³ Adolescent Master Protocol (AMP) of the Pediatric HIV/AIDS Cohort Study (PHACS); Note: 4 participants were not exposed to cART

⁴ Kruskal-Wallis Test

Table 5.

Zero-Inflated Negative Binomial (ZINB) multivariable model exploring the association between (a) age at first protease inhibitor (PI) and decayed-missing-filled-teeth (DMFT) score (n=204), and (b) age at first combination antiretroviral therapy (cART)¹ and number of teeth with bleeding on probing (BOP), 2-6 sites (n=200); both models controlling for age at oral health study entry, HIV viral load and CD4 cell count nadir, among perinatally HIV-infected youth participating in the Oral Health study in the AMP/PHACS²

(a) DMFT		Adj. Ratio Estimate	95%CI	P-value
<i>Age (years) at First PI</i>	< 2 vs 6+	0.58	0.40 – 0.85	0.006
	2-<6 vs 6+	0.69	0.50 – 0.95	0.03
<i>Age (years) at time of oral exam</i>	14-<17 vs < 14	0.98	0.66 – 1.47	0.94
	17-<19 vs < 14	1.14	0.73 – 1.78	0.56
	19+ vs < 14	1.21	0.76 – 1.94	0.43
% RNA > 400 copies/mL ³		1.004 ⁴	1.000 – 1.008	0.06
CD4 nadir < 200 cells/mm		1.10	0.86 – 1.41	0.45
(b) Teeth with 2-6 BOP sites		Adj. Ratio Estimate	95%CI	P-value
<i>Age (years) at First cART</i>	< 2 vs 6+	0.51	0.35 - 0.75	<0.001
	2-<6 vs 6+	0.79	0.58 – 1.09	0.15
<i>Age (years) at time of oral exam</i>	14-<17 vs < 14	0.87	0.61 – 1.24	0.44
	17-<19 vs < 14	0.67	0.44 – 1.01	0.06
	19+ vs < 14	0.66	0.42 – 1.04	0.07
% RNA > 400 copies/mL ³		1.004 ⁴	1.00 – 1.01	0.79
CD4 nadir < 200 cells/mm		1.01	0.78 – 1.29	0.97

¹ cART was defined as any regimen containing at least three drugs from at least two drug classes.

² Adolescent Master Protocol (AMP) of the Pediatric HIV/AIDS Cohort Study (PHACS)

³ > 400 copies/mL (% among lifetime number of VL assays)

⁴ Per one percent increase in HIV RNA unsuppressed viral load