Cochrane
Library

Cochrane Database of Systematic Reviews

Use of hyaluronidase as an adjunct to local anaesthetic eye blocks to reduce intraoperative pain in adults (Review)

Rüsch H, Aravindh K, Bunce C, Bokre D

Rüsch H, Aravindh K, Bunce C, Bokre D.

Use of hyaluronidase as an adjunct to local anaesthetic eye blocks to reduce intraoperative pain in adults.

Cochrane Database of Systematic Reviews 2018, Issue 3. Art. No.: CD010368.

DOI: [10.1002/14651858.CD010368.pub2](https://doi.org/10.1002/14651858.CD010368.pub2).

www.cochranelibrary.com

Use of hyaluronidase as an adjunct to local anaesthetic eye blocks to reduce intraoperative pain in adults
(Review)

Copyright © 2018 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

WILEY

TABLE OF CONTENTS

ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	4
BACKGROUND	6
OBJECTIVES	7
METHODS	7
RESULTS	9
Figure 1.	10
Figure 2.	12
Figure 3.	13
Figure 4.	14
Figure 5.	14
DISCUSSION	15
AUTHORS' CONCLUSIONS	16
ACKNOWLEDGEMENTS	16
REFERENCES	17
CHARACTERISTICS OF STUDIES	19
DATA AND ANALYSES	26
Analysis 1.1. Comparison 1 Hyaluronidase versus control, Outcome 1 Intraoperative pain (reported continuous).	27
Analysis 1.2. Comparison 1 Hyaluronidase versus control, Outcome 2 Intraoperative pain (reported dichotomous).	27
APPENDICES	27
WHAT'S NEW	42
CONTRIBUTIONS OF AUTHORS	42
DECLARATIONS OF INTEREST	43
SOURCES OF SUPPORT	43
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	43
INDEX TERMS	44

[Intervention Review]

Use of hyaluronidase as an adjunct to local anaesthetic eye blocks to reduce intraoperative pain in adults

Heinrich Rüschén¹, Kavitha Aravinth¹, Catey Bunce², Desta Bokre³

¹Department of Anaesthesia, Moorfields Eye Hospital NHS Foundation Trust, London, UK. ²Department of Primary Care & Public Health Sciences, King's College London, London, UK. ³The Joint Library of Ophthalmology, Moorfields Eye Hospital & UCL Institute of Ophthalmology, London, UK

Contact: Heinrich Rüschén, Department of Anaesthesia, Moorfields Eye Hospital NHS Foundation Trust, 162 City Road, London, EC1V 2PD, UK. ruschen@gmail.com.

Editorial group: Cochrane Anaesthesia Group.

Publication status and date: Edited (no change to conclusions), published in Issue 10, 2018.

Citation: Rüschén H, Aravinth K, Bunce C, Bokre D. Use of hyaluronidase as an adjunct to local anaesthetic eye blocks to reduce intraoperative pain in adults. *Cochrane Database of Systematic Reviews* 2018, Issue 3. Art. No.: CD010368. DOI: [10.1002/14651858.CD010368.pub2](https://doi.org/10.1002/14651858.CD010368.pub2).

Copyright © 2018 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Hyaluronidase has been used over many decades as an adjunct to local anaesthetic solution to improve the speed of onset of eye blocks and to provide better akinesia and analgesia. With the evolution of modern eye surgery techniques, fast onset and akinesia are not essential requirements anymore. The assumption that the addition of hyaluronidase to local anaesthetic injections confers better analgesia for the patient needs to be examined. There has been no recent systematic review to provide evidence that hyaluronidase actually improves analgesia.

Objectives

To ascertain if adding hyaluronidase to local anaesthetic solutions for use in ophthalmic anaesthesia in adults results in a reduction of perceived pain during the operation and to assess harms, participant and surgical satisfaction, and economic impact.

Search methods

We carried out systematic searches in Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE, Embase, and four other databases in June 2017. We searched the trial registers at www.ISRCTN.com, ClinicalTrials.gov and www.clinicaltrialsregister.eu for relevant trials. We imposed no language restrictions.

Selection criteria

We included randomized controlled trials (RCTs) that evaluated the effect of hyaluronidase on pain experienced by adults during intraocular surgery using a rating scale.

Data collection and analysis

Two review authors (HR and KA) independently extracted data and assessed methodological quality using standard procedures as expected by Cochrane.

Main results

We included seven trials involving 500 participants that studied the effect of hyaluronidase on intraoperative pain. Four of the seven trials with 289 participants reported the primary outcome in a dichotomous manner, and we proceeded to meta-analyse the findings which showed a moderate heterogeneity that could not be explained ($I^2 = 41\%$). The pooled risk ratio (RR) for these four trials was 0.83 with

the 95% confidence interval ranging from 0.48 to 1.42. The reduction in intraoperative pain scores in the hyaluronidase group were not statistically significant. Among the three trials that reported the primary outcome in a continuous manner, the presence of missing data made it difficult to conduct a meta-analysis. To further explore the data, we imputed standard deviations for the other studies from another included RCT (Sedghipour 2012). However, this resulted in substantial heterogeneity between study estimates ($I^2 = 76\%$). The lack of reported relevant data in two of the three remaining trials made it difficult to assess the direction of effect in a clinical setting.

Overall, there was no statistical difference regarding the intraoperative reduction of pain scores between the hyaluronidase and control group. All seven included trials had a low risk of bias.

According to GRADE, we found the quality of evidence was low and downgraded the trials for serious risk of inconsistency and imprecision. Therefore, the results should be analysed with caution.

Participant satisfaction scores were significantly higher in the hyaluronidase group in two high quality trials with 122 participants. Surgical satisfaction was also superior in two of three high quality trials involving 141 participants. According to GRADE, the quality of evidence was moderate for participant and surgical satisfaction as the trials were downgraded for imprecision due to the small sample sizes. The risk of bias in these trials was low.

There was no reported harm due to the addition of hyaluronidase in any of the studies. No study reported on the cost of hyaluronidase in the context of eye surgery.

Authors' conclusions

The effects of adding hyaluronidase to local anaesthetic fluid on pain outcomes in people undergoing eye surgery are uncertain due to the low quality of evidence available. A well designed RCT is required to address inconsistency and imprecision among the studies and to determine the benefit of hyaluronidase to improve analgesia during eye surgery. Participant and surgical satisfaction is higher with hyaluronidase compared to the control groups, as demonstrated in moderate quality studies. There was no harm attributed to the use of hyaluronidase in any of the studies. Considering that harm was only rarely defined as an outcome measure, and the overall small number of participants, conclusions cannot be drawn about the incidence of harmful effects of hyaluronidase. None of the studies undertook cost calculations with regards to use of hyaluronidase in local anaesthetic eye blocks.

PLAIN LANGUAGE SUMMARY

Addition of hyaluronidase to local anaesthetic eye blocks to reduce pain during eye surgery in adults.

Review question

We reviewed the evidence on the effectiveness of adding hyaluronidase to local anaesthetic eye block solutions (a numbing medicine injected into the eye to block nerves) to reduce pain and increase participant and surgical satisfaction during eye surgery in adults. We also looked for reports on side effects and cost.

Background

Hyaluronidase is an enzyme (a protein that regulates a chemical reaction in the body) that helps the spread of local anaesthetic through the tissues around the eye. It is widely used as an additive to local anaesthetic eye blocks to give more rapid onset of anaesthesia and reduce or block movement of the eye (called akinesia). With modern eye surgery techniques, fast onset and akinesia are no longer essential requirements, and often surgery can be undertaken pain-free with topical (on the surface of the eye) anaesthesia alone. Hyaluronidase has been associated with infrequent side effects. Therefore, the use of hyaluronidase needs to be justified, which was the aim of this review.

Search date

The review is current to 30 June 2017.

Study characteristics

We included seven randomized controlled trials (clinical studies where people are randomly put into one of two or more treatment groups) in our review. These involved 500 adults undergoing eye surgery under local anaesthesia. We looked at any additional effect of adding hyaluronidase to local anaesthetic on the pain experienced during eye surgery. We also looked at participant and surgical satisfaction scores and if any harms were reported after using hyaluronidase in the injection solution. None of the studies reported on costs.

Key results

Of the seven included trials, we pooled the results of four trials (289 participants) as the results were reported in a similar manner. They found that addition of hyaluronidase did not significantly reduce pain during surgery. Among the three remaining trials (211 participants) lack of data reporting in two trials made it difficult to pool the results. The overall result of looking at all these trials together suggests there was no significant reduction of pain with using hyaluronidase in eye nerve blocks.

We found moderate quality evidence from two trials (122 participants) to suggest that addition of hyaluronidase increased participant satisfaction scores. Three studies involving 141 participants looked at surgical satisfaction, which was reported as superior with hyaluronidase in the two larger studies and not significantly different in one small study (19 participants). None of the included studies reported any harmful effects of hyaluronidase.

Quality of evidence

The included trials that reported on pain during surgery were at low risk for bias. The overall quality of evidence was low because of variations in the effect on pain reduction. We contacted all trial authors to request more information on the trials, but the data were not available.

Moderate quality studies reported greater participant and surgical satisfaction with hyaluronidase.

Analgesia alone does not take into account the full spectrum of the beneficial effects of hyaluronidase. Patient comfort with the eye surgery is also likely to be improved by a speedy onset and reduced eye movements due to hyaluronidase.

SUMMARY OF FINDINGS

Summary of findings for the main comparison. Use of hyaluronidase as an adjunct to local anaesthetic eye blocks to reduce intraoperative pain in adults

Use of hyaluronidase as an adjunct to local anaesthetic eye blocks to reduce intraoperative pain in adults

Patients or population: adults (aged ≥ 18 years) undergoing ophthalmic surgery under local anaesthetic eye blocks.

Setting: hospitals in the UK (4), Germany (1), Brazil (1) and Iran (1).

Intervention: local anaesthetic eye blocks containing hyaluronidase.

Comparison: local anaesthetic eye blocks containing no hyaluronidase.

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	N° of participants (studies)	Quality of the evidence (GRADE)	Comments
	Risk with no hyaluronidase	Risk with hyaluronidase				
Intraoperative pain (reported dichotomous) assessed with: analogue rating scales No follow-up - measured on day of surgery.	301 per 1000	250 per 1000 (145 to 428)	RR 0.83 (0.48 to 1.42)	289 (4 RCTs)	⊕⊕⊕⊖ Low ^{1,2}	-
Intraoperative pain (reported continuous) assessed with: analogue rating scales No follow-up - measured on day of surgery.	3 trials looked at effect of hyaluronidase on reduction of intraoperative pain measured by rating scales. Results were reported as continuous data. 2 studies did not provide the SMD, which measures the effect in a clinical setting, the results could not be meta-analysed and hence were reported narratively (Khandwala 2008; Rowley 2000). Among the 3 trials covering 211 participants (Khandwala 2008: Mean difference 0.70; Rowley 2000: Mean difference 0.31; Sedghipour 2012: Mean difference -1.10), only the Sedghipour study with 42 participants, which is a high quality study, showed a statistically significant (at the 5 % level) reduction in pain in the hyaluronidase group (P = 0.04). The remaining 2 studies with 169 participants showed no statistically significant (at the 5 % level) reduction of pain intraoperatively with hyaluronidase (Khandwala 2008: P = 0.5; Rowley 2000: n.s). These studies were also of high quality and low risk of bias. Khandwala and colleagues had an unclear attrition bias as 1/10 participants in the treatment group was dropped after randomization with no clear explanation.		-	211 (3 RCTs)	⊕⊕⊕⊖ Low ³	-
Incidence of harm	None of the studies reported harms in relation to hyaluronidase.		-	(0 studies)	-	-

Participant satisfaction assessed with: scoring system No follow-up - measured on day of surgery.	Significantly better satisfaction in these well designed studies with low risk of bias (Remy 2008; Sedghipour 2012). The studies included 122 participants and showed higher satisfaction scores in the treatment group ($P < 0.05$).	-	122 (2 RCTs)	⊕⊕⊕⊖ Moderate ⁴	-
Surgical satisfaction assessed with: scoring system No follow-up - measured on the day of surgery.	Surgical satisfaction was reportedly superior with hyaluronidase in the larger 2 studies (Remy 2008: $P < 0.001$; Sedghipour 2012: $P = 0.02$) and not significantly different in 1 small study (Khandwala 2008: $P = 0.96$).	-	141 (3 RCTs)	⊕⊕⊕⊖ Moderate ⁴	-
Economic outcomes or cost calculations	None of the included studies reported economic outcomes or cost calculations	-	(0 RCTs)	-	-

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **n.s.:** not statistically significant; **RCT:** randomized controlled trial; **RR:** risk ratio; **SMD:** standardized mean difference.

GRADE Working Group grades of evidence

High quality: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate quality: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low quality: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low quality: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

¹Downgraded one level due to marked heterogeneity with a calculated $I^2 > 50\%$.

²Downgraded one level for imprecision due to wide 95% confidence intervals, reflecting uncertainty in the direction of effect estimate.

³Downgraded one level for imprecision and inconsistency in measurement, lack of data and small sample size.

⁴Downgraded one level because of imprecision secondary to small sample size.

BACKGROUND

Description of the condition

Local anaesthesia for ophthalmic surgery can be provided by regional injection block or topical anaesthesia alone. The most frequently used anaesthetic injections are retrobulbar, peribulbar and sub-Tenon's block.

During a local anaesthetic injection for ophthalmic surgery, the objective is to deliver local anaesthetic fluid to the sensory and motor nerve fibres in the orbit. There is a large variation of techniques, anaesthetic mixtures and instruments in use to achieve this. Some of these techniques have been compared in Cochrane Reviews; for example, peribulbar versus retrobulbar block ([Alhassan 2015](#)), and topical anaesthesia alone versus sub-Tenon's block ([Guay 2015](#)). [Schein 2000](#) undertook a comprehensive systematic review of anaesthetic interventions.

Unfortunately, there is always a proportion of blocks that fail to provide adequate analgesia or akinesia. To improve the quality of the anaesthetic block, various adjuncts to the local anaesthetic fluid have been introduced.

Description of the intervention

[Atkinson 1949](#) first described the addition of hyaluronidase to local anaesthetic fluid with the intention of improving the speed of onset of analgesia and akinesia. Subsequently, hyaluronidase has commonly been added to local anaesthetic injection fluid for this purpose.

How the intervention might work

For any local anaesthetic block to work, the local anaesthetic fluid needs to spread through the orbital cavity to reach the relevant motor and sensor fibres. A complex system of connective tissue membranes divides the orbital space, thereby impeding the spread of local anaesthetic fluid ([Koornneef 1988](#)).

[Buhren 2016](#) described the molecular mechanisms of how hyaluronidase spreads through this connective tissue barrier, in the most recent review on this topic. The main components of connective tissue are the fibrous proteins collagen and elastin as well as proteoglycans located in the extracellular matrix. Glycosamine-glycans attach to proteoglycans in a characteristic manner giving the connective tissue its viscoelastic properties. The most common glycosamine-glycan is hyaluronic acid, it is a linear glycosaminoglycan disaccharide composed of alternating units of N-acetyl-D-glucosamine and D-glucuronic acid via alternating $\beta(1-4)$ and $\beta(1-3)$ glycosidic bonds. Hyaluronidase (hyaluronoglucosaminidase) is an enzyme that facilitates the spread of local anaesthetic fluid through connective tissue by degrading hyaluronic acid into smaller fragments and hydrolyzing the disaccharides at hexosaminidic $\beta(1-4)$ linkages.

[Meyer 1934](#) first extracted hyaluronidase. Preparations contain purified ovine testicular hyaluronidase as a dehydrated sterilized solid for reconstitution before use. Brand names of animal-derived hyaluronidase include Hydase, Vitrase, Amphadase, Wydase and Hyalase. Apart from a preparation of ovine testicular hyaluronidase, a recombinant human hyaluronidase is also available as Hylenex. It is produced by genetically engineered Chinese hamster ovary cells containing a DNA plasmid encoding for a soluble fragment

of human hyaluronidase ([Hylenex® Prescribing Information 2016](#)). The exact chemical structure of this enzyme is unknown. The approximate molecular weight is 61,000 daltons ([Borders 1968](#)).

Hyaluronidase also alters the pH of a local anaesthetic due to the presence of phosphate buffers within the preparation. The pH of plain bupivacaine solution is changed from 5.3 to 6.3 following the addition of hyaluronidase, and it may maintain local anaesthetic solubility during the process of alkalization ([Roberts 1993](#)). This alkalization may also explain any improved anaesthesia and akinesia.

A reduction in time to onset of surgical anaesthesia is considered desirable to facilitate patient throughput. The action of hyaluronidase may promote rapid onset of anaesthesia and akinesia. The minimum and maximum effective doses of hyaluronidase are unknown. The doses used range from 0.75 IU/mL to 300 IU/mL ([Dempsey 1997](#)).

Why it is important to do this review

We conducted this systematic review to explore the uncertainty about the benefits of using hyaluronidase in local anaesthetic mixtures to provide analgesia during eye surgery. There is considerable variation of practice, and the studies in this area show conflicting results.

Furthermore, the use of hyaluronidase increases the cost of the anaesthetic and has been associated with adverse allergic reactions in a small number of cases. For example, [Kempeneers 1992](#), described five people who developed an orbital pseudotumour as a complication of retrobulbar anaesthesia. Allergic reactions can range from local reactions to anaphylactic (systemic allergy) shock. When hyaluronidase is added to a local anaesthetic agent, the wider spread of the local anaesthetic solution also increases its absorption and removal in the bloodstream. This shortens the duration of action of the local anaesthetic and tends to increase the incidence of systemic reactions. Some hyaluronidase products contain bovine ingredients, and due to the theoretical concerns about transmissible spongiform encephalopathies, the World Health Organization (WHO) issued guidelines regarding the use of bovine materials in the manufacture of biological and pharmaceutical products ([WHO 2010](#)).

The absence of hyaluronidase in ophthalmic regional blockade has also been associated with adverse events. An interruption in hyaluronidase supply was associated with a cluster of postoperative diplopia (double vision) [Brown 1999](#). It was postulated that the absence of hyaluronidase caused the local anaesthetic to loculate in close proximity to the extraocular muscles and cause clinically significant myotoxicity. When hyaluronidase was unavailable once again in 2000, [Brown 2001](#) published a repeated cluster of diplopia cases. The omission of hyaluronidase in local anaesthesia fluid leads to clinically important rises in intraocular pressure. This is thought to be due to the decreased removal and dispersal of local anaesthetic fluid from the periocular compartment ([Dempsey 1997](#)).

Therefore, use of hyaluronidase must be justified, and data must be available for clinicians and patients to make an informed decision regarding the efficacy of hyaluronidase addition. The results of this

review should allow justification (or not) for the use of adjuvant hyaluronidase to improve the quality of anaesthesia and analgesia.

OBJECTIVES

To ascertain if adding hyaluronidase to local anaesthetic solutions for use in ophthalmic anaesthesia in adults results in a reduction of perceived pain during the operation and to assess harms, participant and surgical satisfaction and economic impact.

(See [Differences between protocol and review](#)).

METHODS

Criteria for considering studies for this review

Types of studies

We included:

1. Randomized controlled trials (RCTs) and quasi-randomized controlled clinical trials either published or unpublished;
2. Studies if they compared equal volumes and concentrations of local anaesthetic with and without adjuvant hyaluronidase administered with the injection;
3. Studies when other adjuvants such as adrenaline were used, only if the adjuvant was present in both the control and hyaluronidase intervention;

Types of participants

We included:

1. Adults (aged 18 years and older) presenting for ophthalmic surgery under ophthalmic anaesthetic block;
2. Participants receiving sub-Tenon's, peribulbar, retrobulbar or other types of local anaesthetic;
3. Participants receiving sedation but documented this fact;

We excluded:

1. Participants receiving adnexal surgery and any other eye surgery that was not intraocular;
2. Participants who received general anaesthesia;

Types of interventions

Ophthalmic local anaesthetic blocks comparing adjuvant hyaluronidase to an otherwise equal anaesthetic and surgery without hyaluronidase.

We considered any dose of hyaluronidase in the intervention group and any dose or type of local anaesthetic agent.

We included studies that used any number of injections to anaesthetise the eye if the number of injections was equal in the treatment and control groups.

Types of outcome measures

Primary outcomes

1. Intraoperative pain, as measured by analogue rating scales. We excluded studies that reported pain, but did not measure pain formally using analogue rating scales, as they did not provide sufficiently useful information on the outcome. We excluded

studies if they reported that 'supplementary injections' were primarily given to achieve akinesia, for example, if a certain immobility score was not reached.

(See [Differences between protocol and review](#)).

Secondary outcomes

1. Incidence of harm (reported as a narrative).
2. Participant and surgical satisfaction, as documented by scoring systems.
3. Economic outcomes or cost calculations (reported as a narrative).

Search methods for identification of studies

Electronic searches

We carried out systematic searches in:

1. The Cochrane Central Register of Controlled Trials(CENTRAL, 2007 Issue 6; [Appendix 1](#));
2. Ovid MEDLINE (1946 to 30 June 2017; [Appendix 2](#));
3. Ovid Embase (1947 to 30 June 2017; [Appendix 3](#));
4. Web of Science (1900 to 30 June 2017; [Appendix 4](#));
5. Scopus (1823 to 30 June 2017; [Appendix 5](#));
6. CINAHL Plus (EBSCOhost, 1937 to 30 June 2017; [Appendix 6](#));
7. LILACS (1982 to 30 June 2017; [Appendix 7](#));

We broke down our research question into four key searchable concepts: " Eye," "Surgery," " Local Anaesthesia" and "Hyaluronidase".This strategy ensured that we retrieved studies on eye surgery where local anaesthesia was applied along with hyaluronidase. We conducted searches for each concept using free text terms and MeSH terms wherever possible. When we carried out free text searches, we applied synonyms, derivative forms and singular/plural forms for each concept. Detailed search steps are documented in the 'Appendices'.

We applied no language restrictions.

Searching other resources

We searched the reference lists of all eligible trials and reviews and used any trials that fit the inclusion criteria.

We searched the registers at www.controlled-trials.com, www.ISRCTN.com and www.clinicaltrialsregister.eu for relevant trials.

We contacted specialists in the field, authors of the included trials and pharmaceutical manufacturers for any unpublished data.

The search of these other resources was completed by 30 June 2017.

Data collection and analysis

Selection of studies

Two review authors (KA and HR) independently reviewed the trials identified from the search strategy, removed duplicates and documented the reason for each trial being excluded (see [Characteristics of excluded studies](#) table). We resolved any disagreements with the studies by input from a third review author

CB). We presented information regarding methods, participants, setting, interventions and outcomes in the [Characteristics of included studies](#) table. Where studies had multiple publications, we planned to collate the reports of the same study so that each study, rather than each report, was the unit of interest for the review, and such studies had a single identifier with multiple references. (See [Differences between protocol and review](#)).

Data extraction and management

Two review authors (KA and HR) independently extracted and collected data on a paper form. After initial piloting this form was assessed and agreed for usability. A copy of this form is in [Appendix 8](#). We (KA and HR) resolved any discrepancies in data extracted by discussion with a third review author (CB) as a final arbiter. In the case of additional information being required, HR or KA contacted the authors of the relevant trial.

Assessment of risk of bias in included studies

To assess the risk of bias, two review authors (KA and HR) independently assessed the studies included in the review according to the criteria described by [Higgins 2011](#). We assessed the following aspects as being at either 'low risk', 'high risk' or 'unclear risk' of bias. We assessed the risk of bias for the following components of each trial.

1. Random sequence generation (selection bias).
2. Allocation concealment (selection bias).
3. Masking of participants and personnel (performance bias).
4. Masking of outcome assessment (detection bias).
5. Incomplete outcome data (attrition bias).
6. Selective reporting (reporting bias).
7. Other bias.

We included a 'Risk of bias' table as part of the [Characteristics of included studies](#) table based on Cochrane's tool for assessing the risk of bias, from the *Cochrane Handbook for Systematic Reviews of Interventions* (Chapter 8: [Higgins 2011](#)). See [Appendix 8](#) (data collection form) and [Appendix 10](#) ('Risk of bias' table).

Measures of treatment effect

Intraoperative pain

We noted intraoperative pain measured using visual analogue scales (VAS) or verbal rating scales (VRS) when available and interpreted them as continuous data. We used the greatest intraoperative score reported. We planned to use the standardized mean difference (SMD) as an effect measure and 95% confidence intervals (CIs) to allow for the fact that different studies might have used different scales. However, because the only studies detected used the same scaling method, there was no need to use the SMD. If authors documented non-normality of their data, we extracted medians and interquartile ranges and collated this information. (See [Differences between protocol and review](#)).

In the case of absolute numbers of participants experiencing pain where a rating scale was used but reported in a dichotomous manner (data as pain or no pain), we used risk ratios (RR) with 95% CI as a measure of effect. We collated this information and reported it. While there was evidence of moderate heterogeneity ($I^2 = 41\%$) we presented a meta-analysis but urge caution interpreting this.

To avoid multiplicity, we restricted meta-analyses to the primary outcomes but this would not be at the cost of presenting the totality of evidence should there be more RCTs providing information with regards to adverse effects.

Incidence of harm

We would have reported adverse events due to the use of hyaluronidase (e.g. allergic reactions) as a narrative, but as expected, there were no reports of such adverse events, probably because of their rarity and the relatively small number of participants in each trial.

Participant and surgical satisfaction

We reported participant and surgical satisfaction scores narratively. We would also have noted as narrative if other validated tools had been used.

Economic outcomes or cost calculation

We planned to report economic outcomes or cost calculations narratively.

Unit of analysis issues

We anticipated that most trials would involve one eye per participant, and even if both eyes were included, our outcomes were primarily measured at the participant rather than eye level. It was very unlikely that both eyes were operated on simultaneously with different anaesthetic procedures. (See [Differences between protocol and review](#)).

Dealing with missing data

We contacted authors and asked them to provide missing data. We imposed a time limit of two months and follow-up on one occasion. Irrespective of the type of data, we reported dropout rates in the 'Risk of bias' tables within the [Characteristics of included studies](#) table and noted whether or not authors had compared characteristics of participants who had complete data sets against those that did not. We investigated studies that had missing data, whether or not imputing for missing cases impacted greatly on the interpretation of findings. We attempted to impute the data, but the values obtained would not be consistently reflected within the trials due to the variation in sample size.

Assessment of heterogeneity

We assessed all studies for clinical and methodological heterogeneity. We examined the I^2 statistic and its 95% CI to assess inconsistency between studies as recommended by the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011](#)). We used the thresholds advised by [Higgins 2011](#), for the interpretation of the I^2 statistic. We found substantial inconsistency in our primary outcome when assessed as a continuous score ($I^2 = 76\%$) but less when assessed as a dichotomous measure ($I^2 = 41\%$). We attempted to investigate causes for this by exploring factors such as the type and duration of surgery, anaesthetic intervention (hyaluronidase dosage, type and volume of anaesthetic fluid) but the number of studies contributing to the meta-analysis was small. Despite the heterogeneity, we presented a meta-analysed outcome for the dichotomized outcome but urge caution in its interpretation.

Assessment of reporting biases

We assessed publication bias and small study effects in a qualitative manner using a funnel plot. We planned to test for funnel plot asymmetry if there had been a meta-analysis with more than 10 studies included.

Data synthesis

We performed the analysis using Review Manager 5 ([RevMan 2014](#)).

While we found some evidence of heterogeneity (for the dichotomous outcome), we meta-analysed results using a random-effects model as per our original intentions. A random-effects model was chosen because we believe that each trial estimates an intervention effect that follows a distribution across studies.

Subgroup analysis and investigation of heterogeneity

We planned no subgroup analyses.

Sensitivity analysis

We carried out sensitivity analyses to explore the robustness of the results to key methodological decisions that we made in our review. We examined whether or not excluding studies at risk of bias impacted on our findings, and since we had missing data, we attempted to impute and examine whether or not analysing intention-to-treat data differed considerably from the available case meta-analysis.

GRADE assessment of quality of evidence

We adopted the GRADE system postprotocol ([Rüschén 2013](#)) to rate the quality of evidence for each outcome ([Guyatt 2011](#)). GRADE

assessment classifies the quality of evidence into four categories; high, moderate, low and very low. The overall assessment considers the study design, risk of bias, imprecision, inconsistency, indirectness, publication bias, large effect size, dose-response effect and presence of confounding factors to rate the evidence. We applied the principles of GRADE to assess the quality of evidence specific to each outcome in our review.

1. Intraoperative pain, as measured by analogue rating scales.
2. Incidence of harm.
3. Participant and surgical satisfaction, as documented by scoring systems.
4. Economic outcomes or cost calculations.

The GRADE software from [GRADEpro GDT](#) generated the 'Summary of findings' table. The GRADE approach ensures the confidence which one can have in the estimate of effect from the outcomes being assessed in the included studies.

(See [Differences between protocol and review](#)).

RESULTS

Description of studies

See [Characteristics of included studies](#) and [Characteristics of excluded studies](#) tables.

Results of the search

See [Figure 1](#) for the study flow diagram.

Figure 1. Study flow diagram.

We identified 942 references after removal of duplicates. Two review authors (HR and KA) independently read and analysed the abstracts of all references and if needed, full papers. We excluded 779 references as clearly irrelevant to the review. We obtained the full papers for the remaining 163 references and again analysed them independently. We identified 18 trials, and 11 trials were further excluded for reasons documented in the '[Characteristics of excluded studies](#)' table. The review includes seven trials.

Included studies

Design

We included seven RCTs published from 1995 to 2012 with 500 participants ([Bowman 1997](#); [Brydon 1995](#); [Khandwala 2008](#); [Remy 2008](#); [Rowley 2000](#); [Sedghipour 2012](#); [Shiroma 2002](#)). One study was published in Portuguese and translated into English ([Shiroma 2002](#)).

Characteristics of study population

The review included 500 participants. The participants were adults (aged 18 years or older) presenting for ophthalmic surgery undergoing a retrobulbar, peribulbar or sub-Tenon block. The mean age in the studies ranged from 66 to 77 years. Studies were balanced with regards to gender.

Setting

Four studies were conducted in the UK ([Bowman 1997](#); [Brydon 1995](#); [Khandwala 2008](#); [Rowley 2000](#)). The remaining three studies were based in Germany ([Remy 2008](#)), Brazil ([Shiroma 2002](#)), and Iran ([Sedghipour 2012](#)).

Intervention

The seven trials studied the effect of adding hyaluronidase to a local anaesthetic mixture with the primary outcome measure of reduction of intraoperative pain. The participants were divided into a treatment group (hyaluronidase) and a control group (no hyaluronidase). The doses of hyaluronidase used ranged from 15 IU/mL to 150 IU/mL.

All seven trials assessed pain objectively using either the VAS or the VRS and compared a group with hyaluronidase to a group without hyaluronidase ([Bowman 1997](#); [Brydon 1995](#); [Khandwala 2008](#); [Remy 2008](#); [Rowley 2000](#); [Sedghipour 2012](#); [Shiroma 2002](#)).

One trial included three arms in their study looking at effects of no hyaluronidase and effects of hyaluronidase at 50 IU/mL and 150 IU/mL ([Brydon 1995](#)). We combined the results of the groups with different doses of hyaluronidase and compared them with the no hyaluronidase group.

Funding sources

Two trials reported no conflict of interest and received no funding support for the trials ([Khandwala 2008](#); [Remy 2008](#)). There was no clear documentation of reported conflict of interest or funding support from the remaining five trials ([Bowman 1997](#); [Brydon 1995](#); [Rowley 2000](#); [Sedghipour 2012](#); [Shiroma 2002](#)).

We attempted to contact the authors of all the trials for additional data but received no clarification.

For more details about the included trials, see the [Characteristics of included studies](#) table.

Excluded studies

We excluded 11 studies after analysis ([Berg 2001](#); [Crawford 1994](#); [Guise 1999](#); [House 1991](#); [Johansen 1993](#); [Lange 1989](#); [Moharib 2002](#); [Morsman 1992](#); [Ramanathan 1999](#); [Sarvela 1992](#); [Soares 2002](#)).

Three studies reported that "supplementary injections" were primarily given to achieve akinesia, for example, if a certain immobility score was not reached. Pain during surgery was not assessed or reported specifically. ([Crawford 1994](#); [House 1991](#); [Soares 2002](#)).

One study did not mask the relevant part of the trial ([Morsman 1992](#)).

Four studies did not assess pain or discomfort using a rating scale, and the inclusion criteria were ultimately not met ([Berg 2001](#); [Guise 1999](#); [Johansen 1993](#); [Moharib 2002](#)). We excluded these studies because they did not measure the outcome measure of "pain by rating scale". They reported an unstructured description of pain. This is unlikely to provide useful information about the levels of pain during the operation.

One study was published as a poster presentation ([Ramanathan 1999](#)). We contacted the author for more details about randomization, masking and results but none was made available.

We excluded one study because there was no randomization, this was not immediately obvious ([Lange 1989](#)).

We excluded one study because the relevant part of the study was not randomized and only compared different concentrations of hyaluronidase ([Sarvela 1992](#)).

See [Characteristics of excluded studies](#) table.

Awaiting classification

We found no studies awaiting classification.

Ongoing studies

We identified no ongoing studies.

Risk of bias in included studies

We judged the quality of studies according to the methods described in the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011](#)). Overall, the selected studies were of low risk; however, there was a lack of concise information in the methodology of randomization and withdrawal description. Please refer to [Figure 2](#) and [Figure 3](#) for a summary of risk of bias assessment for the selected studies and a 'Risk of bias' graph that represents the studies.

Figure 2. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)
Bowman 1997	?	?	+	+	+	+
Brydon 1995	?	?	+	+	?	+
Khandwala 2008	+	+	?	+	+	+
Remy 2008	?	+	+	+	+	+
Rowley 2000	+	+	+	+	+	+
Sedghipour 2012	+	+	+	+	+	+
Shiroma 2002	?	?	+	+	+	+

Figure 3. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.

Allocation

All seven included studies reported that allocation was randomized, but only three studies provided any detail about the method of random allocation (Khandwala 2008; Rowley 2000; Sedghipour 2012). These three studies were at low risk of allocation bias. The studies used random number tables, stratified lottery system or computer generated randomization. The remaining four studies were at unclear risk of bias.

With regard to allocation concealment, four of the seven studies used "coded syringes" (Khandwala 2008; Remy 2008; Rowley 2000; Sedghipour 2012). We classed these at low risk of bias, but an experienced operator might have recognized the formation of tiny bubbles in the mixture indicating hyaluronidase content.

Most studies implied that participants, personnel and assessors were unaware of the composition of the anaesthetic solution because the syringes were coded by an uninvolved third party, but exact details were rarely given.

In this context, only one study used a placebo control, with inactive Hyalase (Remy 2008).

Blinding

Five studies were classified as low risk for performance and detection bias (Bowman 1997; Khandwala 2008; Rowley 2000; Sedghipour 2012; Shiroma 2002). These studies had described double masking where neither the participant nor the caregiver was aware of the contents of the syringe. The outcome assessors were also masked, which further reduced the risk of bias by masking.

Incomplete outcome data

Withdrawal of participants after randomization was rarely reported in detail.

Six studies were at low risk as there was a description of no withdrawals; therefore, we had more confidence in the intention-to-treat analysis for these studies (Bowman 1997; Brydon 1995; Remy 2008; Rowley 2000; Sedghipour 2012; Shiroma 2002).

Of the included seven studies, one described the withdrawal of participants after randomization (Khandwala 2008). One participant was withdrawn from the treatment group due to incomplete data. No further details were given. Considering the low number of participants in this trial (10 in each of two arms) the exclusion may represent bias.

Selective reporting

We found all seven studies at low risk of reporting bias. Pain was stated as an outcome measure at the start of the trials. We did not attempt to obtain research protocols (Bowman 1997; Brydon 1995; Khandwala 2008; Remy 2008; Rowley 2000; Sedghipour 2012; Shiroma 2002).

Other potential sources of bias

We found no other potential sources of bias.

Effects of interventions

See: [Summary of findings for the main comparison Use of hyaluronidase as an adjunct to local anaesthetic eye blocks to reduce intraoperative pain in adults](#)

Primary outcomes

1. Intraoperative pain, as measured by analogue rating scales

Seven trials with 500 participants looked at the effect of hyaluronidase on the reduction of intraoperative pain. Four of these trials with 289 participants reported intraoperative pain in a dichotomous manner (Bowman 1997; RR 0.82, 95% CI 0.38 to 1.75; Brydon 1995; RR 1.00, 95% CI 0.20 to 5.00; Remy 2008; RR 0.47, 95% CI 0.24 to 0.92; Shiroma 2002; RR 1.45, 95% CI 0.70 to 3.00). We calculated the (RR) as a measure of effect. The I^2 statistic was 41%, which represents moderate heterogeneity. We proceeded to meta-analyse the results and found that the pooled RR was 0.83 (95% CI 0.48 to 1.42). Therefore, there was no statistically significant reduction of pain scores in the hyaluronidase group.

Three studies involving 211 participants reported pain objectively using rating scales and presented continuous data (Khandwala 2008; Rowley 2000; Sedghipour 2012). Sedghipour 2012 described

42 participants in a high quality study and provided SDs for their data. They found a significant reduction of pain in the hyaluronidase group ($P = 0.04$). The other two trials did not report SDs (Khandwala 2008; Rowley 2000). We interchanged the SDs

from Sedghipour 2012, but this resulted in significant heterogeneity among the studies ($I^2 = 76\%$), so we did not perform a meta-analysis. See Figure 4 and Figure 5.

Figure 4. Forest plot of comparison: 1 Hyaluronidase versus control, outcome: 1.1 Intraoperative pain (measured by analogue rating scales; reported continuous).

Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Incomplete outcome data (attrition bias)
- (D) Selective reporting (reporting bias)
- (E) Blinding of participants and personnel (performance bias)
- (F) Blinding of outcome assessment (detection bias)

Figure 5. Forest plot of comparison: 1 Hyaluronidase versus control, outcome: 1.2 Intraoperative pain (measured by analogue rating scales; reported dichotomous).

Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Incomplete outcome data (attrition bias)
- (D) Selective reporting (reporting bias)
- (E) Blinding of participants and personnel (performance bias)
- (F) Blinding of outcome assessment (detection bias)

Looking at the overall studies taking into account the results of the meta-analysis and the individual studies that reported the continuous outcome, there was no statistically significant reduction in intraoperative pain with hyaluronidase in the local anaesthetic mixture. However, this has to be interpreted with caution.

We adopted the GRADEpro method of analysing the quality of evidence and produced [Summary of findings for the main comparison](#). We found the quality of studies was low. We downgraded the quality of evidence due to concerns regarding inconsistency in the direction and magnitude of effect across the studies ($I^2 = 41\%$ and 76%). We looked at the individual studies for factors that could have contributed to the heterogeneity and found that there was no wide variability between characteristics of participants, interventions and outcome measures. We tried to establish if the heterogeneity was due to the dose of hyaluronidase, the volume of injection or number of participants, but there was insufficient data provided to enable a valid analysis. The level of

imprecision was another reason for downgrading the quality of evidence. Only Rowley 2000, provided a rationale for the selected sample size that would yield the specific effect measure.

Secondary outcomes

1. Incidence of harm

None of the included studies measured or reported the incidence of harm from hyaluronidase.

2. Participant and surgical satisfaction, as documented by scoring systems

Two studies analysing 122 participants looking at participant satisfaction reported that the investigator and participant assessment scores were significantly higher in the hyaluronidase group ($P < 0.05$) (Remy 2008; Sedghipour 2012). The studies assessed satisfaction in different ways so prohibiting meta-analysis. Remy 2008 used a five level VAS to assess participant efficacy

and tolerability at the end of surgery and at the final visit, while [Sedghipour 2012](#) captured data as a dichotomous 'satisfied' or 'unsatisfied'. Using the GRADE system to assess the quality of evidence, we found the studies were of moderate quality with low risk of bias.

We found three studies involving 141 participants that measured surgical satisfaction scores ([Khandwala 2008](#); [Remy 2008](#); [Sedghipour 2012](#)). [Sedghipour 2012](#) reported that surgical satisfaction with intraoperative anaesthesia was 85.7% in the hyaluronidase group compared to 52.5% in the control group ($P = 0.02$). [Remy 2008](#) reported a P value of less than 0.001 for surgical satisfaction in the hyaluronidase group. [Khandwala 2008](#) found no difference in the quality of the surgical field between groups ($P = 0.96$). These studies assessed surgeon satisfaction as they had assessed participant satisfaction while [Khandwala 2008](#) simply asked surgeons to rate surgical conditions on a VAS from 0 (worst) to 10 (best). Since each had assessed satisfaction using a different method, no meta-analysis was conducted. There was low risk of bias with regards to the method of randomization with [Sedghipour 2012](#), and there was incomplete outcome data reporting with [Khandwala 2008](#). Overall, we found the quality of evidence with these two studies to be moderate. See [Figure 2](#).

3. Economic outcomes or cost calculations

None of the included studies reported on the economic impact of using hyaluronidase.

DISCUSSION

The use of hyaluronidase in ophthalmic surgery remains a topic of debate. The perceived advantages of adding hyaluronidase include shortened time to onset of the block and improved akinesia, and as investigated in this review: analgesia. The apparent disadvantages of hyaluronidase include the additional cost and possible adverse reactions.

Most modern surgical techniques are no longer essentially dependent on akinesia. Anaesthesia itself is readily provided by eye blocks without hyaluronidase. Even topical anaesthesia is often deemed sufficient during routine cataract surgery. We systematically reviewed the literature on the benefits of hyaluronidase for analgesia in ophthalmic surgery.

Summary of main results

We reviewed evidence from seven RCTs involving 500 participants regarding the reduction of pain during intraocular surgery by adding hyaluronidase to the local anaesthetic fluid. We found that the reduction of intraoperative pain by hyaluronidase was not statistically significant. The quality of evidence was low. We assessed the literature in this field as having a low risk of bias, but we had concerns regarding heterogeneity across the studies. With regards to the outcomes of participant and surgeon satisfaction, the moderate quality studies show an advantage of using hyaluronidase. (See [Summary of findings for the main comparison](#)).

Overall completeness and applicability of evidence

We are confident that our search strategy obtained all available studies. The results of this review are applicable to all adults undergoing intraocular surgery who would want to make an

informed decision regarding the use of hyaluronidase as an adjunct in eye blocks to reduce intraoperative pain. We found that the use of hyaluronidase is beneficial in terms of participant and surgical satisfaction. Such benefit from using hyaluronidase was not statistically significant with regards to intraoperative reduction of pain.

We consider that most of the authors gave priority to akinesia as an outcome measure over analgesia, probably due to the perceived importance for the safe conduct of surgery. This priority has now receded as the majority of surgeons can carry out most operations without depending on fully established akinesia. Profound akinesia will still be necessary for more difficult operations and training situations. Hyaluronidase may be necessary to achieve akinesia in such situations.

Quality of the evidence

We found the overall quality of evidence to be low due to imprecision and inconsistency of the results. There was moderate heterogeneity ($I^2 = 41\%$). Therefore, we downgraded the quality of evidence by one level. We looked for possible causes for the variation such as sample size, dose of hyaluronidase or characteristics of participants but data were sparse.

We found the overall risk of bias in the studies to be low. However, there was an unclear risk with regards to methods of randomization and concealment in a few studies.

As for imprecision, failure to estimate the sample size needed to make an effect by six of the seven studies led to the quality of evidence to be downgraded by one level.

Potential biases in the review process

A potential bias arises from the narrow spectrum of the review question: "Does hyaluronidase improve pain control during eye surgery?" Hyaluronidase is used for a variety of indications. For example, if the speed of onset of anaesthesia is increased by hyaluronidase and the eye is much more akinetic at the beginning of the operation, participant and surgeon comfort will also likely be increased. Therefore, the narrow aspect of analgesia alone does not consider the full spectrum of beneficial effects from hyaluronidase.

This review found only a relatively small number of studies (seven) with a small number of participants (500). However, can systematic reviews with such sparse data be trusted ([Afshari 2017](#))? During this Cochrane Review, we adhered to all essential requirements such as publishing a protocol, incorporating risk of bias assessment, searching for unpublished data and many other review tools as laid out in the Cochrane framework ([Higgins 2011](#)).

According to our protocol, we excluded studies that did not assess pain in a structured manner (using rating scales). Because of this, we excluded unstructured assessments that may have shown results. We consider that results from these trials would have a high risk of reporting bias and therefore, would not produce a reliable, useful effect.

Lack of reported data also led to certain included trials being excluded from the meta-analysis. We were unable to obtain clarification about these issues from authors of the included studies.

Readers of this review may be interested in the incidence of adverse effects of hyaluronidase use. Adverse effects such as allergy to hyaluronidase are extremely rare. None of the trials we analysed reported any adverse events related to hyaluronidase, but we would like to highlight that our review would not have reliably captured the incidence of very rare adverse events due to an overall small number of participants.

We have acknowledged and taken into account the inherent methodological limitation of our systematic review and in our opinion addressed them adequately.

Agreements and disagreements with other studies or reviews

Our findings match that of the review of [Schein 2000](#). Their review reported the same problems with the available literature that we found. Those are;

1. very few studies reported data on the effect of hyaluronidase on pain;
2. high levels of inconsistency among the included studies.

Schein's review was written in 2000, and despite many additional studies having been published since, there is still no certainty on the effect of hyaluronidase use for analgesia.

AUTHORS' CONCLUSIONS

Implications for practice

The requirements for ophthalmic anaesthesia have changed considerably since the early 2000s. Nowadays, the majority of

routine cataract surgery can be conducted pain free under topical anaesthesia alone. Injection blocks are used to provide more profound analgesia for some people and during some operations. The effects of adding hyaluronidase to local anaesthetic fluid on pain outcomes in people undergoing eye surgery are uncertain due to the low quality of the available evidence.

Implications for research

To reach certainty on the question of hyaluronidase use for intraoperative analgesia, future studies should separate the various outcome parameters of speed of onset and akinesia from that of analgesia.

The importance of pain control is different for anterior and posterior segment eye surgery. This should be looked at in a well powered randomized controlled trial.

We found no studies that described the economical impact of using hyaluronidase. In the current climate of financial restrictions, this information would be very valuable, and any future study should incorporate this aspect.

ACKNOWLEDGEMENTS

We thank Jane Cracknell, Managing Editor of the Cochrane Anaesthesia, Critical and Emergency Care Group (ACE), and Karen Hovhannisyan (former ACE trials search co-ordinator). We would like to thank Andrew Smith (Content Editor), Marialena Trivella (Statistical Editor), Vibeke E Horstmann (Statistical Editor), Ana Licina, Jacques Ripart, Mahmoud B Alhassan (Peer Reviewers), Janet Wale (Consumer Editor), and Andrew Smith (Co-ordinating Editor) for their help and editorial advice during the preparation of the protocol ([Rüschén 2013](#)) and the systematic review.

REFERENCES

References to studies included in this review

Bowman 1997 {published data only (unpublished sought but not used)}

Bowman RJ, Newman DK, Richardson EC, Callear AB, Flanagan DW. Is hyaluronidase helpful for peribulbar anaesthesia?. *Eye* 1997;**11**(Pt 3):385-8. [PUBMED: 9373482]

Brydon 1995 {published data only (unpublished sought but not used)}

Brydon CW, Basler M, Kerr WJ. An evaluation of two concentrations of hyaluronidase for supplementation of peribulbar anaesthesia. *Anaesthesia* 1995;**50**(11):998-1000. [PUBMED: 8678264]

Khandwala 2008 {published data only (unpublished sought but not used)}

Khandwala M, Ahmed S, Goel S, Simmons IG, McLure HA. The effect of hyaluronidase on ultrasound-measured dispersal of local anaesthetic following sub-Tenon injection. *Eye* 2008;**22**(8):1065-8. [PUBMED: 17525774]

Remy 2008 {published data only (unpublished sought but not used)}

Remy M, Pinter F, Nentwich MM, Kampik A, Schönfeld CL. Efficacy and safety of hyaluronidase 75 IU as an adjuvant to mepivacaine for retrobulbar anesthesia in cataract surgery. *Journal of Cataract and Refractive Surgery* 2008;**34**(11):1966-9. [PUBMED: 19006746]

Rowley 2000 {published data only (unpublished sought but not used)}

Rowley SA, Hale JE, Finlay RD. Sub-Tenon's local anaesthesia: the effect of hyaluronidase. *British Journal of Ophthalmology* 2000;**84**(4):435-6. [PUBMED: 10729306]

Sedghipour 2012 {published data only (unpublished sought but not used)}

Sedghipour M, Mahdawifard A, Fouladi RF, Gharabaghi D, Rahbani M, Amiraslanzadeh G, et al. Hyaluronidase in subTenon's anaesthesia for phacoemulsification, a double blind randomized clinical trial. *International Journal of Ophthalmology* 2012;**5**(3):389-92. [PUBMED: 22773994]

Shiroma 2002 {published data only (unpublished sought but not used)}

Shiroma HF, Ferreira EM, Isaac DLC, Ghanem VC, Arieta CEL. A comparison of 1% ropivacaine efficacy when associated or not with hyaluronidase in peribulbar anaesthesia in cataract surgery [Comparação da eficácia da ropivacaína 1% quando associada ou não à hialuronidase na anestesia peribulbar para cirurgia de catarata]. *Arquivos Brasileiros de Oftalmologia* 2002;**65**(5):525-8. [0004-2749; lil-322156]

References to studies excluded from this review

Berg 2001 {published data only (unpublished sought but not used)}

Berg AA, Montoya-Pelaez LF. Comparison of lignocaine 2% with adrenaline, bupivacaine 0.5% with or without hyaluronidase

and a mixture of bupivacaine, lignocaine and hyaluronidase for peribulbar block analgesia. *Acta Anaesthesiologica Scandinavica* 2001;**45**(8):961-6. [PUBMED: 11576046]

Crawford 1994 {published data only (unpublished sought but not used)}

Crawford M, Kerr WJ. The effect of hyaluronidase on peribulbar block. *Anaesthesia* 1994;**49**(10):907-8. [PUBMED: 7802194]

Guise 1999 {published and unpublished data}

Guise P, Laurent S. Sub-Tenon's block: the effect of hyaluronidase on speed of onset and block quality. *Anaesthesia and Intensive Care* 1999;**27**(2):179-81. [PUBMED: 10212716]

House 1991 {published data only (unpublished sought but not used)}

House PH, Hollands RH, Schulzer M. Choice of anesthetic agents for peribulbar anesthesia. *Journal of Cataract and Refractive Surgery* 1991;**17**(1):80-3. [PUBMED: 2005563]

Johansen 1993 {published data only (unpublished sought but not used)}

Johansen J, Kjeldgård M, Corydon L. Retrobulbar anaesthesia: a clinical evaluation of four different anaesthetic mixtures. *Acta Ophthalmologica* 1993;**71**(6):787-90. [PUBMED: 8154254]

Lange 1989 {published data only (unpublished sought but not used)}

Lange W, Von Denffer H, Honis M. Comparison of bupivacaine 0.75% with a mixture of bupivacaine 0.75% and mepivacaine 2% in retrobulbar anaesthesia: effects of hyaluronidase. *Fortschritte der Ophthalmologie* 1989;**86**(4):312-5. [PUBMED: 2676792]

Moharib 2002 {published data only (unpublished sought but not used)}

Moharib MM, Mitra S, Rizvi SG. Effect of alkalization and/or hyaluronidase adjuvancy on a local anesthetic mixture for sub-Tenon's ophthalmic block. *Acta Anaesthesiologica Scandinavica* 2002;**46**(5):599-602. [PUBMED: 12027856]

Morsman 1992 {published data only (unpublished sought but not used)}

Morsman CD, Holden R. The effects of adrenaline, hyaluronidase and age on peribulbar anaesthesia. *Eye* 1992;**6**(3):290-2. [PUBMED: 1446762]

Ramanathan 1999 {published data only (unpublished sought but not used)}

Ramanathan US, Sullivan CA, Thompson SM, McCauley D. A study of effect of hyaluronidase on subTenon anaesthesia. *Cataract Surgery and Assessment* 1999;**40**(4):S287.

Sarvela 1992 {published data only (unpublished sought but not used)}

Sarvela J, Nikki P. Hyaluronidase improves regional ophthalmic anaesthesia with etidocaine. *Canadian Journal of Anaesthesia* 1992;**39**(9):920-4. [PUBMED: 1451220]

Soares 2002 {published data only (unpublished sought but not used)}

Soares LF, Escovedo Helayel P, Conceição DB, Oliveira Filho GR. Peribulbar block with the association of 0.5% enantiomeric mixture of bupivacaine and 2% lignocaine: effects of hyaluronidase addition. *Revista Brasileira de Anestesiologia* 2002;**52**(4):420-5. [PUBMED: 19479106]

Additional references

Afshari 2017

Afshari A, Wetterslev J, Smith AF. Can systematic reviews with sparse data be trusted?. *Anaesthesia* 2017;**72**(1):12-6. [PUBMED: 27804113]

Alhassan 2015

Alhassan MB, Kyari F, Ejere HOD. Peribulbar versus retrobulbar anaesthesia for cataract surgery. *Cochrane Database of Systematic Reviews* 2015, Issue 7. [DOI: [10.1002/14651858.CD004083.pub3](https://doi.org/10.1002/14651858.CD004083.pub3)]

Atkinson 1949

Atkinson WS. Use of hyaluronidase with local anaesthesia in ophthalmology; preliminary report. *Archives of Ophthalmology* 1949;**42**(5):628-33. [PUBMED: 15393388]

Borders 1968

Borders CL, Raferty MA. Purification and partial characterization of testicular hyaluronidase. *Journal of Biological Chemistry* 1968;**243**(13):3756-62. [PUBMED: 5658550]

Brown 1999

Brown S, Brooks S, Mazow M, Avilla C, Braverman D, Greenhaw S, et al. Cluster of diplopia cases after periocular anesthesia without hyaluronidase. *Journal of Cataract and Refractive Surgery* 1999;**25**(9):1245-9. [PUBMED: 10476509]

Brown 2001

Brown S, Coats D, Collins M, Underdahl J. Second cluster of strabismus cases after periocular anesthesia without hyaluronidase. *Journal of Cataract and Refractive Surgery* 2001;**27**(11):1872-5. [PUBMED: 11709263]

Buhren 2016

Buhren BA, Schrupf H, Hoff NP, Bölke N, Hilton S, Gerber PA. Hyaluronidase: from clinical applications to molecular and cellular mechanisms. *European Journal of Medical Research* 2016;**21**(5). [PUBMED: 26873038]

Dempsey 1997

Dempsey GA, Barrett PJ, Kirby IJ. Hyaluronidase and peribulbar block. *British Journal of Anaesthesia* 1997;**78**(6):671-4. [PUBMED: 9215017]

GRADEpro GDT [Computer program]

McMaster University (developed by Evidence Prime). GRADEpro GDT. Version accessed August 2017. Hamilton (ON): McMaster University (developed by Evidence Prime), 2015.

Guay 2015

Guay J, Sales K. Sub-Tenon's anaesthesia versus topical anaesthesia for cataract surgery. *Cochrane Database of Systematic Reviews* 2015, Issue 8. [DOI: [10.1002/14651858.CD006291.pub3](https://doi.org/10.1002/14651858.CD006291.pub3)]

Guyatt 2011

Guyatt G, Oxman AD, Akl EA, Kunz R, Vist G, Brozek J, et al. GRADE guidelines: 1. Introduction - GRADE evidence profiles and summary of findings tables. *Journal of Clinical Epidemiology* 2011;**64**(4):383-94. [PUBMED: 21195583]

Higgins 2011

Higgins JPT, Green S, editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Hylenex® Prescribing Information 2016

Halozyme Therapeutics, Inc. Hylenex® recombinant prescribing information, 2016. hylanex.com/downloads/approved-uspi-bl301feb2016.pdf (accessed 27 January 2018).

Kempeneers 1992

Kempeneers A, Dralands L, Ceuppens J. Hyaluronidase induced orbital pseudotumour as complication of retrobulbar anesthesia. *Bulletin de la Societe Belge d'Ophthalmologie* 1992;**243**:159-66. [PUBMED: 1302146]

Koornneef 1988

Koornneef L. Eyelid and orbital fascial attachments and their clinical significance. *Eye* 1988;**2**(2):130-4. [PUBMED: 3197870]

Meyer 1934

Meyer K, Palmer FW. The polysaccharide of the vitreous humour. *Journal of Biochemistry* 1934;**107**:629-34.

RevMan 2014 [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Roberts 1993

Roberts J, MacLeod B, Hollands RH. Improved peribulbar anaesthesia with alkalization and hyaluronidase. *Canadian Journal of Anaesthesia* 1993;**40**(9):835-8. [PUBMED: 8403178]

Schein 2000

Schein OD, Friedman DS, Fleisher LA, Lubomski LH, Magaziner J, Sprintz M, et al. Anaesthesia Management During Cataract Surgery. Evidence Report/Technology Assessment Number 16 AHRQ Publication No. 00-E015 (Archived). Vol. **1**, Rockville (MD): Agency for Healthcare Research and Quality, 2001.

WHO 2010

World Health Organization. WHO guidelines on tissue infectivity distribution in transmissible spongiform encephalopathies, 2010. www.who.int/bloodproducts/tse/WHO%20TSE%20Guidelines%20FINAL-22%20JuneupdatedNL.pdf (accessed 27 January 2018).

References to other published versions of this review

Database of Systematic Reviews 2013, Issue 2. [DOI:
[10.1002/14651858.CD010368](https://doi.org/10.1002/14651858.CD010368)]

Rüsch H 2013

Rüsch H, Adams L, Bunce C. Use of hyaluronidase as an adjunct to local anaesthetic eye blocks. *Cochrane*

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Bowman 1997

Methods	Parallel group, prospective, masked randomized controlled single centre study in the UK. Study dates not stated.
Participants	92 adults (extracapsular cataract extraction phacoemulsification and trabeculectomy) received peribulbar block. Number of participants: hyaluronidase group; 44 control group ; 48. Mean age : hyaluronidase group: 72 years; control group : 75 years.
Interventions	Hyaluronidase group ; lignocaine 2% with adrenaline 1:200,000, + bupivacaine 0.5% + hyaluronidase 150 IU/mL. Control group: lignocaine 2% with adrenaline 1:200,000 + bupivacaine 0.5%. 10 mL peribulbar injection using a standardized technique.
Outcomes	Akinesia, objective analgesia assessed by surgeon, subjective analgesia assessed by participant after surgery. VAS 0 to 10 for subjective and objective pain scores were stratified into a dichotomous pain/no pain.
Notes	Conflict of interest and funding sources not documented.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No details of randomization described.
Allocation concealment (selection bias)	Unclear risk	Masked allocation but no details described
Incomplete outcome data (attrition bias) All outcomes	Low risk	No withdrawals reported.
Selective reporting (reporting bias)	Low risk	All groups were reported on.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Masking of participants described.

Bowman 1997 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Double masking described.
---	----------	---------------------------

Brydon 1995

Methods	Parallel group, randomized double blind design in the UK. Study dates were not stated.
Participants	60 consecutive adults for elective intra-ocular surgery. Number of participants: 20 per group. Results for low dose and higher dose were combined for analysis. Mean age: hyaluronidase (low dose) group: 74 years; hyaluronidase (higher dose) group: 73 years; control group: 72 years.
Interventions	Hyaluronidase (low dose) group: peribulbar block with equal mixture of lignocaine 2% and bupivacaine 0.75% + hyaluronidase 50 IU/mL. Hyaluronidase (higher dose) group: peribulbar block with equal mixture of lignocaine 2% and bupivacaine 0.75% + hyaluronidase 150 IU/mL. Control group: peribulbar block with equal mixture of lignocaine 2% and bupivacaine 0.75%. No sedation and premedication given.
Outcomes	Speed of onset, akinesia, analgesia, top-up frequency, incidence of harm. Analgesia measured by assessing participant's reaction to insertion of superior rectus suture and by direct questioning during procedure. 3 point scoring system used. Akinesia was the primary outcome measure.
Notes	Conflict of interest and funding sources not documented.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Random assignment but no details described.
Allocation concealment (selection bias)	Unclear risk	Masked allocation but no details described.
Incomplete outcome data (attrition bias) All outcomes	Low risk	No withdrawals.
Selective reporting (reporting bias)	Low risk	All prespecified outcomes reported on.
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	"Composition of the local anaesthetic solution was not known to the anaesthetist", but no further details of masking described.

Brydon 1995 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Assessor (surgeon) masked.
---	----------	----------------------------

Khandwala 2008

Methods	Parallel group, prospective randomized controlled double masked, single centre trial in Leeds, the UK. Study dates; not stated.
Participants	20 adults undergoing routine cataract surgery. Data for 1 participant in hyaluronidase group were incomplete and excluded from analysis. Participants were ASA 1-3. Number of participants in analysis: hyaluronidase group: 9; control group: 10. Mean age: hyaluronidase group: (73.8 years; control group: 74 years). Exclusion criteria; refusal, language problems, history of allergy to amide local anaesthetics or hyaluronidase or pre-existing extra ocular muscle palsy.
Interventions	Hyaluronidase group: lignocaine 2% + hyaluronidase 15 IU/mL. Control group: lignocaine 2%. Sub-Tenon's block. Total volume of local anaesthesia 5 mL with no premedication sedation.
Outcomes	Akinesia, depth of anaesthetic fluid spread on ultrasound, surgical conditions, pain during operation measured by visual analogue scale (VAS). SD pain scores unavailable. Attempts to contact authors for more clarification unsuccessful.
Notes	Authors declared no conflict of interest and received no funding from private or public bodies.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer generated randomization.
Allocation concealment (selection bias)	Low risk	Coded syringes.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	1 participant in treatment group excluded after randomization, due to incomplete data.
Selective reporting (reporting bias)	Low risk	All outcomes reported on.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Participants and personnel were masked.

Khandwala 2008 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Assessor masked.
---	----------	------------------

Remy 2008

Methods	Parallel group, prospective randomized double masked placebo controlled trial with a multicentre design in Germany. Study dates: 29 July 2003 to 2 November 2004.
Participants	80 adults undergoing elective cataract surgery with retrobulbar block. No participant dropped out. Number of participants: hyaluronidase group: 40; control group: 40. Mean (SD) age: hyaluronidase group: 76 ± 11 years; control group: 74±10 years. Inclusion criteria; adults aged >18 years, elective surgery, no active ocular disease and informed consent obtained in written form. Exclusion criteria; known intolerance to hyaluronidase, pregnancy, lack of co-operation, history of alcohol or drug abuse, or local anaesthetic complications.
Interventions	Hyaluronidase group: 5 mL 1% mepivacaine + 75 IU/mL hyaluronidase. Control group: 5 mL 1% mepivacaine + placebo (special batch of Hyalase without active ingredient).
Outcomes	Primary end point; complete akinesia after 5 minutes. Secondary end points; akinesia at other times, top-up injections, ptosis, time to anaesthesia, pain (VAS) immediately after surgery and 3 hours postsurgery and efficacy and tolerability for participant and surgeon. Adverse events recorded. First pain assessment point, planned before surgery then reported for immediately after surgery.
Notes	No conflict or financial interest reported by author.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Randomization not described in detail.
Allocation concealment (selection bias)	Low risk	Double masked as per German legal framework.
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participant dropped out.
Selective reporting (reporting bias)	Low risk	All outcome measures were reported on.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Placebo control, double masked.

Remy 2008 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Masked according to federal law.
---	----------	----------------------------------

Rowley 2000

Methods	Parallel group, randomized double masked controlled trial in 1 centre in UK. Study dates: not stated.
Participants	150 adults for elective cataract surgery with sub-Tenon's block. Number of participants: hyaluronidase group: 76; control group: 74. Mean age; in hyaluronidase group; 77.14 years; control group; 76.51 years. Groups similar in terms of age, sex and proportion of blocks administered by each investigator. Exclusion criteria: people with learning difficulties, dementia, profound deafness and known adverse reaction to lignocaine or hyaluronidase.
Interventions	Hyaluronidase group: 3 mL 2% lignocaine/adrenaline + hyaluronidase 30 IU/mL. Control group: 3 mL 2% lignocaine/adrenaline.
Outcomes	Akinesia, post-injection and immediate postoperative pain. Pain measured using VAS (0-10 cm). SD pain scores not available with attempts to obtain more clarification from authors unsuccessful.
Notes	Declaration of funding sources and conflict of interest not reported.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number tables.
Allocation concealment (selection bias)	Low risk	Masked syringes.
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts.
Selective reporting (reporting bias)	Low risk	All outcomes reported accordingly.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Masking of participants not described. Masking of personnel (operative surgeon, independent assistant and nursing staff) was described.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Double masked design.

Sedghipour 2012

Methods	Parallel group, randomized double masked trial in 1 centre in Iran. Study dates: February 2011 to July 2011.
Participants	44 adults initially recruited from a referral eye centre (Nikookari Eye Hospital) to undergo elective cataract surgery (phacoemulsification) under sub-Tenon block. 2 participants did not meet the criteria and were excluded. Number of participants: hyaluronidase group: 21; control group: 21. Mean (SD) ages: hyaluronidase group: 65.62 ± 3.01 years; control group; 67 ± 4.4 years. Groups comparable for gender and age. Exclusion criteria: people with deafness and allergy to lidocaine or hyaluronidase.
Interventions	Hyaluronidase group: 2 mL 2% lidocaine + hyaluronidase 150 IU/mL Control group: 2 mL 2% lidocaine. Ampoules were identical in appearance with a printed code (A or B). Codes disclosed for statistical analysis only at end of study.
Outcomes	Akinesia, participant and surgical satisfaction with yes/no questions, postoperative pain via VAS scoring using a standard VAS chart. Participants were given appropriate explanation on usage of chart.
Notes	No declaration of funding sources or conflict of interest reported.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Consecutive numbers assigned to participants on admission by a staff member not involved in study.
Allocation concealment (selection bias)	Low risk	Coded syringes.
Incomplete outcome data (attrition bias) All outcomes	Low risk	No exclusions after randomization reported.
Selective reporting (reporting bias)	Low risk	All outcomes reported.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Masking of participants, personnel by coded syringes.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Assessor masked by coded syringes.

Shiroma 2002

Methods	Randomized double masked study conducted at State University of Campinas - Unicamp, Brazil. Study dates: not stated.
Participants	57 adults undergoing elective extracapsular cataract extraction on an outpatient basis. Participant's physical statuses described as ASA 1- 3. Number of participants: hyaluronidase group: 29; control group: 28. Sex: 31 men (54.4%); 26 women (45.6%). Age range (overall): 45- 89 years; mean (SD) overall: 67.73 ($\pm$ 10.65). Groups homogeneous in relation to sex, age and physical condition. Peribulbar injection given as a block by double needle injection with 25x7 mm needle, administered 4 mL at lower temporal with super-medial inclination of about 15° and 3 mL (nasal-superior). Anaesthetic solution prepared without knowledge of ophthalmologist who performed the block.
Interventions	Hyaluronidase group: ropivacaine 1% + hyaluronidase 100 IU/mL. Control group: ropivacaine 1%.
Outcomes	Onset time to akinesia, need for supplementary injections and pain assessed by VAS (0-10).
Notes	No declaration of funding sources or conflict of interest reported.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method of randomization not specified.
Allocation concealment (selection bias)	Unclear risk	Allocation concealment not described.
Incomplete outcome data (attrition bias) All outcomes	Low risk	No withdrawals documented, all participants included initially were assessed and reported.
Selective reporting (reporting bias)	Low risk	Primary outcomes reported.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Masking described in participants and personnel.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Assessment masked.

ASA: American Society of Anesthesiology Classification; SD: standard deviation; VAS: visual analogue scale.

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Berg 2001	Pain not assessed formally using a rating scale. Instead, augmentation eye drops given if participant complained of pain during procedure. Therefore, pain not measured but reported.
Crawford 1994	2 participants in the hyaluronidase group received local anaesthetic drops for pain intraoperatively. However, pain not formally measured, therefore, inclusion criteria not met.
Guisse 1999	Pain not formally measured using a rating scale.
House 1991	Pain not measured as per rating scale and top-ups primarily given to achieve akinesia initially.
Johansen 1993	Surgeon assessed quality of analgesia and pain, not measured by asking participant. No formal method of assessing pain described.
Lange 1989	On further inspection, not a randomized trial.
Moharib 2002	Adequate pain relief defined as lack of complaint or response from participant. Did not constitute pain measurement and no formal assessment of pain described in the study.
Morsman 1992	Presence or absence of hyaluronidase not masked.
Ramanathan 1999	Poster presentation, Further details could not be obtained about randomization, masking and results.
Sarvela 1992	Part one of third study was not randomized. Part two of this study compared two different concentrations of hyaluronidase only.
Soares 2002	Intraoperative pain not measured (e.g. by asking participant). It was assumed that the participants would spontaneously voice their pain during operation. Therefore, study did not measure any of our stipulated outcomes.

DATA AND ANALYSES

Comparison 1. Hyaluronidase versus control

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Intraoperative pain (reported continuous)	3		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2 Intraoperative pain (reported dichotomous)	4	289	Risk Ratio (M-H, Random, 95% CI)	0.83 [0.48, 1.42]

Analysis 1.1. Comparison 1 Hyaluronidase versus control, Outcome 1 Intraoperative pain (reported continuous).

Study or subgroup	Hyaluronidase		Control		Std. Mean Difference Random, 95% CI	Weight	Std. Mean Difference Random, 95% CI
	N	Mean(SD)	N	Mean(SD)			
Khandwala 2008	9	1.4 (1.5)	10	0.7 (1.6)		0%	0.44[-0.47,1.36]
Rowley 2000	76	2.3 (1.5)	74	2 (1.6)		0%	0.21[-0.12,0.53]
Sedghipour 2012	21	1.9 (1.5)	21	3 (1.6)		0%	-0.72[-1.35,-0.09]

Favours hyaluronidase -100 -50 0 50 100 Favours control

Analysis 1.2. Comparison 1 Hyaluronidase versus control, Outcome 2 Intraoperative pain (reported dichotomous).

Study or subgroup	Hyaluronidase		Control		Risk Ratio M-H, Random, 95% CI	Weight	Risk Ratio M-H, Random, 95% CI
	n/N	n/N	n/N	n/N			
Remy 2008	9/40	19/40				32.63%	0.47[0.24,0.92]
Bowman 1997	9/44	12/48				28.18%	0.82[0.38,1.75]
Brydon 1995	4/40	2/20				9.63%	1[0.2,5]
Shiroma 2002	12/29	8/28				29.56%	1.45[0.7,3]
Total (95% CI)	153	136				100%	0.83[0.48,1.42]

Total events: 34 (Hyaluronidase), 41 (Control)
Heterogeneity: Tau²=0.12; Chi²=5.04, df=3(P=0.17); I²=40.53%
Test for overall effect: Z=0.69(P=0.49)

Favours hyaluronidase 0.01 0.1 1 10 100 Favours control

APPENDICES

Appendix 1. The Cochrane Central Register of Controlled Trials (CENTRAL, 2017, Issue 6)

Search lines	Search terms	Search results
1	eye* or Ophthalm* or ocular	39,485
	cornea* or retin* or scler* or vitre* or iris or pupil or orbit* or chorod*	33,449
3	intraocular or intra-ocular or extraocular or extra-ocular or monocular or ocu- lo* or oculi or optic*	16,808
4	visual* or vision or sight or see* or view* or blind*	380,089
5	glaucoma or conjuncti* or uveitis or macula* or oedema or edema or strabis- mus or squint or astigmati* or myopi* or hypermetropia or trachoma	33,656
6	#1 or #2 or #3 or #4 or #5	413,088
7	surg* or operat*	203,926
8	transplant* or graft* or extract* or cataract or refractive or oculoplast* or oph- thalmosurg*	92,037

(Continued)

9	#7 or #8	258,017
10	#6 and #9	112,631
11	retrobulbar or peribulbar or sub-tenon's block or sub-tenon's block	806
12	"Nerve Block" or Lidocaine or Mepivacaine or Bupivacaine	19,237
13	(local anaesthe*) or (local anesthe*)	15,573
14	#11 or #12 or #13	25,341
15	#10 and #14	11,101
16	hyaluronidase or hyaluronoglucosaminidase or hyadase or vitrase or Amphadase or wydase or hyalase or hylenex	533
17	#10 and #14 and #16	230
18	Limit to TRIALS database	207

Appendix 2. MEDLINE (Ovid) from 1946 to 30 June 2017

Search lines	Search terms	Search results
1	eye* or Ophthalm* or ocular	578,410
2	cornea* or retina* or scler* or vitre* or iris or pupil or orbit* or chorod*	738,578
3	intraocular or intra-ocular or extraocular or extra-ocular or monocular or ocu- lo* or oculi or optic*	478,576
4	visual* or vision or sight or see* or view* or blind*	2,691,438
5	glaucoma or conjuncti* or uveitis or macula* or oedema or edema or strabis- mus or squint or astigmati* or myopi* or hypermetropia or trachoma	436,501
6	1 or 2 or 3 or 4 or 5	3,928,131
7	surg* or operat*	2,516,135
8	transplant* or graft* or extract* or cataract or refractive or oculoplast* or oph- thalmosurg*	1,578,347
9	7 or 8	3,810,842
10	6 and 9	724,695
11	exp Ophthalmologic Surgical Procedures/ or exp Cataract Extraction/	99,847
12	exp Refractive Surgical Procedures/	54,938
13	exp Corneal Transplantation/	14,186

(Continued)

14	exp Phacoemulsification/ or exp Lens Implantation, Intraocular/	13,574
15	11 or 12 or 13 or 14	99,847
16	10 or 15	753,408
17	retrobulbar or peribulbar or sub-tenon's block or subtenon's block	4698
18	local anaesthe* or local aneshe*	38,258
19	17 or 18	42,504
20	exp Anesthesia, Local/	16,213
21	exp Anesthetics, Local/	99,177
22	exp Lidocaine/ or exp Nerve Block/	30,602
23	exp Mepivacaine/ or exp Bupivacaine/	12,786
24	20 or 21 or 22 or 23	120,407
25	19 or 24	140,352
26	16 and 25	13,782
27	hyaluronidase or hyaluronoglucosaminidase or hyadase or vitrase or Am-phadase or wydase or hyalase or hyleneX	10,943
28	exp Hyaluronoglucosaminidase	8216
29	27 or 28	10,943
30	16 and 25 and 29	319
31	limit 30 to controlled clinical trial or randomised controlled trial	139
32	((single adj (blind* or masked)) or double) adj (blind* or masked)).ab. or (((single adj (blind* or masked)) or double) adj (blind* or masked)).ti.	142,612
33	(randomized or randomly or placebo or trial or (controlled adj study)).ab. or (randomized or randomly or placebo or trial or (controlled adj study)).ti.	1,054,662
34	32 or 33	1,066,944
35	30 and 34	135
36	31 or 35	155

Appendix 3. Embase (Ovid) 1947 to 30 June 2017

Search lines	Search terms	Search results
--------------	--------------	----------------

(Continued)

1	eye* or Ophthalm* or ocular	714,955
2	cornea* or retin* or scler* or vitre* or iris or pupil or orbit* or chorod*	1,053,140
3	intraocular or intra-ocular or extraocular or extra-ocular or monocular or ocu- lo* or oculi or optic*	503,874
4	visual* or vision or sight or see* or view* or blind*	3,707,107
5	glaucoma or conjuncti* or uveitis or macula* or oedema or edema or strabis- mus or squint or astigmati* or myopi* or hypermetropia or trachoma	688,043
6	1 or 2 or 3 or 4 or 5	5,252,878
7	surg* or operat*	4,385,968
8	transplant* or graft* or extract* or cataract or refractive or oculoplast* or oph- thalmosurg*	2,208,147
9	7 or 8	6,014,664
10	6 and 9	1,223,645
11	exp eye surgery/	133,842
12	exp cornea transplantation/	11,680
13	exp cataract extraction/ or extracapsular cataract extraction/	43,300
14	exp refractive surgery/	8947
15	exp lens implant/	20,733
16	exp eyelid reconstruction/	5242
17	11 or 12 or 13 or 14 or 15 or 16	138,278
18	10 or 17	1,241,841
19	retrobulbar or peribulbar or sub-tenon's block or subtenon's block	7897
20	local anaesthe* or local anesthe*	80,577
21	19 or 20	87,361
22	exp retrobulbar drug administration/	1081
23	exp lidocaine/ or exp peribulbar anesthesia/ or exp local anesthetic agent/ or exp bupivacaine/	228,041
24	exp local anesthesia/ or exp nerve block/	68,789
25	exp levobupivacaine/ or exp procaine/	22,051
26	ropivacaine/ or midazolam/ or propofol/	80,182

(Continued)

27	22 or 23 or 24 or 25 or 26	326,465
28	21 or 27	343,082
29	18 and 28	40,436
30	hyaluronidase or hyaluronoglucosaminidase or hyadase or vitrase or Amphadase or wydase or hyalase or hylenex	16,613
31	exp recombinant hyaluronidase/ or exp hyaluronidase/ or exp hyaluronoglucosaminidase/	14,509
32	30 or 31	16,613
33	18 and 28 and 32	709
34	limit 33 to (clinical trial or randomized controlled trial or controlled clinical trial or multicenter study or phase 1 clinical trial or phase 2 clinical trial or phase 3 clinical trial or phase 4 clinical trial)	199
35	(randomi?ed or randomly or placebo or trial or (controlled adj study)).ab. or (randomi?ed or randomly or placebo or trial or (controlled adj study)).ti.	1,391,344
36	((single adj (blind* or masked)) or double) adj (blind* or masked)).ab. or (((single adj (blind* or masked)) or double) adj (blind* or masked)).ti.	185,195
37	35 or 36	1,408,147
38	33 and 37	199
39	34 or 38	251

Appendix 4. Web of Science from 1900 to 30 June 2017

Search lines	Search terms	Search results
1	eye* or Ophthalm* or ocular	476,661
2	cornea* or retin* or scler* or vitre* or iris or pupil or orbit* or chorod*	1,045,927
3	intraocular or intra-ocular or extraocular or extra-ocular or monocular or ocu-lo* or oculi or optic*	1,576,811
4	visual* or vision or sight or see* or view* or blind*	4,121,849
5	glaucoma or conjuncti* or uveitis or macula* or oedema or edema or strabismus or squint or astigmati* or myopi* or hypermetropia or trachoma	425,695
6	#5 OR #4 OR #3 OR #2 OR #1	6,754,964
7	surg* or operat*	3,423,025

(Continued)

8	transplant* or graft* or extract* or cataract or refractive or oculoplast* or ophthalmosurg*	2,193,730
9	#8 OR #7	5,349,001
10	#9 AND #6	1,005,525
11	(Lens Implantat*) OR (cornea transplantat*) OR (cataract extract*) OR (refractive surg*) OR (exp eyelid reconstruct*)	26,003
12	#11 OR #10	1,008,989
13	(retrobulbar or peribulbar or sub-tenon's block or subtenon's block)	4057
14	(retrobulbar or peribulbar or sub-tenon's block or sub-tenon's block)	4056
15	(local anaesthe*) OR (local anesthe*)	41,088
16	(lidocaine OR levobupivacaine OR procaine OR ropivacaine OR midazolam OR propofol) OR TOPIC: (nerve block)	85,921
17	#16 OR #15 OR #14 OR #13	117,842
18	(hyaluronidase or hyaluronoglucosaminidase or hyalase or vitrase or Amphadase or wydase or hyalase or hylenex) OR TOPIC: (recombinant hyaluronidase)	6294
19	#18 AND #17 AND #12	214

Appendix 5. Scopus from 1823 to 30 June 2017

Search lines	Search terms	Search results
1	eye* or Ophthalm* or ocular	1,943,281
2	cornea* or retin* or scler* or vitre* or iris or pupil or orbit* or chorod*	3,136,708
3	intraocular or intra-ocular or extraocular or extra-ocular or monocular or ocu- lo* or oculi or optic	872,852
4	visual* or vision or sight or see* or view* or blind*	13,637,048
5	glaucoma or conjuncti* or uveitis or macula* or oedema or edema or strabis- mus or squint or astigmati* or myopi* or hypermetropia or trachoma	1,292,318
6	surg* or operat*	11,977,249
7	transplant* or graft* or extract* or cataract or refractive or oculoplast* or oph- thalmosurg*	6,168,543
8	retrobulbar or peribulbar or (sub-tenon's block) or (subtenon's block)	15,268
9	(local anaesthe*) or (local anesthe*)	241,430

(Continued)

10	Lidocaine or (Nerve Block) or Mepivacaine or Bupivacaine	334,730
11	hyaluronidase OR hyaluronoglucosaminidase OR hyalase OR vitrase OR am-phadase OR wydase OR hyalase OR hylenex	28,617
12	#1 or #2 or #3 or #4 or #5	16,812,220
13	#6 or #7	16,339,616
14	#8 or #9 or #10	509,491
15	#11 and #12 and #13 and #14	1,857

Appendix 6. CINAHL Plus (EBSCOhost) from 1937 to 30 June 2017

Search lines	Search terms	Search results
1	eye* or Ophthalm* or ocular	62,235
2	cornea* or retina* or scler* or vitre* or iris or pupil or orbit* or chorod*	68,825
3	intraocular or intra-ocular or extraocular or extra-ocular or monocular or ocu-lo* or oculi or optic*	24,333
4	visual* or vision or sight or see* or view* or blind*	426,645
5	glaucoma or conjuncti* or uveitis or macula* or oedema or edema or strabis-mus or squint or astigmati* or myopi* or hypermetropia or trachoma	48,062
6	S1 OR S2 OR S3 OR S4 OR S5	547,031
7	surg* or operat*	513,190
8	transplant* or graft* or extract* or cataract or refractive or oculoplast* or oph-thalmosurg*	162,110
9	S7 OR S8	625,438
10	S6 AND S9	101,771
11	(MH "Surgery, Eye+") OR (MM "Keratomileusis, Laser in Situ") OR (MM "Eye Enucleation")	13,308
12	(MM "Corneal Transplantation")	1060
13	(MH "Cataract Extraction+")	5140
14	(MM "Phacoemulsification")	848
15	S11 OR S12 OR S13 OR S14	13,308
16	S10 OR S15	104,161

(Continued)

17	retrobulbar or peribulbar or sub-tenon's block or subtenon's block	352
18	local anaesthe* or local anesthe*	10,097
19	S17 OR S18	11,261
20	(MM "Nerve Block")	4680
21	(MM "Anesthesia, Local") OR (MH "Anesthetics, Local+")	14,442
22	S20 OR S21	17,375
23	S19 OR S22	20,462
24	S16 AND S23	2377
25	hyaluronidase or hyaluronoglucosaminidase or hyadase or vitrase or Am-phadase or wydase or hyalase or hylenex	371
26	(MM "Hyaluronidase")	129
27	S25 OR S26	371
28	S16 AND S23 AND S27	40

Appendix 7. LILACS from 1982 to 30 June 2017

Search lines	Search terms	Search results
1	eye* or Ophthalm* or ocular or cornea* or retina* or scler* or vitre* or iris or pupil or orbit* or chorod* or intraocular or intra-ocular or extraocular or extra-ocular or monocular or oculo* or oculi or optic* or visual* or vision or sight or see* or view* or blind* or glaucoma or conjuncti* or uveitis or macula* or oedema or edema or strabismus or squint or astigmati* or myopi* or hypermetropia or trachoma	5,238,714
2	surg* or operat* or transplant* or graft* or extract* or cataract or refractive or oculoplast* or ophthalmosurg*	4,831,015
3	retrobulbar or peribulbar or sub-tenon's block or sub-tenon's block or (local anaesthe*) or (local anesthe*) or Lidocaine or (Nerve Block) or Mepivacaine or Bupivacaine	320,196
4	hyaluronidase OR hyaluronoglucosaminidase OR hyadase OR vitrase OR am-phadase OR wydase OR hyalase OR hylenex	8146
5	1 and 2 and 3 and 4 and 5	336

Appendix 8. Data collection form

Cochrane Anaesthesia Review Group

Study Selection, Quality Assessment & Data Extraction Form

Use of hyaluronidase as an adjunct to local anaesthetic eye blocks to reduce intraoperative pain in adults (Review)

Copyright © 2018 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

First author	Journal/Conference Proceedings etc	Year
--------------	------------------------------------	------

Study eligibility

RCT/Quasi/CCT (delete as appropriate)	Relevant participants	Relevant interventions	Relevant outcomes
Yes / No / Unclear	Yes / No / Unclear	Yes / No / Unclear	Yes / No* / Unclear

*** Issue relates to selective reporting when authors may have taken measurements for particular outcomes, but not reported these within the paper(s). Reviewers should contact trialists for information on possible non-reported outcomes & reasons for exclusion from publication. Study should be listed in 'Studies awaiting assessment' until clarified. If no clarification is received after three attempts, study should then be excluded.**

Do not proceed if any of the above answers are 'No'. If study to be included in 'Excluded studies' section of the review, record below the information to be inserted into 'Table of excluded studies'.

Freehand space for comments on study design and treatment:

References to trial

Check other references identified in searches. If there are further references to this trial link the papers now & list below. All references to a trial should be linked under one *Study ID* in RevMan.

Code each paper	Author(s)	Journal/Conference Proceedings etc	Year
A	<i>The paper listed above</i>		
B	<i>Further papers</i>		

(Continued)

Participant characteristics

Further details

Age (mean, median, range, etc)

Sex of participants (numbers / %, etc)

Dose of hyaluronidase

Trial characteristics

Methodological quality

Allocation of intervention

State here method used to generate allocation and reasons for grading

Grade (circle)

Adequate (Random)

Inadequate (e.g. alternate)

Unclear

Concealment of allocation

Process used to prevent foreknowledge of group assignment in a RCT, which should be seen as distinct from masking

State here method used to conceal allocation and reasons for grading

Grade (circle)

Adequate

Inadequate

Unclear

Masking

Person responsible for participants care	Yes / No
Participant	Yes / No
Outcome assessor	Yes / No
Other (please specify)	Yes / No

Intention-to-treat

An intention-to-treat analysis is one in which all the participants in a trial are analysed according to the intervention to which they were allocated, whether they received it or not.

All participants entering trial

15% or fewer excluded

More than 15% excluded

Not analysed as 'intention-to-treat'

Unclear

Were withdrawals described? Yes ? No ? not clear ?

Is attrition reported? Yes ? no ?

Total number randomized

Number in hyaluronidase group

Number in control group

Discuss if appropriate and note reasons for attrition

Is there a possibility of selective outcome reporting Yes? No?

Discuss if appropriate

Data extraction

Outcomes relevant to your review

	Reported in paper (circle)
Intraoperative pain	Yes / No

(Continued)

Adverse Events	Yes / No
----------------	----------

Satisfaction scoring	Yes / No
----------------------	----------

For Continuous data						
Code of paper	Unit of measurement	Intervention group		Control group		Details if outcome only described in text
		n	Mean (SD)	n	Mean (SD)	
Outcomes						
	Intraoperative VAS pain score					
	(include range)					

For Dichotomous data

Code of paper	Outcomes	Intervention group (n) n = number of participants, not number of events	Control group (n) n = number of participants, not number of events
	Intraoperative Pain (where not VAS)		

Other information which you feel is relevant to the results

Indicate if: any data were obtained from the primary author; if results were estimated from graphs etc; or calculated by you using a formula (this should be stated and the formula given). In general if results not reported in paper(s) are obtained this should be made clear here to be cited in review.

Adverse events recorded here

Freehand space for writing actions such as contact with study authors and changes

Surgical specialty: (CIRCLE)

1. Anterior segment.
2. Glaucoma.
3. Vitreoretinal surgery.

References to other trials

Did this report include any references to published reports of potentially eligible trials not already identified for this review?

First author	Journal / Conference	Year of publication
--------------	----------------------	---------------------

Did this report include any references to unpublished data from potentially eligible trials not already identified for this review? If yes, give list contact name and details

(Continued)

Trial characteristics Summary

Further details

Single centre / Multicentre

Country / Countries

How was participant eligibility defined?

How many people were randomized?

Number of participants in each intervention group

Number of participants who received intended treatment

Number of participants who were analysed

Local anaesthetic Drug dose (s) used

Dose of hyaluronidase

Median (range) length of follow-up reported in this paper (state weeks, months or years or if not stated)

Time-points when measurements were taken during the study

Time-points reported in the study

Time-points you are using in RevMan

Trial design

Other

Appendix 9. Characteristics of excluded studies

Reason for exclusion noted as narrative only:

Appendix 10. Risk of bias table

Entry

Judgement

Description

(Continued)

Low risk/high risk/un-
clear risk

1. Random sequence generation
2. Allocation concealment
3. Masking of participants and personnel
4. Masking of outcome assessment
5. Incomplete outcome data
6. Selective reporting
7. Other bias

WHAT'S NEW

Date	Event	Description
4 October 2018	Amended	Acknowledgement section amended to include Co-ordinating Editor

CONTRIBUTIONS OF AUTHORS

Conceiving the review: HR.

Co-ordinating the review: HR.

Undertaking manual searches: KA and HR.

Screening search results: KA and HR.

Organizing retrieval of papers: DB, KA and HR.

Screening retrieved papers against inclusion criteria: KA and HR.

Appraising quality of papers: KA and HR.

Abstracting data from papers: KA and HR.

Writing to authors of papers for additional information: KA and HR.

Providing additional data about papers: HR.

Obtaining and screening data on unpublished studies: HR.

Data management for the review: KA and HR.

Entering data into Review Manager (5): KA and HR.

Review Manager 5 statistical data: KA, HR and CB.

Other statistical analysis not using Review Manager 5: CB.

Double entry of data: KA and HR.

Interpretation of data: KA and HR.

Statistical inferences: CB.

Assessment of risk of bias: KA and HR.

Writing the review: KA and HR.

Securing funding for the review: HR and CB.

Guarantor for the review: HR.

Responsible for reading and checking review before submission: KA and HR.

DECLARATIONS OF INTEREST

HR: no known conflict of interest.

KA: no known conflict of interest.

CB : no known conflict of interest.

DB : no known conflict of interest.

SOURCES OF SUPPORT

Internal sources

- Moorfields Eye Hospital, UK.

Salary and facilities

External sources

- No sources of support supplied

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

We made the following changes to the protocol ([Rüschén 2013](#));

1. This review included only adults, we therefore stated "adults" in the title.
2. The phrase 'to reduce intraoperative pain' was added to the title to comply with PICO to reflect the review question.
3. Lee Adams was initially a registered author, but gave his agreement to be removed from the authors list at the protocol stage.
4. We added Kavitha Aravinth as second author.
5. We added Desta Bokre as fourth author.
6. We further stated in the primary objective that when considering studies for inclusion that we excluded studies that did not measure pain with a rating scale because spontaneous reporting of pain by the participant is not likely to produce reproducible results.
7. We stated the secondary objectives (incidence of harm, participant and surgical satisfaction and economic outcomes or cost calculations) in the [Objectives](#) section.
8. We stated in the protocol that intraoperative pain would be recorded using (VAS); however, we observed that included studies used either (VAS) or (VRS) to rate pain. As both are accepted and validated tools to measure pain, we analysed both these methods of measuring pain in the same manner.
9. We initially proposed to include only studies that described the first eye operation, but during the review process, we found no publication that described if the participants had a first or second eye operation. (See [Types of participants](#) and [Unit of analysis issues](#)).
10. We excluded adnexal surgery and any other eye surgery that was not intraocular. Anaesthetic techniques and surgical interventions for adnexal operations are inherently different from intraocular surgery and produce a very different pain profile.
11. We used GRADE-pro to analyse the quality of evidence and to produce a 'Summary of Findings table'.
12. Collation of references: we added the following to the [Selection of studies](#) section. "Where studies had multiple publications, we planned to collate the reports of the same study so that each study, rather than each report, was the unit of interest for the review and such studies had a single identifier with multiple references.

INDEX TERMS

Medical Subject Headings (MeSH)

Analgesia [*methods]; Analgesics [*administration & dosage]; Anesthesia, Local [methods]; Anesthetics, Local [*administration & dosage]; Hyaluronoglucosaminidase [*administration & dosage]; Intraoperative Complications [*prevention & control]; Pain Measurement; Pain, Procedural [*prevention & control]; Patient Satisfaction; Randomized Controlled Trials as Topic

MeSH check words

Adult; Humans