Surgical correction of coxa vara: Evaluation of neck shaft angle, Hilgenreiner-epiphyseal angle for indication of recurrence

Thammanoon Srisaarn^a, Krits Salang^b, Benjamin Klawson^c, Kitiwan Vipulakorn^b, Ornusa Chalayon^d, Perajit Eamsobhana^{d,*}

^a Department of Orthopaedic Surgery, Faculty of Medicine Phramongkutklao Hospital, Bangkok, Thailand

^b Department of Orthopaedic Surgery, Faculty of Medicine, Khon Kean Hospital, Khon Kean, Thailand

^c Department of Orthopaedic Surgery, Washington University School of Medicine in St. Louis, St. Louis, MO, USA

^d Department of Orthopaedic Surgery, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok, Thailand

ARTICLE INFO

Article history:

Received 12 December 2017

Received in revised form 30 May 2018

Accepted 26 June 2018

Available online 28 June 2018

Keywords:

Coxa vara

Osteotomy

Intramedullary nail

Osteogenesis imperfecta

Fibrous dysplasia

Skeletal dysplasia

ABSTRACT

Background: Coxa vara is a hip deformity in which the femoral neck-shaft angle decreases below its normal value. Standard surgical treatment for this condition is corrective valgus osteotomy. Appropriate correction of the Hilgenreiner-epiphyseal angle is important to prevent recurrence. The purpose of this study is to: 1) evaluate the recurrence of the deformity at the latest follow up; and 2) find the appropriate angle of correction associated with the lowest recurrence.

Methods: 34 hips in 31 patients who underwent surgery for treatment of coxa vara from 2005 to 2014 were included. Patient-reported outcomes, Hilgenreiner-epiphyseal angle, and neck-shaft angle were assessed preoperatively, postoperatively, and at latest follow-up.

Results: The mean age at surgery was 10.99, with a range of 5–30, years. Preoperative neck-shaft angle ranged from 60 to 100 degrees, and Hilgenreiner-epiphyseal angle ranged from 60 to 90 degrees. At the latest follow up, the neck-shaft angle ranged from 120 to 135 degrees and the Hilgenreiner-epiphyseal angle ranged from 22 to 35 degrees ($p < 0.001$). The Harris hip score improved from 47.20 (34–66) to 79.68 (60–100) ($p < 0.001$). There was no recurrence of deformities at the mean follow up of 37.87 months.

Conclusion: Surgical correction of coxa vara in various pathologies can be done successfully with the Hilgenreiner-epiphyseal angle corrected to ≤ 35 degrees or the neck shaft angle corrected to ≥ 120 degrees in order to prevent recurrence of the deformity. Majority of the patients were reported improvement of hip function. However, a longer-term follow up is required to determine further outcomes regarding to recurrence of the deformity.

© 2018

1. Background

Coxa vara is a condition in which the femoral neck-shaft angle decreases below its normal value based on age.^{1–3} The deformity can be primarily caused by a congenital defect of the femoral neck cartilage, or from secondary causes related to trauma, infection, skeletal dysplasia, and pathologic bone disorders.^{1,4}

Common presentations in patients with coxa vara include Trendelenburg gait and, in unilateral cases, leg-length discrepancy.

When left untreated, patients can develop morbidities, such as stress fractures and hip joint degeneration.¹ Because coxa vara occurs infrequently, there have been only a few reports describing its surgical correction. The indications for treatment of coxa vara are: 1) Hilgenreiner-epiphyseal angle greater than 60 degrees; 2) progression of the deformity; and 3) Trendelenburg movement.^{5–10} Goals of treatment are: 1) restoration of the hip mechanical axis; 2) restoration of Hilgenreiner-epiphyseal angle and the neck-shaft angle to prevent the recurrence of deformities; and 3) restored function of the hip abductor.¹¹

The purpose of this study is to: 1) report the outcomes of surgical correction of coxa vara with various pathologies using radiographic measurements and outcome scores; 2) evaluate the recurrence of the deformity at the latest follow up; and 3) find the appropriate angle of correction that may help prevent recurrence.

* Corresponding author at: Department of Orthopaedic Surgery, Faculty of Medicine Siriraj Hospital, Mahidol University, 2 Wanglang Road, Bangkoknoi, Bangkok 10700, Thailand.

E-mail addresses: srisaarn@pmk.ac.th (T. Srisaarn), kwanchanon@gmail.com (K. Salang), btklawson@wustl.edu (B. Klawson), KITVIP@kku.ac.th (K. Vipulakorn), ornusa9@gmail.com (O. Chalayon), peerajite@gmail.com (P. Eamsobhana).

2. Methods

An IRB-approval was obtained prior to data collection for this study. During the 2005–2014 study periods, there were 31 patients (34 hips) with coxa vara, from various etiologies, who underwent surgical correction. Clinic notes and radiographs of these patients were reviewed. The criteria for exclusion were incomplete patient information, incomplete radiographic data, presenting of osteoarthritis, and a follow up of less than 2 years. There were 18 males (19 hips) and 13 females (15 hips) included in this study. The left side was affected in 13 hips, and the right side in 21 hips. Three patients had bilateral correction. A valgus proximal femoral osteotomy was performed in all patients^{12,13} (Fig. 3). In addition, some patients underwent a femoral osteotomy due to femoral shaft deformity¹⁴ (Figs. 1 and 2). The radiographic parameters recorded were Hilgenreiner-epiphyseal angle and the neck-shaft angle, measured on standard AP hip radiographs. All of these parameters were reviewed and evaluated preoperatively, postoperatively, and at the latest follow up. The Harris hip scores were reviewed preoperatively, and at the latest follow up. More than 20% progression of the Hilgenreiner-epiphyseal angle and the neck shaft angle were considered a recurrence of deformity.

3. Results

The mean age at surgery was 10.99 (ranging from 5 to 30) years. The average follow-up period was 37.87 (ranging from 24 to 84) months. The etiologies of the deformity were classified as Osteogenesis Imperfecta (OI) in 9 patients, spondyloepiphyseal dysplasia or multiple epiphyseal dysplasia in 3 patients, malunion femoral neck fracture in 4 patients, congenital coxa vara in 4 patients, polyostotic fibrous dysplasia, fibrous dysplasia /monostotic fibrous dysplasia in 9 patients, and Histiocytosis X in 2 patients (Table 1). Radiographic analysis of Hilgenreiner-epiphyseal angle and the neck-shaft angle showed significant improvement at the final follow up compared with the preoperative baseline for all patients (Table 2). The mean neck-shaft angle improved from 86.09 ± 11.31 degrees preoperatively to

129.64 ± 3.95 degrees at the latest follow-up ($p < 0.001$). The mean Hilgenreiner-epiphyseal angle improved from 66.76 ± 7.99 degrees preoperatively to 29.26 ± 4.37 degrees at the final follow-up ($p < 0.001$). By the last follow-up visit, all bone fragments showed signs of radiographic union. In all osteotomies, surgical correction was maintained at the final follow up with a neck-shaft angle of at least 122 degrees in each case. At the final follow-up visit for each patient, no recurrence of coxa vara was reported. The Harris hip score improved from 47.23 ± 7.46 to 79.68 ± 8.22 ($p < 0.001$). One patient had a delayed union that was treated with bone grafting at 4 months after the first surgery. The osteotomy healed successfully without further incident after the bone grafting. Within the study period, there were no incidences of implant-related complications or infection. Significant improvement in the quality of life following the surgery was reported. Those patients who used wheel chair or axillary crutches before surgery were able to walk independently without a gait aid afterward.

4. Discussion

Patients with coxa vara are generally presented with difficulties in ambulating, specifically leg-length discrepancy and Trendelenburg gait. Abnormal force transferred through the femoral neck can cause stress fractures and osteoarthritis of the hip.^{1,4} Surgical correction plays a vital role to prevent the progression of these sequelae in patients with coxa vara. The surgical indications include Trendelenburg gait, progressively decreasing neck-shaft angle, and a Hilgenreiner-epiphyseal angle of greater than 60 degrees. Pauwels¹⁵ described the concept of osteotomy aiming to convert coxa vara to the normal axis. The treatment goal for coxa vara is to restore a normal biomechanical function of the hip joint by correcting Hilgenreiner-epiphyseal angle and the neck-shaft angle. In pediatric patients, this will re-orientate the physis to a more regular horizontal position, which is less susceptible to shear stress. Several methods have been used for fixation of the femoral osteotomy, including tension band wiring, angle blade plates, dynamic sliding hip screws, and external fixators.^{2,7,8,16}

Fig. 1. Radiographic characteristics of a 12-year-old female patient: (1A) Osteogenesis imperfecta proximal femoral fractures with severe coxa vara; (1B) After healing of the right proximal femur, the neck-shaft angle decreased to 60 degrees, but the left hip had delayed union; (1C) Valgus osteotomy combined with a Sofield-Millar operation using a pediatric dynamic hip screw and titanium elastic nail performed at the left hip, then at the right hip 3 months later. At the 2-year follow-up, the patient could walk independently without aid and had a neck-shaft angle of 130 degrees.

Fig. 2. Radiographic characteristics of a 30-year-old female: (2A) Monostotic fibrous dysplasia with coxa vara and femoral fracture; (2B) Valgus osteotomy combined with a Sofield-Millar operation using gamma nail with plate and screw augmentation was performed at the left hip. At 3-year follow-up, the patient could walk independently without aid and had a neck-shaft angle of 130 degrees.

Fig. 3. Radiographic characteristics of a 18-year-old female patient: (3A) Fibrous dysplasia with coxa vara; valgus osteotomy using a gamma nail was performed at the right hip. At the 3-year follow-up, the patient could walk independent without aid and had a neck-shaft angle of 130 degrees. (3B).

Recurrence of the deformity is the most important problem following corrective surgery and has been reported at rates varying from 30 to 70%.^{2,3,8,10,17–21} Carroll et al⁹ stated that a high recurrence may be related to the altered hip biomechanics. When the proximal femoral physis is in a vertical position, the mechanical forces across the hip joint change, leading to a progression of coxa vara as well as osteoarthritis. Many authors have agreed that appropriate correction of the Hilgenreiner-epiphyseal angle is the most crucial factor in prevention of the recurrence.^{9,18,19} Desai and Johnson¹⁹ claimed that no recurrence occurred in a case series of 12 hips with a postoperative Hilgenreiner-epiphyseal angle of ≤ 35 degrees and a neck-shaft angle of ≥ 130 degrees, irrespective of the patient's age at surgery. Cordes et al¹⁸ studied 14 cases and described excellent clinical results, regardless to the etiology of coxa vara, when the Hilgenreiner-epiphyseal angle of ≤ 40 degrees was achieved. Carroll et al⁹ found that when the Hilgenreiner-epiphyseal angle was corrected to less than 38 degrees, 95% of their studied group had no recurrence of varus deformity. They also found that the recurrence rate was not affected by etiology of deformity, type of implant, type of surgery, and the age at surgery. A study by Burns

and Stevens¹ concurrently emphasized the importance of the correction of the Hilgenreiner-epiphyseal angle to less than 38 degrees.

In this studied group, we were able to improve the mean Hilgenreiner-epiphyseal angle and neck-shaft angle at the last follow up to 29.26 and 129.64 degrees, respectively. The maximum postoperative value of the Hilgenreiner-epiphyseal angle was 35 degrees, and none of these cases had recurrence. Comparing the postoperative images to the final follow-up images, we found no statically significant difference in neck-shaft angles and Hilgenreiner-epiphyseal angles, implying that the correction was maintained throughout the follow-up period of 37.84 months. The postoperative Harris hip scores improved significantly at the final follow up, indicating improved functional activity of the patients after corrective treatment.

Based on our findings, surgical correction of coxa vara in various bone diseases with a Hilgenreiner-epiphyseal angle correction of ≤ 35 degrees and a neck-shaft angle of > 120 degrees produced good short-term results. However, a longer-term observation is necessary to determine further outcomes with respect the recurrence of the deformity.

Table 1
Demographic and clinical data of coxa vara patients.

Case	Gender	Hip	Diagnosis	Age at surgery (year)	Healing time (months)	Follow-up (months)	Ambulatory Status Pre-op	Ambulatory Status post-op	fixation
1	F	Right	OI	12	3	36	Wheel chair	Independent walk	DHS + titanium elastic nail
2	F	Right	OI	13	3	26	Wheel chair	Independent walk	DHS + titanium elastic nail
3	M	Left			5				
4	F	Right	Fibrous dysplasia	9	3	30	crutches	Independent walk	DHS
5	M	Right	Histiocytosis X	5	2	40	crutches	Independent walk	DHS
6	M	Right	OI	5	3	24	crutches	Independent walk	DHS + titanium elastic nail
7	M	Right	Fibrous dysplasia	6	2	28	crutches	Independent walk	DHS
8	F	Left	OI	11	4	30	crutches	Independent walk	DHS
9	F	Right	OI	8	3	40	Independent walk	Independent walk	ABP
10	F	Right	Monostotic fibrous dysplasia	30	3	36	Independent walk	Independent walk	Gamma nail
11	F	Left	polyostotic fibrous dysplasia	28	5	30	Wheel chair	Independent walk	Gamma nail
12	M	Right	Histiocytosis X	5	2	40	crutches	Independent walk	DHS
13	M	Right	OI	10	3	24	crutches	Independent walk	DHS + titanium elastic nail
14	M	Right	Fibrous dysplasia	7	2	28	crutches	Independent walk	DHS
15	M	Left	OI	13	4	30	crutches	Independent walk	DHS + titanium elastic nail
16	F	Right	OI	8	3	40	Independent walk	Independent walk	DHS + titanium elastic nail
17	F	Right	Monostotic fibrous dysplasia	18	3	36	Independent walk	Independent walk	Gamma nail + plate and screws
18	F	Left	polyostotic fibrous dysplasia	15	5	30	Wheel chair	Independent walk	Gamma nail
19	M	Right	Fibrous dysplasia	14	2	28	crutches	Independent walk	DHS
20	M	Right	OI	13	3	24	crutches	Independent walk	DHS + titanium elastic nail
21	M	Right	Fibrous dysplasia	6	2	28	Crutches	Independent walk	DHS
22	M	Right	SED	7	2	40	Independent walk	Independent walk	ABP
23	M	Left	SED	7.4	2	42	Independent walk	Independent walk	ABP
24	M	Left	MED	10.5	3	65	Independent walk	Independent walk	ABP
25	F	Right	MFNF	12	3	44	Crutches	Independent walk	ABP
26	F	Left	MFNF	11.6	2	40	Crutches	Independent walk	ABP
27	M	Right	MFNF	10	3	67	Crutches	Independent walk	ABP
28	M	Right	MFNF	8	3	36	Crutches	Independent walk	ABP
29	M	Left	CCV	11.5	3	41	Independent walk	Independent walk	ABP
30	M	Left	CCV	9	3	84	Independent walk	Independent walk	ABP
31	M	Left	CCV	10	2	50	Independent walk	Independent walk	ABP
32	F	Right	CCV	7.6	3	37	Independent walk	Independent walk	ABP
Mean ± SD		Left		10.99 ± 5.75	3 ± 0.94	37.87 ± 13.40			

Abbreviations: OI: Osteogenesis imperfecta, SED: Spondyloepiphyseal dysplasia, MED: Multiple epiphyseal dysplasia, MFNF: Malunion femoral neck fracture, CCV: Congenital coxa vara, ABP: Angle Blade plate.

Table 2
Radiographic data and ambulatory status of Coxa Vara patients.

Case	NSA (pre-op)	NSA (post-op)	NSA (final follow-up)	HEA (pre-op)	HEA (post-op)	HEA (final follow-up)	HHS (pre-op)	HHS (final follow-up)
1	Right 60° Left 90°	Right 130° Left 130°	Right 130° Left 130°	Right 90° Left 70°	Right 28° Left 30°	Right 27° Left 28°	34	78
2	Right 85° Left 100°	Right 130° Left 130°	Right 130° Left 130°	Right 70° Left 75°	Right 25° Left 22°	Right 28° Left 21°	41	60
3	90°	130°	130°	65°	26°	28°	50	82
4	100°	135°	135°	60°	27°	28°	40	75
5	85°	130°	130°	60°	25°	26°	52	78
6	100°	135°	135°	60°	28°	30°	49	78
7	80°	130°	130°	70°	22°	22°	66	82
8	85°	130°	130°	60°	25°	26°	50	68
9	100°	135°	135°	60°	28°	30°	49	85
10	100°	135°	135°	60°	28°	30°	45	89
11	80°	130°	130°	70°	22°	22°	43	81
12	90°	130°	130°	65°	26°	28°	41	68
13	100°	135°	135°	60°	27°	28°	43	81
14	85°	130°	130°	60°	25°	26°	41	75
15	90°	130°	130°	65°	26°	28°	50	78
16	100°	135°	135°	60°	27°	28°	40	85
17	85°	130°	130°	60°	25°	26°	41	89
18	100°	135°	135°	60°	28°	30°	45	78
19	80°	130°	130°	70°	22°	22°	43	81
20	85°	130°	130°	60°	25°	26°	41	75
21	Rt.80° Lt.85°	131° 135°	130° 135°	75° 80°	30° 30°	31° 31°	55	78
22	85°	125°	130°	75°	35°	35°	50	68
23	65°	125°	125°	73°	28°	30°	60	85
24	69°	123°	124°	70°	30°	32°	50	90
25	70°	120°	123°	62°	33°	35°	43	81
26	85°	124°	124°	61°	35°	35°	41	68
27	65°	125°	125°	65°	35°	35°	55	90
28	92	125	127	65	35	37	41	75
29	93	120	122	68	32	33	60	100
30	95	123	123	66	35	37	60	80
31	73	120	125	86	33	35	45	89
Mean ± SD	86.09 ± 11.13	128.26 ± 6.59	129.64 ± 3.95	66.76 ± 7.99	28.09 ± 3.92	29.26 ± 4.37	47.23 ± 7.46	76.68 ± 8.22
P value	<0.001			<0.001			<0.001	

Abbreviations: NSA: neck-shaft angle; HEA: Hilgenreiner-epiphyseal angle, HHS: Harris hip score.

5. Conclusion

The surgical treatment of femoral coxa vara to achieve Hilgenreiner-epiphyseal angle ≤ 35 degrees and neck-shaft angle > 120 degrees can prevent recurrence regardless of the cause of the deformity. The aim of treatment was to modify the femoral neck anatomy to normalize the forces around the hip joint and reduce the incidence of further stress fracture and osteoarthritis.

Funding disclosure

This was an unfunded study.

Ethical approval

This article does not contain any studies with human participants or animals performed by any of the authors.

Conflict of interest declaration

The authors hereby declare no personal or professional conflicts of interest regarding any aspect of this study.

Informed consent

Informed consent was obtained from all individual participants included in the study.

Acknowledgements

The authors of this study gratefully acknowledge Suchitphon Chanchoo for assistance with data collection and statistical analysis.

References

1. Burns KA, Stevens PM. Coxa vara: another option for fixation. *J Pediatr Orthop B*. 2001;10(October (4)):304–310.
2. Weighill FJ. The treatment of developmental coxa vara by abduction subtrochanteric and intertrochanteric femoral osteotomy with special reference to the role of adductor tenotomy. *Clin Orthop Relat Res*. 1976;(May (116)):116–124.
3. Chotigavanichaya C, Leepakoboon D, Eamsobhana P, Kaewpornawan K. Results of surgical treatment of coxa vara in children: valgus osteotomy with angle blade plate fixation. *J Med Assoc Thai*. 2014;97(September (Suppl. 9)):S78–S82.
4. Key JA. The classic: epiphyseal coxa vara or displacement of the capital epiphysis of the femur in adolescence. 1926. *Clin Orthop Relat Res*. 2013;471(July (7)):2087–2117.
5. Elzohairy MM, Khairy HM. Fixation of intertrochanteric valgus osteotomy with T Plate in treatment of developmental coxa vara. *Clin Orthop Surg*. 2016;8(September (3)):310–315.
6. Aarabi M, Rauch F, Hamdy RC. High prevalence of coxa vara in patients with severe osteogenesis imperfecta. *J Pediatr Orthop*. 2006;26(January– February (1)):24–28.
7. Amstutz HC, Frieberger RH. Coxa vara in children. *Clin Orthop*. 1962;22:73–92.
8. Amstutz HD, Wilson PD. Dysgenesis of the proximal femur (coxa vara) and its surgical management. *J Bone Jt Surg Am*. 1962;44-A(January):1–24.
9. Carroll K, Coleman S, Stevens PM. Coxa vara: surgical outcomes of valgus osteotomies. *J Pediatr Orthop*. 1997;17(March–April (2)):220–224.
10. Weinstein J, Kuo K, Millar E. Congenital coxa vara. A retrospective review. *J Pediatr Orthop*. 1984;4(January (1)):70–77.
11. Serafin J, Szulc W. Coxa vara infantum, hip growth disturbances, etiopathogenesis, and long term results of treatment. *Clin Orthop Relat Res*. 1991;(November (272)):103–113.
12. Eamsobhana P, Kaewpornawan K. Nonunion paediatric femoral neck fracture treatment without open reduction. *Hip Int*. 2016;26(November (6)):608–611.
13. Borden J, Spencer Jr. GE Jr., Herndon CH. Treatment of coxa vara in children by means of a modified osteotomy. *Bone Jt Surg Am*. 1966;48(September (6)):1106–1110.
14. Freeman BH, Bray EW, Meyer LC. Multiple osteotomies with Zickel nail fixation for polyostotic fibrous dysplasia involving the proximal part of the femur. *J Bone Jt Surg Am*. 1987;69(June (5)):691–698.
15. Pauwels F. *Biomechanics of the normal and diseased hip. Theoretical foundation, technique and results of treatment*. Berlin, Heidelberg, New York: Springer-Verlag; 1976:55–80.
16. Dobbs M, Morcuende J. Coxa vara. In: Morrissy RT, Weinstein SL, eds. *Lovell & winter's pediatric orthopaedics*. Philadelphia, PA: Lippincott-Raven Publishers; 2006:1126–1134.
17. Widmann RF, Hresko MT, Kasser JR. Wagner multiple K-wire osteosynthesis to correct coxa vara in the young child: experience with a versatile 'tailor-made' high angle blade plate equivalent. *J Pediatr Orthop B*. 2001;10(January (1)):43–50.
18. Cordes S, Dickens DR, Cole WG. Correction of coxa vara in childhood. The use of Pauwels' Y-shaped osteotomy. *J Bone Jt Surg Br*. 1991;73(January (1)):3–6.
19. Desai SS, Johnson LO. Long term results of valgus osteotomy for congenital coxa vara. *Clin Orthop Relat Res*. 1993;(September (294)):204–210.
20. Livesley PJ, McAllister JC, Catterall A. The treatment of progressive coxa vara in children with bone softening disorders. *Int Orthop*. 1994;18(October (5)):310–312.
21. Fassier F, Sardar Z, Aarabi M, Odent T, Haque T. Results and complications of a surgical technique for correction of coxa vara in children with osteopenic bones. *J Pediatr Orthop*. 2008;28(December (8)):799–805.