Published in final edited form as:

Angew Chem Int Ed Engl. 2019 May 13; 58(20): 6654–6658. doi:10.1002/anie.201901927.

Catalytic Conjunctive Coupling of Carboxylic Acid Derivatives with 9-BBN-Derived Ate Complexes

Chunyin Law, Yan Meng, Seung Moh Koo, and Prof. James P. Morken

Department of Chemistry, Boston College, 2609 Beacon Street, Chestnut Hill, MA 02467, USA

Abstract

β -Boryl carbonyl compounds are produced by a Ni-catalyzed cross-coupling of vinylboron “ate” complexes and acid chloride or acid anhydride electrophiles. Reactions are efficient, being complete in as little as two minutes, and apply to a broad range of substrates.

Graphical Abstract

Conjunctive Coupling: A new route to synthetically useful β -boryl carbonyls is enabled by the Ni-catalyzed addition of alkenyl boron “ate” complexes to activated carboxylic acids.

Keywords

Boron; Cross-Coupling; Nickel; Catalysis

Organoboron compounds can be prepared by transition-metal-catalyzed conjunctive cross-coupling reactions between alkenylboron-derived “ate” complexes and organic electrophiles (eq. 1).^{1,2,3} These reactions can be accomplished in an enantioselective and diastereoselective fashion, but so far have been restricted to hydrocarbon (aryl, alkenyl, alkyl) electrophiles. To extend conjunctive coupling reactions to the preparation of alternatively functionalized motifs, we considered the use of carbonyl-derived electrophiles. Herein, we describe a process that employs carboxylic acid derivatives as the electrophile in conjunctive couplings and produces β -boryl carbonyls as the reaction product.⁴ Considering the synthesis utility of β -boryl carbonyls, compounds that are generally prepared by boron conjugate addition⁵, the alternate route to these materials that is presented herein may have value.

Our preliminary direction in developing a conjunctive-coupling reaction with carbonyl electrophiles was largely inspired by the body of work by Rovis⁶, Gong⁷, Weix⁸, Reisman⁹, Doyle¹⁰, and Yamamoto¹¹ who determined that Ni complexes could undergo oxidative

Dedicated to Professor Amir Hoveyda on the occasion of his 60th birthday.

Supporting information for this article is given via a link at the end of the document.

addition with activated carboxylic acid derivatives thereby enabling either catalytic cross-coupling with nucleophiles or, in the presence of stoichiometric reductant, catalytic cross-electrophile coupling. Importantly, ^{13}C labeling experiments conducted by Rovis and Coates¹² established that (bipyridine)Ni(acyl) complexes were prone to undergo direct transmetalation reaction with alkylzinc reagents; this observation provided optimism that such a system might facilitate catalytic acyl conjunctive coupling reactions, as well.

To develop the conjunctive coupling between carbonyl electrophiles and boron “ate” complexes, we investigated the reaction between butanoyl chloride and various *in situ*-generated vinylboron ate complexes. Whereas preliminary experiments with pinacoloboron-derived reagents were not fruitful, use of 9-BBN borate complex **1** as a coupling partner furnished the coupling product: as depicted in Table 1, it was observed that after one hour of reaction in the presence of 5 mol% $\text{NiBr}_2\cdot\text{glyme}$ and 6 mol% bipyridine (**L1**), followed by oxidation, β -hydroxy carbonyl **2** could be obtained in 65% yield (Table 1, entry 1). Importantly, under these conditions, background reaction between the acid chloride and the “ate” complex was minimal. Extensive ligand analysis (see Supporting Information) revealed that the most generally applicable reaction conditions employ 5,5'-bis(trifluoromethyl)-2,2'-bipyridine (**L3**) as the ligand (Table 1, entry 7). While the more readily available ligand bipyridine (**L1**) operates effectively with aliphatic acid chloride electrophiles, markedly diminished yield was obtained when this ligand was employed with aromatic acid chloride electrophiles. Thus, ligand **L3** was employed for subsequent studies. It was determined that reactions in the presence of **L3** can proceed with very good efficiency, achieving complete conversion of the probe reaction within two minutes at room temperature (entry 8).

To survey the scope of this reaction, a range of 9-BBN derived borate complexes were prepared by hydroboration¹³ of the corresponding alkenes in THF, followed by addition of vinylolithium.¹⁴ The so-formed “ate” complex was treated with pre-mixed $\text{NiBr}_2\cdot\text{glyme}$ and **L3** in the presence of acid chloride electrophiles, followed by stirring at room temperature for 1 h and oxidative workup. As seen in Table 2, the reaction with aliphatic unbranched (i.e., **3**, **6**, **8**) and branched (**4**, **5**, **7**) acyl chlorides afforded products in good yields, although only moderate yield was observed when the sterically demanding pivaloyl chloride (**7**) was employed in the reaction. Acyl chloride derived from TBS-protected lithocholic acid (**20**) was also subjected to the reaction conditions and afforded the desired product in reasonable yield. Regarding the functional group tolerance with respect to acyl chloride electrophiles, whereas benzoyl chloride (**10**) as well as substrates bearing electron-donating substituents in the para-position (**11**, **14**) resulted in good yields, diminished yield was observed with an electron-withdrawing group (**13**) in the same position. Migrating groups derived from various alkenes were also examined in the reaction. Secondary migrating groups such as cyclohexyl group (**16**) provided the desired product in reasonable yield with no observation of byproducts derived from the migration of the bicyclooctyl ligand of 9-BBN.¹⁵ Migrating groups with appended alkene (**21**), and acetal (**22**), as well as silyl ether (**18**) were also tolerated. It was also shown that the 9-BBN borate complex derived from Ph-9-BBN (prepared from phenylmagnesium bromide and 9-BBN-OMe) can be employed in the reaction to give the corresponding product (**23**) in good yield. Lastly, under the current

reaction conditions, propenyllithium-derived ‘ate’ complexes have not furnished coupling products.

The compatibility of other classes of acyl electrophiles in conjunctive cross-coupling was also explored. When anhydrides were subjected to the coupling under the conditions of Table 2, only trace quantities of coupled product were observed. We reasoned that the oxidative addition of Ni to less electrophilic anhydrides may be less facile as compared to the reaction with acid chloride counterparts¹⁶, and this process might be facilitated with a more electron-rich metal center. In line with this hypothesis, the use of unsubstituted bipyridine (**L1**) in place of the bis(trifluoromethyl) version resulted in a much more efficient coupling (Scheme 2). Of note, this reaction extended to a mixed anhydride with coupling occurring with the less hindered carbonyl (inset, Scheme 2).¹⁷

While mechanistic aspects of this process remain under study, preliminary experiments shed light on the rudiments of the underlying features. By subjecting isotopically-labelled boron “ate” complex (**25**)¹⁸ to the reaction conditions (eq. 1, Scheme 3), it was found that the Ni-catalyzed coupling reaction occurs stereospecifically and with *anti* addition of the migrating group and the electrophile to the alkene, giving **26**. This observation is consistent with a mechanism involving nickel-induced metallate shift as opposed to alternate one-electron processes that might involve acyl radical addition to the alkene.¹⁹ To probe whether a Ni⁰-Ni^{II} redox cycle might represent the operative mechanism, the oxidative addition adduct **27** was independently synthesized using the method reported by Gong⁷, and treated with allylbenzene-derived borate complex **1**. As depicted in equations (2) and (3), the stoichiometric experiment furnished the conjunctive coupling product in 43% yield, suggesting that the Ni^{II} oxidative addition adduct is a competent intermediate in the conjunctive cross-coupling reaction. Lastly, as depicted in equation (4), with a chiral, enantiomerically-enriched bis(oxazoline) ligand (**L6**), a modest dependence of reaction enantioselectivity on the identities of electrophile was observed. This observation is suggestive of a scenario in which the electrophile is associated with the metal center before a stereochemistry-determining, and likely irreversible, 1,2-metallate-shift. Thus, observations about the described cross coupling support a mechanism involving oxidative addition of the carbonyl electrophile to a Ni(0) complex²⁰ (**A**→**B**, Scheme 3, box), followed by binding of the ate complex, metallate shift, and reductive elimination to deliver the product.

Although β -(pinacolato)boryl ketones have received ample study in the chemical synthesis community, β -trialkyl(boryl) ketones are novel species and a study of their structure and chemistry was of interest. The ¹¹B NMR resonance of β -boryl ketone **28** (δ 22.0 ppm) exhibits a significant upfield shift relative to a non-functionalized trialkylborane (δ 70–80 ppm).²¹ This feature is consistent with tight coordination between the boron and carbonyl oxygen atom. Reasoning that this chelation would enhance the acidity of the carbonyl α -protons and allow for the formation of a rigid cyclic enolate structure, we examined deprotonation/alkylation reactions of **28** (Scheme 4). Initial attempts with LDA or (*i*-Pr)₂NEt (see SI) provided no evidence of productive deprotonation; however, when **28** was treated with methyllithium followed by iodomethane, alkylation product **29** was obtained with a high level of *anti* diastereomeric purity.²² Remarkably, products derived from nucleophilic addition of CH₃Li to the internally-activated ketone were not detected. To understand the

origin of reactivity and stereoselectivity in the deprotonation/alkylation reaction, we attempted to obtain a crystal structure of **28**. While these attempts were unsuccessful, x-ray quality crystals of the methylamine-derived Schiff base were obtained (**31**) and these reveal internal chelation between boron and the heteroatom. Of note, due to the steric demands of the bicyclo[3.3.1]nonane framework, the substituent (Cy) at the stereogenic carbon atom is directed in an axial orientation and appears to block access to one face of the prochiral α -carbon atom. It appears plausible that an enolate derived from **28** would have a similar conformation and this would allow an electrophile to approach only from the less hindered face, thereby accounting for the excellent *anti* diastereoselectivity.²³

In conclusion, we have described a catalytic cross-coupling between four-coordinate boron “ate” complexes and carbonyl electrophiles. This process provides a simple route to β -trialkylboron carbonyls which exhibit uncommon and synthetically useful reactivity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by a grant from the US National Institutes of Health (NIGMS GM-R35-127140).

References

- [1]. For reports from these labs, see a)Zhang L, Lovinger GJ, Edelstein EK, Szymaniak AA, Chierchia MP, Morken JP, Science 2016, 351, 70; [PubMed: 26721996] b)Lovinger GJ, Aparece MD, Morken JP, J. Am. Chem. Soc 2017, 139, 3153; [PubMed: 28161943] c)Edelstein EK, Namirembe S, Morken JP, J. Am. Chem. Soc 2017, 139, 5027; [PubMed: 28335597] d)Chierchia MP, Law C, Morken JP, Angew. Chem. Int. Ed 2017, 56, 11870;e)Lovinger GJ, Morken JP, J. Am. Chem. Soc 2017, 139, 17293; [PubMed: 29116784] f)Myhill JA, Zhang L, Lovinger GA, Morken JP, Angew. Chem. Int. Ed 2018, 57, 12799;g)Myhill JA, Whilhelmsen CA, Zhang L, Morken JP, J. Am. Chem. Soc 2018, 140, 15181. [PubMed: 30376317]
- [2]. For related catalytic reactions involving a metallate shift, see a)Ishida N, Narumi M, Murakami M, Org. Lett 2008, 10, 1279; [PubMed: 18271597] b)Ishida N, Shimamoto Y, Murakami M, Org. Lett 2009, 11, 5434; [PubMed: 19877621] c)Ishida N, Shinmoto T, Sawano S, Miura T, Murakami M, Bull. Chem. Soc. Jpn 2010, 83, 1380;d)Ishida N, Shimamoto Y, Murakami M, Org. Lett 2010, 12, 3179; [PubMed: 20557031] e)Ishida N, Ikemoto W, Narumi M, Murakami M, Org. Lett 2011, 13, 3008; [PubMed: 21604778] f)i) Panda S, Ready JM, J. Am. Chem. Soc 2017, 139, 6038; [PubMed: 28414430] g)Panda S, Ready JM, J. Am. Chem. Soc 2018, 140, 13242; [PubMed: 30240211] h)Tao Z, Robb KA, Panger JL, Denmark SE, J. Am. Chem. Soc 2018, 140, 15621; [PubMed: 30411878] i)Fawcett A, Biberger T, Aggarwal VK, Nat. Chem 2019, 11, 117. [PubMed: 30532013]
- [3]. For non-catalytic electrophile-induced metallate shifts, see a)Zweifel G, Arzoumanian H, Whitney CC, J. Am. Chem. Soc 1967, 89, 3652;b)Evans DA, Crawford TC, Thomas RC, Walker JA, J. Org. Chem 1976, 41, 3947; [PubMed: 993895] c)Utimoto K, Uchida K, Nozaki H, Tetrahedron Lett 1973, 45, 4527;d)Armstrong RJ, Sandford C, García-Ruiz C, Aggarwal VK, Chem. Commun 2017, 53, 4922;e)Armstrong RJ, García-Ruiz C, Myers EL, Aggarwal VK, Angew. Chem. Int. Ed 2017, 56, 786;f)Silvi M, Sandford C, Aggarwal VK, J. Am. Chem. Soc 2017, 139, 5736; [PubMed: 28402109] g)Kischkewitz M, Okamoto K, Mück-Lichtenfeld C, Studer A, Science 2017, 355, 936 For recent related processes, see [PubMed: 28254937] h)Wilson CM, Ganesh V, Noble A, Aggarwal VK, Angew. Chem. Int. Ed 2017, 56, 16318.i)Ganesh V, Odachowski M, Aggarwal VK, Angew. Chem. Int. Ed 2017, 56, 9752.j)Wang Y, Noble A, Sandford C, Aggarwal VK, Angew. Chem. Int. Ed 2017, 56, 1810.

- [4]. For non-catalyzed metallate shifts with carbonyl electrophiles, see a) Utimoto K, Uchida K, Nozaki H, *Chemistry Lett* 1974, 1493; b) Utimoto K, Uchida K, Nozaki H, *Tetrahedron* 1977, 33, 1949; c) Deng M-Z, Lu D-A, Xu W-H, *J. Chem. Soc., Chem. Commun* 1985, 1478
- [5]. For reports of catalytic enantioselective boron conjugate addition, see a) Lee J-E, Yun J, *Angew. Chem. Int. Ed* 2008, 47, 145; b) Chea H, Sim H-S, Yun J, *Adv. Synth. Catal* 2009, 351, 855; c) Chen I-H, Yin L, Itano W, Kanai M, Shibasaki M, *J. Am. Chem. Soc* 2009, 131, 11664; [PubMed: 19653692] d) Lillo V, Prieto A, Bonet A, Díaz-Requejo MM, Ramírez P J, Pérez J, Fernández E, *Organometallics* 2009, 28, 659; e) Chen I-H, Kanai M, Shibasaki M, *Org. Lett* 2010, 12, 4098; [PubMed: 20722382] f) O'Brien JM, Lee K-S, Hoveyda AH, *J. Am. Chem. Soc* 2010, 132, 10630; [PubMed: 20681680] g) Park JK, Lackey HH, Rexford MD, Kovnir K, Shatruk M, McQuade DT, *Org. Lett* 2010, 12, 5008; [PubMed: 20919706] h) Schiffner J A, Muther K, Oestreich M, *Angew. Chem. Int. Ed* 2010, 49, 1194; i) Kobayashi S, Xu P, Endo T, Ueno M, Kitanosono T, *Angew. Chem. Int. Ed* 2012, 51, 12763; j) Zhao L, Ma Y, He F, Duan W, Chen J, Song C, *J. Org. Chem* 2013, 78, 1677; [PubMed: 23330603] k) Wu H, Garcia JM, Haefner F, Radomkit S, Zhugralin AR, Hoveyda AH, *J. Am. Chem. Soc* 2015, 137, 10585 For a comprehensive review on boron conjugate addition see: [PubMed: 26263513] l) Collins BSL, Wilson CM, Myers EL, Aggarwal VK, *Angew. Chem. Int. Ed* 2017, 56, 11700.
- [6]. Zhang Y, Rovis T, *J. Am. Chem. Soc* 2004, 126, 15964. [PubMed: 15584721]
- [7]. Zhao C, Jia X, Wang X, Gong H, *J. Am. Chem. Soc* 2014, 136, 17645. [PubMed: 25415424]
- [8]. a) Wotal AC, Weix DJ, *Org. Lett* 2012, 14, 1476; [PubMed: 22360350] b) Wotal AC, Ribson RD, Weix DJ, *Organometallics* 2014, 33, 5874. [PubMed: 25364092]
- [9]. Cherney AH, Kadunce NT, Reisman SE, *J. Am. Chem. Soc* 2013, 135, 7442. [PubMed: 23634932]
- [10]. Joe CL, Doyle AG, *Angew. Chem. Int. Ed* 2016, 55, 4040.
- [11]. Sano K, Yamamoto T, Yamamoto A, *Chem. Lett* 1983, 12, 115.
- [12]. Johnson JB, Bercot EA, Rowley JM, Coates GW, Rovis T, *J. Am. Chem. Soc* 2007, 129, 2718. [PubMed: 17295486]
- [13]. For detailed studies on hydroboration using 9-BBN, see: Brown HC, Chen JC, *J. Org. Chem* 1981, 46, 3978.
- [14]. Vinyl lithium was prepared from n-BuLi and tetravinylstannane according to the reported procedure from ref. [1a].
- [15]. Similar chemoselectivity was observed in ref. [1d]. For a discussion of factors affecting the selectivity in migration with mixed borates, see a) Aggarwal VK, Fang GY, Ginesta X, Howells DM, Zaja M, *Pure Appl. Chem* 2006, 78, 215; b) Bottoni A, Lombardo M, Neri A, Trombini C, *J. Org. Chem* 2003, 68, 3397. [PubMed: 12713337]
- [16]. For an example that suggests a correlation between oxidative addition rates and carbonyl electrophilicity, see a) Meng G; Szostak M, *Org. Biomol. Chem* 2016, 14, 5690. [PubMed: 26864384]
- [17]. For a similar example of selective oxidative addition of mixed anhydride with nickel catalyst, see ref. 10.
- [18]. Deuterium-labelled vinyl lithium was prepared according to the reported procedure from Ref. [1a]. See Supplementary Material for details.
- [19]. For reactions involving radical addition to an alkenyl boron "ate" complex: See ref [3f], [3g].
- [20]. Fahey DR, Mahan JE, *J. Am. Chem. Soc* 1977, 99, 2501.
- [21]. For a comprehensive list of 11 BNMR chemical shifts, see: Wrackmeyer B, *Annu. Rep. NMR Spectrosc*, 1988, 20, 61.
- [22]. The relative stereochemistry was determined by comparison of ¹H NMR spectrum with the anti diastereomer prepared through independent synthesis. See Supplementary Material for details.
- [23]. For mechanistically related chelation-controlled enolate alkylations, see a) Fráter G, Müller U, Günther W, *Tetrahedron* 1984, 40, 1269; b) Seebach D, Wasmuth D, *Helv. Chim. Acta* 1980, 63, 197.

Previous Work: Hydrocarbon Electrophiles **This work: Carbonyl Electrophiles** **Scheme 1.**Catalytic Synthesis of β -Boryl Carbonyls by Conjunctive Coupling.

Scheme 2.

Ni-Catalyzed Conjunctive Coupling with Anhydride Electrophiles.

Scheme 3.

Experimental observations about conjunctive coupling with acyl electrophiles and proposed mechanism.

Angew Chem Int Ed Engl. Author manuscript; available in PMC 2020 May 13.

Table 1.Effect of Ligand on Conjunctive Coupling with Acid Chloride Electrophiles.^[a]

Entry	R	Time	Temp (°C)	Ligand	Yield (%)
1	propyl	1 h	22	L1	65
2	propyl	1 h	22	none	<5
3	propyl	1 h	40	L1	48
4	propyl	1 h	22	L2	72
5	Ph	1 h	22	L1	40
6	Ph	1 h	22	L3	75
7	propyl	1 h	22	L3	78
8	propyl	2 min	22	L3	75
9	propyl	1 h	22	L4	32
10	propyl	1 h	22	L5	40

^[a] See text and Supporting Information for experimental details. Yields refer to isolated yield of purified material and are an average of two experiments.

Table 2.

Ni-Catalyzed Conjunctive Coupling with Various Alkenes and Acid Chlorides.^[a]

^[a] See text and Supporting Information for experimental details. Yields refer to isolated yield of purified material and are an average of two experiments.

^[b] Starting material prepared from 9-BBN-OMe and $PhMgBr$.