CORRESPONDENCE

The Investigation of Hematuria

by Prof. Dr. med. Christian Bolenz, Prof. Dr. med. Bernd Schröppel, PD Dr. med. Andreas Eisenhardt, Prof. Dr. med. Bernd J. Schmitz-Dräger, and Prof. Dr. med. Marc-Oliver Grimm
in issue 48/2018

Endometriosis Is a Possible Cause

I read the article on investigating hematuria with great interest (1). Unfortunately, one cause was not mentioned, although it occurs at a rate that is relevant: deeply infiltrating endometriosis of bladder or ureter.

Reliable data on rates hardly exist, but according to estimates, 1–2% of patients with endometriosis have urogenital endometriosis. The symptoms depend on a woman's monthly cycle, but in advanced findings they can occur independently of the monthly cycle. For this reason, this cause should always be considered in premenopausal women with hematuria, and patients should be referred for a gynecological examination.

DOI: 10.3238/arztebl.2019.0191a

References

1. Bolenz C, Schröppel B, Eisenhardt A, Schmitz-Dräger BJ, Grimm MO: The investigation of hematuria. *Dtsch Arztebl Int* 2018; 115: 801–7.

Dr. med. Christoph Therre

Klinisch wissenschaftliches Endometriosezentrum
Albertinenkrankenhaus, Hamburg
christoph.therre@albertinen.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Consider Urogenital Schistosomiasis and Tuberculosis

With regard to the interesting article on investigating hematuria, we would like to focus on infectious causes—which for reasons of space were only mentioned in the table—of apparently “sterile” hematuria, especially in refugees and migrants (1). In this setting, two infectious diseases are relevant whose diagnosis is not always easy: urogenital schistosomiasis and urogenital tuberculosis. A study in apparently healthy minor refugees from sub-Saharan Africa found a prevalence of schistosomiasis of almost 25% (2). Furthermore, since 2014 it has been known that bladder schistosomiasis can also once again be acquired in Europe—namely, in south Corsica (3). The suspicion of schistosomiasis can be supported in a laboratory that has the relevant specialization; it is confirmed microscopically by proving helminthic eggs in microfiltrated urine (collected from 11.00 am to 14.00 pm or collected over 24 hours).

We are also frequently confronted with refugees with urogenital tuberculosis who were not captured by admission screening for tuberculosis. The reason is that

tuberculosis that was acquired outside Europe often takes an extrapulmonary course and is not seen on a chest x-ray (4). In addition to a specific interferon gamma release alpha (IGRA) assay, which often yields positive results in immigrants even without clinical manifestations, the suspicion of productive tuberculosis is supported by a highly positive reaction to the intracutaneous Mendel-Mantoux skin test. The absence of acid-fast bacilli in urine or tissue specimens and false-negative specific polymerase chain reaction may mislead clinicians. The culture, which has to be grown from concentrated morning urine, can take several weeks to become positive. DOI: 10.3238/arztebl.2019.0191b

References

1. Bolenz C, Schröppel B, Eisenhardt A, Schmitz-Dräger BJ, Grimm MO: The investigation of hematuria. *Dtsch Arztebl Int* 2018; 115: 801–7.
2. Theuring S, Friedrich-Jänicke B, Pörtner K, et al.: Screening for infectious diseases among unaccompanied minor refugees in Berlin, 2014–2015. *Eur J Epidemiol* 2016; 31: 707–10.
3. Holtfreter MC, Moné H, Müller-Stöver I, Mouahid G, Richter J: Schistosoma haematobium infections acquired in Corsica, France, August 2013. *Euro Surveill* 2014; 19: pii: 20821.
4. Singh DD, Vogel M, Müller-Stöver I, et al.: TB or not TB? Difficulties in the diagnosis of tuberculosis in HIV-negative immigrants to Germany. *Eur J Med Res* 2011; 16: 381–4.

Prof. Dott. Joachim Richter, Gabriela Equihua-Martinez,

Dr. med. Kirsten Müller, Dr. med. Andreas K. Lindner

Institut für Tropenmedizin und Internationale Gesundheit,
Charité-Universitätsmedizin Berlin
Joachim.Richter@charite.de

Conflict of interest statement

Prof. Richter has received reimbursement of travel and accommodation expenses from Fiocruz, Brazil. The remaining signatories declare that no conflict of interest exists.

Consumption of Certain Types of Mushrooms

My thanks go to the authors of the clear article on micro- and macrohematuria (1). Allow me one additional point preceding the diagnostic evaluation: if patients notice that their urine is red—and they think it contains blood—they will be very concerned and therefore seek medical advice.

If subsequently urinalysis for blood yields a negative result in the practice setting or outpatient setting, and if no erythrocytes are seen on the microscope's field of vision, the doctor will consider myoglobinuria, porphyria, or, in particular, hemoglobinuria after excessive exercise (for example, “march” hemoglobinuria). What is also generally known is the fact that consumption of beetroot can stain urine red because beetroot contains betaine.

What is less well known, however, is the fact that ingestion of certain types of fungi/mushrooms can stain urine red. The first to mention in this context is the saffron milk cap mushroom (*Lactarius deliciosus*), which is very popular with mushroom pickers/foragers and which surfaces in late summer and autumn around pines.

The mushroom is in the family of Russulaceae. But all other types of mushrooms whose gills—when cut—release bright red-orange latex that eventually turns wine red and which are all edible (such as the *Lactarius salmonicolor* mushroom and the very common false saffron milk cap [*Lactarius deterrimus*] and others)—can cause red or brownish-red staining of the urine.

The reason for this is the azulene (sesquiterpene) content in the fruiting bodies of these mushrooms. During long years of working as an assistant and senior physician, I was able on several occasions to calm down worried patients by asking if they had eaten beetroot or *Lactarius* mushrooms on the previous day and spare them further diagnostic evaluation.

DOI: 10.3238/arztebl.2019.0191c

Reference

1. Bolenz C, Schröppel B, Eisenhardt A, Schmitz-Dräger BJ, Grimm MO: The investigation of hematuria. *Dtsch Arztebl Int* 2018; 115: 801–7.

Prof. Dr. med Siegmard Berndt

Toxikologe der Deutschen Gesellschaft für Mykologie e. V., Paderborn
 drs.berndt@t-online.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Algorithm Deviates From the Guideline

The review article entitled “The Investigation of Hematuria” provides well founded recommendations for urological practices for how to investigate macro- or microhematuria (1). These insights should be rapidly integrated into future guidelines and implemented accordingly in specialist practices.

At the primary care level (general practitioners) with its own epidemiological classification numbers/performance indicators, however, the recently revised guideline from the German College of General Practitioners and Family Physicians (DEGAM) provides an algorithm to inform and direct medical practice (2) that deviates from the recommendations in the article in the context of non-visible hematuria (“microhematuria”). In this guideline, the recommendation is for risk assessment on the basis of the medical history, focusing on rapid urine testing—opportunistic screening is rejected in accordance with international recommendations—on shared decision making between patient and doctor for further diagnostic evaluation. Patient information complements the guideline and is intended to convey the fundamentals in a clear fashion, so as to enable considered decision making.

(Increased) cooperation of the specialist disciplines on the basis of scientifically sound guidelines would make a sustained positive contribution to the quality of care.

DOI: 10.3238/arztebl.2019.0192a

References

1. Bolenz C, Schröppel B, Eisenhardt A, Schmitz-Dräger BJ, Grimm MO: The investigation of hematuria. *Dtsch Arztebl Int* 2018; 115: 801–7.
2. AWMF: Leitlinie „Hämaturie, nicht-sichtbare“. 053 – 028. Aktualisierte Fassung, Stand 2018. www.awmf.org/leitlinien/detail/II/053-028.html (last accessed on 19 February 2019).

Dr. med. Armin Mainz

Korbach
 dr.mainz@docduo.de

Conflict of interest statement

Dr. Mainz has received honoraria from the specialist society “German College of General Practitioners and Family Physicians” for developing guidelines.

In Reply:

We thank the correspondents for their contributions and additional comments with regard to investigating hematuria. It is important to consider—in addition to the potential causes of hematuria listed in Figure 1 (in the article)—further, even rare, causes. These include gynecological disorders, such as endometriosis, which can manifest in the urogenital tract, or urogenital atrophy, which may be accompanied by microhematuria that is reversible by applying topical/local estrogen. Furthermore, the medical history should help to exclude pseudohematuria as a result of the consumption of certain foods, as well as owing to false-positive results from the often highly sensitive rapid urine tests (1).

The revised version of the practice recommendations of the German College of General Practitioners and Family Physicians (DEGAM) has recently been published (2). The recommendations are based on rapid urine testing using a test strip. On the basis of comorbidities and established risk factors, the recommendation is to stratify patients for further investigation, in order to avoid overdiagnosis. In tandem, patient information on hematuria was developed and made available. This information may empower patients to make their own decision regarding referral to a urologist and undergoing further diagnostic tests.

We wish to add to Dr. Mainz’s comments that we do not recommend “opportunistic screening” for hematuria. But if we already are in possession of positive findings from test strips then we think that—in view if the high rate of false-positives mentioned earlier—it is important and logical to confirm the hematuria by using sediment analysis before undertaking further diagnostic tests. Pertinent diagnoses should not be missed even if their prevalence may be low. According to recent study data, in many patients, non-invasive ultrasound diagnostics—if indicated in combination with urethrocytostomy—provides a high degree of certainty in ruling out malignant causes (3). A graded/stepwise, risk adapted approach can help reduce overdiagnosis using complex medical apparatus and incurring unnecessarily high costs.

DOI: 10.3238/arztebl.2019.0192b

References

1. Rao PK, Gao T, Pohl M, Jones JS: Dipstick pseudohematuria: unnecessary consultation and evaluation. *J Urol* 2010; 183: 560–4.

2. Mainz A: Nicht-sichtbare Hämaturie – weniger ist mehr! Neue S1-Handlungsempfehlungen der Deutschen Gesellschaft für Allgemeinmedizin und Familienmedizin (DEGAM). Z Allg Med 2014; 90: 58.
3. Tan WS, Sarpong R, Khetrapal P, et al.: Can renal and bladder ultrasound replace CT urogram in patients investigated for microscopic hematuria? J Urol 2018; pii: S0022–5347.
4. Bolenz C, Schröppel B, Eisenhardt A, Schmitz-Dräger BJ, Grimm MO: The investigation of hematuria. Dtsch Arztebl Int 2018; 115: 801–7.

Prof. Dr. med. Christian Bolenz

Klinik für Urologie und Kinderurologie, Universitätsklinikum Ulm
Christian.Bolenz@uniklinik-ulm.de

Prof. Dr. med. Bernd Schröppel

Klinik für Innere Medizin 1, Sektion Nephrologie, Universitätsklinikum Ulm

PD Dr. med. Andreas Eisenhardt

Praxisklinik Urologie Rhein Ruhr, Mülheim an der Ruhr
Klinik für Urologie, Kinderurologie und Uroonkologie, Universitätsklinikum Essen

Prof. Dr. med. Bernd J. Schmitz-Dräger

Sektion Urologische Onkologie, Urologische Klinik St. Theresienkrankenhaus, Nürnberg und Urologische Klinik und Kinderklinik der Universität Erlangen

Prof. Dr. med. Marc-Oliver Grimm

Klinik und Poliklinik für Urologie, Universitätsklinikum Jena

Conflict of interest statement

The authors declare that no conflict of interest exists.

CLINICAL SNAPSHOT

Giant Intraductal Papillary Mucinous Neoplasia (IPMN) of the Pancreas with Malignant Degeneration

Sonography in a 75-year-old man with cingulate upper abdominal pain showed an greatly enlarged pancreas with multiple cystic elements. Magnetic resonance cholangiopancreatography (MRCP) identified a huge (6 × 7 × 16 cm) mucinous mass arising from the main pancreatic duct (arrows). Magnetic resonance imaging (MRI) revealed several contrast-enhancing nodular tumor elements, some of them large, leading to the diagnosis of an extraordinarily large main duct intraductal papillary mucinous neoplasia (IPMN) of the pancreas of with suspected

malignant degeneration. Although cytology of solid tumor samples harvested by endosonographically guided aspiration showed no sign of malignancy, the high risk of degeneration of large main duct IPMNs (circa 70%) prompted us to carry out complete pancreatectomy, which the patient tolerated well. Pathological examination revealed extensive intraductal proliferation of neoplastic duct epithelium with mucus production (IPMN) of oncocytic type with extended associated invasive carcinoma (T3). This experience confirms that large main duct IPMNs should be treated surgically owing to the high likelihood of malignant degeneration, even in cases of benign cytology.

PD Dr. med. Kai Nassenstein, Universitätsklinikum Essen, Institut für Diagnostische und Interventionelle Radiologie und Neuroradiologie, Essen, kai.nassenstein@uk-essen.de

Prof. Dr. med. Peter Markus, Elisabeth-Krankenhaus Essen, Klinik für Allgemein-, Viszeral- und Unfallchirurgie, Essen

PD Dr. med. Brigitte Schumacher, Elisabeth-Krankenhaus Essen, Klinik für Innere Medizin und Gastroenterologie, Essen

Conflict of interest statement: The authors declare that no conflict of interest exists.

Translated from the original German by David Roseveare

Cite this as: Nassenstein K, Markus P, Schumacher B: Giant intraductal papillary mucinous neoplasia (IPMN) of the pancreas with malignant degeneration. Dtsch Arztebl Int 2019; 116: 193. DOI: 10.3238/arztebl.2019.0193