Published in final edited form as:

Neurosci Lett. 2019 May 14; 701: 65–70. doi:10.1016/j.neulet.2019.02.005.

Pubertal hormones increase hippocampal expression of $\alpha 4\beta\delta$ GABA_A receptors

Nicole Keating¹, Nicole Zeak, and Sheryl S. Smith*

Department of Physiology and Pharmacology, SUNY Downstate Medical Center, 450 Clarkson Ave., Brooklyn, NY, 11203, USA

Abstract

CA1 hippocampal expression of $\alpha 4\beta\delta$ GABA_A receptors (GABARs) increases at the onset of puberty in female mice, an effect dependent upon the decline in hippocampal levels of the neurosteroid THP (3 α -OH-5 α -pregnan-20-one) which occurs at this time. The present study further characterized the mechanisms underlying $\alpha 4\beta\delta$ expression, assessed in vivo. Blockade of pubertal levels of 17 β -estradiol (E₂) (formestane, 0.5 mg/kg, i.p. 3 d) reduced $\alpha 4$ and δ expression by 75–80% ($P < 0.05$) in CA1 hippocampus of female mice, assessed using Western blot techniques. Conversely, E₂ administration increased $\alpha 4$ and δ expression by 50–100% in adults, an effect enhanced by more than 2-fold by concomitant administration of the 5 α -reductase blocker finasteride (50 mg/kg, i.p., 3d, $P < 0.05$), suggesting that both declining THP levels and increasing E₂ levels before puberty trigger $\alpha 4\beta\delta$ expression. This effect was blocked by ICI 182,780 (20 mg/kg, s.c., 3 d), a selective blocker of E₂ receptor- α (ER- α). These results suggest that both the rise in circulating levels of E₂ and the decline in hippocampal THP levels at the onset of puberty trigger maximal levels of $\alpha 4\beta\delta$ expression in the CA1 hippocampus.

Keywords

Allopregnanolone; alpha-4; Delta; Estradiol; GABA_A receptor; CA1 hippocampus

1. Introduction

The $\alpha 4\beta\delta$ GABA_A receptor (GABAR) mediates a tonic inhibition in hippocampus which is sensitive to modulation by neurosteroids [65]. Regulation of the expression of this receptor is sensitive to ovarian hormone fluctuations [31,34,37,57]. Previous findings from this laboratory have shown that the onset of puberty in female mice results in increased expression of $\alpha 4\beta\delta$ GABARs on the dendrites and dendritic spines of pyramidal cells in CA1 hippocampus [57,60] for a period of ~10 d [6]. This increase in receptor expression was mimicked by administration of a 5 α -reductase blocker to reduce levels of the neurosteroid THP (3 α -OH-5 α -pregnan-20-one or allopregnanolone) and create a “THP withdrawal state” in pre-pubertal female mice [57]. This suggests that the decline in hippocampal levels of THP at puberty serves to trigger $\alpha 4\beta\delta$ GABAR expression.

*Corresponding author. sheryl.smith@downstate.edu (S.S. Smith).

¹Present address: Regeneron Pharmaceuticals, 777 Old Saw Mill River Road, Tarrytown, NY 10591.

Expression levels of $\alpha 4$ and δ are also increased on CA1 hippocampal pyramidal cells on the proestrous stage of the estrous cycle when circulating levels of E_2 peak [51]. Circulating levels of E_2 rise ~5 d before the onset of puberty in female mice [4]. Thus, this peri-pubertal E_2 rise is a potential mediator of the pubertal increase in $\alpha 4\beta\delta$ GABARs, where it could act independently or synergistically with THP withdrawal. E_2 can act via two different estrogen receptors (ERs), ER α and ER β [30,40]. Both of these receptor subtypes are localized to hippocampal neurons [41] where they mediate genomic actions for a variety of functions. In particular, ER α triggers activation of brain derived neurotrophic factor (BDNF), which has been shown to increase expression of the GABAR $\alpha 4$ subunit [49]. Thus, we also examined the role of ER α in mediating $\alpha 4\beta\delta$ GABAR expression. In addition, we tested whether mimicking the pubertal hormonal milieu could increase $\alpha 4$ and δ expression at times other than puberty, during pre-pubertal and adult life stages.

2. Materials and methods

2.1. Animals

Female C57/BL6 mice (Jackson Labs) were housed in a reverse light:dark cycle (12 :12, lights off at 1100 hs) and euthanized or injected during the light phase of the cycle (2 h before dark, 0900 hs). Mice were evaluated as to pubertal stage by vaginal opening during the light phase. For the following experiments, pre-pubertal (PND 27–32) and pubertal (PND 40–44) mice were used as well as adult female mice (3 months of age) in the estrous stage of the estrous cycle. Estrous cycle stage was determined by the vaginal cytology in adult animals with established regular cycles [51]. All procedures were in accordance with the SUNY Downstate Institutional Animal Care and Use Committee.

2.2. Western blot [57]

Crude membranes from hippocampus were prepared as follows: After removal and hemi-section of the brain, the hippocampus was dissected from each hemisphere on ice. The cortex was peeled back laterally and the underlying hippocampus rolled out with the microspatula. Samples were homogenized using a polytron on ice in 0.0625M Tris, pH 6.8, 2% SDS, 10% glycerol, 5% dithiothreitol (DTT), 1mM EDTA and a cocktail of protease inhibitors (GE healthcare). Crude membranes were prepared after centrifugation at 1000xg for 15 min to remove cellular debris and then collecting the pellet (P2) after centrifugation of the supernatant (SI) at 200,000xg for 30 min.

Samples were first normalized according to protein content using standard techniques and protein concentrations used were used in the linear range (4–10 mg); then, proteins were separated using SDS gel electrophoresis, transferred to nitrocellulose membranes and probed with selective antibodies for the rat $\alpha 4$ (67 kDa, SC-7355, Santa Cruz Biotechnology, 1:20 or 1:100) or δ (54 kDa, Novus Biologicals, 1:1000). Bands were detected with a highly sensitive chemiluminescence substrate (Pierce Supersignal West Femto substrate) for visualization and quantified using Adobe Photoshop. The results were standardized to a Glyceraldehyde 3-phosphate dehydrogenase (GAPDH, 36 kDa) housekeeping protein.

2.3. Drugs

The ovarian hormone 17 β -estradiol (E₂) was injected at a dose of 8 μ g/kg, i.p., to adult female mice for 3 d beginning on the day of estrus. In some cases the 5 α -reductase inhibitor finasteride (Fin, 50 mg/kg, i.p.) was injected for 3 d alone or concomitant with E₂ to pre-pubertal or adult mice to create a THP withdrawal state [63]. In other cases, the aromatase blocker formestane (Form) was administered for 3 d (0.5 mg/kg, i.p., 3d) in order to prevent formation of endogenous E₂. In the final experiments, adult mice were treated for 3 d with a blocker of estrogen receptor- α (ER α) ICI 182,780 (ICI, 20 mg/kg, s.c.) concomitant with E₂ plus finasteride. In all cases the vehicle was oil, and vehicle-injected mice served as controls. All steroids and steroid blockers were obtained from Steraloids (Newport, RI).

2.4. Statistics

Comparisons of the degree of change across groups for all experimental procedures were analyzed with a student's unpaired *t*-test (2 groups) or one-way analysis of variance (ANOVA, 3+ groups). Post-hoc comparisons for the ANOVA were made with a Tukey's test. For all tests, the level of significance was $P < 0.05$. Complete statistics for all experiments are included in the figure legends.

3. Results

3.1. Peri-pubertal increases in circulating E₂ levels contribute to the pubertal increase in hippocampal $\alpha 4$ and δ GABAR subunit expression

Because previous reports suggest that E₂ may increase $\alpha 4$ and δ expression depending on the existing hormonal milieu [70], we tested whether the pubertal rise in circulating E₂ [4] played a role in the increase in $\alpha 4\beta\delta$ expression at this time. To this end, we used the aromatase blocker formestane (0.5 mg/kg, i.p.) which has been shown to reduce E₂ formation and mimic the ovariectomized state [17]. Formestane was administered across a 48 h period (3 injections) because we have previously shown that this time-course is maximal for $\alpha 4\beta\delta$ GABAR expression [31]. Formestane treatment of pubertal mice reduced $\alpha 4$ and δ expression by 75–80% ($P < 0.05$, Fig. 1) suggesting that peri-pubertal increases in E₂ are necessary for maximal expression of $\alpha 4$ and δ subunits in CA1 hippocampus at puberty.

3.2. THP withdrawal increases $\alpha 4$ and δ GABAR subunit expression in the pre-pubertal hippocampus: an E₂-dependent effect

Circulating E₂ levels rise 5 d before the onset of puberty [4] when circulating and hippocampal levels of THP are also elevated [57]. We used finasteride to reduce THP levels to simulate the pubertal THP withdrawal state in order to test whether THP withdrawal in pre-pubertal mice would increase $\alpha 4$ and δ expression and, further, whether pre-pubertal E₂ could play a role in their expression. Finasteride (50 mg/kg, i.p., 3d) was injected using a dosing regimen shown to reduce hippocampal THP levels by ~50% [63]. In some cases, late prepubertal mice (~5 d before vaginal opening) were also injected with the aromatase inhibitor formestane concomitant with finasteride to test the effect of reducing E₂ levels on $\alpha 4$ and δ expression. Finasteride treatment increased $\alpha 4$ and δ expression by ~200%, an

effect prevented by formestane (Fig. 2) suggesting that both E₂ and THP withdrawal are necessary for maximal increases in $\alpha 4\beta\delta$ expression in pre-pubertal hippocampus.

3.3. E₂ administration increases $\alpha 4$ and δ GABAR subunit expression in adult hippocampus

In order to further investigate the effects of pubertal hormones, we tested whether E₂ could increase $\alpha 4$ and δ expression in adult hippocampus. To this end, we administered E₂ (8 μ g/kg, i.p., 3d) to adult, female mice in the stage of estrus. This treatment increased δ and $\alpha 4$ expression by ~50 and 100% compared to vehicle-treated values ($P < 0.05$, Fig. 3), suggesting that E₂ alone can alter expression of $\alpha 4\beta\delta$ GABARs.

3.4. E₂ plus THP withdrawal produce peak levels of $\alpha 4$ and δ GABAR subunit expression in adult hippocampus

In order to determine the effects of THP withdrawal on $\alpha 4\beta\delta$ GABAR expression in the adult, we administered finasteride (50 mg/kg, i.p., 3d) to adult female mice in estrus. THP withdrawal produced by finasteride increased $\alpha 4$ and δ expression by 100% and 30%, respectively ($P < 0.05$, Fig. 3). However, administration of E₂ concomitant with finasteride increased $\alpha 4$ and δ expression significantly more, by 250% and 125% ($P < 0.05$, Fig. 3), suggesting that maximal levels of $\alpha 4$ and δ expression require both E₂ and THP withdrawal.

3.5. E₂ effects on $\alpha 4$ and δ GABAR subunit expression are mediated by estrogen receptor- α

E₂ can act through two different nuclear receptors (ER α and ER β) [30,40] which have genomic effects, as well as via membrane effects. In order to determine whether E₂ effects on $\alpha 4\beta\delta$ GABAR expression were due to actions mediated by ER α , which mediates most effects of the hormone, we administered the selective ER α blocker [32] ICI182,780 (ICI) concomitant with treatment with E₂ and finasteride. ICI was injected using a dosing regimen (20 mg/kg, s.c., 3 d) that has been shown to block effects of E₂ on uterine weight in a manner similar to ER α knock-out [22]. ICI treatment prevented the increase in $\alpha 4$ and δ expression produced by E₂ plus finasteride (Fig. 4), suggesting that E₂ acts through ER α to increase $\alpha 4$ and δ GABAR subunit expression.

4. Discussion

4.1. Summary

Our findings suggest that the pubertal increase in $\alpha 4\beta\delta$ GABARs is due to both declining levels of THP as well as peri-pubertal elevations in circulating E₂. In addition, we show that use of these ovarian hormones to mimic the changes which occur at puberty can also increase $\alpha 4$ and δ expression to maximal levels at times other than puberty. The effect of E₂ was mediated by ER α and potentiated by the effects of THP withdrawal.

4.2. Pubertal hormones

Our results suggest that optimal levels of $\alpha 4$ expression are achieved with sustained exposure to E₂ in combination with THP withdrawal, mimicking the ovarian milieu of the

peri-pubertal period. The onset of puberty in the female mouse is preceded by increases in circulating E₂, with two-fold increases observed 5 d before vaginal opening which persist across the beginning of the pubertal period [4].

In addition, both brain and circulating levels of THP are increased during the late pre-pubertal period and decrease by ~70% at the onset of puberty in the female mouse [57]. Early studies established that this change in steroid levels is due to altered enzyme activity in the ovary [16,38]. THP is formed from P via two enzymatic conversions mediated by 5 α -reductase and 3 α -hydroxysteroid oxidoreductase (3 α -HSD) [13,28,43]. In addition to an ovarian site of production, this steroid can be produced in the adrenal gland, with release into the general circulation before puberty onset [11]. Recent studies also suggest that THP can be produced de novo directly in the brain from cholesterol via side chain cleavage enzyme; thus, it is also classified as a neurosteroid [13]. This can occur in a variety of CNS sites, in both glia and neurons, including the CA1 hippocampal pyramidal cell [3]. In the mouse, THP levels peak during the dark phase of the circadian cycle [14]. This peri-pubertal pattern of ovarian hormonal fluctuations is also observed in the human: E₂ levels rise prior to puberty onset and circulating levels of THP fall at puberty onset, when they fluctuate in a circadian manner [7,25,39].

4.3. E₂ and $\alpha 4\beta\delta$ GABAR expression

The present results suggest that E₂ alone increases $\alpha 4$ and δ expression in addition to enhancing the effect of THP withdrawal. In our previous studies, increased $\alpha 4\beta\delta$ GABAR expression initiated at puberty was maintained for 10 d before undergoing cyclic expression patterns in female CA1 hippocampus [6,57]. This prolonged pubertal period has been noted before, where high circulating E₂ levels are maintained for many days after puberty onset despite estrous cycle variations in the vaginal cytology [23]. Thus, this pubertal plateau in E₂ levels may underlie the prolonged period of $\alpha 4\beta\delta$ expression at puberty.

Consistent with the present results, a number of studies have confirmed the role of E₂ in increasing expression of $\alpha 4$ and/or δ expression, which has been noted in cell lines [47,70] as well as in the hippocampus, where both ER α and β are localized [32], after increases in exogenous or endogenous E₂ levels [40]. The role of E₂ in increasing $\alpha 4\beta\delta$ GABAR expression at puberty and in the in vitro model is likely via its increase in brain-derived neurotrophic factor (BDNF), which activates the $\alpha 4$ promoter via expression of early growth response factor-3 (EGR-3) [50]. BDNF also increases membrane trafficking of δ -GABARs via TrkB receptors (tropomyosin receptor kinase B) [27].

E₂ increases both BDNF protein and mRNA in CA1 and CA3 regions of the hippocampus in ovariectomized rats [55,69]. BDNF immunoreactivity is also increased on the proestrous stage of the estrous cycle, during the peak in circulating levels of E₂ in the dendrites of CA1 and CA3 pyramidal cells [56]. More recent studies have established that E₂ increases phosphorylated TrkB receptors, via ER α , as well as ER β , in hippocampal neurons [64]. ER α generates genomic events via classic nuclear steroid receptors [45], but can also mediate membrane-initiated signaling to activate kinase cascades in a more rapid time-course [5]. These genomic effects of E₂ likely underlie the E₂ (ER α)-mediated increases in $\alpha 4$ expression observed in the present study.

4.4. THP withdrawal and $\alpha 4\beta 6$ GABAR expression

THP withdrawal was shown in the present study to result in peak levels of $\alpha 4$ and δ expression in the presence of high circulating levels of E_2 . Our previous study showed that at puberty onset, the declining levels of THP trigger expression of $\alpha 4\beta 6$ GABARs in CA1 hippocampus [57]. Administration of replacement THP prevents this increase in receptor expression, confirming the pivotal role of THP withdrawal in regulating $\alpha 4\beta 6$ GABAR expression. Other studies have reported that declining levels of THP or its parent compound progesterone, either administered exogenously or as a result of the ovarian cycle and pregnancy, also increase expression of $\alpha 4$ or $\alpha 4\beta 6$ GABARs in a variety of CNS sites [20,34,36,62].

In the present study, $\alpha 4$ and δ expression were assessed in the whole hippocampus which would include both the CA3 hippocampus and the dentate gyrus in addition to the CA1 hippocampus. Unlike the CA1 and CA3 hippocampus, the dentate gyrus expresses high levels of $\alpha 4\beta 6$ GABARs at times other than puberty [8,9]. However, $\alpha 4\beta 6$ expression is increased at puberty in the CA3 [46] and, preliminary data suggest, also in the dentate gyrus. Therefore, the pubertal hormone conditions used in the present study to increase $\alpha 4$ and δ expression may be relevant for all sub-regions of the hippocampus.

4.5. $\alpha 4\beta 6$ GABAR expression as a compensatory response

The underlying mechanism for increases in $\alpha 4\beta 6$ GABARs after THP withdrawal may be a result of a compensatory change in inhibitory tone as a result of the loss of this GABA modulatory neurosteroid. $\alpha 4\beta 6$ GABARs generate a tonic inhibition due to their extrasynaptic location [66], high sensitivity to GABA [11,42] and relative lack of desensitization under steady state conditions [11]. Their relative plasticity suggests that their expression may serve a compensatory role; their expression is increased in certain seizure states [19], during chronic administration of the stimulant methamphetamine [59], in response to traumatic brain injury [44] and excitotoxic challenge [54]. Their increase after THP withdrawal, as demonstrated here, may also represent a compensatory increase in response to declining levels of this positive GABAR modulator which would decrease inhibition. One possibility is that removal of THP, which inhibits L-type voltage-gated calcium channels (VGCCs) [15], then increases Ca^{++} influx, which can increase expression and trafficking of $\alpha 4\beta 6$ GABARs [52,53]. Conversely, $\alpha 4\beta 6$ expression is decreased during pregnancy when circulating levels of THP are elevated [36]. This increase in $\alpha 4\beta 6$ expression across hormonal transitions (i.e., the estrous cycle, pregnancy and puberty onset) reduces seizure susceptibility [35,67].

4.6. THP administration and $\alpha 4\beta 6$ GABAR expression

In contrast to THP withdrawal, 48 h administration of THP also increases expression of $\alpha 4\beta 6$ GABARs [1,31,58] by increasing surface expression of the receptor, with no effect on internalization [31]. This effect was mediated by PKC, including PKC- δ , targeting serine 443 in $\alpha 4$ [1,31]. The reason that both sustained exposure to and withdrawal from THP increase $\alpha 4\beta 6$ expression is not clear, but may be related to the diverse effects of the steroid on triggering phosphorylation [1] and inhibiting VGCCs [15] which would increase receptor expression after administration or withdrawal, respectively.

4.7. $\alpha 4\beta 8$ GABAR expression and puberty: effects on anxiety and panic response

Anxiety disorders, which are more prevalent in females, are most likely to emerge during or after puberty [21]. Our previous findings suggest that $\alpha 4\beta 8$ GABAR expression at puberty onset are associated with alterations in mood. At the onset of puberty, when $\alpha 4\beta 8$ GABARs are increased, the stress steroid THP [48] produces anxiety [57], due to a paradoxical decrease in $\alpha 4\beta 8$ -generated current, in contrast to its typical anti-anxiety effect [10]. This effect is dependent upon the polarity of the GABA-generated Cl⁻ current [18,57] and is not observed in $\alpha 4$ knock-out mice. In addition, estrous cycle-dependent changes in $\alpha 4\beta 8$ expression on midbrain interneurons have been shown to generate panic responses, suggesting a role for this receptor in panic disorder [33] as well as stress-induced anxiety.

4.8. $\alpha 4\beta 8$ GABAR expression and puberty: effects on cognition and synaptic pruning

More recent studies from this laboratory [60,61] have shown that the expression of $\alpha 4\beta 8$ on the dendritic shaft and spines of CA1 pyramidal cells at puberty impairs cognition acutely and prevents induction of long-term potentiation, an in vitro model of learning. This outcome may define the end of an optimal period for learning at the onset of puberty which has been described in the human for certain tasks such as learning a second language [26]. The impact of puberty onset on learning is complex, however, as it also is a time when stress can enhance cognition [23], which may be related to corticosterone effects or to the effects of THP [60], which is also released after stress [48].

One of the long-term consequences of $\alpha 4\beta 8$ expression on the CA1 spines during adolescence is to trigger synaptic pruning [2], a phenomenon where dendritic spines are removed from many areas of the CNS [24,29,68], and is thought to be essential for optimal cognition in adulthood [12]. In the absence of synaptic pruning, as observed in the $\alpha 4^{-/-}$ mouse, spine density is increased and cognitive flexibility is impaired [2].

4.9. Conclusions

The results from this study demonstrate that pubertal increases in expression of $\alpha 4\beta 8$ GABARs are due to both peri-pubertal increases in circulating levels of E₂ and declining levels of THP ("THP withdrawal"). These effects were mediated by ER α . Because $\alpha 4\beta 8$ GABARs are associated with changes in anxiety and synaptic function, the results from the present study may be relevant for understanding the processes which contribute to neuropathologies associated with mood and cognition.

Acknowledgements

The authors are grateful to Y. Ruderman for helpful technical assistance.

Funding

This work was supported by the National Institutes of Health [grant numbers R01-MH100561 and R01-MH115900] to SSS.

Abbreviations:

3 α -HSD 3 α -hydroxysteroid oxidoreductase

BDNF	brain-derived neurotrophic factor
E₂	17 β -estradiol
ER-α	E ₂ receptor- α
Fin	finasteride
Form	formestane
GABAR	GABA _A receptor
GAPDH	glyceraldehyde 3-phosphate dehydrogenase
ICI	ICI 182,780 (i.e., fulvestrant)
PKC	protein kinase C
PND	post-natal day
THP	3 α -OH-5 α -pregnan-20-one (allopregnanolone)
TrkB	tropomyosin receptor kinase B
VGCC	voltage-gated calcium channel

References

- [1]. Abramian AM, Comenencia-Ortiz E, Vithlani M, Tretter EV, Sieghart W, Davies PA, Moss SJ, Protein kinase C phosphorylation regulates membrane insertion of GABAA receptor subtypes that mediate tonic inhibition, *J. Biol. Chem.* 285 (2010) 41795–41805. [PubMed: 20940303]
- [2]. Afroz S, Parato J, Shen H, Smith SS, Synaptic pruning in the female hippocampus is triggered at puberty by extrasynaptic GABAA receptors on dendritic spines, *Elife* 5 (2016).
- [3]. Agis-Balboa RC, Pinna G, Zhubi A, Maloku E, Veldic M, Costa E, Guidotti A, Characterization of brain neurons that express enzymes mediating neurosteroid biosynthesis, *Proc. Natl. Acad. Sci.* 103 (2006) 14602–14607. [PubMed: 16984997]
- [4]. Ahima RS, Dushay J, Flier SN, Prabakaran D, Flier JS, Leptin accelerates the onset of puberty in normal female mice, *J. Clin. Invest.* 99 (1997) 391–395. [PubMed: 9022071]
- [5]. Akama KT, McEwen BS, Estrogen stimulates postsynaptic density-95 rapid protein synthesis via the Akt/protein kinase B pathway, *J. Neurosci.* 23 (2003) 2333–2339. [PubMed: 12657692]
- [6]. Aoki C, Sabaliauskas N, Chowdhury T, Min JY, Colacino AR, Laurino K, Barbarich-Marsteller NC, Adolescent female rats exhibiting activity-based anorexia express elevated levels of GABA(A) receptor alpha4 and delta subunits at the plasma membrane of hippocampal CA1 spines, *Synapse* 66 (2012) 391–407. [PubMed: 22213233]
- [7]. Apter D, Hermanson E, Update on female pubertal development, *Curr. Opin. Obstet. Gynecol.* 14 (2002) 475–481.
- [8]. Benke D, Mertens S, Trzeciak A, Gillessen D, Mohler H, Identification and immunohistochemical mapping of GABAA receptor subtypes containing the delta-subunit in rat brain, *FEBS Lett.* 283 (1991) 145–149. [PubMed: 1645294]
- [9]. Benke D, Michel C, Mohler H, GABAA receptors containing the $\alpha 4$ -subunit: prevalence, distribution, pharmacology, and subunit architecture in situ, *J. Neurochem.* 69 (1997) 806–814. [PubMed: 9231742]
- [10]. Bitran D, Dugan M, Renda P, Ellis R, Foley M, Anxiolytic effects of the neuroactive steroid pregnanolone (3 α -OH-5 β -pregnan-20-one) after microinjection in the dorsal hippocampus and lateral septum, *Brain Res.* 850 (1999) 217–224. [PubMed: 10629767]

- [11]. Brown N, Kerby J, Bonnert TP, Whiting PJ, Wafford KA, Pharmacological characterization of a novel cell line expressing human $\alpha(4)\beta(3)\delta$ GABA(A) receptors, *Br. J. Pharmacol.* 136 (2002) 965–974. [PubMed: 12145096]
- [12]. Chechik G, Meilijson I, Ruppin E, Synaptic pruning in development: a computational account, *Neural Comput.* 10 (1998) 1759–1777. [PubMed: 9744896]
- [13]. Compagnone NA, Mellon SH, Neurosteroids: biosynthesis and function of these novel neuromodulators, *Front. Neuroendocrinol.* 21 (2000) 1–56. [PubMed: 10662535]
- [14]. Corpechot C, Collins B, Carey M, Tsouros A, Robel P, Fry J, Brain neurosteroids during the mouse oestrous cycle, *Brain Res.* 766 (1997) 276–280. [PubMed: 9359616]
- [15]. Earl DE, Tietz EI, Inhibition of recombinant L-type voltage-gated calcium channels by positive allosteric modulators of GABAA receptors, *J. Pharmacol. Exp. Ther.* 337 (2011) 301–311. [PubMed: 21262851]
- [16]. Fadalti M, Petraglia F, Luisi S, Bernardi F, Casarosa E, Ferrari E, Luisi M, Saggese G, Genazzani AR, Bernasconi S, Changes of serum allopregnanolone levels in the first 2 years of life and during pubertal development, *Pediatr. Res.* 46 (1999) 323–327. [PubMed: 10473049]
- [17]. Flood P, Daniel D, Pronociceptive actions of isoflurane: a protective role for estrogen, *Anesthesiology* 99 (2003) 476–479. [PubMed: 12883422]
- [18]. Gong QH, Smith SS, Characterization of neurosteroid effects on hyperpolarizing current at $\alpha(4)\beta(2)\delta$ GABA receptors, *Psychopharmacology (Berl.)* (2014).
- [19]. Grabenstatter HL, Cogswell M, Cruz Del Angel Y, Carlsen J, Gonzalez MI, Raol YH, Russek SJ, Brooks-Kayal AR, Effect of spontaneous seizures on GABAA receptor $\alpha(4)$ subunit expression in an animal model of temporal lobe epilepsy, *Epilepsia* 55 (2014) 1826–1833. [PubMed: 25223733]
- [20]. Griffiths J, Lovick T, Withdrawal from progesterone increases expression of $\alpha(4)$, $\beta(1)$ and δ GABA(A) receptor subunits in neurons in the periaqueductal gray matter in female Wistar rats, *J. Comp. Neurol.* 486 (2005) 89–97. [PubMed: 15834956]
- [21]. Hayward C, Sanborn K, Puberty and the emergence of gender differences in psychopathology, *J. Adol. Health* 30 (2002) 49–58.
- [22]. He B, Rhodes-Brower S, Miller MR, Munson AE, Germolec DR, Walker VR, Korach KS, Meade BJ, Octamethylcyclotetrasiloxane exhibits estrogenic activity in mice via ER α , *Toxicol. Appl. Pharmacol.* 192 (2003) 254–261. [PubMed: 14575643]
- [23]. Hodes GE, Shors TJ, Distinctive stress effects on learning during puberty, *Horm. Behav.* 48 (2005) 163–171. [PubMed: 15885691]
- [24]. Huttenlocher PR, Synaptic density in human frontal cortex - developmental changes and effects of aging, *Brain Res.* 163 (1979) 195–205. [PubMed: 427544]
- [25]. Janfaza M, Sherman TI, Larmore KA, Brown-Dawson J, Klein KO, Estradiol levels and secretory dynamics in normal girls and boys as determined by an ultrasensitive bioassay: a 10 year experience, *J. Pediatr. Endocrinol. Metab.* 19 (2006) 901–909. [PubMed: 16995570]
- [26]. Johnson JS, Newport EL, Critical period effects in second language learning: the influence of maturational state on the acquisition of English as a second language, *Cogn. Psychol.* 21 (1989) 60–99. [PubMed: 2920538]
- [27]. Joshi S, Kapur J, Slow intracellular accumulation of GABA(A) receptor δ subunit is modulated by brain-derived neurotrophic factor, *Neuroscience* 164 (2009) 507–519. [PubMed: 19665523]
- [28]. Karavolas HJ, Hodges DR, Neuroendocrine metabolism of progesterone and related progestins, *Ciba Found. Symp.* 153 (1990) 22–44. [PubMed: 2292213]
- [29]. Koss WA, Belden CE, Hristov AD, Juraska JM, Dendritic remodeling in the adolescent medial prefrontal cortex and the basolateral amygdala of male and female rats, *Synapse* 68 (2014) 61–72. [PubMed: 24105875]
- [30]. Kuiper GG, Carlsson B, Grandien K, Enmark E, Haggblad J, Nilsson S, Gustafsson JA, Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptors α and β , *Endocrinology* 138 (1997) 863–870. [PubMed: 9048584]
- [31]. Kuver A, Shen H, Smith SS, Regulation of the surface expression of $\alpha(4)\beta(2)\delta$ GABA(A) receptors by high efficacy states, *Brain Res.* 1463 (2012) 1–20. [PubMed: 22609410]

- [32]. Long X, Nephew KP, Fulvestrant (ICI 182,780)-dependent interacting proteins mediate immobilization and degradation of estrogen receptor- α , *J. Biol. Chem.* 281 (2006) 9607–9615. [PubMed: 16459337]
- [33]. Lovick T, Panic disorder: a malfunction of multiple transmitter control systems within the midbrain periaqueductal gray matter? *Neuroscientist* 6 (2000) 48–59.
- [34]. Lovick TA, Griffiths JL, Dunn SM, Martin IL, Changes in GABA(A) receptor subunit expression in the midbrain during the oestrous cycle in Wistar rats, *Neuroscience* 131 (2005) 397–405. [PubMed: 15708482]
- [35]. Maguire J, Ferando I, Simonsen C, Mody I, Excitability changes related to GABAA receptor plasticity during pregnancy, *J. Neurosci.* 29 (2009) 9592–9601. [PubMed: 19641122]
- [36]. Maguire JL, Mody I, GABA(A)R plasticity during pregnancy: relevance to postpartum depression, *Neuron* 59 (2009) 207–213.
- [37]. Maguire JL, Stell BM, Rafizadeh M, Mody I, Ovarian cycle-linked changes in GABA(A) receptors mediating tonic inhibition alter seizure susceptibility and anxiety, *Nat. Neurosci.* 8 (2005) 797–804. [PubMed: 15895085]
- [38]. Mannan MA, O'Shaughnessy PJ, Ovarian steroid metabolism during post-natal development in the normal mouse and in the adult hypogonadal (hpg) mouse, *J. Reprod. Fertil.* 82 (1988) 727–734. [PubMed: 3361507]
- [39]. McCartney CR, et al., Obesity and sex steroid changes across puberty: evidence for marked hyperandrogenemia in pre- and early pubertal obese girls, *J. Clin. Endocrinol. Metab.* 92 (2007) 430–436. [PubMed: 17118995]
- [40]. McEwen BS, Alves SE, Bulloch K, Weiland NG, Ovarian steroids and the brain: implications for cognition and aging, *Neurology* 48 (1997) S8–15.
- [41]. McEwen BS, Milner TA, Hippocampal formation: shedding light on the influence of sex and stress on the brain, *Brain Res. Rev.* 55 (2007) 343–355. [PubMed: 17395265]
- [42]. Meera P, Wallner M, Otis T, Molecular basis for the high THIP/gaboxadol sensitivity of extrasynaptic GABA-A receptors, *J. Neurophysiol.* 106 (2011) 2057–2011. [PubMed: 21795619]
- [43]. Mellon SH, Vaudry H, Biosynthesis of neurosteroids and regulation of their synthesis, *Int. Rev. Neurobiol.* 46 (2001) 33–78. [PubMed: 11599305]
- [44]. Mchedlishvili Z, Lepsveridze E, Xu H, Kharlamov EA, Lu B, Kelly KM, Increase of GABAA receptor-mediated tonic inhibition in dentate granule cells after traumatic brain injury, *Neurobiol. Dis.* 38 (2010) 464–475. [PubMed: 20304069]
- [45]. Nilsson S, Makela S, Treuter E, Tujague M, Thomsen J, Andersson G, Enmark E, Pettersson K, Warner M, Gustafsson JA, Mechanisms of estrogen action, *Physiol. Rev.* 81 (2001) 1535–1565. [PubMed: 11581496]
- [46]. Parato J, Shen H, Smith SS, α 4 β delta GABAA receptors trigger synaptic pruning and reduce dendritic length of female mouse CA3 hippocampal pyramidal cells at puberty, *Neuroscience* 398 (2019) 23–36. [PubMed: 30496825]
- [47]. Pierson RC, Lyons AM, Greenfield LJJ, Gonadal steroids regulate GABA-A receptor subunit mRNA expression in NT2-N cells, *Brain Res. Mol. Brain Res.* 138 (2005) 105–115. [PubMed: 16000229]
- [48]. Purdy RH, Morrow AL, Moore PH Jr, Paul SM, Stress-induced elevations of gamma-aminobutyric acid type A receptor-active steroids in the rat, *PNAS* 88 (1991) 4553–4557. [PubMed: 1852011]
- [49]. Roberts DS, Hu Y, Lund IV, Brooks-Kayal AR, Russek SJ, Brain-derived neurotrophic factor (BDNF)-induced synthesis of early growth response factor 3 (Egr3) controls the levels of type A GABA receptor α 4 subunits in hippocampal neurons, *J. Biol. Chem.* 281 (2006) 29431–29435. [PubMed: 16901909]
- [50]. Roberts DS, Raol YH, Bandyopadhyay S, Lund IV, Budreck EC, Passini MA, Wolfe JH, Brooks-Kayal AR, Russek SJ, Egr3 stimulation of GABARA4 promoter activity as a mechanism for seizure-induced up-regulation of GABA(A) receptor α 4 subunit expression, *Proc. Natl. Acad. Sci.* 102 (2005) 11894–11899. [PubMed: 16091474]
- [51]. Sabaliauskas N, Shen H, Molla J, Gong QH, Kuver A, Aoki C, Smith SS, Neurosteroid effects at α 4 β delta GABAA receptors alter spatial learning and synaptic plasticity in CA1

hippocampus across the estrous cycle of the mouse, *Brain Res.* 1621 (2015) 170–186. [PubMed: 25542386]

- [52]. Saliba RS, Gu Z, Yan Z, Moss SJ, Blocking L-type voltage-gated Ca²⁺ channels with dihydropyridines reduces gamma-aminobutyric acid type A receptor expression and synaptic inhibition, *J. Biol. Chem.* 284 (2009) 32544–32550. [PubMed: 19778903]
- [53]. Saliba RS, Kretschmannova K, Moss SJ, Activity-dependent phosphorylation of GABA_A receptors regulates receptor insertion and tonic current, *EMBO J.* 31 (2012) 2937–2951. [PubMed: 22531784]
- [54]. Santhakumar V, Jones RT, Mody I, Developmental regulation and neuroprotective effects of striatal tonic GABA-A currents, *Neuroscience* 167 (2010) 644–655. [PubMed: 20206233]
- [55]. Scharfman HE, Hintz TM, Gomez J, Stormes KA, Barouk S, Malthankar-Phatak GH, McCloskey DP, Luine VN, MacLusky NJ, Changes in hippocampal function of ovariectomized rats after sequential low doses of estradiol to simulate the preovulatory estrogen surge, *Eur. J. Neurosci.* 26 (2007) 2595–2612. [PubMed: 17970745]
- [56]. Scharfman HE, Mercurio TC, Goodman JH, Wilson MA, MacLusky NJ, Hippocampal excitability increases during the estrous cycle in the rat: a potential role for brain-derived neurotrophic factor, *J. Neurosci.* 23 (2003) 11641–11652. [PubMed: 14684866]
- [57]. Shen H, Gong QH, Aoki C, Yuan M, Ruderman Y, Dattilo M, Williams K, Smith SS, Reversal of neurosteroid effects at alpha4-beta2-delta GABA-A receptors triggers anxiety at puberty, *Nat. Neurosci.* 10 (2007) 469–477. [PubMed: 17351635]
- [58]. Shen H, Gong QH, Yuan M, Smith SS, Short-term steroid treatment increases delta GABA-A receptor subunit expression in rat CA1 hippocampus: pharmacological and behavioral effects, *Neuropharmacology* 49 (2005) 573–586. [PubMed: 15950994]
- [59]. Shen H, Mohammad A, Ramroop J, Smith SS, A stress steroid triggers anxiety via increased expression of alpha4betadelta GABA receptors in methamphetamine dependence, *Neuroscience* 254 (2013) 452–475. [PubMed: 23994152]
- [60]. Shen H, Sabaliauskas N, Sherpa A, Fenton AA, Stelzer A, A. C, Smith SS, A critical role for $\alpha 4\beta 8$ GABA-A receptors in shaping learning deficits at puberty in mice, *Science* 327 (2010) 1515–1518. [PubMed: 20299596]
- [61]. Shen H, Sabaliauskas N, Yang L, Aoki C, Smith SS, Role of alpha4-containing GABA receptors in limiting synaptic plasticity and spatial learning of female mice during the pubertal period, *Brain Res.* (2016).
- [62]. Smith SS, Gong QH, Hsu FC, Markowitz RS, French-Mullen JMH, Li X, GABA_A receptor $\alpha 4$ subunit suppression prevents withdrawal properties of an endogenous steroid, *Nature* 392 (1998) 926–929. [PubMed: 9582073]
- [63]. Smith SS, Ruderman Y, Frye CA, Homanics GE, Y. M, Steroid withdrawal in the mouse results in anxiogenic effects of 3alpha,5alpha-THP: A possible model of premenstrual dysphoric disorder, *Psychopharmacology* 186 (2006) 323–333. [PubMed: 16193334]
- [64]. Spencer-Segal JL, Tsuda MC, Mattel L, Waters EM, Romeo RD, Milner TA, McEwen BS, Ogawa S, Estradiol acts via estrogen receptors alpha and beta on pathways important for synaptic plasticity in the mouse hippocampal formation, *Neuroscience* 202 (2012) 131–146. [PubMed: 22133892]
- [65]. Stell BM, Brickley SG, Tang CY, Farrant M, Mody I, Neuroactive steroids reduce neuronal excitability by selectively enhancing tonic inhibition mediated by delta subunit-containing GABA-A receptors, *Proc. Natl. Acad. Sci.* 100 (2003) 14439–14444. [PubMed: 14623958]
- [66]. Wei W, Zhang N, Peng Z, Houser CR, Mody I, Perisynaptic localization of delta subunit-containing GABA(A) receptors and their activation by GABA spillover in the mouse dentate gyrus, *J. Neurosci.* 23 (2003) 10650–10661. [PubMed: 14627650]
- [67]. Yang L, Shen H, Merlin LR, Smith SS, Pubertal expression of alpha4betadelta GABA_A receptors reduces seizure-like discharges in CA1 Hippocampus, *Sci. Rep.* 6 (2016) 31928. [PubMed: 27561815]
- [68]. Yildirim M, Mapp OM, Janssen WG, Yin W, Morrison JH, Gore AC, Postpubertal decrease in hippocampal dendritic spines of female rats, *Exp. Neurol.* 210 (2008) 339–348. [PubMed: 18096161]

- [69]. Zhou J, Zhang H, Cohen RS, Pandey SC, Effects of estrogen treatment on expression of brain-derived neurotrophic factor and cAMP response element-binding protein expression and phosphorylation in rat amygdaloid and hippocampal structures, *Neuroendocrinology* 81 (2005) 294–310. [PubMed: 16179807]
- [70]. Zhou X, Smith SS, Steroid requirements for regulation of the alpha-4 subunit of the GABA-A receptor in an in vitro model, *Neurosci.Lett.* 411 (2007) 61–66. [PubMed: 17081691]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Fig. 1. Increases in pubertal expression of GABA_A receptor $\alpha 4$ and δ subunits in hippocampus is reduced by blockade of estrogen synthesis.

In order to determine if elevated levels of the ovarian steroid 17 β -estradiol (E₂) during the pubertal period contribute to the increase in hippocampal $\alpha 4$ and δ expression, the aromatase blocker formestane (Form) was administered in vivo (0.5 mg/kg, i.p., 3d). A, Representative Western blot. B, Summary figure: Form decreased hippocampal expression of $\alpha 4$ and δ subunits at puberty in female mice. $\alpha 4$, $t(10)=3.35$, $*P=0.007$; δ , $t(10)=4.28$, $*P=0.0016$. N=6, performed in triplicate.

Fig. 2. Inducing “THP withdrawal” in pre-pubertal mice increases expression of hippocampal α4 and δ subunits: An effect prevented by blockade of estrogen synthesis.

“THP withdrawal” was induced in pre-pubertal mice by blocking THP formation with finasteride (Fin) administered in some cases with formestane (Form) to block E₂ synthesis. A, Representative Western blot. B, Summary figure. α4, F(2,15)=20.5, *P < 0.0001; δ, F(2,15)=15.7, P=0.0002. *P < 0.05 vs. vehicle-treated control. N=6, performed in triplicate.

Fig. 3. Mimicking the pubertal hormonal milieu produces maximal expression of $\alpha 4$ and δ in the adult, female mouse.

E₂ (8 μ g/kg, i.p.) and/or finasteride (Fin) were administered to adult mice for 3 d beginning on estrus. A, Representative Western blots show that both drugs significantly increased $\alpha 4$ and δ expression, but together produced optimal expression of these subunits. B, Summary figure. $\alpha 4$, $F(3,20)=4.62$, $*P=0.013$; δ , $F(3,20)=3.23$, $P=0.0442$. $*P < 0.05$ vs. vehicle-treated control, $**P < 0.01$ vs. other groups. $N=8$ mice/group, performed in triplicate.

Fig. 4. E₂ increases expression of GABAR $\alpha 4$ and δ subunits via ER- α .

E₂ (8 μ g/kg, i.p.) and finasteride (Fin) or vehicle (Con) were administered to adult mice for 3 d in some cases along with a blocker of estrogen receptor- α (ER α) ICI 182,780 (ICI, 20 mg/kg, s.c.). A, Representative Western blots show that blockade of ER α prevented the E₂+Fin-induced increases in $\alpha 4$ and δ expression. B, Summary figure. $\alpha 4$, $F(2,21)=14.0$, $*P=0.0001$; δ , $F(2,21)=23.2$, $P < 0.0001$. $*P < 0.05$ vs. vehicle-treated control, $**P < 0.01$ vs. other groups. N=8 mice/group, performed in triplicate.