Published in final edited form as:

Addict Behav. 2018 December ; 87: 260–266. doi:10.1016/j.addbeh.2018.07.020.

Development and validation of a contextual behavioral distress intolerance task in cigarette smokers

Samantha G. Farris^{a,b,c,*}, Angelo M. DiBello^d, and Michael J. Zvolensky^{e,f}

^aAlpert Medical School of Brown University, Providence, RI, United States

^bThe Miriam Hospital, Providence, RI, United States

^cRutgers, the State University of New Jersey, Piscataway, NJ, United States

^dBrown University School of Public Health, Providence, RI, United States

^eUniversity of Houston, Houston, TX, United States

^fUniversity of Texas MD Anderson Cancer Center, Houston, TX, United States

Abstract

Introduction: Distress intolerance, an individual's perceived or actual inability to withstand negative emotional or physical distress, contributes to the maintenance of smoking. However, there is limited understanding of the contextual factors that impact distress intolerance in general or among smokers specifically. This study aimed to adapt and test a computerized behavioral persistence task that requires re-typing a passage while adhering to specific instructions (Contextual-Frustration Intolerance Typing Task [C-FiTT]). C-FiTT was designed to model contextual factors that influence distress intolerance, negative affect, and smoking urges.

Method: Daily smokers ($n = 550$) were recruited through the use of Qualtrics Panels. Using a $2 \times 2 + 1$ experimental design, participants were randomly assigned to one of four C-FiTT conditions that crossed task difficulty (low or high difficulty) with passage content (neutral or tobacco withdrawal text), or a neutral control group.

Results: C-FiTT produced an average persistence time of 94.1 ± 114.3 s and 64.7% of participants self-terminated the task. C-FiTT also produced small to medium sized-increases in negative affect and smoking urges. Between-condition comparisons indicated that the high-difficulty C-FiTT produced shorter behavioral persistence, greater self-termination likelihood, and larger increases in negative affect and smoking urges. The combination of high-difficulty and withdrawal content resulted in the shortest persistence time, 100% self-termination rate, and largest increases in negative affect and smoking urges, compared to other conditions

*Corresponding author at: Alpert Medical School of Brown University, Department of Psychiatry and Human Behavior, 345 Blackstone Blvd, Butler Hospital, Providence, RI 02906, United States, samantha_farris@brown.edu (S.G. Farris).

Declarations of interest

None

Appendix A. Supplementary data

Supplementary data to this article can be found online at <mailto:https://doi.org/10.1016/j.addbeh.2018.07.020>.

Conclusions: Findings provide initial evidence for the validity of C-FiTT in smokers within the context of tobacco withdrawal at low and high levels of task difficulty. Avenues for refinement and use of C-FiTT are discussed.

Keywords

Distress intolerance; Negative affect; Persistence; Tobacco withdrawal; Measurement

1 Introduction

Distress intolerance (DI) reflects one's perceived or behavioral in-capacity to withstand negative emotional or physical distress states (Leyro, Zvolensky, & Bernstein, 2010; Zvolensky, Leyro, Bernstein, & Vujanovic, 2010). DI is linked to the persistence of smoking (Leventhal & Zvolensky, 2015) including more severe tobacco dependence (Leyro, Bernstein, Vujanovic, McLeish, & Zvolensky, 2011), smoking to alleviate negative affect (Trujillo et al., 2017), and greater nicotine consumption following nicotine deprivation (Bold, Yoon, Chapman, & McCarthy, 2013; Perkins, Karelitz, Giedgowd, Conklin, & Sayette, 2010). Certain aspects of DI, like intolerance for physical distress, are associated with poor smoking cessation outcomes (e.g., Kahler, McHugh, Metrik, Spillane, & Rohsenow, 2013), perhaps due to increased sensitivity to nicotine deprivation and heightened withdrawal severity (Farris, Zvolensky, Otto, & Leyro, 2015).

DI is a higher order construct including intolerance for negative emotional states (e.g., frustration, anxiety) or physical sensations (e.g., pain, bodily discomfort; Zvolensky et al., 2010). One's *perceived* intolerance for distress can be evaluated with self-report measures that tap intolerance for general (Corstorphine, Mountford, Tomlinson, Waller, & Meyer, 2007; McHugh & Otto, 2012) or specific emotional distress (Buhr & Dugas, 2002; Harrington, 2005) and physical discomfort (Schmidt, Richey, & Fitzpatrick, 2006). Measures of *behavioral* intolerance for distress involve evaluating the duration of an individual's capacity to withstand an aversive task or stimulus, with shorter persistence being indicative of higher behavioral DI (Leyro et al., 2010). Several tasks have been developed and used to index behavioral DI, including persistence during exposure to thermal pain (Neufeld & Thomas, 1977; Rhudy & Meagher, 2003) or respiratory distress (Brown, Lejuez, Kahler, & Strong, 2002; Hajek, Belcher, & Stapleton, 1987; Marshall et al., 2008). Exposure to stressful/frustrating mental tasks (Lejuez, Kahler, & Brown, 2003; Quinn, Brandon, & Copeland, 1996; Strong et al., 2003) or to negative emotional pictures (Veilleux, Pollert, Zielinski, Shaver, & Hill, 2017) are used to measure intolerance for mental and emotional distress, respectively.

Notably, the convergent validity of self-report and behavioral indices of DI is typically low (McHugh et al., 2011) and DI (especially behavioral indices) can be influenced by contextual factors (e.g., Szuhany & Otto, 2015) including acute nicotine deprivation (Bernstein, Trafton, Ilgen, & Zvolensky, 2008) and task-difficulty (Daughters et al., 2005; MacPherson, Stipelman, Duplinsky, Brown, & Lejuez, 2008). Moreover, DI indices have discrepant predictive validity in terms of smoking-relevant outcomes, like short-term abstinence and puff topography (Farris et al., 2015; Perkins, Giedgowd, Karelitz, Conklin, &

Lerman, 2012). Measures of DI that are specific to smoking (relative to general) are posited to have increased predictive validity of smoking-relevant outcomes. For example, one self-report measure taps perceived DI for distress experienced during smoking abstinence (e.g., withdrawal; Sirota et al., 2010). However, no smoking-specific behavioral measures of DI have been developed.

To address these limitations, we developed the Contextual-Frustration Intolerance Typing Task (C-FiTT) to model contextual influences on persistence of goal-directed behavior, an index of DI. This task was developed based on theoretical principles of self-regulation and self-control (Baumeister, Bratslavsky, Muraven, & Tice, 1998). Self-regulation can be defined as goal-directed behavior that, in part, affords individuals the capacity to reduce discrepancy between standards and actual states, including thoughts, feelings, and behaviors (Hofmann, Schmeichel, & Baddeley, 2012). Self-regulation is posited to be a resource that can be depleted or conserved, much like physical energy (Muraven, Shmueli, & Burkley, 2006); thus demands on self-regulation resources (e.g., a challenging task) will limit availability for subsequent goal-directed behavior which can increase vulnerability for failure in self-regulation (Baumeister et al., 1998; Baumeister & Vonasch, 2015; Heatherton & Wagner, 2011). Failure of self-regulation is linked to the persistence of a wide range of health behaviors, including cigarette smoking (Baumeister & Vonasch, 2015). Self-regulation has been modeled utilizing a computer based re-typing task that requires individuals to retype a passage under conditions that are mentally taxing (instruction to type without use of letters “a” or “n” or spaces) which depletes self-regulation resources (Muraven et al., 2006). Similarly, behavioral DI marks one’s inability to persist in goal-directed activity in the face of distress, and is commonly assessed with a challenging task requiring persistence to complete a goal (e.g., complete task). Drawing upon this work, we modified the previously developed computer based re-typing task (Muraven et al., 2006) and adapted it to assess DI, such that an individual with higher DI would experience greater frustration in response to the mental demand of the typing task, and be less capable of persisting and completing the task relative to someone lower in DI.

In the current study, we tested the feasibility and initial validity of the C-FiTT in two different contexts – task difficulty (omitting frequent and infrequent letters: low versus high difficulty) and content (typing tobacco withdrawal versus neutral text). In order to evaluate construct validity, we examined the association between the C-FiTT persistence with an established measure of DI, the Distress Tolerance Scale (DTS; Simons & Gaher, 2005). Additionally, we evaluated persistence time and DTS in terms of negative affect reactivity to the task, where evidence for DI as an individual difference variable would be seen if (a) persistence time accounted for significant variance in negative affect reactivity and (b) both DI measures performed consistently in predicting negative affect reactivity. Finally, we tested the effect of C-FiTT versions (i.e., the contextual manipulation) on behavioral DI and task responding. There were three specific hypotheses regarding predictive validity: (1) Exposure to the high-difficulty task would produce higher DI (shorter persistence time; greater likelihood self-terminating task) and greater task responding (increases in negative affect and smoking urges), relative to the low-difficulty task; (2) Exposure to the withdrawal content would produce higher DI and greater task responding, relative to the neutral

content; and (3) Exposure to the high-difficulty task with withdrawal content would produce the highest DI and greatest task responding, relative to other conditions.

2 Material and methods

2.1 Participants

Participants ($N = 550$; $M_{age} = 44.6$, $SD = 13.5$; 52.2% female) were daily smokers recruited from all 50 states in the United States for an anonymous study on smoking and health. Data were collected online and participants were recruited through the Qualtrics Panels service in July of 2015. Potential participants were targeted based on the following pre-selected eligibility criteria: 18 years of age; a qualifying rate of 80% (satisfactory completion of 80% of surveys in which they participated); and a daily smoker. Interested individuals completed a study screening assessment and were eligible if they met the following criteria: (a) daily smoking for 1 year, (b) smoking 5 cigarettes/day, and (c) cigarettes were primary tobacco product of use. Participants were excluded if they reported having reduced their smoking rate by more than half in the past 1 month. See Table 1 for sample characteristics.

2.2 Measures

Items from the Smoking History Questionnaire (SHQ; Brown et al., 2002) were used to describe the sample in terms of smoking characteristics. Two additional items were used to assess same-day smoking (cigarettes smoked today and minutes since last cigarette). The Fagerstrom Test for Cigarette Dependence (FTCD; Fagerström, 2012) was used to assess level of tobacco dependence, with higher scores reflecting higher dependence (possible range 0–10). Internal consistency was low in the current sample ($\alpha = 0.55$), which is consistently found with this measure (Korte, Capron, Zvolensky, & Schmidt, 2013). The Patient Health Questionnaire (PHQ; Spitzer, Kroenke, & Williams, 1999) is a self-report assessment that was used to assess the severity of current depressive ($\alpha = 0.92$) and anxiety symptoms ($\alpha = 0.88$). The 14-item version of the Distress Tolerance Scale (DTS; Simons & Gaher, 2005) is a self-report measure used to assess perceived tolerance for emotional distress. Items are rated on a 5-point Likert-type scales ranging from 1 (strongly agree) to 5 (strongly disagree), assessing respondents' perceived ability to experience and endure negative emotional states. A mean score is computed (possible range is 1–5) with higher scores reflecting greater tolerance (lower intolerance) for distress. This scale has good psychometric properties, including high internal consistency and convergent validity (Simons & Gaher, 2005) and has been validated in daily cigarette smokers (Leyro et al., 2011). Internal consistency was $\alpha = 0.95$ in the current sample.

The *Positive and Negative Affect Schedule* (PANAS; Watson, Clark, & Tellegen, 1988) was used to assess state negative affect pre-post C-FiTT via the 10-item subscale (possible range 10–50). Internal consistency was $\alpha = 0.96$ (pre) and $\alpha = 0.95$ (post). The Questionnaire of Smoking Urges-Brief (QSU-B; Cox, Tiffany, & Christen, 2001) is a 10-item self-report measure of smoking urges that was completed pre-post C-FiTT, where items are rated on 0–100 scale and summed such that higher ratings indicate greater intensity of smoking urges (possible range 0–1000). Internal consistency was $\alpha = 0.96$ (pre) and $\alpha = 0.98$ (post).

2.2.1. C-FiTT task—C-FiTT was programed on Qualtrics. Participants were informed that they were going to complete a typing task to measure how well they “pay attention to details”. Administration involved visually presenting individuals with a passage of typed text with instructions to retype the 57-word passage. Participants completed a practice exercise before the real typing trial, and were encouraged to do the best they can without making any mistakes. Individuals were required to correctly retype the passage to complete the task. If an incorrectly re-typed response was submitted, an error message was displayed (“*The passage was incorrectly re-typed. Please try again*”), and individuals were required to re-start the typing. Participants were also instructed that they had the option to self-terminate the task: “*If after several attempts you feel that you are unable to continue, you can type the words ‘I Quit’ at any point to discontinue this task.*” DI was assessed by total trial length (persistence in seconds) and whether the task was self-terminated. Duration of time spent on the C-FiTT was programed to automatically record (in seconds). Participants were unaware that time was being recorded.

2.2.2. C-FiTT context—Two contextual factors were manipulated: task content and difficulty. The task content included a passage to read/re-type that was related to mountains (neutral) or tobacco withdrawal (context-specific). The withdrawal content was used to prime respondents to think about distress experienced during a cessation attempt, to increase the salience of the distressing stimuli. To manipulate task difficulty, the re-typing instructional set prompted individuals to omit the space bar and certain letters while typing that either occur frequently (“a” and “n”, high difficulty) or infrequently (“x” and “z”, low difficulty) in the passage. The difficulty associated with omitting typed letters and spaces increases cognitive load, depletes self-regulatory resources and produces negative affect (Muraven et al., 2006). The number of omitted characters were approximately matched for the low difficulty ($n = 3$) and high difficulty condition ($n = 51-53$). See supplemental online materials for additional C-FiTT information.

2.3. Procedures

A $2 \times 2 + 1$ experimental study design was utilized wherein participants were randomly assigned, stratified by biological sex, that crossed C-FiTT passage content (neutral v context-specific) $\times$ task difficulty (low v high difficulty). The combination of these conditions resulted in four experimental groups: (1) Neutral Context + Low-difficulty; (2) Neutral Context + High-difficulty; (3) Withdrawal Context + Low-difficulty; and (4) Withdrawal Context + High-difficulty. The design also included a neutral control condition, in which participants were asked to retype the neutral content without omitting any letters. Ratings of negative affect and smoking urges were completed immediately before and after C-FiTT. All participants were shown a written debriefing statement at the completion of the study. Participants received compensation in the form of Qualtrics credits, which can be used to purchase gift cards or other items through the Qualtrics Panels portal. Researchers are not informed of the specific number of credits that participants receive.

2.4. Data analytic approach

Four attentional check questions were imbedded in the survey (e.g., The correct answer for this question is “strongly agree”). Cases were retained if all four questions were correctly

answered. Additional cases were excluded if persistence time was < 10 s given that C-FiTT requires typing a passage of text and spending < 10 s on the task likely reflects an invalid test, and if time spent on the C-FiTT was > 600 s, which was informed by prior task persistence work (Chow & Lau, 2015; Leyro et al., 2010). See Fig. 1 for the CONSORT.

Data were analyzed in SAS 9.4. Preliminary analyses using chi-square analyses and one-way between-subject ANOVAs with post hoc comparisons were used to compare conditions on baseline factors, as a check of randomization. Descriptive statistics were utilized to examine feasibility of the C-FiTT, relative to the control condition. Within-condition mean changes in negative affect and smoking urges resulting from C-FiTT and the control condition were examined as indicators of initial task validity. Next, bivariate correlations between the DTS and persistence time were tested to examine construct validity.

Additionally, a multiple regression model was constructed wherein indicators of DI (persistence time from C-FiTT and DTS) were tested as predictors of post-task negative affect, controlling for pre-task negative affect. Subsequent analyses using OLS and logistic regression were conducted to examine the effect of the contextual manipulation on behavioral DI and task responding. Specifically, we tested the main effects of task difficulty and passage context to evaluate between-group differences with respect to: (1) task persistence (seconds), (2) likelihood of self-terminating C-FiTT (0 = completed; 1 = self-terminated), (3) changes in negative affect; and (4) changes in smoking urges. For between-condition comparisons, the main effects of task difficulty were examined using two dummy codes to reflect differences between the high-difficulty condition vs. control condition and low-difficulty condition vs. control condition. Follow-up tests using a third dummy code were used to examine differences between the high-difficulty condition vs. low-difficulty condition. The main effects of passage content were evaluated using one dummy code to reflect the difference between the withdrawal content vs. neutral content. Finally, the effect of the high-difficulty task with withdrawal content was examined using a single dummy code reflecting the difference between that group and the three other experimental groups. Model covariates included participant sex (0 = male; 1 = female), age, number of same-day cigarettes smoked, and tobacco dependence.¹

3 Results

3.1. Descriptive overview

Descriptive statistics for the sample are presented in Table 1. Mean $\pm$ standard deviation is presented. There were no significant differences between groups in terms of any demographic, smoking, or psychological symptoms.

3.2. Initial feasibility and validity of C-FiTT

A descriptive summary of the full sample as well as specific conditions in terms of smoking history, psychological symptom severity, and distress intolerance is presented in Table 2.

¹Post-hoc analyses revealed that models were unchanged when controlling for symptoms of depressive and anxiety symptoms severity. These results are available upon request and are not presented here given that the incremental validity of C-FiTT beyond psychological symptoms is beyond the scope of this initial validation study.

3.2.1. Invalid cases—As indicated in Fig. 1, of the 490 participants who completed a C-FiTT task (which excludes the 129 participants randomized to the control condition), 8.2% of cases demonstrated persistence time < 10 s. An additional 2.0% of cases demonstrated persistence time = 600 s. Thus, the C-FiTT resulted in invalid data for 10.2% of cases, and these cases were excluded. In comparison, the control condition resulted in 7.8% invalid cases. The excluded participants were significantly younger in age, had lower levels of tobacco dependence and smoked fewer cigarettes per day (p 's < 0.01) relative to cases that were retained. No other significant differences were observed.

3.2.2. Behavioral persistence—Among participants who completed any version of the C-FiTT ($n = 431$), the average persistence time was 94.1 ± 114.3 s and 64.7% of participants self-terminated the task. In contrast, average persistence time among participants in the control condition was 75.3 ± 94.7 s, which was not statistically different from the average C-FiTT persistence time ($t = -1.83$, $p = .068$). Approximately half (53.8%) of participants self-terminated the control task, which was significantly lower than the self-termination rate from C-FiTT ($\chi^2 = 4.77$, $p = .029$).

3.2.3. Construct validity—C-FiTT persistence time was not significantly correlated with the DTS ($r = 0.01$). Regarding the association between DI indices (persistence time and DTS) and emotional reactivity to C-FiTT, pre-task negative affect, and anxiety and depressive symptoms were entered as covarying factors. Persistence time and DTS each accounted for unique predictive variance in negative affect following the task. Shorter persistence time predicted significantly larger increases in negative affect ($b = -0.006$, $t = -3.46$, $p = .001$). Similarly, lower DTS scores predicted significantly larger increases in negative affect ($b = -0.823$, $t = -3.33$, $p = .001$).

3.3. Negative affect and smoking urges

C-FiTT produced increases in negative affect (small-medium effects; Table 3). The largest increases in negative affect were produced by the high-difficulty, withdrawal content, and the combination high-difficulty/withdrawal content versions of C-FiTT. The control condition produced a small-sized increase in negative affect. There were also small increases in smoking urges following C-FiTT, though these effects were evident only from the high-difficulty and combination high-difficulty/withdrawal content C-FiTT. The control condition did not produce changes in smoking urges.

3.4. Effects of contextual manipulation on C-FiTT responding

Between-condition differences were examined to determine the extent to which the contextual manipulation influenced behavioral DI, negative affect and smoking urges. Within-condition effect sizes are presented in Table 4. Between-condition effect sizes are presented in Table 5.

3.4.1. Behavioral persistence—Compared to the low-difficulty C-FiTT, the high-difficulty C-FiTT resulted in significantly shorter persistence time and a significantly greater likelihood of self-termination of C-FiTT. The combination high-difficulty/withdrawal content C-FiTT, compared to other C-FiTT conditions, also resulted in significantly shorter

task persistence and a significantly higher self-termination rate. There were no significant differences in task persistence for the neutral versus withdrawal content C-FiTT, although the withdrawal content C-FiTT resulted in a significantly greater likelihood of self-termination.

3.4.2. Negative affect—The high-difficulty C-FiTT produced significantly greater increases in negative affect compared the control condition and low-difficulty C- FiTT. The combination high-difficulty/withdrawal content C-FiTT compared to other C-FiTT versions produced significantly greater increases in negative affect. The content manipulation alone did not produce significant differences in negative affect change.

3.4.3. Smoking urges—The high-difficulty C-FiTT produced significantly greater increases in smoking urges compared to the control and low-difficulty C-FiTT conditions. The combination high-difficulty/withdrawal content C-FiTT compared to other C-FiTT versions, produced significantly greater increases in smoking urges. The content manipulation alone did not produce significant differences in smoking urge change.

4. Discussion

We found evidence for the initial feasibility and validity of C-FiTT. Four versions of the C-FiTT were administered and yielded sufficient variability in behavioral DI indicated by persistence time (in seconds) and self-termination of the task. Shorter task persistence was significantly related to greater emotional reactivity to the task. Of note, persistence time was not significantly correlated with the DTS, which is often the case when mixed-method approaches are used to measure of DI (McHugh et al., 2011) and is consistent with the multidimensional conceptualization of DI (Zvolensky et al., 2010). C-FiTT also produced small to medium sized-increases in negative affect and smoking urges. Findings also suggest that behavioral DI is influenced by contextual factors, particularly in the context of task difficulty manipulation. The high-difficulty C-FiTT produced significantly shorter task persistence and greater likelihood of self-terminating the task compared to the low-difficulty version. The self-termination rate in the high-difficulty condition was nearly double that of the low-difficulty condition (85.2% versus 45.5%). In addition, the high-difficulty version of C-FiTT resulted in significantly greater increases in negative affect and smoking urges relative to the low-difficulty and control conditions.

Contrary to expectation, the contextual priming for nicotine with-drawal yielded minimal independent effects on behavioral DI, negative affect, or smoking urges. This finding is surprising given that task salience influences physical distress intolerance (e.g., Branstetter-Rost, Cushing, & Douleh, 2009) and intolerance for *withdrawal* would theoretically be more salient in smokers, relative to general non-specific distress (Sirota, Rohsenow, Dolan, Martin, & Kahler, 2013). It is possible that the withdrawal text was not a sufficiently “potent” provocation. It also is possible that nicotine withdrawal priming was less salient to participants in the study given that they were not recruited on the basis of motivation for smoking cessation. Future refinement and/or modification to the C-FiTT content would be useful.

Interestingly, as hypothesized, the combination high-difficulty/ withdrawal version of C-FiTT produced significantly higher DI, a 100% self-termination rate, and greater task responding (increases in negative affect and smoking urges) relative to the other experimental conditions. Given that tobacco withdrawal produces acute physiological and emotional distress that is most severe during early phases of a quit attempt (Hughes, 2007), the combination high-difficulty/withdrawal version of C-FiTT may most closely resemble the context of a quit attempt in that the task requires persistence in an effortful task while being presented with withdrawal-relevant information.

Findings should be viewed in terms of their limitations, particularly of relevance for internet-based studies. This study relied largely on self-reported assessments, and participants were not monitored during their completion of the C-FiTT which could introduce bias due to effort and attention. The C-FiTT was also not programed to require a minimum or maximum time duration, although this is easily programmable and should be considered in future work to minimize occurrence of invalid data. We also did not provide reward for successful completion of the task, which could influence effort, although use of rewards has a potentially confounding effect on construct validity (Leyro et al., 2010) which was avoided in our study design. Additionally, the control condition had a high rate of self-termination perhaps reflecting low effort, thus the utility of this condition as a control was limited. Finally, the sample was primarily white and well-educated, thus generalizability of these findings to more heterogeneous sample of smokers may be limited.

Overall, there are several innovative features of the C-FiTT that enhance the contextual utility of this task: it is (a) theoretically-based; (b) the first smoking-specific behavioral DI task and can be used to model contextual factors relevant to smoking cessation; (c) easily accessible through an online, web-based platform which increases feasibility for use and dissemination; and (d) highly flexible, in that there are multiple versions of the task and the task can be modified with alternative clinically-relevant content (e.g., other substance use cues, specific emotional states). These are design features that enable researchers to administer a specific task version that meets the unique needs of the target sample or research focus. The concurrent validity of the C-FiTT in relation to other behavioral measures of DI and self-report smoking-specific DI (e.g., Intolerance for Smoking Abstinence Questionnaire; Sirota et al., 2010) is needed, in addition to the evaluation of additional validity indices (i.e., discriminant, known-groups, predictive, incremental validity). If future work further strengthens and supports the validity of this task, the C-FiTT could enhance future assessment and monitoring of DI across changing contexts.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

- Baumeister RF, Bratslavsky E, Muraven M, & Tice DM (1998). Ego depletion: Is the active self a limited resource? *Journal of Personality and Social Psychology*, 74(5), 1252–1265. 10.1037/0022-3514.74.5.1252. [PubMed: 9599441]

- Baumeister RF, & Vonasch AJ (2015). Uses of self-regulation to facilitate and restrain addictive behavior. *Addictive Behaviors*, 44, 3–8. <https://doi.org/10.1016/j.addbeh.2014.09.011>. [PubMed: 25267213]
- Bernstein A, Trafton J, Ilgen M, & Zvolensky MJ (2008). An evaluation of the role of smoking context on a biobehavioral index of distress tolerance. *Addictive Behaviors*, 33(11), 1409–1415. <https://doi.org/10.1016/j.addbeh.2008.06.003>. [PubMed: 18657912]
- Bold KW, Yoon H, Chapman GB, & McCarthy DE (2013). Factors predicting smoking in a laboratory-based smoking-choice task. *Experimental and Clinical Psychopharmacology*, 21(2), 133–143. 10.1037/a0031559. [PubMed: 23421357]
- Branstetter-Rost A, Cushing C, & Douleh T (2009). Personal values and pain tolerance: Does a values intervention add to acceptance? *The Journal of Pain*, 10(8), 887–892. 10.1016/j.jpain.2009.01.001. [PubMed: 19398384]
- Brown RA, Lejuez CW, Kahler CW, & Strong DR (2002). Distress tolerance and duration of past smoking cessation attempts. *Journal of Abnormal Psychology*, 111(1), 180–185. (Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11866171>). [PubMed: 11866171]
- Buhr K, & Dugas MJ (2002). The intolerance of uncertainty scale: Psychometric properties of the English version. *Behaviour Research and Therapy*, 40(8), 931–945. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12186356>. [PubMed: 12186356]
- Chow JT, & Lau S (2015). Nature gives us strength: Exposure to nature counteracts ego-depletion. *The Journal of Social Psychology*, 155(1), 70–85. [PubMed: 25310796]
- Corstorphine E, Mountford V, Tomlinson S, Waller G, & Meyer C (2007). Distress tolerance in the eating disorders. *Eating Behaviors*, 8(1), 91–97. 10.1016/j.eatbeh.2006.02.003. [PubMed: 17174856]
- Cox LS, Tiffany ST, & Christen AG (2001). Evaluation of the brief questionnaire of smoking urges (QSU-brief) in laboratory and clinical settings. *Nicotine & Tobacco Research*, 3(1), 7–16. 10.1080/14622200020032051. [PubMed: 11260806]
- Daughters S, Lejuez C, Bornovalova M, Kahler C, Strong D, & Brown R (2005). Distress tolerance as a predictor of early treatment dropout in a residential substance abuse treatment facility. *Journal of Abnormal Psychology*, 114(4), 729–734. [PubMed: 16351393]
- Fagerstrom KO (2012). Determinants of tobacco use and renaming the FTND to the Fagerstrom test for cigarette dependence. *Nicotine & Tobacco Research*, 14(1), 75–78. [PubMed: 22025545]
- Farris SG, Zvolensky MJ, Otto MW, & Leyro TM (2015). The role of distress intolerance for panic and nicotine withdrawal symptoms during a biological challenge. *Journal of Psychopharmacology*, 29(7), 783–791. 10.1177/0269881115575536. [PubMed: 25762651]
- Hajek P, Belcher M, & Stapleton J (1987). Breath-holding endurance as a predictor of success in smoking cessation. *Addictive Behaviors*, 12(3), 285–288. [PubMed: 3661283]
- Harrington N (2005). The frustration discomfort scale: Development and psychometric properties. *Clinical Psychology & Psychotherapy*, 12(5), 374–387. 10.1002/cpp.465.
- Heatherton TF, & Wagner DD (2011). Cognitive neuroscience of self-regulation failure. *Trends in Cognitive Sciences*, 15(3), 132–139. 10.1016/j.tics.2010.12.005. [PubMed: 21273114]
- Hofmann W, Schmeichel BJ, & Baddeley AD (2012). Executive functions and self regulation. *Trends in Cognitive Sciences*, 16(3), 174–180. 10.1016/j.tics.2012.01.006. [PubMed: 22336729]
- Hughes JR (2007). Effects of abstinence from tobacco: Valid symptoms and time course. *Nicotine & Tobacco Research*, 9(3), 315–327. 10.1080/14622200701188919. [PubMed: 17365764]
- Kahler CW, McHugh RK, Metrik J, Spillane NS, & Rohsenow DJ (2013). Breath holding duration and self-reported smoking abstinence intolerance as predictors of smoking lapse behavior in a laboratory analog task. *Nicotine & Tobacco Research*, 15(6), 1151–1154. 10.1093/ntr/nts231. [PubMed: 23132658]
- Korte KJ, Capron DW, Zvolensky M, & Schmidt NB (2013). The Fagerstrom test for nicotine dependence: Do revisions in the item scoring enhance the psychometric properties? *Addictive Behaviors*, 38(3), 1757–1763. 10.1016/j.addbeh.2012.10.013. [PubMed: 23254226]
- Lejuez CW, Kahler CW, & Brown RA (2003). A modified computer version of the paced auditory serial addition task (PASAT) as a laboratory-based stressor. *The Behavior Therapist*, 26, 290–293.

- Leventhal AM, & Zvolensky MJ (2015). Anxiety, depression, and cigarette smoking: A transdiagnostic vulnerability framework to understanding emotion-smoking comorbidity. *Psychological Bulletin*, 141(1), 176–212. 10.1037/bul0000003. [PubMed: 25365764]
- Leyro TM, Bernstein A, Vujanovic AA, McLeish AC, & Zvolensky MJ (2011). Distress tolerance scale: A confirmatory factor analysis among daily cigarette smokers. *Journal of Psychopathology and Behavioral Assessment*, 33(1), 47–57. 10.1007/s10862-010-9197-2. [PubMed: 23935238]
- Leyro TM, Zvolensky MJ, & Bernstein A (2010). Distress tolerance and psychopathological symptoms and disorders: A review of the empirical literature among adults. *Psychological Bulletin*, 136(4), 576–600. 10.1037/a0019712. [PubMed: 20565169]
- MacPherson L, Stipelman BA, Duplinsky M, Brown RA, & Lejuez CW (2008). Distress tolerance and pre-smoking treatment attrition: Examination of moderating relationships. *Addictive Behaviors*, 33(11), 1385–1393. 10.1016/j.addbeh.2008.07.001. [PubMed: 18706768]
- Marshall EC, Zvolensky MJ, Vujanovic AA, Gregor K, Gibson LE, & Leyro TM (2008). Panic reactivity to voluntary hyperventilation challenge predicts distress tolerance to bodily sensations among daily cigarette smokers. *Experimental and Clinical Psychopharmacology*, 16(4), 313–321. 10.1037/a0012752. [PubMed: 18729685]
- McHugh RK, Daughters SB, Lejuez CW, Murray HW, Hearon BA, Gorka SM, & Otto MW (2011). Shared variance among self-report and behavioral measures of distress intolerance. *Cognitive Therapy and Research*, 35, 266–275. 10.1007/s10608-010-9295-1. [PubMed: 23894216]
- McHugh RK, & Otto MW (2012). Refining the measurement of distress intolerance. *Behavior Therapy*, 43(3), 641–651. 10.1016/j.beth.2011.12.001. [PubMed: 22697451]
- Muraven M, Shmueli D, & Burkley E (2006). Conserving self-control strength. *Journal of Personality and Social Psychology*, 91(3), 524–537. 10.1037/0022-3514.91.3.524. [PubMed: 16938035]
- Neufeld RWJ, & Thomas P (1977). Effects of perceived efficacy of a prophylactic controlling mechanism on self-control under pain stimulation. *Canadian Journal of Behavioral Science*, 9, 224–232.
- Perkins KA, Giedgowd GE, Karelitz JL, Conklin CA, & Lerman C (2012). Smoking in response to negative mood in men versus women as a function of distress tolerance. 14(12), 1418–1425. 10.1093/ntr/nts075.
- Perkins KA, Karelitz JL, Giedgowd GE, Conklin CA, & Sayette MA (2010). Differences in negative mood-induced smoking reinforcement due to distress tolerance, anxiety sensitivity, and depression history. *Psychopharmacology*, 210(1), 25–34. 10.1007/s00213-010-1811-1. [PubMed: 20217051]
- Quinn EP, Brandon TH, & Copeland AL (1996). Is task persistence related to smoking and substance abuse? The application of learned industriousness theory to addictive behaviors. *Experimental and Clinical Psychopharmacology*, 4(2), 186–190. Retrieved from <http://psycnet.apa.org/index.cfm?fa=buy.optionToBuy&id=1996-00322-008>.
- Rhudy JL, & Meagher MW (2003). Negative affect: Effects on an evaluative measure of human pain. *Pain*, 104(3), 617–626. (Retrieved from) <http://www.ncbi.nlm.nih.gov/pubmed/12927634>. [PubMed: 12927634]
- Schmidt NB, Richey JA, & Fitzpatrick KK (2006). Discomfort intolerance: Development of a construct and measure relevant to panic disorder. *Journal of Anxiety Disorders*, 20(3), 263–280. 10.1016/j.janxdis.2005.02.002. [PubMed: 16564432]
- Simons JS, & Gaher RM (2005). The distress tolerance scale: Development and validation of a self-report measure. *Motivation and Emotion*, 29(2), 83–102. 10.1007/s11031-005-7955-3.
- Sirota AD, Rohsenow DJ, Dolan SL, Martin RA, & Kahler CW (2013). Intolerance for discomfort among smokers: Comparison of smoking-specific and nonspecific measures to smoking history and patterns. *Addictive Behaviors*, 38(3), 1782–1787. <https://doi.org/10.1016/j.addbeh.2012.10.009>. [PubMed: 23254229]
- Sirota AD, Rohsenow DJ, MacKinnon SV, Martin RA, Eaton CA, Kaplan GB, ... Swift RM (2010). Intolerance for smoking abstinence questionnaire: Psychometric properties and relationship to tobacco dependence and abstinence. *Addictive Behaviors*, 35(7), 686–693. 10.1016/j.addbeh.2010.02.014. [PubMed: 20381260]
- Spitzer RL, Kroenke K, & Williams JB (1999). Validation and utility of a self-report version of PRIME-MD: The PHQ primary care study. Primary care evaluation of mental disorders. Patient

health questionnaire. JAMA, 282(18), 1737–1744. (Retrieved from) <http://www.ncbi.nlm.nih.gov/pubmed/10568646>. [PubMed: 10568646]

- Strong DR, Lejuez CW, Daughters S, Marinello M, Kahler CW, & Brown RA (2003). The computerized mirror tracing task (Version 1) [Unpublished manual]. Retrieved from
- Szuhany KL, & Otto MW (2015). Contextual influences on distress intolerance: Priming effects on behavioral persistence. *Cognitive Therapy and Research*, 39(4), 499–507. 10.1007/s10608-015-9672-x. [PubMed: 26366022]
- Trujillo MA, Khoddam R, Greenberg JB, Dyal SR, Ameringer KJ, Zvolensky MJ, & Leventhal AM (2017). Distress tolerance as a correlate of tobacco dependence and motivation: Incremental relations over and above anxiety and depressive symptoms. *Behavioral Medicine (Washington, D.C.)*, 43(2), 120–128. 10.1080/08964289.2015.1110559.
- Veilleux JC, Pollert GA, Zielinski MJ, Shaver JA, & Hill MA (2017). Behavioral assessment of the negative emotion aspect of distress tolerance. *Assessment*, 10731911166898110.1177/1073191116689819.
- Watson D, Clark LA, & Tellegen A (1988). Development and validation of brief measures of positive and negative affect: The PANAS scales. *Journal of Personality and Social Psychology*, 54(6), 1063–1070.
- Zvolensky MJ, Leyro TM, Bernstein A, & Vujanovic AA (2010). Historical perspectives, theory, and measurement of distress tolerance In Zvolensky MJ, Bernstein A, & Vujanovic AA (Eds.). *Distress tolerance: Theory, research, and clinical applications*. Guilford: New York, NY.

HIGHLIGHTS

- C-FiTT is a new task that considers the impact of contextual factors on DI
- C-FiTT produces increases in negative affect and smoking urges
- DI is increased when smokers are exposed to a high versus low-difficulty task
- DI is highest when smokers are primed for withdrawal paired with high-difficulty task

Fig. 1.
CONSORT diagram

Table 1

Sample demographics.

	Mean/n	SD/%
Sex		
Male	263	47.8
Female	287	52.2
Age	44.6	13.5
Race		
White	490	89.1
Black	25	4.5
Asian	13	2.4
American Indian/Alaska Native	10	1.8
Native Hawaiian or Other Pacific Islander	3	0.5
Other	9	1.6
Ethnicity		
Non-Hispanic	513	93.3
Hispanic	37	6.7
Marital Status		
Married	284	51.6
Never married	141	25.6
Divorced	94	17.1
Widowed	20	3.6
Separated	11	2.0
Educational Attainment		
Less than high school	2	0.4
Part of high school	9	1.6
High school diploma or the equivalent	150	27.3
Some undergraduate college	146	26.5
Associate degree	74	13.5
Bachelor's degree	117	21.3
Master's degree	41	7.5
Professional or Doctoral Degree	11	2.0
Employment Status		
Employed full time	275	50.0
Homemaker	66	12.0
Retired	60	10.9
Employed part time	52	9.5
Unemployed	47	8.5
Disability	40	7.3
Student	10	1.8

Table 2

Descriptive overview.

	Full Sample (n = 550)	Control (n = 119)	Neutral + Low Difficulty (n = 120)	Neutral + High Difficulty (n = 102)	Withdrawal + Low Difficulty (n = 102)	Withdrawal + High Difficulty (n = 107)
Smoking History						
Tobacco Dependence (FTCD)	5.4 ± 2.0	5.6 ± 2.0	5.4 ± 2.1	5.6 ± 1.9	5.2 ± 2.1	5.3 ± 1.8
Cigarettes/day	17.2 ± 8.5	18.5 ± 9.1	17.3 ± 8.2	17.0 ± 8.5	16.2 ± 8.2	16.8 ± 8.5
Years smoking	26.1 ± 14.2	27.3 ± 12.6	25.3 ± 15.2	24.9 ± 13.6	26.0 ± 14.8	27.1 ± 14.8
Number/cigarettes (same-day)	9.4 ± 6.8	9.7 ± 7.4	9.3 ± 6.5	9.0 ± 6.0	9.2 ± 7.0	9.5 ± 6.9
Time (minutes) since last cigarette ^a	25.5 ± 50.5	26.0 ± 49.9	26.0 ± 35.8	28.5 ± 76.5	24.1 ± 51.7	23.1 ± 29.3
Psychological Symptom Severity						
Depressive symptom severity (PHQ)	7.0 ± 6.9	7.8 ± 7.0	6.3 ± 6.3	6.8 ± 6.6	6.5 ± 7.2	7.6 ± 7.5
Anxiety symptom severity (PHQ)	4.8 ± 3.9	5.3 ± 3.9	4.3 ± 3.7	5.0 ± 4.0	4.3 ± 3.9	5.0 ± 4.0
Distress intolerance						
Distress Tolerance Scale (DTS)	3.3 ± 1.0	3.3 ± 1.0	3.2 ± 1.1	3.2 ± 1.0	3.2 ± 1.1	3.3 ± 1.0

Note: One-way between subject ANOVAs, using post hoc Tukey corrections, revealed no significant differences between the groups on any of the baseline variables.

^a (n = 11 cases reported smoking zero same-day cigarettes thus were not included in average time since last cigarette).

Table 3

Descriptive summary of C-FiTT outcomes.

Condition	Persistence		Self-Termination n (%)	Negative Affect		Smoking Urges	
	M ± SD	Skew (Kurtosis)		Pre	Post	Pre	Post
Control (n = 119)	75.3 ± 94.7	2.41 (5.75)	64 (53.8)	14.2 ± 7.3	15.3 ± 8.2	442.6 ± 312.7	424.8 ± 345.3
Neutral + Low Difficulty (n = 120)	125.4 ± 128.3	1.21 (0.611)	49 (40.8)	14.6 ± 8.4	15.6 ± 8.8	417.0 ± 301.7	404.5 ± 345.7
Neutral + High Difficulty (n = 102)	69.9 ± 85.3	2.15 (4.19)	71 (69.6)	14.7 ± 7.9	16.7 ± 9.1	447.1 ± 312.4	495.8 ± 362.1
Withdrawal + Low Difficulty (n = 102)	128.7 ± 140.9	1.52 (1.76)	52 (51.0)	15.3 ± 8.9	16.1 ± 9.1	431.4 ± 303.2	428.6 ± 338.1
Withdrawal + High Difficulty (n = 107)	49.1 ± 61.4	2.96 (9.34)	107 (100.0)	15.3 ± 8.2	18.3 ± 9.6	418.4 ± 298.0	473.9 ± 345.0

Table 4
Acute changes in negative affect and smoking urges from C-FiTT (Within-Condition Effects).

Condition	Negative Affect			Smoking Urges		
	<i>t</i>	<i>p</i>	<i>d</i>	<i>t</i>	<i>p</i>	<i>d</i>
Control	2.62	0.01	0.25	-1.20	0.233	-0.11
Low Difficulty	3.55	0.001	0.24	-0.76	0.446	-0.05
High Difficulty	6.40	< 0.001	0.45	4.29	< 0.001	0.30
Neutral Text	4.90	< 0.001	0.33	1.53	0.129	0.10
Withdrawal Text	5.28	< 0.001	0.40	2.10	0.036	0.15
High Difficulty + Withdrawal	4.95	< 0.001	0.49	3.18	0.002	0.32

Table 5

Between-condition effects.

Condition comparisons	Task persistence			Self-termination			Negative affect			Smoking urges		
	<i>t</i>	<i>p</i>	<i>d</i>	OR	<i>p</i>	<i>V</i>	<i>t</i>	<i>p</i>	<i>d</i>	<i>t</i>	<i>p</i>	<i>d</i>
Control vs. Low Difficulty	4.28	< 0.001	0.45	1.39	0.145	-0.13	-0.28	0.780	0.07	0.51	0.610	0.03
Control vs. High Difficulty	1.31	0.190	0.19	4.93	< 0.001	0.16	2.78	< 0.001	0.26	3.70	< 0.001	0.17
Low Difficulty vs. High Difficulty	6.62	< 0.001	0.65	6.88	< 0.001	0.42	3.51	< 0.001	0.19	3.67	< 0.001	0.20
Neutral vs. Withdrawal Content	1.13	0.261	0.03	4.08	< 0.001	0.23	1.20	0.232	0.13	1.20	0.233	0.08
High Difficulty/Withdrawal vs. Others	-4.71	< 0.001	0.65	77.55 ^a	< 0.001	0.43	3.53	< 0.001	0.24	2.37	0.018	0.10

^aDenotes value is Chi-square.