Published in final edited form as:

Soc Sci Med. 2021 February ; 271: 112038. doi:10.1016/j.socscimed.2018.11.018.

Affective Response to Physical Activity as an Intermediate Phenotype

Harold H. Lee¹, Jessica A. Emerson¹, Lauren Connell Bohlen², David M. Williams¹

¹Department of Behavioral and Social Sciences, Brown University School of Public Health, Providence, RI, USA

²Department of Kinesiology, University of Rhode Island, Kingston, RI, USA

Abstract

Over the past seventy years, biomedical and epidemiological research has shown that regular physical activity (PA) is critical for physical and mental health. Despite this knowledge, physical inactivity is the fourth leading risk factor for global mortality, accounting for 9% (5.3 million) of premature deaths annually. We suggest this mismatch between knowing about the risks of PA and engaging in regular PA can be reconciled by focusing less on expected health benefits of PA and more on how people feel during PA. Specifically, in this position paper, we argue that affective response (feeling good versus bad) to PA is an intermediate phenotype that can explain significant variance in PA behavior and is, in turn, a function of genetic variability. In making this argument, we first review empirical evidence showing that affective response to PA predicts future physical activity behavior. Second, we systematically review research on single nucleotide morphisms (SNPs) that are associated with affective response to PA. Investigating affective response to PA as an intermediate phenotype will allow future researchers to move beyond asking “*What* SNPs are associated with PA?”, and begin asking “*How* do these SNPs influence PA?”, thus ultimately optimizing the translation of knowledge gained from genomic data to intervention development.

Keywords

Affect; exercise; intermediate phenotype; physical activity; motivation; mechanism

1. Introduction

Over the past seventy years, biomedical and epidemiological research has shown that regular physical activity (PA) is critical for physical and mental health (Garber et al., 2011; Haskell et al., 2007; Horstmann, 1950; Schuch et al., 2011; Wennesland et al., 1959). Despite this knowledge, physical inactivity is the fourth leading risk factor for global mortality, accounting for 9% (5.3 million) of premature deaths globally (Lee et al., 2012).

The low rates of PA are a function of a wide range of environmental factors, but a substantial proportion (48–71%) of variance in habitual PA can also be attributed to genomic variability

(Aaltonen et al., 2010; Bouchard & Perusse, 1994; Cai et al., 2006; Carlsson et al., 2006; Eriksson et al., 2006; Fisher et al., 2010; Franks et al., 2005; Joosen et al., 2005; Maia et al., 2002; B. D. Mitchell et al., 2003; Simonen et al., 2002; Stubbe et al., 2006). Research identifying the specific genetic variants, or single nucleotide polymorphisms (SNPs), has been stagnant in the past decade (chiefly due to small sample sizes). However, large-scale genomic data collection efforts, such as NIH's precision medicine initiative cohort (i.e., "All of Us": (<https://allofus.nih.gov/>), may identify what SNPs are associated with PA (Collins & Varmus, 2015).

To optimize the translation of discoveries from large-scale genomic data to behavioral intervention development, it is necessary to identify intermediate phenotypes—i.e., traits that mediate the effects of the genome on the behavioral phenotype, in this case PA. Accordingly, the goal of this paper is to examine affective response to PA (i.e., how one feels during PA) as a potential intermediate phenotype. Similar intermediate phenotypes have been hypothesized and received empirical support in the context of other health behaviors, including subjective and affective responses to the consumption of alcohol (Ray & Hutchison, 2004; Ray et al., 2010), caffeine (Alsene et al., 2003; Dlugos et al., 2007; Kendler & Prescott, 1999; Loke, 1988), amphetamines (Hart et al., 2011; Lott et al., 2005; Lott et al., 2006), and nicotine (Hoft et al., 2011; Hutchison et al., 2007; Sherva et al., 2008; Zeiger et al., 2007). If affective response to PA proves to be a viable intermediate phenotype, it can advance understanding of PA behavior and potentially provide new opportunities for PA promotion intervention.

In proposing affective response to PA as an intermediate phenotype, we will first explicate the general concept of an intermediate phenotype and introduce its criteria. We will then examine whether affective response to PA is a conceptually plausible intermediate phenotype for understanding the effects of genes on PA behavior. Specifically, we will summarize findings from a recent systematic review that examined the effect of affective response to PA on future PA behavior. We will then conduct our own systematic review to examine genetic variants that are associated with affective response to PA. Finally, we will discuss strengths and limitations of the literature and implications of applying an intermediate phenotype approach to affective response to PA.

2. Definition of intermediate phenotype

An intermediate phenotype is a phenotype that is posited to mediate the effects of the genotype on a more distal phenotype such as an outcome or behavior of interest. For a variable to be considered an intermediate phenotype, the variable should be predictive of the distal phenotype of interest, and the variable should be influenced by genetic variation.

The term *intermediate phenotype* is often interchangeably used with *endophenotype*—i.e., "measurable components unseen by the unaided eye along the pathway between disease and distal genotype"—which originated from genetic research in schizophrenia (Gottesman & Gould, 2003). There are subtle differences between the two concepts (for further discussion of this distinction see (Goldman & Ducci, 2007)), but the pragmatic utility of both approaches for public health research is that they shift the focus away from the clinical

outcome—e.g., disease or behavioral outcome proximal to the disease—and toward psychobiological variables that are mechanistically closer to the genotype. By way of analogy, knowledge regarding psychobiological mediators between genetic variation and health behaviors may provide additional insights into the causes of negative health behaviors (e.g., PA, smoking, drinking, and drug abuse) in the same way that knowledge regarding psychosocial mediators between socioeconomic status and health behaviors provide insights into alleviation of health disparities. In other words, knowledge regarding intermediary mechanistic processes between genome and behaviors are crucial as they allow researchers to move beyond *predicting* behaviors based on genomic profile and start investigating *how to change* behaviors.

3. Affective response to physical activity as a potential intermediate phenotype

Affective response to PA is a conceptually plausible intermediate phenotype because it may have a causal influence on PA behavior. Consistent with the ancient principle of psychological hedonism, people tend to repeat behaviors that feel good and avoid behaviors that feel bad (Bentham, 1789/2007; Hobbes, 1789/2008; Kahneman et al., 1997; Mill, 1861/2012; Usener, 1887/2010; Williams, 2018). When applied to PA behavior, those who experience a more positive affective response to PA are more likely to repeat it in the future, and thus more likely to engage in regular PA (Williams, 2008).

Affective response to PA is defined herein as the shift in core affective valence, ranging from good/pleasure to bad/displeasure (Russell, 1980), from immediately *prior to* PA to *during* PA. We focus on *core affective valence* rather than discreet moods or emotions because core affective valence is pervasive in human experience, both within and outside the context of various moods and emotions, and has a strong theoretical basis regarding its relationship to behavior—i.e., psychological hedonism. We focus on affective response *during* PA (i.e., change in affect from prior to PA to during PA) rather than affective response following PA (i.e., change in affect from prior to PA to after PA), because the proximal consequences of a behavior more robustly predict future behavior compared to more distal consequences (Hall, 1976). Moreover, affective response following PA may be better conceptualized as how one feels as a result of completing a session of PA, rather than affective response to PA behavior itself.

Despite frequent media messages that “exercise feels good” (Backhouse et al., 2007) and evidence suggesting that most people feel good after PA (Berger & Motl, 2000; Reed & Ones, 2006), numerous studies have shown that affective response *during* moderate intensity PA—the intensity of PA on which national recommendations are based (USDHHS, 2008)—varies greatly from person to person. Some people tend to feel bad during PA, other typically feel good, whereas others feel neither good nor bad (Backhouse et al., 2007; Ekkekakis et al., 2005; Parfitt et al., 2006; Rose & Parfitt, 2007; Van Landuyt et al., 2000; Welch et al., 2007). Some of the variance in affective response to moderate intensity PA may involve situational factors, such as the physical and/or social setting (Dunton et al., 2015; Gellert et al., 2011), or individual factors such as fitness level or overweight status (Ekkekakis et al.,

2010; Ekkekakis et al., 2016). We posit that a significant proportion of the variability in affective response to moderate intensity PA may be a function of genetic variance, as is the case for affective response to the consumption of alcohol (Ray & Hutchison, 2004; Ray et al., 2010), caffeine (Alsene et al., 2003; Dlugos et al., 2007; Kendler & Prescott, 1999; Loke, 1988), amphetamines (Hart et al., 2011; Lott et al., 2005; Lott et al., 2006), and nicotine (Hoft et al., 2011; Hutchison et al., 2007; Sherva et al., 2008; Zeiger et al., 2007).

In the next two sections, we review the available empirical evidence that directly bears on affective response to PA as a potential intermediate phenotype: whether it is a function of genetic variability (Figure 1, A-path) and whether it has downstream causal effects on PA behavior (Figure 1, B-path). We examine the latter hypothesis first because there is a more robust literature on this topic.

4. Evidence for affective response to physical activity as a predictor of physical activity behavior

Rhodes and Kates (2015) performed a systematic review of evidence on affective response to PA as a potential predictor of PA behavior. The authors identified 24 studies that examined some version of this relationship based on literature published up to July 2014. Among the 24 studies, four specifically examined the relationship between affective response to PA and its association with PA behavior, of which three were longitudinal (Kwan & Bryan, 2010; Williams et al., 2008; Williams et al., 2012) and one was cross-sectional (Schneider et al., 2009).

All four studies examining affective response to PA were predictive of PA behavior, despite heterogeneity in study designs, intensity of PA, and assessments. For example, in a cross-sectional design, Schneider et al. (2009) assessed affective response to moderate (80 % of ventilatory threshold) and high intensity (above ventilatory threshold) cycling, and its association with accelerometry-measured moderate to vigorous PA among 124 adolescents of mixed PA status (i.e., inactive and physically active). Affective response was assessed before and during exercise at minutes 10, 20, and 40 using the Feeling Scale, a one-item measure of affective valence (i.e., feeling good versus bad; Hardy & Rejeski, 1989). In a longitudinal observational study, Kwan and Bryan (2010) assessed affective response to a 30-min session of PA at moderate intensity (i.e., 65 % of VO_{2max}) and its relationship with change in self-reported PA 3 months later among 127 adults. Affective response was assessed before and during exercise at minutes 5, 10, 20, and 30 using the Physical Activity Affect Scale, a 12-item questionnaire that measures four subscales: positive affect, negative affect, tranquility, and fatigue (Lox et al., 2000). In another longitudinal observational study, Williams et al. (2008) assessed affective response to moderate intensity PA (i.e., 64–76 % of maximum predicted heart rate) and its relationship with change in self-reported PA six and 12 months later among 31 sedentary adults. Affective response was assessed every two minutes during a 30-minute treadmill walk using the Feeling Scale. Finally, Williams et al. (2012) evaluated affective response to during a 10-min moderate intensity walking bout (i.e., 2.5–4.0 mph on a treadmill) and its relationship with change in self-reported PA 6 months later among 146 adults, again using the Feeling Scale to assess affective response before and

during exercise at minutes 2, 5, and 8 during a 10-minute treadmill walk. Despite the heterogeneity in procedures (e.g., operationalization of moderate intensity) and assessment of PA, all four studies that examined the association between affective response to PA and future PA showed a positive association, ranging from $r=.18$ to $r=.51$. That is, participants who reported more positive shifts in affective response to PA tended to engage in more future PA.

Of note, all four studies were considered high quality research based on evaluation of the risk of bias by Rhodes and Kates (2015) using a modified version of the checklist created by Downs and Black (1998). For example, affect experienced prior to the PA session was controlled in all four studies, and three of the four studies controlled for past PA behavior (Kwan & Bryan, 2010; Williams et al., 2008; Williams et al., 2012).

Studies that examined the association between affective response after PA and future PA behavior had overall null effects (Rhodes & Kates, 2015). Taken together, the existing research generally supports the overall premise that affective response to PA is predictive of PA behavior (B-path, Figure 1), although there are some limitations which we address below in the discussion. Next, though, we turn to the evidence for genetic influence on affective response to PA.

5. Evidence for genetic influence on affective response to physical activity

In order for affective response to PA to be an intermediate phenotype, genetic variation should explain differences in the putative intermediate phenotype. Schutte et al. (2016) estimated the heritability of affective response during and after PA among adolescents ($n=226$ twins and $n=38$ siblings) and found that genetic factors explained 15% of the between-person variance in affective response to PA during a graded exercise test. While this study suggests low to moderate heritability of affective response to PA, it does not provide insight into which particular gene loci account for the differences in affective response to PA.

To identify the specific genetic variants associated with affective response to PA, we conducted a systematic review. Specifically, in accordance with the PRISMA checklist (Moher et al., 2009), we searched PubMed and PsycINFO using search terms specific to genetic variants, affective response, and PA behavior. Consistent with a previous systematic review of genetic variants (Bruneau et al., 2016), search terms used for genetic variants were “genetic variants,” “genotype,” “single nucleotide polymorphism,” “allele,” and “polymorphism”. Consistent with a previous systematic review of affective response to PA (Rhodes & Kates, 2015), search terms used for affect included “affect,” “emotion,” “mood,” “activation,” “circumplex model,” “exercise induced feeling states,” “subjective exercise experience,” and “pleasant feeling state”. Consistent with a previous systematic review in the PA literature (Johnson et al., 2014), search terms used for PA were “physical activity,” “exercise,” “running,” “bicycle,” “treadmill,” “speed training,” “training duration,” “training frequency,” “training intensity,” and “aerobic endurance”.

To be eligible for review, studies must (i) assess genotype, (ii) include a measure of affective response to PA, and (iii) report an association between the two variables. We excluded articles that examined affect that was unrelated to PA (e.g., depressive symptoms), affective response after PA, and non-human studies. Finally, we only included peer-reviewed articles published in English.

The literature search included studies published between January 1, 1970 and November 1, 2017. Title and abstract screening was done by the lead author and selected papers (n=19) were reviewed by co-authors to determine the final articles included in the systematic review.

Out of 634 articles from PubMed and PsychInfo, 19 articles were relevant based on the title and abstract screening, and only two articles met inclusion criteria after full-text review (Figure 1). Among the 19 articles, three papers were excluded because they were non-human studies and 14 papers were excluded because they did not assess affective response to PA. Of note, three of the excluded studies examined pre to post changes in mood resulting from a 4-week PA intervention (Dotson et al., 2016; Gujral et al., 2014; Hopkins et al., 2012) but did not examine affective response during PA. The two included studies were Bryan et al. (2007) and Karoly et al. (2012).

Bryan et al. (2007) investigated the association between the SNP in brain derived neurotrophic factor (BDNF) and affective response to treadmill walking. The human *BDNF* gene, located on chromosome 11p14.1, encodes a BDNF protein that is widely expressed in the mammalian adult brain and plays an essential role in the growth and survival of neuronal cells in the peripheral and central nervous system (Hofer et al., 1990; Murer et al., 2001). In the *BDNF* SNP, rs6265, some people carry the G allele, while approximately 1% of African, 19%, of European Caucasians, 21% of Hispanic, and 62% of Asians are A-allele carriers (Sherry et al., 2001). Investigators asked 64 healthy participants (32 men and 32 women; mean age = 23.8 years) walk on a treadmill at moderate intensity (65% VO_{2max}) for 30 minutes and assessed affective response during the treadmill walk. Compared to A-allele-carriers (AG/AA), the GG-allele carriers for *BDNF*rs6265 showed a more negative affective response to PA.

Karoly et al. (2012) investigated 14 SNPs and their associations with physiological and subjective responses to aerobic PA among 238 sedentary adults. Investigators assessed affective response to treadmill walking at moderate intensity (65% VO_{2max}) for 30 minutes. Among the 14 SNPs assessed, two SNPs (rs8044769 and rs3751812) in the *Fat Mass and Obesity-associated gene* (FTO) were associated with affective response to PA. Specifically, C-allele-carriers (relative to TT-allele-carriers) of rs8044769 and T-allele-carriers (relative to GG-allele-carriers) of rs3751812 had greater increase in positive affect over the course of the 30-minute treadmill walk. The other 12 SNPs were not associated with affective response to PA.

Taken together, findings from our systematic review generally support the premise that affective response to PA is a function of genetic variability (Figure 1, a-path). However, as

with affective response to PA as a predictor of PA behavior, there are some limitations to the existing research, which we address below.

6. Discussion

We examined whether affective response to PA is a conceptually plausible intermediate phenotype that mediates genetic effects on PA behavior. Specifically, we summarized a recent systematic review that examined the effect of affective response to PA on PA behavior (Rhodes & Kates, 2015), and we conducted our own systematic review to examine genetic variants that are associated with affective response to PA.

The systematic review that examined the relationship between affective response to PA and its impact on future PA (B-path, Figure 1) (Rhodes & Kates, 2015) revealed four studies showing relationships between affective response and PA behavior (Kwan & Bryan, 2010; Schneider et al., 2009; Williams et al., 2008; Williams et al., 2012) in the small to large effect size. More research is needed, however. Specifically, it will be important to examine whether the same association exists among diverse populations in terms of age, ethnicity, culture, and health status. Additionally, in the previous studies, affective response to PA was measured in the laboratory by research assistants, which may bias participants' responses due to factors such as social desirability (Paulhus, 1991). To improve the ecological validity—i.e., the extent to which the outcome of a trial can be generalized to real-world contexts— affective response to PA should be studied in a more natural setting (e.g., outdoors, gym, or park), which can be achieved through the use of ecological momentary assessment. Since the systematic review by Rhodes and Kates (2015), at least one study has used ecological momentary assessment to investigate the effect of affective response to PA on future PA, again showing support for the hypothesized association (Williams et al., 2016). Finally, it should be noted that the aforementioned research does not allow us to infer causality. That is, even when researchers control for prior PA, an unmeasured third variable could confound the findings by influencing both affective response to PA and future PA level.

Regarding the genetic influence on PA behavior, the present systematic review revealed two studies that investigated genetic variants associated with affective response to PA (Figure 1, A-path). In one study, the G allele of *BDNF*rs6265 was associated with negative affective response to PA (Bryan et al., 2007). In previous research the G allele of rs6265 was also associated with low intrinsic motivation for PA (Hooper et al., 2014), lower adherence in response to a PA promotion intervention (Bryan et al., 2013), and obesity (Gong et al., 2013; Gunstad et al., 2006; León-Mimila et al., 2013; J. A. Mitchell et al., 2013; Shugart et al., 2009; Thorleifsson et al., 2009; Wen et al., 2012; Wu et al., 2010; Xi et al., 2013; Zhao et al., 2009). Further, PA can increase serum BDNF, which may create a feedback loop (*BDNF*→PA) that could magnify the genetic effects of the *BDNF* phenotype on the PA behavior phenotype (Currie et al., 2009; J. Zoladz et al., 2008; J. A. Zoladz et al., 1998). However, more research is needed to study the *BDNF* genetic variant given the relatively small sample size ($n < 100$) of the Bryan et al (2007) study.

In the study by Karoly et al (2012), people who are C-allele-carriers of rs8044769 and T-allele-carriers of rs3751812 on the *FTO* gene had a more positive affective response to PA.

Prior studies have shown that C-allele-carriers of rs8044769 and T-allele-carriers of rs3751812 have higher body mass index (Hassanein et al., 2010; Thorleifsson et al., 2009; Wing et al., 2009). The specific molecular-physiological mechanism by which the FTO gene might affect brain function and adiposity remains the object of intense scientific investigation.

Overall, the quantity of studies supporting genetic influence on PA is much lower than emerging human genetic research examining subjective responses to alcohol (Ray & Hutchison, 2004; Ray et al., 2010), caffeine (Alsene et al., 2003; Dlugos et al., 2007; Kendler & Prescott, 1999; Loke, 1988), amphetamine (Hart et al., 2011; Lott et al., 2005; Lott et al., 2006), and nicotine (Hoft et al., 2011; Hutchison et al., 2007; Sherva et al., 2008; Zeiger et al., 2007) consumption.

In sum, there is preliminary evidence supporting affective response to PA as an intermediate phenotype; however, more research is needed. The advantage of the intermediate phenotype when applied to affective response to PA can be best illustrated when considering the interaction between genetic and environmental determinants of PA behavior. Indeed, the depiction in Figure 1 is simplified because it does not account for gene-environment interactions. While the genetic predispositions of modern humans do not differ significantly from recent ancestors (e.g., grandparents and great-grandparents), their environments differ substantially. For example, before modern conveniences like motor vehicles, washing machines, and televisions, a genetic predisposition for a more negative response to PA was less likely to lead to an overall inactive lifestyle because most people had to engage in a significant amount of PA to function in society and in occupational contexts. On the other hand, a predisposition for a negative affective response to PA may be more likely to lead to low rates of PA in modern environments, which allow for sedentary lifestyles. Gene-environment interactions may be even more pronounced when considering variability in socio-economic factors such as unsafe sidewalks, low neighborhood attractiveness, and lack of access to parks, playgrounds, and fitness clubs. In sum, the effects of affective response to PA on PA behavior are moderated by various environmental factors.

The investigation of the genetic underpinnings of affective response to PA has the potential to accelerate optimal translation of discoveries from large-scale genomic data to behavioral intervention development. Genome wide association (GWA) studies examining PA will likely increase (e.g., NIH's precision medicine initiative cohort, "All of Us"), and a large number of PA-related SNPs will be identified soon. However, without a theoretical context, GWA studies reveal little about the mechanisms of genetic effects on behavior that may serve as intervention targets. To realize this translation, we need to understand *how* genetic variation influences PA among humans.

An intermediate phenotype is, relative to the target behavior, a more proximal and malleable variable that can be targeted by the health promotion intervention. For example, if genetic influences on PA behavior are mediated by variation in affective response to PA, this may point towards the need for targeting and tailoring interventions that consider differences in affective response. Specifically, those who respond more negatively to PA due to genomic profile may be encouraged to exercise at a self-paced intensity rather than prescribed

moderate intensity (Williams et al, 2015). While no studies to date have used such an approach, the conceptual plausibility of affective response to PA as an intermediate phenotype for intervention development has gained significant traction among researchers in recent years (Bryan et al., 2017; Yanovski & Yanovski, 2018).

7. Conclusion

With recent advancements in technology for ecological momentary assessment devices (e.g., smartphones), objective assessments of PA (e.g., accelerometers) and the decreasing cost of genotyping, the investigation of the genetic underpinnings of the affective response to PA is timely and feasible. Investigating intermediate phenotypes will allow researchers to move from *predicting* PA based on genomic information to *understanding how* SNPs influence PA behavior, thus ultimately optimizing the translation of knowledge gained from genomic data to intervention development.

Acknowledgments

The authors are thankful for Drs. John McGeary, Jeanne McCaffery, Angela Bryan, Beth Bock, and Shira Dunsiger for their feedback on the earlier version of this paper. Harold Lee was supported by National Institutes of Health, National Heart Lung and Blood Institute (1F31HL140817). Jessica Emerson was supported by National Institutes of Health, National Cancer Institute (5F31CA206245). Lauren Connell was supported by National Institutes of Health, National Heart Lung and Blood Institute (T32 HL076134).

References

- Aaltonen S, Ortega-Alonso A, Kujala UM, & Kaprio J, 2010 A longitudinal study on genetic and environmental influences on leisure time physical activity in the Finnish Twin Cohort. *Twin Research and Human Genetics: the Official Journal of the International Society for Twin Studies*, 13, 475–481. [PubMed: 20874470]
- Alsene K, Deckert J, Sand P, & de Wit H, 2003 Association between A2a receptor gene polymorphisms and caffeine-induced anxiety. *Neuropsychopharmacology*, 28, 1694. [PubMed: 12825092]
- Backhouse SH, Ekkekakis P, Biddle SJ, Foskett A, & Williams C, 2007 Exercise makes people feel better but people are inactive: Paradox or artifact? *Journal of Sport and Exercise Psychology*, 29, 498. [PubMed: 17968050]
- Bentham J, 1789/2007 *An introduction to the principles of morals and legislation* New York: Dover Publication.
- Berger BG, & Motl RW, 2000 Exercise and mood: A selective review and synthesis of research employing the profile of mood states. *Journal of Applied Sport Psychology*, 12, 69–92.
- Bouchard C, & Perusse L, 1994 Heredity, activity level, fitness, and health, in: Stephens T, (Ed.), *Physical activity, fitness, and health: International Proceedings and Consensus Statement*, Human Kinetics Publishers, pp. 106–118.
- Bruneau ML, Johnson BT, Huedo-Medina TB, Larson KA, Ash GI, & Pescatello LS, 2016 The blood pressure response to acute and chronic aerobic exercise: A meta-analysis of candidate gene association studies. *Journal of Science and Medicine in Sport*, 19, 424–431. [PubMed: 26122461]
- Bryan AD, Hutchison KE, Seals DR, & Allen DL, 2007 A transdisciplinary model integrating genetic, physiological, and psychological correlates of voluntary exercise. *Health Psychology*, 26, 30–39. [PubMed: 17209695]
- Bryan AD, Magnan RE, Hooper AE, Ciccolo JT, Marcus B, & Hutchison KE, 2013 Colorado stride (COSTRIDE): testing genetic and physiological moderators of response to an intervention to increase physical activity. *International Journal of Behavioral Nutrition and Physical Activity*, 10, 139.

- Bryan AD, Jakicic JM, Hunter CM, Evans ME, Yanovski SZ, & Epstein LH (2017). Behavioral and psychological phenotyping of physical activity and sedentary behavior: implications for weight management. *Obesity*, 25(10), 1653–1659. [PubMed: 28948719]
- Cai G, Cole SA, Butte N, Bacino C, Diego V, Tan K, et al., 2006 A quantitative trait locus on chromosome 18q for physical activity and dietary intake in Hispanic children. *Obesity*, 14, 1596–1604. [PubMed: 17030971]
- Carlsson S, Andersson T, Lichtenstein P, Michaelsson K, & Ahlbom A, 2006 Genetic effects on physical activity: results from the Swedish Twin Registry. *Medicine Science Sports and Exercise*, 38, 1396–1401.
- Collins FS, & Varmus H, 2015 A new initiative on precision medicine. *New England Journal of Medicine*, 372, 793–795.
- Currie J, Ramsbottom R, Ludlow H, Nevill A, & Gilder M, 2009 Cardio-respiratory fitness, habitual physical activity and serum brain derived neurotrophic factor (BDNF) in men and women. *Neuroscience Letters*, 451, 152–155. [PubMed: 19133315]
- Dlugos A, Freitag C, Hohoff C, McDonald J, Cook EH, Deckert J, et al., 2007 Norepinephrine transporter gene variation modulates acute response to D-amphetamine. *Biological Psychiatry*, 61, 1296–1305. [PubMed: 17239355]
- Dotson VM, Hsu F-C, Langa TY, McDonough CW, King AC, Cohen RA, et al., 2016 Genetic moderators of the impact of physical activity on depressive symptoms. *Journal of Frailty Aging*, 5, 6. [PubMed: 26980363]
- Downs SH, & Black N, 1998 The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *Journal of Epidemiology & Community Health*, 52, 377–384. [PubMed: 9764259]
- Dunton GF, Liao Y, Intille S, Huh J, & Leventhal A, 2015 Momentary assessment of contextual influences on affective response during physical activity. *Health Psychology*, 34, 1145. [PubMed: 26053885]
- Ekkekakis P, Hall EE, & Petruzzello SJ, 2005 Variation and homogeneity in affective responses to physical activity of varying intensities: An alternative perspective on dose–response based on evolutionary considerations. *Journal of Sports Sciences*, 23, 477–500. [PubMed: 16194996]
- Ekkekakis P, Lind E, & Vazou S, 2010 Affective responses to increasing levels of exercise intensity in normal-weight, overweight, and obese middle-aged women. *Obesity*, 18, 79–85. [PubMed: 19556979]
- Ekkekakis P, Vazou S, Bixby W, & Georgiadis E, 2016 The mysterious case of the public health guideline that is (almost) entirely ignored: call for a research agenda on the causes of the extreme avoidance of physical activity in obesity. *Obesity Reviews*
- Eriksson M, Rasmussen F, & Tynelius P, 2006 Genetic factors in physical activity and the equal environment assumption--the Swedish young male twins study. *Behavior Genetics*, 36, 238–247. [PubMed: 16502139]
- Fisher A, van Jaarsveld CH, Llewellyn CH, & Wardle J, 2010 Environmental influences on children's physical activity: quantitative estimates using a twin design. *PLoS One*, 5, e10110. [PubMed: 20422046]
- Franks PW, Ravussin E, Hanson RL, Harper IT, Allison DB, Knowler WC, et al., 2005 Habitual physical activity in children: the role of genes and the environment. *The American Journal of Clinical Nutrition*, 82, 901–908. [PubMed: 16210723]
- Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, et al., 2011 American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Medicine and Science in Sports and Exercise*, 43, 1334–1359.
- Gellert P, Ziegelmann JP, Warner LM, & Schwarzer R, 2011 Physical activity intervention in older adults: does a participating partner make a difference? *European Journal of Ageing*, 8, 211. [PubMed: 28798651]

- Goldman D, & Ducci F, 2007 Deconstruction of vulnerability to complex diseases: enhanced effect sizes and power of intermediate phenotypes. *The Scientific World Journal*, 7, 124–130. [PubMed: 17982585]
- Gong J, Schumacher F, Lim U, Hindorff LA, Haessler J, Buyske S, et al., 2013 Fine mapping and identification of BMI loci in African Americans. *The American Journal of Human Genetics*, 93, 661–671. [PubMed: 24094743]
- Gottesman II, & Gould TD, 2003 The endophenotype concept in psychiatry: etymology and strategic intentions. *American Journal of Psychiatry*
- Gujral S, Manuck SB, Ferrell RE, Flory JD, & Erickson KI, 2014 The BDNF Val66Met polymorphism does not moderate the effect of self-reported physical activity on depressive symptoms in midlife. *Psychiatry Research*, 218, 93–97.
- Gunstad J, Schofield P, Paul RH, Spitznagel MB, Cohen RA, Williams LM, et al., 2006 BDNF Val66Met polymorphism is associated with body mass index in healthy adults. *Neuropsychobiology*, 53, 153–156. [PubMed: 16707914]
- Hall JF, 1976 Classical conditioning and instrumental learning: A contemporary approach: Lippincott Williams & Wilkins.
- Hardy CJ, & Rejeski WJ (1989). Not what, but how one feels: The measurement of affect during exercise. *Journal of Sport and Exercise Psychology*, 11, 304–317.
- Hart AB, de Wit H, & Palmer AA, 2011 Genetic factors modulating the response to stimulant drugs in humans, in: Cryan JF, & Reif A (Eds), *Behavioral Neurogenetics*, Springer, pp. 537–577
- Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, et al., 2007 Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Medicine and Science in Sports and Exercise*, 39, 1423–1434.
- Hassanein MT, Lyon HN, Nguyen TT, Akyzbekova EL, Waters K, Lettre G, et al., 2010 Fine mapping of the association with obesity at the FTO locus in African-derived populations. *Human Molecular Genetics*, 19, 2907–2916. [PubMed: 20430937]
- Hobbes T, 1789/2008 *Leviathan* New York, NY: Simon & Schuster Inc.
- Hofer M, Pagliusi SR, Hohn A, Leibrock J, & Barde Y, 1990 Regional distribution of brain-derived neurotrophic factor mRNA in the adult mouse brain. *The EMBO journal*, 9, 2459. [PubMed: 2369898]
- Hoft NR, Stitzel JA, Hutchison KE, & Ehringer MA, 2011 CHRNA2 promoter region: association with subjective effects to nicotine and gene expression differences. *Genes, Brain and Behavior*, 10, 176–185.
- Hooper AEC, Bryan AD, & Hagger MS, 2014 What keeps a body moving? The brain-derived neurotrophic factor val66met polymorphism and intrinsic motivation to exercise in humans. *Journal of Behavioral Medicine*, 37, 1180–1192. [PubMed: 24805993]
- Hopkins ME, Davis FC, VanTieghem MR, Whalen PJ, & Bucci DJ, 2012 Differential effects of acute and regular physical exercise on cognition and affect. *Neuroscience*, 215, 59–68. [PubMed: 22554780]
- Horstmann DM, 1950 Acute poliomyelitis: relation of physical activity at the time of onset to the course of the disease. *Journal of the American Medical Association*, 142, 236–241. [PubMed: 15400610]
- Hutchison KE, Allen DL, Filbey FM, Jepson C, Lerman C, Benowitz NL, et al., 2007 CHRNA4 and tobacco dependence: from gene regulation to treatment outcome. *Archives of General Psychiatry*, 64, 1078–1086. [PubMed: 17768273]
- Johnson BT, MacDonald HV, Bruneau ML Jr, Goldsby TU, Brown JC, Huedo-Medina TB, et al., 2014 Methodological quality of meta-analyses on the blood pressure response to exercise: A review. *Journal of Hypertension*, 32, 706–723. [PubMed: 24463936]
- Joosen AM, Gielen M, Vlietinck R, & Westerterp KR, 2005 Genetic analysis of physical activity in twins. *The American Journal of Clinical Nutrition*, 82, 1253–1259. [PubMed: 16332658]
- Kahneman D, Wakker PP, & Sarin R, 1997 Back to Bentham? Explorations of experienced utility. *Quarterly Journal of Economics*, 112, 375–405.

- Karoly HC, Stevens CJ, Magnan RE, Harlaar N, Hutchison KE, & Bryan AD, 2012 Genetic influences on physiological and subjective responses to an aerobic exercise session among sedentary adults. *Journal of Cancer Epidemiology*, 2012, 540563. [PubMed: 22899923]
- Kendler KS, & Prescott CA, 1999 Caffeine intake, tolerance, and withdrawal in women: a population-based twin study. *American Journal of Psychiatry*, 156, 223–228.
- Kwan BM, & Bryan A, 2010 In-task and post-task affective response to exercise: Translating exercise intentions into behaviour. *British Journal of Health Psychology*, 15, 115–131. [PubMed: 19397847]
- Lee I-M, Shiroma EJ, Lobelo F, Puska P, Blair SN, Katzmarzyk PT, et al., 2012 Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *The Lancet*, 380, 219–229.
- León-Mimila P, Villamil-Ramírez H, Villalobos-Comparán M, Villarreal-Molina T, Romero-Hidalgo S, López-Contreras B, et al., 2013 Contribution of common genetic variants to obesity and obesity-related traits in mexican children and adults. *PLoS One*, 8, e70640. [PubMed: 23950976]
- Loke WH, 1988 Effects of caffeine on mood and memory. *Physiology & Behavior*, 44, 367–372. [PubMed: 3222359]
- Lott DC, Kim S-J, Cook EH, & De Wit H, 2005 Dopamine transporter gene associated with diminished subjective response to amphetamine. *Neuropsychopharmacology*, 30, 602–609. [PubMed: 15602501]
- Lott DC, Kim SJ, Cook EH, & Wit H, 2006 Serotonin transporter genotype and acute subjective response to amphetamine. *The American Journal on Addictions*, 15, 327–335. [PubMed: 16966188]
- Lox CL, Jackson S, Tuholski SW, Wasley D, & Treasure DC (2000). Revisiting the measurement of exercise-induced feeling states: The Physical Activity Affect Scale (PAAS). *Measurement in Physical Education and Exercise Science*, 4, 79–95.
- Maia JA, Thomis M, & Beunen G, 2002 Genetic factors in physical activity levels: a twin study. *American Journal of Preventive Medicine*, 23, 87–91. [PubMed: 12133742]
- Mill JS, 1861/2012 *Utilitarianism* Lexington, KY, USA: Renaissance Classics.
- Mitchell BD, Rainwater DL, Hsueh WC, Kennedy AJ, Stern MP, & Maccluer JW, 2003 Familial aggregation of nutrient intake and physical activity: results from the San Antonio Family Heart Study. *Annals of Epidemiology*, 13, 128–135. [PubMed: 12559672]
- Mitchell JA, Hakonarson H, Rebbeck TR, & Grant SF, 2013 Obesity-susceptibility loci and the tails of the pediatric BMI distribution. *Obesity*, 21, 1256–1260. [PubMed: 23408508]
- Moher D, Liberati A, Tetzlaff J, & Altman DG, 2009 Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Annals of Internal Medicine*, 151, 264–269. [PubMed: 19622511]
- Murer M, Yan Q, & Raisman-Vozari R, 2001 Brain-derived neurotrophic factor in the control human brain, and in Alzheimer's disease and Parkinson's disease. *Progress in Neurobiology*, 63, 71–124. [PubMed: 11040419]
- Parfitt G, Rose EA, & Burgess WM, 2006 The psychological and physiological responses of sedentary individuals to prescribed and preferred intensity exercise. *British Journal of Health Psychology*, 11, 39–53. [PubMed: 16480554]
- Paulhus DL, 1991 Measurement and control of response bias, in: Robinson JP, Shaver PR & Wrightsman LS (Eds.), *Measures of Personality and Social Psychological Attitudes* Academic Press, San Diego, CA, pp. 17–59.
- Ray LA, & Hutchison KE, 2004 A Polymorphism of the μ -Opioid Receptor Gene (OPRM1) and Sensitivity to the Effects of Alcohol in Humans. *Alcoholism: Clinical and Experimental Research*, 28, 1789–1795.
- Ray LA, Miranda R Jr, Tidey JW, McGeary JE, MacKillop J, Gwaltney CJ, et al., 2010 Polymorphisms of the μ -opioid receptor and dopamine D₄ receptor genes and subjective responses to alcohol in the natural environment. *Journal of Abnormal Psychology*, 119, 115. [PubMed: 20141248]
- Reed J, & Ones DS, 2006 The effect of acute aerobic exercise on positive activated affect: A meta-analysis. *Psychology of Sport and Exercise*, 7, 477–514.

- Rhodes RE, & Kates A, 2015 Can the Affective Response to Exercise Predict Future Motives and Physical Activity Behavior? A Systematic Review of Published Evidence. *Annals of Behavioral Medicine*, 49, 715–731. [PubMed: 25921307]
- Rose EA, & Parfitt G, 2007 A quantitative analysis and qualitative explanation of the individual differences in affective responses to prescribed and self-selected exercise intensities. *Journal of Sport and Exercise Psychology*, 29, 281–309. [PubMed: 17876968]
- Russell JA, 1980 A circumplex model of affect. *Journal of Personality and Social Psychology*, 39, 1161.
- Schneider M, Dunn A, & Cooper D, 2009 Affect, exercise, and physical activity among healthy adolescents. *Journal of Sport and Exercise Psychology*, 31, 706–723. [PubMed: 20384008]
- Schuch FB, Vasconcelos-Moreno MP, & Fleck MP, 2011 The impact of exercise on Quality of Life within exercise and depression trials: A systematic review. *Mental Health and Physical Activity*, 4, 43–48.
- Schutte NM, Nederend I, Hudziak JJ, Bartels M, & de Geus EJ, 2016 Heritability of the affective response to exercise and its correlation to exercise behavior. *Psychology of Sport and Exercise*
- Sherry ST, Ward M-H, Kholodov M, Baker J, Phan L, Smigielski EM, et al., 2001 dbSNP: the NCBI database of genetic variation. *Nucleic Acids Research*, 29, 308–311. [PubMed: 11125122]
- Sherva R, Wilhelmsen K, Pomerleau CS, Chasse SA, Rice JP, Snedecor SM, et al., 2008 Association of a single nucleotide polymorphism in neuronal acetylcholine receptor subunit alpha 5 (CHRNA5) with smoking status and with 'pleasurable buzz' during early experimentation with smoking. *Addiction*, 103, 1544–1552. [PubMed: 18783506]
- Shugart YY, Chen L, Day IN, Lewis SJ, Timpson NJ, Yuan W, et al., 2009 Two British women studies replicated the association between the Val66Met polymorphism in the brain-derived neurotrophic factor (BDNF) and BMI. *European Journal of Human Genetics*, 17, 1050–1055. [PubMed: 19209189]
- Simonen RL, Perusse L, Rankinen T, Rice T, Rao D, & Bouchard C, 2002 Familial aggregation of physical activity levels in the Quebec Family Study. *Medicine and Science in Sports and Exercise*, 34, 1137–1142. [PubMed: 12131254]
- Stubbe JH, Boomsma DI, Vink JM, Cornes BK, Martin NG, Skytthe A, et al., 2006 Genetic influences on exercise participation in 37,051 twin pairs from seven countries. *PLoS One*, 1, e22. [PubMed: 17183649]
- Thorleifsson G, Walters GB, Gudbjartsson DF, Steinthorsdottir V, Sulem P, Helgadóttir A, et al., 2009 Genome-wide association yields new sequence variants at seven loci that associate with measures of obesity. *Nature Genetics*, 41, 18–24. [PubMed: 19079260]
- USDHHS, 2008 Physical Activity Guidelines Advisory Committee Report, 2008 Washington, DC: U.S. Department of Health and Human Services.
- Usener H, 1887/2010 *Epicurea*: Cambridge University Press.
- Van Landuyt LM, Ekkekakis P, Hall EE, & Petruzzello SJ, 2000 Throwing the mountains into the lakes: On the perils of nomothetic conceptions of the exercise-affect relationship. *Journal of Sport and Exercise Psychology*, 22, 208–234.
- Welch AS, Hulley A, Ferguson C, & Beauchamp MR, 2007 Affective responses of inactive women to a maximal incremental exercise test: A test of the dual-mode model. *Psychology of Sport and Exercise*, 8, 401–423.
- Wen W, Cho Y-S, Zheng W, Dorajoo R, Kato N, Qi L, et al., 2012 Meta-analysis identifies common variants associated with body mass index in east Asians. *Nature Genetics*, 44, 307–311. [PubMed: 22344219]
- Wennesland R, Brown E, Hopper J Jr, Hodges J Jr, Guttentag O, Scott K, et al., 1959 Red cell, plasma and blood volume in healthy men measured by radiochromium (Cr51) cell tagging and hematocrit: influence of age, somatotype and habits of physical activity on the variance after regression of volumes to height and weight combined. *Journal of Clinical Investigation*, 38, 1065.
- Williams DM, 2008 Exercise, affect, and adherence: an integrated model and a case for self-paced exercise. *Journal of Sport & Exercise Psychology*, 30, 471. [PubMed: 18971508]

- Williams DM, 2018 Psychological hedonism, hedonic motivation, and health-related behavior, in Williams DM, Rhodes RE, & Conner MT (Eds.), *Affective Determinants of Health Behavior* New York: Oxford University Press.
- Williams DM, Dunsiger S, Ciccolo JT, Lewis BA, Albrecht AE, & Marcus BH, 2008 Acute Affective Response to a Moderate-intensity Exercise Stimulus Predicts Physical Activity Participation 6 and 12 Months Later. *Psychology of Sport and Exercise*, 9, 231–245. [PubMed: 18496608]
- Williams DM, Dunsiger S, Emerson JA, Gwaltney CJ, Monti PM, & Miranda R Jr, 2016 Self-paced exercise, affective response, and exercise adherence: a preliminary investigation using ecological momentary assessment. *Journal of Sport and Exercise Psychology*, 38, 282–291. [PubMed: 27383469]
- Williams DM, Dunsiger S, Jennings EG, & Marcus BH, 2012 Does affective valence during and immediately following a 10-min walk predict concurrent and future physical activity? *Annals of Behavioral Medicine*, 44, 43–51. [PubMed: 22532005]
- Wing MR, Ziegler J, Langefeld CD, Ng MC, Haffner SM, Norris JM, et al., 2009 Analysis of FTO gene variants with measures of obesity and glucose homeostasis in the IRAS Family Study. *Human Genetics*, 125, 615–626. [PubMed: 19322589]
- Wu L, Xi B, Zhang M, Shen Y, Zhao X, Cheng H, et al., 2010 Associations of six single nucleotide polymorphisms in obesity-related genes with BMI and risk of obesity in Chinese children. *Diabetes*, 59, 3085–3089. [PubMed: 20843981]
- Xi B, Cheng H, Shen Y, Chandak GR, Zhao X, Hou D, et al., 2013 Study of 11 BMI-associated loci identified in GWAS for associations with central obesity in the Chinese children. *PLoS One*, 8, e56472. [PubMed: 23424664]
- Yanovski SZ, & Yanovski JA (2018). Toward precision approaches for the prevention and treatment of obesity. *JAMA*, 319(3), 223–224. [PubMed: 29340687]
- Zeiger JS, Haberstick BC, Schlaepfer I, Collins AC, Corley RP, Crowley TJ, et al., 2007 The neuronal nicotinic receptor subunit genes (CHRNA6 and CHRNB3) are associated with subjective responses to tobacco. *Human Molecular Genetics*, 17, 724–734. [PubMed: 18055561]
- Zhao J, Bradfield JP, Li M, Wang K, Zhang H, Kim CE, et al., 2009 The Role of Obesity-associated Loci Identified in Genome-wide Association Studies in the Determination of Pediatric BMI. *Obesity*, 17, 2254–2257. [PubMed: 19478790]
- Zoladz J, Pilc A, Majerczak J, Grandys M, Zapart-Bukowska J, & Duda K, 2008 Endurance training increases plasma brain-derived neurotrophic factor concentration in young healthy men. *Journal of Physiological Pharmacology*, 59, 119–132.
- Zoladz JA, Duda K, & Majerczak J, 1998 Oxygen uptake does not increase linearly at high power outputs during incremental exercise test in humans. *European Journal of Applied Physiology and Occupational Physiology*, 77, 445–451. [PubMed: 9562296]

Figure 1.
Affective Response to Physical Activity as an Intermediate Phenotype

Figure 2.
Flow Diagram of Study Selection for Systematic Review of Research on the Genetic
Influence on Affective Response to Physical Activity Behavior