Published in final edited form as:

Org Lett. 2019 May 03; 21(9): 3346–3351. doi:10.1021/acs.orglett.9b01097.

Deaminative Reductive Arylation Enabled by Nickel/Photoredox Dual Catalysis

Jun Yi^{†,‡,§}, Shorouk O. Badir^{†,§}, Lisa Marie Kammer[†], María Ribagorda[†], and Gary A. Molander^{*,†}

[†]Roy and Diana Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, 231 South 34th Street, Philadelphia, Pennsylvania 19104-6323, United States

[‡]Jiangsu Laboratory of Advanced Functional Materials, School of Chemistry and Materials Engineering, Changshu Institute of Technology, Changshu 215500, China

Abstract

Described is a cross-electrophilic, deaminative coupling strategy harnessing Katritzky salts as a new species of electrophile in Ni/photoredox dual catalytic reductive cross-coupling reactions. Distinguishing features of this arylation protocol include its mild reaction conditions, high chemoselectivity, and adaptability to a variety of complex substrates [i.e., pyridinium salts derived from amines and partners derived from (hetero)aryl bromides].

Graphical Abstract

Aliphatic primary amines are a privileged class of compounds prevalent in natural products, valued synthetic intermediates, and pharmaceutical drugs such as Tamiflu, Linagliptin, Amlodipine, and Sitagliptin.¹ In an elegant report, the Watson group recently demonstrated the use of Katritzky salts, formed *via* a simple condensation of the corresponding amines with a bench-stable, commercially-available pyrylium salt, as alkyl radical precursors in cross-coupling with arylboronic acids.^{2,3} Subsequently, Glorius, Aggarwal, Shi, Gryko, and Liu further demonstrated the utility of these redox-active amines in C—H arylation,⁴ borylation,⁵ alkynylation,⁶ allylation,⁷ and dicarbofunctionalization, respectively.⁸ More recently, Watson and co-workers disclosed a deaminative alkyl-alkyl Negishi cross-coupling with alkylzinc halides.⁹ The growing interest in this area highlights the challenges associated

*Corresponding Author: gmolandr@sas.upenn.edu.

[§]These authors contributed equally to this work.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

The authors declare no competing financial interest.

with C—N bond activation and presents new opportunities to address these limitations, in particular with regard to the development of cross-electrophilic, deaminative strategies to forge C(sp³)-hybridized centers.

In recent years, nickel/photoredox dual catalysis has emerged as a powerful tool to construct C—C bonds *via* a single-electron transmetalation pathway.¹⁰ In this context, the unique characteristics of this reaction manifold favors the formation of tetrahedral carbon centers without the need of harsh nucleophilic organometallic reagents or elevated temperature.¹¹ Given our and other groups' long-standing interest in reductive nickel transformations and our deep understanding of photoredox catalysis,¹³ we envisioned the application of Katritzky salts as a new species of alkyl radical precursors in a reductive, cross-electrophilic coupling with (hetero)aryl bromides – *representing one of the few synthetic methods employing a net reductive photoredox/Ni dual catalytic transformation*.¹⁴

Given the accessible redox potential of Katritzky salts ($E_{1/2} = -0.93$ V vs. SCE),⁷ the organic dye 4CzIPN (reduced photocatalyst $E_{1/2} = -1.21$ V vs. SCE) was examined and proved effective at delivering the desired coupled product (Table 1). Other photocatalysts were also tested (entries 2 and 3 in Table 1). Not surprisingly, [Ir(dtbbpy)(ppy)₂]PF₆ provided quantitative conversion to the desired product, likely because of its strongly reductive nature ($E_{1/2} = -1.51$ V vs SCE).^{4,10b} Considering the low cost, ease of preparation, and excellent reactivity of 4CzIPN,¹⁵ it was chosen as the photocatalyst of choice. Although DMA provided equally good results in this cross-coupling (entry 6 in Table 1), THF was chosen because of its low boiling point. However, it should be noted that the use of DMA for electron-rich substrates proved effective at times (see Supporting Information). As anticipated, control experiments proved that all components of the reaction were necessary for the dual catalytic system to proceed (entries 7–10). Finally, in an attempt to provide an easier reaction set-up, the optimization was carried out using the precomplex NiBr₂•dtbbpy.

With suitable reaction conditions in hand, the scope of this deaminative cross-coupling was examined (Scheme 2). To this end, a wide array of aryl bromides bearing electron-withdrawing substituents at the *para*-position exhibited excellent reactivity (**3–10**, **16**). Electron-rich aryl bromides were also compatible substrates (**11** and **12**). Importantly, aryl bromides bearing an additional handle for further elaboration in Chan-Lam and Suzuki couplings, such as pinacol boronic ester (**17**) and aryl chlorides (**6** and **13**), reacted to afford the corresponding products in good yields.

Given the mildly basic conditions of this protocol, a variety of sensitive functional groups, including lactone **15** and sulfonamide **16**, were compatible structural motifs. Substitution at the *meta*-position (**13**, **15**, **17**, and **18**) and *ortho*-position (**14** and **18**) was explored; whereby efficient cross-coupling events took place. Moreover, electron deficient difluoroaryl bromide **18** and several heteroaryl systems (**19–22**) were successfully employed. Because of their notoriously unstable nature as the corresponding boronic acids (owing to protodeboronation under strongly basic conditions),¹⁶ the analogous substructural partners had provided sluggish reactivity in previous reports.²

Given their importance in pharmaceutical settings,¹⁷ a variety of nitrogen-containing heteroaryl bromides, including pyrimidine (**20** and **26**), quinoline (**23**, **27** and **28**), isoquinoline **24**, pyridine (**25** and **29**), and indole (**30**) electrophiles were effectively incorporated under this reaction manifold. It is worth highlighting that heteroaromatics (**23** and **24**), previously well-behaved under photoinduced deamination C—H arylation conditions,³ reacted in a chemoselective manner. This highlights the complementary advantages of this protocol to existing Minisci transformations.^{4,18}

Next, attention was turned to the scope of the alkylpyridinium salts, where a wide array of functional groups was tolerated. Katritzky salts bearing a free hydroxyl group (**39** and **40**) did not inhibit the reaction, presenting orthogonal reactivity to C—O bond activation previously reported by MacMillan.¹⁹ Other pyridinium salts bearing an ester handle (**41**) and nitrogen heterocyclic structural motifs (**33** and **34**) afforded the desired products in excellent yields. It is worth noting that both cyclic (**31–36**, **39–42**) and acyclic (**37**, **38**, **43**, and **44**) alkylpyridinium salts reacted efficiently under the reaction conditions.

Finally, to demonstrate the applicability of this protocol to the synthesis of bioactive molecules, we prepared the corresponding alkylpyridinium salt from Mexiletine, a voltage-gated sodium channel blocker used as an antiarrhythmic.²⁰ Utilizing two different aryl bromides, success was achieved at delivering the cross-coupled products **43** and **44** in excellent yields. Finally, a complex aryl bromide derived from Loratadine,²¹ used in the treatment of allergies, was compatible under the reaction conditions (**45**).

To highlight the benefits of employing 4CzIPN as an inexpensive organic photocatalyst [$\sim$ \$5 mmol⁻¹], a transformation was successfully performed on gram-scale, whereby the desired heteroaryl coupled product **27** was obtained in 90% yield.

Taking further advantage of the inherent chemoselectivity of this cross-coupling as described earlier, the quinoline scaffold of **27** was further elaborated at the C2 position to obtain **46** and **47** in good yields, utilizing Minisci-type photoinduced deaminative and decarboxylative strategies previously reported by Glorius and Dhar, respectively.^{4,18c} The ease with which a wide array of aryl- and heteroaryl systems can be fashioned showcases the significant potential of deaminative-based C—C bond construction in industrial settings.

To probe the reaction pathway, we conducted a series of mechanistic experiments. In the presence of the radical scavenger TEMPO [(2,2,6,6-tetramethylpiperidin-1-yl)oxyl], the arylation was completely inhibited, with full recovery of the aryl bromide (Scheme 4A). This is suggestive of the involvement of radical species under this metallaphotoredox manifold. This hypothesis was further corroborated by isolation of **48**, a byproduct likely formed *via* a radical-radical coupling of the persistent dihydropyridine radical intermediate **A** with a transient secondary alkyl radical during the course of the reaction⁶ (Scheme 4B).

To gain further understanding of the role of the photocatalyst, fluorescence quenching studies were performed (Figure 1). As anticipated, the Stern-Volmer relationship confirmed that both triethylamine and the Katritzky salt were effectively quenched by the photoexcited 4CzIPN. Although these results could support an operative single-electron-transfer (SET)

reduction of the alkylpyridinium salt by the photocatalyst, we cannot rule out the possibility of a direct SET event from a low-valent nickel intermediate as proposed by Watson and Lei.^{2,14a}

Based on these results and precedented literature reports detailing the reductive cleavage of Katritzky salts,^{2–8} a plausible mechanism is depicted in Scheme 5. First, visible-light irradiation of 4CzIPN generates a potent excited-state oxidant ($E_{1/2} = +1.35$ V vs. SCE),²² which is sufficiently quenched by triethylamine **D** ($E_{1/2} = +1.0$ V vs SCE),^{10b} thus forming the reduced state of the photocatalyst. Upon generation of alkyl radical **IV** from the Katritzky salt ($E_{1/2} = -0.93$ V vs. SCE)⁷ via SET from the photoexcited 4CzIPN or a low-valent nickel intermediate, the radical is intercepted by a ligated Ni(0) complex **V** to afford a Ni(I) intermediate. Subsequently, **VI** undergoes oxidative addition with the (hetero)aryl bromide to afford the corresponding high-valent Ni(III) complex **VII**. Reductive elimination would then occur to yield the desired coupled product **3** and Ni—Br species **VIII**. Reduction of **VIII** by the ground state of the reduced photocatalyst **III** ($E_{1/2} = -1.21$ V vs. SCE) concurrently completes both catalytic cycles.

The developed arylation protocol does exhibit a limitation owing to the inherent stability of the dihydropyridine radical intermediate **A** generated (Scheme 5). Thus, primary alkylpyridinium salts preferentially undergo reoxidation to deliver the corresponding pyridine byproduct.⁶ Considering the sheer number of existing, complementary cross-coupling approaches for primary alkyl radicals developed by our group and others,²² we foresee the current work as an important, complementary approach to the challenges associated with the activation of inert C—N bonds in recently described systems that appear restricted to aryl iodides and use excess metal reductants at elevated temperatures.²³

In conclusion, a cross-electrophilic, reductive deaminative arylation protocol under photoredox conditions has been developed, using triethylamine as the terminal reductant. Eliminating the need for stoichiometric metal reductants, elevated temperatures, and/or strong bases, an exceptional array of functional groups and complex structural scaffolds can be incorporated, including those from bioactive molecules. This arylation protocol is scalable, operationally simple, and utilizes alkylamines as abundant feedstocks.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

The authors are grateful for the financial support provided by the NSF (CHE-1664818) and the NIGMS (R01 GM 113878 to G.M.). J. Y. acknowledges funding from the National Natural Science Foundation of China (Grant No. 21602017), the Natural Science Foundation of Jiangsu Province (Grant No. BK20160405), the University Science Research Project of Jiangsu Province (16KJB150001), and China Scholarship Council (201708320115). M. R. acknowledges support from the Ministerio de Educación, Cultura y Deporte, Subprograma Estatal de Movilidad and Fulbright grant visiting scholar program. We thank Mr. Shuai Zheng (UPenn) for stimulating discussions. We thank Dr. Charles W. Ross, III (UPenn) for his assistance in obtaining HRMS data. Kessil is acknowledged for providing the lights used in this study.

REFERENCES

- (1). (a) McGrath NA; Brichacek M; Njardarson JT A Graphical Journey of Innovative Organic Architectures That Have Improved Our Lives. *J. Chem. Educ* 2010, 87, 1348–1349; (b) Ruiz-Castillo P; Buchwald SL Applications of Palladium-Catalyzed C–N Cross-Coupling Reactions. *Chem. Rev* 2016, 116, 12564–12649; [PubMed: 27689804] (c) Liu Y; Ge H Site-selective C–H arylation of primary aliphatic amines enabled by a catalytic transient directing group. *Nat. Chem* 2017, 9, 26–32.
- (2). Basch CH; Liao J; Xu J; Piane JJ; Watson MP Harnessing Alkyl Amines as Electrophiles for Nickel-Catalyzed Cross Couplings via C–N Bond Activation *J. Am. Chem. Soc* 2017, 139, 5313–5316. [PubMed: 28359153]
- (3). Aryl pyridium salt see: Moser D; Duan Y; Wang F; Ma Y; O'Neill MJ; Cornella J Selective Functionalization of Aminoheterocycles by a Pyrylium Salt *Angew. Chem. Int. Ed* 2018, 57, 11035–11039.
- (4). Klauck FJR; James MJ; Glorius F Deaminative Strategy for the Visible-Light-Mediated Generation of Alkyl Radicals, *Angew. Chem. Int. Ed* 2017, 56, 12336–12339.
- (5). (a) Wu J; He L; Noble A; Aggarwal VK Photoinduced Deaminative Borylation of Alkylamines *J. Am. Chem. Soc* 2018, 140, 10700–10704; [PubMed: 30091912] (b) Sandfort F; Strieth-Kalthoff F; Klauck FJR; James MJ; Glorius F Deaminative Borylation of Aliphatic Amines Enabled by Visible Light Excitation of an Electron Donor–Acceptor Complex *Chem. - Eur. J* 2018, 24, 17210–17214; [PubMed: 30290050] (c) Hu J; Wang G; Li S; Shi Z Selective C–N Borylation of Alkyl Amines Promoted by Lewis Base, *Angew. Chem. Int. Ed* 2018, 57, 15227–15231;
- (6). Ociepa M; Turkowska J; Gryko D Redox-Activated Amines in C(sp³)–C(sp) and C(sp³)–C(sp²) Bond Formation Enabled by Metal-Free Photoredox Catalysis *ACS Catal* 2018, 8, 11362–11367.
- (7). Zhang M-M; Liu F Visible-light-mediated allylation of alkyl radicals with allylic sulfones via a deaminative strategy *Org. Chem. Front* 2018, 5, 3443–3446.
- (8). (a) Klauck FJR; Yoon H; James MJ; Lautens M; Glorius F Visible-Light-Mediated Deaminative Three-Component Dicarbofunctionalization of Styrenes with Benzylic Radicals *ACS Catal* 2019, 9, 236–241; (b) Jiang X; Zhang MM; Xiong W; Lu LQ; Xiao WJ Deaminative (Carbonylative) Alkyl-Heck-type Reactions Enabled by Photocatalytic C–N Bond Activation *Angew. Chem. Int. Ed* 2019, 58, 2402–2406; (c) Wu J; Grant PS; Li X; Noble A; Aggarwal VK Catalyst-Free Deaminative Functionalizations of Primary Amines by Photoinduced Single-Electron Transfer *Angew. Chem. Int. Ed* 2019, 58, DOI:10.1002/anie.201814452.
- (9). Plunkett S; Bash CH; Santana SO; Watson MP Harnessing Alkylpyridinium Salts as Electrophiles in Deaminative Alkyl–Alkyl Cross-Couplings *J. Am. Chem. Soc* 2019, 141, 2257–2262. [PubMed: 30682254]
- (10). For reviews on Ni/photoredox dual catalysis: Tellis JC; Kelly CB; Primer DN; Jouffroy M; Patel NR; Molander GA Single-Electron Transmetalation via Photoredox/Nickel Dual Catalysis: Unlocking a New Paradigm for sp³–sp² Cross-Coupling *Acc. Chem. Res* 2016, 49, 1429–1439; Prier CK; Rankic DA; MacMillan DWC Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis *Chem. Rev* 2013, 113, 5322–5363; [PubMed: 23509883] Matsui JK; Lang SB; Heitz DR; Molander GA Photoredox-Mediated Routes to Radicals: The Value of Catalytic Radical Generation in Synthetic Methods Development *ACS Catal* 2017, 7, 2563–2575. [PubMed: 28413692]
- (11). (a) Tellis JC; Primer DN; Molander GA Dual catalysis. Single-electron transmetalation in organoboron cross-coupling by photoredox/nickel dual catalysis *Science* 2014, 345, 433–436; [PubMed: 24903560] (b) Perry IB; Brewer TF; Sarver PJ; Schultz DM; DiRocco DA; MacMillan DWC Direct arylation of strong aliphatic C–H bonds *Nature* 2018, 560, 70–75; [PubMed: 30068953] (c) Le C; Liang Y; Evans RW; Li X; MacMillan DWC Selective sp³ C–H alkylation via polarity-match-based cross-coupling *Nature* 2017, 547, 79–83. [PubMed: 28636596] (d) Johnston CP; Smith RT; Allmendinger S; MacMillan DWC Metallaphotoredox-catalysed sp³–sp³ crosscoupling of carboxylic acids with alkyl halides *Nature* 2016, 536, 322–325; [PubMed: 27535536] (e) Shaw MH; Shurtleff VW; Terrett JA; Cuthbertson JD; MacMillan DWC Native functionality in triple catalytic cross-coupling: sp³ C–H bonds as latent nucleophiles *Science* 2016, 352, 1304–1308; [PubMed: 27127237] (f) Zuo Z; Ahneman DT; Chu L; Terrett JA; Doyle

- AG; MacMillan DWC Dual catalysis. Merging photoredox with nickel catalysis: coupling of α -carboxyl sp^3 -carbons with aryl halides Science 2014, 345, 437–440. [PubMed: 24903563]
- (12). (a) Molander GA; Traister KM; O'Neill BT Engaging Nonaromatic, Heterocyclic Tosylates in Reductive Cross-Coupling with Aryl and Heteroaryl Bromides J. Org. Chem 2015, 80, 2907–2911; [PubMed: 25711834] (b) Molander GA; Traister KM; O'Neill BT Reductive Cross-Coupling of Nonaromatic, Heterocyclic Bromides with Aryl and Heteroaryl Bromides J. Org. Chem 2014, 79, 5771–5780; [PubMed: 24892751] (c) Molander GA; Wisniewski SR; Traister KM Reductive Cross-Coupling of 3-Bromo-2,1-borazaronaphthalenes with Alkyl Iodides. Org. Lett 2014, 16, 3692–3695; [PubMed: 24977641] (d) Garcia KJ; Gilbert MM; Weix DJ Nickel-Catalyzed Addition of Aryl Bromides to Aldehydes To Form Hindered Secondary Alcohols J. Am. Chem. Soc 2019, 141, 1823–1827; [PubMed: 30693771] (e) Olivares AM; Weix DJ Multimetallic Ni- and Pd-Catalyzed Cross-Electrophile Coupling To Form Highly Substituted 1,3-Dienes J. Am. Chem. Soc 2018, 140, 2446–2449; [PubMed: 29420028] (f) Huang L; Olivares AM; Weix DJ Reductive Decarboxylative Alkynylation of N-Hydroxyphthalimide Esters with Bromoalkynes Angew. Chem. Int. Ed 2017, 56, 11901–11905; (g) Ackerman LKG; Lovell MM; Weix DJ Multimetallic catalysed cross-coupling of aryl bromides with aryl triflates Nature 2015, 524, 454–457; [PubMed: 26280337] (h) Heinz C; Lutz JP; Simmons EM; Miller MM; Ewing WR; Doyle AG Ni-Catalyzed Carbon–Carbon Bond-Forming Reductive Amination J. Am. Chem. Soc 2018, 140, 2292–2300; [PubMed: 29341599] (i) Hofstra JL; Cherney AH; Ordner CM; Reisman SE Synthesis of Enantioenriched Allylic Silanes via Nickel-Catalyzed Reductive Cross-Coupling J. Am. Chem. Soc 2018, 140, 139–142; [PubMed: 29202243] (j) Poremba KE; Kadunce NT; Suzuki N; Cherney AH; Reisman SE Nickel-Catalyzed Asymmetric Reductive Cross-Coupling To Access 1,1-Diaryllkanes J. Am. Chem. Soc 2017, 139, 5684–5687; [PubMed: 28406620] (k) Ye Y; Chen H; Sessler JL; Gong H Zn-Mediated Fragmentation of Tertiary Alkyl Oxalates Enabling Formation of Alkylated and Arylated Quaternary Carbon Centers J. Am. Chem. Soc. 2019, 141, 820–824; [PubMed: 30571912] (l) Wang X; Wang S; Xue W; Gong H Nickel-Catalyzed Reductive Coupling of Aryl Bromides with Tertiary Alkyl Halides J. Am. Chem. Soc 2015, 137, 11562–11565; [PubMed: 26325479] (m) Zhao C; Jia X; Wang X; Gong H Ni-Catalyzed Reductive Coupling of Alkyl Acids with Unactivated Tertiary Alkyl and Glycosyl Halides J. Am. Chem. Soc 2014, 136, 17645–17651; [PubMed: 25415424] (n) García-Domínguez A; Li Z; Nevado C Nickel-Catalyzed Reductive Dicarbofunctionalization of Alkenes J. Am. Chem. Soc 2017, 139, 6835–6838. [PubMed: 28489351]
- (13). Gutierrez O; Tellis JC; Primer DN; Molander GA Nickel-Catalyzed Cross-Coupling of Photoredox-Generated Radicals: Uncovering a General Manifold for Stereoconvergence in Nickel-Catalyzed Cross-Couplings J. Am. Chem. Soc 2015, 137, 4896–4899. [PubMed: 25836634]
- (14). (a) Duan Z; Li W; Lei A Nickel-Catalyzed Reductive Cross-Coupling of Aryl Bromides with Alkyl Bromides: Et₃N as the Terminal Reductant Org. Lett 2016, 18, 4012–4015; [PubMed: 27472556] (b) Peng L; Li Z; Yin G Photochemical Nickel-Catalyzed Reductive Migratory Cross-Coupling of Alkyl Bromides with Aryl Bromides Org. Lett 2018, 20, 1880–1883; [PubMed: 29561162] (c) Paul A; Smith MD; Vannucci AK Photoredox-Assisted Reductive Cross-Coupling: Mechanistic Insight into Catalytic Aryl–Alkyl Cross-Couplings J. Org. Chem 2017, 82, 1996–2003. [PubMed: 28112920]
- (15). Luo J; Zhang J Donor–Acceptor Fluorophores for Visible-Light-Promoted Organic Synthesis: Photoredox/Ni Dual Catalytic C(sp³)–C(sp²) Cross-Coupling ACS Catal, 2016, 6, 873–877.
- (16). Chen L; Sanchez DR; Zhang B; Carrow BP “Cationic” Suzuki–Miyaura Coupling with Acutely Base-Sensitive Boronic Acids J. Am. Chem. Soc 2017, 139, 12418–12421. [PubMed: 28862445]
- (17). Welsch ME; Snyder SA; Stockwell BR Privileged Scaffolds for Library Design and Drug Discovery Curr. Opin. Chem. Biol 2010, 14, 347–361. [PubMed: 20303320]
- (18). (a) Kammer LM; Rahman A; Opatz T A Visible Light-Driven Minisci-Type Reaction with N-Hydroxyphthalimide Esters Molecules 2018, 23, 764; (b) Matsui JK; Primer DN; Molander GA Metal-free C–H alkylation of heteroarenes with alkyltrifluoroborates: a general protocol for 1°, 2° and 3° alkylation Chem. Sci 2017, 8, 3512–3522; [PubMed: 28507725] (c) Sherwood TC; Li N; Yazdani AN; Murali Dhar TG Organocatalyzed, Visible-Light Photoredox-Mediated, One-Pot Minisci Reaction Using Carboxylic Acids via N-(Acylxy)phthalimides J. Org. Chem 2018, 83, 3000–3012; [PubMed: 29420898] (d) Cheng W-M; Shang R; Fu Y Photoredox/Brønsted Acid

Co-Catalysis Enabling Decarboxylative Coupling of Amino Acid and Peptide Redox-Active Esters with NHeteroarenes ACS Catal 2017, 7, 907–911.

- (19). Terrett JA; Cuthbertson JD; Shurleff VW; MacMillan DWC. Switching on elusive organometallic mechanisms with photoredox catalysis Nature 2015, 524, 330–334. [PubMed: 26266976]
- (20). Koszelewski D; Pressnitz D; Clay D; Kroutil W Deracemization of Mexiletine Biocatalyzed by ω -Transaminases Org. Lett 2009, 11, 4810–4812. [PubMed: 19785441]
- (21). Geha RS; Meltzer EO Desloratadine: A new, nonsedating, oral antihistamine J. Allergy Clin. Immunol 2001, 107, 751–76. [PubMed: 11295678]
- (22). (a)Jouffroy M; Primer DN; Molander GA Base-Free Photoredox/Nickel Dual-Catalytic Cross-Coupling of Ammonium Alkylsilicates. J. Am. Chem. Soc 2016, 138, 475–478; [PubMed: 26704168] (b)Jana R; Pathak TP; Sigman MS Advances in Transition Metal (Pd,Ni,Fe)-Catalyzed Cross-Coupling Reactions Using Alkyl-organometallics as Reaction Partners Chem. Rev 2011, 111, 1417–1492.
- (23). (a)Li Chunxiao; Han Jianlin; Mao Yu; Pan, Yi Ni-Catalyzed Deamination Cross-Electrophile Coupling of Katritzky Salts with Halides via C–N Bond Activation ChemRxiv. Preprint DOI: 10.26434/chemrxiv.7638164.v1;(b)Yue H; Zhu C; Shen L; Geng Q; Hock KJ; Yuan T; Cavallo L; Rueping M Nickel-catalyzed C–N bond activation: activated primary amines as alkylating reagents in reductive cross-coupling Chem. Sci, 2019, Advance Article. DOI: 10.1039/c9sc00783k.

A. Alkyl Feedstocks Utilized in Reductive Nickel Transformations: **B. Katritzky Pyridinium Salts in Nickel-Catalyzed Arylation:** **C. This work: Reductive Nickel/Photoredox Deaminative Arylation with ArBr:**

÷ Commodity chemical
÷ Air stable reductant
÷ Mild, basic conditions

÷ Cheap & metal-free [PC]

Scheme 1.
Aliphatic Amines as Radical Precursors in Cross-Coupling

Scheme 2. Scope of (Hetero)Aryl Bromides and Alkylpyridinium Salts^a^aAs in Table 1 (entry 1), 0.5 mmol scale. ^bUsing DMA as a solvent, 48 h.

Scheme 3.
Sequential Gram-Scale Deaminative Cross-Coupling and Minisci C-H Arylation

Scheme 4.

Support for Involvement of Radical Intermediates

^aAs in Table 1 (entry 1), 0.3 mmol scale. ^bNMR yield was calculated based on the alkylpyridinium reagent (theoretical yield of 0.225 mmol).

Scheme 5.
Reductive Nickel/Photoredox Dual Catalysis: Mechanistic Rationale

Figure 1.
Stern-Volmer Fluorescence Quenching Studies of 4CzIPN ($c = 2.0$ M) in THF

Table 1.

Optimization of the Reaction Conditions^a

entry	deviation from standard conditions	3 (%) ^b
1	None	93 (86)
2	[Ir{dF(CF ₃) ₂ ppy}(dtbbpy)]PF ₆	87
3	[Ir(dtbbpy)(ppy) ₂]PF ₆	99
4	3.0 equiv Hantzsch ester	33
5	CH ₃ CN	73
6	DMA	92
7	No light (dark)	0
8	No photocatalyst	0
9	No Ni catalyst	0
10	No amine	0

^a 1 (0.45 mmol), 2 (0.3 mmol), 4CzIPN (3 mol %), NiBr₂(dtbbpy) (5 mol %) in dry, degassed solvent (3.0 mL, 0.1 M) under blue LED irradiation for 24 h. ^b GCMS yield using 1,3,5-trimethoxybenzene as internal standard. Isolated yield in parenthesis.