Cochrane
Library

Cochrane Database of Systematic Reviews

Interventions for treating oro-antral communications and fistulae due to dental procedures (Review)

Kiran Kumar Krishanappa S, Eachempati P, Kumbargere Nagraj S, Shetty NY, Moe S, Aggarwal H, Mathew RJ

Kiran Kumar Krishanappa S, Eachempati P, Kumbargere Nagraj S, Shetty NY, Moe S, Aggarwal H, Mathew RJ.
Interventions for treating oro-antral communications and fistulae due to dental procedures.
Cochrane Database of Systematic Reviews 2018, Issue 8. Art. No.: CD011784.
DOI: [10.1002/14651858.CD011784.pub3](https://doi.org/10.1002/14651858.CD011784.pub3).

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	3
BACKGROUND	5
OBJECTIVES	6
METHODS	6
RESULTS	9
Figure 1.	10
Figure 2.	12
Figure 3.	13
DISCUSSION	13
AUTHORS' CONCLUSIONS	14
ACKNOWLEDGEMENTS	14
REFERENCES	16
CHARACTERISTICS OF STUDIES	18
DATA AND ANALYSES	21
Analysis 1.1. Comparison 1 PBFPF versus BF, Outcome 1 Successful (complete) closure of the oro-antral communication.	22
ADDITIONAL TABLES	22
APPENDICES	23
WHAT'S NEW	24
CONTRIBUTIONS OF AUTHORS	25
DECLARATIONS OF INTEREST	25
SOURCES OF SUPPORT	25
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	26
INDEX TERMS	26

[Intervention Review]

Interventions for treating oro-antral communications and fistulae due to dental procedures

Salian Kiran Kumar Krishanappa¹, Prashanti Eachempati¹, Sumanth Kumbargere Nagraj², Naresh Yedthare Shetty³, Soe Moe⁴, Himanshi Aggarwal⁵, Rebecca J Mathew⁶

¹Department of Prosthodontics, Faculty of Dentistry, Melaka-Manipal Medical College, Manipal Academy of Higher Education (MAHE), Melaka, Malaysia. ²Department of Oral Medicine and Oral Radiology, Faculty of Dentistry, Melaka-Manipal Medical College, Manipal Academy of Higher Education (MAHE), Manipal, Melaka, Malaysia. ³School of Dentistry Oral & Maxillofacial Surgery, International Medical University - IMU, Bukit Jalil, Malaysia. ⁴Community Medicine, Melaka Manipal Medical College, Melaka, Malaysia. ⁵Department of Prosthodontics, Saraswati Dental College and Hospital, Lucknow, India. ⁶South Asian Cochrane Network & Center, Prof. BV Moses Center for Evidence-Informed Health Care and Health Policy, Christian Medical College, Vellore, India

Contact address: Salian Kiran Kumar Krishanappa, Department of Prosthodontics, Faculty of Dentistry, Melaka-Manipal Medical College, Manipal Academy of Higher Education (MAHE), Jalan Batu Hampar, Melaka, 75150, Malaysia. kiran.kumar@manipal.edu.my, kiru.82@gmail.com.

Editorial group: Cochrane Oral Health Group.

Publication status and date: New search for studies and content updated (no change to conclusions), published in Issue 8, 2018.

Citation: Kiran Kumar Krishanappa S, Eachempati P, Kumbargere Nagraj S, Shetty NY, Moe S, Aggarwal H, Mathew RJ. Interventions for treating oro-antral communications and fistulae due to dental procedures. *Cochrane Database of Systematic Reviews* 2018, Issue 8. Art. No.: CD011784. DOI: [10.1002/14651858.CD011784.pub3](https://doi.org/10.1002/14651858.CD011784.pub3).

Copyright © 2018 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

An oro-antral communication is an unnatural opening between the oral cavity and maxillary sinus. When it fails to close spontaneously, it remains patent and is epithelialized to develop into an oro-antral fistula. Various surgical and non-surgical techniques have been used for treating the condition. Surgical procedures include flaps, grafts and other techniques like re-implantation of third molars. Non-surgical techniques include allogenic materials and xenografts. This is an update of a review first published in May 2016.

Objectives

To assess the effectiveness and safety of various interventions for the treatment of oro-antral communications and fistulae due to dental procedures.

Search methods

Cochrane Oral Health's Information Specialist searched the following databases: Cochrane Oral Health's Trials Register (to 23 May 2018), the Cochrane Central Register of Controlled Trials (CENTRAL) (the Cochrane Library, 2018, Issue 4), MEDLINE Ovid (1946 to 23 May 2018), and Embase Ovid (1980 to 23 May 2018). The US National Institutes of Health Trials Registry (ClinicalTrials.gov) and the World Health Organization International Clinical Trials Registry Platform were searched for ongoing trials. No restrictions were placed on the language or date of publication when searching the electronic databases. We also searched the reference lists of included and excluded trials for any randomised controlled trials (RCTs).

Selection criteria

We included RCTs evaluating any intervention for treating oro-antral communications or oro-antral fistulae due to dental procedures. We excluded quasi-RCTs and cross-over trials. We excluded studies on participants who had oro-antral communications, fistulae or both related to Caldwell-Luc procedure or surgical excision of tumours.

Data collection and analysis

Two review authors independently selected trials. Two review authors assessed trial risk of bias and extracted data independently. We estimated risk ratios (RR) for dichotomous data, with 95% confidence intervals (CI). We assessed the overall quality of the evidence using the GRADE approach.

Main results

We included only one study in this review, which compared two surgical interventions: pedicled buccal fat pad flap and buccal flap for the treatment of oro-antral communications. The study involved 20 participants. The risk of bias was unclear. The relevant outcome reported in this trial was successful (complete) closure of oro-antral communication.

The quality of the evidence for the primary outcome was very low. The study did not find evidence of a difference between interventions for the successful (complete) closure of an oro-antral communication (RR 1.00, 95% CI 0.83 to 1.20) one month after the surgery. All oro-antral communications in both groups were successfully closed so there were no adverse effects due to treatment failure.

We did not find trials evaluating any other intervention for treating oro-antral communications or fistulae due to dental procedures.

Authors' conclusions

We found very low quality evidence from a single small study that compared pedicled buccal fat pad and buccal flap. The evidence was insufficient to judge whether there is a difference in the effectiveness of these interventions as all oro-antral communications in the study were successfully closed by one month after surgery. Large, well-conducted RCTs investigating different interventions for the treatment of oro-antral communications and fistulae caused by dental procedures are needed to inform clinical practice.

PLAIN LANGUAGE SUMMARY

Treatment of communications between the oral cavity and the maxillary sinus due to dental procedures

Review question

What evidence is available for the safe and effective treatment of openings between the mouth and main sinus caused by dental procedures?

Background

The floor of the main sinus near the nose is thin and lies directly above the roots of the teeth at the back of the mouth. Sometimes following infection or dental treatment, this structure becomes damaged and openings or channels between the mouth and the sinus are formed. These are known as oro-antral communications (OAC). If the OAC is left open (then described as an oro-antral fistula (OAF), it may become permanent, leading to long-lasting sinus infections. This condition can be treated surgically by using flaps, grafts and other techniques; or non-surgically using a variety of methods and materials. There is little evidence for the most effective and safe treatments for closing OACs and OAFs and clinicians who treat these conditions have identified an urgent need for this. This is an update of a review first published in May 2016.

Study characteristics

We searched various databases until 23 May 2018. Only one study, which was conducted in Iran, is included in our review. The study ran for two years and involved 20 people with OAC aged between 25 and 56 years. Participants were divided into two groups and two surgical treatments were compared for treating oro-antral communications; one group was treated with pedicled buccal fat pad flap (PBFPP) and the other with buccal flap (BF).

Key results and quality of evidence

The study did not find evidence of a difference between PBFPP and BF in terms of successful (complete) closure of OAC. Both interventions resulted in successful closure by one month after surgery. The study did not therefore report any adverse effects of treatment failure. It may not be possible to generalise these findings because the quality of the evidence was very low, due to unclear risk of bias and the small numbers studied in the single included trial.

Conclusion

The evidence currently available is insufficient to draw reliable conclusions regarding the effects of interventions used to treat OAC or fistulae due to dental procedures. More well-designed and well-reported trials evaluating different interventions are needed to provide reliable evidence to inform clinical decisions.

SUMMARY OF FINDINGS

Summary of findings for the main comparison. Pedicled buccal fat pad flap compared to buccal flap for treatment of oro-antral communications and fistulae due to dental procedures

Pedicled buccal fat pad flap compared to buccal flap for treatment of oro-antral communications and fistulae due to dental procedures

Patient or population: people with oro-antral communication/fistulae

Settings: hospital

Intervention: pedicled buccal fat pad flap

Comparison: buccal flap

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Buccal flap	Pedicled buccal fat pad flap				
Successful (complete) closure of oro-antral communication			RR 1.00 (0.83 to 1.20)	20 (1 study ¹)	⊕⊕⊕⊕ very low ^{2,3}	Both groups showed 100% improvement in the closure of OAC after one month. Hence RR is one and the risk difference (absolute effect) is zero.
Adverse effects of treatment failure (such as graft necrosis and rejection or chronic sinusitis)						None reported as there were no treatment failures.

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR:** Risk ratio

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ [Nezafati 2012](#)

- ² Results are imprecise as the single included study has relatively few participants and few events and thus have wide confidence intervals around the estimate of the effect. Hence downgraded by two levels for imprecision.
- ³ Downgraded by one level for risk of bias due to unclear allocation concealment, selective reporting and other bias.

BACKGROUND

Description of the condition

Oro-antral communications (OAC) are pathological conditions characterised by the existence of an unnatural opening ('communication') between the oral cavity and maxillary sinus due to loss of soft and hard tissues that normally separate these compartments. The term 'oro-antral communications' has been used synonymously with the terms 'oro-antral perforation', 'antro-oral communication', 'oro-antral fistula', 'oro-sinus communication' (Visscher 2010), and 'antro-alveolar fistula' (Eneroth 1961). Although the above-mentioned terms are often used synonymously, there is a difference between an oro-antral communication and an oro-antral fistula: only when the communication becomes epithelialized and remains patent is it referred to as oro-antral fistula (OAF) (Batra 2010; Sandhya 2013). OAC is most commonly encountered during maxillary posterior teeth extraction due to the anatomical proximity between root apices and the maxillary antrum (Logan 2003; Nezafati 2012). Its frequency ranges between 0.31% and 4.7% following the extraction of upper teeth (Gacic 2009). It can also occur as a result of iatrogenic complications while performing dental procedures such as surgical removal of cysts (Abuabara 2006; Borgonovo 2012; Dym 2012). It can also be caused as a complication due to infection of the antral filling used to stabilise zygomatic complex fracture (Goodger 2004). Various pathological lesions of the maxillary sinus like mucormycosis (Nilesh 2018), periodontal infections and trauma can also result in the formation of an OAC (Franco-Carro 2011).

Signs and symptoms

Symptoms have been classified based on whether the OAC is acute or chronic (OAF) (Malik 2008).

Acute OAC

- Epistaxis
- Escape of fluid from mouth to nose
- Excruciating pain in and around the region of affected sinus
- Escape of air from mouth to nose on sucking, inhaling or puffing the cheeks
- Enhanced column of air causing alteration in vocal resonance and subsequently change in the voice

Chronic OAC

- Negligible pain as the fistula becomes established and allows the free escape of fluids
- Development of an antral polyp seen as a bluish red lump extruding through the fistula
- Postnasal drip accompanied by unpleasant taste, nocturnal cough, hoarseness of voice, ear ache or catarrhal deafness
- Persistent mucopurulent, foul, unilateral nasal discharge from the affected nostril especially when head is lowered

Complications

Chronic communication between oral cavity and maxillary sinus can act as a pathway for further bacterial and fungal penetration (Borgonovo 2012). Sinusitis has been reported to occur in 60% of cases on the fourth day after sinus exposure (Watzak 2005). Long-standing OAF can cause a general systemic toxæmic condition

leading to fever, malaise, morning anorexia, frontal and parietal headache, anosmia and cacosmia (Malik 2008).

Description of the intervention

Clinical decision-making about how to treat an OAC/OAF depends on multiple factors that include the size of the communication, time of diagnosis and presence of infection. Furthermore, the selection of treatment strategy is influenced by the amount and condition of tissue available for repair and the possible placement of dental implants in the future (Visscher 2010; Dym 2012).

Communications of 1 to 2 mm diameter heal spontaneously by the formation of blood clot in the absence of any infection (Liversedge 2002). Interventions for closure of the OACs can be broadly categorised into surgical, non-surgical and pharmacological interventions.

Surgical interventions

Surgical interventions have been further divided into flaps, grafts and other techniques (Zide 1992; Kitagawa 2003; Visscher 2010; Borja 2011; Visscher 2011; Saleh 2013; Blythe 2016; Demetoglu 2018).

- **Soft tissue flaps:** some of the traditional methods include buccal advancement flaps, palatal rotational flaps, palatal transposition flaps and tongue flaps. Other techniques include local flaps such as a combination of buccal and palatal flap, pedicled buccal fat pad flap, Bichat's fat pad graft and acellular dermal graft.
- **Grafts:** autogenous grafts from chin, retromolar area, zygoma, iliac crest, interseptal and inter-radicular areas, plasma-rich fibrin membrane, cryoplatelet gel and septal cartilage have been advocated to close OAC. Xenografts (with flap closure) such as lyophilised porcine dermis, porcine collagen membrane, bovine bone and guided tissue regeneration (GTR) using bovine barrier membranes have also been used. Allogeneous grafts such as lyophilised fibrin glue and GTR using allogeneous barrier membranes have been reported for treating OAC.
- **Other techniques:** re-implantation of third molar, gingival suturing, metal plates, foils and polymethylmethacrylate plates by approximation of buccal and palatal flap, hydroxylapatite blocks and haemostatic gauze have been tried.

Non-surgical interventions

Non-surgical interventions (Grzesiak-Janias 2001; Thoma 2006; Gacic 2009; Visscher 2010; Buric 2012).

- **Allogeneous materials** (without flap closure) such as fibrin glue, dura; synthetic bone graft materials such as polylactic acid/glycolic acid (PLGA)-coated porous beta tri-calcium phosphate, prolamine occlusion gel and absorbable polyglactin/polydioxanone implant are some of the non-surgical interventions used to manage OAC.
- **Xenografts** (without flap closure) such as porcine dermis and collagen.
- **Other methods** such as acrylic splints, laser light, root analogues and N-butyl cyanoacrylate gel have been tried. Biostimulation with laser light has also been used for closure of OAC (Grzesiak-Janias 2001).

Pharmacological interventions

Used as an adjuvant to surgical and non-surgical interventions. The most commonly used drugs include antibiotics and nasal decongestants.

- **Antibiotics:** a combination of antibiotics such as amoxicillin and clavulanate potassium 875 mg, clindamycin 300 mg 4 times daily, or moxifloxacin 400 mg) have been used in treatment of OAC. (Dym 2012).
- **Nasal decongestants:** can be used as adjuvants to healing of OAC/OAFs if the patient has any sinus infection (Dym 2012).

How the intervention might work

Surgical interventions

Surgical interventions are mostly based on mobilising the tissue and advancing the resultant flap into the defect (Batra 2010).

- **Soft tissue flaps:** a small OAC can be closed immediately by suturing the gingiva, but when this does not provide adequate closure, a soft tissue flap is indicated (Visscher 2010; Dym 2012). In the case of fully developed fistulae, the epithelium lining must be removed in order to facilitate healing (Moore 1991). A buccal advancement flap can be used in small OACs, when the alveolar ridge is very resorbed and the location of the fistula is more mesial (Visscher 2010). The palatal rotational flaps can be used in OAC larger than 1 cm in diameter (Visscher 2010). A modified palatal flap has been proposed that involves only the mucous membrane, leaving the submucosa and periosteum intact to reduce the complication (denudation of bone) of a palatal rotational flap (Kale 2010).
- **Grafts:** these are recommended for the closure of chronic OAF when soft tissue flap closure fails or when augmentation of the alveolar ridge in conjunction with closure is desired (Waldrop 1993). The use of autogenous, allogeneous or xenografts helps to correct the residual bone defects during closure of OACs (Scala 2007). Foils and plates form a mechanical barrier encouraging growth of healthy tissue for the closure of OAC (Steiner 2008). Plasma-rich fibrin membrane is a natural fibrin-based biomaterial which stimulates tissue regeneration and promotes cell migration in the site of interest (Assad 2017).
- **Other techniques:** cryoplatelet gel and GTR accelerate tissue healing and also promote bone reconstruction by release of osteo-inductive growth factors in cases of large OAC (Waldrop 1993).

Non-surgical interventions

Non-surgical interventions promote closure of OAC without the need for a soft tissue flap (Grzesiak-Janias 2001; Buric 2013). These interventions involve minimum tissue handling, hence reducing post-surgical trauma during healing (Choi 2006; Buric 2012; Buric 2013).

- **Allogeneous materials:** glues, adhesives and sealants have the structural ability to enhance the coagulation process and to create a mechanical barrier at the site of tissue breakdown that aids the closure of OAC (Buric 2013). Synthetic absorbable implants are press fitted directly into the defect to obtain the direct closure of the OAC (Buric 2012).
- **Xenografts:** prolamine occlusion gel is directly injected into the perforation, which hardens to form a barrier (Visscher 2010).

- **Other methods:** acrylic splints act as mechanical barriers in people who are immunocompromised to facilitate healing of OAC. Splints may also be appropriate in cases of large defects that do not respond to other treatment modalities.

Pharmacological interventions

- **Antibiotics:** these are needed to control infections of the sinus thereby helping with better healing of the oro-antral communication (Von Wowern 1982).
- **Nasal decongestants, sprays (steroidal and non-steroidal) or a combination:** should be used preoperatively to reduce the inflammation of the sinus mucosa thereby aiding a tension-free closure of soft tissue flap over intact bone (Kamadajja 2008; Borgonovo 2012).

Why it is important to do this review

Cochrane Oral Health undertook an extensive prioritisation exercise in 2014 to identify a core portfolio of titles that were the most clinically important ones to maintain on *The Cochrane Library* (Worthington 2015); this review was identified as a priority title by an oral and maxillofacial surgery expert panel (Cochrane OHG priority review portfolio).

There are many techniques suggested for the treatment of OAC/OAF encountered during dental procedures. Until our review in 2016, there had been no systematic review to summarise the effects of various interventions available to treat OAC/OAF and provide evidence to guide dental practice. Considering the different complex interventions available to treat OAC/OAF in dental patients, it is important to identify the best intervention strategies to help clinicians to treat people with OAC/OAF efficiently and improve patient comfort. We hoped our systematic review would influence the implementation of different approaches and trigger the development of new interventions based on current best evidence, as well as direct further research on interventions with questionable effectiveness and unclear consequences.

OBJECTIVES

To assess the effectiveness and safety of various interventions for the treatment of oro-antral communications and fistulae caused by dental procedures.

METHODS

Criteria for considering studies for this review

Types of studies

We included randomised controlled trials (RCTs) evaluating any intervention for treating oro-antral communications and oro-antral fistulae due to dental procedures. We excluded quasi-RCTs and cross-over trials.

Types of participants

Inclusion criteria

People of any age with oro-antral communications and fistulae caused by dental procedures.

Exclusion criteria

People with oro-antral communications or fistulae created due to Caldwell-Luc procedure (fenestration of the anterior wall of the maxillary sinus and the surgical drainage of this sinus into the nose via an antrostomy) or surgical excision of tumours.

Types of interventions

Any surgical or non-surgical technique or material used for the management of oro-antral communications or oro-antral fistulae due to dental procedures. We intended to make the following comparisons.

- Surgical technique A versus B
- Non-surgical technique A versus B
- Surgical technique versus non-surgical technique
- Surgical technique versus no treatment
- Non-surgical technique versus no treatment

Types of outcome measures

Primary outcomes

- Complete closure of OACs, OAFs, or both, as assessed clinically and with participant-reported outcomes, by absence of associated signs and symptoms such as nasal regurgitation, pain, inflammation, etc.

Secondary outcomes

- Adverse effects, such as graft necrosis and rejection, chronic sinusitis due to failure of OACs, OAFs, or both.
- Time required for healing.

Search methods for identification of studies

Electronic searches

Cochrane Oral Health's Information Specialist conducted systematic searches in the following databases for randomised controlled trials and controlled clinical trials. There were no language, publication year or publication status restrictions:

- Cochrane Oral Health's Trials Register (searched 23 May 2018) ([Appendix 1](#));
- Cochrane Central Register of Controlled Trials (CENTRAL; 2018, Issue 4) in the Cochrane Library (searched 23 May 2018) ([Appendix 2](#));
- MEDLINE Ovid (1946 to 23 May 2018) ([Appendix 3](#));
- Embase Ovid (1980 to 23 May 2018) ([Appendix 4](#)).

Subject strategies were modelled on the search strategy designed for MEDLINE Ovid. Where appropriate, they were combined with subject strategy adaptations of the highly sensitive search strategy designed by Cochrane for identifying randomised controlled trials and controlled clinical trials as described in the *Cochrane Handbook for Systematic Reviews of Interventions* Chapter 6 ([Lefebvre 2011](#)).

Searching other resources

The following trial registries were searched for ongoing studies:

- US National Institutes of Health Ongoing Trials Register ClinicalTrials.gov ([clinicaltrials.gov](#); searched 23 May 2018) ([Appendix 5](#));

- World Health Organization International Clinical Trials Registry Platform ([apps.who.int/trialsearch](#); searched 23 May 2018) ([Appendix 6](#)).

We searched the reference lists of included and excluded trials for any RCTs. We did not perform a separate search for adverse effects of interventions used, we considered adverse effects described in included studies only.

Data collection and analysis

Selection of studies

Two review authors (HA, RJM) independently screened the titles and abstracts from the electronic searches to identify potentially eligible studies that required further evaluation to determine whether they met the inclusion criteria for this review. The search was designed to be sensitive and include controlled clinical trials, these were filtered out early in the selection process if they were not randomised. We obtained full-text copies of all eligible and potentially eligible studies. Two review authors (SK, SN) evaluated these to identify the studies that actually met all the inclusion criteria. From this group, we intended to record the studies that did not meet the inclusion criteria in the 'Characteristics of excluded studies' table, noting the reasons for exclusion. However, we had no studies in this category. We resolved disagreements by discussion. When resolution was not possible, we consulted an arbiter (KNS). We assessed articles in languages other than English by their abstracts, where possible, and if they appeared to be potentially eligible, we intended to translate the full text of the article. However, we did not need to do any translations.

Data extraction and management

Two review authors (SK, SN) independently extracted the data. The review authors were not blinded to the authors of the included study. We resolved disagreements by discussion and when necessary, we consulted a third review author (EP) in order to reach consensus. We extracted data using a customised data extraction form, which was first pilot tested using a sample of the included study. All the items in the data extraction form were designed following guidance from the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011](#)). We entered study details in Review Manager 5 ([RevMan 2014](#)).

We recorded the following details.

- Publication details such as year of publication, language
- Demographic details of the report
- Inclusion and exclusion criteria
- Type of trial, sample size, method of randomisation, allocation concealment, blinding, method of assessing the outcomes and drop-outs
- Type of intervention
- Details of the outcomes reported
- Duration of follow-up
- Results of the intervention
- Funding details

We emailed to the author of the included study where clarification of details or any additional data were required.

Assessment of risk of bias in included studies

We independently assessed the risk of bias in the included trial for seven domains: sequence generation; allocation concealment; performance bias; detection bias; incomplete outcome data; selective outcome reporting; and other biases. For each of these components, we assigned a judgment regarding the risk of bias as either 'high', 'low' or 'unclear', based on guidance in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011).

We contacted the trial authors if details were missing in the publication or were unclear. We resolved disagreements through consensus. We recorded our judgements and justifications in 'Risk of bias' tables for each included study and generated a 'Risk of bias' summary figure. We used these judgements while grading the overall quality of evidence for outcomes in the 'Summary of findings' table.

We summarised the risk of bias according to Higgins 2011, as follows:

Risk of bias	Interpretation	In outcome	In included studies
Low risk of bias	Plausible bias unlikely to seriously alter the results	Low risk of bias for all key domains	Most information is from studies at low risk of bias
Unclear risk of bias	Plausible bias that raises some doubt about the results	Unclear risk of bias for ≥ 1 key domains	Most information is from studies at low or unclear risk of bias
High risk of bias	Plausible bias that seriously weakens confidence in the results	High risk of bias for ≥ 1 key domains	The proportion of information from studies at high risk of bias is sufficient to affect the interpretation of results

Measures of treatment effect

Data for complete closure of OAC was dichotomous. We expressed this effect estimate as a risk ratio (RR) together with 95% confidence intervals (CI). If continuous data had been present, we would have expressed it as mean and standard deviation.

We planned to use standardised mean difference (SMD) if studies used different scales to measure the same outcome. If data expressed were in ordinal scales, we planned to explore the possibility of converting them to dichotomous outcomes. If outcomes were reported both at baseline and at follow-up or at trial endpoints, we intended to extract both the mean change from baseline and the standard deviation of this mean for each treatment group, as well as the same for endpoint data. We had planned to prefer end scores. However, as there was only one included study we did not encounter any of the above.

Unit of analysis issues

We did not find any cluster-randomised trials or split-mouth designs for this condition. In case of a trial with multiple treatment arms, we planned to combine groups to create a single pair-wise comparison as recommended in Higgins 2011. However, we did not find any trial with multiple treatment groups.

If the search found trials with repeated observations on participants, we would have followed the method as described in Section 9.3.4 of the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). In case of multiple treatment attempts per participant, we planned to use the number of participants randomised to calculate the CIs (Higgins 2011). In trials where adverse effects were described as counts, we intended to follow the method described in Section 9.2.5 of the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). However, we did not encounter these situations in our included study.

Dealing with missing data

We tried to contact study authors to obtain missing data. If missing statistics had been present, we would have imputed data by using the formulas as described in section 7.7.3.3 of the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). As the included study had drop-outs, we used what the paper reported and dealt with it in the 'Risk of bias' assessment.

Assessment of heterogeneity

We intended to assess heterogeneity between the trials by examining the forest plot to check for overlapping CIs, using the χ^2 test for heterogeneity with a 10% level of significance to detect inconsistency in study results that were not due to random error (chance), and the I^2 statistic to denote the percentage of inconsistency in results due to inter-trial variability that exceeded chance. In general, we interpret an I^2 value of 50% or greater to denote significant heterogeneity (Higgins 2003). We acknowledge that this cut-off is arbitrary. Therefore, we intended to interpret I^2 values between 0% to 40% as possibly insignificant, 30% to 60% as possibly significant, 50% to 90% as possibly substantial and 75% to 100% as substantial; depending on whether the inconsistency in results was due to differences in the direction of effects estimates between trials rather than due to differences in the magnitude of effect estimates favouring an intervention; and the strength of the evidence for heterogeneity from the P value for the χ^2 test for heterogeneity (Deeks 2011). However, we did not need to assess for heterogeneity in the present review.

Assessment of reporting biases

As there is only one study included in the review, we did not assess the possible presence of reporting bias by testing for asymmetry in a funnel plot. If there had been a sufficient number of trials (more than 10), we would have done statistical analysis using the methods described by Egger 1997.

Data synthesis

We analysed the data using Review Manager 5 software ([RevMan 2014](#)). If the data available from the studies had similar comparisons and outcomes, we planned to undertake meta-analysis. We planned to use a random-effects model because in this approach, the CIs for the mean intervention effect will be wider than those obtained using a fixed-effect approach, leading to a more conservative interpretation. We planned to use all change scores or end scores when available, and combine change and end scores where necessary using the criteria in Section 9.4.5.2 of the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011](#)). We planned to report the results from studies not suitable for inclusion in a meta-analysis in additional tables. However, none of them was done as there was only one included study in the review.

Subgroup analysis and investigation of heterogeneity

We did not conduct any subgroup analysis as there was only one included trial. In future updates, we may consider the following.

- Size of the defect
- Immune status of the participant
- End score versus change score

Sensitivity analysis

As there was only one included study, we did not undertake sensitivity analysis.

Summary of findings

We used the GRADE approach to interpret findings ([Schünemann 2011](#)). We used GRADE Profiler software ([GRADE 2004](#)); and imported data from Review Manager 5 ([RevMan 2014](#)) to create 'Summary of findings' tables for each comparison included in the review. These tables provided information concerning the overall

quality of the evidence from the trials, the magnitude of effect of the interventions examined and the sum of available data on the primary and secondary outcomes. The GRADE approach considers 'quality' to be a judgement of the extent to which we can be confident that the estimates of effect are correct ([Schünemann 2011](#)). A body of evidence from RCTs is initially graded as high and downgraded by one or two levels on each of five domains after full consideration of: risk of bias, directness (or applicability) of the evidence, consistency of results, precision of results and possibility of publication bias. A quality level of 'high' reflects confidence that the true effect lies close to that of the estimate of the effect for an outcome. A judgement of 'moderate' quality indicates that the true effect is likely to be close to the estimate of the effect, but acknowledges the possibility that it could be substantially different. 'Low' and 'very low' quality evidence limit our confidence in the effect estimate ([Balslem 2011](#)).

RESULTS

Description of studies

See [Characteristics of included studies](#) and [Characteristics of studies awaiting classification](#) table.

Results of the search

The electronic search strategies identified 273 records from English and other language databases. We had 164 records after de-duplication. From the 164 records, we discarded 155 after screening the abstracts as they were not relevant; and requested full-text copies of nine studies. From the nine studies, six were rejected, as they were not RCTs. One trial was excluded as it was a non-inferiority trial, which used retrospective surgical data for comparison. The remaining two studies were assessed for eligibility. One study awaits classification. Therefore we included only one RCT in the review. See [Figure 1](#) for details of the selection process and [Table 1](#) for summary of the search process.

Figure 1. Study flow diagram

Figure 1. (Continued)

1 study included quantitative synthesis

Included studies

Trial design

[Nezafati 2012](#), the included trial, was conducted in Iran. It was a single centre trial with parallel group design. The trial was funded by Tabriz University of Medical Sciences, Iran.

Participants

The trial included a total of 22 adults aged 25 to 56 years, and did not state how many participants were male and how many were female. Consenting patients who had established oro-antral communications which were closed at the time of surgery were recruited in the trial. Two of the 22 participants did not report for follow-up and were excluded from the study.

Interventions

The trial compared pedicled buccal fat pad flap with buccal flap. The trial duration was two years.

Outcomes

The reported relevant outcome was successful (complete) closure of oro-antral communication (measured at one month). The

authors reported that both the groups achieved successful closure of OAC. We presented this outcome as dichotomous data in our analysis.

Excluded studies

We excluded one study because it was a non-inferiority trial in which retrospective surgical data was used for comparison ([Bos 2015](#)).

Studies awaiting classification

We categorised one study as 'awaiting classification' ([Gacic 2009](#)). The trial compared resorbable PLGA-coated beta-TCP root analogues, haemostatic gauze and buccal flaps. We are unclear whether the study is an RCT or not as the authors did not explain the randomisation method and there is missing information regarding the numbers in each group. We have contacted the study authors for clarification. For details, see [Characteristics of studies awaiting classification](#).

Risk of bias in included studies

See [Figure 2](#).

Figure 2. Risk of bias summary: review authors' judgements about each risk of bias item

Allocation

Sequence generation was judged as low risk as the trial authors used Rand List 1.2 software to generate the randomisation sequence; however, the trial authors did not describe the method

of allocation concealment and hence was judged as unclear risk of selection bias.

Blinding

The study was at low risk of performance and detection bias since the participants and the dentist who gathered the data were not aware of the type of surgery.

Incomplete outcome data

The study was at low risk of bias as there are only two drop-outs, one from each group.

Selective reporting

Although a study protocol was not available to compare the outcomes, as the trial was not registered in the Iranian Clinical Trial Registry, we judged the study as being at low risk of reporting bias as the expected outcomes were appropriately reported.

Other potential sources of bias

The study was at unclear risk of bias due to the following aspects:

1. Some of the participants whose surgery was not in the first two days of the trial had an additional sinus lavage with saline to reduce the chances of contamination.
2. The length of time participants had OAC is not mentioned.
3. A table of baseline characteristics is not provided.

Overall risk of bias

The overall risk of bias is judged unclear.

Effects of interventions

See: [Summary of findings for the main comparison](#) Pedicled buccal fat pad flap compared to buccal flap for treatment of oro-antral communications and fistulae due to dental procedures

Pedicled buccal fat pad flap (PBFPF) versus buccal flap (BF)

1.1 Successful (complete) closure of the oro-antral communication

There was no difference between BF and PBFPF for the complete closure of oro-antral communication (RR 1.00, 95% CI 0.83 to 1.20; one RCT, 20 participants; [Analysis 1.1](#); [Figure 3](#)).

Figure 3. Forest plot of comparison: 1 PBFPF versus BF, outcome: 1.1 Successful (complete) closure of the oro-antral communication

Secondary outcomes

The study did not report any adverse effects or the time required for healing.

Subgroup and sensitivity analyses

We did not conduct subgroup analysis and sensitivity analysis as there was only a single included trial in this review.

DISCUSSION

Summary of main results

The main objective of the review was to assess effectiveness and safety of various interventions for the treatment of oro-antral communications and fistulae due to dental procedures. We found only one study, which compared pedicled buccal fat pad flap (PBFPF) with buccal flap (BF) ([Nezafati 2012](#)). The trial provided the data for the primary outcome of successful (complete) closure of oro-antral communication (OAC).

The authors did not report any adverse events. We assessed the body of evidence using [GRADE 2004](#), which incorporates risk of bias, the directness of the evidence, the consistency of the results, the precision of the estimates and the possibility of publication bias. The quality of the evidence is very low.

The main results of this review are as follows.

- The single trial included was judged as unclear for risk of bias.
- PBFPF compared to BF showed no difference in the closure of OAC (very low quality of evidence) ([Summary of findings for the main comparison](#)).

Overall completeness and applicability of evidence

Completeness

In spite of our comprehensive search of multiple databases, we found only one trial assessing the effectiveness of PBFPF compared to BF for treating OAC due to dental procedures with a total sample size of 20. Data from this trial was insufficient to provide precise results, as it had insufficient sample size for relevant comparisons and was judged unclear for risk of bias. The trial did not report any definite adverse events with interventions.

We found one trial evaluating resorbable PLGA-coated β -TCP root analogues, haemostatic gauze and buccal flaps for the closure of the OAC, and if it is included in future updates of this review, it may provide additional evidence on effects of interventions for treating OAC due to dental procedures.

We are not aware of any ongoing trials assessing any of the interventions proposed or commonly used for treating OAC and oral-antral fistula (OAF). We did not find any trials assessing the

effectiveness of interventions on children. The evidence base is thus incomplete and insufficient to draw robust conclusions about the most effective intervention for treating OAC due to dental procedures.

Applicability

The results obtained from this Cochrane review are insufficient to determine the relative efficacy of PBFPF and BF for treatment of oro-antral communications or fistulae. The study awaiting classification has compared two non-surgical techniques (PLGA-coated β -TCP and haemostatic gauze) with a surgical technique (buccal flap). Even if the efficacy of these interventions is confirmed in future trials, it is uncertain whether these techniques can be universally applied to different cases of OAC/OAF with varying defect sizes. More RCTs involving surgical and non-surgical interventions with emphasis on the size and type of defect need to be done in order to draw definitive conclusions on the applicability of the interventions.

Quality of the evidence

We have included one parallel-arm RCT comparing PBFPF with BF in 20 adult participants after two dropouts. The quality of evidence was assessed for the primary outcome, namely successful (complete) closure of oro-antral communication. We downgraded the quality of evidence by one level for unclear risk of bias and by two levels for imprecision. Overall, the evidence is very low quality. The results do not allow us to draw any firm conclusions regarding whether one of these interventions is more effective than the other.

Potential biases in the review process

We have taken steps to minimise the bias in every step of the review. All the mentioned databases, conference proceedings and trial registries have been searched to include all the relevant reports. We tried to contact the authors for missing data through emails. If the reports were very old, we tried to get the contact details of the authors through peer contacts, Google search and university/hospital web sites where they were previously affiliated. We tried our best to follow the methodology stated in the protocol. We used standard methods described in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011), and ensured compliance with the Cochrane standards for the conduct of new reviews of interventions (MECIR 2011). In spite of our comprehensive search strategies, we cannot rule out publication bias occurring due to non-identification of unpublished trials.

Agreements and disagreements with other studies or reviews

We found one systematic review and meta-analysis on the frequency and treatment of oro-antral communications (Borja 2011). The purpose of the review was to determine the optimum surgical treatment for OAC and to understand the postoperative complications associated with the procedure. The literature search was carried out for the period between 1983 and 2008. Fifteen non-RCTs were included in this systematic review. The review author concluded that Bichat's fat pad grafts showed excellent results for closure of OAC and low rate of failure and good patient satisfaction postoperatively. The inclusion criteria for Borja 2011 and our review vary considerably and hence the results are not comparable.

AUTHORS' CONCLUSIONS

Implications for practice

There is very low quality evidence to inform clinicians and patients how best to treat oro-antral communications. There was no RCT evidence available for treating oro-antral fistulae. We found a single small study that compared surgical techniques pedicled buccal fat pad flap versus buccal flap and did not show evidence of a difference in effectiveness for treating OAC caused by dental procedures. All oro-antral communications were closed by one month after surgery so it was not possible to assess adverse effects from treatment failure. We did not find any evidence on any other interventions for treating this condition.

Implications for research

Further research should be done in interventions for treating oro-antral communications and fistulae due to dental procedures by conducting well-planned RCTs with more clarity and uniformity in the variables. In designing such clinical trials, the following should be considered.

Evidence: Trials should evaluate all the outcomes mentioned in this review. Future research on adverse effects of different interventions should include longer follow-up. Furthermore, reports on clinical trials would be improved by following CONSORT group guidelines.

Population: Inclusion criteria for clinical trials should be well defined. The trials should include different sizes of OAC/OAF. Outcome measures should be clearly defined for different age groups.

Intervention: More interventional studies should be conducted on both surgical and non-surgical interventions.

Comparison: Comparisons between two different interventions (surgical versus non-surgical or non-surgical versus non-surgical) can be considered in future trials.

Outcome: Along with complete closure of OAC and OAF, patient-reported outcomes in terms of satisfaction and quality of life should be evaluated in the trials.

ACKNOWLEDGEMENTS

For this update, we are thankful to Cochrane Oral Health, especially Anne Littlewood, Information Specialist; Laura MacDonald, Managing Editor; and Anne-Marie Glenny, Deputy Co-ordinating Editor. For the original review, we are extremely thankful to Anne Littlewood, Information Specialist; Helen Wakeford, Deputy Managing Editor; Joanne Weldon, Research/Editorial Support Co-ordinator; Philip Riley, Editor; Luisa M Fernandez Mauleffinch and Laura MacDonald, Managing Editors; and Helen Worthington, Co-ordinating Editor. We also thank Paul Coulthard, Ruth Floate, Edmund Bailey, Leonardo Righesso and Anirudha Agnihotry for their comments on the original review. We thank Prof. Datuk Dr. Abdul Razzak, Pro VC, Manipal University, Melaka campus, for his constant encouragement to take up Cochrane reviews; Prof. Dr. Jaspal Singh Sahota, CE, Melaka Manipal campus for his support; Prof. Adinegara Lutfi Abas, Dean, Faculty of Medicine, MMMC; Prof. Abdul Rashid Hj Ismail, Dean, Faculty of Dentistry, MMMC; and Prof. Dr. Ravi Kant, Vice Chancellor, King George's Medical University UP, Lucknow, India for all the suggestions and help.

during the review preparation. We thank Prof. Noorliza Mastura Ismail, Head of the Department of Community Dentistry, Faculty of Dentistry, MMMC for proofreading the review and helping us with language correction. Our sincere gratitude and thanks to Dr. Prathap Tharyan, Director of the South Asian Cochrane Centre,

CMC, Vellore for his motivation. We are indebted to Ms. Shazana MS, Chief Librarian, Melaka Manipal Medical College, and Ms. Janet Lear, School of Dentistry, University of Manchester for helping us procure full-text articles.

REFERENCES

References to studies included in this review

Nezafati 2012 {published data only}

Nezafati S, Vafaii A, Ghojzadeh M. Comparison of pedicled buccal fat pad flap with buccal flap for closure of oro-antral communication. *International Journal of Oral and Maxillofacial Surgery* 2012;**41**(5):624-8. [PUBMED: 22192386]

References to studies excluded from this review

Bos 2015 {published data only}

Bos RRM, Visscher SH, Minnen van B. Is biodegradable PU foam as effective as surgery for closure of oroantral communications? A prospective clinical trial. *International Journal of Oral and Maxillofacial Surgery* 2015;**44**(S1):e191.

References to studies awaiting assessment

Gacic 2009 {published data only}

Gacic B, Todorovic L, Kokovic V, Danilovic V, Stojcev-Stajcic L, Drazic R, et al. The closure of oroantral communications with resorbable PLGA-coated beta-TCP root analogs, haemostatic gauze, or buccal flaps: a prospective study. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontics* 2009;**108**(6):844-50. [PUBMED: 19875318]

Additional references

Abuabara 2006

Abuabara A, Cortez AL, Passeri LA, de Moraes M, Moreira RW. Evaluation of different treatments for oroantral/oronasal communications: experience of 112 cases. *International Journal of Oral and Maxillofacial Surgery* 2006;**35**(2):155-8. [PUBMED: 15955666]

Assad 2017

Mounzer A, Waseem B, Alhajj Nasser AM. Closure of oroantral communication using platelet-rich fibrin: a report of two cases. *Annals of Maxillofacial Surgery* 2017;**7**(1):117-9.

Balshem 2011

Balshem H, Helfand M, Schünemann HJ, Oxman AD, Kunz R, Brozek J, et al. GRADE guidelines: 3. Rating the quality of evidence. *Journal of Clinical Epidemiology* 2011;**64**(4):401-6. [PUBMED: 21208779]

Batra 2010

Batra H, Jindal G, Kaur S. Evaluation of different treatment modalities for closure of oro-antral communications and formulation of a rational approach. *Journal of Maxillofacial & Oral Surgery* 2010;**9**(1):13-8. [PUBMED: 23139559]

Blythe 2016

Blythe JN, Moffat C, Bakshi A, Brennan PA. Novel approach in the treatment of a persistent oroantral fistula using an acellular dermal graft tissue: technical note. *British Journal of Oral & Maxillofacial Surgery* 2016;**54**(6):700-1. [PUBMED: 26656268]

Borgonovo 2012

Borgonovo AE, Berardinelli FV, Favale M, Maiorana C. Surgical options in oroantral fistula treatment. *Open Dentistry Journal* 2012;**6**:94-8. [PUBMED: 22715347]

Borja 2011

Franco-Carro B, Barona-Dorado C, Martínez-González MJ, Rubio-Alonso LJ, Martínez-González JM. Meta-analytic study on the frequency and treatment of oral antral communications. *Medicina Oral, Patología Oral y Cirugía Bucal* 2011;**16**(5):682-7. [PUBMED: 20711106]

Buric 2012

Buric N, Jovanovic G, Krasic D, Tijanac M, Buric M, Tarana S, et al. The use of absorbable polyglactin/polydioxanone implant (Ethisorb®) in non-surgical closure of oro-antral communication. *Journal of Cranio-maxillo-facial Surgery* 2012;**40**(1):71-7. [PUBMED: 21377886]

Buric 2013

Buric N. Use of N-butyl cyanoacrylate with metacryloxisulfolane (Glubran 2) surgical glue for flapless closure of oroantral communication. *Implant Dentistry* 2013;**22**(3):238-43. [PUBMED: 23442603]

Choi 2006

Choi BH, Zhu SJ, Jung JH, Lee SH, Huh JY. The use of autologous fibrin glue for closing sinus membrane perforations during sinus lifts. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontics* 2006;**101**(2):150-4. [PUBMED: 16448914]

Deeks 2011

Deeks JJ, Higgins JPT, Altman DG. Chapter 9: Analysing data and undertaking meta-analyses. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Demetoglu 2018

Demetoglu U, Ocak H, Bilge S. Closure of oroantral communication with plasma-rich fibrin membrane. *Journal of Craniofacial Surgery* 2018;**29**(4):e367-70. [PUBMED: 29485557]

Dym 2012

Dym H, Wolf JC. Oroantral communication. *Oral and Maxillofacial Surgery Clinics of North America* 2012;**24**(2):239-47. [PUBMED: 22503070]

Egger 1997

Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;**315**(7109):629-34. [PUBMED: 9310563]

Eneroth 1961

Eneroth CM, Martensson G. Closure of antro-alveolar fistulae. *Acta Oto-laryngologica* 1961;**53**:477-85. [PUBMED: 13696774]

Franco-Carro 2011

Franco-Carro B, Barona-Dorado C, Martínez-González MJ, Rubio-Alonso LJ, Martínez-González JM. Meta-analytic study on the frequency and treatment of oral antral communications. *Medicina Oral, Patología Oral, Cirugía Bucal* 2011;**16**(5):682-7. [PUBMED: 20711106]

Goodger 2004

Goodger NM, Halfpenny W, Williams S. An unusual cause of oro-antral fistula. *Dental Update* 2004;**31**(1):31-3. [PUBMED: 15000006]

GRADE 2004 [Computer program]

Brozek J, Oxman A, Schunemann H. GRADEpro. Version 3.6 for Windows. Grade Working Group, 2012.

Grzesiak-Janias 2001

Grzesiak-Janias G, Janias A. Conservative closure of antro-oral communication stimulated with laser light. *Journal of Clinical Laser Medicine & Surgery* 2001;**19**(4):181-4. [PUBMED: 11523860]

Higgins 2003

Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327**(7414):557-60. [12958120]

Higgins 2011

Higgins JPT, Green S (editors). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0. [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Kale 2010

Kale TP, Urolagin S, Khurana V, Kotrashetti SM. Treatment of oro antral fistula using palatal flap - A case report and technical note. *Journal of International Oral Health* 2010;**2**(3):78-82.

Kamadajaja 2008

Kamadajaja DB. The role of proper treatment of maxillary sinusitis in persistent oroantral communication. *Dental Journal* 2008;**41**(3):128-31.

Kitagawa 2003

Kitagawa Y, Sano K, Nakamura M, Ogasawara T. Use of third molar transplantation for closure of the oroantral communication after tooth extraction: a report of 2 cases. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontics* 2003;**95**(4):409-15. [PUBMED: 12686925]

Lefebvre 2011

Lefebvre C, Manheimer E, Glanville J. Chapter 6: Searching for studies. In: Higgins JP, Green S, editor(s). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Liversedge 2002

Liversedge RL, Wong K. Use of the buccal fat pad in maxillary and sinus grafting of the severely atrophic maxilla preparatory to implant reconstruction of the partially or completely

edentulous patient: technical note. *International Journal of Oral & Maxillofacial Implants* 2002;**17**(3):424-8. [PUBMED: 12074460]

Logan 2003

Logan RM, Coates EA. Non-surgical management of an oro-antral fistula in a patient with HIV infection. *Australian Dental Journal* 2003;**48**(4):255-8. [PUBMED: 14738129]

Malik 2008

Malik NA. Textbook of Oral and Maxillofacial Surgery. Second Edition. New Delhi: Jaypee Brothers Medical Publishers, 2008.

MECIR 2011

Chandler J, Churchill R, Higgins J, Lasserson T, Tovey D. Methodological Expectations of Cochrane Intervention Reviews (MECIR). Methodological standards for the conduct of new Cochrane Intervention Reviews. Version 2.3. Available at: http://editorial-unit.cochrane.org/sites/editorial-unit.cochrane.org/files/uploads/MECIR_conduct_standards%202.3%2002122013_0.pdf 2 December 2013 (accessed on 15 August 2015).

Moore 1991

Moore JR, Gillbe GV. Principles of Oral Surgery. 4th Edition. Manchester University Press, 1991.

Nilesh 2018

Nilesh K, Vande AV. Mucormycosis of maxilla following tooth extraction in immunocompetent patients: reports and review. *Journal of Clinical and Experimental Dentistry* 2018;**10**(3):e300-5. [PUBMED: 29721234]

RevMan 2014 [Computer program]

Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Saleh 2013

Saleh EA, Issa IA. Closure of large oroantral fistulas using septal cartilage. *Otolaryngology--head and neck surgery: official journal of American Academy of Otolaryngology-Head and Neck Surgery* 2013;**148**(6):1048-50. [PUBMED: 23520068]

Sandhya 2013

Sandhya G, Reddy PB, Kumar KA, Sridhar Reddy B, Prasad N, Kiran G. Surgical management of oro-antral communications using resorbable GTR membrane and FDMB sandwich technique: a clinical study. *Journal of Maxillofacial and Oral Surgery* 2013;**12**(3):254-59. [PUBMED: 24431851]

Scala 2007

Scala M, Gipponi M, Pasetti S, Dellachá E, Ligorio M, Villa G, et al. Clinical applications of autologous cryoplatelet gel for the reconstruction of the maxillary sinus. A new approach for the treatment of chronic oro-sinus fistula. *In Vivo (Athens, Greece)* 2007;**21**(3):541-7. [PUBMED: 17591367]

Schünemann 2011

Schünemann HJ, Oxman AD, Vist GE, Higgins JPT, Deeks JJ, Glasziou P, et al. Chapter 12: Interpreting results drawing

conclusions. In: Higgins JPT, Green S(editors). *Cochrane Handbook for Systematic Reviews of Intervention* Version 5.0.1 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Steiner 2008

Steiner M, Gould AR, Madion DC, Abraham MS, Loeser JG. Metal plates and foils for closure of oroantral fistulae. *Journal of Oral and Maxillofacial Surgery* 2008;**66**(7):1551-5. [PUBMED: 18571051]

Thoma 2006

Thoma K, Pajarola GF, Gratz KW, Schmidlin PR. Bioabsorbable root analogue for closure of oroantral communications after tooth extraction: a prospective case-cohort study. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontics* 2006;**101**(5):558-64. [PUBMED: 16632265]

Visscher 2010

Visscher SH, Van Minnen B, Bos RR. Closure of oroantral communications: a review of the literature. *Journal of Oral and Maxillofacial Surgery: Official Journal of the American Association of Oral and Maxillofacial Surgeons* 2010;**68**(6):1384-91. [PUBMED: 20227153]

Visscher 2011

Visscher SH, Van Roon MR, Sluiter WJ, Van Minnen B, Bos RR. Retrospective study on the treatment outcome of surgical closure of oroantral communications. *Journal of Oral and Maxillofacial Surgery* 2011;**69**(12):2956-61. [PUBMED: 21752508]

Von Wowern 1982

Von Wowern N. Closure of oroantral fistula with buccal flap: Rehrmann versus Môczár. *International Journal of Oral Surgery* 1982;**11**(3):156-65. [PUBMED: 6813275]

Waldrop 1993

Waldrop TC, Semba SE. Closure of oroantral communication using guided tissue regeneration and an absorbable gelatin membrane. *Journal of Periodontology* 1993;**64**(11):1061-6. [PUBMED: 8295091]

Watzak 2005

Watzak G, Tepper G, Zechner W, Monov G, Busenlechner D, Watzek G. Bony press-fit closure of oro-antral fistulas: a technique for pre-sinus lift repair and secondary closure. *Journal of Oral and Maxillofacial Surgery* 2005;**63**(9):1288-94. [PUBMED: 16122592]

Worthington 2015

Worthington H, Clarkson J, Weldon J. Priority oral health research identification for clinical decision-making. *Evidence-based Dentistry* 2015;**16**(3):69-71.

Zide 1992

Zide MF, Karas ND. Hydroxylapatite block closure of oroantral fistulas: report of case. *Acta Stomatologica Croatica* 1992;**50**(1):71-5. [PUBMED: 1309241]

References to other published versions of this review

Kiran Kumar Krishanappa 2016

Kiran Kumar Krishanappa S, Prashanti E, Sumanth KN, Naresh S, Moe S, Aggarwal H, Mathew RJ. Interventions for treating oro-antral communications and fistulae due to dental procedures. *Cochrane Database of Systematic Reviews* 2016, Issue 5. [DOI: [10.1002/14651858.CD011784.pub2](https://doi.org/10.1002/14651858.CD011784.pub2)]

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Nezafati 2012

Methods	<p>Trial design: double-blinded randomised clinical trial</p> <p>Trial location: Tabriz University of Medical Sciences</p> <p>Number of centres: one</p> <p>Recruitment period (duration): two years (2006 to 2008)</p> <p>Funding source: Tabriz University of Medical Sciences</p>
Participants	<p>Participants with chief complaint of oro-antral communication</p> <p>Total number: 20 participants aged 25 to 56 years (actual enrolment was 22 but 2 participants, one from each group, did not attend for follow-up and they were excluded)</p> <p>Inclusion criteria</p> <ul style="list-style-type: none"> Oro-antral communication <p>Exclusion criteria</p>

Nezafati 2012 (Continued)

- Systemic disease affecting healing
- Smoking, drug and alcohol abuse
- History of sinus disease
- History of sinus surgery
- Intra-antral foreign body
- People with need for Caldwell-Luc Procedure

Number randomised: 22

Number evaluated: 20 (2 dropouts - one from each group)

Interventions	<p>Total number of intervention groups: two</p> <p>Group 1: buccal flap</p> <p>Group 2: pedicled buccal fat pad flap</p> <p>Duration of treatment: from first incision to the last suture measured with a chronometer.</p> <p>Group 1: buccal flap – 23.61 ± 7.02 minutes</p> <p>Group 2: pedicled buccal fat pad flap – 26.01 ± 10.49 minutes</p>
Outcomes	<p>Relevant outcome reported and used in our review:</p> <ul style="list-style-type: none"> • Successful (complete) closure of oro-antral communication (OAC) <p>Other outcomes reported in the trial:</p> <ul style="list-style-type: none"> • Mean pain score • Proportion of postoperative swelling • Reduction in maximum mouth opening (MMO)
Notes	<p>Sample size calculation: not reported</p> <p>Adverse effects: none reported</p> <p>Health-related quality of life: not reported</p> <p>Key conclusions of the study authors: both methods were equally successful for the closure of OAC as tested by negative nose blowing one month later indicating closure.</p> <p>PBFPF group had a higher mean pain score, more pronounced swelling, and reduced mouth opening at 2 and 7 days after surgery compared to control group. Maximum mouth opening was no different between groups at 1 month.</p> <p>Correspondence required: yes, author was contacted via email on 8/07/2015 and 25/07/2015 with the following queries, however there was no response.</p> <ul style="list-style-type: none"> • Number randomised • Method of allocation concealment • Baseline characteristics <p>We contacted trial authors regarding duration of OAC, allocation concealment and baseline characteristics (8/07/2015 and 25/07/2015). No response was received.</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Nezafati 2012 *(Continued)*

Random sequence generation (selection bias)	Low risk	Quote from article: "After completing the consent form, the patients were randomly divided into two groups using Rand List 1.2 software"
Allocation concealment (selection bias)	Unclear risk	Not mentioned in the trial
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote from article: "The patients and the dentist who gathered the data were not aware of the type of surgery"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote from article: "The patients and the dentist who gathered the data were not aware of the type of surgery"
Incomplete outcome data (attrition bias) All outcomes	Low risk	The study was at low risk as there were only two drop-outs, one from each group.
Selective reporting (reporting bias)	Low risk	Study trial not registered in Iranian National Trial Registry (we searched Iranian Clinical Trial Registry on 15 April 2016) so study protocol was not available to compare the outcomes; however, expected outcomes seemed to be reported appropriately.
Other bias	Unclear risk	<ol style="list-style-type: none"> 1. Some of the participants whose surgery was not in the first two days of the trial had an additional sinus lavage with saline to reduce the chances of contamination. 2. The length of time participants had OAC is not mentioned. 3. A table of baseline characteristics is not provided.

Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
Bos 2015	Non-inferiority trial with retrospective surgical data as comparison

Characteristics of studies awaiting assessment *[ordered by study ID]*
Gacic 2009

Methods	<p>Trial design (including number of arms): not clear, but states participants were randomly allocated to interventions</p> <p>Location and setting: University of Belgrade</p> <p>Number of centres: one</p> <p>Recruitment period (duration): not available</p> <p>Funding source: PLGA-coated β-TCP and instruments for making root analogue provided by Degradable Solutions Schlieren, Switzerland</p>
Participants	<p>Total number: 30</p> <p>Inclusion criteria</p>

Gacic 2009 (Continued)

- Oro-antral communication following extraction of tooth

Exclusion criteria

- Smokers
- Pregnant or lactating women
- Patient under medication

Number randomised: not available

Method of randomisation: not available

Method of allocation concealment: not available

Method of blinding: not available

Number evaluated: not available

Interventions	<p>Total number of intervention groups: three</p> <ul style="list-style-type: none"> • Group 1: PLGA-coated β-TCP • Group 2: haemostatic gauze • Group 3: buccal flap <p>Duration of treatment: not available</p>
Outcomes	<ul style="list-style-type: none"> • Closure OAC • Vestibular depth • Pain intensity • Swelling • Histologic analysis of the surgical site after implant placement: to check for necrosis/inflammation
Notes	<p>Sample size calculation: not mentioned</p> <p>Adverse effects: not available</p> <p>Health-related quality of life: not available</p> <p>Key conclusions of the study authors: closure of OAC with PLGA-coated β-TCP or haemostatic gauze are reliable and minimally invasive methods. They showed minimal atrophy of alveolar bone, swelling and pain compared to buccal flap technique.</p> <p>Correspondence required: Yes, author was contacted via email on 8/05/2018, 30/5/2018 and 1/7/2018 with the following queries; however, there was no response:</p> <ul style="list-style-type: none"> • Trial design • Number randomised • Method of randomisation • Method of allocation concealment • Method of blinding • Number evaluated

DATA AND ANALYSES

Comparison 1. PBFPF versus BF

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Successful (complete) closure of the oro-antral communication	1	20	Risk Ratio (M-H, Fixed, 95% CI)	1.0 [0.83, 1.20]

Analysis 1.1. Comparison 1 PBFPF versus BF, Outcome 1 Successful (complete) closure of the oro-antral communication.

ADDITIONAL TABLES

Table 1. Summary of the search results

May 2018 searches carried out by Anne Littlewood, Information Specialist, Cochrane Oral Health			
Database	Version/issue	Date of search	Records retrieved
Cochrane Oral Health Trials Register	New to authors since 3 July 2015	23.05.18	14
Cochrane Central Register of Controlled Trials (CENTRAL), <i>The Cochrane Library</i>	Issue 6, 2015 to Issue 4, 2018	23.05.18	24
MEDLINE Ovid (inc ePub ahead of print, pre-indexed etc)	3 July 2015 to 23 May 2018	23.05.18	35 (with filter)
Embase Ovid	3 July 2015 to 23 May 2018	23.05.18	29 (with filter)
US National Institutes of Health Trials Registry (ClinicalTrials.gov)	New to authors since 3 July 2015	23.05.18	0
WHO International Clinical Trials Registry Platform	New to authors since 3 July 2015	23.05.18	1

Total number retrieved from electronic searches: 170 (July 2015) + 103 (May 2018) = 273

Total retrieved duplicates removed: 104 (July 2015) + 60 (May 2018) = 164

Table 1. Summary of the search results (Continued)

Total sent to authors for this search: 60

APPENDICES

Appendix 1. Cochrane Oral Health Trials Register search strategy

1. (((oroantral* or oro-antral*) and (fistula* or communicat*))) AND (INREGISTER)
2. (((orosinusual or oro-sinusual) and (fistula* or communicat*))) AND (INREGISTER)
3. ((antral and perforation*)) AND (INREGISTER)
4. ("antro-alveolar fistula*") AND (INREGISTER)
5. ((antrooral or antro-oral)) AND (INREGISTER)
6. (((alveolo-sinusual or palatal-sinusual or vestibulo-sinusual) and fistula*)) AND (INREGISTER)
7. (("maxillary sinus" or "maxillary antrum")) AND (INREGISTER)
8. ((fistula* or perforat*)) AND (INREGISTER)
9. (#7 and #8) AND (INREGISTER)
10. (#1 or #2 or #3 or #4 or #5 or #6 or #9) AND (INREGISTER)

Appendix 2. The Cochrane Central Register of Controlled Trials (CENTRAL) search strategy

- #1 [mh ^"oroantral fistula"]
- #2 ((oroantral* or oro-antral*) and (fistula* or communicat*))
- #3 ((orosinusual or oro-sinusual) and (fistula* or communicat*))
- #4 (antral near/2 perforation*)
- #5 (antro-alveolar next fistula*)
- #6 (antrooral or antro-oral)
- #7 ((alveolo-sinusual or palatal-sinusual or vestibulo-sinusual) next fistula*)
- #8 [mh ^"Maxillary sinus"]
- #9 ("maxillary sinus" or "maxillary antrum")
- #10 #8 or #9
- #11 (fistula* or perforat*)
- #12 #10 and #11
- #13 #1 or #2 or #3 or #4 or #5 or #6 or #7 or #12

Appendix 3. MEDLINE Ovid search strategy

1. Oroantral fistula/
2. ((oroantral\$ or oro-antral\$) and (fistula\$ or communicat\$)).mp.
3. ((orosinusual or oro-sinusual) and (fistula\$ or communicat\$)).mp.
4. (antral adj2 perforation\$).mp.
5. (antro-alveolar adj fistula\$).mp.
6. (antrooral or antro-oral).mp.
7. ((alveolo-sinusual or palatal-sinusual or vestibulo-sinusual) adj fistula\$).mp.
8. Maxillary sinus/
9. ("maxillary sinus" or "maxillary antrum").mp.
10. 8 or 9
11. (fistula or perforat\$).mp.
12. 10 and 11
13. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 12

The above subject search was linked to the Cochrane Highly Sensitive Search Strategy (CHSSS) for identifying randomised trials in MEDLINE: sensitivity maximising version (2008 revision) as referenced in Chapter 6.4.11.1 and detailed in box 6.4.c of *The Cochrane Handbook for Systematic Reviews of Interventions*, Version 5.1.0 [updated March 2011] ([Lefebvre 2011](#)).

1. randomized controlled trial.pt.
2. controlled clinical trial.pt.
3. randomized.ab.
4. placebo.ab.
5. drug therapy.fs.
6. randomly.ab.

7. trial.ab.
8. groups.ab.
9. or/1-8
10. exp animals/ not humans.sh.
11. 9 not 10

Appendix 4. Embase Ovid search strategy

1. Oroantral fistula/
2. ((oroantral\$ or oro-antral\$) and (fistula\$ or communicat\$)).mp.
3. ((orosinusal or oro-sinusal) and (fistula\$ or communicat\$)).mp.
4. (antral adj2 perforation\$).mp.
5. (antro-alveolar adj fistula\$).mp.
6. (antrooral or antro-oral).mp.
7. ((alveolo-sinusal or palatal-sinusal or vestibulo-sinusal) adj fistula\$).mp.
8. Maxillary sinus/
9. ("maxillary sinus" or "maxillary antrum").mp.
- 10.8 or 9
- 11.(fistula or perforat\$).mp.
- 12.10 and 11
- 13.1 or 2 or 3 or 4 or 5 or 6 or 7 or 12

This subject search was linked to an adapted version of the Cochrane Centralised Search Project filter for identifying RCTs in Embase Ovid (see <http://www.cochranelibrary.com/help/central-creation-details.html> for information:)

1. Randomized controlled trial/
2. Controlled clinical study/
3. Random\$.ti,ab.
4. randomization/
5. intermethod comparison/
6. placebo.ti,ab.
7. (compare or compared or comparison).ti.
8. ((evaluated or evaluate or evaluating or assessed or assess) and (compare or compared or comparing or comparison)).ab.
9. (open adj label).ti,ab.
10. ((double or single or doubly or singly) adj (blind or blinded or blindly)).ti,ab.
11. double blind procedure/
12. parallel group\$1.ti,ab.
13. (crossover or cross over).ti,ab.
14. ((assign\$ or match or matched or allocation) adj5 (alternate or group\$1 or intervention\$1 or patient\$1 or subject\$1 or participant \$1)).ti,ab.
15. (assigned or allocated).ti,ab.
16. (controlled adj7 (study or design or trial)).ti,ab.
17. (volunteer or volunteers).ti,ab.
18. trial.ti.
19. or/1-18
20. (exp animal/ or animal.hw. or nonhuman/) not (exp human/ or human cell/ or (human or humans).ti.)
21. 19 not 20

Appendix 5. The US National Institutes of Health Trials Registry (ClinicalTrials.gov) search strategy

oroantral communication
oroantral fistula

Appendix 6. World Health Organization International Clinical Trials Registry Platform search strategy

oroantral

WHAT'S NEW

Date	Event	Description
29 June 2018	New search has been performed	We updated our search to 23 May 2018. We made minor changes to update the Background.
19 June 2018	New citation required but conclusions have not changed	No new studies identified

CONTRIBUTIONS OF AUTHORS

- Salian Kiran Kumar Krishanappa: drafting the protocol, selection of trials, data extraction and entering data into Review Manager 5, data analysis, drafting the final review and updating the review.
- Eachempati Prashanti, Kumbargere Nagraj Sumanth: arbiter, drafting the protocol, carrying out analysis, drafting final review and updating the review.
- Shetty Naresh: drafting the protocol, selection of trials, data extraction and entering data into RevMan and drafting the final review.
- Soe Moe: drafting the protocol, carrying out and interpreting analysis and drafting the final review.
- Himanshi Aggarwal: undertaking searches, selection of trials, drafting final review.
- Rebecca J Mathew: undertaking searches, selection of trials, carrying out and interpreting analysis and drafting final review.

DECLARATIONS OF INTEREST

Salian Kiran Kumar Krishanappa: none known
 Eachempati Prashanti: none known
 Kumbargere N Sumanth: none known
 Shetty Naresh: none known
 Soe Moe: none known
 Himanshi Aggarwal: none known
 Rebecca J Mathew: none known

SOURCES OF SUPPORT

Internal sources

- Faculty of Dentistry, Melaka Manipal Medical College, Melaka, Malaysia.

Library support and training
- South Asian Cochrane Network & Center, Prof. BV Moses Center for Evidence-Informed Health Care and Health Policy, Christian Medical College, Vellore, India.

Training in protocol writing
- The School of Dentistry, The University of Manchester, UK.

External sources

- National Institute for Health Research (NIHR), UK.

This project was supported by the NIHR, via Cochrane Infrastructure funding to the Cochrane Oral Health Group. The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the Systematic Reviews Programme, NIHR, NHS or the Department of Health.

- Cochrane Oral Health Group Global Alliance, Other.

The production of Cochrane Oral Health reviews has been supported financially by our Global Alliance since 2011 (oralhealth.cochrane.org/partnerships-alliances). Contributors over the past year have been the American Association of Public Health Dentistry, USA; AS-Akademie, Germany; the British Association for the Study of Community Dentistry, UK; the British Society of Paediatric Dentistry, UK; the Canadian Dental Hygienists Association, Canada; the Centre for Dental Education and Research at All India Institute of Medical Sciences, India; the National Center for Dental Hygiene Research & Practice, USA; New York University College of Dentistry, USA; Swiss Society of Endodontology, Switzerland.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

There are no differences between the protocol and review.

INDEX TERMS

Medical Subject Headings (MeSH)

Adipose Tissue [*transplantation]; Dental Care [*adverse effects]; Oroantral Fistula [etiology] [*surgery]; Randomized Controlled Trials as Topic; Surgical Flaps [*transplantation]

MeSH check words

Adult; Humans; Middle Aged