Cochrane
Library

Cochrane Database of Systematic Reviews

Biochemical tests of placental function versus ultrasound assessment of fetal size for stillbirth and small-for-gestational-age infants (Review)

Heazell AEP, Hayes DJL, Whitworth M, Takwoingi Y, Bayliss SE, Davenport C

Heazell AEP, Hayes DJL, Whitworth M, Takwoingi Y, Bayliss SE, Davenport C.

Biochemical tests of placental function versus ultrasound assessment of fetal size for stillbirth and small-for-gestational-age infants.

Cochrane Database of Systematic Reviews 2019, Issue 5. Art. No.: CD012245.

DOI: [10.1002/14651858.CD012245.pub2](https://doi.org/10.1002/14651858.CD012245.pub2).

www.cochranelibrary.com

Biochemical tests of placental function versus ultrasound assessment of fetal size for stillbirth and small-for-gestational-age infants (Review)

Copyright © 2019 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

WILEY

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	4
BACKGROUND	7
Figure 1.	8
OBJECTIVES	9
METHODS	9
RESULTS	12
Figure 2.	13
Figure 3.	15
Figure 4.	16
Figure 5.	17
Figure 6.	19
Figure 7.	20
Figure 8.	22
Figure 9.	24
Figure 10.	26
Figure 11.	27
Figure 12.	28
Figure 13.	30
DISCUSSION	31
AUTHORS' CONCLUSIONS	32
ACKNOWLEDGEMENTS	33
REFERENCES	34
CHARACTERISTICS OF STUDIES	59
DATA	198
Test 1. EFW and SGA.	198
Test 2. Placental grading and SGA.	199
Test 3. hPL and SGA.	199
Test 4. E3 and SGA.	199
Test 5. Urinary E3 and SGA.	199
Test 6. PlGF and SGA.	199
Test 7. UA and SGA.	199
Test 8. E3 AND/OR hPL and SGA.	199
Test 9. EFW for SGA - 10th centile threshold only.	199
Test 10. hPL for SGA - 10th centile threshold only.	199
Test 11. EFW and SGA3.	199
Test 13. PlGF and SGA3.	199
Test 14. hPL and stillbirth.	199
Test 15. Urinary E3 and stillbirth.	199
Test 16. PlGF and stillbirth.	200
Test 17. UA and stillbirth.	200
Test 18. Placental grading and stillbirth.	200
Test 19. SGA data from studies with both SGA infants and stillbirths.	200
Test 20. Stillbirth data from studies with both SGA infants and stillbirths.	200
Test 22. PlGF or EFW and SGA.	200
Test 23. PlGF or EFW and SGA3.	200
ADDITIONAL TABLES	201
APPENDICES	214
Figure 14.	224

Figure 15.	225
Figure 16.	226
Figure 17.	227
Figure 18.	229
Figure 19.	230
Figure 20.	231
Figure 21.	234
WHAT'S NEW	234
CONTRIBUTIONS OF AUTHORS	235
DECLARATIONS OF INTEREST	235
SOURCES OF SUPPORT	235
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	235
INDEX TERMS	236

[Diagnostic Test Accuracy Review]

Biochemical tests of placental function versus ultrasound assessment of fetal size for stillbirth and small-for-gestational-age infants

Alexander EP Heazell¹, Dexter JL Hayes¹, Melissa Whitworth¹, Yemisi Takwoingi², Susan E Bayliss², Clare Davenport²

¹Maternal and Fetal Health Research Centre, University of Manchester, Manchester, UK. ²Institute of Applied Health Research, University of Birmingham, Birmingham, UK

Contact address: Alexander EP Heazell, Maternal and Fetal Health Research Centre, University of Manchester, 5th floor (Research), St Mary's Hospital, Oxford Road, Manchester, M13 9WL, UK. alexander.heazell@manchester.ac.uk, alex_heazell@talk21.com.

Editorial group: Cochrane Pregnancy and Childbirth Group.

Publication status and date: Edited (no change to conclusions), published in Issue 6, 2019.

Citation: Heazell AEP, Hayes DJL, Whitworth M, Takwoingi Y, Bayliss SE, Davenport C. Biochemical tests of placental function versus ultrasound assessment of fetal size for stillbirth and small-for-gestational-age infants. *Cochrane Database of Systematic Reviews* 2019, Issue 5. Art. No.: CD012245. DOI: [10.1002/14651858.CD012245.pub2](https://doi.org/10.1002/14651858.CD012245.pub2).

Copyright © 2019 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Stillbirth affects 2.6 million pregnancies worldwide each year. Whilst the majority of cases occur in low- and middle-income countries, stillbirth remains an important clinical issue for high-income countries (HICs) - with both the UK and the USA reporting rates above the mean for HICs. In HICs, the most frequently reported association with stillbirth is placental dysfunction. Placental dysfunction may be evident clinically as fetal growth restriction (FGR) and small-for-dates infants. It can be caused by placental abruption or hypertensive disorders of pregnancy and many other disorders and factors

Placental abnormalities are noted in 11% to 65% of stillbirths. Identification of FGR is difficult in utero. Small-for-gestational age (SGA), as assessed after birth, is the most commonly used surrogate measure for this outcome. The degree of SGA is associated with the likelihood of FGR; 30% of infants with a birthweight < 10th centile are thought to be FGR, while 70% of infants with a birthweight < 3rd centile are thought to be FGR. Critically, SGA is the most significant antenatal risk factor for a stillborn infant. Correct identification of SGA infants is associated with a reduction in the perinatal mortality rate. However, currently used tests, such as measurement of symphysis-fundal height, have a low reported sensitivity and specificity for the identification of SGA infants.

Objectives

The primary objective was to assess and compare the diagnostic accuracy of ultrasound assessment of fetal growth by estimated fetal weight (EFW) and placental biomarkers alone and in any combination used after 24 weeks of pregnancy in the identification of placental dysfunction as evidenced by either stillbirth, or birth of a SGA infant. Secondary objectives were to investigate the effect of clinical and methodological factors on test performance.

Search methods

We developed full search strategies with no language or date restrictions. The following sources were searched: MEDLINE, MEDLINE In Process and Embase via Ovid, Cochrane (Wiley) CENTRAL, Science Citation Index (Web of Science), CINAHL (EBSCO) with search strategies adapted for each database as required; ISRCTN Registry, UK Clinical Trials Gateway, WHO International Clinical Trials Portal and ClinicalTrials.gov for ongoing studies; specialist abstract and conference proceeding resources (British Library's ZETOC and Web of Science Conference Proceedings Citation Index). Search last conducted in October 2016.

Selection criteria

We included studies of pregnant women of any age with a gestation of at least 24 weeks if relevant outcomes of pregnancy (live birth/stillbirth; SGA infant) were assessed. Studies were included irrespective of whether pregnant women were deemed to be low or high

risk for complications or were of mixed populations (low and high risk). Pregnancies complicated by fetal abnormalities and multi-fetal pregnancies were excluded as they have a higher risk of stillbirth from non-placental causes. With regard to biochemical tests, we included assays performed using any technique and at any threshold used to determine test positivity.

Data collection and analysis

We extracted the numbers of true positive, false positive, false negative, and true negative test results from each study. We assessed risk of bias and applicability using the QUADAS-2 tool. Meta-analyses were performed using the hierarchical summary ROC model to estimate and compare test accuracy.

Main results

We included 91 studies that evaluated seven tests — blood tests for human placental lactogen (hPL), oestriol, placental growth factor (PLGF) and uric acid, ultrasound EFW and placental grading and urinary oestriol — in a total of 175,426 pregnant women, in which 15,471 pregnancies ended in the birth of a small baby and 740 pregnancies which ended in stillbirth. The quality of included studies was variable with most domains at low risk of bias although 59% of studies were deemed to be of unclear risk of bias for the reference standard domain. Fifty-three per cent of studies were of high concern for applicability due to inclusion of only high- or low-risk women.

Using all available data for SGA (86 studies; 159,490 pregnancies involving 15,471 SGA infants), there was evidence of a difference in accuracy ($P < 0.0001$) between the seven tests for detecting pregnancies that are SGA at birth. Ultrasound EFW was the most accurate test for detecting SGA at birth with a diagnostic odds ratio (DOR) of 21.3 (95% CI 13.1 to 34.6); hPL was the most accurate biochemical test with a DOR of 4.78 (95% CI 3.21 to 7.13). In a hypothetical cohort of 1000 pregnant women, at the median specificity of 0.88 and median prevalence of 19%, EFW, hPL, oestriol, urinary oestriol, uric acid, PLGF and placental grading will miss 50 (95% CI 32 to 68), 116 (97 to 133), 124 (108 to 137), 127 (95 to 152), 139 (118 to 154), 144 (118 to 161), and 144 (122 to 161) SGA infants, respectively. For the detection of pregnancies ending in stillbirth (21 studies; 100,687 pregnancies involving 740 stillbirths), in an indirect comparison of the four biochemical tests, PLGF was the most accurate test with a DOR of 49.2 (95% CI 12.7 to 191). In a hypothetical cohort of 1000 pregnant women, at the median specificity of 0.78 and median prevalence of 1.7%, PLGF, hPL, urinary oestriol and uric acid will miss 2 (95% CI 0 to 4), 4 (2 to 8), 6 (6 to 7) and 8 (3 to 13) stillbirths, respectively. No studies assessed the accuracy of ultrasound EFW for detection of pregnancy ending in stillbirth.

Authors' conclusions

Biochemical markers of placental dysfunction used alone have insufficient accuracy to identify pregnancies ending in SGA or stillbirth. Studies combining U and placental biomarkers are needed to determine whether this approach improves diagnostic accuracy over the use of ultrasound estimation of fetal size or biochemical markers of placental dysfunction used alone. Many of the studies included in this review were carried out between 1974 and 2016. Studies of placental substances were mostly carried out before 1991 and after 2013; earlier studies may not reflect developments in test technology.

PLAIN LANGUAGE SUMMARY

Blood tests in late pregnancy to identify small babies and those at risk of stillbirth

Background

Placental dysfunction describes when the placenta does not meet the demands of the growing baby; it may result in a baby that is smaller than expected or is stillborn. Currently, it is not easy to detect placental dysfunction before birth; ultrasound scans are most often used to identify small babies. However, tests can measure substances made by the placenta in mothers' blood and urine which may detect a placenta that is not functioning well. We aimed to find the best test to identify placental dysfunction.

What we did

We searched for studies in October 2016 and identified and total of 24,059 studies - with 91 of those studies providing us with information that we could include in this review. We looked at ultrasound scanning and six different tests of placental substances, including proteins and hormones. These studies involved 175,426 women in total of which 15,471 pregnancies ended in the birth of a small baby and 740 pregnancies which ended in stillbirth.

What we found

Of the 91 included studies, 86 had information on small babies, of which 18 also looked at stillbirth; another five studies only looked at stillbirth. The most accurate test for detecting a small baby was ultrasound scan to estimate a baby's weight. Of the substances measured in mother's blood, human placental lactogen (hPL), a hormone produced by the placenta during pregnancy, was the most accurate. There was only one study which looked at both ultrasound scanning and measurement of a placental substance. Placental growth factor (PLGF) was the most accurate test of a placental substance to identify a baby that would be stillborn; there were no studies of ultrasound scanning to detect a baby that would be stillborn. Tests of placental substances were better at identifying a baby at risk of stillbirth than detecting a small baby.

Other important information to consider

Biochemical tests of placental function versus ultrasound assessment of fetal size for stillbirth and small-for-gestational-age infants (Review)

Copyright © 2019 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Many of the studies included in this review were carried out between 1974 and 2016. Studies of placental substances were mostly carried out before 1991 and after 2013; earlier studies may not reflect developments in test technology. More studies are needed to find out whether a combination of ultrasound scans and mother's blood tests could improve identification of pregnancies which end in the birth of a small baby or in a stillborn baby. No studies were identified for this review that looked at the accuracy of ultrasound and blood tests used together.

SUMMARY OF FINDINGS

Summary of findings 1. Summary of findings table

Review question	To assess and compare the diagnostic accuracy of ultrasound assessment of fetal growth and placental biomarkers alone and in any combination used after 24 weeks of pregnancy in the identification of placental dysfunction as evidenced by either stillbirth or born small-for-gestational age (SGA).					
Population	Pregnant women of any reproductive age after 24 weeks' gestation with relevant outcomes of pregnancy recorded.					
Settings	All settings					
Numbers of studies, pregnancies, SGA births, and stillbirths	Ninety-one studies were included; 86 studies involving 159,490 pregnancies with 15,471 SGA infants and 23 studies involving 115,911 pregnancies with 851 stillbirths (18 of these also looked at SGA).					
Index tests	Human placental lactogen (hPL), oestriol (in blood or urine), placental grading, placental growth factor (PLGF), ultrasound estimated fetal weight (EFW), uric acid					
Reference standards	Small for gestational age (SGA), stillbirth					
Study limitations	Full-text papers of some studies were unobtainable; comparisons of testing at different gestational ages were not possible; comparisons of tests in the same population were not possible; broad definitions of patient risk were used.					
Conclusions	Biochemical markers of placental dysfunction alone are not sufficiently accurate to identify pregnancies ending in SGA or stillbirth. Studies combining ultrasound and placental biomarkers are needed to determine whether this approach improves diagnostic accuracy.					
Test strategy	Studies	Number of women (cases)	Sensitivity (95% CI)^a	Specificity (95% CI)^a	Number of missed cases in a hypothetical cohort of 1000 pregnant women^b	Number of false positives in a hypothetical cohort of 1000 pregnant women^b
SGA defined as birthweight $\leq$ 10th centile						
Comparison of ultrasound EFW and biochemical tests						
Ultrasound EFW	32	51,702 (6169)	0.74 (0.64 to 0.83)	0.88	50 (32 to 68)	97
Placental grading	12	4940 (520)	0.24 (0.15 to 0.36)		144 (122 to 161)	

hPL	20	3486 (624)	0.39 (0.30 to 0.49)		116 (97 to 133)	
Serum oestriol	9	2773 (373)	0.35 (0.28 to 0.43)		124 (108 to 137)	
Urinary oestriol	9	92,406 (7076)	0.33 (0.20 to 0.50)		127 (95 to 152)	
PIGF	8	6997 (1029)	0.24 (0.15 to 0.38)		144 (118 to 161)	
Uric acid	8	2884 (605)	0.27 (0.19 to 0.38)		139 (118 to 154)	
Combination of biochemical tests						
Serum oestriol and hPL	1	88 (9)	0.56 (0.21 to 0.86)	0.95 (0.88 to 0.99)	45 (14 to 91)	45
SGA3 defined as birthweight < 3rd centile						
EFW ^c	3	5678 (212)	0.66 (0.56 to 0.76)	0.87 (0.80 to 0.91)	16 (11 to 20)	124
PIGF	2	1861 (101)	-	-	-	
EFW & PIGF	1	343 (52)	0.69 (0.55 to 0.81)	0.72 (0.67 to 0.77)	42 (35 to 50)	237
Stillbirth						
Comparison of biochemical tests^d						
hPL	6	544 (36)	0.76 (0.55 to 0.90)	0.78	4 (2 to 8)	216
Urinary oestriol	7	92,186 (651)	0.62 (0.58 to 0.66)		6 (6 to 7)	
PIGF	4	5894 (16)	0.93 (0.78 to 0.98)		2 (0 to 4)	
Uric acid	4	2063 (37)	0.53 (0.21 to 0.83)		8 (3 to 13)	
Ultrasound						
Placental grading	3	15,236 (114)	-	-	-	-

^aFor SGA and stillbirth, the sensitivities were estimated from the SROC curves at fixed values of specificity that correspond to the median of the specificities reported in the studies included in each comparative meta-analysis. For SGA3, the estimates are the pooled sensitivity and specificity from the only meta-analysis that was possible. All other estimates are the sensitivity and specificity from a single study.

^bTo calculate the number of missed cases and false positives for SGA and stillbirth in a hypothetical cohort of 1000 pregnant women, we used the median prevalence of 19% and 1.7% from the studies included in the comparative meta-analysis of SGA and stillbirth, respectively. For SGA3, we used the median prevalence of 4.6% to calculate the number of missed cases from the pooled estimates of sensitivity and specificity from the only meta-analysis that was possible. For all single studies, we used the prevalence derived from the study.

^cMeta-analysis was not possible for EFW. For the three studies, the sensitivities were between 0.58 and 0.77, and the specificities were between 0.79 and 0.91.

^dThis comparison was limited to only biochemical tests because there were several tests and only three placental grading studies. Including placental grading increased model complexity and made the comparative meta-analysis model impossible to fit. Therefore, placental grading was excluded. The sensitivities of the three placental grading studies ranged between 0.35 and 0.69, and the specificities between 0.14 and 0.94.

BACKGROUND

Stillbirth affects 2.6 million pregnancies worldwide each year (Lawn 2016). Whilst the majority of cases occur in low- and middle-income countries, stillbirth remains an important clinical issue for high-income countries (HICs) - with both the UK and the USA reporting rates above the mean for HICs (Flenady 2016).

In HICs, the most frequently reported association with stillbirth is placental dysfunction, which may be clinically evident as fetal growth restriction (FGR), small-for-gestational-age (SGA) infants, placental abruption or hypertensive disorders of pregnancy. Placental abnormalities are noted in 11% to 65% of stillbirths (Ptacek 2014). Identification of FGR is difficult in utero and even after birth, with SGA being most commonly used as a surrogate measure (Worton 2014). The degree of SGA is associated with the likelihood of FGR; 30% of infants with a birthweight < 10th centile are thought to be FGR, while 70% of infants with a birthweight < 3rd centile are thought to be FGR. Critically, SGA is the most significant antenatal risk factor for a stillborn infant (Flenady 2011; Gardosi 2013; McCowan 2007). Correct identification of SGA infants is associated with a reduction in the perinatal mortality rate (Gardosi 2013). However, currently used tests, such as measurement of symphysis-fundal height, have a low reported sensitivity and specificity for the identification of SGA infants (RCOG 2014).

Due to the importance of the placenta in FGR and stillbirth there is growing interest in antenatal placental evaluation in an attempt to identify pregnancies at increased risk of stillbirth or fetal compromise (Heazell 2015a). A systematic review of biochemical tests of placental function found insufficient evidence to conclude whether these interventions had any effect on perinatal mortality or fetal compromise (Heazell 2015b). In contrast, a single trial of placental grading assessed by ultrasound demonstrated reduced perinatal mortality (Proud 1987). Systematic reviews of other methods employed to identify fetal compromise such as ultrasound assessment of fetal growth or umbilical artery Doppler (measurement of blood flow through the umbilical artery) in late pregnancy have also found insufficient evidence to conclude whether these interventions reduce perinatal mortality in a low-risk maternity population (Alfirevic 2015; Bricker 2015), although both are effective in women deemed to be at high risk of pregnancy complications (Alfirevic 2017). The efficacy of umbilical artery Doppler in high-risk populations may be due to its prognostic accuracy; a systematic review found this test predicted SGA infants with a positive likelihood ratio of 3.76 and stillbirth with a positive likelihood ratio of 4.37 (Morris 2011).

Two components are necessary to reduce perinatal mortality and minimise unwarranted intervention. Firstly, the test must accurately identify fetal compromise and secondly, the intervention must be effective in preventing the adverse outcome. There is now strong evidence that planned delivery (by induction of labour) after 37 weeks of pregnancy is associated with a reduction in perinatal mortality (Stock 2012). Therefore, the most accurate test to identify fetal compromise needs to be determined so that it may be combined with planned delivery where appropriate.

Target condition being diagnosed

The target condition of interest is placental dysfunction – which describes the condition in which the placenta does not meet the demands of the fetus (Heazell 2015a). As with other organ

dysfunction, there are multiple pathways that can result in placental dysfunction including vascular, inflammatory, infective and genetic disorders. These various processes may lead to changes in placental structure and/or function that may lead to two clinical outcomes i) stillbirth or ii) the birth of an SGA infant. As placental dysfunction cannot easily be quantified, this review will use these two clinical outcomes as the target conditions of interest.

Index test(s)

This review evaluated tests used in late pregnancy (after 24 weeks) to identify pregnancies with placental dysfunction to inform decisions to continue with the pregnancy or institute intervention. Tests that were included in this review assessed placental structure or biochemical function by one or more of ultrasound scan or measurement of placental products in maternal blood (plasma or serum) or urine.

Biochemical tests of placental function measure placental products (proteins, peptides, metabolites, hormones) in maternal biofluids (serum, plasma, urine); it is hypothesised that levels of such products in maternal fluids reflect endocrine and metabolic functions of the placenta. Many placental products can be detected in maternal biofluids including protein hormones: human chorionic gonadotrophin (hCG), human placental lactogen (hPL), human placental growth hormone (hPGH), placental growth factor (PlGF), placental protein-13 (PP-13), pregnancy specific glycoproteins and steroid hormones including oestrogens and progesterone with their related metabolites. Ultrasonography has been used to measure the size, shape, and echotexture of the placenta; the majority of such studies have used 2D ultrasound to evaluate placental morphology, although newer studies have utilised 3D techniques.

Clinical pathway

Antenatal care differs between countries; the clinical pathway described here applies to the UK and follows guidance from the Royal College of Obstetricians and Gynaecologists (RCOG 2014) and the National Institute for Health and Social Care Excellence (NICE 2008).

Prior test(s)

Currently, in the UK women are grouped into high risk and low risk for SGA in early pregnancy at the booking-visit by assessing a woman's past medical history, obstetric history and risk factors for an SGA infant (RCOG 2014). All women are offered screening for Down's syndrome (which is currently based on measurement of nuchal translucency by ultrasound scan and measurement of serum analytes between 11 and 13 + 6 weeks of pregnancy) and for fetal anomaly (by ultrasound scan from 18 to 20 + 6 weeks).

In clinical practice, placental dysfunction is suspected by identification of an SGA infant. However, testing for SGA currently depends upon the risk status of the woman (RCOG 2014). The National Institute for Health and Care Excellence do not recommend routine measurement of fetal growth by ultrasound scan in late pregnancy (NICE 2008). Fetal growth is assessed in women deemed to be at low risk of an SGA infant by measurement of symphysis-fundal height with a tape measure (RCOG 2014). Women at increased risk of SGA are recommended to have a uterine artery Doppler (to assess blood flow through both uterine arteries) at 20 weeks' gestation and regular scans to measure fetal

biometry with assessment of liquor volume and umbilical artery Doppler. Umbilical artery Doppler is the most frequently employed test to predict fetal outcome; the relationship between umbilical artery Doppler indices and placental function is not clear. In addition to recommendations for the diagnosis and management

of an SGA fetus, ultrasound assessment of fetal growth, liquor volume and umbilical artery Doppler are recommended following maternal presentation with reduced fetal movements, as this may be a symptom of placental insufficiency (RCOG 2011). The current clinical pathway is shown in Figure 1.

Figure 1. Current clinical pathway and three proposed uses of a placental function test. Currently, women are screened for a small-for-gestational age fetus (as a proxy for placental dysfunction) using symphysis-fundal height and maternal awareness of fetal movements. Women deemed to be at increased risk are screened using ultrasound measurement of fetal biometry. We propose three different clinical pathways for placental function tests. Firstly, they could be used as an additional test when Doppler measurements are normal. They could be used in combination with currently used tests, and finally they could be used as a triage test to differentiate infants who are constitutionally small from those with placental dysfunction. Although treatment decisions would be tailored to individual cases, a positive test would be expected to lead to increased surveillance or intervention (planned delivery) and a negative test would lead to continuing with the pregnancy.

When an SGA infant is identified by tests, clinical management is dependent upon gestation. Prior to 37 weeks' gestation, identification of SGA prompts further assessment of fetal well-being, primarily by measurement of Doppler waveforms in the umbilical artery, but may also include the middle cerebral artery and ductus venosus. Delivery is recommended when evidence of fetal compromise is identified (RCOG 2014). After 39 weeks' gestation, delivery of the baby may be offered as this is associated with a reduction in stillbirth, avoids potential hazards of early term

birth (MacKay 2010), and is not associated with an increase in obstetric intervention (Stock 2012).

There are currently no routinely used measures of placental function after 16 weeks of pregnancy. There is evidence that measurement of placental analytes as part of screening for aneuploidy may identify fetuses at high risk of early-onset FGR (Smith 2002; Smith 2006). Assessment of these analytes is incorporated into the current clinical pathway (RCOG 2014); women with low pregnancy-associated plasma protein A (PAPP-A) levels are managed as high risk for SGA. Therefore, we wish to focus

on placental tests performed in late pregnancy (after 24 weeks' gestation).

Role of index test(s)

Due to the established use of ultrasound in obstetric practice, we envisage that additional tests of placental function would most likely be added to an ultrasound measurement of fetal size rather than replacing it (Figure 1); this is certainly true of the intervention trials of placental assessment (by biochemical tests) that have been conducted (Duenholter 1976; Heazell 2013; Sharf 1984). It is hypothesised that the addition of a placental function test to an ultrasound scan would improve identification of an SGA infant and consequently focus intervention on those pregnancies at the greatest risk of stillbirth or fetal compromise, thereby reducing the burden of perinatal mortality and morbidity. It is also possible that a placental function test could be used to triage infants who were SGA to identify which were constitutionally small and which had placental dysfunction. This would allow the pregnancy to continue in otherwise healthy constitutionally small infants, reducing unnecessary intervention.

The importance of specific aspects of test performance will depend upon the context in which it is used in late pregnancy. From the perspective of reducing perinatal mortality and morbidity in a high-risk population (e.g. triage of women with a small baby on ultrasound scan or women presenting with reduced fetal movements, Figure 1), a false negative test would be more harmful than a false positive test as pregnant women may be deprived of further monitoring or intervention which may mitigate some of the increased risk. In women at low risk of SGA or stillbirth then a false negative test would mean that the mother continued upon the pathway of care she would have otherwise received, whereas a false positive result would mean she may be exposed to monitoring or intervention which was unnecessary, which may have negative medical and economic consequences. Thus, it is important to consider the clinical group being studied and how this impacts upon test performance.

Alternative test(s)

Presently, there are no tests in widespread clinical use that directly assess placental biochemical function. Umbilical artery Doppler is often used in clinical practice to identify placental dysfunction but has not been included in this analysis as a systematic review and meta-analysis has already been conducted (Morris 2011).

Rationale

There are several tests of placental structure and function. Systematic reviews of the measurement of biochemical placental factors and the effectiveness of ultrasound in late pregnancy found that few tests of placental structure or function have been evaluated in robust intervention studies (Bricker 2015; Heazell 2015a). This review aims to identify and evaluate tests of placental structure and function, not restricted to those evaluated in intervention studies, to determine which measurement(s) have the greatest diagnostic accuracy for detection of placental dysfunction leading to stillbirth and SGA. The most accurate test(s) can then be taken forward into intervention studies to determine whether performing investigations can reduce perinatal morbidity or mortality.

OBJECTIVES

The primary objective of this review was to assess and compare the diagnostic accuracy of ultrasound assessment of fetal growth and placental biomarkers alone and in any combination used after 24 weeks of pregnancy in the identification of placental dysfunction as evidenced by either stillbirth or born small-for-gestational age (SGA). Accuracy is described by the proportion of fetuses who are subsequently stillborn or who have an SGA baby detected by a positive test result (the presence of placental dysfunction) (sensitivity) and by the proportion of fetuses that have an uncomplicated pregnancy following a negative index test result (absence of placental dysfunction) (specificity).

Secondary objectives

We investigated the effect of clinical (patient and test characteristics) and methodological factors (study design, threshold used to define SGA) on test performance. The clinical factors include patient group (low-risk or high-risk pregnancies), gestation at measurement, ethnicity, maternal age and method of testing. With regard to methodological variation, studies may include an intervention (delivery or additional fetal surveillance for index test positive cases) which impacts on the outcome; therefore we assessed whether this is a source of heterogeneity.

METHODS

Criteria for considering studies for this review

Types of studies

Presently, there are no effective interventions to reverse placental dysfunction *in utero*. This means that an intervention cannot be employed to reverse the small-for-gestational age (SGA) phenotype following a positive test result. Delivery may be indicated, although at earlier gestations this does not affect perinatal mortality (GRIT 2003).

We included prospective and retrospective cross-sectional or cohort studies in which all women received one or more index tests and the outcome of their pregnancy was known. Case-control studies were excluded.

We included studies which measured index tests on one occasion (cross-sectional design).

We excluded studies where it was not possible to derive a 2 x 2 table of the number of true positives, false positives, false negatives and true negatives, or studies that reported preliminary experimental findings, i.e. laboratory-based studies.

Participants

We included studies of pregnant women after 24 weeks' gestation that recorded relevant outcomes of pregnancy (live birth/stillbirth; SGA infant).

We included studies of pregnant women of any reproductive age, who were deemed to be low or high risk for complications (e.g. who had pre-existing medical disorders or previous stillbirth) or studies of mixed populations (of low and high risk for complications).

We excluded pregnancies complicated by fetal abnormalities, as they often have a higher risk of stillbirth from non-placental causes. We excluded studies of women with multi-fetal pregnancies.

Index tests

We included, but were not restricted to, the following index tests of placental biochemical function, placental structure or assessment of fetal biometry to identify an SGA infant:

- human placental lactogen (hPL) in maternal urine/blood;
- oestriol in maternal urine/blood;
- placental growth factor (PlGF) in maternal blood;
- ultrasound assessment of placental echogenicity;
- ultrasound assessment of fetal size.

With regard to biochemical tests, we included assays that were performed using different techniques, including: immunoassay, enzyme-linked immunosorbent assay (ELISA), chromatography or point of care test in any combination and at any threshold used to determine test positivity. Examples of current commercially available tests are listed in [Appendix 1](#).

Target conditions

The target conditions were stillbirth and delivery of a SGA infant at the centile or threshold used by each study, as clinical manifestations of placental dysfunction.

Reference standards

The outcome of pregnancy was considered as the reference standard. A “positive” result was either i) a stillbirth – an infant born with no signs of life after 24 weeks’ gestation, or ii) a birthweight classified as SGA. A “negative” result was a live birth after 24 weeks’ gestation or a birthweight classified as appropriate for gestational age.

The classification of SGA was determined according to the definition used in the study. Where possible, the definition of an infant with a birthweight $\leq$ 10th centile using a customised birthweight calculator was used ([Clausson 2001](#)). Where this was not possible, the definition of SGA from the manuscript was used and recorded. The effects of different definitions of SGA were addressed as a potential source of heterogeneity.

Search methods for identification of studies

We conducted a comprehensive search for existing systematic reviews and primary studies relevant to the prevention of adverse pregnancy outcome in women at increased risk of stillbirth by detecting placental dysfunction. A scoping search was undertaken in the bibliographic databases MEDLINE, MEDLINE In Process, Embase, the Cochrane Library (CDSR, DARE, HTA, NHS EED and Central Register of Controlled Trials (CENTRAL) databases), HTA and relevant web sites in order to identify existing reviews and to gauge the nature and number of relevant studies to inform the protocol.

Electronic searches

We developed full search strategies based on the scoping searches, expert advice, and consultation with the Cochrane Pregnancy and Childbirth Group’s Information Specialist. Search strategies included a combination of text words and index terms.

Methodological search filters for diagnostic test accuracy were avoided as they have been shown to miss relevant studies ([Whiting 2011a](#)). We did not apply any language or date restrictions. We searched the following sources:

- bibliographic databases - MEDLINE, MEDLINE In Process and Embase via Ovid, Cochrane (Wiley) CENTRAL, Science Citation Index (Web of Science), CINAHL (EBSCO) with search strategies adapted for each database as required;
- [ISRCTN Registry](#), [UK Clinical Trials Gateway](#), WHO International Clinical Trials Portal (ICTRP) and [ClinicalTrials.gov](#) for ongoing studies;
- specialist abstract and conference proceeding resources (British Library’s ZETOC and Web of Science Conference Proceedings Citation Index).

Searches were last conducted on 27 October 2016. The full search strategy is provided in [Appendix 2](#).

Searching other resources

We checked citation lists of included studies and relevant reviews. We examined grey literature by searching websites of companies producing biochemical tests of placental function ([Alere 2015](#); [Perkin Elmer 2015](#); [Roche 2015](#)). We also undertook consultation with experts in the field to access relevant unpublished data.

Data collection and analysis

We used the [methods](#) advocated by the *Cochrane Screening and Diagnostic Test Methods Group*.

Selection of studies

Two review authors (DH and AH) independently screened the titles and abstracts of all studies identified by the search strategy. We obtained full-text versions of all potentially relevant studies. Two review authors (DH and AH) independently assessed studies for inclusion using pre-specified inclusion criteria stated earlier. We included studies of pregnant women after 24 weeks’ gestation that recorded relevant outcomes of pregnancy (live birth/stillbirth; SGA infant), and presented data to construct a 2 x 2 table. We resolved any disagreement between the two review authors or by discussion with a third party (CD) if needed. Reasons for study exclusion were documented.

Data extraction and management

We developed a customised form to ensure reproducible collection of data items. Data collection was piloted on five manuscripts then reviewed by the review authors. Data were extracted independently by two review authors (DH and AH). We resolved discrepancies, where they occurred, through discussion or if required we consulted a third author (CD or MW), drawing on clinical and methodological expertise in the team as appropriate to the content of the query. We extracted characteristics of participants, index tests or test combinations (including thresholds used), and details of the reference standard in terms of pregnancy outcome (live birth or stillbirth) and whether the infant was SGA. For studies that reported data at multiple thresholds for a test, we extracted a 2 x 2 table at each reported threshold. Where possible we recorded the frequency of obstetric intervention and infant admission to neonatal intensive care. If reported, we also recorded data on outcomes including harms of testing, need for further testing, and

the effects of the test. We did not address women's experiences of testing, caregivers' satisfaction with testing or economic evaluation of testing as this is beyond the scope of this review.

We attempted to contact the authors of included studies where information considered key to assessment of methodological quality, investigation of heterogeneity, or completion of a 2 x 2 table was unclear or missing. Studies published only as conference abstracts were followed up to identify whether a subsequent full paper had been published.

Assessment of methodological quality

We used the QUADAS-2 tool (Whiting 2011b) to assess the risk of bias and applicability of included studies. We tailored the tool to our review question using the operational criteria detailed in Appendix 3 to answer signalling questions and make the overall judgement of risk of bias and applicability concerns for each domain of the tool. Two review authors (DH and AH) assessed each included study separately. We resolved differences in assessment through discussion and if required, by discussion with a third person (CD). We assessed each criteria in QUADAS-2 as "yes", "no" or "unclear" and summarised the results graphically or in tables.

We included all signalling questions of QUADAS-2 assessment including the time interval between testing and the outcome and any intervention as these may alter the outcome. We have operationalised the domains of the QUADAS-2 tool for the clinical context of this review. For example, the domain concerning patient selection was amended to reflect review exclusion criteria including women with multiple pregnancies or with fetal abnormalities. However, other criteria that might be expected to alter the accuracy of tests in universal populations (e.g. ethnicity, maternal age and income) would be inappropriate exclusions. Studies restricted to specific high-risk groups, e.g. maternal hypertension, will reduce the applicability of review findings. We also tailored the target condition domain to assess the quality of measures of SGA used in studies, some of which may not be related to gestation, e.g. low birthweight (< 2.5 kg). Studies using a threshold which alters with sex and gestation, e.g. individualised birthweight centile, were rated more highly than those which did not.

Statistical analysis and data synthesis

From here on we use the term SGA for SGA defined as birthweight ≤ 10th centile, and SGA3 for SGA defined as birthweight < 3rd centile. We performed separate analyses for each target condition (SGA, SGA3 and stillbirth). For each test and target condition, estimates of sensitivity and specificity from each study were plotted in receiver operating characteristic (ROC) space and forest plots for preliminary investigations of the data. Since studies used different thresholds to determine test positivity, we performed meta-analyses using the hierarchical summary ROC (HSROC) model (Rutter 2001) to estimate and compare the SROC curves of the tests. Methods that allow joint synthesis of sensitivities and specificities at multiple thresholds have been proposed, but are not yet used routinely in practice and require further evaluation before they can be used in diagnostic test accuracy reviews (Ensor 2018; Riley 2015; Steinhauser 2016). As such, where a study reported multiple thresholds for a test, we selected the threshold most frequently reported across studies so that only one 2 x 2 table was included in a meta-analysis. In separate analyses of each test, where studies reported common thresholds for the test, we estimated summary

sensitivities and specificities using functions of HSROC model parameters.

Before performing meta-analyses to compare test accuracy, we performed meta-analysis of each test separately for preliminary investigation of the shape of the SROC curve of each test and to explore if assuming common variances across tests for the random effects would be reasonable. The main test comparison was an indirect comparison pooling all relevant studies that assessed at least one of the index tests. In secondary analyses, we performed direct comparisons by restricting the analyses to only studies that compared tests head-to-head in the same study population. This analytical strategy was adopted because of the paucity of comparative studies of diagnostic accuracy (Takwoingi 2013). We limited the indirect comparison to only tests with at least four studies because of potential model complexity given the number of tests included and number of model parameters to be estimated. For direct comparisons, we performed pairwise comparisons of tests. Test comparisons were performed by adding a covariate for test type to the HSROC model to estimate differences in accuracy, threshold, and/or shape of SROC curves. When there were adequate data, we also allowed the variance parameters for accuracy and threshold to depend on test type, i.e. differences in accuracy and threshold modelled as random effects. We assessed the statistical significance of differences between tests using likelihood ratio tests comparing models with and without the covariate terms. The NLMIXED procedure in the SAS software package (version 9.4; SAS Institute, Cary, NC, USA) was used for meta-analyses.

To quantify differences in accuracy between tests, we computed ratios of diagnostic odds ratios when SROC curves were symmetric or a common shape was assumed. Using the estimate statement within NLMIXED, we also estimated sensitivities along the SROC curves at fixed values of specificity that correspond to the median and interquartile range of specificities from the studies included in the comparative meta-analysis. We used these values along with the median and interquartile range of the prevalence estimated from the studies to compute numbers of missed cases and false positives in a hypothetical cohort of 1000 pregnant women. We used these frequencies to illustrate the accuracy of the tests in absolute terms.

Investigations of heterogeneity

We initially examined heterogeneity between studies by visually inspecting forest plots of sensitivity and specificity and SROC plots. Where a sufficient number of studies assessed the same index test and there were at least four studies per subgroup of a categorical covariate, we performed meta-regression by adding the potential source of heterogeneity as a covariate to the HSROC model. We assessed the effect of the covariate on test accuracy by using likelihood ratio tests to compare models with and without the covariate terms.

Sensitivity analyses

We planned to perform sensitivity analyses by restricting analyses to studies that:

- were without an intervention that may have altered outcome;
- were at low risk of bias in each of the four domains of the QUADAS-2 tool;

- specifically described histological evidence of placental insufficiency.

These analyses were not possible due to limited data.

Assessment of reporting bias

We did not undertake any formal assessment of reporting bias in our review due to current uncertainty about how to assess reporting bias in diagnostic accuracy reviews, especially in the presence of heterogeneity ([Macaskill 2010](#)).

RESULTS

Results of the search

Our literature searches identified a total of 24,059 papers before duplicates were removed ([Figure 2](#)). After initial screening based on title and abstract, we obtained full-text copies of 471 papers, of which 91 were included. The included studies evaluated seven tests — blood tests for human placental lactogen (hPL), oestriol, placental growth factor (PLGF) and uric acid, ultrasound EFW and placental grading and urinary oestriol — in a total of 175,426 pregnant women, in which 15,471 pregnancies ended in the birth of a small baby and 740 pregnancies which ended in stillbirth.

Figure 2. PRISMA flow diagram for selection of studies.

Figure 2. (Continued)

In total, seven different tests were evaluated, as well as two combinations of these tests. Included studies are described in [Table 1](#). We described the studies that were excluded after full-text assessment in [Characteristics of excluded studies](#). Some papers could not be obtained, index tests used in these papers are stated in [Characteristics of studies awaiting classification](#) where possible.

Methodological quality of included studies

[Figure 3](#) shows the summary risk of bias and applicability concerns for included studies. Risk of bias and applicability concerns for individual index tests are in [Appendix 5](#). The quality of included studies was judged to be mostly high, with most domains at low risk of bias, although risk of bias for the reference standard domain

was mostly unclear (59% of studies) due to definitions of small-for-gestational age (SGA) not being reported or uncertainty as to whether index test results were blinded. In terms of applicability concerns; 53% of studies were at high concern for patient selection, mostly due to inclusion of only high- or low-risk women rather than an unselected population of pregnant women where both are likely to be included. Fourteen studies (15%) were of high concern regarding the reference standard due to differing definitions of SGA, for example if gestation was not taken into account or centiles were based on the study population only. Eighty-four per cent of studies were judged to be at low risk of bias for flow and timing as testing was mostly performed with an appropriate interval between testing and delivery.

Figure 3. Risk of bias and applicability concerns graph: review authors' judgements about each domain presented as percentages across included studies. E3 = oestriol; EFW = estimated fetal weight; hPL = human placental lactogen; PlGF = placental growth factor. Each bar shows the number of studies in each category for a domain. The index test domain was evaluated separately for each test. Of the 91 included studies, 9 studies evaluated more than one test and so the numbers of studies shown for the 7 index test domains do not add up to 91 across tests.

Findings

Small-for-gestational age (SGA) (birthweight $\leq$ 10th centile)

The findings are summarised in [Summary of findings 1](#). SGA was the outcome assessed in 86 studies involving 159,490 pregnancies (included 15,471 SGA infants). The study specific estimates of

sensitivity and specificity for each test are shown for structural tests (ultrasound assessment of fetal size and placental grading) in [Figure 4](#), and for biochemical tests (human placental lactogen (hPL), serum oestriol, urinary oestriol, placental growth factor (PlGF), uric acid, and a combination of serum oestriol and hPL) in [Figure 5](#).

Figure 4. Forest plot of structural tests for identifying small-for-gestational age (birthweight $\leq$ tenth centile) infants. EFW = estimated fetal weight; FN = false negative; FP = false positive; SGA = small-for-gestational-age; TN = true negative; TP = true positive. For each test, the forest plot shows the estimates of sensitivity and specificity from each study, the threshold used to define test positivity, risk of SGA pregnancy and whether there was an intervention that may have altered outcome. Studies are sorted by threshold, risk and intervention status.

EFW and SGA

Placental grading and SGA

Figure 5. Forest plot of biochemical tests for identifying small-for-gestational-age (birthweight $\leq$ tenth centile) infants. E3 = oestriol; FN = false negative; FP = false positive; hPL = human placental lactogen; PlGF = placental growth factor; SGA = small-for-gestational-age; TN = true negative; TP = true positive; UA = uric acid. For each test, the forest plot shows the estimates of sensitivity and specificity from each study, the threshold used to define test positivity, risk of SGA pregnancy and whether there was an intervention that may have altered outcome. Studies are sorted by threshold, risk and intervention status.

hPL and SGA

Study	TP	FP	FN	TN	Threshold	Risk	Intervention	Sensitivity (95% CI)	Specificity (95% CI)
Nisbet 1982	25	13	46	82	-2SD	High	Unknown	0.35 [0.24, 0.47]	0.86 [0.78, 0.93]
Howell 1985	12	44	38	407	2.5µg/ml	Mixed	Unknown	0.24 [0.13, 0.38]	0.90 [0.87, 0.93]
Obiekwe 1983	5	50	24	443	4.8µg/ml	Mixed	Unknown	0.17 [0.06, 0.36]	0.90 [0.87, 0.92]
Granat 1977	10	4	0	15	4µg/ml	High	No	1.00 [0.69, 1.00]	0.79 [0.54, 0.94]
Zhang 1990	25	42	13	41	4µg/ml	High	No	0.66 [0.49, 0.80]	0.49 [0.38, 0.61]
Gohari 1978	25	11	13	62	5µg/ml	High	Unknown	0.66 [0.49, 0.80]	0.85 [0.75, 0.92]
Nice 2016	10	5	13	49	<0.8MoM	Mixed	No	0.43 [0.23, 0.66]	0.91 [0.80, 0.97]
Altman 1978	3	4	3	0	Abnormal value	High	Unknown	0.50 [0.12, 0.88]	0.00 [0.00, 0.60]
Westergaard 1984	15	38	13	326	Abnormal value	Mixed	Unknown	0.54 [0.34, 0.72]	0.90 [0.86, 0.93]
Kunz 1976	8	16	7	52	Below fifth centile	High	Unknown	0.53 [0.27, 0.79]	0.76 [0.65, 0.86]
Klebe 1990	1	0	2	10	Below fifth centile	High	Unknown	0.33 [0.01, 0.91]	1.00 [0.69, 1.00]
Marin 1979	11	6	2	28	Below fifth centile	Mixed	Unknown	0.85 [0.55, 0.98]	0.82 [0.65, 0.93]
Odendaal 1981	35	16	8	18	Below tenth centile	High	No	0.81 [0.67, 0.92]	0.53 [0.35, 0.70]
Steiner 1991	25	17	43	28	Below tenth centile	High	Unknown	0.37 [0.25, 0.49]	0.62 [0.47, 0.76]
Trudinger 1979	10	4	15	30	Below tenth centile	High	Unknown	0.40 [0.21, 0.61]	0.88 [0.73, 0.97]
Sagen 1984	28	6	12	28	Below tenth centile	High	Yes	0.70 [0.53, 0.83]	0.82 [0.65, 0.93]
Siebert 1974	1	2	10	54	Below tenth centile	Low	Yes	0.09 [0.00, 0.41]	0.96 [0.88, 1.00]
Chard 1985	13	29	26	324	Below tenth centile	Mixed	Unknown	0.33 [0.19, 0.50]	0.92 [0.88, 0.94]
Lilford 1983	15	41	37	429	Below tenth centile	Mixed	Unknown	0.29 [0.17, 0.43]	0.91 [0.88, 0.94]
Sperml 1989	5	3	17	85	Below tenth centile	Mixed	Unknown	0.23 [0.08, 0.45]	0.97 [0.90, 0.99]

E3 and SGA

Study	TP	FP	FN	TN	Threshold	Risk	Intervention	Sensitivity (95% CI)	Specificity (95% CI)
Palo 1987	9	0	31	50	-2SD	High	Unknown	0.23 [0.11, 0.38]	1.00 [0.93, 1.00]
Nisbet 1982	9	4	37	53	-2SD	High	Unknown	0.20 [0.09, 0.34]	0.93 [0.83, 0.98]
Nielsen 1985	11	40	50	917	Below 2.5th centile	Mixed	No	0.18 [0.09, 0.30]	0.96 [0.94, 0.97]
Sperml 1989	2	3	20	85	Below fifth centile	Mixed	Unknown	0.09 [0.01, 0.29]	0.97 [0.90, 0.99]
Odendaal 1981	13	3	17	20	Below tenth centile	High	No	0.43 [0.25, 0.63]	0.87 [0.66, 0.97]
Cedard 1979	9	13	8	34	Below tenth centile	High	Unknown	0.53 [0.28, 0.77]	0.72 [0.57, 0.84]
Sagen 1984	22	6	18	28	Below tenth centile	High	Yes	0.55 [0.38, 0.71]	0.82 [0.65, 0.93]
Gerhard 1986	18	71	60	720	Below tenth centile	Mixed	No	0.23 [0.14, 0.34]	0.91 [0.89, 0.93]
Chard 1985	11	32	28	321	Below tenth centile	Mixed	Unknown	0.28 [0.15, 0.45]	0.91 [0.87, 0.94]

Urinary E3 and SGA

Study	TP	FP	FN	TN	Threshold	Risk	Intervention	Sensitivity (95% CI)	Specificity (95% CI)
Weerasinghe 1977	36	98	9	184	-2SD	High	Unknown	0.80 [0.65, 0.90]	0.65 [0.59, 0.71]
Steiner 1991	9	17	59	28	8mg at 30w to 12mg at 40w (per 24h)	High	Unknown	0.13 [0.06, 0.24]	0.62 [0.47, 0.76]
Beischer 1991	1454	5621	3936	61051	8mg at 30w to 12mg at 40w (per 24h)	High	Yes	0.27 [0.26, 0.28]	0.92 [0.91, 0.92]
Fliegner 1979	22	57	15	235	8mg at 30w to 12mg at 40w (per 24h)	Mixed	No	0.59 [0.42, 0.75]	0.80 [0.75, 0.85]
Oats 1979	497	2064	894	15664	8mg at 30w to 12mg at 40w (per 24h)	Mixed	Unknown	0.36 [0.33, 0.38]	0.98 [0.88, 0.99]
Chew 1976	0	0	15	28	Below 2.5th centile	High	No	0.00 [0.00, 0.22]	1.00 [0.88, 1.00]
Kunz 1976	10	12	5	56	Below fifth centile	High	Unknown	0.67 [0.38, 0.88]	0.82 [0.71, 0.91]
Odendaal 1981	6	1	22	17	Below tenth centile	High	No	0.21 [0.08, 0.41]	0.94 [0.73, 1.00]
Campbell 1972	46	34	41	163	Unknown	High	Unknown	0.53 [0.42, 0.64]	0.83 [0.77, 0.88]

PlGF and SGA

Study	TP	FP	FN	TN	Threshold	Risk	Intervention	Sensitivity (95% CI)	Specificity (95% CI)
Shawkat 2015	18	43	88	112	12pg/ml	High	No	0.17 [0.10, 0.26]	0.72 [0.65, 0.79]
Benton 2016	87	70	72	182	12pg/ml	High	No	0.55 [0.47, 0.63]	0.72 [0.66, 0.78]
Nice 2016	4	4	19	49	12pg/ml	Mixed	No	0.17 [0.05, 0.39]	0.92 [0.82, 0.98]
Molvarec 2013	16	25	6	42	3.9:1	High	No	0.73 [0.50, 0.89]	0.63 [0.50, 0.74]
Chaiworapongsa 2013	38	184	70	977	<0.3MoM	Mixed	Unknown	0.35 [0.26, 0.45]	0.84 [0.82, 0.86]
Griffin 2015	47	40	145	360	Below fifth centile	High	Unknown	0.24 [0.19, 0.31]	0.90 [0.87, 0.93]
Kienast 2016	29	80	11	226	Below fifth centile	Mixed	Unknown	0.72 [0.56, 0.85]	0.74 [0.69, 0.79]
Valino 2016	48	150	331	3424	Below tenth centile	Mixed	Yes	0.13 [0.09, 0.16]	0.96 [0.95, 0.96]

UA and SGA

Study	TP	FP	FN	TN	Threshold	Risk	Intervention	Sensitivity (95% CI)	Specificity (95% CI)
Hawkins 2012	168	621	56	461	+1SD	High	No	0.75 [0.69, 0.81]	0.43 [0.40, 0.46]
Amini 2014	14	89	32	269	+1SD	Mixed	Unknown	0.30 [0.18, 0.46]	0.75 [0.70, 0.80]
Bellomo 2011	36	34	7	86	309µmol/l	High	Unknown	0.84 [0.69, 0.93]	0.72 [0.63, 0.80]
Williams 2002	14	42	73	327	450µmol/l	High	No	0.16 [0.09, 0.26]	0.89 [0.85, 0.92]
Jauniaux 1996	7	4	9	21	4mg/dl	High	Unknown	0.44 [0.20, 0.70]	0.84 [0.64, 0.95]
Odendaal 1997	8	13	92	83	520µmol/l	High	No	0.08 [0.04, 0.15]	0.86 [0.78, 0.93]
Voto 1988	10	47	20	138	6mg%	High	Unknown	0.33 [0.17, 0.53]	0.75 [0.68, 0.81]
Yassae 2003	41	12	18	32	6mg/dl	High	Unknown	0.69 [0.56, 0.81]	0.73 [0.57, 0.85]

E3 AND/OR hPL and SGA

Study	TP	FP	FN	TN	Threshold	Risk	Intervention	Sensitivity (95% CI)	Specificity (95% CI)
Lenstrup 1982	5	4	4	75	Below tenth centile	Mixed	No	0.56 [0.21, 0.86]	0.95 [0.88, 0.99]

1) *Ultrasound assessment of fetal size (estimated fetal weight (EFW))*

There were 32 studies of estimated fetal weight (EFW) with a total of 51,702 pregnancies involving 6169 SGA infants (Figure 4). Of these, 20 studies (Baird 2016; Barel 2016; Ben-Haroush 2007; Callec 2015; Chauhan 1999; Chauhan 1999a; Chauhan 2003; Christensen 2015; Freire 2010; Gabbay-Benziv 2016; Geerts 2016; Gupta 2008; Hammad 2015; Hendrix 2000; Roma 2015; Sekar 2016; Sovio 2015; Takeuchi 1985; Turitz 2014; Weiner 2016) used the Hadlock formula, two (Laurin 1987; Palo 1989) used the Eik-Nes formula, and four (Berkowitz 1988; Chervenak 1984; Ott 1984; Skovron 1991) used the Shepard formula to estimate fetal weight. The formula used in the remaining six studies (Bikmetova 2013; Griffin 2015; Hatfield 2010; MacLeod 2013; Mahran 1988; Valino 2016) was not stated. There were two, 10 and 19 studies in low-, high- and mixed-risk cohorts; the patient group was unknown in one study. Six (18.8%) studies (Berkowitz 1988; Callec 2015; Chauhan 2003; Roma 2015;

Turitz 2014; Valino 2016) intervened based on test results, which may have altered the outcome of the pregnancy, while index test results did not affect management in four (12.5%) studies (Hammad 2015; Hendrix 2000; Sekar 2016; Weiner 2016). The remaining 22 (68.8%) studies did not provide information on whether the test results led to intervention(s). Sovio 2015 blinded clinicians to the results of the universal ultrasonography and Weiner 2016 blinded clinicians to results of all ultrasound methods other than the one they conducted, but in the majority of studies clinicians either were not blinded to test results or this was not reported. The same ultrasound threshold (10th centile) was used to determine test positivity in 25 studies, one study did not report the threshold used, and each of the six remaining studies used a different threshold (Figure 4). For the 10th centile ultrasound threshold, the summary sensitivity (95% CI) and specificity (95%) were 0.54 (0.43 to 0.65) and 0.95 (0.92 to 0.97) from meta-analysis of the 25 studies (47,057 pregnancies involving 5650 SGA infants) (Figure 6).

Figure 6. Summary ROC plot of ultrasound estimated fetal weight (EFW) at a tenth centile threshold for identifying small-for-gestational-age (birthweight $\leq$ tenth centile) infants. The study points are shown using different symbols for the formulas used. The study points were scaled according to the precision of sensitivity and specificity in the studies. The solid circle (summary point) represents the summary estimate of sensitivity and specificity for all 25 studies, and is surrounded by a dotted line representing the 95% confidence region and a dashed line representing the 95% prediction region. The 95% prediction region is the region within which one is 95% certain the results of a new study will lie.

2) Ultrasound placental grading

Twelve studies assessed placental grading (by ultrasound echogenicity) for identifying SGA infants (Figure 4). The studies included 4940 pregnancies involving 520 SGA infants. There were

two, six and four studies in low-, high- and mixed-risk cohorts, respectively. One study (McKenna 2005) intervened based on test results, a grade III placenta was used to identify pregnancies to be induced for suspected fetal compromise, and one study did not

(Chen 2012a). Three studies fall beneath the diagonal, two of these are small studies of nine and 55 participants, respectively (Altmann 1978; Estel 1989), while the other study examined placental grading in late pregnancy (36 to 38 weeks) (Miller 1988). The remaining

10 studies did not report this information. All the studies used Grannum grading based on placental calcification and set a grade III threshold; the summary sensitivity (95% CI) and specificity (95%) were 0.38 (0.23 to 0.55) and 0.79 (0.62 to 0.90) (Figure 7).

Figure 7. Summary ROC plot of placental grading for identifying small-for-gestational-age (birthweight $\leq$ tenth centile) infants. The study points are shown using different symbols for different risk groups. The study points were scaled according to the precision of sensitivity and specificity in the studies. The solid circle (summary point) represents the summary estimate of sensitivity and specificity for all 12 studies, and is surrounded by a dotted line representing the 95% confidence region and a dashed line representing the 95% prediction region. The 95% prediction region is the region within which one is 95% certain the results of a new study will lie.

3) Human placental lactogen (hPL)

Figure 5 shows the 20 studies that assessed hPL. A total of 3486 pregnancies involving 624 SGA infants were included. There were one, 11, and eight studies in low-, high- and mixed-risk cohorts, respectively. Most of the studies (14/20, 70%) did not report whether or not interventions were affected by index test results, four (20%) studies ([Granat 1977](#); [Nice 2016](#); [Odendaal 1981](#);

[Zhang 1990](#)) did not intervene based on test results, and in two (10%) studies ([Sagen 1984](#); [Siebert 1974](#)) index test results led to interventions. The studies used different test thresholds (Figure 5), but eight studies used the 10th centile test threshold. The meta-analysis of the eight studies (1124 pregnancies involving 303 SGA infants) gave summary sensitivity (95% CI) and specificity (95% CI) of 0.38 (0.23 to 0.55) and 0.88 (0.78 to 0.94) (Figure 8).

Figure 8. Summary ROC plot of human placental lactogen (hPL) for identifying small-for-gestational-age (birthweight $\leq$ tenth centile) infants. The study points are shown using different symbols for different risk groups. The study points were scaled according to the precision of sensitivity and specificity in the studies. The solid circle (summary point) represents the summary estimate of sensitivity and specificity for all nine studies, and is surrounded by a dotted line representing the 95% confidence region and a dashed line representing the 95% prediction region. The 95% prediction region is the region within which one is 95% certain the results of a new study will lie.

4) Serum oestriol

Nine studies of 2773 pregnancies involving 373 SGA infants were included (Figure 5). Five of the studies were in high-risk cohorts while the remaining four were in mixed-risk cohorts. Five studies

(Cedard 1979; Chard 1985; Nisbet 1982; Palo 1987; Spornol 1989) did not state whether they intervened based on index test results, three studies (Gerhard 1986; Nielsen 1985; Odendaal 1981) did not intervene based on index test results, while one study (Sagen

1984) did, described above. Five studies (Cedard 1979; Chard 1985; Gerhard 1986; Odendaal 1981; Sagen 1984) used the 10th centile test threshold and included a total of 1248 pregnancies (involving 204 SGA infants). Based on the five studies, the summary sensitivity (95% CI) and specificity (95% CI) were 0.39 (0.27 to 0.54) and 0.86 (0.79, 0.91).

5) Urinary oestriol

Urinary oestriol was assessed in nine studies with a total of 92,406 pregnancies involving 7076 SGA infants (Figure 5). Seven of the studies were in high-risk cohorts and two were in mixed-risk cohorts. Five studies (Campbell 1972; Kunz 1976; Oats 1979; Steiner 1991; Weerasinghe 1977) did not state whether or not index test results led to interventions, three studies (Chew 1976; Fliegner 1979; Odendaal 1981) did not intervene based on index test results and one study (Beischer 1991) stated that oestriol levels were an indicator for early delivery, although no numbers were given. Studies used different index test positivity thresholds (Figure 5) but four studies (Beischer 1991; Fliegner 1979; Oats 1979; Steiner 1991) used a threshold of 8 mg per 24 hours (at 30 weeks) to 12 mg per 24 hours (at 40 weeks). The summary sensitivity (95% CI) and specificity (95% CI) from the four studies (84,737 pregnancies involving 6886 SGA infants) were 0.31 (0.18, 0.49) and 0.84 (0.72, 0.91).

6) Placental growth factor (PIGF)

The seven studies of PIGF are shown in Figure 5. A total of 6405 pregnancies (involving 837 SGA infants) were included in the studies. Three of the studies were in high-risk cohorts while four were in mixed-risk cohorts. Two studies (Chaiworapongsa 2013; Kienast 2016) did not state whether index test results led to interventions, four studies (Benton 2016; Molvarec 2013; Nice 2016; Shawkat 2015) did not intervene based on test results and one study (Valino 2016) used the results of the EFW scan to determine whether suspected SGA pregnancies should be delivered by caesarean section. Three studies (Benton 2016; Nice 2016; Shawkat 2015) used a 12 pg/mL threshold, while each of the four remaining studies used a different threshold (Figure 5). Due to the limited number of studies at a common threshold and substantial heterogeneity, meta-analysis to obtain summary estimates of sensitivity and specificity was not performed.

7) Uric acid

Eight studies with a total of 2884 pregnancies (involving 605 SGA infants) assessed uric acid (Figure 5). The studies were in high-risk cohorts except for one study in a mixed-risk cohort. Five studies (Amini 2014; Bellomo 2011; Jauniaux 1996; Voto 1988; Yassaee 2003) did not state whether interventions were made based on index test results and three studies (Hawkins 2012; Odendaal 1997; Williams 2002) had no interventions. The studies used various thresholds (Figure 5).

8) Combination of serum oestriol and human placental lactogen (hPL)

Lenstrup 1982 (88 pregnancies) assessed this combination in a mixed-risk cohort (Figure 5). The sensitivity (95% CI) and specificity (95%) were 0.56 (0.21 to 0.86) and 0.95 (0.88 to 0.99). The specific measurement used as a threshold in this study was not stated; a low hPL and/or E3 level was classified as a positive test result. In this study, one patient was hospitalised after a cardiotocography (CTG) investigation due to a low plasma oestriol level.

Comparative analyses of seven tests for identifying small-for-gestational-age infants (SGA) (birthweight $\leq$ 10th centile)

Using all available data (86 studies), we compared the accuracy of EFW, hPL, serum oestriol, urinary oestriol, PIGF and uric acid in a single model. Based on the preliminary assessments and likelihood ratio tests comparing different HSROC meta-regression models, in the final model fitted, we modelled differences in accuracy and threshold as random effects with symmetric curves for all tests, i.e. parallel SROC curves such that each curve can be described using a diagnostic odds ratio (DOR) (Figure 9, Table 2). There was a statistically significant ($P < 0.0001$) difference in accuracy between the tests, with EFW being more accurate than all the other tests. For example, the DOR (95% CI) of EFW was 21.3 (13.1 to 34.6) while that of hPL, the biochemical test with the highest DOR, was 4.78 (3.21 to 7.13). Comparing EFW to hPL, the ratio of DORs (95% CI) was 4.45 (2.38 to 8.25) with statistical evidence of a difference in accuracy ($P < 0.0001$). Pairwise comparisons of the seven tests are shown in Table 2.

Figure 9. Summary ROC plot of structural and biochemical tests for identifying small-for-gestational-age (birthweight $\leq$ tenth centile) infants. E3 = oestriol; EFW = estimated fetal weight; hPL = human placental lactogen; PlGF = placental growth factor; SGA = small-for-gestational-age; UA = uric acid. The curve for each test is drawn within the range of estimates of specificity from the studies included for the test. Compared to the other curves, the curve for EFW lies closest to the top left hand corner (ideal position where sensitivity and specificity both equal 1). The position of the curves for UA and placental grading is very similar. The SROC curve for UA is the green curve lying above the pink curve for placental grading.

The sensitivities estimated along the SROC curves at fixed values of specificity are shown in Table 3 for different values of prevalence. Table 3 also shows the numbers of missed SGA infants and false positives in a hypothetical cohort of 1000 pregnant women. At the

median specificity of 0.88 and median prevalence of 19%, EFW, hPL, oestriol, urinary oestriol, uric acid, PlGF, and placental grading will miss 49, 116, 123, 128, 139, 144, and 145 SGA infants, respectively.

Ten studies (Altmann 1978; Chard 1985; Geerts 2016; Kunz 1976; Nice 2016; Nisbet 1982; Sagen 1984; Spornol 1989; Steiner 1991; Valino 2016) evaluated two tests and one study (Odendaal 1981) evaluated three tests (hPL, serum oestriol, and urinary oestriol). Of

the 11 studies, five (Chard 1985; Nisbet 1982; Odendaal 1981; Sagen 1984; Spornol 1989) evaluated hPL and serum oestriol (Figure 10). From the comparative meta-analysis of the five studies, the DOR (95% CI) for hPL was 5.60 (2.84 to 11.0) and that of serum oestriol was 4.06 (1.81 to 9.07); ratio of the DORs was 1.29 (0.58 to 2.86), $P = 0.40$. Due to limited data we did not perform meta-analyses for other pair-wise comparisons but summarised individual study results in Appendix 6.

Figure 10. Summary ROC plot of direct comparisons of human placental lactogen and oestriol for identifying small-for-gestational-age (birthweight $\leq$ tenth centile) infants. E3 = oestriol; hPL = human placental lactogen; SGA = small-for-gestational-age. Each symbol represents the pair of sensitivity and specificity for one test from a study. The pair of points for the two tests from a study are connected by a dotted line. The size of each symbol was scaled according to the precision of sensitivity and specificity in the study. Each summary curve was restricted to the range of specificities for each test from the five studies that evaluated both tests in the same patients.

Small-for-gestational age (birthweight < third centile) (SGA3)

SGA3 was evaluated in four studies ([Chaiworapongsa 2013](#); [Griffin 2015](#); [Roma 2015](#); [Sovio 2015](#)) involving 6953 pregnancies (235 cases). The four studies assessed PIGF and/or EFW in high- or mixed-risk cohorts ([Figure 11](#)). One study ([Griffin 2015](#)) of 592

pregnancies evaluated EFW, PIGF, and a combination of the two. The other three studies (5678 pregnancies) assessed EFW and used the 10th centile as the threshold. The summary sensitivity (95% CI) and specificity (95% CI) were 0.66 (0.56 to 0.76) and 0.87 (0.80 to 0.91). The two PIGF studies (1861 pregnancies) used

different thresholds (< 0.3 MoM (multiples of the median) and fifth centile); sensitivities were 0.52 (0.31 to 0.73) and 0.37 (0.26 to 0.49) and specificities were 0.83 (0.81 to 0.85) and 0.89 (0.86 to 0.91)

respectively. The sensitivity (95% CI) and specificity (95% CI) from the single study of the PLGF and EFW combination were 0.69 (0.55 to 0.81) and 0.72 (0.67 to 0.77).

Figure 11. Forest plot of ultrasound estimated fetal weight (EFW) and placental growth factor (PLGF) for identifying small-for-gestational age (birthweight <third centile) infants. FN = false negative; FP = false positive; SGA3 = small-for-gestational-age birthweight <third centile; TN = true negative; TP = true positive. For each test, the forest plot shows the estimates of sensitivity and specificity from each study, the threshold used to define test positivity, risk of SGA pregnancy and whether there was an intervention that may have altered outcome. Studies are sorted by threshold, risk and intervention status.

EFW and SGA3

Study	TP	FP	FN	TN	Threshold	Risk	Intervention	Sensitivity (95% CI)	Specificity (95% CI)
Griffin 2015	44	108	32	402	Below tenth centile	High	Unknown	0.58 [0.46, 0.69]	0.79 [0.75, 0.82]
Sovio 2015	67	495	20	3395	Below tenth centile	Mixed	Unknown	0.77 [0.67, 0.85]	0.87 [0.86, 0.88]
Roma 2015	30	91	19	975	Below tenth centile	Mixed	Yes	0.61 [0.46, 0.75]	0.91 [0.90, 0.93]

PLGF and SGA3

Study	TP	FP	FN	TN	Threshold	Risk	Intervention	Sensitivity (95% CI)	Specificity (95% CI)
Chaiworapongsa 2013	12	210	11	1036	<0.3MoM	Mixed	Unknown	0.52 [0.31, 0.73]	0.83 [0.81, 0.85]
Griffin 2015	29	58	49	456	Below fifth centile	High	Unknown	0.37 [0.26, 0.49]	0.89 [0.86, 0.91]

Stillbirth

1) Ultrasound placental grading

Three studies assessed placental grading for stillbirth (Figure 12) with a total of 15,2236 pregnancies involving 114 stillbirths. Two of the studies (Altmann 1978; Chen 2012) were in high-risk cohorts

and the third (Chen 2015) was in a low-risk cohort. Chen 2015 contributed most of the data (15,122/15,236, 99%). The sensitivity and specificity of placental grading in this study was 0.35 (0.26 to 0.46) and 0.94 (0.93 to 0.94). None of the studies stated whether or not there were any interventions based on test results and all used a grade III threshold.

Figure 12. Forest plot of structural and biochemical tests for predicting stillbirth. E3 = oestriol; FN = false negative; FP = false positive; hPL = human placental lactogen; PlGF = placental growth factor; TN = true negative; TP = true positive; UA = uric acid. For each test, the forest plot shows the estimates of sensitivity and specificity from each study, the threshold used to define test positivity, risk of stillbirth and whether there was an intervention that may have altered outcome. Studies are sorted by threshold, risk and intervention.

hPL and stillbirth

Study	TP	FP	FN	TN	Threshold	Risk	Intervention	Sensitivity (95% CI)	Specificity (95% CI)
Ylikorkala 1973	12	68	2	117	Below 2.5th centile	High	Unknown	0.86 [0.57, 0.98]	0.63 [0.56, 0.70]
Siebert 1974	1	2	1	16	Below tenth centile	Low	Yes	0.50 [0.01, 0.99]	0.89 [0.65, 0.99]
Altmann 1978	2	1	1	6	Abnormal value	High	Unknown	0.67 [0.09, 0.99]	0.86 [0.42, 1.00]
Trudinger 1979	1	13	0	45	Below tenth centile	High	Unknown	1.00 [0.03, 1.00]	0.78 [0.65, 0.87]
Leader 1980	5	18	3	109	4mg/ml	High	No	0.63 [0.24, 0.91]	0.86 [0.79, 0.91]
Zhang 1990	8	59	0	54	4µg/ml	High	No	1.00 [0.63, 1.00]	0.48 [0.38, 0.57]

Urinary E3 and stillbirth

Study	TP	FP	FN	TN	Threshold	Risk	Intervention	Sensitivity (95% CI)	Specificity (95% CI)
Elliott 1970	2	8	0	12	-2SD	High	Yes	1.00 [0.16, 1.00]	0.60 [0.36, 0.81]
Campbell 1972	6	74	5	199	Unknown	High	Unknown	0.55 [0.23, 0.83]	0.73 [0.67, 0.78]
Chew 1976	0	0	6	37	-2SD	High	No	0.00 [0.00, 0.46]	1.00 [0.91, 1.00]
Weerasinghe 1977	3	131	0	193	-2SD	High	Unknown	1.00 [0.29, 1.00]	0.60 [0.54, 0.65]
Fliegner 1979	5	74	0	250	8mg at 30w to 12mg at 40w (per 24h)	Mixed	No	1.00 [0.48, 1.00]	0.77 [0.72, 0.82]
Oats 1979	90	2471	82	16476	8mg at 30w to 12mg at 40w (per 24h)	Mixed	Unknown	0.52 [0.45, 0.60]	0.87 [0.86, 0.87]
Beischer 1991	163	6912	289	64698	8mg at 30w to 12mg at 40w (per 24h)	High	Yes	0.36 [0.32, 0.41]	0.90 [0.90, 0.91]

PlGF and stillbirth

Study	TP	FP	FN	TN	Threshold	Risk	Intervention	Sensitivity (95% CI)	Specificity (95% CI)
Chaiworapongsa 2013	4	71	1	1193	<0.12MoM	Mixed	Unknown	0.80 [0.28, 0.99]	0.94 [0.93, 0.96]
Shawkat 2015	2	59	1	199	12pg/ml	High	No	0.67 [0.09, 0.99]	0.77 [0.72, 0.82]
Benton 2016	6	151	1	253	12pg/ml	High	No	0.86 [0.42, 1.00]	0.63 [0.58, 0.67]
Valino 2016	1	197	0	3755	Below tenth centile	Mixed	Yes	1.00 [0.03, 1.00]	0.95 [0.94, 0.96]

UA and stillbirth

Study	TP	FP	FN	TN	Threshold	Risk	Intervention	Sensitivity (95% CI)	Specificity (95% CI)
Redman 1976	1	32	1	247	360µmol/l	Unknown	No	0.50 [0.01, 0.99]	0.89 [0.84, 0.92]
Odendaal 1997	3	18	15	160	520µmol/l	High	No	0.17 [0.04, 0.41]	0.90 [0.84, 0.94]
Yassae 2003	12	41	0	50	6mg/dl	High	Unknown	1.00 [0.74, 1.00]	0.55 [0.44, 0.65]
Hawkins 2012	3	883	2	595	+1SD	High	No	0.60 [0.15, 0.95]	0.40 [0.38, 0.43]

Placental grading and stillbirth

Study	TP	FP	FN	TN	Threshold	Risk	Intervention	Sensitivity (95% CI)	Specificity (95% CI)
Altmann 1978	1	6	1	1	Grade III	High	Unknown	0.50 [0.01, 0.99]	0.14 [0.00, 0.58]
Chen 2012	9	35	4	57	Grade III	High	Unknown	0.69 [0.39, 0.91]	0.62 [0.51, 0.72]
Chen 2015	35	939	64	14084	Grade III	Low	Unknown	0.35 [0.26, 0.46]	0.94 [0.93, 0.94]

2) Human placental lactogen

Six studies of 544 pregnancies involving 36 stillbirths were included (Figure 12). One study (Siebert 1974) was in a low-risk cohort while the others were in high-risk cohorts. Three studies (Altmann 1978; Trudinger 1979; Ylikorkala 1973) did not report whether or not there were interventions based on index test results, two studies (Leader 1980; Zhang 1990) did not intervene while one study (Siebert 1974) intervened based on test results; both false positive results were caesarean sections prompted by falling hPL values. The studies used various thresholds with sensitivities ranging from 0.50 to 1.00, and specificities from 0.48 to 0.89 (Figure 12).

3) Urinary oestriol

Seven urinary oestriol studies included 92,186 pregnancies involving 651 stillbirths (Figure 12). Five studies were in high-risk cohorts and two were in mixed-risk cohorts. Three studies (Campbell 1972; Oats 1979; Weerasinghe 1977) did not report whether there were any interventions based on index test results, two studies (Chew 1976; Fliegner 1979) did not intervene due to index test results while in two studies (Beischer 1991; Elliott 1970) index test results affected pregnancy management; in the Elliott study there were five caesarean sections that were performed due to greatly reduced oestriol excretion. The studies used different

thresholds with sensitivities ranging from 0 to 1, and specificities from 0.60 to 1.00.

4) Placental growth factor

A total of 5894 pregnancies (involving 16 stillbirths) were included in three studies (Figure 12). Two studies were in high-risk cohorts and two were in mixed-risk cohorts. One study (Chaiworapongsa 2013) did not report whether or not index test results could lead to interventions, two studies (Benton 2016; Shawkat 2015) did not intervene based on index test results and one study (Valino 2016) did intervene, described earlier. The studies used different thresholds with sensitivities ranging from 0.67 to 1.00, and specificities from 0.63 to 0.95.

5) Uric acid

Four studies included 2063 pregnancies (involving 37 stillbirths) (Figure 12). Three of the studies were in high-risk cohorts and the remaining study was in a mixed-risk cohort. Intervention status was unknown for one study (Yassae 2003) while there were no interventions in three studies (Hawkins 2012; Odendaal 1997; Redman 1976). None of the studies used the same threshold. The sensitivities ranged from 0.17 to 1.00, and specificities ranged from 0.40 to 0.90.

Comparative analyses of four biochemical tests for predicting stillbirth

In an indirect test comparison based on 21 studies (including 100,687 pregnancies involving 740 stillbirths), we compared the accuracy of hPL, urinary oestriol, PlGF and uric acid. As there were several tests and only three placental grading studies, to reduce model complexity, we did not include this test in the model. We fitted a model with symmetric curves for the four biochemical tests

(Figure 13) and the DORs and ratio of DORs are shown in Table 4. There was a statistically significant ($P < 0.0001$) difference in accuracy between the tests, with PlGF being the most accurate. For example, the DOR (95% CI) of PlGF was 49.2 (12.7 to 191) while that of urinary oestriol was 5.83 (4.91 to 6.92). The ratio of the DORs (95% CI) of PlGF and urinary oestriol was 8.44 (2.15 to 33.1) with statistical evidence of a difference in accuracy ($P = 0.004$). Pair-wise comparisons of the four tests are shown in Table 5.

Figure 13. Summary ROC plot of biochemical tests for predicting stillbirth. E3 = Oestriol; hPL = human placental lactogen; PlGF = placental growth factor; SGA = small-for-gestational-age; UA =uric acid. The SROC curves for the four tests are parallel. The curve for each test is drawn within the range of estimates of specificity from the studies included for the test. Compared to the other curves, the curve for PlGF lies closest to the top left hand corner (ideal position where sensitivity and specificity both equal 1).

Using sensitivities estimated at fixed values of specificity, [Table 5](#) shows the numbers of missed stillbirths and false positives in a hypothetical cohort of 1000 pregnant women. At the median specificity of 0.78 and median prevalence of 1.7%, PlGF, hPL,

urinary oestriol and uric acid will miss 2, 5, 7 and 8 stillbirths, respectively.

There were no comparative studies of biochemical tests; one study (Altmann 1978) evaluated hPL and placental grading in a small cohort of 10 pregnancies.

Studies with evaluations of both small-for-gestational-age infants (birthweight $\leq$ 10th centile) and stillbirth

Biochemical tests were assessed for both SGA and stillbirth in 18 studies (Appendix 7). In addition, Chen 2012 assessed placental grading for both outcomes. As it is only possible to show one outcome per study on the forest plot, we chose to display only biochemical tests for clarity. Results for Altmann 1978 and Chen 2012 can be seen in Figure 4 for SGA and in Figure 12 for stillbirth.

Investigation of heterogeneity

Where the number of studies allowed, we investigated the effect of the formula used to estimate fetal weight based on ultrasound examination on test accuracy in a meta-regression. We compared the 20 studies (41,104 pregnancies) that used the Hadlock formula with the four studies (1710 pregnancies) that used the Shepard formula. The DORs (95% CI) for the Shepard and Hadlock formulas were 28.3 (6.60 to 121) and 24.2 (12.3 to 47.7). There was no statistical evidence of a difference in accuracy; the ratio of DORs (95%) was 1.17 (0.23 to 5.79), $P = 0.55$. Due to limited data, we were unable to formally investigate the effect of other potential sources of heterogeneity as outlined in the Secondary objectives.

DISCUSSION

Summary of main results

Small-for-gestational age (SGA) (birthweight $\leq$ 10th centile)

The main findings of this review are that ultrasound estimated fetal weight (EFW) is the most accurate test available to detect an SGA infant, but still only detects about 50% of infants born SGA. There are more studies of ultrasound EFW than any biochemical marker of placental dysfunction in this review. Overall, ultrasound EFW is better at predicting SGA < 3rd centile than SGA < 10th centile, which suggests that the more severe the phenotype of SGA, the better the accuracy of ultrasound EFW to identify that baby as SGA. Analysis of the biochemical tests found that human placental lactogen (hPL) was best for identifying an infant that would be SGA at birth, although it was inferior to the performance of ultrasound EFW.

Stillbirth

Fewer studies investigated the prediction of pregnancies that would end in stillbirth and only three studies examined abnormal placental structure or function directly (Benton 2016, Jauniaux 1996, Ylikorkala 1973) by pathological examination.

Placental histopathology data were available for 213 women from the Benton 2016 study; 55 of 94 women with low placental growth factor (PIGF) had Grannum grade II or III placentas, as did one of the 119 with normal PIGF values. Jauniaux 1996 found extended vascular lesions after histopathological examination in 12 complicated cases and Ylikorkala 1973 found evidence of degenerative placental changes and dysmature placenta in 50% of cases; we did not use data from the entire cohort so cannot know numbers from these but there was no relationship found between placental calcification and serum hPL level.

There were no studies that presented data to allow estimation of the diagnostic accuracy of EFW measurement for stillbirth; Sovio 2015 included stillbirths but data were presented as part of a severe adverse perinatal outcome group meaning that stillbirth could not be assessed separately. Included studies measured hPL, PIGF, placental grading, uric acid, and urinary E3. Of the biochemical tests for detecting pregnancies that would end in stillbirth, PIGF had the best diagnostic accuracy.

Critically, very few studies investigated the accuracy of different tests in the same study population, which would allow direct comparison of diagnostic test accuracy, and only one study (Lenstrup 1982) investigated combinations of biomarkers. Indirect comparison suggests that biochemical tests of placental function performed better in identification of pregnancies that would end in a stillbirth compared to detection of a pregnancy that would end in the birth of an SGA infant.

Strengths and weaknesses of the review

This review is the first comprehensive review of biochemical markers in maternal blood or serum in late pregnancy in comparison with the performance of ultrasound to detect pregnancies that end in stillbirth and SGA births. We examined papers from different countries and published in languages other than English, allowing us to cover a wide range of populations of pregnant women. We contacted authors where necessary to clarify or obtain data (although this was not always successful) and were able to include unpublished data in some cases. However, some papers were also unobtainable, meaning it is possible that some usable data may not have been included (see Characteristics of studies awaiting classification). Furthermore, the nomenclature of some of the biochemical factors has changed meaning that the search strategies may not have included all possible studies.

There are some limitations which need to be considered.

- Limitations in the availability and variability in reporting of data prevented us from undertaking comparisons and investigations of heterogeneity as originally set out in the protocol. Measurements were taken at many different gestational ages and these were often not comparable. Studies often took measurements over a fixed range of time rather than at one time point so there were few studies that used the exact same gestation; a comparison of studies that tested before and after term (37 weeks) was not possible, due to the amount of overlap of this time point.
- Studies used many different measurement techniques for the same index tests, for example measurement of different isoforms of the biochemical factor of interest, which could have had an impact on test performance. Definitions of index test positivity thresholds for individual tests varied and were often not prespecified; they were frequently based on optimally performing thresholds in the study population itself rather than on externally validated thresholds. This meant that it was not practical to perform a formal comparison of all measurements at the 10th centile, for example, as exact measurements differed across studies. If there were common thresholds between studies then these are the ones for which data were extracted, especially if these were in clinical use (e.g. < 12 pg/mL for PIGF). Defined externally-validated thresholds are needed before recommendations for clinical practice can be made.

- Few papers made direct comparisons between index tests (comparison of index test in the same population) or investigated the accuracy of testing strategies (combining several index tests for the same woman). Due to differences between study populations, such as risk factors, it is unclear how useful comparisons between studies are.
- Due to variation between study populations we decided to use fairly broad definitions of patient risk; high risk was defined as all women at high risk due to various factors (pre-eclampsia, hypertension, suspected fetal growth restriction (FGR), history of SGA or stillbirth), mixed risk was where some but not all women were affected by these conditions (including unselected populations), and low-risk was where all complications were excluded. It was not possible to look at test performances in only hypertensive women or only women with suspected FGR, for example.
- With the exception of uric acid, the studies of biochemical markers were performed in two distinct time periods, before 1991 (for hPL and E3) and after 2013 (for PlGF), whereas the accuracy of ultrasound scan has been studied continuously since the 1970s. Earlier studies tended to be smaller and less rigorously conducted and reported as required by contemporary standards of study reporting. In addition, methods of biochemical analysis have also developed over the time frame of the review from radioimmunoassay to enzyme-linked immunosorbent assay (ELISA) or chemiluminescent assays.

Applicability of findings to the review question

Ultrasound EFW is a well-established technique in contemporary obstetric practice and is at the centre of screening for FGR (RCOG 2013) and strategies to reduce stillbirth (NHS England 2016). Many of the studies were conducted in women who were at increased risk of FGR or stillbirth, so these findings cannot necessarily be generalised to a mixed- or low-risk population. This is particularly important because the risk status of the woman alters the importance of specific aspects of test performance. In mothers at high risk of stillbirth, a false negative result would potentially deprive the woman of additional monitoring or intervention which may mitigate the increased risk of adverse outcome, thus sensitivity would be prioritised over specificity. However, women in low- or mixed-risk populations, a false positive result may increase monitoring and/or intervention which is unnecessary and could have short- and long-term consequences (Peters 2018). Unfortunately, limitations in data meant that the impact of the potential sources of heterogeneity described above (including risk status of the mother) on test accuracy could not be explored. We were not able to find all of the information we had intended to, particularly about the outcome of testing and potential harms of testing because data were unavailable. The search was completed in October 2016 and it is possible that new studies, particularly regarding PlGF, have been added since then.

AUTHORS' CONCLUSIONS

Implications for practice

Ultrasound estimated fetal weight (EFW) appears to be the most accurate method to identify a baby that will be small-for-gestational age (SGA) at birth. However, clinicians should be aware that the estimates of sensitivity (the range of sensitivity for included studies was 0.07 to 0.93) that ultrasound EFW does not detect a

significant proportion of babies who have a birthweight < 10th centile. Importantly, EFW has a high specificity (pooled estimate 0.95) which avoids implications of false positive results (e.g. maternal anxiety, further testing, or unwarranted intervention). This review could not find a significant difference between two of the formulae used to derive EFW on diagnostic accuracy (Shepard or Hadlock). This review also suggests that biochemical markers of placental function cannot be used alone to identify which pregnancies will end with the birth of an SGA baby. Biochemical markers show promise in identifying babies who go on to be stillborn, although data are largely confined to women at increased risk of stillbirth. Furthermore, the performance of biochemical markers in comparison to ultrasound EFW for stillbirth is unknown.

Implications for research

Further research studies of ultrasound are required to determine whether an EFW under a specific threshold, e.g. 10th centile identifies pregnancies that end in stillbirth. This could be achieved by including stillbirth as an outcome measure in all studies; stillbirths were frequently excluded from many of the studies examining the accuracy of ultrasound EFW. While individual studies of the accuracy of ultrasound EFW are likely to be underpowered to evaluate the prediction of a pregnancy that ends in stillbirth, this aim could be achieved by meta-analysis of such studies. Further studies are also needed of biochemical markers alone and in combination with ultrasound EFW. Such studies would facilitate direct comparison of test performance and whether a combination of tests would be more effective than a single means of assessment for detection of SGA infants at birth and pregnancies that end in stillbirth.

Ideally, future studies should have an adequate sample size to study their outcome of interest or when this is unlikely to be possible, e.g. in studies of stillbirth a range of important pregnancy outcomes should be reported to facilitate future meta-analyses. Similarly, studies should also have standardised reporting using the STARD guidelines (Bossuyt 2015) to ensure that information can be extracted for future meta-analyses. In addition, researchers should consider using direct assessment of placental pathology as the reference standard. For practical reasons, this review used two clinical endpoints which are associated with placental pathology. However, the use of clinical endpoints rather than evidence of placental pathology could account for the apparent suboptimal test performance. For example, 70% of SGA infants (< 10th percentile) are "constitutionally small", with no evidence of underlying placental dysfunction, so would not be expected to have a positive test result. This may also explain why biochemical markers of placental function appear to have a greater accuracy in predicting stillbirth compared to SGA, as stillbirth is more frequently associated with placental abnormalities (in up to 60% of cases). Further studies are needed to determine whether ultrasound EFW, biochemical markers of placental function, and other measurements such as fetal and maternal Doppler measurements accurately identify placental pathology.

Initial research studies may wish to focus on women at highest risk of placental dysfunction, and by association the birth of an SGA infant and stillbirth, e.g. women who are suspected to have a SGA infant by symphysis-fundal height (SFH) measurement, or who present with reduced fetal movements or have hypertension. Studies could then explore the diagnostic accuracy of biochemical factors alone, or in association with ultrasound measurements, in a

low- or mixed-risk population where the background risk of adverse outcome is lower.

Test evaluation studies need to consider the implications of revealing index test results as this could have adverse consequences such as increased intervention by induction of labour or increased frequency of birth by caesarean section and admission to neonatal units, all of which may impact upon mothers' experience of care and fetal outcome. The clinical role of biochemical markers of placental dysfunction needs to be revisited using robust study designs with adequate sample sizes and standardised reporting. Ultimately, clinical efficacy will need to be demonstrated by intervention studies before any test is adopted into clinical practice.

ACKNOWLEDGEMENTS

Alexander Heazell is supported by a Clinician Scientist Fellowship (CS-2013-009) from the National Institute for Health Research (NIHR).

The authors would like to thank the patient participant involvement group chaired by Claire Storey for the Placental Assessment Predicting Pregnancy Outcome (PAPPO) study for their assistance with preparing the Plain language summary. We are also very grateful to everyone who has helped with translations at any point during this review: Laura Avagliano, Lauren Baker, Sandor Bako, Janine Dretske, Rui Duarte, Filip Ericsson, Claudia Ravaldi, Michael Robinson, Katalin Wilkinson and Yu-Tian Xiao.

As part of the pre-publication editorial process, this review has been commented on by three peers (an editor and two referees who are external to the editorial team). The authors are grateful to the following peer reviewers for their time and comments: Professor GC Smith, Helen Sassoon.

This project was supported by the National Institute for Health Research, via Cochrane Infrastructure funding to Cochrane Pregnancy and Childbirth. The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the Systematic Reviews Programme, NIHR, NHS or the Department of Health.

REFERENCES

References to studies included in this review

Altmann 1978 {published data only}

Altmann P, Janisch H, Muller-Tyl E, Reinold E, Spona J, Havelec L. Serum oestriol: a parameter of the function of the feto-placental unit. *Wiener Klinische Wochenschrift* 1978;**90**:121-7.

Amini 2014 {published data only}

Amini E, Sheikh M, Hantoushzadeh S, Shariat M, Abdollahi A, Kashanian M. Maternal hyperuricemia in normotensive singleton pregnancy, a prenatal finding with continuous perinatal and postnatal effects, a prospective cohort study. *BMC Pregnancy & Childbirth* 2014;**14**:104.

Baird 2016 {published data only}

Baird SM, Davies-Tuck M, Coombs P, Knight M, Wallace EM. Detection of the growth-restricted fetus: which centile charts?. *Sonography* 2016;**3**:81-6.

Barel 2016 {published data only}

Barel O, Maymon R, Elovits M, Smorgick N, Tovbin J, Vaknin Z. Evaluation of fetal weight estimation formulas in assessing small-for-gestational-age fetuses. *Ultraschall in der Medizin* 2016;**37**:283-9.

Beischer 1991 {published data only}

Beischer N, Brown J, Parkinson P, Walstab J. Urinary oestriol assay for monitoring fetoplacental function. *Australian & New Zealand Journal of Obstetrics & Gynaecology* 1991;**31**:1-8.

Bellomo 2011 {published data only}

Bellomo G, Venanzi S, Saronio P, Verdura C, Narducci PL. Prognostic significance of serum uric acid in women with gestational hypertension. *Hypertension* 2011;**58**:704-8.

Ben-Haroush 2007 {published data only}

Ben-Haroush A, Yogeve Y, Hod M, Bar J. Predictive value of a single early fetal weight estimate in normal pregnancies. *European Journal of Obstetrics Gynecology and Reproductive Biology* 2007;**130**:187-92.

Benton 2016 {published data only}

Benton SJ, McCowan LM, Heazell AE, Grynspan D, Hutcheon JA, Senger C, et al. Placental growth factor as a marker of fetal growth restriction caused by placental dysfunction. *Placenta* 2016;**42**:1-8.

Berkowitz 1988 {published data only}

Berkowitz GS, Chitkara U, Rosenberg J, Cogswell C, Walker B, Lahman EA, et al. Sonographic estimation of fetal weight and Doppler analysis of umbilical artery velocimetry in the prediction of intrauterine growth retardation: a prospective study. *American Journal of Obstetrics & Gynecology* 1988; Vol. 158:1149-53.

Bikmetova 2013 {published data only}

Bikmetova E, Trishkin A, Artymuk N. Fetal growth restriction: Informativity of modern diagnosis methods. *Journal of Perinatal Medicine* 2013;**41**:85.

Callec 2015 {published data only}

Callec R, Lamy C, Perdriolle-Galet E, Patte C, Heude B, Morel O. Impact on obstetric outcome of third-trimester screening for small-for-gestational-age fetuses. *Ultrasound in Obstetrics & Gynecology* 2015;**46**:216-20.

Campbell 1972 {published data only}

Campbell S, Kurjak A. Comparison between urinary oestrogen assay and serial ultrasonic cephalometry in assessment of fetal growth retardation. *British Medical Journal* 1972;**4**:336-40.

Cedard 1979 {published data only}

Cedard L, Bedin M, Leblond J, Tanguy G, Kaminski M. Maternal plasma total oestriol and dehydroepiandrosterone sulfate loading test as indicators of feto-placental function or placental sulfatase deficiency. *Journal of Steroid Biochemistry* 1979;**11**:501-7.

Chaiworapongsa 2013 {published data only}

Chaiworapongsa T, Romero R, Korzeniewski SJ, Kusanovic JP, Soto E, Lam J, et al. Maternal plasma concentrations of angiogenic/antiangiogenic factors in the third trimester of pregnancy to identify the patient at risk for stillbirth at or near term and severe late preeclampsia. *American Journal of Obstetrics & Gynecology* 2013;**208**:287.e1-287.e15.

Chard 1985 {published data only}

Chard T, Sturdee J, Cockrill B, Obiekwe BC. Which is the best placental function test? A comparison of placental lactogen and unconjugated oestriol in the prediction of intrauterine growth retardation. *European Journal of Obstetrics, Gynecology, & Reproductive Biology* 1985;**19**:13-7.

Chauhan 1999 {published data only}

Chauhan SP, Scardo JA, Hendrix NW, Magann EF, Morrison JC. Accuracy of sonographically estimated fetal weight with and without oligohydramnios. A case-control study. *Journal of Reproductive Medicine* 1999;**44**:969-73.

Chauhan 1999a {published data only}

Chauhan SP, Scardo JA, Magann EF, Devoe LD, Hendrix NW, Martin JN Jr. Detection of growth-restricted fetuses in preeclampsia: a case-control study. *Obstetrics & Gynecology* 1999;**93**:687-91.

Chauhan 2003 {published data only}

Chauhan SP, Magann EF, Velthuis S, Nunn SL, Reynolds D, Scardo JA, et al. Detection of fetal growth restriction in patients with chronic hypertension: is it feasible?. *Journal of Maternal-Fetal & Neonatal Medicine* 2003;**14**:324-8.

Chen 2012 {published data only}

Chen KH, Chen LR. PP096. The effect of preterm placental calcification on uteroplacental blood flow, fetal growth and

perinatal outcome in hypertension complicating pregnancy. *Pregnancy Hypertension* 2012;**2**:292.

Chen 2012a {published data only}

Chen KH, Chen LR, Lee YH. The role of preterm placental calcification in high-risk pregnancy as a predictor of poor uteroplacental blood flow and adverse pregnancy outcome. *Ultrasound in Medicine & Biology* 2012;**38**(6):1011-8.

Chen 2015 {published data only}

Chen KH, Seow KM, Chen LR. The role of preterm placental calcification on assessing risks of stillbirth. *Placenta* 2015;**36**:1039-44.

Chervenak 1984 {published data only}

Chervenak FA, Romero R, Berkowitz RL, Scott D, Tortora M, Hobbins JC. Use of sonographic estimated fetal weight in the prediction of intrauterine growth retardation. *American Journal of Perinatology* 1984; Vol. 1, issue 4:298-301.

Chew 1976 {published data only}

Chew PC, Ratnam SS. Plasma estradiol-17 beta as an index of fetoplacental function. *International Journal of Gynaecology & Obstetrics* 1976;**14**:445-8.

Chitlange 1990 {published data only}

Chitlange SM, Hazari KT, Joshi JV, Shah RK, Mehta AC. Ultrasonographically observed preterm grade III placenta and perinatal outcome. *International Journal of Gynaecology & Obstetrics* 1990;**31**:325-8.

Christensen 2015 {published data only}

Christensen KM, Heilbrun ME, Patel N, Woodward PJ, Kennedy A. Estimated fetal weight and birth weight associated with isolated single umbilical artery: the University of Utah experience. *Ultrasound Quarterly* 2015;**31**:19-22.

Elliott 1970 {published data only}

Elliott PM. Urinary oestriol excretion in retarded intrauterine fetal growth. *Australian & New Zealand Journal of Obstetrics & Gynaecology* 1970;**10**(1):18-21.

Estel 1989 {published data only}

Estel C, Eichhorn KH. [The correlation of reduced amniotic fluid volume, placental maturity and morphologic placental changes in placental insufficiency]. *Zentralblatt fur Gynakologie* 1989;**111**:891-6.

Fliegner 1979 {published data only}

Fliegner JR. Comparison of urinary pregnanediol and estriol excretion as indexes of placental function. *Obstetrics & Gynecology* 1979;**53**(1):93-8.

Freire 2010 {published data only}

Freire DM, Cecatti JG, Paiva CS. Correlation between estimated fetal weight by ultrasound and neonatal weight [Correlacao entre peso fetal estimado por ultrassonografia e peso neonatal]. *Revista Brasileira de Ginecologia e Obstetrica* 2010; Vol. 32:4-10.

Gabbay-Benziv 2016 {published data only}

Gabbay-Benziv R, Aviram A, Bardin R, Ashwal E, Melamed N, Hiersch L, et al. Prediction of small for gestational age: accuracy of different sonographic fetal weight estimation formulas. *Fetal Diagnosis & Therapy* 2016;**40**:205-13.

Geerts 2016 {published data only}

Geerts L, Van der ME, Theron A, Rademan K. Placental insufficiency among high-risk pregnancies with a normal umbilical artery resistance index after 32 weeks. *International Journal of Gynaecology & Obstetrics* 2016;**135**:38-42.

Gerhard 1986 {published data only}

Gerhard I, Fitzer C, Klinga K, Rahman N, Runnebaum B. Estrogen screening in evaluation of fetal outcome and infant's development. *Journal of Perinatal Medicine* 1986;**14**:279-91.

Gohari 1978 {published data only}

Gohari P, Hobbins JC, Berkowitz R. Use of hPL in the diagnosis of intrauterine growth retardation. *Obstetrics & Gynecology* 1978;**52**:127-8.

Granat 1977 {published data only}

Granat M, Sharf M, Diengott D, Spindel A, Kahana L, Elrad H. Further investigation on the predictive value of human placental lactogen in high-risk pregnancies. *American Journal of Obstetrics and Gynecology* 1977; Vol. 129:647-54.

Griffin 2015 {published data only}

Griffin M, Seed PT, Webster L, Myers J, MacKillip L, Simpson N, et al. Diagnostic accuracy of placental growth factor and ultrasound parameters to predict the small-for-gestational-age infant in women presenting with reduced symphysis-fundus height. *Ultrasound in Obstetrics & Gynecology* 2015;**46**:182-90.

Gupta 2008 {published data only}

Gupta LM, Gaston L, Chauhan SP. Detection of fetal growth restriction with preterm severe preeclampsia: experience at two tertiary centers. *American Journal of Perinatology* 2008;**25**:247-9.

Hammad 2015 {published data only}

Hammad IA, Chauhan SP, Mlynarczyk M, Rabie N, Goodie C, Chang E, et al. Uncomplicated pregnancies and ultrasounds for fetal growth restriction: a pilot randomized clinical trial. *AJP Reports* 2015;**6**:e83-90.

Hatfield 2010 {published data only}

Hatfield T, Caughey AB, Lagrew DC, Heintz R, Chung JH. The use of ultrasound to detect small-for-gestational-age infants in patients with elevated human chorionic gonadotropin on maternal serum screening. *American Journal of Obstetrics and Gynecology* 2010; Vol. 27:173-80.

Hawkins 2012 {published and unpublished data}

Hawkins TL, Roberts JM, Mangos GJ, Davis GK, Roberts LM, Brown MA. Plasma uric acid remains a marker of poor outcome in hypertensive pregnancy: a retrospective cohort study. *BJOG: an international journal of obstetrics and gynaecology* 2012;**119**:484-92.

Hendrix 2000 {published data only}

Hendrix NW, Grady CS, Chauhan SP. Clinical vs. sonographic estimate of birth weight in term parturients: a randomized clinical trial. *Journal of Reproductive Medicine for the Obstetrician and Gynecologist* 2000;**45**:317-22.

Howell 1985 {published data only}

Howell RJ, Perry LA, Ghoglay NS. Placental protein 12 (PP12): a new test for the prediction of the small-for-gestational-age infant. *British Journal of Obstetrics and Gynaecology* 1985;**92**:1141-4.

Jauniaux 1996 {published data only}

Jauniaux E, Gulbis B, Tunkel S, Ramsay B, Campbell S, Meuris S. Maternal serum testing for alpha-fetoprotein and human chorionic gonadotropin in high-risk pregnancies. *Prenatal Diagnosis* 1996;**16**:1129-35.

Kazzi 1983a {published data only}

Kazzi GM, Gross TL, Sokol RJ, Kazzi NJ. Detection of intrauterine growth retardation: a new use for sonographic placental grading. *American Journal of Obstetrics and Gynecology* 1983;**145**:733-7.

Kienast 2016 {published data only}

Kienast C, Moya W, Rodriguez O, Jijon A, Geipel A. Predictive value of angiogenic factors, clinical risk factors and uterine artery Doppler for pre-eclampsia and fetal growth restriction in second and third trimester pregnancies in an Ecuadorian population. *Journal of Maternal-Fetal & Neonatal Medicine* 2016;**29**:537-43.

Klebe 1990 {published data only}

Klebe JG, Marushak A, Bock J. Human placental lactogenic hormone as a parameter for placental function in renal transplanted women. *Acta Obstetrica et Gynecologica Scandinavica* 1990;**69**:41-3.

Kunz 1976 {published data only}

Kunz J, Keller PJ. Ultrasound and biochemical findings in intrauterine growth retardation. *Journal of Perinatal Medicine* 1976;**4**:85-94.

Laurin 1987 {published data only}

Laurin J, Persson PH. Ultrasound screening for detection of intra-uterine growth retardation. *Acta Obstetrica et Gynecologica Scandinavica* 1987; Vol. 66:493-500.

Leader 1980 {published data only}

Leader LR, Baillie P. Comparison of fetal movements and human placental lactogen as predictors of fetal outcome. A prospective study. *South African Medical Journal* 1980;**58**:609-10.

Lenstrup 1982 {published data only}

Lenstrup C. Predictive value of a single unstressed antepartum cardiotocogram in apparently uncomplicated pregnancy. Introduction of a new cardiotocography score. *Acta Obstetrica et Gynecologica Scandinavica* 1982; Vol. 61:177-82.

Lilford 1983 {published data only}

Lilford RJ, Obiekwe BC, Chard T. Maternal blood levels of human placental lactogen in the prediction of fetal growth retardation: choosing a cut-off point between normal and abnormal. *British Journal of Obstetrics and Gynaecology* 1983;**90**:511-5.

MacLeod 2013 {published data only}

MacLeod K, Sandercombe N, Barrett A, Heffernan E, Kalema H, Ngonzi J, et al. Clinical relevance of fetal weight estimation in Southwest Uganda. *Archives of Disease in Childhood: Fetal and Neonatal Edition* 2013;**98**:A15.

Mahran 1988 {published data only}

Mahran M, Omran M. The impact of diagnostic ultrasound on the prediction of intrauterine growth retardation in developing countries. *International Journal of Gynaecology and Obstetrics* 1988;**26**:375-8.

Marin 1979 {published data only}

Marin RD, Hood W. Significance of amniotic fluid glucose in late pregnancy. *Australian & New Zealand Journal of Obstetrics & Gynaecology* 1979;**19**:91-4.

McKenna 2005 {published data only}

McKenna D, Tharmaratnam S, Mahsud S, Dornan J. Ultrasonic evidence of placental calcification at 36 weeks' gestation: maternal and fetal outcomes. *Acta Obstetrica et Gynecologica Scandinavica* 2005;**84**:7-10.

Miller 1988 {published data only}

Miller JM Jr, Brown HL, Kissling GA, Gabert HA. The relationship of placental grade to fetal size and growth at term. *American Journal of Perinatology* 1988;**5**:19-21.

Molvarec 2013 {published data only}

Molvarec A, Gullai N, Stenczer B, Fugedi G, Nagy B, Rigo J. Comparison of placental growth factor and fetal flow Doppler ultrasonography to identify fetal adverse outcomes in women with hypertensive disorders of pregnancy: an observational study. *BMC Pregnancy and Childbirth* 2013;**13**:161.

Montan 1986 {published data only}

Montan S, Jorgensen C, Svalenius E, Ingemarsson I. Placental grading with ultrasound in hypertensive and normotensive pregnancies. A prospective, consecutive study. *Acta Obstetrica et Gynecologica Scandinavica* 1986; Vol. 65:477-80.

Nice 2016 {published and unpublished data}

Nice D, Hayden K, Higgins L, Garrod A, Johnstone E, Heazell A. Human placental lactogen and placental growth factor differentiate adverse pregnancy outcome from healthy outcomes in high-risk pregnancies. *BJOG: an international journal of obstetrics and gynaecology* 2016;**123**:44-5.

Nielsen 1985 {published data only}

Nielsen PV, Schultz-Larsen P, Schioeler V. Screening in pregnancy with unconjugated estriol compared with total estriol. *Acta Obstetrica et Gynecologica Scandinavica* 1985;**64**:297-301.

Nisbet 1982 {published data only}

Nisbet AD, Horne CH, Jandial V, Bremner RD, Cruickshank N, Sutcliffe RG. Measurement of plasma placental proteins and estriol in the detection of intrauterine growth retardation. *European Journal of Obstetrics, Gynecology, & Reproductive Biology* 1982;**13**:333-42.

Oats 1979 {published data only}

Oats JJ, Beischer NA. The recurrence rate and significance of low oestriol excretion in successive pregnancies. *British Journal of Obstetrics and Gynaecology* 1979;**86**:15-8.

Obiekwe 1983 {published data only}

Obiekwe BC, Chard T. A comparative study of the clinical use of four placental proteins in the third trimester. *Journal of Perinatal Medicine* 1983;**11**:121-6.

Odendaal 1981 {published data only}

Odendaal HJ, Malan C, Oosthuizen J. Hormonal placental functions and intrauterine growth retardation in patients with positive contraction stress tests. *South African Medical Journal* 1981;**59**:822-4.

Odendaal 1997 {published data only}

Odenaal HJ, Pienaar ME. Are high uric acid levels in patients with early preeclampsia an indication for delivery?. *South African Medical Journal* 1997;**87**:213-8.

Ott 1984 {published data only}

Ott WJ, Doyle S. Ultrasonic diagnosis of altered fetal growth by use of a normal ultrasonic fetal weight curve. *Obstetrics & Gynecology* 1984;**63**:201-4.

Palo 1987 {published data only}

Palo P, Erkkola R, Irjala K, Taina E. Serum free estriol inefficient in the detection of intrauterine growth retardation. *Annales Chirurgiae et Gynaecologiae* 1987;**Suppl 202**:20-2.

Palo 1989 {published data only}

Palo P, Erkkola R, Piironen O, Ruotsalainen P. Accuracy of ultrasonic fetal weight estimation and detection of small for gestational age fetuses. *American Journal of Perinatology* 1989; Vol. 6:400-4.

Patterson 1983 {published data only}

Patterson RM, Hayashi RH, Cavazos D. Ultrasonographically observed early placental maturation and perinatal outcome. *American Journal of Obstetrics and Gynecology* 1983;**147**:773-7.

Redman 1976 {published data only}

Redman CW, Beckin LJ, Bonnar J, Wilkinson RH. Plasma urate measurements in predicting fetal death in hypertensive pregnancy. *Lancet* 1976;**307**:1370-3.

Roma 2015 {published data only}

Roma E, Arnau A, Berdala R, Bergos C, Montesinos J, Figueras F. Ultrasound screening for fetal growth restriction at 36 vs 32weeks' gestation: a randomized trial (ROUTE). *Ultrasound in Obstetrics & Gynecology* 2015;**46**:391-7.

Sagen 1984 {published data only}

Sagen N, Nilsen ST, Kim HC, Koller O, Bergsjø P. The predictive value of total estriol; HPL and Hb on perinatal outcome in severe pre-eclampsia. *Acta Obstetrica et Gynecologica Scandinavica* 1984; Vol. 63:603-8.

Sekar 2016 {published data only}

Sekar R, Khatun M, Barrett HL, Duncombe G. A prospective pilot study in assessing the accuracy of ultrasound estimated fetal weight prior to delivery. *Australian & New Zealand Journal of Obstetrics & Gynaecology* 2016;**56**:49-53.

Shawkat 2015 {published and unpublished data}

Shawkat E, Johnstone E, Nice D, Sayce A, Hayden K, Myers J. [105-POS] : Clinical evaluation of placental growth factor (PLGF) for the management of suspected placental pathology in high-risk pregnancies. *Pregnancy Hypertension* 2015;**5**:57.

Siebert 1974 {published data only}

Siebert W, Meitinger C, Vogt W, Sandel P. [HPL serum levels in intrauterine retardation of fetal growth during late pregnancy (author's transl)]. *Geburtshilfe und Frauenheilkunde* 1974;**34**:520-4.

Skovron 1991 {published data only}

Skovron ML, Berkowitz GS, Lapinski RH, Kim JM, Chitkara U. Evaluation of early third-trimester ultrasound screening for intrauterine growth retardation. *Journal of Ultrasound in Medicine* 1991;**10**:153-9.

Sovio 2015 {published data only}

Sovio U, Smith G, Dacey A, Pasupathy D, White I. Screening for fetal growth restriction (FGR) using universal third trimester ultrasonography: A prospective cohort study of 3,977 nulliparous women. *American Journal of Obstetrics and Gynecology* 2015;**1**:S92.

Spernal 1989 {published data only}

Spernal R, Hecher K, Szalay S. [The value of fetal blood flow measurements in intrauterine growth retardation in comparison with E3 and human placental lactogen determinations]. *Geburtshilfe und Frauenheilkunde* 1989;**49**:463-5.

Steiner 1991 {published data only}

Steiner H, Schaffer H, Lassmann R, Staudach A, Batka M. [Comparison of biochemical and Doppler sonographic monitoring of high-risk pregnancies]. *Geburtshilfe und Frauenheilkunde* 1991;**51**:540-3.

Takeuchi 1985 {published data only}

Takeuchi K. [Antenatal assessment of intrauterine growth retardation by ultrasonic fetal biometry and non-stress fetal heart rate testing]. *Nippon Sanka Fujinka Gakkai Zasshi - Acta Obstetrica et Gynaecologica Japonica* 1985;**37**:2376-84.

Trudinger 1979 {published data only}

Trudinger BJ, Gordon YB, Grudzinskas JG, Hull MG, Lewis PJ, Arrans ME. Fetal breathing movements and other tests of fetal wellbeing: a comparative evaluation. *British Medical Journal* 1979;**2**:577-9.

Turitz 2014 {published data only}

Turitz AL. Isolated abdominal circumference <5% or estimated fetal weight 10 to 19% as predictors of small for gestational age infants. *American Journal of Perinatology* 2014; Vol. 31:469-76.

Valino 2016 {published data only}

Valino N, Giunta G, Gallo DM, Akolekar R, Nicolaides KH. Biophysical and biochemical markers at 35-37 weeks' gestation in the prediction of adverse perinatal outcome. *Ultrasound in Obstetrics & Gynecology* 2016;**47**:203-9.

Voto 1988 {published data only}

Voto LS, Ilia R, Darbon-Grosso A, Uranga Imaz F, Marguiles M. Uric acid levels: a useful index of the severity of preeclampsia and perinatal prognosis. *Journal of Perinatal Medicine* 1988;**16**:123-6.

Walker 2010 {published data only}

Walker GM, Hindmarsh, PC, Geary M, Kingdom JC. Sonographic maturation of the placenta at 30 to 34 weeks is not associated with second trimester markers of placental insufficiency in low-risk pregnancies. *Journal of Obstetrics and Gynaecology Canada* 2010;**32**(12):1134-9.

Weerasinghe 1977 {published data only}

Weerasinghe DS, Legge M, Aickin DR. The clinical significance of low urine oestriol values. *New Zealand Medical Journal* 1977;**86**:383-4.

Weiner 2016 {published data only}

Weiner E, Fainstein N, Mizrahi Y, Elyashiv O, Mevorach-Zussman N, Bar J, et al. Comparison between three methods for the detection of macrosomia and growth restriction in patients presenting in active labor-a prospective study. *American Journal of Obstetrics and Gynecology* 2016;**1**:S225-6.

Westergaard 1984 {published data only}

Westergaard JG, Teisner B, Hau J, Grudzinskas JG. Placental protein measurements in complicated pregnancies. I. Intrauterine growth retardation. *British Journal of Obstetrics and Gynaecology* 1984;**91**:1216-23.

Williams 2002 {published data only}

Williams KP, Galerneau F. The role of serum uric acid as a prognostic indicator of the severity of maternal and fetal complications in hypertensive pregnancies. *Journal of Obstetrics & Gynaecology Canada: JOGC* 2002;**24**:628-32.

Yassaee 2003 {published data only}

Yassaee F. Hyperuricaemia and perinatal outcomes in patients with severe preeclampsia. *Iranian Journal of Medical Sciences* 2003;**28**(4):198-9.

Ylikorkala 1973 {published data only}

Ylikorkala O. Maternal serum HPL levels in normal and complicated pregnancy as an index of placental function. *Acta Obstetrica et Gynecologica Scandinavica - Supplement* 1973;**26**:1-52.

Zhang 1990 {published data only}

Zhang WY. [Single radial immunodiffusion of human placental lactogen and its clinical uses]. *Chung-Hua Fu Chan Ko Tsa Chih [Chinese Journal of Obstetrics & Gynecology]* 1990;**25**:278-81, 316.

References to studies excluded from this review
Adekanle 2013 {published data only}

Adekanle DA, Oparinde DP, Atiba AS, Akintayo AA. Effect of different modes of delivery on cord blood oxidative stress markers. *International Journal of Biomedical Science* 2013;**9**:249-54.

Agboola 1978 {published data only}

Agboola A. The effect of low haematocrit on serum human placental lactogen values. *British Journal of Obstetrics & Gynaecology* 1978; Vol. 85, issue 10:758-60.

Aggarwal 2006 {published data only}

Aggarwal PK, Jain V, Sakhuja V, Karumanchi SA, Jha V. Low urinary placental growth factor is a marker of pre-eclampsia. *Kidney International* 2006;**69**:621-4.

Agorastos 2014 {published data only}

Agorastos T. Biomarkers of endothelial dysfunction in preeclampsia and neonatal morbidity: a case-control study. *European Journal of Obstetrics, Gynecology, and Reproductive Biology* 2014;**175**:119-23.

Ahmad 1979 {published data only}

Ahmad T, Rahat T. Diagnostic evaluation of pregnancy urinary oestriol for foeto-placental function. *JPMA - Journal of the Pakistan Medical Association* 1979; Vol. 29, issue 10:209-11.

Aickin 1983 {published data only}

Aickin DR, Duff GB, Evans JJ, Legge M. Antenatal biochemical screening to predict low birthweight infants. *British Journal of Obstetrics and Gynaecology* 1983;**90**:129-33.

Alahakoon 2014 {published data only}

Alahakoon TI, Zhang WY, Trudinger B, Lee V. Discordant clinical presentations of preeclampsia and intrauterine fetal growth restriction with similar pro and anti-angiogenic profiles. *Reproductive Sciences* 2014;**1**:185A.

Al-Amin 2015 {published data only}

Al-Amin A, Hingston T, Mayall P, Araujo E, Fabricio Da Silva C, Friedman D. The utility of ultrasound in late pregnancy compared with clinical evaluation in detecting small and large for gestational age fetuses in low-risk pregnancies. *Journal of Maternal-Fetal and Neonatal Medicine* 2015;**28**:1495-9.

Alberry 2009 {published data only}

Alberry MS, Maddocks DG, Hadi MA, Metawi H, Hunt LP, Abdel-Fattah SA, et al. Quantification of cell free fetal DNA in maternal plasma in normal pregnancies and in pregnancies with placental dysfunction. *American Journal of Obstetrics and Gynecology* 2009;**200**:98.e1-6.

Algeri 2013 {published data only}

Algeri P, Ornaghi S, Bernasconi D, Cappellini F, Signorini S, Brambilla P, et al. The role of placental dysfunction in the pathogenesis of IUGR. *American Journal of Obstetrics and Gynecology* 2013;**208** (1 Suppl 1):S192-3.

Alvarez-Fernandez 2014 {published data only}

Alvarez-Fernandez I, Prieto B, Rodriguez V, Ruano Y, Escudero AI, Alvarez FV. New biomarkers in diagnosis of early onset preeclampsia and imminent delivery prognosis. *Clinical Chemistry & Laboratory Medicine* 2014; Vol. 52:1159-68.

Alwasel 2013 {published data only}

Alwasel SH, Harrath AH, Aljarallah JS, Abotalib Z, Osmond C, Al Omar SY, et al. The velocity of fetal growth is associated with the breadth of the placental surface, but not with the length. *American Journal of Human Biology* 2013;**25**:534-7.

Anastasakis 2008 {published data only}

Anastasakis E, Papantoniou N, Daskalakis G, Mesogitis S, Antsaklis A. Screening for pre-eclampsia by oxidative stress markers and uteroplacental blood flow. *Journal of Obstetrics and Gynaecology* 2008;**28**:285-9.

Anderson 1978 {published data only}

Anderson SG. Real-time sonography in obstetrics. *Obstetrics & Gynecology* 1978;**51**:284-7.

Arabin 1993 {published data only}

Arabin B, Snyders R, Nicolaidis KH, Versmold HK, Weitzel HK, Giffel J, et al. Systematic antepartum fetal evaluation ("Safe"). A concept for diagnosis of fetal function in threatened hypoxia. *Geburtshilfe und Frauenheilkunde* 1993; Vol. 53, issue 12:835-42.

Arabin 1995 {published data only}

Arabin B, Ragosch V, Mohnhaupt A. From biochemical to biophysical placental function tests in fetal surveillance. *American Journal of Perinatology* 1995; Vol. 12, issue 3:168-71.

Arias 1977 {published data only}

Arias F. The diagnosis and management of intrauterine growth retardation. *Obstetrics & Gynecology* 1977;**49**:293-8.

Ariyuki 1995 {published data only}

Ariyuki Y, Hata T, Kitao M. Evaluation of perinatal outcome using individualized growth assessment: comparison with conventional methods. *Pediatrics* 1995;**96**:36-42.

Atzeni 2012 {published data only}

Atzeni I, Paoletti AM, Deiana SF, Meloni A, Guerriero S, Parodo G, et al. Placental growth factor (PLGF): Correlations with placental function. *International Journal of Gynecology and Obstetrics* 2012;**119**:S782-3.

Aviram 2015 {published data only}

Aviram A, Yogev Y, Bardin R, Meizner I, Wiznitzer A, Hadar E. Small for gestational age newborns - Does pre-recognition make a difference in pregnancy outcome?. *Journal of Maternal-fetal & Neonatal Medicine* 2015;**28**:1520-4.

Axelsson 1978 {published data only}

Axelsson O, Nilsson BA, Johansson ED. Assessment of placental function in uncomplicated and complicated late pregnancy by estimation of unconjugated oestrogens in plasma after an intravenous injection of dehydroepiandrosterone sulphate. *Acta Endocrinologica* 1978;**89**:359-71.

Baeza Valenzuela 1995 {published data only}

Baeza Valenzuela A, Garcia Mendez A. Premature aging of the placenta. Ultrasonic diagnosis. [Spanish]. *Ginecologia y Obstetricia de Mexico* 1995;**63**:287-92.

Bahado-Singh 1998 {published data only}

Bahado-Singh RO, Dashe J, Deren O, Daftary G, Copel JA, Ehrenkranz RA. Prenatal prediction of neonatal outcome in the extremely low-birth-weight infant. *American Journal of Obstetrics and Gynecology* 1998;**178**:462-8.

Bainbridge 2008 {published data only}

Bainbridge SA, Roberts JM. Uric acid as a pathogenic factor in preeclampsia. *Placenta* 2008;**29** Suppl A:S67-S72.

Bakketeig 1984 {published data only}

Bakketeig LS, Eik-Nes SH, Jacobsen G. Randomised controlled trial of ultrasonographic screening in pregnancy. *Lancet* 1984;**2**:207-11.

Baltajian 2016 {published data only}

Baltajian K, Bajracharya S, Salahuddin S, Berg AH, Geahchan C, Wenger JB, et al. Sequential plasma angiogenic factors levels in women with suspected preeclampsia. *American Journal of Obstetrics & Gynecology* 2016;**215**:e1-89.e10.

Barden 1999 {published data only}

Barden AE, Beilin LJ, Ritchie J, Walters BN, Graham D, Michael CA. Is proteinuric pre-eclampsia a different disease in primigravida and multigravida?. *Clinical Science* 1999;**97**:475-83.

Bardien 2016 {published data only}

Bardien N, Whitehead CL, Tong S, Ugoni A, McDonald S, Walker SP. Placental insufficiency in fetuses that slow in growth but are born appropriate for gestational age: a prospective longitudinal study. *PLOS One* 2016;**11**:e0142788.

Baron 1996 {published data only}

Baron F, Graham D. Age-independent assessment of abnormal fetal growth in the third trimester. *Journal of Ultrasound in Medicine* 1996;**15**:381-4.

Barrilleaux 2007 {published data only}

Barrilleaux PS, Magann EF, Chauhan SP, York BM, Philibert L, Lewis DF. Amniotic fluid index as a predictor of adverse perinatal outcome in the HELLP syndrome. *Journal of Reproductive Medicine* 2007;**52**:293-8.

Bartha 2003 {published data only}

Bartha JL, Romero-Carmona R, Escobar-Llompant M, Paloma-Castro O, Comino-Delgado R. Human chorionic gonadotropin and vascular endothelial growth factor in normal and complicated pregnancies. *Obstetrics and Gynecology* 2003;**102**:995-9.

Bashir 1982 {published data only}

Bashir T, Brash JH, Buchan PC, Bevis DC, Clayden AD. Screening for the small-for-dates: a clinical and biochemical appraisal. *European Journal of Obstetrics, Gynecology, & Reproductive Biology* 1982;**13**:61-6.

Bastek 2009 {published data only}

Bastek JA, Parr E, Wang E, Elovitz MA, Srinivas SK. Limitations of ultrasound in diagnosing intrauterine growth restriction in severe preeclampsia. *Journal of Maternal-fetal & Neonatal Medicine* 2009;**22**:1039-44.

Battaglia 1995 {published data only}

Battaglia C, Artini PG, Ballestri M, Bonucchi D, Galli PA, Bencini S, et al. Hemodynamic, hematological and hemorrhological evaluation of post-term pregnancy. *Acta Obstetrica et Gynecologica Scandinavica* 1995;**74**:336-40.

Beischer 1975 {published data only}

Beischer NA. Low oestriol excretion: incidence, significance and treatment in an obstetric population. *Medical Journal of Australia* 1975;**2**:379-81.

Bell 1967 {published data only}

Bell ET, Loraine JA, McEwan HP, Charles D. Serial hormone assays in patients with uteroplacental insufficiency. *American Journal of Obstetrics and Gynecology* 1967;**97**:562-70.

Benton 2010 {published data only}

Benton S, Hu Y, Fang X, Knudsen AB, Kronborg CS, Vittinghus E, et al. Placental growth factor as a diagnostic for early onset pre-eclampsia and normotensive intrauterine growth restriction of placental origin. *Placenta* 2010; Vol. 1:S24.

Benton 2011 {published data only}

Benton S, Hu Y, Xie F, Kupfer K, Lee SW, Magee L, et al. Can placental growth factor identify placental intra-uterine growth restriction in fetuses who are antenatally identified as being small for gestational age?. *Placenta* 2011;**32**(9):A100.

Benton 2011a {published data only}

Benton S, Hu Y, Xie F, Kupfer K, Lee SW, Magee L, et al. Placental growth factor quantified on rapid assay as a diagnostic test for early-onset pre-eclampsia. *Placenta* 2011;**32**(9):A84.

Benton 2012 {published data only}

Benton S, Knudsen U, Hu Y, Vittinghus E, Allen J, Konborg C, et al. Angiogenic factor imbalance is not specific to the maternal syndrome of pre-eclampsia. *Reproductive Sciences* 2012;**1**:317A-8A.

Benton 2012a {published data only}

Benton SJ, Hu X, Xie F, Kupfer K, Lee SW, Magee LA, et al. Can placental growth factor in maternal circulation identify fetuses with placental intrauterine growth restriction?. *American Journal of Obstetrics and Gynecology* 2012;**206**(2):163.

Benton 2014 {published data only}

Benton S, Hu Y, Magee L, Von Dadelszen P. Angiogenic factors and the prediction of complications in a high-risk pregnancy cohort. *Reproductive Sciences* 2014;**1**:186A-187A.

Benton 2014a {published data only}

Benton S, Yockell-Lelievre J, Grynspan D, Magee L, Hu Y, Gruslin A, et al. Low maternal placental growth factor is associated with abnormal placental morphology in fetuses with suspected intrauterine growth restriction. *Placenta* 2014;**35**(9):A44.

Benz 1980 {published data only}

Benz J, Ingold W, Landolt A, Keller PJ. [Hormonal diagnosis in fetal intrauterine retardation]. *Gynakologische Rundschau* 1980;**20** Suppl 1:74-6.

Berendtsen 1985 {published data only}

Berendtsen H, Klunder KB, Norskov O. Biparietal measurements, human placental-lactogenic hormone and oestriol in assessment of foetal condition. *Ugeskrift for Laeger* 1985; Vol. 147, issue 38:2984-6.

Bergsjö 1973 {published data only}

Bergsjö P, Bakke T, Salamonsen LA, Stoa KF, Thorsen T. Urinary oestriol in pregnancy, daily fluctuation, and correlation with fetal growth. *Journal of Obstetrics & Gynaecology of the British Commonwealth* 1973;**80**:305-10.

Berle 1973 {published data only}

Berle P. [Value of hormone analysis for the assessment of placenta function in late pregnancy]. *Deutsche Medizinische Wochenschrift* 1973;**98**:2146-50.

Berle 1973a {published data only}

Berle P. Maternal serum levels of human placental lactogen in placental insufficiency. *Geburtshilfe und Frauenheilkunde* 1973; Vol. 33, issue 6:455-9.

Berle 1973b {published data only}

Berle P, van Leyen H, Rossler H. [Proceedings: Diagnosis of placenta insufficiency by determination of placental lactogen in maternal blood]. *Archiv für Gynakologie* 1973;**214**:213-5.

Berle 1975 {published data only}

Berle P, Haselmayer B, Plambeck H. [Comparative studies on CTG and hormonal parameters in the diagnosis of placenta insufficiency]. *Archiv für Gynakologie* 1975;**219**:426-8.

Bernardes 2013 {published data only}

Bernardes LS, Francisco RP, Dourado RM, Doro GF, Zugaib M. Ultrasound evaluation of the placenta in fetuses of IUGR. *Placenta* 2013;**34**:A92.

Bernatavicius 2013 {published data only}

Bernatavicius G, Roberts S, Garrod A, Whitworth MK, Johnstone ED, Gillham JC, et al. A feasibility study for a randomised controlled trial of management of reduced fetal movements after 36 weeks gestation. *Archives of Disease in Childhood: Fetal and Neonatal Edition* 2013;**98**:35.

Bersinger 2004 {published data only}

Bersinger NA, Odegard RA. Second- and third-trimester serum levels of placental proteins in preeclampsia and small-for-gestational age pregnancies. *Acta Obstetrica et Gynecologica Scandinavica* 2004;**83**:37-45.

Bersinger 2005 {published data only}

Bersinger NA, Odegard RA. Serum levels of macrophage colony stimulating, vascular endothelial, and placenta growth factor in relation to later clinical onset of pre-eclampsia and a small-for-gestational age birth. *American Journal of Reproductive Immunology* 2005;**54**:77-83.

Bhansali 1975 {published data only}

Bhansali KG. Placental function tests by blood analysis. *Journal of the National Medical Association* 1975;**67**:137-9.

Bian 1992 {published data only}

Bian XM. [Choice of ultrasound fetal weight estimation formulae]. *Chung-Hua i Hsueh Tsa Chih [Chinese Medical Journal]* 1992;**72**:677-9, 703.

Biberoglu 2016 {published data only}

Biberoglu E, Kirbas A, Daglar K, Biberoglu K, Timur H, Demirtas C, et al. Serum angiogenic profile in abnormal placentation. *Journal of Maternal-Fetal & Neonatal Medicine* 2016;**29**:3193-7.

Bieglmayer 1981 {published data only}

Bieglmayer C, Spona J. [Monitoring of at-risk pregnancies through E3 and HPL]. *Acta Universitatis Palackianae Olomucensis Facultatis Medicae* 1981;**101**:38-42.

Bila 1980 {published data only}

Bila S, Sulovic V, Genbacev O. [Human placental lactogen in the blood in fetal growth retardation]. *Srpski Arhiv Za Celokupno Lekarstvo* 1980;**108**:317-24.

Bitzer 1985 {published data only}

Bitzer J, Hendry M, Richter R. [Detection of intrauterine growth retardation within the scope of routine prenatal care]. *Geburtshilfe und Frauenheilkunde* 1985; Vol. 45, issue 2:79-82.

Blaskova 1977 {published data only}

Blaskova O, Sasko A, Pont'uch A, Chabada J. The value of estriol determination, and cytologic, ultrasound, and amnioscopic investigation in high-risk pregnancy with a view to placental insufficiency (author's transl). [Slovak]. *Ceskoslovenska Gynekologie* 1977;**42**:411-4.

Bligh 2015 {published data only}

Bligh L, Al Solai A, Greer RM, Kumar S. Intrapartum fetal compromise is associated with lower maternal placental growth factor (PLGF) levels at term. *BJOG: an international journal of obstetrics and gynaecology* 2015;**122**:162-3.

Blitz 2016 {published data only}

Blitz MJ, Rochelson B, Vohra N. Maternal serum analyses as predictors of fetal growth restriction with different degrees of placental vascular dysfunction. *Clinics in Laboratory Medicine* 2016;**36**:353-67.

Blumenfeld 2007 {published data only}

Blumenfeld Y, Lee H, Pullen K, Wong A, Pettit K, Taslimi M. Ultrasound accuracy of estimated fetal weight in pregnancies complicated by intra-uterine growth restriction. *American Journal of Obstetrics and Gynecology* 2007;**197**:S59.

Bobrow 2002 {published data only}

Bobrow CS, Holmes RP, Muttukrishna S, Mohan A, Groome N, Murphy DJ, et al. Maternal serum activin A, inhibin A, and follistatin in pregnancies with appropriately grown and small-for-gestational-age fetuses classified by umbilical artery Doppler ultrasound. *American Journal of Obstetrics and Gynecology* 2002;**186**:283-7.

Bock 1976 {published data only}

Bock JE, Gaede P, Trolle D. Human placental lactogen hormone in serum from pregnant women with rhesus (anti-D) isoimmunization. *Acta Obstetrica et Gynecologica Scandinavica - Supplement* 1976; Vol. 53:14-9.

Boij 2012 {published data only}

Boij R, Svensson J, Nilsson-Ekdahl K, Sandholm K, Lindahl TL, Palonek E, et al. Biomarkers of coagulation, inflammation, and angiogenesis are independently associated with preeclampsia. *American Journal of Reproductive Immunology* 2012;**68**:258-70.

Borges 2005 {published data only}

Borges MH, Zamudio S, Pullockaran J, Belliappa S, Albieri A, Catalano PM, et al. Fetal circulating growth factors in diabetic and normal pregnancy: relationship to placental and fetal growth. *Journal of the Society for Gynecologic Investigation* 2005; Vol. 12:299A.

Botasheva 2016 {published data only}

Botasheva TL, Linde VA, Ermolova NV, Palieva NV, Sargsyan OD, Barinova VV. Peculiarities of angiogenic factors and cytokines in the physiological and complicated pregnancy, depending on the sex of the fetus. *Journal of Maternal-Fetal and Neonatal Medicine* 2016;**29**:134.

Boucoiran 2012 {published data only}

Boucoiran I, Thissier-Levy S, Wu Y, Wei SQ, Zhong-Cheng L, Delvin E, et al. Risk for preeclampsia and intrauterine growth restriction: Effective value of PLGF, Sflt-1 and Inhibin A in singleton and multiple pregnancies. *American Journal of Obstetrics and Gynecology* 2012;**1**:S336-7.

Branconi 1981 {published data only}

Branconi F, Cariati E, Cappelli G, Paladini S, Scarselli G. Plasma hormone assay in pregnancy. *International Journal of Biological Research in Pregnancy* 1981;**2**:146-8.

Brush 1970 {published data only}

Brush MG, Maxwell R, Scherer J, Taylor RW, Tye G. Placental failure in the small-for-dates syndrome. *Proceedings of the Royal Society of Medicine* 1970; Vol. 63:1098-9.

Bukowski 2014 {published data only}

Bukowski R, Hansen NI, Willinger M, Reddy UM, Parker CB, Pinar H, et al. Fetal growth and risk of stillbirth: a population-based case-control study. *PLOS Medicine* 2014;**11**:e1001633.

Butcher 2012 {published data only}

Butcher I, Whatmore A, Bonshek C, Shaikh G, Dunn W, Brown M, et al. Metabolic profiles identify markers of catch-up growth in children born small for gestational age (SGA). *Archives of Disease in Childhood* 2012;**97**:A102.

Buyon 2011 {published data only}

Buyon JP, Garabet L, Kim M, Reeves ER, Guerra MM, Lockshin MD, et al. Favorable prognosis in a large, prospective multicenter study of lupus pregnancies. *Arthritis and Rheumatism* 2011;**63**(S10):1707.

Cage 2009 {published data only}

Cage CR, Kreither M, Coussons-Read M, Cole S, Brandt C, Lobel M. Pregnancy-related stress influences maternal corticotropin-releasing factor and estradiol concentrations affecting birth outcome. *Brain, Behavior, and Immunity* 2009;**23**:S29-30.

Calabrese 2012 {published data only}

Calabrese S, Cardellicchio M, Mazzocco M, Taricco E, Martinelli A, Cetin I. Placental growth factor (PLGF) maternal circulating levels in normal pregnancies and in pregnancies at risk of developing placental insufficiency complications. *Reproductive Sciences* 2012; Vol. 1:211A-2A.

Calderon 2011 {published data only}

Calderon AC, Berezowski AT, Marcolin AC, Martins WP, Duarte G, Cavalli RC. Ultrasonography in pregnant women with antiphospholipid syndrome using salicylic acid and heparin. *Archives of Gynecology & Obstetrics* 2011;**284**:79-84.

Campobasso 1967 {published data only}

Campobasso M, Pontrelli VC, Longo M. [Chorionic gonadotropins as an index of placental function in normal and pathological pregnancy]. *Minerva Ginecologica* 1967;**19**:133-8.

Camus-Bablon 1990 {published data only}

Camus-Bablon F, Cohen RM, Berk MA, Perisutti G, Hunter K, Frohman LA. Alterations in circulating human chorionic gonadotropin free alpha-subunit in insulin-dependent diabetic pregnancy: correlation with maternal characteristics. *Journal of Clinical Endocrinology & Metabolism* 1990; Vol. 71:46-52.

Carne 1987 {published data only}

Carne RJ, Drew JH. Infant development following the use of intravenous nutrition to women with persistently low urinary oestriol excretion. *Australian & New Zealand Journal of Obstetrics & Gynaecology* 1987;**27**:30-6.

Castren 1966 {published data only}

Castren O, Rauramo L, Pekkarinen A. Excretion of oestriol and pregnanediol in placental insufficiency. *Acta Obstetrica et Gynecologica Scandinavica* 1966;**45**(Suppl 9):66-7.

Cavazza 2015 {published data only}

Cavazza MC, Pinto L, Graca L. Intrauterine growth restriction- a three-year experience at a tertiary care hospital in Portugal. *Journal of Perinatal Medicine. Conference: 12th World Congress of Perinatal Medicine* 2015;**43**:639.

Ceccarello 1980 {published data only}

Ceccarello PL, Calcagnile F, Varagnolo C, Castello C, Destro F. [Fetal ultrasonic biometry of the small-for-date fetus]. *Minerva Ginecologica* 1980;**32**:107-12.

Cefalo 2005 {published data only}

Cefalo RC. [Commentary on] Value of biochemical markers for outcome in term infants with asphyxia. *Obstetrical & Gynecological Survey* 2005;**60**:293-5.

Cetin 2014 {published data only}

Cetin I, Mazzocco M, Giardini V, Calabrese S, Algeri P, Martinelli A, et al. Placental growth factor for the prediction of fetal outcomes in pregnancies at risk and in IUGR. *Journal of Maternal-Fetal and Neonatal Medicine* 2014; Vol. 27:135.

Cetin 2016 {published data only}

Cetin I, Mazzocco MI, Giardini V, Cardellicchio M, Calabrese S, Algeri P, et al. PLGF in a clinical setting of pregnancies at risk of preeclampsia and/or intrauterine growth restriction. *Journal of Maternal-Fetal and Neonatal Medicine* 2016;**30**(2):144-9.

Chaiworapongsa 2008 {published data only}

Chaiworapongsa T, Espinoza J, Gotsch F, Kim YM, Kim GJ, Goncalves LF, et al. The maternal plasma soluble vascular endothelial growth factor receptor-1 concentration is elevated in SGA and the magnitude of the increase relates to Doppler abnormalities in the maternal and fetal circulation. *Journal of Maternal-Fetal & Neonatal Medicine* 2008;**21**:25-40.

Chaiworapongsa 2012 {published data only}

Chaiworapongsa T, Romero R, Korzeniewski SJ, Kusanovic JP, Soto E, Hernandez-Andrade E, et al. Prediction of stillbirth and late-onset preeclampsia. *Reproductive Sciences* 2012;**1**:90A-1A.

Chaiworapongsa 2013a {published data only}

Chaiworapongsa T, Whitten AE, Romero R, Korzeniewski SJ, Schwartz A, Chaemsaitong P, et al. Biomarkers identify mothers with small for gestational age fetuses who will develop preeclampsia or require an indicated early preterm delivery. *Journal of Perinatal Medicine* 2013;**41**(S1):1.

Chaiworapongsa 2013b {published data only}

Chaiworapongsa T, Whitten AE, Romero R, Korzeniewski SJ, Schwartz A, Cortez J C, et al. Biomarkers in maternal blood predict neonatal morbidity. *Journal of Perinatal Medicine* 2013;**41**(S1):1.

Chambers 1989 {published data only}

Chambers SE, Hoskins PR, Haddad NG, Johnstone FD, McDicken WN, Muir BB. A comparison of fetal abdominal circumference measurements and Doppler ultrasound in the prediction of small-for-dates babies and fetal compromise. *British Journal of Obstetrics and Gynaecology* 1989;**96**:803-8.

Chang 1993 {published data only}

Chang TC, Robson SC, Spencer JA, Gallivan S. Identification of fetal growth retardation: comparison of Doppler waveform indices and serial ultrasound measurements of abdominal circumference and fetal weight. *Obstetrics & Gynecology* 1993;**82**:230-6.

Chang 1994 {published data only}

Chang TC, Robson SC, Spencer JA, Gallivan S. Prediction of perinatal morbidity at term in small fetuses: comparison of fetal

growth and Doppler ultrasound. *British Journal of Obstetrics and Gynaecology* 1994;**101**:422-7.

Chapman 1978 {published data only}

Chapman MG, Jones WR. Pregnancy-specific beta-1 glycoprotein (SP-1) in normal and abnormal pregnancy. *Australian & New Zealand Journal of Obstetrics & Gynaecology* 1978;**18**:172-5.

Chapman 1981 {published data only}

Chapman MG, O'Shea RT, Jones WR, Hillier R. Pregnancy-specific beta 1-glycoprotein as a screening test for at-risk pregnancies. *American Journal of Obstetrics and Gynecology* 1981; Vol. 141:499-502.

Chappell 2002 {published data only}

Chappell LC, Seed PT, Briley A, Kelly FJ, Hunt BJ, Charnock-Jones DS, et al. A longitudinal study of biochemical variables in women at risk of preeclampsia. *American Journal of Obstetrics and Gynecology* 2002;**187**:127-36.

Chard 1982 {published data only}

Chard T. Human placental lactogen in the monitoring of high-risk pregnancy. *Ricerca in Clinica e in Laboratorio* 1982;**12**:207-20.

Chauhan 2012 {published data only}

Chauhan S, Tajik P, Boers K, Wyk L, Mol BW, Scherjon S. Differentiating newborns with birth weight < vs > 3 percentiles for gestational age: Secondary analysis of a randomized clinical trial (DIGITAT). *American Journal of Obstetrics and Gynecology* 2012; Vol. 1:S177-s178.

Chawengsettakul 2015 {published data only}

Chawengsettakul S, Russameecharoen K, Wanitpongpan P. Fetal cardiac function measured by myocardial performance index of small-for-gestational age fetuses. *Journal of Obstetrics and Gynaecology Research* 2015;**41**:222-8.

Chew 2014 {published data only}

Chew C, Crocker I, Kiss O, Allibone A, Tower C, Bruce IN. Placental growth factor is reduced in lupus patients with adverse pregnancy outcomes. *Rheumatology* 2014;**53**:i182.

Church 2016 {published data only}

Church E, Bellis A. Impact of education of pregnant women regarding fetal movements and improved detection and assessment of small-for-gestational-age babies over a year in a District General Hospital. *BJOG: an international journal of obstetrics and gynaecology* 2016;**123**:88.

Clelia 2013 {published data only}

Cabo Fustaret MC, Escobar A, Illia R, Uranga M, Rivas C, Lobenstein G, et al. NT-Pro-BNP: Correlation with adverse outcome markers in hypertensive gestational syndromes. *Pregnancy Hypertension* 2013;**3**(2):84-5.

Clowse 2011 {published data only}

Clowse ME, Criscione-Schreiber LG, Pisetsky DS. Predictors of preterm birth and preeclampsia in systemic lupus erythematosus. *Arthritis and Rheumatism* 2011;**63**(S1):2278.

Cody 2013 {published data only}

Cody F, Unterscheider J, Daly S, Geary MP, Kennelly MM, McAuliffe FM, et al. Impact of maternal obesity on accuracy of sonographic fetal weight estimation in IUGR. *Archives of Disease in Childhood: Fetal and Neonatal Edition* 2013;**98**(S1):A30.

Cody 2016 {published data only}

Cody F, Unterscheider J, Daly S, Geary MP, Kennelly MM, McAuliffe FM, et al. The effect of maternal obesity on sonographic fetal weight estimation and perinatal outcome in pregnancies complicated by fetal growth restriction. *Journal of Clinical Ultrasound* 2016;**44**:34-9.

Cooley 2011 {published data only}

Cooley SM, Donnelly JC, Walsh T, McMahon C, Gillan J, Geary MP. The impact of ultrasonographic placental architecture on antenatal course, labor and delivery in a low-risk primigravid population. *Journal of Maternal-Fetal & Neonatal Medicine* 2011;**24**:493-7.

Cordano 1988 {published data only}

Cordano MC, Comte E, Bessis R, Papiernik E. Longitudinal follow-up of 100 patients at risk of intrauterine growth retardation: comparison of diagnosis in two periods. *Journal of Perinatal Medicine* 1988;**16**:61-6.

Craig 1996 {published data only}

Craig SD, Beach ML, Harvey-Wilkes KB, D'Alton ME. Ultrasound predictors of neonatal outcome in intrauterine growth restriction. *American Journal of Perinatology* 1996; Vol. 13, issue 8:465-71.

Crane 1979 {published data only}

Crane JP, Kopta MM. Prediction of intrauterine growth retardation via ultrasonically measured head/abdominal circumference ratios. *Obstetrics & Gynecology* 1979;**54**:597-601.

Crawford 1985 {published data only}

Crawford DC, Fenton DW, Price WL. Ultrasonic tissue characterization of the placenta: Is it of clinical value?. *Journal of Clinical Ultrasound* 1985;**13**:533-7.

D'Anna 2000 {published data only}

D'Anna R, Baviera G, Scilipoti A, Leonardi I, Leo R. The clinical utility of serum uric acid measurements in pre-eclampsia and transient hypertension in pregnancy. *Panminerva Medica* 2000;**42**:101-3.

Daikoku 1979 {published data only}

Daikoku NH, Tyson JE, Graf C, Scott R, Smith B, Johnson JW, et al. The relative significance of human placental lactogen in the diagnosis of retarded fetal growth. *American Journal of Obstetrics & Gynecology* 1979; Vol. 135, issue 4:516-21.

Darling 2014 {published data only}

Darling AM, McDonald CR, Conroy AL, Hayford KT, Liles WC, Wang M, et al. Angiogenic and inflammatory biomarkers in midpregnancy and small-for-gestational-age outcomes in Tanzania. *American Journal of Obstetrics and Gynecology* 2014;**211**:509.e1-8.

Dave 2016 {published data only}

Dave A, Maru L, Jain A. LDH (lactate dehydrogenase): a biochemical marker for the prediction of adverse outcomes in pre-eclampsia and eclampsia. *Journal of Obstetrics & Gynaecology of India* 2016;**66**:23-9.

Dawood 1976 {published data only}

Dawood MY. Circulating maternal serum progesterone in high-risk pregnancies. *American Journal of Obstetrics and Gynecology* 1976;**125**:832-40.

Del Moral 2015 {published data only}

Del Moral R, Gozalo M, Spies K, Joigneau L, Martinez R. Diagnosis and follow-up of in patients diagnosed with intrauterine growth restriction in our center. *Journal of Perinatal Medicine. Conference: 12th World Congress of Perinatal Medicine* 2015;**43**:1.

De Marchi 1977 {published data only}

De Marchi C, Santi F, Zauli F, Roberti L, Perolo F. [Fetal growth studied by echographic evaluation and by placental lactogen and estradiol levels]. *Rivista Italiana di Ginecologia* 1977;**58**:287-93.

Deter 2016 {published data only}

Deter RL, Levytska K, Melamed N, Lee W, Kingdom JC. Classifying neonatal growth outcomes: use of birth weight, placental evaluation and individualized growth assessment. *Journal of Maternal-Fetal & Neonatal Medicine* 2016;**29**:3939-49.

Di Lorenzo 2013 {published data only}

Di Lorenzo G, Monasta L, Ceccarello M, Cecotti V, D'Ottavio G. Third trimester abdominal circumference, estimated fetal weight and uterine artery doppler for the identification of newborns small and large for gestational age. *European Journal of Obstetrics, Gynecology, & Reproductive Biology* 2013;**166**:133-8.

Dombrowski 1992 {published data only}

Dombrowski MP, Saleh AA, Berry SM, Cotton DB. Perinatal outcome with sonographically thin placentas. *Journal of Maternal-Fetal Medicine* 1992;**1**:137-9.

Dombrowski 1992a {published data only}

Dombrowski MP, Wolfe HM, Saleh A, Evans MI, O'Brien J. The sonographically thick placenta: a predictor of increased perinatal morbidity and mortality. *Ultrasound in Obstetrics & Gynecology* 1992;**2**:252-5.

Ducarme 2012 {published data only}

Ducarme G, Seguro E, Chesnoy V, Davitian C, Luton D. [Estimation of fetal weight by external abdominal measurements and fundal height measurement near term for the detection of intra-uterine growth retardation]. *Gynecologie, Obstetrique & Fertilité* 2012;**40**:642-6.

Duff 1986 {published data only}

Duff GB. The realities of screening for the small for dates fetus using ultrasound measurement. *Australian & New Zealand Journal of Obstetrics & Gynaecology* 1986;**26**:102-5.

Dutton 2012 {published data only}

Dutton P, Warrander L, Bernatavicius G, Kroll J, Gaze D, Jones R, et al. Placentally-derived factors in women with reduced fetal movements and their relationship to pregnancy outcome. *BJOG: an international journal of obstetrics and gynaecology* 2012;**119**(6):e3.

Dutton 2012a {published data only}

Dutton PJ, Warrander LK, Roberts SA, Bernatavicius G, Byrd LM, Gaze D, et al. Predictors of poor perinatal outcome following maternal perception of reduced fetal movements - a prospective cohort study. *PLOS One* 2012;**7**:7.

Eik-Nes 1984 {published data only}

Eik-Nes SH, Okland O, Aure JC, Ulstein M. Ultrasound screening in pregnancy: a randomised controlled trial. *Lancet* 1984; Vol. 1:1347.

El-Ahmady 1997 {published data only}

El-Ahmady O, Halim AB, Saleh AA, Ismail MT, Amin AF. Serum alpha-fetoprotein (AFP) and beta-human chorionic gonadotropin (beta-hCG) in normal and high risk pregnancy. *Clinical Chemistry and Enzymology Communications* 1997;**7**:341-7.

Elchalal 2000 {published data only}

Elchalal U, Ezra Y, Levi Y, Bar-Oz B, Yanai N, Intrator O, et al. Sonographically thick placenta: a marker for increased perinatal risk--a prospective cross-sectional study. *Placenta* 2000;**21**:268-72.

Ernst 2016 {published data only}

Ernst SA, Brand T, Zeeb H. Antenatal detection of intrauterine growth restriction: A case-control study. *European Journal of Epidemiology* 2016;**31**:S50-1.

Fadigas 2015 {published data only}

Fadigas C, Peeva G, Mendez O, Poon L C, Nicolaides KH. Prediction of small-for-gestational-age neonates: screening by placental growth factor and soluble fms-like tyrosine kinase-1 at 35-37 weeks. *Ultrasound in Obstetrics & Gynecology* 2015;**46**:191-7.

Falkner 1995 {published data only}

Falkner F. Ultrasonography and fetal growth: key perinatal factors. *Journal of Perinatology* 1995;**15**:114-8.

Ferrazzi 1986 {published data only}

Ferrazzi E, Nicolini U, Kustermann A, Pardi G. Routine obstetric ultrasound: effectiveness of cross-sectional screening for fetal growth retardation. *Journal of Clinical Ultrasound* 1986;**14**:17-22.

Fioretti 1986 {published data only}

Fioretti P, Melis GB, Paoletti AM. Ultrasound and human placental lactogen in the diagnosis of intrauterine growth retardation. *New Trends in Gynaecology and Obstetrics* 1986;**2**:59-64.

Fischer-Rasmussen 1971 {published data only}

Fischer-Rasmussen W. Plasma oestrogens and the fetal outcome. *Acta Obstetrica et Gynecologica Scandinavica* 1971; Vol. 50, issue 4:301-9.

Fisteag-Kiprono 2006 {published data only}

Fisteag-Kiprono L, Neiger R, Sonek JD, Croom CS, McKenna DS, Ventolini G. Perinatal outcome associated with sonographically detected globular placenta. *Journal of Reproductive Medicine* 2006;**51**:563-6.

Forger 2016 {published data only}

Forger F, Baumann M, Risch L, Raio L, Surbek D, Wiedemann U, et al. Angiogenic placental factors during pregnancy in rheumatoid arthritis. *Annals of the Rheumatic Diseases* 2016;**75**:474.

Fotiou 2015 {published data only}

Fotiou M, Michaelidou AM, Athanasiadis AP, Menexes G, Symeonidou M, Koulourida V, et al. Second trimester amniotic fluid glucose, uric acid, phosphate, potassium, and sodium concentrations in relation to maternal pre-pregnancy BMI and birth weight centiles. *Journal of Maternal-Fetal & Neonatal Medicine* 2015;**28**:910-5.

Furuhashi 1984 {published data only}

Furuhashi N, Tachibana Y, Shinkawa O, Hiruta M, Takahashi T, Tanaka M. Simultaneous and serial measurement of serum levels of human placental lactogen, beta-human chorionic gonadotropin and unconjugated estriol levels in pregnant women. *Tohoku Journal of Experimental Medicine* 1984;**144**:211-5.

Gabbay-Benziv 2016a {published data only}

Gabbay-Benziv R, Aviram A, Ashwal E, Hiersch L, Melamed N, Hadar E, et al. Sonographic prediction of small for gestational age-which formula is more accurate?. *American Journal of Obstetrics and Gynecology* 2016;**1**:S356.

Gabbay-Benziv 2017 {published data only}

Gabbay-Benziv R, Aviram A, Hadar E, Chen R, Bardin R, Wiznitzer A, et al. Pregnancy outcome after false diagnosis of fetal growth restriction. *Journal of Maternal-Fetal and Neonatal Medicine* 2017;**30**(16):1916-9.

Gaillard 2014 {published data only}

Gaillard R, Steegers EA, de Jongste JC, Hofman A, Jaddoe VW. Tracking of fetal growth characteristics during different trimesters and the risks of adverse birth outcomes. *International Journal of Epidemiology* 2014;**43**:1140-53.

Gao 2008 {published data only}

Gao T, Zablith NR, Burns DH, Skinner CD, Koski KG. Second trimester amniotic fluid transferrin and uric acid predict infant birth outcomes. *Prenatal Diagnosis* 2008;**28**:810-4.

Garcia-Flores 2015 {published data only}

Garcia-Flores J, Cruceyra M, Canameres M, Garicano A, Nieto O, Lopez A, et al. Fetal limb soft tissue assessment for prediction of birth weight and umbilical cord blood analytes in gestational diabetes. *Prenatal Diagnosis* 2015;**35**:1187-96.

Garoff 1976 {published data only}

Garoff L. Prediction of fetal outcome by urinary estriol, maternal serum placental lactogen, and alpha-fetoprotein in diabetes and hepatosis of pregnancy. *Obstetrics & Gynecology* 1976;**48**:659-66.

Gaziano 1988 {published data only}

Gaziano E, Knox GE, Wager GP, Bendel RP, Boyce DJ, Olson J. The predictability of the small-for-gestational-age infant by real-time ultrasound-derived measurements combined with pulsed Doppler umbilical artery velocimetry. *American Journal of Obstetrics and Gynecology* 1988; Vol. 158, issue 6:1431-9.

Geerts 2007 {published data only}

Geerts L, Odendaal HJ. Severe early onset pre-eclampsia: prognostic value of ultrasound and Doppler assessment. *Journal of Perinatology* 2007;**27**:335-42.

Gerhard 1987 {published data only}

Gerhard I, Vollmar B, Runnebaum B, Klinga K, Haller U, Kubli F. Weight percentile at birth. II. Prediction by endocrinological and sonographic measurements. *European Journal of Obstetrics, Gynecology, & Reproductive Biology* 1987;**26**:313-28.

Gernand 2015 {published data only}

Gernand AD, Schulze KJ, Nanayakkara-Bind A, Arguello M, Shamim AA, Ali H, et al. Effects of prenatal multiple micronutrient supplementation on fetal growth factors: a cluster-randomized, controlled trial in rural Bangladesh. *PLOS One* 2015;**10**(10):e0137269.

Gherzi 1981 {published data only}

Gherzi R, Pastore M, Labate D. [Monitoring of HPL, estriol and estetrol in high-risk pregnancy]. *Minerva Ginecologica* 1981;**33**:911-6.

Giambanco 1986 {published data only}

Giambanco V, Germana CF, Esposito S. Ultrasonographic assessment of fetal growth retardation. [Italian]. *Ultrasuoni in Ostetricia e Ginecologia* 1986;**4**:284-8.

Giardini 2014 {published data only}

Giardini V, Algeri P, Todyrenchuk L, Pellizzoni F, Callegari C, D'Arcangelo F, et al. A longitudinal study of serum placental growth factor levels in normal pregnancy and women at high risk for preeclampsia and/or intrauterine growth restriction. *Reproductive Sciences* 2014;**1**:191A-2A.

Gloning 1991 {published data only}

Gloning KP, Papaioannou A, Kuss E. The validity of CTG and estriol as predictors of intrauterine death. [German]. *Archives of Gynecology and Obstetrics* 1991;**250**:1184-5.

Goetzinger 2013 {published data only}

Goetzinger KR, Tuuli MG, Odibo AO, Roehl KA, Macones GA, Cahill AG. Screening for fetal growth disorders by clinical exam in the era of obesity. *Journal of Perinatology* 2013;**33**:352-7.

Goldenberg 1993 {published data only}

Goldenberg RL, Davis RO, Cliver CP, Cutter GR, Hoffman HJ, Dubard MB, et al. Maternal risk factors and their influence

on fetal anthropometric measurements. *American Journal of Obstetrics and Gynecology* 1993;**168**:1197-205.

Goldenberg 1997 {published data only}

Goldenberg RL, Cliver SP, Neggers Y, Copper RL, DuBard MD, Davis RO, et al. The relationship between maternal characteristics and fetal and neonatal anthropometric measurements in women delivering at term: a summary. *Acta Obstetrica et Gynecologica Scandinavica. Supplement* 1997;**165**:8-13.

Gomez-Roig 2015 {published data only}

Gomez-Roig MD, Mazarico E, Sabria J, Parra J, Oton L, Vela A. Use of placental growth factor and uterine artery doppler pulsatility index in pregnancies involving intrauterine fetal growth restriction or preeclampsia to predict perinatal outcomes. *Gynecologic and Obstetric Investigation* 2015;**80**:99-105.

Gordon 1978 {published data only}

Gordon YB, Lewis JD, Pendlebury DJ. Is measurement of placental function and maternal weight worth while?. *Lancet* 1978;**1**:1001-3.

Grantz 2016 {published data only}

Grantz K, Kim S, Grobman W, Newman R, Owen J, Skupski D, et al. Does the addition of fetal growth velocity improve the precision of sonographic birthweight estimation?. *American Journal of Obstetrics and Gynecology* 2016;**1**:S78.

Gravett 2015 {published data only}

Gravett M. Maternal serum biomarkers for assessment of preeclampsia: Glycosylated fibronectin as a point-of-care biomarker. *Journal of Perinatal Medicine. Conference: 12th World Congress of Perinatal Medicine* 2015;**43**(S1):73.

Griffin 2014 {published data only}

Griffin M, Duckworth S, Webster L, Seed P, Chappell L, Shennan A. Comparison of PlGF and other biomarkers against current ultrasound parameters for predicting delivery of small for gestational age (SGA) infants in women with suspected preeclampsia: the PELICAN study. *BJOG: an international journal of obstetrics and gynaecology* 2014;**121**(7):e6.

Gris 2015 {published data only}

Gris JC, Bouvier S, Lavigne G, Nouvellon E, Mercier E, Galanaud JP, et al. Obstetric antiphospholipid syndrome: Early Imwh induced variations of angiogenic factors predict pregnancy loss. *Journal of Thrombosis and Haemostasis* 2015;**13**:167.

Habib 2002 {published data only}

Habib FA. Prediction of low birth weight infants from ultrasound measurement of placental diameter and placental thickness. *Annals of Saudi Medicine* 2002;**22**:312-4.

Hargreaves 2011 {published data only}

Hargreaves K, Cameron M, Edwards H, Gray R, Deane K. Is the use of symphysis-fundal height measurement and ultrasound examination effective in detecting small or large fetuses?. *Journal of Obstetrics & Gynaecology* 2011;**31**:380-3.

Harper 2014 {published data only}

Harper LM, Jauk VC, Owen J, Biggio RR. The utility of ultrasound surveillance of fluid and growth in obese women. *American Journal of Obstetrics and Gynecology* 2014;**211**:524.e1-8.

Hassan 1987 {published data only}

Hassan MM, Bottoms SF, Mariona FG, Syner FN, Simkowski KM, Sokol RJ. The use of clinical, biochemical, and ultrasound parameters for the diagnosis of intrauterine growth retardation. *American Journal of Perinatology* 1987;**4**:191-4.

Hawkins 2014 {published data only}

Hawkins LK, Schnettler WT, Modest AM, Hacker MR, Rodriguez D. Association of third-trimester abdominal circumference with provider-initiated preterm delivery. *Journal of Maternal-Fetal & Neonatal Medicine* 2014;**27**:1228-31.

Heazell 2014 {published data only}

Heazell A. A kick in the right direction-reduced fetal movements and stillbirth prevention. *BMC Pregnancy and Childbirth* 2014;**15**(Supp 1):A7.

Henrichs 2016 {published data only}

Henrichs J, Verfaillie V, Viester L, Westerneng M, Molewijk B, Franx A, et al. Effectiveness and cost-effectiveness of routine third trimester ultrasound screening for intrauterine growth restriction: study protocol of a nationwide stepped wedge cluster-randomized trial in The Netherlands (The IRIS Study). *BMC Pregnancy & Childbirth* 2016;**16**(1):310.

Hensleigh 1977 {published data only}

Hensleigh PA, Cheatum SG, Spellacy N. Oxytocinase and human placental lactogen for the prediction of intrauterine growth retardation. *American Journal of Obstetrics and Gynecology* 1977; Vol. 129:675-8.

Herraiz 2014 {published data only}

Herraiz I, Droge LA, Gomez-Montes E, Henrich W, Galindo A, Verloren S. Characterization of the soluble fms-like tyrosine kinase-1 to placental growth factor ratio in pregnancies complicated by fetal growth restriction. *Obstetrics & Gynecology* 2014;**124**:265-73.

Hinkle 2015 {published data only}

Hinkle SN, Johns AM, Albert PS, Kim S, Grantz KL. Longitudinal changes in gestational weight gain and the association with intrauterine fetal growth. *European Journal of Obstetrics Gynecology and Reproductive Biology* 2015;**190**:41-7.

Hughes 1980 {published data only}

Hughes G, Bischof P, Wilson G, Smith R, Kloppe A. Tests of fetal wellbeing in the third trimester of pregnancy. *British Journal of Obstetrics and Gynaecology* 1980;**87**:650-6.

Husse 2014 {published data only}

Husse S, Gottschlich A, Schrey S, Stepan H, Hoffmann J. [Predictive value of the sFlt1/PlGF ratio for the diagnosis of preeclampsia in high-risk patients]. *Zeitschrift für Geburtshilfe und Neonatologie* 2014;**218**:34-41.

Jabeen 1999 {published data only}

Jabeen S, Sabrina, Ahmad S. Screening for intra-uterine growth retardation. *Journal of the College of Physicians and Surgeons Pakistan* 1999;**9**:17-9.

James-Todd 2015 {published data only}

James-Todd T, Cohen A, Wenger J, Brown FM. Placental-like growth factor (PIGF) and infant birth weight in women with preexisting diabetes. *Diabetes* 2015;**64**:A378.

Johnson 2011 {published data only}

Johnson RJ, Kanbay M, Kang DH, Sanchez-Lozada LG, Feig D. Uric acid: A clinically useful marker to distinguish preeclampsia from gestational hypertension. *Hypertension* 2011;**58**:548-9.

Johnstone 2015 {published data only}

Johnstone ED, Higgins L, Myers J, Simcos L, Heazell A. Exclusion of early onset fetal growth restriction using a combination placental and uterine artery doppler screening. *Reproductive Sciences* 2015;**22**:383A-4A.

Karjalainen 1975 {published data only}

Karjalainen O, Stenman U, Widholm O. Urinary oestriol and serum activities of human placental lactogen and heat stable alkaline phosphatase as indicators of fetoplacental function. *Annales Chirurgiae et Gynaecologiae Fenniae* 1975;**64**:50-4.

Karlsen 2016 {published data only}

Karlsen HO, Johnsen SL, Rasmussen S, Kiserud T. Prediction of adverse perinatal outcome of small-for-gestational-age pregnancy using size centiles and conditional growth centiles. *Ultrasound in Obstetrics & Gynecology* 2016;**48**:217-23.

Kase 2012 {published data only}

Kase B, Blackwell SC. Ultrasound derived estimated fetal weight using customized standards: Does it improve prediction of adverse pregnancy outcomes?. *American Journal of Obstetrics and Gynecology* 2012;**1**:S158.

Kase 2012a {published data only}

Kase BA, Carreno CA, Blackwell SC. Customized estimated fetal weight: a novel antenatal tool to diagnose abnormal fetal growth. *American Journal of Obstetrics and Gynecology* 2012;**207**:218.e1-5.

Kazzi 1983 {published data only}

Kazzi GM, Gross TL, Sokol RJ. Fetal biparietal diameter an placental grade: predictors of intrauterine growth retardation. *Obstetrics & Gynecology* 1983;**62**:755-9.

Khalil 2015 {published data only}

Khalil AA, Morales-Rosello J, Elsaddig M, Khan N, Papageorgiou A, Bhide A, et al. The association between fetal Doppler and admission to neonatal unit at term. *American Journal of Obstetrics and Gynecology* 2015;**213**:57.e1-7.

Khalil 2016 {published data only}

Khalil A, Morales-Rosello J, Townsend R, Morlando M, Papageorgiou A, Bhide A, et al. Value of third-trimester cerebroplacental ratio and uterine artery Doppler indices as predictors of stillbirth and perinatal loss.[Erratum appears

in *Ultrasound Obstet Gynecol.* 2016 Apr;**47**(4):526; PMID: 27062979]. *Ultrasound in Obstetrics & Gynecology* 2016;**47**:74-80.

Kihaile 1988 {published data only}

Kihaile PE. Ultrasonic grey-level histograms of prenatal placenta and its relation to fetal well-being. *Yonago Acta Medica* 1988;**31**:139-46.

Kim 2009 {published data only}

Kim YN, Lee DS, Jeong DH, Sung MS, Kim KT. The relationship of the level of circulating antiangiogenic factors to the clinical manifestations of preeclampsia. *Prenatal Diagnosis* 2009;**29**(5):464-70.

Kim 2014 {published data only}

Kim JW, Kim YH, Song TB. The usefulness of gestation corrected hyperuricemia as predictors of the recurrence of preeclampsia and obstetric outcomes on subsequent pregnancy. *Journal of Maternal-Fetal and Neonatal Medicine* 2014;**27**:355-6.

Kim 2016 {published data only}

Kim MY, Buyon JP, Guerra MM, Rana S, Zhang D, Laskin CA, et al. Angiogenic factor imbalance early in pregnancy predicts adverse outcomes in patients with lupus and antiphospholipid antibodies: results of the PROMISSE study. *American Journal of Obstetrics and Gynecology* 2016;**214**:108.e1-108.e14.

Kjos 2015 {published data only}

Kjos K, Ghashghaei R, Klisser K, Woelkers D. Prediction of adverse outcomes with mid-pregnancy serum placental growth factor levels and uterine artery Dopplers in high risk pregnancies. *Pregnancy Hypertension* 2015;**5**(1):80.

Kneitel 2016 {published data only}

Kneitel AW, Treadwell MC, O'Brien LM. Effects of maternal sleep apnea on fetal growth. *American Journal of Obstetrics and Gynecology* 2016;**1**:S250.

Kolovetsiou-Kreiner 2014 {published data only}

Kolovetsiou-Kreiner V, Stern EC, Mayer-Pickel K, Lakovschek I, Ulrich D, Lang U, et al. Predictive value of gestational angiogenic biomarkers SFLT-1 and PIGF in high-risk women. *Reproductive Sciences* 2014;**1**:300A.

Kolovetsiou-Kreiner 2015 {published data only}

Kolovetsiou-Kreiner V, Stern EC, Mayer-Pickel K, Lakovschek I, Lang U, Cervar-Zivkovic M. Repeated controls of gestational angiogenic biomarkers in high-risk pregnancies-do they derive a clinical benefit?. *Journal of Perinatal Medicine. Conference: 12th World Congress of Perinatal Medicine* 2015;**43**:300A.

Krochik 2010 {published data only}

Krochik AG, Chaler EA, Maceiras M, Aspres N, Mazza CS. Presence of early risk markers of metabolic syndrome in prepubertal children with a history of intrauterine growth restriction. [Spanish]. *Archivos Argentinos de Pediatría* 2010;**108**:10-6.

Kulkarni 1981 {published data only}

Kulkarni BD, O'Leary JA, Takagi RL, Avila TD, Jabamoni R. Plasma estrogens in the assessment of fetoplacental function. *Clinical Biochemistry* 1981;**14**:108-12.

Kulkarni 2010 {published data only}

Kulkarni AV, Mehendale SS, Yadav HR, Kilari AS, Taralekar VS, Joshi SR. Circulating angiogenic factors and their association with birth outcomes in preeclampsia. *Hypertension Research - Clinical & Experimental* 2010;**33**:561-7.

Kullander 1982 {published data only}

Kullander S, Marsal K, Persson PH. Human placental lactogen and ultrasonic screening for the detection of placental insufficiency. *Contributions to Gynecology & Obstetrics* 1982;**9**:129-44.

Kundu 1978 {published data only}

Kundu N, Carmody PJ, Didolkar SM, Petersen LP. Sequential determination of serum human placental lactogen, estriol, and estetrol for assessment of fetal morbidity. *Obstetrics & Gynecology* 1978;**52**:513-20.

Kunzig 1975 {published data only}

Kunzig HJ, Geiger W, Schlensker KH. [Serum estriol and HPL concentration as well as echographic maturity diagnosis as parameters of placental function]. *Archiv fur Gynakologie* 1975;**219**:443-5.

Kunzig 1980 {published data only}

Kunzig HJ. [Diagnosis and monitoring of risk pregnancies. Serum estrone, estradiol-17 beta and estradiol determination in normal and risk pregnancies. Methods and results of radioimmunologic studies]. *Fortschritte der Medizin* 1980;**98**:626-31.

Lai 2014 {published data only}

Lai J, Garcia-Tizon Larroca S, Peeva G, Poon LC, Wright D, Nicolaides KH. Competing risks model in screening for preeclampsia by serum placental growth factor and soluble fms-like tyrosine kinase-1 at 30-33 weeks' gestation. *Fetal Diagnosis and Therapy* 2014;**35**(4):240-8.

Larkin 2015 {published data only}

Larkin J, Chauhan S, Simhan H. Small for gestational age: The differential mortality when detected versus undetected antenatally. *American Journal of Obstetrics and Gynecology* 2015;**1**:S44.

Larsen 1992 {published data only}

Larsen T, Larsen JF, Petersen S, Greisen G. Detection of small-for-gestational-age fetuses by ultrasound screening in a high risk population: a randomized controlled study. *British Journal of Obstetrics and Gynaecology* 1992;**99**:469-74.

Larsen 1997 {published data only}

Larsen T, Greisen G, Petersen S. Intrauterine growth correlation to postnatal growth--influence of risk factors and complications in pregnancy. *Early Human Development* 1997;**47**:157-65.

Laurin 1987a {published data only}

Laurin J, Persson PH, Fernlund P. The efficacy of biochemical assays in assessing an IUGR risk-group, preselected with ultrasound fetometry. *European Journal of Obstetrics, Gynecology, & Reproductive Biology* 1987;**24**:177-87.

Lean 2016 {published data only}

Lean S, Heazell A, Peacock L, Lodge K, Roberts S, Jones R. Manchester Advanced Maternal Age Study (MAMAS): A prospective observational cohort study to identify risk factors and markers of adverse pregnancy outcome. *BJOG: an international journal of obstetrics and gynaecology* 2016;**123**:41.

Leanos-Miranda 2013 {published data only}

Leanos-Miranda A, Campos-Galicia I, Ramirez-Valenzuela KL, Chinolla-Arellano ZL, Isordia-Salas I. Circulating angiogenic factors and urinary prolactin as predictors of adverse outcomes in women with preeclampsia. *Hypertension* 2013;**61**:1118-25.

Lechner 1987 {published data only}

Lechner W, Heim K, Zech J, Daxenbichler G, Marth C. The relation between saliva estriol levels in pregnancy and infant birth weight. *Archives of Gynecology and Obstetrics* 1987;**241**:9-12.

Levine 2005 {published data only}

Levine RJ, Thadhani R, Qian C, Lam C, Lim KH, Yu KF, et al. Urinary placental growth factor and risk of preeclampsia. *JAMA* 2005;**293**:77-85.

Li 2014 {published data only}

Li SW, Ling Y, Jin S, Lin YF, Chen ZJ, Hu CX, et al. Expression of soluble vascular endothelial growth factor receptor-1 and placental growth factor in fetal growth restriction cases and intervention effect of tetramethylpyrazine. *Asian Pacific Journal of Tropical Medicine* 2014;**7**:663-7.

Little 2016 {published data only}

Little J, Chew C, Kither H, Kiss O, Allibone A, Bruce IN, et al. Circulating placental growth factor, measured at 28 weeks of gestation, may hold limited prognostic value in predicting poor pregnancy outcome in patients with systemic lupus erythematosus. *BJOG: an international journal of obstetrics and gynaecology* 2016;**123**:51-2.

Lobmaier 2014 {published data only}

Lobmaier SM, Figueras F, Mercade I, Perello M, Peguero A, Crovetto F, et al. Angiogenic factors vs Doppler surveillance in the prediction of adverse outcome among late-pregnancy small-for-gestational-age fetuses. *Ultrasound in Obstetrics & Gynecology* 2014;**43**:533-40.

London 1983 {published data only}

London R, Vallejos J, Baer D. Serum unconjugated estriol concentrations before and after the oxytocin challenge test: An index of fetal well-being. *American Journal of Obstetrics and Gynecology* 1983;**146**:630-2.

MacDonald 1983 {published data only}

MacDonald DJ, Scott JM, Gemmell RS, Mack DS. A prospective study of three biochemical fetoplacental tests: serum human

placental lactogen, pregnancy-specific beta 1-glycoprotein, and urinary estrogens, and their relationship to placental insufficiency. *American Journal of Obstetrics and Gynecology* 1983;**147**:430-6.

Macmillian 1976 {published data only}

Macmillian DR, Hawkins R, Collier RN. Chorionic somatomammotrophin as index of fetal growth. *Archives of Disease in Childhood* 1976;**51**(2):120-3.

Maly 1987 {published data only}

Maly Z, Cupr Z, Kadrnkova M. [Evaluation of fetal hypotrophy by examining placental structure using ultrasonography and urinary estriol levels]. *Ceskoslovenska Gynecologie* 1987;**52**:179-82.

March 2015 {published data only}

March M I, Geahchan C, Wenger J, Raghuraman N, Berg A, Haddow H, et al. Circulating angiogenic factors and the risk of adverse outcomes among Haitian women with preeclampsia. *PLOS One* 2015;**10**(5):e0126815.

Margossian 2016 {published data only}

Margossian A, Boisson-Gaudin C, Subtil F, Rudigoz RC, Dubernard G, Allias F, et al. [Intra-uterine growth restriction impact on maternal serum concentration of PLGF (placental growth factor): A case control study]. *Gynecologie, Obstetrique & Fertilité* 2016;**44**:23-8.

Markestad 1997 {published data only}

Markestad T. Prediction of fetal growth based on maternal serum concentrations of human chorionic gonadotropin, human placental lactogen and estriol. *Acta Obstetrica et Gynecologica Scandinavica, Supplement* 1997;**76**:50-5.

Martins 2005 {published data only}

Martins MM, Tedesco JJ. Early diagnosis of intra-uterine growth restriction by ultrasonographic estimation of fetal weight. *Revista Da Associacao Medica Brasileira* 2005; Vol. 51, issue 1:41-5.

Masoura 2014 {published data only}

Masoura S, Kalogiannidis I, Makedou K, Theodoridis T, Koiou K, Gerou S, et al. Biomarkers of endothelial dysfunction in preeclampsia and neonatal morbidity: a case-control study. [Erratum appears in *Eur J Obstet Gynecol Reprod Biol.* 2014 Sep;180:209]. *European Journal of Obstetrics, Gynecology, & Reproductive Biology* 2014;**175**:119-23.

Matthews 2017 {published data only}

Matthews KC, Williamson J, Gupta S, Lam-Rachlin J, Saltzman DH, Rebarber A, et al. The effect of a sonographic estimated fetal weight on the risk of cesarean delivery in macrosomic and small for gestational-age infants. *Journal of Maternal-Fetal and Neonatal Medicine* 2017;**30**(10):1172-6.

Mazzocco 2014 {published data only}

Mazzocco MI, Calabrese S, Cardellicchio M, Martinelli A, Cetin I. Could PLGF be a useful marker to predict fetal outcomes in pregnancies at risk?. *Reproductive Sciences* 2014;**1**:304A-5A.

McKenna 2003 {published data only}

McKenna D, Tharmaratnam S, Mahsud S, Bailie C, Harper A, Dornan J. A randomized trial using ultrasound to identify the high-risk fetus in a low-risk population. *Obstetrics & Gynecology* 2003;**101**:626-32.

Melamed 2015 {published data only}

Melamed N, Kingdom J. Sonographic weight estimation of small for gestational age fetuses: Is the optimal model related to fetal body proportions?. *American Journal of Obstetrics and Gynecology* 2015;**1**:S263-4.

Melamed 2016 {published data only}

Melamed N, Pittini A, Kingdom J, Barrett J. Sonographic factors distinguishing late intrauterine growth restriction from late small for gestational age fetuses. *American Journal of Obstetrics and Gynecology* 2016;**1**:S104-5.

Melamed 2016a {published data only}

Melamed N, Ryan G, Windrim R, Toi A, Kingdom J. Choice of formula and accuracy of fetal weight estimation in small-for-gestational-age fetuses. *Journal of Ultrasound in Medicine* 2016;**35**:71-82.

Merriam 2014 {published data only}

Merriam A, Jain V, Shlossman P, Hoffman M. Does abdominal circumference less than the 5 percentile improve detection of fetal growth restriction?. *American Journal of Obstetrics and Gynecology* 2014;**1**:S102.

Mertens 1975 {published data only}

Mertens H, Puder H. [Diagnosis of placental insufficiency through HPL studies]. *Archiv fur Gynakologie* 1975; Vol. 219:428-30.

Mirza 2015 {published data only}

Mirza FG, Bauer ST, Van Der Veer A, Simpson LL. Gastroschisis: Incidence and prediction of growth restriction. *Journal of Perinatal Medicine* 2015;**43**:605-8.

Miwa 2014 {published data only}

Miwa I, Sase M, Torii M, Sanai H, Nakamura Y, Ueda K. A thick placenta: a predictor of adverse pregnancy outcomes. *Springerplus* 2014;**3**:353.

Mlynarczyk 2015 {published data only}

Mlynarczyk M, Chauhan S, Baydoun H, Wilkes C, Earhart K, Goodier C, et al. Fetal growth restriction < 5% versus 5-9%: Multi-center study for comparison of neonatal morbidity (ULTRA TOT). *American Journal of Obstetrics and Gynecology* 2015;**1**:S109.

Mlynarczyk 2015a {published data only}

Mlynarczyk M, Chauhan SP, Wilkes CM, Earhart KR, Lee NM, Owens M, et al. Accuracy of sonographic estimation of fetal weight (SEFW) as a predictor of birth weight (BW) in growth-restricted fetuses; A large multicenter study. *Ultrasound in Medicine and Biology* 2015;**1**:S131.

Mone 2016 {published data only}

Mone F, Hartigan L, Ali F, Neville G, Mahony R, Carroll S, et al. How accurately do we predict birthweight in the small-for-gestational-age fetus?. *BJOG: an international journal of obstetrics and gynaecology* 2016;**123**:76.

Moore 2012 {published data only}

Moore A, Young H, Keller J, Ojo L, Yan J, Simas TM, et al. Angiogenic biomarkers for the prediction of pregnancy complications in women with suspected preeclampsia. *American Journal of Obstetrics and Gynecology* 2012;**1**:S326-7.

Morrison 1980 {published data only}

Morrison I, Green P, Oomen B. The role of human placental lactogen assays in antepartum fetal assessment. *American Journal of Obstetrics & Gynecology* 1980; Vol. 136, issue 8:1055-60.

Muraguchi 1981 {published data only}

Muraguchi K, Takahashi K, Suzuki M, Ikeno N. [The concentration of serum unconjugated estradiol, estriol and estetrol in pregnant women, and the significance of these hormones in pregnancy (author's transl)]. *Nippon Naibunpi Gakkai Zasshi - Folia Endocrinologica Japonica* 1981;**57**:974-82.

Myatt 2013 {published data only}

Myatt L. Association of maternal biomarkers with obstetric outcomes. *Reproductive Sciences* 2013; Vol. 1:330A.

Nadal 2015 {published data only}

Nadal A, Parra-Saavedra M, Crovetto F, Triunfo S, Savchev S, Peguero A, et al. A comprehensive analysis of placental insufficiency in late-onset small for gestational age births. *Laboratory Investigation* 2015;**95**:299A.

Nair 2016 {published data only}

Nair A, Savitha C. Estimation of serum uric acid as an indicator of severity of preeclampsia and perinatal outcome. *Journal of Obstetrics and Gynecology of India* 2016;**67**(2):1-10.

Nelson 2015 {published data only}

Nelson DB, Martin R, Twickler DM, Santiago-Munoz PC, McIntire DD, Dashe JS. Sonographic detection and clinical importance of growth restriction in pregnancies with gastroschisis sonographic detection and clinical importance of growth restriction in pregnancies with gastroschisis. *Journal of Ultrasound in Medicine* 2015;**34**(12):2217-23.

Nice 2014 {published data only}

Nice D, Hayden K, Higgins L, Johnstone E, Heazell A. Human placental lactogen and placental growth factor can differentiate small for gestational age from appropriately grown infants. *Placenta* 2014;**35**(9):A19.

Nice 2014a {published data only}

Nice DB, Hayden K, Higgins L, Johnstone E, Heazell AE. The measurement of placental biomarkers in the detection of compromised pregnancies. *Clinical Chemistry and Laboratory Medicine* 2014;**52**(11):eA372.

Nieder 1976 {published data only}

Nieder J, Kapitza W. [The determination of human placental lactogen (HPL) for hormonal supervision in late pregnancy]. *Zentralblatt fur Gynakologie* 1976;**98**:1129-36.

Nielsen 1981 {published data only}

Nielsen PV, Schioler V. Ratio of human placental lactogenic hormone (hPL) in amniotic fluid/maternal serum. *Acta Obstetrica et Gynecologica Scandinavica* 1981; Vol. 60, issue 1:9-12.

Niknafs 2001 {published data only}

Niknafs P, Sibbald J. Accuracy of single ultrasound parameters in detection of fetal growth restriction. *American Journal of Perinatology* 2001; Vol. 18, issue 6:325-34.

O'Connor 2015 {published data only}

O'Connor H, Unterscheider J, Daly S, Geary M, Kennelly M, McAuliffe F, et al. Comparison of asymmetric versus symmetric IUGR - results from a national prospective trial. *American Journal of Obstetrics and Gynecology* 2015;**212**(1):S173-S174.

Obiekwe 1982 {published data only}

Obiekwe BC, Chard T. Human chorionic gonadotropin levels in maternal blood in late pregnancy: relation to birthweight, sex and condition of the infant at birth. *British Journal of Obstetrics & Gynaecology* 1982;**89**:543-6.

Odibo 2014 {published data only}

Odibo AO. Routine ultrasound examination at 41 weeks of gestation does not improve perinatal outcomes. *BJOG: an international journal of obstetrics and gynaecology* 2014;**121**:1116.

Okonofua 1986 {published data only}

Okonofua FE, Ayangade SO, Chan RC, O'Brien PM. A prospective comparison of clinical and ultrasonic methods of predicting normal and abnormal fetal growth. *International Journal of Gynaecology & Obstetrics* 1986;**24**:447-51.

Pal 2015 {published data only}

Pal A, Rajoria L. Hyperuricemia and preeclampsia: evaluation of uric acid as predictor of adverse perinatal outcome. *International Journal of Gynecology and Obstetrics* 2015;**131**:E244.

Palomaki 2015 {published data only}

Palomaki GE, Haddow JE, Haddow H, Salahuddin S, Geahchan C, Cerdeira AS, et al. Modeling risk for adverse outcomes in women with suspected preterm preeclampsia using angiogenic factor measurements. *Pregnancy Hypertension* 2015;**5**(1):12.

Palomaki 2015a {published data only}

Palomaki GE, Haddow JE, Haddow HR, Salahuddin S, Geahchan C, Cerdeira AS, et al. Modeling risk for severe adverse outcomes using angiogenic factor measurements in women with suspected preterm preeclampsia. *Prenatal Diagnosis* 2015;**35**:386-93.

Papastefanou 2014 {published data only}

Papastefanou I, Pilalis A, Chrelias C, Kassanos D, Souka AP. Screening for birth weight deviations by second and third trimester ultrasound scan. *Prenatal Diagnosis* 2014;**34**:759-64.

Papastefanou 2015 {published data only}

Papastefanou I, Souka AP, Eleftheriades M, Pilalis A, Chrelias C, Kassanos D. Predicting fetal growth deviation in parous women: combining the birth weight of the previous pregnancy and third trimester ultrasound scan. *Journal of Perinatal Medicine* 2015; Vol. 43, issue 4:485-92.

Parra Saavedra 2015 {published data only}

Parra Saavedra MA, Triunfo S, Crovetto F, Gratacos E, Figueras F. OP14.02: Ultrasound and Doppler evaluation at routine third trimester scan to detect late pregnancy (SGA) in adequate-for-gestational age (AGA) pregnancies. *Journal of Perinatal Medicine* 2015;**46**(S1):93-4.

Parrish 2010 {published data only}

Parrish M, Griffin M, Morris R, Darby M, Owens MY, Martin JN Jr. Hyperuricemia facilitates the prediction of maternal and perinatal adverse outcome in patients with severe/superimposed preeclampsia. *Journal of Maternal-Fetal & Neonatal Medicine* 2010;**23**:1451-5.

Partap 2015 {published data only}

Partap U, Sovio U, Smith G. Fetal growth and the risk of spontaneous preterm birth. *American Journal of Obstetrics and Gynecology* 2015;**1**:S208.

Pavelka 1982 {published data only}

Pavelka R, Schmid R, Reinold E. Evaluation of various monitoring techniques in late pregnancy to detect poor intrauterine fetal growth. *Gynecologic & Obstetric Investigation* 1982; Vol. 13:65-75.

Pecks 2015 {published data only}

Pecks U, Kleine-Eggebrecht N, Winkler BS, Mohaupt M, Escher G, Rath W. Maternal lipid-and steroid hormone concentrations during the course of pregnancy and in pregnancy pathologies. *Pregnancy Hypertension* 2015;**5**(1):99-100.

Peixoto 2016 {published data only}

Peixoto AB, Rodrigues da Cunha Caldas TM, Godoy Silva TA, Silva Gomes Caetano MS, Martins WP, Martins Santana EF, et al. Assessment of ultrasound and Doppler parameters in the third trimester of pregnancy as predictors of adverse perinatal outcome in unselected pregnancies. *Ginekologia Polska* 2016;**87**(7):510-5.

Perez-Cruz 2015 {published data only}

Perez-Cruz M, Cruz-Lemini M, Fernandez MT, Parra JA, Bartrons J, Gomez-Roig MD, et al. Fetal cardiac function in late-onset intrauterine growth restriction vs small-for-gestational age, as defined by estimated fetal weight, cerebroplacental ratio and uterine artery Doppler. *Ultrasound in Obstetrics & Gynecology* 2015;**46**:465-71.

Perry 1986 {published data only}

Perry L, Hickson R, Obiekwe BC, Chard T. Maternal oestriol levels reflect placental function rather than foetal function. *Acta Endocrinologica* 1986;**111**:563-6.

Persson 1978 {published data only}

Persson PH, Marsal K. Monitoring of fetuses with retarded BPD growth. *Acta Obstetrica et Gynecologica Scandinavica. Supplement* 1978; Vol. 78:49-55.

Persson 1980 {published data only}

Persson PH, Grennert L, Gennser G, Eneroth P. Fetal biparietal diameter and maternal plasma concentrations of placental lactogen, chorionic gonadotrophin, oestriol, and alpha-fetoprotein in normal and pathological pregnancies. *British Journal of Obstetrics and Gynaecology* 1980;**87**:25-32.

Peyronnet 2016 {published data only}

Peyronnet V, Kayem G, Mandelbrot L, Sibiude J. [Detection of small for gestational age fetuses during third trimester ultrasound. A monocentric observational study]. *Gynecologie, Obstetrique & Fertilité* 2016;**44**:531-6.

Pfeiffer 1990 {published data only}

Pfeiffer KH. [Detection of fetal growth retardation by instrumental and biochemical monitoring methods]. *Zeitschrift für Geburtshilfe und Perinatologie* 1990;**194**:99-103.

Pinheiro 2014 {published data only}

Pinheiro CC, Rayol P, Gozzani L, Reis LM, Zampieri G, Dias CB, et al. The relationship of angiogenic factors to maternal and neonatal manifestations of early-onset and late-onset preeclampsia. *Prenatal Diagnosis* 2014;**34**:1084-92.

Pledger 1984 {published data only}

Pledger DR, Belfield A, Calder AA, Wallace AM. The predictive value of three pregnancy-associated proteins in the detection of the light-for-dates baby. *British Journal of Obstetrics and Gynaecology* 1984; Vol. 91, issue 9:870-4.

Pluta 1979 {published data only}

Pluta M, Hardt W, Schmidt-Gollwitzer K, Schmidt-Gollwitzer M. Radioimmunoassay of serum SP 1 and HPL in normal and abnormal pregnancies. *Archives of Gynecology* 1979;**227**:327-36.

Ponce 1995 {published data only}

Ponce J, Boguna JM, Salvador C, Borrás M, Laila JM. Value of hyperuricemia as biological marker in hypertensive disorders of pregnancy. [Spanish]. *Progresos en Obstetricia y Ginecologia* 1995;**38**:244-50.

Powers 2010 {published data only}

Powers R, Roberts J, Plymire D, Pucci D, Datwyler S, Laird D, et al. Low maternal PlGF across pregnancy identifies a subset of women with preterm preeclampsia; Type 1 vs. type 2 preeclampsia?. *Pregnancy Hypertension* 2010;**1**:S14.

Prakash 2012 {published data only}

Prakash S, Sharma N, Kumari P, Kumar A. Serum uric acid as marker for diagnosing preeclampsia. *International Journal of Pharmaceutical Sciences and Research* 2012;**3**:2669-75.

Qublan 2005 {published data only}

Qublan HS, Ammarin V, Bataineh O, Al-Shraideh Z, Tahat Y, Awamleh I, et al. Lactic dehydrogenase as a biochemical marker of adverse pregnancy outcome in severe pre-eclampsia. *Medical Science Monitor* 2005;**11**:CR393-7.

Raghuramulu 1978 {published data only}

Raghuramulu N. Plasma placental lactogen in pregnancy. *Nutrition & Metabolism* 1978;**22**:160-6.

Rajasingam 2009 {published data only}

Rajasingam D, Seed PT, Briley AL, Shennan AH, Poston L. A prospective study of pregnancy outcome and biomarkers of oxidative stress in nulliparous obese women. *American Journal of Obstetrics & Gynecology* 2009;**200**:395.e1-9.

Rasanen 2015 {published data only}

Rasanen J, Quinn MJ, Laurie A, Bean E, Roberts CT, Nagalla SR, et al. Maternal serum glycosylated fibronectin as a point-of-care biomarker for assessment of preeclampsia. *American Journal of Obstetrics and Gynecology* 2015;**212**:82.e1-82e9.

Reck 1987 {published data only}

Reck G, Kronitz B, Breckwoldt M. [Significance of the estriol profile as an endogenous function test of the fetoplacental unit]. *Geburtshilfe und Frauenheilkunde* 1987;**47**:774-80.

Riss 1982 {published data only}

Riss P, Bartl W. Placental function, fetal distress, and the fetal/placental weight ratio in normal and gestotic pregnancies. *International Journal of Biological Research in Pregnancy* 1982;**3**:10-3.

Ris-Stalpers 2012 {published data only}

Ris-Stalpers C, Hassani Lahsinoui H, Boussata S, Am Van Der Post J. Placental growth factor as a diagnostic and prognostic test for placental complications of pregnancy. *Pregnancy Hypertension* 2012;**2**(3):212.

Rizos 2013 {published data only}

Rizos D, Eleftheriades M, Karampas G, Rizou M, Haliassos A, Hassiakos D, et al. Placental growth factor and soluble fms-like tyrosine kinase-1 are useful markers for the prediction of preeclampsia but not for small for gestational age neonates: a longitudinal study. *European Journal of Obstetrics, Gynecology, & Reproductive Biology* 2013;**171**:225-30.

Rocca 1995 {published data only}

Rocca MM, Said MS, Khamis MY, Ghanem IA, Karkour TA. The value of Doppler study of the umbilical artery in predicting perinatal outcome in pre-eclamptic patients. *Journal of Obstetrics and Gynaecology* 1995;**21**:427-31.

Romero 2008 {published data only}

Romero R, Nien JK, Espinoza J, Todem D, Fu W, Chung H, et al. A longitudinal study of angiogenic (placental growth factor) and anti-angiogenic (soluble endoglin and soluble vascular endothelial growth factor receptor-1) factors in normal pregnancy and patients destined to develop preeclampsia and deliver a small for gestational age neonate. *Journal of Maternal-Fetal & Neonatal Medicine* 2008;**21**:9-23.

Ronin-Walknowska 1984 {published data only}

Ronin-Walknowska E, Holmgren PA, von Schoultz B, Stigbrand T. Placental alkaline phosphatase compared with human placental lactogen and oestriol in high-risk pregnancies. *Gynecologic & Obstetric Investigation* 1984;**18**:206-11.

Rosendahl 1988 {published data only}

Rosendahl H, Kivinen S. Routine ultrasound screening for early detection of small for gestational age fetuses. *Obstetrics and Gynecology* 1988;**71**:518-21.

Rosendahl 1991 {published data only}

Rosendahl H, Kivinen S. Detection of small for gestational age fetuses by the combination of clinical risk factors and ultrasonography. *European Journal of Obstetrics, Gynecology, and Reproductive Biology* 1991;**39**:7-11.

Rothenbacher 2016 {published data only}

Rothenbacher D, Braig S, Muller M, Koenig W, Reister F, Genuneit J. Maternal uric acid serum concentrations, renal function at delivery and health-related pregnancy outcomes in neonates: The Ulm SPATZ Health Study. *European Journal of Epidemiology* 2016;**31**:S166.

Ruozi Berretta 1967 {published data only}

Ruozi Berretta L, Avitto P. [Estriol and pregnanediol as indices of placental function in normal and pathological pregnancy]. *Minerva Ginecologica* 1967;**19**:138-41.

Sabbagha 1979 {published data only}

Sabbagha RE, Tamura RK. Antenatal ultrasound diagnosis of intrauterine growth retardation. *Perinatology Neonatology* 1979;**3**:33-6.

Salahuddin 2016 {published data only}

Salahuddin S, Wenger JB, Zhang D, Thadhani R, Karumanchi SA, Rana S. KRYPTOR-automated angiogenic factor assays and risk of preeclampsia-related adverse outcomes. *Hypertension in Pregnancy* 2016;**35**:330-45.

Salas 1993 {published data only}

Salas SP, Rosso P, Espinoza R, Robert JA, Valdes G, Donoso E. Maternal plasma volume expansion and hormonal changes in women with idiopathic fetal growth retardation. *Obstetrics and Gynecology* 1993;**81**:1029-33.

Salas 1998 {published data only}

Salas SP, Rosso P. Plasma volume, renal function, and hormonal levels in pregnant women with idiopathic fetal growth restriction or preeclampsia. *Hypertension in Pregnancy* 1998;**17**:69-79.

Salas 2006 {published data only}

Salas SP, Marshall G, Gutierrez BL, Rosso P. Time course of maternal plasma volume and hormonal changes in women with preeclampsia or fetal growth restriction. *Hypertension* 2006;**47**:203-8.

Saleh 2015 {published data only}

Saleh L, Verdonk K, Danser AH, Steegers EA, Russcher H, Van Den Meiracker AH, et al. The preratio study: Is the SFLT-1/PLGF

- ratio a suitable marker to diagnose preeclampsia and to predict adverse maternal/neonatal pregnancy outcome?. *Journal of Hypertension* 2015;**33**:e347-8.
- Salkie 1977** {published data only}
Salkie ML, Hannah CL. Maternal serum hyaluronidase activity in pregnancy. *Enzyme* 1977;**22**:52-9.
- Samanta 1989** {published data only}
Samanta B, Dutta GP. Study on blood level of human placental lactogen in abnormal pregnancy. *Journal of the Indian Medical Association* 1989;**87**:205-7.
- Sanchez Fernandez 2015** {published data only}
Sanchez Fernandez M, De La Fuente Pedrosa ER, Garcia Cotes AE, Sanchez Barroso MT, Mozas Moreno J, Jimenez-Moleon JJ. Ultrasound scan error associated factors on fetal weight measurement. *Journal of Perinatal Medicine* 2015;**43**:908.
- Sarandakou 1989** {published data only}
Sarandakou A, Rizos D, Phocas I. Comparative study of four hormonal parameters in pregnancies with low-birthweight infants. *Clinical Chemistry and Enzymology Communications* 1989;**1**:159-69.
- Sato 1974** {published data only}
Sato Y, Arai S, Takeuchi S, Hiroi M. [Feto-placental function tests based on the ratio of alpha-fetoprotein and HPL of the maternal blood]. *Hormon to Rinsho - Clinical Endocrinology* 1974;**22**:1061-5.
- Secher 1986** {published data only}
Secher NJ, Hansen PK, Lenstrup C, Eriksen PS. Controlled trial of ultrasound screening for light for gestational age (LGA) infants in late pregnancy. *European Journal of Obstetrics, Gynecology, and Reproductive Biology* 1986;**23**:307-13.
- Secher 1987** {published data only}
Secher NJ, Kern Hansen P, Lenstrup C, Sindberg Eriksen P, Morsing G. A randomized study of fetal abdominal diameter and fetal weight estimation for detection of light-for-gestation infants in low-risk pregnancies. *British Journal of Obstetrics and Gynaecology* 1987;**94**:105-9.
- Sekar 2015** {published data only}
Sekar R, Duncombe G. Fetalweight study - Single centre experience. *BJOG: an international journal of obstetrics and gynaecology* 2015;**122**:111.
- Selbing 1984** {published data only}
Selbing A, Wichman K, Ryden G. Screening for detection of intra-uterine growth retardation by means of ultrasound. *Acta Obstetrica et Gynecologica Scandinavica* 1984;**63**:543-8.
- Semczuk-Sikora 2007** {published data only}
Semczuk-Sikora A, Krzyzanowski A, Stachowicz N, Robak J, Kraczkowski J, Kwiatek M, et al. [Maternal serum concentration of angiogenic factors: PIGF, VEGF and VEGFR-1 and placental volume in pregnancies complicated by intrauterine growth restriction]. *Ginekologia Polska* 2007;**78**:783-6.
- Shaarawy 2001** {published data only}
Shaarawy M, El Meleigy M, Rasheed K. Maternal serum transforming growth factor beta-2 in preeclampsia and eclampsia, a potential biomarker for the assessment of disease severity and fetal outcome. *Journal of the Society for Gynecologic Investigation* 2001;**8**:27-31.
- Shah 1996** {published data only}
Shah DM, Reed G. Parameters associated with adverse perinatal outcome in hypertensive pregnancies. *Journal of Human Hypertension* 1996;**10**:511-5.
- Sharf 1984** {published data only}
Sharf M, Eibschitz I, Hakim M, Degani S, Rosner B. Is serum free estriol measurement essential in the management of hypertensive disorders during pregnancy?. *European Journal of Obstetrics, Gynecology, and Reproductive Biology* 1984;**17**:365-75.
- Sheth 2016** {published data only}
Sheth T, Glantz JC. Third-trimester fetal biometry and neonatal outcomes in term and preterm deliveries. *Journal of Ultrasound in Medicine* 2016;**35**:103-10.
- Shibata 2005** {published data only}
Shibata E, Rajakumar A, Powers RW, Larkin RW, Gilmour C, Bodnar LM, et al. Soluble fms-like tyrosine kinase 1 is increased in preeclampsia but not in normotensive pregnancies with small-for-gestational-age neonates: relationship to circulating placental growth factor. *Journal of Clinical Endocrinology & Metabolism* 2005;**90**:4895-903.
- Sibiude 2012** {published data only}
Sibiude J, Guibourdenche J, Dionne MD, Le Ray C, Anselem O, Serreau R, et al. Placental growth factor for the prediction of adverse outcomes in patients with suspected preeclampsia or intrauterine growth restriction. *PLOS One* 2012;**7**:e50208.
- Sichinava 2014** {published data only}
Sichinava L, Gugushvili N. Fetal growth restriction and placental growth factor. *Journal of Maternal-Fetal and Neonatal Medicine* 2014;**27**:120.
- Singer 1970** {published data only}
Singer W, Desjardins P, Friesen HG. Human placental lactogen. An index of placental function. *Obstetrics and Gynecology* 1970;**36**:222-32.
- Smith 2014** {published data only}
Smith NA, Bukowski R, Thomas AM, Cantonwine D, Zera C, Robinson JN. Identification of pathologically small fetuses using customized, ultrasound and population-based growth norms. *Ultrasound in Obstetrics and Gynecology* 2014;**44**:595-9.
- Smith-Bindman 2002** {published data only}
Smith-Bindman R, Chu PW, Ecker JL, Feldstein VA, Filly RA, Bacchetti P. US evaluation of fetal growth: prediction of neonatal outcomes. *Radiology* 2002;**223**:153-61.

Smith-Bindman 2003 {published data only}

Smith-Bindman R, Chu PW, Ecker J, Feldstein VA, Filly RA, Bacchetti P. Adverse birth outcomes in relation to prenatal sonographic measurements of fetal size. *Journal of Ultrasound in Medicine* 2003;**22**:347-56; quiz 357-8.

Soler 1975 {published data only}

Soler NG, Nicholson HO, Malins JM. Serial determinations of human placental lactogen in the management of diabetic pregnancy. *Lancet* 1975;**2**:54-7.

Sood 1988 {published data only}

Sood M, Hingorani V, Kashyap N, Kumar S, Berry M, Bhargava S. Ultrasonic measurement of foetal parameters in normal pregnancy & in intrauterine growth retardation. *Indian Journal of Medical Research* 1988;**87**:453-8.

Sorensen 2000 {published data only}

Sorensen S, von Tabouillot D, Schioler V, Greisen G, Petersen S, Larsen T. Serial measurements of serum human placental lactogen (hPL) and serial ultrasound examinations in the evaluation of fetal growth. *Early Human Development* 2000;**60**:25-34.

Souka 2012 {published data only}

Souka AP, Papastefanou I, Pilalis A, Michalitsi V, Kassanos D. Performance of third-trimester ultrasound for prediction of small-for-gestational-age neonates and evaluation of contingency screening policies. *Ultrasound in Obstetrics and Gynecology* 2012;**39**:535-42.

Souka 2013 {published data only}

Souka AP, Papastefanou I, Pilalis A, Michalitsi V, Panagopoulos P, Kassanos D. Performance of the ultrasound examination in the early and late third trimester for the prediction of birth weight deviations. *Prenatal Diagnosis* 2013;**33**:915-20.

Sovio 2014 {published data only}

Sovio U, Smith G, Dacey A. Level 1 evidence for the diagnostic effectiveness of routine sonography as a screening test for small for gestational age (SGA) infants. *American Journal of Obstetrics and Gynecology* 2014;**1**:S408.

Spellacy 1967 {published data only}

Spellacy WN, Cohen WD, Carlson KL. Human placental lactogen levels as a measure of placental function. *American Journal of Obstetrics and Gynecology* 1967;**97**:560-1.

Spellacy 1975 {published data only}

Spellacy WN, Buhi WC, Birk SA. The effectiveness of human placental lactogen measurements as an adjunct in decreasing perinatal deaths. Results of a retrospective and a randomized controlled prospective study. *American Journal of Obstetrics and Gynecology* 1975; Vol. 121, issue 6:835-44.

Spellacy 1976 {published data only}

Spellacy WN, Buhi WC, Birk SA. Human placental lactogen and intrauterine growth retardation. *Obstetrics and Gynecology* 1976;**47**:446-8.

Spona 1971 {published data only}

Spona J, Janisch H. Serum placental lactogen (HPL) as index of placental function. *Acta Endocrinologica* 1971;**68**:401-12.

Spona 1972 {published data only}

Spona J, Janisch H. [Diagnosis of placental insufficiency by radioimmunoassay of serum HCG and HPL]. *Wiener Klinische Wochenschrift* 1972;**84**:385-9.

Stefanelli 2014 {published data only}

Stefanelli S, Groom KM. The accuracy of ultrasound-estimated fetal weight in extremely preterm infants: a comparison of small for gestational age and appropriate for gestational age. *Australian & New Zealand Journal of Obstetrics & Gynaecology* 2014;**54**:126-31.

Stefanidis 1998 {published data only}

Stefanidis K, Solomou E, Mouzakioti E, Stefanos T, Farmakides G. Comparison of somatomedin-C (SMC/IGF-I), human placental lactogen and Doppler velocimetry between appropriate and small-for-gestational-age pregnancies. *Clinical & Experimental Obstetrics & Gynecology* 1998;**25**:20-2.

Strizhakov 2013 {published data only}

Strizhakov A, Timokhina E, Ignatko I. The value of pathogenetic factors of fetal growth retardation in the early diagnosis, prevention and treatment of the placental insufficiency. *Journal of Perinatal Medicine* 2013;**41**:1.

Strom 1983 {published data only}

Strom H, Berg B, Jacobson L. Plasma estriol in late pregnancy in relation to fetal outcome. *Acta Obstetrica et Gynecologica Scandinavica* 1983;**62**:355-7.

Sucak 2010 {published data only}

Sucak A, Kanat-Pektas M, Gungor T, Mollamahmutoglu L. Leptin levels and antihypertensive treatment in preeclampsia. *Singapore Medical Journal* 2010;**51**:39-43.

Sudik 1982 {published data only}

Sudik R. [Results of an ultrasonic screening programme in the detection of intrauterine growth retardation]. *Zeitschrift fur Geburtshilfe und Perinatologie* 1982;**186**:119-24.

Sundrani 2013 {published data only}

Sundrani D, Khot V, Pisal H, Mehendale S, Wagh G, Joshi A, et al. Gestation dependant changes in angiogenic factors and their associations with fetal growth measures in normotensive pregnancy. *PLOS One* 2013;**8**:e54153.

Tajik 2012 {published data only}

Tajik P, Boers K, Wyk L, Mol B, Sisco S. Evaluation of markers guiding management decision for intrauterine growth restriction: A sub-analysis of a randomized trial, DIGITAT. *American Journal of Obstetrics and Gynecology* 2012, issue 1:S199-S200.

Takeuchi 1988 {published data only}

Takeuchi M, Morikawa H, Ueda Y, Mochizuki M. [Studies on the roles of insulin-like growth factor-1 (IGF-1)/somatomedin C

(SMC) during pregnancy]. *Nippon Naibunpi Gakkai Zasshi - Folia Endocrinologica Japonica* 1988;**64**:489-505.

Tammemae 2016 {published data only}

Tammemae L, Angerjas T, Szirko F. 5 Years of ERT-quality assurance in East-Tallinn Central Hospital, SGA. *Journal of Maternal-Fetal and Neonatal Medicine* 2016;**29**:306-7.

Tayama 1983 {published data only}

Tayama C, Ichimaru S, Ito M, Nakayana M, Maeyama M, Miyakawa I. Unconjugated estradiol, estriol and total estriol in maternal peripheral vein, cord vein, and cord artery serum at delivery in pregnancies with intrauterine growth retardation. *Endocrinologia Japonica* 1983;**30**:155-62.

Taylor 2003 {published data only}

Taylor RN, Grimwood J, Taylor RS, McMaster MT, Fisher SJ, North RA. Longitudinal serum concentrations of placental growth factor: evidence for abnormal placental angiogenesis in pathologic pregnancies. *American Journal of Obstetrics and Gynecology* 2003;**188**:177-82.

Teoh 1971 {published data only}

Teoh ES, Spellacy WN, Buhi WC. Human chorionic somatomammotrophin (HCS): a new index of placental function. *Journal of Obstetrics & Gynaecology of the British Commonwealth* 1971;**78**:673-85.

Tonari 1987 {published data only}

Tonari M. Studies on the relation between fetal heart rate change, placental finding and fetal outcome. [Japanese]. *Acta Obstetrica et Gynaecologica Japonica* 1987;**39**:751-9.

Torok 1987 {published data only}

Torok A, Csernus V, Csaba I. Diagnostic value of serum estriol determination in placental insufficiency. [Hungarian]. *Orvosi Hetilap* 1987;**128**(52):2731-2, 2735.

Triunfo 2014 {published data only}

Triunfo S, Lobmaier S, Parra-Saavedra M, Crovetto F, Peguero A, Nadal A, et al. Angiogenic factors at diagnosis of late-onset small-for-gestational age and histological placental underperfusion. *Placenta* 2014;**35**:398-403.

Triunfo 2016 {published data only}

Triunfo S, Parra-Saavedra M, Rodriguez-Sureda V, Crovetto F, Dominguez C, Gratacos E, et al. Angiogenic factors and Doppler evaluation in normally growing fetuses at routine third-trimester scan: prediction of subsequent low birth weight. *Fetal Diagnosis and Therapy* 2016;**40**:13-20.

Triunfo 2016a {published data only}

Triunfo S, Crovetto F, Scaccocchio E, Parra-Saavedra M, Gratacos E, Figueras F. Contingent versus routine third-trimester screening for late fetal growth restriction. *Ultrasound in Obstetrics and Gynecology* 2016;**47**:81-8.

Tsiakkas 2015 {published data only}

Tsiakkas A, Duvdevani N, Wright A, Wright D, Nicolaides KH. Serum placental growth factor in the three trimesters of

pregnancy: effects of maternal characteristics and medical history. *Ultrasound in Obstetrics & Gynecology* 2015;**45**:591-8.

Tsiakkas 2016 {published data only}

Tsiakkas A, Saiid Y, Wright A, Wright D, Nicolaides KH. Competing risks model in screening for preeclampsia by maternal factors and biomarkers at 30-34 weeks' gestation. *American Journal of Obstetrics and Gynecology* 2016;**215**:e1-e17.

Turpin 2015 {published data only}

Turpin CA, Sakyi SA, Owiredo WK, Ephraim RK, Anto EO. Association between adverse pregnancy outcome and imbalance in angiogenic regulators and oxidative stress biomarkers in gestational hypertension and preeclampsia. *BMC Pregnancy and Childbirth* 2015;**15**:189.

Van Rijn 2015 {published data only}

Van Rijn BB, Maguire P, Clarke S, Roland A, Reading IC, Cameron IT, et al. Salivary uric acid levels in women destined to develop preeclampsia: Longitudinal prospective study. *Reproductive Sciences* 2015;**22**:93A-4A.

Varma 1979 {published data only}

Varma TR, Taylor H, Bridges C. Ultrasound assessment of fetal growth. *British Journal of Obstetrics and Gynaecology* 1979;**86**:623-32.

Varma 1982 {published data only}

Varma TR. Serum uric acid levels as an index of fetal prognosis in pregnancies complicated by preexisting hypertension and preeclampsia of pregnancy. *International Journal of Gynaecology and Obstetrics* 1982;**20**:401-8.

Vatten 2012 {published data only}

Vatten LJ, Asvold BO, Eskild A. Angiogenic factors in maternal circulation and preeclampsia with or without fetal growth restriction. *Acta Obstetrica et Gynecologica Scandinavica* 2012;**91**:1388-94.

Vinayagam 2015 {published data only}

Vinayagam V, Bobby Z, Habeebullah S, Chaturvedula L, Bharadwaj SK. Impaired angiogenesis and pregnancy outcome in patients with hypertensive disorders of pregnancy: A pilot study in an Indian population. *Indian Journal of Clinical Biochemistry* 2015;**30**:S117.

Wallner 2007 {published data only}

Wallner W, Sengenberger R, Strick R, Strissel PL, Meurer B, Beckmann MW, et al. Angiogenic growth factors in maternal and fetal serum in pregnancies complicated by intrauterine growth restriction. *Clinical Science* 2007;**112**:51-7.

Watson 1973 {published data only}

Watson D, Siddiqui SA, Stafford JE, Gibbard S, Hewitt V. A comparative study of five laboratory tests for foeto-placental dysfunction in late pregnancy. *Journal of Clinical Pathology* 1973;**26**:249-300.

Weissbach 1985 {published data only}

Weissbach A, Freymann E, Hubl W, Seefried W, Neef B, Thiele HJ. Enzymeimmunoassay of unconjugated estriol in serum and saliva during pregnancy. *Experimental & Clinical Endocrinology* 1985;**86**:178-84.

Whigham 1980 {published data only}

Whigham KA, Howie PW, Shah MM, Prentice CR. Factor VIII related antigen/coagulant activity ratio as a predictor of fetal growth retardation: a comparison with hormone and uric acid measurements. *British Journal of Obstetrics and Gynaecology* 1980;**87**:797-803.

White 2016 {published data only}

White SW, Marsh JA, Lye SJ, Briollais L, Newnham JP, Pennell CE. Improving customized fetal biometry by longitudinal modelling. *Journal of Maternal-Fetal & Neonatal Medicine* 2016;**29**:1888-94.

Woelkers 2016 {published data only}

Woelkers DA, Von Dadelszen P, Sibai B. Placenta Growth Factor (PLGF) predicts time to delivery in women with signs or symptoms of early preterm preeclampsia. *American Journal of Obstetrics and Gynecology* 2016;**1**:S25-6.

Woo 2016 {published data only}

Woo I, Ingles S, Sriprasert I, Paulson R, Stanczyk F, Chung K. The role of angiogenic markers in adverse perinatal outcomes: Fresh vs frozen embryo transfers. *Fertility and Sterility* 2016;**1**:e7.

Woods 2015 {published data only}

Woods A, Dekker G. Placental growth factor measurement as a marker of subsequent disease and harm in placental insufficiency conditions. *Pregnancy Hypertension* 2015;**5**(1):50.

Wurz 1983 {published data only}

Wurz H, Luben G, Bohn H, Kunzig HJ, Geiger W. Concentration of placental protein 10 (PP10) in maternal serum and amniotic fluid throughout normal gestation and in pregnancy complicated by fetal growth retardation. *Archives of Gynecology* 1983;**233**:165-74.

Xing 2016 {published data only}

Xing Y, Chang RJ, Du XN, Chen D. Placental growth factor, an index for detection of women with preeclampsia. *International Journal of Clinical and Experimental Medicine* 2016;**9**:11348-54.

Xu 2015 {published data only}

Xu Y, Lek N, Cheung YB, Biswas A, Su LL, Kwek KY, et al. Unconditional and conditional standards for fetal abdominal circumference and estimated fetal weight in an ethnic Chinese population: a birth cohort study. *BMC Pregnancy & Childbirth* 2015;**15**:141.

Yamaguchi 1979 {published data only}

Yamaguchi H, Nishiyama Y, Nose Y, Sugiyama Y. [Usefulness of serum HPL determination as an indicator of placental function and fetal growth]. *Horumon to Rinsho - Clinical Endocrinology* 1979;**27**:191-6.

Yanaihara 1984 {published data only}

Yanaihara T, Hirato K, Seo F, Mitsukawa G, Kojima S, Maruyama S, et al. [Prenatal diagnosis of IUGR by assessing the multiple hormone concentration in maternal peripheral blood]. *Nippon Sanka Fujinka Gakkai Zasshi - Acta Obstetrica et Gynaecologica Japonica* 1984;**36**:182-8.

Zail 1975 {published data only}

Zail SS, Safro IL. Serum human placental lactogen levels in intra-uterine fetal growth retardation. *South African Medical Journal. Suid-Afrikaanse Tydskrif Vir Geneeskunde* 1975;**49**:2022-4.

Zera 2011 {published data only}

Zera C, Smith NA, Robinson JN, Bukowski RK. Improved identification of pregnancies at risk for stillbirth with ultrasound based estimates of individualized fetal growth potential. *American Journal of Obstetrics and Gynecology* 2011;**204**:S150-1.

Zhang 2011 {published data only}

Zhang J, Mikolajczyk R, Grewal J, Neta G, Klebanoff M. Prenatal application of the individualized fetal growth reference. *American Journal of Epidemiology* 2011; Vol. 173, issue 5:539-43.

Zhao 2010 {published data only}

Zhao W, Qiao J, Zhang Q, Zhao Y, Chen Q. Levels of antiangiogenic factors in preeclamptic pregnancies. *Growth Factors* 2010;**28**:293-8.

Zlatnik 1979 {published data only}

Zlatnik FJ, Varner MW, Hauser KS. Human placental lactogen: a predictor of perinatal outcome?. *Obstetrics and Gynecology* 1979;**54**:205-10.

Zuckerman 1974 {published data only}

Zuckerman H, Gendler L, Schwarz M, Harpaz S. Human placental lactogen as an index of placental function. *Israel Journal of Medical Sciences* 1974;**10**:490-4.

References to studies awaiting assessment
Bracali 1968 {published data only}

Bracali R, Corrado F. Clinical value of urinary estriol as an index of placental function and of fetal vitality in some pathological conditions of pregnancy. [Italian]. *Quaderni di Clinica Ostetrica e Ginecologica* 1968;**23**:1534-52.

Fuks 1990 {published data only}

Fuks MA, Milovanov AP, Chekhonatskaia ML. Prognostic value of placental maturity staging for pregnancy outcome of the fetus and newborn infant. *Akusherstvo i Ginekologiya* 1990;**7**(3):19-22.

Jain 2000 {published data only}

Jain A, Kumar G, Gupta AK. Relationship of placental grades to gestational age and early grade III placenta and its relation with growth retardation. *Ultrasound International* 2000;**6**:112-20.

Khan 2004 {published data only}

Khan DB, Bari V, Chishty IA. Ultrasound in the diagnosis and management of intrauterine growth retardation. *Journal of the College of Physicians and Surgeons Pakistan* 2004;**14**:601-4.

Ruseva 1983 {published data only}

Ruseva R, Marinov B. [Prognostic potentials of certain parameters in fetal growth retardation]. *Akusherstvo i Ginekologija* 1983;**22**:367-72.

Ruseva 1985 {published data only}

Ruseva R, Katsulov A, Marinov B, Atanasov B, Milkov V. [The placenta. Ultrasonic and hormonal changes in fetal growth retardation]. *Akusherstvo i Ginekologija* 1985;**24**:16-9.

Ruseva 1985a {published data only}

Ruseva R, Khadzhiev A, Marinov B. [Ultrasonic study of placental intrauterine growth]. *Akusherstvo i Ginekologija* 1985;**24**:75-80.

Ruseva 1988 {published data only}

Ruseva R. [Echographic and hormonal parameters in fetal growth retardation]. *Akusherstvo i Ginekologija* 1988;**27**:15-8.

Serban 1971 {published data only}

Serban MD. [Investigation of placental functional values. II. The placental lactogenic hormone (HPL)]. *Studii Si Cercetari de Endocrinologie* 1971;**22**:159-60.

Additional references

Alere 2015

Alere. Alere Triage® PIGF Test. <http://www.alere.com/ww/en/product-details/triage-plgf-test.html> (accessed 27th August 2015).

Alfirevic 2015

Alfirevic Z, Stampalija T, Medley N. Fetal and umbilical Doppler ultrasound in normal pregnancy. *Cochrane Database of Systematic Reviews* 2015, Issue 4. [DOI: [10.1002/14651858.CD001450.pub4](https://doi.org/10.1002/14651858.CD001450.pub4)]

Alfirevic 2017

Alfirevic Z, Stampalija T, Dowswell T. Fetal and umbilical Doppler ultrasound in high-risk pregnancies. *Cochrane Database of Systematic Reviews* 2017, Issue 6. [DOI: [10.1002/14651858.CD007529.pub4](https://doi.org/10.1002/14651858.CD007529.pub4)]

Bossuyt 2015

Bossuyt PM, Reitsma JB, Bruns DE, Gatsonis CA, Glasziou PP, Irwig L, et al. STARD 2015: an updated list of essential items for reporting diagnostic accuracy studies. *BMJ* 2015;**351**:h5527.

Bricker 2015

Bricker L, Medley N, Pratt JJ. Routine ultrasound in late pregnancy (after 24 weeks' gestation). *Cochrane Database of Systematic Reviews* 2015, Issue 6. [DOI: [10.1002/14651858.CD001451.pub4](https://doi.org/10.1002/14651858.CD001451.pub4)]

Clausson 2001

Clausson B, Gardosi J, Francis A, Cnattingius S. Perinatal outcome in SGA births defined by customised versus population-based birthweight standards. *BJOG: an international journal of obstetrics and gynaecology* 2001;**108**(8):830-4.

Duenholter 1976

Duenholter JH, Whalley PJ, MacDonald PC. An analysis of the utility of plasma immunoreactive estrogen measurements in determining delivery time of gravidas with a fetus considered at high risk. *American Journal of Obstetrics and Gynecology* 1976;**125**(7):889-98.

Ensor 2018

Ensor J, Deeks JJ, Martin EC, Riley RD. Meta-analysis of test accuracy studies using imputation for partial reporting of multiple thresholds. *Research Synthesis Methods* 2018;**9**(1):100-15.

Flenady 2011

Flenady V, Koopmans L, Middleton P, Frøen JF, Smith GC, Gibbons K, et al. Major risk factors for stillbirth in high-income countries: a systematic review and meta-analysis. *Lancet* 2011;**377**(9774):1331-40.

Flenady 2016

Flenady V, Wojcieszek AM, Middleton P, Ellwood D, Erwich JJ, Coory M, et al. Stillbirths: recall to action in high-income countries. *Lancet* 2016; Vol. 387, issue 10019:13-9.

Gardosi 2013

Gardosi J, Madurasinghe V, Williams M, Malik A, Francis A. Maternal and fetal risk factors for stillbirth: population based study. *BMJ* 2013;**346**:f108.

GRIT 2003

GRIT Study Group. A randomised trial of timed delivery for the compromised preterm fetus: short term outcomes and Bayesian interpretation. *BJOG: an international journal of obstetrics and gynaecology* 2003;**110**(1):27-32.

Heazell 2013

Heazell AE, Bernatavicius G, Roberts SA, Garrod A, Whitworth MK, Johnstone ED, et al. A randomised controlled trial comparing standard or intensive management of reduced fetal movements after 36 weeks gestation--a feasibility study. *BMC Pregnancy Childbirth* 2013;**13**:95. [DOI: [10.1186/1471-2393-13-95](https://doi.org/10.1186/1471-2393-13-95)]

Heazell 2015a

Heazell AE, Worton SA, Higgins LE, Ingram E, Johnstone ED, Jones RL, et al. IFPA Gábor Than Award Lecture: Recognition of placental failure is key to saving babies' lives. *Placenta* 2015;**36**(Suppl 1):S20-8.

Heazell 2015b

Heazell AE, Whitworth MK, Duley L, Thornton JG. Use of biochemical tests of placental function for improving pregnancy outcome. *Cochrane Database of Systematic Reviews* 2015, Issue 11. [DOI: [10.1002/14651858.CD011202.pub2](https://doi.org/10.1002/14651858.CD011202.pub2)]

Lawn 2016

Lawn JE, Blencowe H, Waiswa P, Amouzou A, Mathers C, Hogan D, et al. Stillbirths: rates, risk factors, and acceleration towards 2030. *Lancet* 2016;**387**(10018):587-603.

Macaskill 2010

Macaskill P, Gatsonis C, Deeks JJ, Harbord RM, Takwoingi Y. Chapter 10: analysing and presenting results. In: Deeks JJ, Bossuyt PM, Gatsonis C editor(s). *Cochrane handbook for systematic reviews of diagnostic test accuracy*. Version 1.0. The Cochrane Collaboration, 2010.

MacKay 2010

MacKay DF, Smith CG, Dobbie R, Pell JP. Gestational age at delivery and special educational need: retrospective cohort study of 407,503 schoolchildren. *PLOS Medicine* 2010;**7**(6):e1000289.

McCowan 2007

McCowan LM, George-Haddad M, Stacey T, Thompson JM. Fetal growth restriction and other risk factors for stillbirth in a New Zealand setting. *Australian & New Zealand Journal of Obstetrics & Gynaecology* 2007;**47**(6):450-6.

Morris 2011

Morris RK, Malin G, Robson SC, Kleijnen J, Zamora J, Khan KS. Fetal umbilical artery Doppler to predict compromise of fetal/neonatal wellbeing in a high-risk population: systematic review and bivariate meta-analysis. *Ultrasound in Obstetrics & Gynecology* 2011;**37**(2):135-42.

NHS England 2016

NHS England. Saving Babies' Lives. A care bundle for reducing stillbirth. NHS England, 2016.

NICE 2008

National Institute for Health and Clinical Excellence. Clinical Guideline 62 - Antenatal Care. National Institute for Health and Clinical Excellence, 2008.

Perkin Elmer 2015

Perkin Elmer. The potential of PP13 in pre-eclampsia early detection. http://www.perkinelmer.co.uk/Content/TechnicalInfo/TCH_PP13NonUS.pdf (accessed 27th August 2015).

Peters 2018

Peters LL, Thornton C, de Jonge A, Khashan A, Tracy M, Downe S, et al. The effect of medical and operative birth interventions on child health outcomes in the first 28 days and up to 5 years of age: A linked data population-based cohort study. *Birth* 2018;**45**(4):347-57.

Proud 1987

Proud J, Grant AM. Third trimester placental grading by ultrasonography as a test of fetal wellbeing. *British Medical Journal* 1987;**294**(6588):1641-4.

Ptacek 2014

Ptacek I, Sebire NJ, Man JA, Brownbill P, Heazell AE. Systematic review of placental pathology reported in association with stillbirth. *Placenta* 2014;**35**(8):552-62.

RCOG 2011

Royal College of Obstetricians and Gynaecologists. Reduced Fetal Movements: Guideline 57. Royal College of Obstetricians and Gynaecologists, 2011.

RCOG 2013

Royal College of Obstetricians and Gynaecologists. Small-for-Gestational-Age Fetus, Investigation and Management (Green-top Guideline No. 31). Royal College of Obstetricians and Gynaecologists, 2013.

RCOG 2014

Royal College of Obstetricians and Gynaecologists. The Investigation and Management of the Small-for-Gestational-Age Fetus - Guideline 31. Royal College of Obstetricians and Gynaecologists, 2014.

Riley 2015

Riley RD, Ahmed I, Ensor J, Takwoingi Y, Kirkham A, Morris RK, et al. Meta-analysis of test accuracy studies: an exploratory method for investigating the impact of missing thresholds. *Systematic Reviews* 2015;**4**:12.

Roche 2015

Roche. Elecsys® Preeclampsia (sFlt-1 & PlGF). <http://www.cobas.com/home/product/clinical-and-immunochemistry-testing/elecsys-preeclampsia-assays-sFlt-1-PlGF.html> (accessed 27th August 2015).

Rutter 2001

Rutter CM, Gatsonis CA. A hierarchical regression approach to meta-analysis of diagnostic test accuracy evaluations. *Statistics in Medicine* 2001;**20**(19):2865-84.

Smith 2002

Smith GC, Stenhouse EJ, Crossley JA, Aitken DA, Cameron AD, Connor JM. Early pregnancy levels of pregnancy-associated plasma protein a and the risk of intrauterine growth restriction, premature birth, preeclampsia, and stillbirth. *Journal of Clinical Endocrinology and Metabolism* 2002;**87**(4):1762-7.

Smith 2006

Smith GC, Shah I, Crossley JA, Aitken DA, Pell JP, Nelson SM, et al. Pregnancy-associated plasma protein A and alpha-fetoprotein and prediction of adverse perinatal outcome. *Obstetrics and Gynecology* 2006;**107**(1):161-6.

Steinhauser 2016

Steinhauser S, Schumacher M, Rücker G. Modelling multiple thresholds in meta-analysis of diagnostic test accuracy studies. *BMC Medical Research Methodology* 2016;**16**(1):97.

Stock 2012

Stock SJ, Ferguson E, Duffy A, Ford I, Chalmers J, Norman JE. Outcomes of elective induction of labour compared with

expectant management: population based study. *BMJ* 2012;**344**:e2838.

Takwoingi 2013

Takwoingi Y, Leeflang MM, Deeks JJ. Empirical evidence of the importance of comparative studies of diagnostic test accuracy. *Annals of Internal Medicine* 2013;**158**(7):544-54.

Whiting 2011a

Whiting P, Westwood M, Beynon R, Burke M, Sterne JA, Glanville J. Inclusion of methodological filters in searches for diagnostic test accuracy studies misses relevant studies. *Journal of Clinical Epidemiology* 2011;**64**(6):602-7.

Whiting 2011b

Whiting PF, Rutjes AW, Westwood ME, Mallett S, Deeks JJ, Reitsma JB, et al. QUADAS-2: a revised tool for the quality

assessment of diagnostic accuracy studies. *Annals of Internal Medicine* 2011;**155**(8):529-36.

Worton 2014

Worton S, Sibley CP, Heazell AE. Understanding the placental aetiology of fetal growth restriction; could this lead to personalized management strategies?. *Fetal and Maternal Medicine Review* 2014;**25**(2):95-116.

References to other published versions of this review

Heazell 2016

Heazell AE, Hayes DJ, Whitworth M, Takwoingi Y, Bayliss SE, Davenport C. Diagnostic accuracy of biochemical tests of placental function versus ultrasound assessment of fetal size for stillbirth and small-for-gestational-age infants. *Cochrane Database of Systematic Reviews* 2016, Issue 6. [DOI: [10.1002/14651858.CD012245](https://doi.org/10.1002/14651858.CD012245)]

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Altmann 1978

Study characteristics			
Patient sampling	Case reports of 10 high-risk pregnancies		
Patient characteristics and setting	Sample size: 10 women Gestation at sampling: > 26 weeks Risk: high risk		
Index tests	hPL measured in serum, values classifies as normal/border-line/abnormal, from a reference group of 242 pregnant women. Grade III used as threshold for placental grading.		
Target condition and reference standard(s)	SGA defined as birthweight under the 10th percentile for gestational age		
Flow and timing			
Comparative			
Notes	Paper translated from German		
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		

Altmann 1978 *(Continued)*

Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test hPL			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Unclear
DOMAIN 2: Index Test Placental grading			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Unclear		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Unclear	

Amini 2014

Study characteristics	
Patient sampling	Prospective multicentric cohort study of singleton pregnancies between 28 and 42 weeks.
Patient characteristics and setting	Sample size: 404 Gestation at sampling: 24 hours before delivery Risk: mixed (some exclusions due to hypertension)

Amini 2014 (Continued)

Setting: Vali-Asr and Akbar-Abadi teaching hospitals of Tehran University of Medical Sciences, Iran

NICU admission: 79 neonates required NICU admission, 31 from women with hyperuricaemia

Index tests	Blood samples taken within 24 hours preceding delivery and uric acid levels were determined using the enzymatic colorimetric method. Hyperuricemia defined as serum uric acid level 1 SD greater than the appropriate for gestational age as defined by Lind and colleagues.		
Target condition and reference standard(s)	SGA defined as birthweight < 10th percentile for gestational age according to Fenton growth charts		
Flow and timing	559 women were asked to participate, 404 met inclusion criteria		
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test Uric acid			
Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear		
If a threshold was used, was it pre-specified?	Yes		
		Unclear	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			

Amini 2014 (Continued)

Was there an appropriate interval between index test and reference standard?	No
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
Low	

Baird 2016

Study characteristics			
Patient sampling	Retrospective cohort study, consecutive enrolment of women clinically suspected of FGR		
Patient characteristics and setting	Sample size: 107 Gestation at sampling: 35-38 weeks (within 2 weeks of birth) Risk: high risk (suspected FGR, previous FGR, maternal medical conditions, decreased fetal movements) Setting: university teaching hospital in Victoria, Australia Mode of delivery: 45.8% normal vaginal delivery, 14.1% instrumental delivery, 12.1% elective caesarean section, 28% emergency caesarean section		
Index tests	EFW calculated using Hadlock charts		
Target condition and reference standard(s)	SGA defined as birthweight < 10th centile for gestational age using the most recent Australian birthweight centiles		
Flow and timing	107 pregnancies met the inclusion criteria over a 12-month study period; all were included		
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High

Baird 2016 (Continued)

DOMAIN 2: Index Test EFW

Were the index test results interpreted without knowledge of the results of the reference standard?	No		
If a threshold was used, was it pre-specified?	Yes		
		Unclear	Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Barel 2016
Study characteristics

Patient sampling	Women referred to the gynaecologic ultrasound unit for SEFW 1 week prior to delivery
Patient characteristics and setting	Sample size: 14,089 Gestation at sampling: 1 week before delivery (24-41 weeks) Risk: mixed Setting: Assaf Harofe Medical Centre
Index tests	
Target condition and reference standard(s)	SGA defined as less than the 10th percentile
Flow and timing	
Comparative	
Notes	53 cases (4.3%) of SGA were delivered before 34 weeks of gestation

Barel 2016 (Continued)

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Beischer 1991
Study characteristics

Patient sampling	Consecutive patients from 1971-1984 and 1985-1989
Patient characteristics and setting	Sample size: 72,062 Gestation at sampling: 30-34 weeks

Beischer 1991 (Continued)

Risk: mixed

Setting: Mercy Maternity Hospital, Melbourne

Mode of delivery: 13.6% delivered by caesarean section (12.7% of normal UE3 group and 21.5% of low UE3 group)

Index tests	24-hour urinary oestriol excretion, threshold 10th centile according to gestational age
Target condition and reference standard(s)	FGR defined as birthweight below 10th centile
Flow and timing	51,427 patients tested from 1971-1984 and 20,635 tested from 1985-1989. Not all patients who delivered were tested due to emergency admissions, premature deliveries, and administrative failures (there were 85,000 total deliveries).
Comparative	
Notes	High oestriol levels were an indicator for early delivery

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test Urinary E3			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	High
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			

Beischer 1991 (Continued)

Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	No
Low	

Bellomo 2011

Study characteristics	
Patient sampling	Prospective cohort of women admitted for suspected hypertension during pregnancy
Patient characteristics and setting	Sample size: 163 Gestation at sampling: 30.4+/- 4.1 weeks Risk: high (all suspected hypertension, 44.7% developed pre-eclampsia) Mode of delivery: 39% delivered by caesarean section Setting: San Giovanni Battista Hospital, Foligno, Italy
Index tests	Uric acid measured in serum, hyperuricemia defined as > 309 umol/L
Target condition and reference standard(s)	SGA defined as those weighing less than the 10th centile based on nationwide derived centile charts for singleton births
Flow and timing	Some patients did not complete the study due to not meeting the BP criteria on entry, elevated proteinuria, incomplete data, insufficient BP recordings, or withdrawal of consent.
Comparative	
Notes	

Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High

Bellomo 2011 (Continued)

DOMAIN 2: Index Test Uric acid

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Unclear		
		Unclear	Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	No		
		Unclear	

Ben-Haroush 2007
Study characteristics

Patient sampling	Women with healthy singleton pregnancies recruited at time of delivery
Patient characteristics and setting	Sample size: 259 Gestation at sampling: 28-34 weeks Risk: low (no obstetric complications, hypertensive and diabetic pregnancies were excluded)
Index tests	Estimated fetal weight calculated using Hadlock's formula and converted to percentiles using locally developed growth charts
Target condition and reference standard(s)	SGA defined as < 10th centile for gestational age
Flow and timing	259 women were included in the study
Comparative	
Notes	

Ben-Haroush 2007 *(Continued)*
Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		Low	High
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Unclear	

Benton 2016
Study characteristics

Patient sampling	Prospective cohort of suspected FGR pregnancies from 3 centres
Patient characteristics and setting	Sample size: 411 Age: 18-45

Benton 2016 (Continued)

Gestation at sampling: 20-41 weeks (IQR 29.8, 36)

Risk: high (suspected FGR, hypertension and pre-eclampsia excluded)

Setting: 3 sites in Canada, New Zealand, UK

Mode of delivery: 92 inductions, 38 caesarean, 7 instrumental

NICU admission: 36 admission, 26 of these from the low PLGF group

Index tests	PLGF measured using Triage immunoassay, very low PLGF defined as < 12 pg/mL
Target condition and reference standard(s)	SGA defined as below the 3rd centile. Birthweight centile determined by the Canadian standard for multi-ethnicity.
Flow and timing	Women recruited from inpatient and outpatient centres at BC Women's Hospital, Vancouver and Ottawa Hospital; cohort of FGR pregnancies from Auckland, New Zealand with banked maternal blood samples; cohort of FGR pregnancies from the UK (PELICAN-FGR Study)
Comparative	
Notes	Placental pathology data were available for 213 women; 55 of 94 with a low PLGF were grade II or III, as was one of the 119 with a normal PLGF.

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test PLGF			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		

Benton 2016 (Continued)

	Low	High
DOMAIN 4: Flow and Timing		
Was there an appropriate interval between index test and reference standard?	Yes	
Did all patients receive the same reference standard?	Yes	
Were all patients included in the analysis?	Yes	
	Low	

Berkowitz 1988

Study characteristics				
Patient sampling		Women with singleton pregnancies selected on the basis of known risk factors or clinical suspicion of IUGR		
Patient characteristics and setting		Sample size: 168 Gestation at sampling: 30 to 42 weeks (all last measurements used) Risk: high (clinical suspicion of IUGR, previous infant with IU-GR, complications associated with IUGR, smoking, alcohol/drug abuse, postdates) Setting: Mount Sinai Medical Centre, New York		
Index tests		EFW calculated using Shepard formula		
Target condition and reference standard(s)		SGA defined as below 10th centile for gestational age at birth on the basis of the nomogram by Brenner and colleagues		
Flow and timing				
Comparative				
Notes				
Methodological quality				
Item	Authors' judgement	Risk of bias	Applicability concerns	
DOMAIN 1: Patient Selection				
Was a consecutive or random sample of patients enrolled?	Yes			
Was a case-control design avoided?	Yes			
Did the study avoid inappropriate exclusions?	Yes			

Berkowitz 1988 (Continued)

		Low	High
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Bikmetova 2013

Study characteristics	
Patient sampling	Retrospective cohort study
Patient characteristics and setting	Sample size: 518 Gestation at sampling: 3rd trimester Risk: unknown
Index tests	Ultrasound formula not known
Target condition and reference standard(s)	IUGR definition unknown
Flow and timing	
Comparative	
Notes	
Methodological quality	

Bikmetova 2013 (Continued)

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Unclear
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear		
If a threshold was used, was it pre-specified?	Unclear		
		Unclear	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Unclear		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Unclear
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Unclear		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Unclear	

Callec 2015

Study characteristics	
Patient sampling	Prospective cohort study across 2 centres between 2003 and 2006, diabetes and illiteracy excluded, as well as intention to deliver outside the hospital or to move outside the region within 3 years.
Patient characteristics and setting	Sample size: 1897 Gestation at sampling: 30-35 weeks Risk: mixed

Biochemical tests of placental function versus ultrasound assessment of fetal size for stillbirth and small-for-gestational-age infants (Review)

72

Callec 2015 (Continued)

Setting: EDEN study, 2 university maternity centres, France

Mode of delivery: 1404 normal vaginal deliveries, 195 instrumental, 298 caesarean sections

NICU admission: 128 admissions, 20 of these from the EFW < 10 group

Index tests	EFW measured using Hadlock formula
Target condition and reference standard(s)	SGA defined as birthweight below the 10th centile
Flow and timing	2002 women originally included in the cohort; 80 were lost to follow-up, declined to continue participation, or experienced fetal death. 1 woman with a stillbirth was also excluded.
Comparative	
Notes	Some elective caesarean sections and inductions were performed due to reduced fetal growth; there were higher rates in the FP compared with the FN group.

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		Unclear	Low
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear		
If a threshold was used, was it pre-specified?	Yes		
		Unclear	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			

Callec 2015 (Continued)

Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
Low	

Campbell 1972

Study characteristics			
Patient sampling	All women at risk of FGR		
Patient characteristics and setting	Sample size: 284 Gestation at sampling: unknown Risk: high (268 women clinically suspected of having a small uterus, 16 diabetic or with a bad obstetric history) Setting: Queen Charlotte's Maternity Hospital Mode of delivery: 43 instrumental deliveries, 26 EmCS, 29 EICS		
Index tests	Oestrogens measured as oestriol in 26 cases and total oestrogens in the rest. One or no abnormal measurements classed as normal, 2+ classed as abnormal; measurements did not have to be consecutive.		
Target condition and reference standard(s)	SGA defined as a birthweight below the 5th centile		
Flow and timing			
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Unclear		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test Urinary E3			

Campbell 1972 (Continued)

Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear
If a threshold was used, was it pre-specified?	Yes
	Low High
DOMAIN 3: Reference Standard	
Is the reference standards likely to correctly classify the target condition?	Yes
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear
	Low High
DOMAIN 4: Flow and Timing	
Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
	Low

Cedard 1979

Study characteristics	
Patient sampling	All patients were clinically suspected of IUGR due to lower than expected SFH on 2 consecutive visits
Patient characteristics and setting	Sample size: 64 Gestation at sampling: 37-39 weeks Risk: high (all clinically suspected of IUGR due to lower than expected SFH on 2 consecutive visits) Setting: Maternite de Port-Royal, France
Index tests	Oestriol measured in plasma using the oestriol RIA kit IM 82, levels for normal pregnancies established from 301 values obtained from 88 judged to be free of complications and low defined as < 10th percentile
Target condition and reference standard(s)	SGA defined as below the 10th centile, birthweights calculated using Lubchenko curve
Flow and timing	Plasma total oestriol was studied in 222 pregnancies, 64 of these had measurements at 37-39 weeks
Comparative	

Cedard 1979 (Continued)

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test E3			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Chaiworapongsa 2013
Study characteristics

Patient sampling	Prospective cohort between November 2003 and August 2006
Patient characteristics and setting	Sample size: 1269

Chaiworapongsa 2013 (Continued)

Gestation at sampling: 30-34 weeks

Risk: mixed (PE excluded in early pregnancy, otherwise mixed)

Setting: Sotero del Rio Hospital, Santiago, Chile

Index tests	PIGF/sVEGFR-1 and PIGF/sEng
Target condition and reference standard(s)	SGA defined as birthweight below 10th centile for gestational age according to the Chilean birthweight distribution of a Hispanic population. Stillbirth defined as death of a fetus before delivery that was not a consequence of an induced termination of pregnancy (including intrapartum and antepartum stillbirth)
Flow and timing	Patients were excluded if they delivered < 34 weeks (n = 29) and did not have a plasma sample collected at 30-34 weeks (n = 326)

Comparative

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test PIGF			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	No		
		High	Unclear
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			

Chaiworapongsa 2013 (Continued)

Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	No
Low	

Chard 1985

Study characteristics			
Patient sampling	Antenatal patients with 3 or more blood samples at weekly intervals from the 36th week onwards		
Patient characteristics and setting	Sample size: 392 Gestation at sampling: 36 weeks onwards Risk: mixed (144 with varying degrees of pre-eclampsia) Setting: Solihull Hospital, Birmingham UK		
Index tests	hPL and oestriol measured in serum using well-established commercial kits		
Target condition and reference standard(s)	IUGR defined as a delivered weight below the 10th percentile for the group (2775 g)		
Flow and timing			
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test hPL			
Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear		

Chard 1985 (Continued)

If a threshold was used, was it pre-specified?	Yes		
		Unclear	Low
DOMAIN 2: Index Test E3			
Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear		
If a threshold was used, was it pre-specified?	Yes		
		Unclear	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	High
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Chauhan 1999

Study characteristics			
Patient sampling	162 patients with oligohydramnios and 162 patients with adequate amniotic fluid		
Patient characteristics and setting	Sample size: 324 Gestation at sampling: 3rd trimester, mean 34 weeks Risk: mixed Setting: Spartanburg Regional Medical Center, Jackson, Mississippi		
Index tests	EFW, Hadlock formula		
Target condition and reference standard(s)	SGA defined as below 10th centile for gestational age		
Flow and timing			

Chauhan 1999 (Continued)

Comparative

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Chauhan 1999a
Study characteristics

Patient sampling	287 women with pre-eclampsia and 287 healthy controls
------------------	---

Chauhan 1999a (Continued)

Patient characteristics and setting

Sample size: 574

Gestation at sampling: average 35 weeks

Risk: mixed

Setting: Spartanburg Regional Medical Center, Jackson, Mississippi

Index tests

Target condition and reference standard(s)

SGA defined as below the 10th centile for gestational age

Flow and timing

Comparative

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		

Chauhan 1999a (Continued)

Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
Low	

Chauhan 2003

Study characteristics			
Patient sampling	Retrospective identification of all pregnant patients with hypertension delivered during a 5-year period at 3 centres.		
Patient characteristics and setting	Sample size: 264 Gestation at sampling: within 21 days of delivery Risk: high (all women had chronic hypertension) Setting: centres in Australia (1) and the USA (2)		
Index tests	EFW estimated using Hadlock formula		
Target condition and reference standard(s)	IUGR defined as birthweight < 10% for gestational age using the fetal growth curve by Alexander and colleagues in the USA and an Australian growth curve in the other centre		
Flow and timing	All women with chronic hypertension defined according to ACOG criteria who delivered within 5-year period, had a reliable gestational age, and a SEFW within 3 weeks of delivery. Known fetal anomalies, multiple gestations, gestational hypertension, pre-eclampsia, diabetes mellitus were excluded.		
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test EFW			

Chauhan 2003 (Continued)

Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear		
If a threshold was used, was it pre-specified?	Yes		
		Unclear	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Chen 2012

Study characteristics			
Patient sampling	Prospective cohort study of 105 women with hypertension		
Patient characteristics and setting	Sample size: 105 Gestation at sampling: 28-36 weeks Risk: high		
Index tests			
Target condition and reference standard(s)	IUGR defined as poor fetal growth. Perinatal death.		
Flow and timing			
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns

Chen 2012 (Continued)

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low

DOMAIN 2: Index Test Placental grading

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Unclear		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	High

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Chen 2012a
Study characteristics

Patient sampling	High-risk women
Patient characteristics and setting	Sample size: 113 Gestation at sampling: 28-36 weeks Risk: high Setting: tertiary hospital with an average of 200 or more deliveries per month

Chen 2012a (Continued)

Index tests	Grannum grade III used as threshold		
Target condition and reference standard(s)	Low birthweight defined as below 2500 g, gestational age not taken into account		
Flow and timing			
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test Placental grading			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Unclear		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	High
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Chen 2015

Study characteristics

Patient sampling	Prospective cohort study of patients in a routine obstetric clinic
Patient characteristics and setting	<p>Sample size: 15,122</p> <p>Gestation at sampling: 28 weeks</p> <p>Risk: low (hypertension, diabetes mellitus, placenta praevia, anaemia excluded)</p> <p>Setting: tertiary teaching hospital</p>
Index tests	Placental grading classified using Grannum grading, measured with ultrasound
Target condition and reference standard(s)	Stillbirth between 28 and 41 weeks' gestation
Flow and timing	19,338 women received examinations at the clinics; 17,991 of these were eligible for further analysis; 15,112 of these met the inclusion criteria
Comparative	
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test Placental grading			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		

Chen 2015 (Continued)

Were the reference standard results interpreted without knowledge of the results of the index tests? Unclear

Low
Low
DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all patients receive the same reference standard? Yes

Were all patients included in the analysis? Yes

Low
Chervenak 1984
Study characteristics

Patient sampling Patients referred with a clinical suspicion of IUGR

Patient characteristics and setting **Sample size:** 179
Gestation at sampling: within 15 days of delivery
Risk: high

Index tests EFW calculated using Shepard formula

Target condition and reference standard(s) IUGR defined as birthweight 2 SD below the mean

Flow and timing

Comparative

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled? Yes

Was a case-control design avoided? Yes

Did the study avoid inappropriate exclusions? Yes

Low
High
DOMAIN 2: Index Test EFW

Chervenak 1984 (Continued)

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes	
If a threshold was used, was it pre-specified?	Yes	
	Low	Low
DOMAIN 3: Reference Standard		
Is the reference standards likely to correctly classify the target condition?	Yes	
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear	
	Unclear	Unclear
DOMAIN 4: Flow and Timing		
Was there an appropriate interval between index test and reference standard?	Yes	
Did all patients receive the same reference standard?	Yes	
Were all patients included in the analysis?	Yes	
	Low	

Chew 1976

Study characteristics	
Patient sampling	Patients with high-risk pregnancies
Patient characteristics and setting	Sample size: 43 Gestation at sampling: 30-40 weeks Risk: high (hypertension, poor obstetric history, suspected IUGR) Setting: Kandang Kerbau Hospital, Singapore
Index tests	Urinary oestriol measured as oestradiol-17-beta
Target condition and reference standard(s)	
Flow and timing	Some samples taken after fetal death
Comparative	
Notes	
Methodological quality	

Chew 1976 (Continued)

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Unclear		
		Low	High
DOMAIN 2: Index Test Urinary E3			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	High
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Unclear		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Chitlange 1990

Study characteristics	
Patient sampling	270 pregnant women with single uncomplicated pregnancies
Patient characteristics and setting	Sample size: 270 Gestation at sampling: 31-34 weeks Risk: low (only normal antenatal women were selected) Setting: Nowrisjee Wadia Maternity Hospital

Chitlange 1990 (Continued)

Mode of delivery: 3 emergency caesarean sections due to intra-partum fetal distress

Index tests	Placental grading
Target condition and reference standard(s)	SGA defined as birthweight < 2500 g (combined data of IUGR, BW < 2 kg, and LBW, 2 kg- to 2.49 kg). No adjustments for gestational age
Flow and timing	
Comparative	
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Unclear		
Was a case-control design avoided?	Unclear		
Did the study avoid inappropriate exclusions?	Yes		
		Unclear	Low
DOMAIN 2: Index Test Placental grading			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Unclear		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	High
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		

Chitlange 1990 (Continued)

Low
Christensen 2015
Study characteristics

Patient sampling	Retrospective chart review cohort study from 2000 to 2009 in a single academic centre
Patient characteristics and setting	Sample size: 157 Gestation at sampling: 3rd trimester Risk: high (all pregnancies with SUA) Setting: University of Utah School of Medicine
Index tests	EFW calculated using the Hadlock equation
Target condition and reference standard(s)	SGA defined as birthweight < 2500 g, gestational age was not taken into account
Flow and timing	425 pregnancies identified, 234 of these were excluded (anomalies present in 165, 35 multiple gestations, birth information unavailable in 27, 7 ended in stillbirth).
Comparative	
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear		
If a threshold was used, was it pre-specified?	Yes		
		Unclear	Low
DOMAIN 3: Reference Standard			

Christensen 2015 (Continued)

Is the reference standards likely to correctly classify the target condition?	Yes
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear
	Low
	High
DOMAIN 4: Flow and Timing	
Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
	Low

Elliott 1970

Study characteristics	
Patient sampling	22 pregnancies where all delivered babies with a birthweight lower than 2500g after 37 weeks, part of a series of 104 pregnancies
Patient characteristics and setting	Sample size: 22 Gestation at sampling: 3rd trimester Risk: high (all fetuses weighed under 2500 g) Setting: King George V Memorial Hospital, Sydney
Index tests	Oestriol was measured using either Brown and Coyle's method or the semi-automatic method of Brown. Low was defined as a reduction of more than the SD. Measurements were taken at daily intervals and the values used for diagnosis were the means of the values from the final week before pregnancy
Target condition and reference standard(s)	Stillbirth; intrauterine death after 37 weeks
Flow and timing	
Comparative	
Notes	We used data for the incidence of stillbirth in this SGA cohort. 5 elective caesarean sections were performed for greatly reduced oestriol excretion.
Methodological quality	
Item	Authors' judgement Risk of bias Applicability concerns

Elliott 1970 (Continued)

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?	Yes
Was a case-control design avoided?	Yes
Did the study avoid inappropriate exclusions?	Yes
	Low High

DOMAIN 2: Index Test Urinary E3

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes
If a threshold was used, was it pre-specified?	Yes
	Low High

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes
Were the reference standard results interpreted without knowledge of the results of the index tests?	No
	Low Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
	Low

Estel 1989
Study characteristics

Patient sampling	All women with reduced amniotic fluid measured with ultrasound between 38 and 40 weeks
Patient characteristics and setting	Sample size: 41 Gestation at sampling: 38-40 weeks Risk: high (reduced amniotic fluid)
Index tests	Placental grading measured using ultrasound, grade III used as threshold

Estel 1989 (Continued)

Target condition and reference standard(s)	SGA defined as below the 10th weight percentile		
Flow and timing			
Comparative			
Notes	From translation notes		
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Unclear		
		Low	High
DOMAIN 2: Index Test Placental grading			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Unclear		
		Unclear	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Unclear		
Did all patients receive the same reference standard?	Unclear		
Were all patients included in the analysis?	Yes		
		Unclear	

Fliegner 1979
Study characteristics

Patient sampling	Patients in whom simultaneous serial oestriol and pregnanediol measurements were performed after 30 weeks
Patient characteristics and setting	<p>Sample size: 329</p> <p>Gestation at sampling: after 30 weeks</p> <p>Risk: mixed (unselected population)</p> <p>Setting: Royal Women's Hospital, Melbourne, Australia</p> <p>Mode of delivery: 76 inductions, 42 caesarean sections (22 normal E3, 20 subnormal)</p>
Index tests	Urinary oestriol assays were measured by the method of Brown and colleagues and considered to be subnormal if below a line joining 8 mg/24 hours at 30 weeks' gestation to 12 mg/24 hours at 40 weeks.
Target condition and reference standard(s)	FGR, diagnosed when the infant's weight was less than the 10th percentile for gestational age as seen in patients in the community
Flow and timing	Some pregnancies were terminated if hypertension or pre-eclampsia developed, or if a clinical diagnosis of placental insufficiency was supported by a failure to obtain clear amniotic fluid. Oestriol values were not used to influence treatment.
Comparative	
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test Urinary E3			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			

Fliegner 1979 (Continued)

Is the reference standards likely to correctly classify the target condition?	Yes
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes
	Low
	Low
DOMAIN 4: Flow and Timing	
Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
	Low

Freire 2010

Study characteristics				
Patient sampling		Cross-sectional study of 122 pregnant women who had EFW calculated by ultrasonography up to 7 days before delivery		
Patient characteristics and setting		Sample size: 122 Gestation at sampling: 29-41 weeks Risk: mixed (unselected population, some prior caesareans) Setting: Joao Pessoa, Paraiba, Brazil		
Index tests		Ultrasound EFW, Hadlock formula		
Target condition and reference standard(s)		SGA defined using Alexander curve		
Flow and timing				
Comparative				
Notes				
Methodological quality				
Item	Authors' judgement	Risk of bias	Applicability concerns	
DOMAIN 1: Patient Selection				
Was a consecutive or random sample of patients enrolled?	Yes			
Was a case-control design avoided?	Yes			

Freire 2010 (Continued)

Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Unclear		
		Unclear	Unclear
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Unclear		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Gabbay-Benziv 2016

Study characteristics			
Patient sampling	Retrospective cohort study of all women presenting with sonographic EFW performed within 3 days prior to delivery, July 2007-December 2014		
Patient characteristics and setting	Sample size: 6126 Gestation at sampling: within 3 days of delivery Risk: mixed (malformations and abnormalities, multiple births, stillbirths, missing measurements excluded) Setting: single tertiary university-affiliated medical centre		
Index tests	EFW calculated using Hadlock formula		

Gabbay-Benziv 2016 (Continued)

Target condition and reference standard(s)	SGA defined as an actual birth weight below the 10th percentile for gestational age, using local growth reference values controlled for gestational age and sex		
Flow and timing	6322 women had fetal weight estimation performed within 3 days of delivery, 133 were excluded due to anomalies and abnormalities, 63 were exclude due to lack of measurements.		
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Geerts 2016

Study characteristics

Patient sampling	Prospective study of women who had been referred for umbilical artery RI assessment after 32 weeks of pregnancy and the RI found to be normal between February 11th and October 21st 2013
Patient characteristics and setting	<p>Sample size: 210</p> <p>Gestation at sampling: after 32 weeks</p> <p>Risk: high (all women referred due to: reduced SFH, hypertension, diabetes, previous fetal loss, previous abruption or FGR)</p> <p>Setting: Tygerberg Hospital (a secondary and tertiary referral centre), Cape Town, South Africa, February 11th to October 21st 2013</p>
Index tests	EFW calculated using Hadlock formula. Grade II used as threshold
Target condition and reference standard(s)	SGA defined as birthweight below 10th centile for gestational age
Flow and timing	228 eligible patients were enrolled during the study period, 18 were excluded (3 did not meet criteria, anomalies detected in 10, 3 left before being scanned, 2 withdrew consent)
Comparative	
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test Placental grading			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		

Geerts 2016 (Continued)

If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Gerhard 1986

Study characteristics			
Patient sampling	Prospective study including all women who visited the outpatient department for the first time before 20 weeks of pregnancy		
Patient characteristics and setting	Sample size: 869 Gestation at sampling: 28-40 weeks Risk: mixed (unselected population) Setting: University Women's Hospital, Heidelberg, Germany Mode of delivery: 130 caesarean sections, 678 spontaneous onset of labour, 61 instrumental NICU admission: 122 admissions (20% sensitivity and 91% specificity of oestriol screening for predicting NICU admission)		
Index tests	Serum assay		
Target condition and reference standard(s)	SGA defined as birthweight < 10th percentile for age and sex		
Flow and timing	1140 women were included in the study, 260 were eliminated on account of miscarriages, insufficient examinations, and missed appointments, 11 multiple births were not studied.		
Comparative			

Gerhard 1986 (Continued)

Notes

Test results were examined up until the time of delivery but did not affect management

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test E3			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	No		
		Low	

Gohari 1978
Study characteristics

Gohari 1978 (Continued)

Patient sampling	All patients studies as they either exhibited subnormal uterine growth clinically or had pregnancy complications often associated with IUGR		
Patient characteristics and setting	Sample size: 111 Gestation at sampling: after 30 weeks Risk: high (suspected IUGR) Setting: Department of Obstetrics and Gynaecology, Yale University School of Medicine, New Haven, Connecticut		
Index tests	hPL measured using the Placgest immunodiffusion technique or radioimmunoassay		
Target condition and reference standard(s)	SGA diagnosed if birthweight was in or below the 10th percentile of mean weight for gestation, gestational age was determined by Dubowitz examination		
Flow and timing			
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test hPL			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Unclear
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		

Gohari 1978 (Continued)

	Unclear	Low
DOMAIN 4: Flow and Timing		
Was there an appropriate interval between index test and reference standard?	Yes	
Did all patients receive the same reference standard?	Yes	
Were all patients included in the analysis?	Yes	
	Low	

Granat 1977

Study characteristics	
Patient sampling	29 severely hypertensive patients from a cohort of 373 women who had hPL measurements in the 3rd trimester
Patient characteristics and setting	Sample size: 29 Gestation at sampling: after 30 weeks Risk: high (severe hypertension) Setting: Rothschild Univeristy Hospital, Israel Mode of delivery: 10 deliveries by caesarean section
Index tests	Serum hPL measured using a radioimmunoassay technique, low hPL defined as < 1 SD below the normal means
Target condition and reference standard(s)	SFD defined as those infants whose weights were below the 10th percentile for gestational age, according to Battaglia and colleagues.
Flow and timing	
Comparative	
Notes	2 x 2 table could only be extracted for the severely hypertensive group

Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		

Granat 1977 (Continued)

	Low	High
DOMAIN 2: Index Test hPL		
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes	
If a threshold was used, was it pre-specified?	Yes	
	Low	Low
DOMAIN 3: Reference Standard		
Is the reference standards likely to correctly classify the target condition?	Yes	
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear	
	Unclear	Low
DOMAIN 4: Flow and Timing		
Was there an appropriate interval between index test and reference standard?	Yes	
Did all patients receive the same reference standard?	Yes	
Were all patients included in the analysis?	Yes	
	Low	

Griffin 2015

Study characteristics	
Patient sampling	Women with singleton pregnancies and reduced SFH across 11 sites in the UK and Canada
Patient characteristics and setting	Sample size: 592 Gestation at sampling: 24-37 weeks Risk: high Setting: PELICAN FGR study Mode of delivery: 68.2% spontaneous vaginal delivery, 15% assisted vaginal delivery, 16.7% caesarean section
Index tests	PIGF measured by plasma assay, EFW formula unclear
Target condition and reference standard(s)	SGA defined using customised BW centiles
Flow and timing	9 women excluded from the analysis due to lack of PIGF/outcome/ultrasound data

Griffin 2015 (Continued)

Comparative

Notes	The EFW and PLGF test included AFI measurements but these did not have much of an effect on the overall results
-------	---

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 2: Index Test PLGF			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	No		

Griffin 2015 (Continued)

Unclear
Gupta 2008
Study characteristics

Patient sampling	Retrospective cohort
Patient characteristics and setting	Sample size: 38 Gestation at sampling: 28-32 weeks Risk: high (all patients with severe preterm pre-eclampsia) Mode of delivery: 19 normal vaginal deliveries, 19 caesarean sections
Index tests	EFW, formula unknown
Target condition and reference standard(s)	SGA defined as birthweight below the 10th centile for gestational age
Flow and timing	
Comparative	
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			

Gupta 2008 (Continued)

Is the reference standards likely to correctly classify the target condition?	Yes
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear
	Unclear
	Low
DOMAIN 4: Flow and Timing	
Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
	Low

Hammad 2015

Study characteristics			
Patient sampling	71 women with uncomplicated pregnancies and confirmed gestational dates recruited across 3 sites and randomised to an additional ultrasound group		
Patient characteristics and setting	Sample size: 71 Gestation at sampling: after 30 weeks Risk: low (autoimmune disorders, diabetes, hypertension, history of IUGR, preterm birth, stillbirth, or pre-eclampsia excluded) Mode of delivery: 53 spontaneous vaginal deliveries, 2 operative deliveries, 16 caesarean sections NICU admission: 2 admissions		
Index tests	EFW using Hadlock formula		
Target condition and reference standard(s)	SGA defined as a birthweight < 10th centile using Alexander curves		
Flow and timing	234 women were approached for randomisation, 149 were recruited and 97% of these had follow-up data.		
Comparative			
Notes	The extra ultrasound group was used, this had the latest measurements taken (36 to 37 weeks)		
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns

Hammad 2015 (Continued)

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?	Yes
Was a case-control design avoided?	Yes
Did the study avoid inappropriate exclusions?	Yes
	Low High

DOMAIN 2: Index Test EFW

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes
If a threshold was used, was it pre-specified?	Yes
	Low Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes
Were the reference standard results interpreted without knowledge of the results of the index tests?	No
	Low Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
	Low

Hatfield 2010
Study characteristics

Patient sampling	Retrospective cohort study from 1999 to 2007
Patient characteristics and setting	Sample size: 659 Gestation at sampling: after 26 weeks Risk: high (all patients had elevated hCG levels) Setting: Saddleback Memorial Medical Centre
Index tests	EFW < 10th centile, formula unknown

Hatfield 2010 (Continued)

Target condition and reference standard(s)

SGA defined as below the 10th centile for gestational age

Flow and timing

Comparative

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear		
If a threshold was used, was it pre-specified?	Unclear		
		High	High
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Unclear		
		Low	

Hawkins 2012
Study characteristics

Patient sampling	Retrospective analysis of 2 databases of hypertensive pregnancies
Patient characteristics and setting	<p>Sample size: 1306</p> <p>Gestation at sampling: after 34 weeks (value closest to delivery was used)</p> <p>Risk: high (all hypertensive pregnancies, excluding chronic hypertension and renal disease)</p> <p>Setting: St George Hospital, Australia</p> <p>NICU admissions: 226 of 1880 NICU or SCN transfer</p>
Index tests	Uric acid measured in serum, values used to determine hyperuricaemia were corrected for gestational age. Elevated uric acid defined as being 1 SD above the gestation-specific mean.
Target condition and reference standard(s)	SGA defined as a birthweight below the 10th centile
Flow and timing	n =1880 after databases were combined and duplicates were excluded.
Comparative	
Notes	The presence of hyperuricaemia was not a component in the diagnosis of pre-eclampsia nor in the decision to deliver any pregnancy. Authors were contacted for data.

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test Uric acid			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Unclear
DOMAIN 3: Reference Standard			

Hawkins 2012 (Continued)

Is the reference standards likely to correctly classify the target condition?	Yes
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes
	Low
	Low
DOMAIN 4: Flow and Timing	
Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
	Low

Hendrix 2000

Study characteristics			
Patient sampling	Randomised clinical trial of singleton pregnancies with reliable gestational age of 37 weeks or more.		
Patient characteristics and setting	Sample size: 367 Gestation at sampling: > 37 weeks Risk: mixed (unselected population) Setting: Spartanburg Regional Medical Centre, South Carolina & Medical College of Georgia, Augusta, Georgia		
Index tests	Ultrasound EFW calculated using Hadlock formula		
Target condition and reference standard(s)	Birthweight < 2500 g, not adjusted for gestational age		
Flow and timing	758 of 2541 eligible patients were randomised to groups for the trial, 367 assigned to the sonographic estimate group		
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		

Hendrix 2000 (Continued)

Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	High
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	No		
		Low	

Howell 1985

Study characteristics			
Patient sampling	501 unselected women with a singleton pregnancy		
Patient characteristics and setting	Sample size: 501 Gestation at sampling: 36-41 weeks Risk: mixed (unselected population; 45 patients with pre-eclampsia, 2 with hypertension, 4 with antepartum haemorrhage, 2 diabetics) Setting: St Bartholomew's Hospital Medical College & London Hospital Medical College		
Index tests	hPL measured by serum radioimmunoassay, 10th centile used as threshold		

Howell 1985 (Continued)

Target condition and reference standard(s)

Low birthweight defined as below the 10th centile for the whole population

Flow and timing

Comparative

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test hPL			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Jauniaux 1996
Study characteristics

Patient sampling	Women with abnormal uterine artery Doppler features and/or an increased pulsatility index
Patient characteristics and setting	Sample size: 41 Gestation at sampling: 25-28 weeks Risk: high (abnormal Doppler, increased pulsatility)
Index tests	Serum levels of uric acid measured by an enzymatic method using uricase, > 4.0 mg/dL considered to be elevated
Target condition and reference standard(s)	IUGR defined as birthweight < 10th centile for local standards
Flow and timing	Measurements taken from 41 women who consented
Comparative	
Notes	Placental histopathological examination revealed extended vascular lesions in 12 complicated cases. These lesions were often combined in cases complicated by PIH and IUGR and by placental abruption.

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Unclear
DOMAIN 2: Index Test Uric acid			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		

Jauniaux 1996 (Continued)

Unclear Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes

Low

Kazzi 1983a
Study characteristics

Patient sampling	Prospective observational study of placental grade and fetal maturity
Patient characteristics and setting	Sample size: 109 Gestation at sampling: within 7 days of delivery Risk: high (pregnancies with birthweight < 2700 g) Setting: Cleveland Metropolitan General Hospital
Index tests	Placental grading determined according to Grannum classification, grade III used as threshold
Target condition and reference standard(s)	SGA defined as a birthweight less than the 10th percentile for gestational age
Flow and timing	109 pregnancies from a cohort of 224 women with birthweight < 2700 g who were examined sonographically within 7 days of delivery
Comparative	
Notes	2700 g was chosen as it was the maximum weight at which infants could be considered SGA

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		

Kazzi 1983a (Continued)

	Low	High
DOMAIN 2: Index Test Placental grading		
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes	
If a threshold was used, was it pre-specified?	Yes	
	Low	Low
DOMAIN 3: Reference Standard		
Is the reference standards likely to correctly classify the target condition?	Yes	
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear	
	Unclear	Low
DOMAIN 4: Flow and Timing		
Was there an appropriate interval between index test and reference standard?	Yes	
Did all patients receive the same reference standard?	Yes	
Were all patients included in the analysis?	No	
	High	

Kienast 2016

Study characteristics	
Patient sampling	Prospective cohort of women who attended for routine obstetrical care between April and December 2010
Patient characteristics and setting	Sample size: 346 Gestation at sampling: 28-32 weeks Risk: mixed (unselected population although there is a high incidence of PE in the highlands of Ecuador) Setting: Hospital Isidro Ayora, Quito, Ecuador
Index tests	Plasma sFlt-1 and PlGF measured using a commercial Roche Elys System. Threshold was not pre-specified; the optimal cutoff in terms of maximising both sensitivity and specificity was reported.
Target condition and reference standard(s)	FGR was defined as a birthweight less than the 10th percentile of a reference group

Kienast 2016 (Continued)

Flow and timing

Comparative

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test PIGF			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	No		
		High	Unclear
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Klebe 1990
Study characteristics

Klebe 1990 (Continued)

Patient sampling	Patients with renal transplants from 4 different centres		
Patient characteristics and setting	Sample size: 13 Gestation at sampling: after 32 weeks Risk: high Setting: 4 centres in Denmark		
Index tests	hPL measured in serum, 5th centile used as threshold		
Target condition and reference standard(s)	SGA defined as outside of normal reference area (below 5th percentile); unclear if this threshold was prespecified		
Flow and timing			
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Unclear
DOMAIN 2: Index Test hPL			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Unclear		
		Unclear	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	High
DOMAIN 4: Flow and Timing			

Klebe 1990 (Continued)

Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
Low	

Kunz 1976

Study characteristics			
Patient sampling	Patients presently hospitalised because of, or with history of, suspected placental insufficiency		
Patient characteristics and setting	Sample size: 83 Gestation at sampling: third trimester Risk: high (suspected placental insufficiency or a case history)		
Index tests	hPL and oestriol determinations both performed in serum. 10 measurements of oestriol were taken but only the last 2 values were used for diagnosis. Biochemical parameters were considered to be pathological if they were below 95% of the normal values, with 2 values being outside this normal range or with 3 or more values being continuously below the 95% normal range.		
Target condition and reference standard(s)	SGA defined if birthweight was below the 10th percentile on the Lubchenko curve		
Flow and timing	Twin pregnancies and those with doubtful duration were excluded		
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test hPL			

Kunz 1976 (Continued)

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	High
DOMAIN 2: Index Test Urinary E3			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	High
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Laurin 1987

Study characteristics	
Patient sampling	All singleton births without major malformations in Malmö in 1983
Patient characteristics and setting	Sample size: 2205 Gestation at sampling: 32 weeks Risk: mixed (94% of the population, malformations and multiple births excluded) Setting: Malmö General Hospital, Sweden
Index tests	EFW using formula of Eik-Nes, IUGR defined as a -15% deviation for gestational age

Laurin 1987 (Continued)

Target condition and reference standard(s)	SGA defined as a birthweight - 2 SD for gestational age		
Flow and timing	2322 singleton births in Malmö in 1983, 2205 of which participated in at least 2 ultrasound examinations. 137 of these were lost to follow-up.		
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Leader 1980
Study characteristics

Patient sampling	Patients admitted antenatally with hypertension, suspicion of IU-GR or postmaturity.
Patient characteristics and setting	Sample size: 135 Gestation at sampling: 30-42 weeks Risk: high (hypertension, suspected IUGR, postmaturity) Setting: Groote Schuur Hospital, Cape Town
Index tests	hPL estimations done by radioimmunoassay using the Amersham hPL kit, levels were known but not used to influence treatment
Target condition and reference standard(s)	Stillbirth after 30 weeks
Flow and timing	
Comparative	
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test hPL			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	No		

Leader 1980 (Continued)

	Low	Low
DOMAIN 4: Flow and Timing		
Was there an appropriate interval between index test and reference standard?	Yes	
Did all patients receive the same reference standard?	Yes	
Were all patients included in the analysis?	Yes	
	Low	

Lenstrup 1982

Study characteristics			
Patient sampling	Consecutive series of patients examined with CTG in the 35th-36th week of pregnancy		
Patient characteristics and setting	Sample size: 88 Gestation at sampling: 35-36 weeks Risk: mixed (unselected population) Setting: Herlev Hospital, Denmark NICU admission: 4 admissions, 3 of these from the reduced variability group		
Index tests	Oestriol and hPL analyses were performed on blood, threshold unclear ('low levels')		
Target condition and reference standard(s)	SFD defined as birthweight below 10th centile for Danish children		
Flow and timing			
Comparative			
Notes	1 patient was hospitalised after CTG because of a low plasma oestriol level and low EFW. Some incorrect data in Table 5; 2 normal E3/hPL reduced variability and normal for date assumed to be 6 instead.		
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		

Lenstrup 1982 (Continued)

Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test hPL			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Unclear		
		Unclear	Unclear
DOMAIN 2: Index Test E3			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Unclear		
		Unclear	Unclear
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	No		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Lilford 1983

Study characteristics			
Patient sampling	Unselected women with a singleton pregnancy		
Patient characteristics and setting	Sample size: 527 Gestation at sampling: 36-40 weeks Risk: mixed (unselected population; 27 women with pre-eclampsia, 7 with antepartum haemorrhage)		

Lilford 1983 (Continued)

Index tests	Serum assay for hPL, 10th centile used as threshold. The effect of changing centiles on the sensitivity and specificity was looked at.		
Target condition and reference standard(s)	FGR defined as a birthweight < 2740 g; the 10th centile for birthweight in the sample studied		
Flow and timing	Stillbirths were excluded from the calculations as they all occurred before the onset of labour and birthweight would be unrelated to the remainder of the group.		
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test hPL			
Were the index test results interpreted without knowledge of the results of the reference standard?	No		
If a threshold was used, was it pre-specified?	No		
		High	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	High
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		

Lilford 1983 (Continued)

Low

MacLeod 2013
Study characteristics

Patient sampling	Prospective study of term, cephalic, singleton pregnancies
Patient characteristics and setting	Sample size: 90 Gestation at sampling: term Risk: mixed (unselected population) Setting: Mbarara Regional Referral Hospital, Uganda
Index tests	Ultrasound EFW, Hadlock formula
Target condition and reference standard(s)	Low birthweight defined as < 2500 g, not adjusted for gestational age
Flow and timing	
Comparative	
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		

MacLeod 2013 (Continued)

Were the reference standard results interpreted without knowledge of the results of the index tests? Unclear

Unclear

High

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all patients receive the same reference standard? Yes

Were all patients included in the analysis? Yes

Low

Mahran 1988
Study characteristics

Patient sampling Patients selected at random from obstetric population who attended the Ultrasound Unit in 1983, only those who had sure dates confirmed by early ultrasonic examination.

Patient characteristics and setting **Sample size:** 828
Gestation at sampling: after 24 weeks
Risk: mixed (unselected population)

Index tests Ultrasound EFW, below 2 SD of normal parameters

Target condition and reference standard(s) SGA defined as birthweight below the 10th centile, evaluated by tables constructed by Thomson and colleagues

Flow and timing

Comparative

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled? Yes

Was a case-control design avoided? Yes

Did the study avoid inappropriate exclusions? Yes

Low

Low

Mahran 1988 (Continued)

DOMAIN 2: Index Test EFW

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes
If a threshold was used, was it pre-specified?	Yes
	Low
	Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear
	Low
	Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
	Low

Marin 1979
Study characteristics

Patient sampling	Antenatal patients with amniotic fluid samples
Patient characteristics and setting	Sample size: 47 Gestation at sampling: after 33 weeks Risk: mixed
Index tests	Serum hPL assayed using radioimmunoassay kits, 5th centile used as threshold
Target condition and reference standard(s)	10th centile for gestational age used to define IUGR
Flow and timing	
Comparative	
Notes	

Methodological quality

Marin 1979 (Continued)

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test hPL			
Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear		
If a threshold was used, was it pre-specified?	Unclear		
		Unclear	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Unclear		
		Unclear	

McKenna 2005

Study characteristics	
Patient sampling	Prospective cohort of singleton pregnancies with known gestational age
Patient characteristics and setting	Sample size: 1802 Gestation at sampling: 36 weeks Risk: low (known maternal medical problems, obstetric complications in present or prior pregnancy, fetal abnormalities excluded)

McKenna 2005 (Continued)

Setting: Royal Jubilee Maternity Hospital, Belfast

Mode of delivery: 1190 normal vaginal deliveries, 71 inductions

NICU admission: 33 admissions, 1 of these from the grade III group

Index tests	Placental maturity was determined using the Grannum classification, grade III used as threshold
Target condition and reference standard(s)	Low birthweight defined as below the 10th centile at birth
Flow and timing	
Comparative	
Notes	Some pregnancies were induced for suspected fetal compromise, grade III placentas helped to identify these pregnancies

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test Placental grading			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	No		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		

McKenna 2005 (Continued)

Were all patients included in the analysis? Yes

Low

Miller 1988
Study characteristics

Patient sampling Patients who delivered beyond 37 weeks with known dates, early examination, confirmatory ultrasound, and who delivered within 7 days of study were selected from all referrals

Patient characteristics and setting
Sample size: 246
Gestation at sampling: 36-38 weeks
Risk: mixed (unselected apart from no preterm delivery)
Setting: Louisiana State University Medical Centre

Index tests Placental grade established from the most mature view of the placenta and assigned in accordance with established criteria, grade III used as threshold

Target condition and reference standard(s) SGA classified as birthweight below 10th centile

Flow and timing

Comparative

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?	Yes		
--	-----	--	--

Was a case-control design avoided?	Yes		
------------------------------------	-----	--	--

Did the study avoid inappropriate exclusions?	Yes		
---	-----	--	--

Low

Low

DOMAIN 2: Index Test Placental grading

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
---	-----	--	--

If a threshold was used, was it pre-specified?	Yes		
--	-----	--	--

Low

Low

Miller 1988 (Continued)

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear
	Low
	Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
	Low

Molvarec 2013
Study characteristics

Patient sampling	Study group selected from groups of hypertensive women based on availability of Doppler ultrasound results
Patient characteristics and setting	Sample size: 89 Gestation at sampling: mean 32 weeks Risk: high (all women had hypertensive disorders of pregnancy) Mode of delivery: 78 caesarean sections
Index tests	PIGF measured using the Alere Triage test, 12 pg/mL used as threshold
Target condition and reference standard(s)	SGA defined as birthweight below the 10th percentile for gestational age and sex according to a Hungarian birthweight percentile table. All neonates with SGA had an asymmetric size, indicating that they had intrauterine growth restriction and were not constitutionally small.
Flow and timing	
Comparative	
Notes	PIGF values were not an indicator for early delivery

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

Molvarec 2013 *(Continued)*
DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Unclear	High

DOMAIN 2: Index Test PIGF

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Low	Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Montan 1986
Study characteristics

Patient sampling	Prospective study of 645 consecutive pregnancies over a 4-month period
Patient characteristics and setting	Sample size: 307 Gestation at sampling: 37 weeks Risk: mixed (unselected pregnancies, 40 admitted due to pregnancy induced hypertension) Setting: University Hospital of Lund

Montan 1986 (Continued)

Index tests	Placental grading was made according to the classification of Grannum and colleagues, grade III used as threshold		
Target condition and reference standard(s)	SGA not defined		
Flow and timing	654 women were recruited for the study, of which 621 had a scan in weeks 32 to 33. 146 gave birth before weeks 38 to 29, and 96 objected to further examinations.		
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test All tests			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 2: Index Test Placental grading			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Low	Unclear

Montan 1986 (Continued)

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
Low	

Nice 2016
Study characteristics

Patient sampling	Blood collected from healthy pregnancies, women with reduced fetal movements, or a suspected SGA fetus after 28 weeks of gestation
Patient characteristics and setting	Sample size: 77 Gestation at sampling: after 28 weeks Risk: mixed Setting: St. Mary's Hospital, Manchester UK NICU admission: 12
Index tests	hPL measured by ELISA (threshold 0.8 MoM), PLGF by Alere Triage, ELISA, and Roche automated immunoassay (threshold 12 pg/mL)
Target condition and reference standard(s)	SGA defined as birthweight < 10th centile
Flow and timing	
Comparative	
Notes	Authors contacted for data

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Unclear	Low

Nice 2016 (Continued)

DOMAIN 2: Index Test hPL

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low

DOMAIN 2: Index Test PlGF

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Nielsen 1985
Study characteristics

Patient sampling	All births that took place in a geographically well-defined area in the course of 1 year that met the inclusion criteria (expected date of delivery, all measurements, measurements not known to the physicians)
Patient characteristics and setting	Sample size: 1018 Gestation at sampling: 26th and 35th Risk: mixed (unselected pregnancies)

Nielsen 1985 (Continued)

Index tests	Oestriol measured with a radioimmunoassay kit, 2.5th centile used as a threshold		
Target condition and reference standard(s)	SGA defined as bodyweight below the 10th centile, referring to the Danish National Board of Health		
Flow and timing	1702 births took place; 1660 of these were registered on a form (8 outside hospital and 34 not registered, these were omitted). 13 twin pregnancies, 153 with uncertain date of expected delivery, 213 without measurements, 263 where monitoring was prescribed were all excluded.		
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test E3			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		

Nielsen 1985 (Continued)

Were all patients included in the analysis? Yes

Low

Nisbet 1982
Study characteristics

Patient sampling	Patients with singleton pregnancies attending routine antenatal clinic
Patient characteristics and setting	Sample size: 166 Gestation at sampling: 3rd trimester (34.3 +/- 3.1) Risk: high (all considered clinically to have small for dates fetuses) Setting: Aberdeen Maternity Hospital
Index tests	Measurements taken from serum or plasma, oestriol was only measured when there was sufficient sample volume. - 2 SD used as threshold
Target condition and reference standard(s)	IUGR defined as birthweight below the 10th centile according to gestation, sex, and parity.
Flow and timing	166 patients were included in the study, only 103 could be assayed for oestriol (25 SGA and 38 AGA not included)
Comparative	
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		High	Low
DOMAIN 2: Index Test hPL			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		

Nisbet 1982 (Continued)

		Low	Low
DOMAIN 2: Index Test E3			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	No		
		Unclear	

Oats 1979

Study characteristics	
Patient sampling	Patients in a 7-year period
Patient characteristics and setting	Sample size: 19,119 Gestation at sampling: from 30 weeks Risk: mixed (unselected population) Setting: Mercy Maternity Hospital, Melbourne
Index tests	Urinary oestriol excretion was measured by the method of Brown and colleagues, which accounts for gestation. Low E3 patients are from 2 groups, persistently low (2 or more consecutive low values) and transiently low (only 1 low value out of the total number of measurements). Values were regarded as low when below a line joining 8 mg/24 hours at 30 weeks and 12 mg/24 hours at 40 + weeks.
Target condition and reference standard(s)	SFD defined as birthweight < 10th centile for gestational age in the community. Stillbirths defined according to the criteria used in the Commonwealth of Australia.

Oats 1979 (Continued)

Flow and timing

Comparative

Notes

The test was performed as a routine at 30 and 36 weeks and at other times when indicated by complications of pregnancy or by poor past obstetrical history. Study looks at 400 patients with low oestriol in a subsequent pregnancy but also gives results for the whole study population so these were used.

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test Urinary E3			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	High
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Obiekwe 1983
Study characteristics

Patient sampling	Unselected women with singleton pregnancies
Patient characteristics and setting	Sample size: 522 Gestation at sampling: 36-40 weeks Risk: mixed (unselected population; 27 women with pre-eclampsia, 7 antepartum haemorrhage)
Index tests	hPL measured in serum by immunoassay, 10th centile used as threshold.
Target condition and reference standard(s)	IUGR defined as birthweight below 10th centile of study population (2740 g)
Flow and timing	Stillbirths (n = 5) excluded from original cohort
Comparative	
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test hPL			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Unclear		
		Unclear	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	High

Obiekwe 1983 (Continued)

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
Low	

Odendaal 1981
Study characteristics

Patient sampling	Patients with positive stress tests
Patient characteristics and setting	Sample size: 148 Gestation at sampling: 25-43 weeks Risk: high (positive stress tests, various indications) Setting: Tygerberg Hospital, South Africa
Index tests	hPL and oestriol immunoassay kit used to measure both, 10th centile used to define low levels. Centiles derived from curves used by Tygerberg hospital derived from 432 patients.
Target condition and reference standard(s)	SGA defined as birthweight below the 10th percentile for the specific duration of pregnancy, gestational age estimated using a Dubowitz score and weight charts used to assess growth
Flow and timing	Serum oestriol and hPL studies were not performed on all patients as these tests were not initially available; intrauterine deaths were also not included in the estimations of IUGR.
Comparative	
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High

Odendaal 1981 *(Continued)*
DOMAIN 2: Index Test hPL

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low

DOMAIN 2: Index Test E3

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low

DOMAIN 2: Index Test Urinary E3

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Yes		
		Low	Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	No		
		Unclear	

Odendaal 1997
Study characteristics

Patient sampling	Prospective study of women with severe pre-eclampsia.
Patient characteristics and setting	Sample size: 196

Odendaal 1997 (Continued)

Gestation at sampling: the week before delivery (blood taken twice a week from admission until delivery, last sample used)

Risk: high (all women with severe pre-eclampsia, all were delivered once 34 weeks was reached)

Setting: Tygerberg Hospital, South Africa - a tertiary hospital to which many patients with severe pre-eclampsia are referred

Index tests	Plasma uric acid was assessed by the automated urokinase method on a Technician SMAC machine, high levels denoted as 1 SD above the mean
Target condition and reference standard(s)	Tygerberg Hospital growth curves were used to assess whether newborns were SGA
Flow and timing	229 women were included but 33 delivered before 28 weeks and so their weights for gestational age could not be assessed.
Comparative	
Notes	Patients were delivered once a gestational age of 34 weeks has been reached or when maternal reasons or abnormal fetal heart rate patterns were an indication for earlier delivery.

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test Uric acid			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low

Odendaal 1997 (Continued)

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	No
Low	

Ott 1984
Study characteristics

Patient sampling	All patients undergoing ultrasonic examination within 72 hours before delivery
Patient characteristics and setting	Sample size: 595 Gestation at sampling: after 30 weeks Risk: mixed (the study population was composed of both high- and low-risk patients) Setting: St Mary's Health Centre, St. Louis, Missouri
Index tests	Ultrasound EFW calculated using the formula of Shepard and colleagues
Target condition and reference standard(s)	SGA classified based on birthweight from gestational age percentile obtained from the normogram of Altman and Coles
Flow and timing	
Comparative	
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low

DOMAIN 2: Index Test EFW

Ott 1984 (Continued)

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes	
If a threshold was used, was it pre-specified?	Yes	
	Low	Low
DOMAIN 3: Reference Standard		
Is the reference standards likely to correctly classify the target condition?	Yes	
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear	
	Low	Low
DOMAIN 4: Flow and Timing		
Was there an appropriate interval between index test and reference standard?	Yes	
Did all patients receive the same reference standard?	Yes	
Were all patients included in the analysis?	Yes	
	Low	

Palo 1987

Study characteristics		
Patient sampling	Admissions due to short SFH measurement	
Patient characteristics and setting	Sample size: 90 Gestation at sampling: 1 week before delivery (28-40 weeks) Risk: high (reduced SFH) Setting: University Central Hospital, Turku	
Index tests	Oestriol measured by radioimmunoassay according to Kaihola, - 2 SD used as threshold	
Target condition and reference standard(s)	IUGR defined as below 10th percentile by weight	
Flow and timing	The study combined prospective and retrospective data; prospective population derived from mothers with reduced SFH plus 'controls' (women without IUGR after screening the whole cohort), retrospective part was mothers with low E3 only so was not included	
Comparative		
Notes		

Palo 1987 (Continued)

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test E3			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Palo 1989
Study characteristics

Patient sampling	Prospective study, women admitted based on clinical suspicion of poor fetal growth.
Patient characteristics and setting	Sample size: 186 Gestation at sampling: 31-42 weeks (mean 38.6)

Palo 1989 (Continued)

Risk: high

Setting: Maternity Clinic of University Central Hospital, Turku

Index tests	EFW calculations were performed using the calculations of Eik-Nes and colleagues.
Target condition and reference standard(s)	SGA defined as birthweight below the 10th centile
Flow and timing	206 women admitted, 20 excluded because exact BPD or AD data were not available
Comparative	
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		

Palo 1989 (Continued)

Were all patients included in the analysis? Yes

Low

Patterson 1983
Study characteristics

Patient sampling Retrospective selection of patients that demonstrated a Grade II or III placenta (Spanish-surnamed patients only)

Patient characteristics and setting
Sample size: 398
Gestation at sampling: 26-39 weeks
Risk: high
Setting: Medical Center Hospital, San Antonio, Texas

Index tests Placental grade assessed according to the criteria of Grannum and associates

Target condition and reference standard(s) SGA defined as birthweight below the 10th centile for gestational age.

Flow and timing 431 patients were identified, 398 of these underwent delivery at Medical Center Hospital and were included in the study.

Comparative

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?	No		
--	----	--	--

Was a case-control design avoided?	Yes		
------------------------------------	-----	--	--

Did the study avoid inappropriate exclusions?	Yes		
---	-----	--	--

High

High

DOMAIN 2: Index Test Placental grading

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
---	-----	--	--

If a threshold was used, was it pre-specified?	Yes		
--	-----	--	--

Low

Low

Patterson 1983 (Continued)

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear
	Unclear
	Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
	Low

Redman 1976
Study characteristics

Patient sampling	All patients with hypertension. Patients do not comprise a single cohort since new patients entered the study up to 32 weeks of gestation and premature delivery removed other patients from the later periods of the study.
Patient characteristics and setting	Sample size: 281 Gestation at sampling: 36 weeks Risk: high
Index tests	Plasma urate was assayed by the routine automated hydroxylamine method
Target condition and reference standard(s)	Stillbirths defined as fetuses born dead after the 24th week of gestation
Flow and timing	Hypertension was confirmed in 332 pregnant patients, 238 of which were participating in a trial of antihypertensive treatment. Some women delivered early; hyperuricaemia was never used as an indicator. 281 women had uric acid measurements at 36 weeks.
Comparative	
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

Redman 1976 (Continued)

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?	No		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High

DOMAIN 2: Index Test Uric acid

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Roma 2015
Study characteristics

Patient sampling	Women were non-selectively enrolled into a randomised trial. Some exclusions based on medical history and history of FGR/PE/stillbirth.
Patient characteristics and setting	Sample size: 1115 Gestation at sampling: 36 weeks Risk: mixed Mode of delivery: 400 inductions, 86 instrumental deliveries and 47 caesarean sections (both for non reassuring fetal status)

Roma 2015 (Continued)

NICU admission: 1 admission

Index tests	EFW calculated using Hadlock formula
Target condition and reference standard(s)	SGA defined as a birthweight below the 10th centile
Flow and timing	1314 women were assigned to US at 36 weeks, 1115 were analysed (1 stillbirth, 13 preterm births, 185 lost to follow up)
Comparative	
Notes	Suspected cases of SGA prompted weekly monitoring and elective induction at 37 + 1 weeks if UA-PI values were abnormal (> 95th centile); otherwise, monitoring was carried out every 2 weeks and delivery was induced at 40 + 1 weeks. Data for US at 36 weeks were used as the last test before delivery.

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		

Roma 2015 *(Continued)*

Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
Low	

Sagen 1984

Study characteristics	
Patient sampling	Women with severe pre-eclampsia
Patient characteristics and setting	Sample size: 74 Gestation at sampling: 1-3 days before birth Risk: high Mode of delivery: 54 caesarean sections, 15 instrumental deliveries, remaining 5 were IUDs
Index tests	Oestriol measured in plasma using a radioimmunoassay, hPL measured in plasma using an immunoassay. Threshold defined as the 10th centile from a reference group of 40 healthy pregnant women. Measurements taken twice a day but only the last measurement before delivery was used.
Target condition and reference standard(s)	Low birthweight defined as under the 10th centile, weight of the newborn related to a centile scale based on 416,756 liveborn infants without congenital malformations and after normal singleton pregnancy in Norway from 1967-1977
Flow and timing	
Comparative	
Notes	Indication for delivery was based on case history, clinical findings, and results of hormonal, biochemical, ultrasonographic, and cardiotocographic tests

Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High

DOMAIN 2: Index Test hPL

Sagen 1984 (Continued)

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 2: Index Test E3			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	No		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Sekar 2016
Study characteristics

Patient sampling	Prospective study of all women booked for induction of labour or elective caesarean section during February to December 2013
Patient characteristics and setting	Sample size: 150 Gestation at sampling: 1 week prior to delivery Risk: mixed
Index tests	EFW calculated using Hadlock formula
Target condition and reference standard(s)	SGA defined as birthweight below the 10th centile for gestational age. Doctors and women were both aware of EFWs

Sekar 2016 (Continued)

Flow and timing

Comparative

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	No		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Shawkat 2015
Study characteristics

Shawkat 2015 (Continued)

Patient sampling	High-risk cohort of women < 35 weeks gestation		
Patient characteristics and setting	Sample size: 261 Gestation at sampling: test nearest delivery Risk: high (suspected FGR, superimposed PE)		
Index tests	PIGF measured using Alere Triage; threshold used was very low (< 12 pg/mL)		
Target condition and reference standard(s)	SGA defined as < 10th centile, fetal death		
Flow and timing			
Comparative			
Notes	Surveillance was adjusted according to test results		
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test PIGF			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	No		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		

Shawkat 2015 (Continued)

Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
Low	

Siebert 1974

Study characteristics	
Patient sampling	A group of 166 pregnant women
Patient characteristics and setting	Sample size: 67 Gestation at sampling: 1-3 days before birth Risk: low
Index tests	Serum assay, some multiple measurements and samples analysed in duplicate
Target condition and reference standard(s)	SGA, intrauterine death
Flow and timing	166 women were recruited but data could only be extracted for the 67 normal pregnancies where hPL data were given for all SGA babies, and 20 diabetic patients for intrauterine death
Comparative	
Notes	Test results facilitated management - 2 caesareans were performed in the diabetic pregnancy group (excluded) due to falling values

Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test hPL			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Unclear		

Siebert 1974 (Continued)

		Unclear	Unclear
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	No		
		High	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Skovron 1991

Study characteristics			
Patient sampling	Patients with a singleton gestation who had ultrasound for determination of fetal size between 26 and 34 weeks, 1985-1987. Gestational diabetes, placenta praevia, premature labour excluded.		
Patient characteristics and setting	Sample size: 768 Gestation at sampling: 26-34 weeks Risk: mixed		
Index tests	Ultrasound EFW calculated using the formula of Shepard and colleagues.		
Target condition and reference standard(s)	SGA defined as below the 10th centile of birthweight for gestational age and sex using the nomogram developed by Brenner and colleagues.		
Flow and timing	996 patients recruited, 768 met the inclusion criteria (37 excluded because of incomplete ultrasound data)		
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns

Skovron 1991 (Continued)

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low

DOMAIN 2: Index Test EFW

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	No		
		Low	

Sovio 2015
Study characteristics

Patient sampling	Prospective cohort of nulliparous women attending for their dating ultrasound scan between January 2008 and July 2012.
Patient characteristics and setting	Sample size: 3977 Gestation at sampling: 28 or 36 weeks (last scan before delivery) Risk: mixed (the only exclusion criterion was multiple pregnancy) Mode of delivery: 1924 normal vaginal deliveries, 949 instrumental, 1089 caesarean sections (data missing for mode of delivery for 15 births)

Sovio 2015 (Continued)

NICU admission: 229 admissions

Index tests	Ultrasound EFW calculated using Hadlock equations
Target condition and reference standard(s)	SGA defined as a birthweight of less than the 10th percentile for sex and gestational age, calculated from a UK reference
Flow and timing	8028 women eligible, 4512 enrolled, 3977 attended all third trimester scans and delivered a liveborn infant after 26 weeks.
Comparative	
Notes	We used universal ultrasonography as these were all single measurements (the last scan before birth) and the data for selective ultrasonography includes women who did not have a scan. Results of routine clinical scans were reported but results of research scans were masked.

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		

Sovio 2015 (Continued)

Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	No
Unclear	

Spernol 1989
Study characteristics

Patient sampling	Women screened prospectively; 30 with suspected IUGR, 5 with infants below the 10th centile for weight who were not suspected SGA, 75 low-risk pregnancies.
Patient characteristics and setting	Sample size: 110 Gestation at sampling: unclear, likely to be 3rd trimester Risk: mixed
Index tests	
Target condition and reference standard(s)	SGA defined as birthweight below the 10th centile
Flow and timing	
Comparative	
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Unclear		
Did the study avoid inappropriate exclusions?	Unclear		
		Unclear	Unclear
DOMAIN 2: Index Test hPL			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low

Spornol 1989 (Continued)

DOMAIN 2: Index Test E3

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low

DOMAIN 3: Reference Standard

Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Steiner 1991
Study characteristics

Patient sampling	113 pregnant women with suspected IUGR
Patient characteristics and setting	Sample size: 113 Gestation at sampling: 28-40 weeks Risk: high (all suspected IUGR)
Index tests	Serum assay for hPL, urinary assay for E3; mean value from 3 measurements was used for diagnosis. Threshold unclear.
Target condition and reference standard(s)	IUGR defined as a birthweight below the 10th centile
Flow and timing	From translation notes - no details
Comparative	
Notes	

Methodological quality

Steiner 1991 (Continued)

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test hPL			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Unclear		
		Unclear	Unclear
DOMAIN 2: Index Test Urinary E3			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Unclear		
		Unclear	Unclear
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Unclear		
Did all patients receive the same reference standard?	Unclear		
Were all patients included in the analysis?	Unclear		
		Unclear	

Takeuchi 1985
Study characteristics

Patient sampling	Prospective cohort
Patient characteristics and setting	Sample size: 210 Gestation at sampling: within 7 days of delivery (36-41 weeks) Risk: mixed Mode of delivery: 20 out of 60 'light-for-dates' fetuses were delivered by CS, 8 for fetal distress
Index tests	
Target condition and reference standard(s)	Mean BW of 'light-for-dates' fetuses was 2186 +/- 452 g so was likely adjusted for gestation
Flow and timing	36-41 weeks' gestation was chosen as it is closest to delivery but not all women had measurements at this time.
Comparative	
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Unclear		
		Unclear	Low
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		

Takeuchi 1985 (Continued)

Unclear Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all patients receive the same reference standard? Yes

Were all patients included in the analysis? No

Unclear

Trudinger 1979
Study characteristics

Patient sampling Consecutive series of inpatients studied within 14 days of delivery where the fetus was considered to be at high risk

Patient characteristics and setting **Sample size:** 59
Gestation at sampling: after 33 weeks
Risk: high (31 hypertensive, 20 suspected FGR, 2 antepartum haemorrhage, 3 previous stillbirth, 1 diabetes mellitus, 3 other)

Index tests hPL measured using radioimmunoassay, 10th centile for a pregnancy of the same maturity used as a threshold

Target condition and reference standard(s) The clinician was aware of the E3 and ultrasound results but not hPL. SGA below 10th centile for gestational age.

Flow and timing

Comparative

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled? Yes

Was a case-control design avoided? Yes

Did the study avoid inappropriate exclusions? Yes

Low High

DOMAIN 2: Index Test hPL

Trudinger 1979 *(Continued)*

Were the index test results interpreted without knowledge of the results of the reference standard?	Yes	
If a threshold was used, was it pre-specified?	Yes	
	Low	Low
DOMAIN 3: Reference Standard		
Is the reference standards likely to correctly classify the target condition?	Yes	
Were the reference standard results interpreted without knowledge of the results of the index tests?	No	
	Unclear	Low
DOMAIN 4: Flow and Timing		
Was there an appropriate interval between index test and reference standard?	Yes	
Did all patients receive the same reference standard?	Yes	
Were all patients included in the analysis?	Yes	
	Low	

Turitz 2014

Study characteristics		
Patient sampling	Retrospective cohort of all singleton pregnancies presenting for at least 1 growth ultrasound between 26 and 36 weeks gestational age, January 2008 and December 2011	
Patient characteristics and setting	Sample size: 10,642 Gestation at sampling: 26-36 weeks Risk: mixed (all pregnancies; only exclusions were fetal anomalies, multiple pregnancies, twins with 1 fetal loss)	
Index tests	EFW centiles calculated using Hadlock formula C	
Target condition and reference standard(s)	SGA defined as birthweight below the 10th centile using the Alexander curve	
Flow and timing		
Comparative		
Notes	Fetuses with growth restriction were delivered at 37 weeks, or sooner as indicated for abnormal fetal testing or Doppler studies.	

Turitz 2014 (Continued)

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Valino 2016
Study characteristics

Patient sampling	Prospective study of screening for adverse obstetric outcomes in women attending for a routine third trimester hospital visit
Patient characteristics and setting	Sample size: 3953 Gestation at sampling: after 35 weeks

Valino 2016 (Continued)

Risk: mixed (normal population, no exclusion criteria, some women with pre-eclampsia)

Setting: King's Hospital, London & Medway Maritime Hospital, Gillingham, Kent

Mode of delivery: 3016 normal vaginal deliveries, 436 elective caesarean sections, 500 emergency caesarean sections (mode of delivery not given for 1 stillbirth)

NICU admission: 232 admissions, 13 of these from the EFW < 5 group

Index tests	PLGF measured in serum using Roche kit, EFW formula unclear; 5th centile used as a threshold for both tests
Target condition and reference standard(s)	SGA defined as birthweight below the 10th centile after correcting for gestational age at delivery
Flow and timing	Included women are those who consented, had data available on all biomarkers, and resulted in the live birth or stillbirth of a phenotypically normal baby at > 24 weeks' gestation.
Comparative	
Notes	The results of the scan were made available to the obstetricians who would have taken specific actions of further monitoring and delivery of the cases of SGA.

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 2: Index Test PLGF			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		

Valino 2016 (Continued)

If a threshold was used, was it pre-specified?	Yes		
		Low	Unclear
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	No		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Voto 1988

Study characteristics			
Patient sampling	Hypertensive pregnant women during the third trimester of pregnancy		
Patient characteristics and setting	Sample size: 215 Gestation at sampling: third trimester Risk: high (all hypertensive)		
Index tests	Threshold > 6 mg%		
Target condition and reference standard(s)	Definition of low birthweight not explicitly stated but it was adjusted for gestational age		
Flow and timing			
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			

Voto 1988 (Continued)

Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test Uric acid			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Unclear		
		Unclear	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Walker 2010
Study characteristics

Patient sampling	Only Caucasian women were included (represent approximately 60% of pregnant women at University College London Hospitals)
Patient characteristics and setting	Sample size: 1238 Gestation at sampling: 30-34 weeks Risk: mixed (exclusion criteria were having a single anatomically normal fetus, normal health status, and uncomplicated obstetrical history) Mode of delivery: 300 caesarean sections

Walker 2010 (Continued)

Index tests	Placental grading measured using Grannum classification, grade II used as a threshold		
Target condition and reference standard(s)	SGA defined as a birthweight below the 10th centile using UK-WHO growth charts		
Flow and timing	1650 women were recruited to the study, of whom 1238 had the 30-34 week ultrasound examination and detailed pregnancy and perinatal outcome available		
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	No		
		High	Low
DOMAIN 2: Index Test Placental grading			
Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear		
If a threshold was used, was it pre-specified?	Yes		
		Unclear	Unclear
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Low	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Unclear		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		

Walker 2010 (Continued)

Low

Weerasinghe 1977
Study characteristics

Patient sampling	327 patients with antenatal complications, including pre-eclampsia, hypertension, clinical FGR, antepartum haemorrhage, threatened abortion, diabetes mellitus, and premature labour.
Patient characteristics and setting	Sample size: 327 Gestation at sampling: between 30 weeks and term Risk: high
Index tests	Urinary E3 measured using Lever's method; 920 assays were obtained from 327 patients and a low index test result was defined as 1 or more low values (-2 SD of mean values)
Target condition and reference standard(s)	SGA defined as a birthweight below 2500 g; no adjustment for gestational age
Flow and timing	
Comparative	
Notes	

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test Urinary E3			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	High
DOMAIN 3: Reference Standard			

Weerasinghe 1977 (Continued)

Is the reference standards likely to correctly classify the target condition?	Unclear
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear
	Unclear
	High
DOMAIN 4: Flow and Timing	
Was there an appropriate interval between index test and reference standard?	Yes
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
	Low

Weiner 2016

Study characteristics			
Patient sampling	All women were recruited in the active phase of labour (mean cervical dilatation at enrolment 5.5 +/- 2.1 cm).		
Patient characteristics and setting	Sample size: 405 Gestation at sampling: after 37 weeks Risk: mixed Setting: Wolfson Medical Centre, Holon, Israel		
Index tests	EFW calculated using Hadlock formula, sensitivity and specificity were calculated for detection (+/- 10%)		
Target condition and reference standard(s)	FGR defined as actual birthweight below the 10th centile		
Flow and timing			
Comparative			
Notes			
Methodological quality			
Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		

Weiner 2016 (Continued)

Did the study avoid inappropriate exclusions?	Yes		
		Low	Unclear
DOMAIN 2: Index Test EFW			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	High
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	Yes		
		Low	

Westergaard 1984

Study characteristics	
Patient sampling	Women that were considered to have a normal singleton pregnancy plus women with a normal singleton pregnancy and abnormal past obstetric history
Patient characteristics and setting	Sample size: 392 Gestation at sampling: 35 weeks Risk: mixed
Index tests	hPL measured by electroimmunoassay, confidence limits of normal ranges were derived from 3648 samples from 721 normal pregnancies.
Target condition and reference standard(s)	IUGR defined as birthweight below the 10th centile for gestational age in the Odense populations and assessment based on phenotypic features

Westergaard 1984 (Continued)

Flow and timing

816 women were invited to participate, of which 611 were recruited. 24 infants with birthweights under the 10th centile but no obvious phenotypic features of IUGR were excluded from the analysis. hPL was determined in 392 women at 35 weeks

Comparative

Notes

Unclear if results were blinded and/or affected management

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	Low
DOMAIN 2: Index Test hPL			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Unclear		
		Unclear	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	No		
		High	

Williams 2002
Study characteristics

Patient sampling	All hypertensive women who presented for care between 1992 and 1996
Patient characteristics and setting	<p>Sample size: 456</p> <p>Gestation at sampling: after 20 weeks</p> <p>Risk: high (patients with gestational hypertension, pre-eclampsia, eclampsia; diabetes and chronic hypertension excluded)</p> <p>Setting: British Columbia Women's Hospital</p>
Index tests	450 umol/L used as a threshold
Target condition and reference standard(s)	SGA infants were defined as being less than the 10th percentile based on the Canadian birthweight percentile figures
Flow and timing	
Comparative	
Notes	<p>Data for combined GH and PE from all tables add to 452 (258 GH and 194 PE) except table 5 (258 GH and 198 PE) but this is the only place that presents totals of SGA so these numbers have to be used.</p> <p>Management was not based on uric acid levels but unclear as to whether this was because measurements were blinded or not.</p>

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test Uric acid			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		

Williams 2002 (Continued)

Were the reference standard results interpreted without knowledge of the results of the index tests? Unclear

Unclear

Low

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard? Yes

Did all patients receive the same reference standard? Yes

Were all patients included in the analysis? Yes

Low

Yassaee 2003
Study characteristics

Patient sampling Cohort study of women with severe pre-eclampsia between 1986 and 2001

Patient characteristics and setting **Sample size:** 103
Gestation at sampling: unknown
Risk: high
Mode of delivery: 59 caesarean sections
Setting: Taleghani Hospital, Tehran, Iran

Index tests No description of when tests were performed

Target condition and reference standard(s) Unclear whether index test results were known. IUGR not defined.

Flow and timing

Comparative

Notes

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
------	--------------------	--------------	------------------------

DOMAIN 1: Patient Selection

Was a consecutive or random sample of patients enrolled? Yes

Was a case-control design avoided? Yes

Did the study avoid inappropriate exclusions? Yes

Yassaee 2003 (Continued)

	Low	High
DOMAIN 2: Index Test Uric acid		
Were the index test results interpreted without knowledge of the results of the reference standard?	Unclear	
If a threshold was used, was it pre-specified?	Yes	
	Unclear	High
DOMAIN 3: Reference Standard		
Is the reference standards likely to correctly classify the target condition?	Yes	
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear	
	Unclear	Unclear
DOMAIN 4: Flow and Timing		
Was there an appropriate interval between index test and reference standard?	Unclear	
Did all patients receive the same reference standard?	Yes	
Were all patients included in the analysis?	Yes	
	Unclear	

Ylikorkala 1973

Study characteristics	
Patient sampling	Series of pregnancies between 1971 and 1972
Patient characteristics and setting	Sample size: 199 Gestation at sampling: third trimester Risk: high (mixture of hypertensive, preeclamptic, diabetic, previous IUD) Setting: Department of Obstetrics and Gynaecology, Oulu University
Index tests	hPL determined using a double antibody radioimmunoassay (HCS Sclavo-Sorin kit), 2.5th centile used as a threshold (calculated according to Herrera 1958)
Target condition and reference standard(s)	SGA defined as birthweight below the 10th centile for gestational age
Flow and timing	687 patients in the series, only data from high-risk patients could be used

Ylikorkala 1973 (Continued)

Comparative

Notes

Placentas were examined microscopically and 46 were found to have degenerative placental changes, 32 had a dysmature placenta, and 78 had normal placental structure. No relationship was found between microscopic calcifications and serum hPL level.

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test hPL			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	Low
DOMAIN 4: Flow and Timing			
Was there an appropriate interval between index test and reference standard?	Yes		
Did all patients receive the same reference standard?	Yes		
Were all patients included in the analysis?	No		
		Low	

Zhang 1990
Study characteristics

Patient sampling	381 pregnant women recruited
Patient characteristics and setting	Sample size: 121 Gestation at sampling: after 36 weeks Risk: high (mixture of hypertension and postterm pregnancy)
Index tests	hPL analysed in serum; threshold - 2 SD for each week of gestation, 4.0 mg/L for 36-41 weeks
Target condition and reference standard(s)	Low birthweight defined as < 2501 g, no adjustments made for gestational age
Flow and timing	Patients were tested at random so not all recruited patients were included, hPL measured using hPL-SRID method
Comparative	
Notes	2 x 2 table derived from high-risk patients only

Methodological quality

Item	Authors' judgement	Risk of bias	Applicability concerns
DOMAIN 1: Patient Selection			
Was a consecutive or random sample of patients enrolled?	Yes		
Was a case-control design avoided?	Yes		
Did the study avoid inappropriate exclusions?	Yes		
		Low	High
DOMAIN 2: Index Test hPL			
Were the index test results interpreted without knowledge of the results of the reference standard?	Yes		
If a threshold was used, was it pre-specified?	Yes		
		Low	Low
DOMAIN 3: Reference Standard			
Is the reference standards likely to correctly classify the target condition?	Yes		
Were the reference standard results interpreted without knowledge of the results of the index tests?	Unclear		
		Unclear	High

Zhang 1990 (Continued)

DOMAIN 4: Flow and Timing

Was there an appropriate interval between index test and reference standard?	Unclear
Did all patients receive the same reference standard?	Yes
Were all patients included in the analysis?	Yes
Low	

ACOG: American College of Obstetricians and Gynecologists; **AFI:** amniotic fluid index; **AGA:** appropriate-for-gestational age; **BP:** blood pressure; **BPD:** biparietal diameter; **BW:** birthweight; **CTG:** cardiotocography; **EFW:** estimated fetal weight; **ELISA:** enzyme-linked immunosorbent assay; **FGR:** fetal growth restriction; **FN:** false negative; **FP:** false positive; **hCG:** human chorionic gonadotrophin; **hPL:** human placental lactogen; **IQR:** interquartile range; **IUD/IUFD:** intrauterine death/intrauterine fetal death; **IUGR:** Intrauterine growth restriction; **NICU:** neonatal intensive care unit; **PE:** pre-eclampsia; **PIGF:** placental growth factor; **PIH:** pregnancy induced hypertension; **RI:** resistance index; **SCN:** special care nursery; **SD:** standard deviation; **SEFW:** sonographic estimated fetal weight; **SFD:** small-for-dates; **SFH:** symphysis fundal height **SGA:** small-for-gestational age; **SUA:** single umbilical artery; **sVEGFR-1:** soluble vascular endothelial growth factor receptor-1.

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Adekanle 2013	Relevant reference standard not recorded
Agboola 1978	2 x 2 table could not be extracted
Aggarwal 2006	Relevant reference standard not recorded
Agorastos 2014	Not a prospective or retrospective cross-sectional or cohort study
Ahmad 1979	Relevant reference standard not recorded
Aickin 1983	2 x 2 table could not be extracted
Al-Amin 2015	Relevant index test not included
Alahakoon 2014	Data presented in another paper
Alberry 2009	2 x 2 table could not be extracted
Algeri 2013	Not a prospective or retrospective cross-sectional or cohort study
Alvarez-Fernandez 2014	2 x 2 table could not be extracted
Alwasel 2013	Relevant reference standard not recorded
Anastasakis 2008	Participants do not match population of interest
Anderson 1978	Relevant index test not included
Arabin 1993	2 x 2 table could not be extracted
Arabin 1995	2 x 2 table could not be extracted

Study	Reason for exclusion
Arias 1977	Relevant index test not included
Ariyuki 1995	Participants do not match population of interest
Atzeni 2012	2 x 2 table could not be extracted
Aviram 2015	2 x 2 table could not be extracted
Axelsson 1978	Relevant reference standard not recorded
Baeza Valenzuela 1995	2 x 2 table could not be extracted
Bahado-Singh 1998	Relevant reference standard not recorded
Bainbridge 2008	2 x 2 table could not be extracted
Bakketeig 1984	2 x 2 table could not be extracted
Baltajian 2016	2 x 2 table could not be extracted
Barden 1999	2 x 2 table could not be extracted
Bardien 2016	Relevant reference standard not recorded
Baron 1996	Index test performed continuously over time
Barrilleaux 2007	Relevant reference standard not recorded
Bartha 2003	Not a prospective or retrospective cross-sectional or cohort study
Bashir 1982	Relevant index test not included
Bastek 2009	Not a prospective or retrospective cross-sectional or cohort study
Battaglia 1995	Relevant reference standard not recorded
Beischer 1975	Data presented in another study
Bell 1967	Not a prospective or retrospective cross-sectional or cohort study
Benton 2010	Not a prospective or retrospective cross-sectional or cohort study
Benton 2011	Not a prospective or retrospective cross-sectional or cohort study
Benton 2011a	Not a prospective or retrospective cross-sectional or cohort study
Benton 2012	Not a prospective or retrospective cross-sectional or cohort study
Benton 2012a	Not a prospective or retrospective cross-sectional or cohort study
Benton 2014	Participants do not match population of interest
Benton 2014a	Relevant reference standard not recorded
Benz 1980	Not a prospective or retrospective cross-sectional or cohort study

Study	Reason for exclusion
Berendtsen 1985	Index test measured continuously over time
Bergsjø 1973	2 x 2 table could not be extracted
Berle 1973	2 x 2 table could not be extracted
Berle 1973a	2 x 2 table could not be extracted
Berle 1973b	Not a prospective or retrospective cross-sectional or cohort study
Berle 1975	2 x 2 table could not be extracted
Bernardes 2013	Relevant reference standard not recorded
Bernatavicius 2013	Participants do not match population of interest
Bersinger 2004	Not a prospective or retrospective cross-sectional or cohort study
Bersinger 2005	2 x 2 table could not be extracted
Bhansali 1975	Review
Bian 1992	2 x 2 table could not be extracted
Biberoglu 2016	Relevant reference standard not recorded
Bieglmayer 1981	2 x 2 table could not be extracted
Bila 1980	2 x 2 table could not be extracted
Bitzer 1985	2 x 2 table could not be extracted
Blaskova 1977	Relevant index test not included
Bligh 2015	Relevant reference standard not recorded
Blitz 2016	Review
Blumenfeld 2007	2 x 2 table could not be extracted
Bobrow 2002	Not a prospective or retrospective cross-sectional or cohort study
Bock 1976	2 x 2 table could not be extracted
Boij 2012	2 x 2 table could not be extracted
Borges 2005	2 x 2 table could not be extracted
Botasheva 2016	2 x 2 table could not be extracted
Boucoiran 2012	2 x 2 table could not be extracted
Branconi 1981	2 x 2 table could not be extracted
Brush 1970	Participants do not match population of interest

Study	Reason for exclusion
Bukowski 2014	Not a prospective or retrospective cross-sectional or cohort study
Butcher 2012	Participants do not match population of interest
Buyon 2011	Relevant reference standard not recorded
Cage 2009	2 x 2 table could not be extracted
Calabrese 2012	Relevant reference standard not recorded
Calderon 2011	2 x 2 table could not be extracted
Campobasso 1967	Participants do not match population of interest
Camus-Bablon 1990	Relevant reference standard not recorded
Carne 1987	Data presented in another study
Castren 1966	Relevant reference standard not recorded
Cavazza 2015	2 x 2 table could not be extracted
Ceccarello 1980	Relevant index test not included
Cefalo 2005	Commentary
Cetin 2014	2 x 2 table could not be extracted
Cetin 2016	Relevant reference standard not recorded
Chaiworapongsa 2008	2 x 2 table could not be extracted
Chaiworapongsa 2012	Data presented in another study
Chaiworapongsa 2013a	2 x 2 table could not be extracted
Chaiworapongsa 2013b	2 x 2 table could not be extracted
Chambers 1989	Relevant index test not included
Chang 1993	Relevant index test not included
Chang 1994	Not a prospective or retrospective cross-sectional or cohort study
Chapman 1978	2 x 2 table could not be extracted
Chapman 1981	Participants do not match population of interest
Chappell 2002	2 x 2 table could not be extracted
Chard 1982	Review
Chauhan 2012	2 x 2 table could not be extracted
Chawengsettakul 2015	Relevant reference standard not recorded

Study	Reason for exclusion
Chew 2014	Relevant reference standard not recorded
Church 2016	Not a prospective or retrospective cross-sectional or cohort study
Clelia 2013	Relevant reference standard not recorded
Clowse 2011	Relevant reference standard not recorded
Cody 2013	2 x 2 table could not be extracted
Cody 2016	2 x 2 table could not be extracted
Cooley 2011	2 x 2 table could not be extracted
Cordano 1988	Participants do not match population of interest
Craig 1996	2 x 2 table could not be extracted
Crane 1979	2 x 2 table could not be extracted
Crawford 1985	2 x 2 table could not be extracted
D'Anna 2000	Relevant reference standard not recorded
Daikoku 1979	2 x 2 table could not be extracted
Darling 2014	Participants do not match population of interest
Dave 2016	Relevant index test not included
Dawood 1976	Participants do not match population of interest
De Marchi 1977	Participants do not match population of interest
Del Moral 2015	Relevant reference standard not recorded
Deter 2016	Relevant reference standard not recorded
Di Lorenzo 2013	2 x 2 table could not be extracted
Dombrowski 1992	Participants do not match population of interest
Dombrowski 1992a	Participants do not match population of interest
Ducarme 2012	2 x 2 table could not be extracted
Duff 1986	Relevant index test not included
Dutton 2012	Relevant reference standard not recorded
Dutton 2012a	Relevant reference standard not recorded
Eik-Nes 1984	Participants do not match population of interest
El-Ahmady 1997	2 x 2 table could not be extracted

Study	Reason for exclusion
Elchalal 2000	Participants do not match population of interest
Ernst 2016	2 x 2 table could not be extracted
Fadigas 2015	2 x 2 table could not be extracted
Falkner 1995	Relevant index test not included
Ferrazzi 1986	Relevant index test not included
Fioretti 1986	2 x 2 table could not be extracted
Fischer-Rasmussen 1971	2 x 2 table could not be extracted
Fisteag-Kiprono 2006	Relevant reference standard not recorded
Forger 2016	2 x 2 table could not be extracted
Fotiou 2015	2 x 2 table could not be extracted
Furuhashi 1984	2 x 2 table could not be extracted
Gabbay-Benziv 2016a	2 x 2 table could not be extracted
Gabbay-Benziv 2017	Abstract for another study
Gaillard 2014	2 x 2 table could not be extracted
Gao 2008	2 x 2 table could not be extracted
Garcia-Flores 2015	Relevant reference standard not recorded
Garoff 1976	2 x 2 table could not be extracted
Gaziano 1988	Participants do not match population of interest
Geerts 2007	Relevant reference standard not recorded
Gerhard 1987	2 x 2 table could not be extracted
Gernand 2015	2 x 2 table could not be extracted
Gherzi 1981	Index test measured continuously over time
Giambanco 1986	Review
Giardini 2014	Relevant reference standard not recorded
Gloning 1991	Relevant index test not included
Goetzinger 2013	Relevant index test not included
Goldenberg 1993	Relevant reference standard not recorded
Goldenberg 1997	Relevant reference standard not recorded

Study	Reason for exclusion
Gomez-Roig 2015	2 x 2 table could not be extracted
Gordon 1978	Relevant reference standard not recorded
Grantz 2016	Relevant reference standard not recorded
Gravett 2015	2 x 2 table could not be extracted
Griffin 2014	Data presented in another study
Gris 2015	2 x 2 table could not be extracted
Habib 2002	2 x 2 table could not be extracted
Hargreaves 2011	2 x 2 table could not be extracted
Harper 2014	Not a prospective or retrospective cross-sectional or cohort study
Hassan 1987	Relevant index test not included
Hawkins 2014	Relevant index test not included
Heazell 2014	Review
Henrichs 2016	Not a prospective or retrospective cross-sectional or cohort study
Hensleigh 1977	Participants do not match population of interest
Herraiz 2014	Not a prospective or retrospective cross-sectional or cohort study
Hinkle 2015	2 x 2 table could not be extracted
Hughes 1980	2 x 2 table could not be extracted
Husse 2014	2 x 2 table could not be extracted
Jabeen 1999	Participants do not match population of interest
James-Todd 2015	Relevant reference standard not recorded
Johnson 2011	Relevant reference standard not recorded
Johnstone 2015	Participants do not match population of interest
Karjalainen 1975	Relevant reference standard not recorded
Karlsen 2016	2 x 2 table could not be extracted
Kase 2012	2 x 2 table could not be extracted
Kase 2012a	2 x 2 table could not be extracted
Kazzi 1983	Participants do not match population of interest
Khalil 2015	2 x 2 table could not be extracted

Study	Reason for exclusion
Khalil 2016	2 x 2 table could not be extracted
Kihaile 1988	Participants do not match population of interest
Kim 2009	2 x 2 table could not be extracted
Kim 2014	2 x 2 table could not be extracted
Kim 2016	Participants do not match population of interest
Kjos 2015	2 x 2 table could not be extracted
Kneitel 2016	2 x 2 table could not be extracted
Kolovetsiou-Kreiner 2014	2 x 2 table could not be extracted
Kolovetsiou-Kreiner 2015	2 x 2 table could not be extracted
Krochik 2010	Participants do not match population of interest
Kulkarni 1981	2 x 2 table could not be extracted
Kulkarni 2010	2 x 2 table could not be extracted
Kullander 1982	Relevant index test not included
Kundu 1978	2 x 2 table could not be extracted
Kunzig 1975	2 x 2 table could not be extracted
Kunzig 1980	2 x 2 table could not be extracted
Lai 2014	Relevant reference standard not recorded
Larkin 2015	2 x 2 table could not be extracted
Larsen 1992	2 x 2 table could not be extracted
Larsen 1997	Relevant reference standard not recorded
Laurin 1987a	2 x 2 table could not be extracted
Lean 2016	2 x 2 table could not be extracted
Leanos-Miranda 2013	2 x 2 table could not be extracted
Lechner 1987	2 x 2 table could not be extracted
Levine 2005	Not a prospective or retrospective cross-sectional or cohort study
Li 2014	2 x 2 table could not be extracted
Little 2016	2 x 2 table could not be extracted
Lobmaier 2014	2 x 2 table could not be extracted

Study	Reason for exclusion
London 1983	Relevant reference standard not recorded
MacDonald 1983	Not a prospective or retrospective cross-sectional or cohort study
Macmillian 1976	2 x 2 table could not be extracted
Maly 1987	Relevant reference standard not recorded
March 2015	2 x 2 table could not be extracted
Margossian 2016	2 x 2 table could not be extracted
Markestad 1997	Not a prospective or retrospective cross-sectional or cohort study
Martins 2005	Relevant index test not included
Masoura 2014	2 x 2 table could not be extracted
Matthews 2017	2 x 2 table could not be extracted
Mazzocco 2014	Relevant reference standard not recorded
McKenna 2003	Relevant index test not included
Melamed 2015	2 x 2 table could not be extracted
Melamed 2016	2 x 2 table could not be extracted
Melamed 2016a	2 x 2 table could not be extracted
Merriam 2014	Participants do not match population of interest
Mertens 1975	Participants do not match population of interest
Mirza 2015	Participants do not match population of interest
Miwa 2014	Relevant reference standard not recorded
Mlynarczyk 2015	2 x 2 table could not be extracted
Mlynarczyk 2015a	2 x 2 table could not be extracted
Mone 2016	2 x 2 table could not be extracted
Moore 2012	Relevant reference standard not recorded
Morrison 1980	Participants do not match population of interest
Muraguchi 1981	Participants do not match population of interest
Myatt 2013	2 x 2 table could not be extracted
Nadal 2015	2 x 2 table could not be extracted
Nair 2016	2 x 2 table could not be extracted

Study	Reason for exclusion
Nelson 2015	Participants do not match population of interest
Nice 2014	Not a prospective or retrospective cross-sectional or cohort study
Nice 2014a	Not a prospective or retrospective cross-sectional or cohort study
Nieder 1976	2 x 2 table could not be extracted
Nielsen 1981	2 x 2 table could not be extracted
Niknafs 2001	Relevant index test not included
O'Connor 2015	Participants do not match population of interest
Obiekwe 1982	Relevant index test not included
Odibo 2014	2 x 2 table could not be extracted
Okonofua 1986	Participants do not match population of interest
Pal 2015	Participants do not match population of interest
Palomaki 2015	Participants do not match population of interest
Palomaki 2015a	Participants do not match population of interest
Papastefanou 2014	2 x 2 table could not be extracted
Papastefanou 2015	2 x 2 table could not be extracted
Parra Saavedra 2015	Relevant reference standard not recorded
Parrish 2010	2 x 2 table could not be extracted
Partap 2015	Relevant reference standard not recorded
Pavelka 1982	Participants do not match population of interest
Pecks 2015	2 x 2 table could not be extracted
Peixoto 2016	2 x 2 table could not be extracted
Perez-Cruz 2015	2 x 2 table could not be extracted
Perry 1986	2 x 2 table could not be extracted
Persson 1978	2 x 2 table could not be extracted
Persson 1980	2 x 2 table could not be extracted
Peyronnet 2016	2 x 2 table could not be extracted
Pfeiffer 1990	2 x 2 table could not be extracted
Pinheiro 2014	2 x 2 table could not be extracted

Study	Reason for exclusion
Pledger 1984	Not a prospective or retrospective cross-sectional or cohort study
Pluta 1979	Participants do not match population of interest
Ponce 1995	2 x 2 table could not be extracted
Powers 2010	Not a prospective or retrospective cross-sectional or cohort study
Prakash 2012	2 x 2 table could not be extracted
Qublan 2005	2 x 2 table could not be extracted
Raghuramulu 1978	2 x 2 table could not be extracted
Rajasingam 2009	2 x 2 table could not be extracted
Rasanen 2015	Relevant reference standard not recorded
Reck 1987	Relevant index test not included
Ris-Stalpers 2012	2 x 2 table could not be extracted
Riss 1982	Relevant reference standard not recorded
Rizos 2013	Not a prospective or retrospective cross-sectional or cohort study
Rocca 1995	2 x 2 table could not be extracted
Romero 2008	Not a prospective or retrospective cross-sectional or cohort study
Ronin-Walknowska 1984	2 x 2 table could not be extracted
Rosendahl 1988	Relevant index test not included
Rosendahl 1991	Relevant index test not included
Rothenbacher 2016	2 x 2 table could not be extracted
Ruozi Berretta 1967	Participants do not match population of interest
Sabbagha 1979	Review
Salahuddin 2016	2 x 2 table could not be extracted
Salas 1993	2 x 2 table could not be extracted
Salas 1998	2 x 2 table could not be extracted
Salas 2006	2 x 2 table could not be extracted
Saleh 2015	Relevant reference standard not recorded
Salkie 1977	2 x 2 table could not be extracted
Samanta 1989	2 x 2 table could not be extracted

Study	Reason for exclusion
Sanchez Fernandez 2015	2 x 2 table could not be extracted
Sarandakou 1989	2 x 2 table could not be extracted
Sato 1974	Participants do not match population of interest
Secher 1986	2 x 2 table could not be extracted
Secher 1987	2 x 2 table could not be extracted
Sekar 2015	2 x 2 table could not be extracted
Selbing 1984	Relevant index test not included
Semczuk-Sikora 2007	2 x 2 table could not be extracted
Shaarawy 2001	2 x 2 table could not be extracted
Shah 1996	2 x 2 table could not be extracted
Sharf 1984	Participants do not match population of interest
Sheth 2016	Relevant reference standard not recorded
Shibata 2005	Not a prospective or retrospective cross-sectional or cohort study
Sibiude 2012	2 x 2 table could not be extracted
Sichinava 2014	2 x 2 table could not be extracted
Singer 1970	2 x 2 table could not be extracted
Smith 2014	Not a prospective or retrospective cross-sectional or cohort study
Smith-Bindman 2002	Participants do not match population of interest
Smith-Bindman 2003	Participants do not match population of interest
Soler 1975	Index test performed continuously over time
Sood 1988	Not a prospective or retrospective cross-sectional or cohort study
Sorensen 2000	2 x 2 table could not be extracted
Souka 2012	2 x 2 table could not be extracted
Souka 2013	2 x 2 table could not be extracted
Sovio 2014	Abstract for an included study
Spellacy 1967	Not a prospective or retrospective cross-sectional or cohort study
Spellacy 1975	Not a prospective or retrospective cross-sectional or cohort study
Spellacy 1976	Not a prospective or retrospective cross-sectional or cohort study

Study	Reason for exclusion
Spona 1971	2 x 2 table could not be extracted
Spona 1972	2 x 2 table could not be extracted
Stefanelli 2014	Participants do not match population of interest
Stefanidis 1998	Not a prospective or retrospective cross-sectional or cohort study
Strizhakov 2013	2 x 2 table could not be extracted
Strom 1983	Not a prospective or retrospective cross-sectional or cohort study
Sucak 2010	2 x 2 table could not be extracted
Sudik 1982	Participants do not match population of interest
Sundrani 2013	2 x 2 table could not be extracted
Tajik 2012	2 x 2 table could not be extracted
Takeuchi 1988	2 x 2 table could not be extracted
Tammemae 2016	2 x 2 table could not be extracted
Tayama 1983	2 x 2 table could not be extracted
Taylor 2003	2 x 2 table could not be extracted
Teoh 1971	Index test performed continuously over time
Tonari 1987	2 x 2 table could not be extracted
Torok 1987	Index test performed continuously over time
Triunfo 2014	2 x 2 table could not be extracted
Triunfo 2016	2 x 2 table could not be extracted
Triunfo 2016a	2 x 2 table could not be extracted
Tsiakkas 2015	Relevant reference standard not recorded
Tsiakkas 2016	Relevant reference standard not recorded
Turpin 2015	2 x 2 table could not be extracted
Van Rijn 2015	2 x 2 table could not be extracted
Varma 1979	Relevant index test not included
Varma 1982	2 x 2 table could not be extracted
Vatten 2012	Not a prospective or retrospective cross-sectional or cohort study
Vinayagam 2015	2 x 2 table could not be extracted

Study	Reason for exclusion
Wallner 2007	2 x 2 table could not be extracted
Watson 1973	2 x 2 table could not be extracted
Weissbach 1985	2 x 2 table could not be extracted
Whigham 1980	2 x 2 table could not be extracted
White 2016	Relevant reference standard not recorded
Woelkers 2016	Relevant reference standard not recorded
Woo 2016	Relevant index test not included
Woods 2015	2 x 2 table could not be extracted
Wurz 1983	Participants do not match population of interest
Xing 2016	2 x 2 table could not be extracted
Xu 2015	2 x 2 table could not be extracted
Yamaguchi 1979	2 x 2 table could not be extracted
Yanaihara 1984	Not a prospective or retrospective cross-sectional or cohort study
Zail 1975	2 x 2 table could not be extracted
Zera 2011	2 x 2 table could not be extracted
Zhang 2011	2 x 2 table could not be extracted
Zhao 2010	Not a prospective or retrospective cross-sectional or cohort study
Zlatnik 1979	Participants do not match population of interest
Zuckerman 1974	Participants do not match population of interest

Characteristics of studies awaiting classification *[ordered by study ID]*

Bracali 1968

Study characteristics	
Patient sampling	Unknown
Patient characteristics and setting	Unknown
Index tests	Urinary oestriol
Target condition and reference standard(s)	Unknown
Flow and timing	Unknown

Bracali 1968 *(Continued)*

Comparative

Notes

Fuks 1990
Study characteristics

Patient sampling	Unknown
Patient characteristics and setting	Unknown
Index tests	Placental grading
Target condition and reference standard(s)	Fetal and neonatal outcomes - unclear
Flow and timing	Unknown
Comparative	
Notes	

Jain 2000
Study characteristics

Patient sampling	Unknown
Patient characteristics and setting	Unknown
Index tests	Placental grading
Target condition and reference standard(s)	IUGR
Flow and timing	Unknown
Comparative	
Notes	

Khan 2004
Study characteristics

Patient sampling	Unknown
Patient characteristics and setting	Unknown
Index tests	Ultrasound EFW?

Khan 2004 *(Continued)*

Target condition and reference standard(s)	IUGR
Flow and timing	Unknown
Comparative	
Notes	

Ruseva 1983
Study characteristics

Patient sampling	Unknown
Patient characteristics and setting	Unknown
Index tests	Unknown
Target condition and reference standard(s)	Unknown
Flow and timing	Unknown
Comparative	
Notes	

Ruseva 1985
Study characteristics

Patient sampling	Unknown
Patient characteristics and setting	Unknown
Index tests	Ultrasound EFW
Target condition and reference standard(s)	Unknown
Flow and timing	Unknown
Comparative	
Notes	

Ruseva 1985a
Study characteristics

Patient sampling	Unknown
------------------	---------

Ruseva 1985a *(Continued)*

Patient characteristics and setting	Unknown
Index tests	Ultrasound EFW
Target condition and reference standard(s)	Unknown
Flow and timing	Unknown
Comparative	
Notes	

Ruseva 1988
Study characteristics

Patient sampling	Unknown
Patient characteristics and setting	Unknown
Index tests	Ultrasound EFW
Target condition and reference standard(s)	Unknown
Flow and timing	Unknown
Comparative	
Notes	

Serban 1971
Study characteristics

Patient sampling	Unknown
Patient characteristics and setting	Unknown
Index tests	hPL
Target condition and reference standard(s)	Unknown
Flow and timing	Unknown
Comparative	
Notes	

EFW: estimated fetal weight; **hPL:** human placental lactogen; **IUGR:** Intrauterine growth restriction

DATA

Presented below are all the data for all of the tests entered into the review.

Table Tests. Data tables by test

Test	No. of studies	No. of participants
1 EFW and SGA	32	51702
2 Placental grading and SGA	12	4940
3 hPL and SGA	20	3486
4 E3 and SGA	9	2773
5 Urinary E3 and SGA	9	92406
6 PlGF and SGA	8	6997
7 UA and SGA	8	2884
8 E3 AND/OR hPL and SGA	1	88
9 EFW for SGA - 10th centile threshold only	25	47057
10 hPL for SGA - 10th centile threshold only	8	1414
11 EFW and SGA3	3	5678
13 PlGF and SGA3	2	1861
14 hPL and stillbirth	6	544
15 Urinary E3 and stillbirth	7	92186
16 PlGF and stillbirth	4	5894
17 UA and stillbirth	4	2063
18 Placental grading and stillbirth	3	15236
19 SGA data from studies with both SGA infants and stillbirths	17	99920
20 Stillbirth data from studies with both SGA infants and stillbirths	17	100050
22 PlGF or EFW and SGA	1	343
23 PlGF or EFW and SGA3	1	343

Test 1. EFW and SGA.

Test 2. Placental grading and SGA.**Test 3. hPL and SGA.****Test 4. E3 and SGA.****Test 5. Urinary E3 and SGA.****Test 6. PlGF and SGA.****Test 7. UA and SGA.****Test 8. E3 AND/OR hPL and SGA.****Test 9. EFW for SGA - 10th centile threshold only.****Test 10. hPL for SGA - 10th centile threshold only.****Test 11. EFW and SGA3.****Test 13. PlGF and SGA3.****Test 14. hPL and stillbirth.****Test 15. Urinary E3 and stillbirth.**

Test 16. PlGF and stillbirth.

Test 17. UA and stillbirth.

Test 18. Placental grading and stillbirth.

Test 19. SGA data from studies with both SGA infants and stillbirths.

Test 20. Stillbirth data from studies with both SGA infants and stillbirths.

Test 22. PlGF or EFW and SGA.

Test 23. PlGF or EFW and SGA3.

ADDITIONAL TABLES

Table 1. Characteristics of included studies table

Study	Target condition	Test	Sample size	Cases	Threshold	EFW formula	Risk	Interventions	Gestational age	Low RoB all
Altmann 1978	SGA	hPL	10	6	Abnormal value		High	Unknown		No
Altmann 1978	SGA	Placental grading	9	6	Grade III		High	Unknown		No
Altmann 1978	Stillbirth	hPL	10	3	Abnormal value		High	Unknown		No
Altmann 1978	Stillbirth	Placental grading	9	2	Grade III		High	Unknown		No
Amini 2014	SGA	UA	404	46	+1 SD		Mixed	Unknown		No
Baird 2016	SGA	EFW	107	78	Below 10th centile	Hadlock	High	Unknown		No
Barel 2016	SGA	EFW	14089	1218	Below 10th centile	Hadlock	Mixed	Unknown		No
Beischer 1991	SGA	Urinary E3	72062	5390	8 mg per 24 hour (30 w) to 12 mg/24 hours (40 w)		High	Yes	Before 37 weeks	No
Beischer 1991	Stillbirth	Urinary E3	72062	452	8 mg per 24 hours (30 w) to 12 mg/24 hours (40 w)		High	Yes	Before 37 weeks	No
Bellomo 2011	SGA	UA	163	43	309 µmol/L		High	Unknown	Before 37 weeks	No
Ben-Haroush 2007	SGA	EFW	259	19	Below 10th centile	Hadlock	Low	Unknown	Before 37 weeks	No
Benton 2016	SGA	PIGF	411	159	12 pg/mL		High	No		Yes
Benton 2016	Stillbirth	PIGF	411	7	12 pg/mL		High	No		Yes
Berkowitz 1988	SGA	EFW	168	42	Below 10th centile	Shepard	High	Yes		No
Bikmetova 2013	SGA	EFW	518	185	Unknown	Unknown	Unknown	Unknown		No
Callec 2015	SGA	EFW	1897	156	Below 10th centile	Hadlock	Mixed	Yes	Before 37 weeks	No

Table 1. Characteristics of included studies table *(Continued)*

Campbell 1972	SGA	Urinary E3	284	87	Unknown		High	Unknown	Yes
Campbell 1972	Stillbirth	Urinary E3	284	11	Unknown		High	Unknown	Yes
Cedard 1979	SGA	E3	64	17	Below 10th centile		High	Unknown	37 weeks onwards No
Chaiworapongsa 2013	SGA	PIGF	1269	108	< 0.3 MoM		Mixed	Unknown	Before 37 weeks No
Chaiworapongsa 2013	SGA3	PIGF	1269	23	< 0.3 MoM		Mixed	Unknown	Before 37 weeks No
Chaiworapongsa 2013	Stillbirth	PIGF	1269	5	< 0.12 MoM		Mixed	Unknown	Before 37 weeks No
Chard 1985	SGA	E3	392	39	Below 10th centile		Mixed	Unknown	No
Chard 1985	SGA	hPL	392	39	Below 10th centile		Mixed	Unknown	No
Chauhan 1999	SGA	EFW	324	44	Below 10th centile	Hadlock	Mixed	Unknown	No
Chauhan 1999a	SGA	EFW	574	59	Below 10th centile	Hadlock	Mixed	Unknown	No
Chauhan 2003	SGA	EFW	264	58	Below 10th centile	Hadlock	High	Yes	No
Chen 2012	SGA	Placental grading	105	36	Grade III		High	Unknown	No
Chen 2012	Stillbirth	Placental grading	105	13	Grade III		High	Unknown	No
Chen 2012a	SGA	Placental grading	113	23	Grade III		High	No	Before 37 weeks Yes
Chen 2015	Stillbirth	Placental grading	15122	99	Grade III		Low	Unknown	Before 37 weeks Yes
Chervenak 1984	SGA	EFW	179	17	L99CL	Shepard	High	Unknown	No
Chew 1976	SGA	Urinary E3	43	15	Below 2.5th centile		High	No	Yes

Table 1. Characteristics of included studies table (Continued)

Chew 1976	Stillbirth	Urinary E3	43	6	-2SD		High	No		Yes
Chitlange 1990	SGA	Placental grading	270	72	Grade III		Low	Unknown	Before 37 weeks	No
Christensen 2015	SGA	EFW	157	7	Below 10th centile	Hadlock	High	Unknown		No
Elliott 1970	Stillbirth	Urinary E3	22	2	-2 SD		High	Yes		Yes
Estel 1989	SGA	Placental grading	55	21	Grade III		High	Unknown	37 weeks onwards	No
Fliegner 1979	SGA	Urinary E3	329	37	8 mg per 24 hours (30 w) to 12 mg/24 hours (40 w)		Mixed	No		Yes
Fliegner 1979	Stillbirth	Urinary E3	329	5	8 mg per 24 hours (30 w) to 12 mg/24 hours (40 w)		Mixed	No		Yes
Freire 2010	SGA	EFW	122	21	Below 10th centile	Hadlock	Mixed	Unknown		No
Gabbay-Benziv 2016	SGA	EFW	6126	638	Below 10th centile	Hadlock	Mixed	Unknown		No
Geerts 2016	SGA	EFW	210	60	Below 10th centile	Hadlock	High	Unknown		Yes
Geerts 2016	SGA	Placental grading	188	51	Grade III	Hadlock	High	Unknown		Yes
Gerhard 1986	SGA	E3	869	78	Below 10th centile		Mixed	No	Before 37 weeks	Yes
Gohari 1978	SGA	hPL	111	38	5 µg/mL		High	Unknown		No
Granat 1977	SGA	hPL	29	10	4 µg/mL		High	No		No
Griffin 2015	SGA	EFW	586	192	Below 10th centile	Unknown	High	Unknown	Before 37 weeks	No
Griffin 2015	SGA3	EFW	586	78	Below 10th centile	Unknown	High	Unknown	Before 37 weeks	No

Table 1. Characteristics of included studies table (Continued)

Griffin 2015	SGA	PIGF	592	192	Below 5th centile	Unknown	High	Unknown	Before 37 weeks	
Griffin 2015	SGA3	PIGF	592	78	Below 5th centile	Unknown	High	Unknown	Before 37 weeks	No
Griffin 2015	SGA	PIGF or EFW	343	115	Below 10th centile/below 5th centile	Unknown	High	Unknown	Before 37 weeks	
Griffin 2015	SGA3	PIGF or EFW	343	52	Below 10th centile/below 5th centile	Unknown	High	Unknown	Before 37 weeks	No
Gupta 2008	SGA	EFW	38	15	Below 10th centile	Hadlock	High	Unknown		No
Hammad 2015	SGA	EFW	71	9	Below 10th centile	Hadlock	Low	No		No
Hatfield 2010	SGA	EFW	659	48	Below 10th centile	Unknown	High	Unknown		No
Hawkins 2012	SGA	UA	1306	224	+1 SD		High	No		Yes
Hawkins 2012	Stillbirth	UA	1483	5	+ 1SD		High	No		Yes
Hendrix 2000	SGA	EFW	367	22	SEFW < 2500 g	Hadlock	Mixed	No		No
Howell 1985	SGA	hPL	501	50	2.5 µg/mL		Mixed	Unknown		Yes
Jauniaux 1996	SGA	UA	41	16	4 mg/dL		High	Unknown	Before 37 weeks	No
Kazzi 1983a	SGA	Placental grading	109	42	Grade III		Mixed	Unknown		No
Kienast 2016	SGA	PIGF	346	40	Below 5th centile		Mixed	Unknown	Before 37 weeks	No
Klebe 1990	SGA	hPL	13	3	Below 10th centile		High	Unknown		
Kunz 1976	SGA	hPL	83	15	Below 5th centile		High	Unknown		No
Kunz 1976	SGA	Urinary E3	83	15	Below 5th centile		High	Unknown		No

Table 1. Characteristics of included studies table (Continued)

Laurin 1987	SGA	EFW	2068	78	Predicted BW deviation for GA of -15% or more	Eik-Nes	Mixed	Unknown	Before 37 weeks	No
Leader 1980	Stillbirth	hPL	135	8	4 mg/mL		High	No		Yes
Lenstrup 1982	SGA	E3 and/or hPL	88	9	Below 10th centile		Mixed	No	Before 37 weeks	No
Lilford 1983	SGA	hPL	522	52	Below 10th centile		Mixed	Unknown		No
MacLeod 2013	SGA	EFW	90	8	< 2500 g (+/- 10%)	Unknown	Mixed	Unknown		No
Mahran 1988	SGA	EFW	828	98	-2 SD	Unknown	Mixed	Unknown		No
Marin 1979	SGA	hPL	47	13	Below 5th centile		Mixed	Unknown		No
McKenna 2005	SGA	Placental grading	1902	109	Grade III		Low	Yes	Before 37 weeks	No
Miller 1988	SGA	Placental grading	246	29	Grade III		Mixed	Unknown		Yes
Molvarec 2013	SGA	PIGF	89	22	3.9:1		High	No	Before 37 weeks	No
Montan 1986	SGA	Placental grading	307	6	Grade III		Mixed	Unknown	37 weeks onwards	No
Nice 2016	SGA	hPL	77	23	< 0.8 MoM		Mixed	No		No
Nice 2016	SGA	PIGF	76	23	12 pg/mL		Mixed	No		No
Nielsen 1985	SGA	E3	1018	61	Below 2.5th centile		Mixed	No	Before 37 weeks	Yes
Nisbet 1982	SGA	E3	103	46	-2 SD		High	Unknown		No
Nisbet 1982	SGA	hPL	166	71	-2 SD		High	Unknown		No
Oats 1979	SGA	Urinary E3	19119	1391	8 mg/24 hours (30 w) to 12 mg/24 hours (40 w)		Mixed	Unknown		No

Table 1. Characteristics of included studies table (Continued)

Oats 1979	Stillbirth	Urinary E3	19119	172	8 mg/24 hours (30 w) to 12 mg/24 hours (40 w)		Mixed	Unknown		No
Obiekwe 1983	SGA	hPL	522	29	4.8 µg/mL		Mixed	Unknown	37 weeks onwards	No
Odendaal 1981	SGA	E3	53	30	Below 10th centile		High	No		No
Odendaal 1981	SGA	hPL	77	43	Below 10th centile		High	No		No
Odendaal 1981	SGA	Urinary E3	46	28	Below 10th centile		High	No		No
Odendaal 1997	SGA	UA	196	100	520 µmol/L		High	No	Before 37 weeks	No
Odendaal 1997	Stillbirth	UA	196	18	520 µmol/L		High	No	Before 37 weeks	No
Ott 1984	SGA	EFW	595	111	-1.5 SD	Shepard	Mixed	Unknown	Before 37 weeks	No
Palo 1987	SGA	E3	90	40	-2 SD		High	Unknown		Yes
Palo 1989	SGA	EFW	186	97	Below 10th centile	Eik-Nes	High	Unknown		No
Patterson 1983	SGA	Placental grading	398	21	Grade III		High	Unknown		No
Redman 1976	Stillbirth	UA	281	2	360 µmol/L		Unknown	No	Before 37 weeks	No
Roma 2015	SGA	EFW	1115	134	Below 10th centile	Hadlock	Mixed	Yes	Before 37 weeks	No
Roma 2015	SGA3	EFW	1115	49	Below 10th centile	Hadlock	Mixed	Yes	Before 37 weeks	No
Sagen 1984	SGA	E3	74	40	Below 10th centile		High	Yes		No
Sagen 1984	SGA	hPL	74	40	Below 10th centile		High	Yes		No

Table 1. Characteristics of included studies table *(Continued)*

Sekar 2016	SGA	EFW	150	15	Below 10th centile	Hadlock	Mixed	No		No
Shawkat 2015	SGA	PIGF	261	106	12 pg/mL		High	No		Yes
Shawkat 2015	Stillbirth	PIGF	261	3	12 pg/mL		High	No		Yes
Siebert 1974	SGA	hPL	67	11	Below 10th centile		Low	Yes		No
Siebert 1974	Stillbirth	hPL	20	2	Below 10th centile		Low	Yes		No
Skovron 1991	SGA	EFW	768	69	Below 10th centile	Shepard	Mixed	Unknown	Before 37 weeks	No
Sovio 2015	SGA	EFW	3977	352	Below 10th centile	Hadlock	Mixed	Unknown	Before 37 weeks	No
Sovio 2015	SGA3	EFW	3977	87	Below 10th centile	Hadlock	Mixed	Unknown	Before 37 weeks	No
Spernol 1989	SGA	E3	110	22	Below 5th centile		Mixed	Unknown		No
Spernol 1989	SGA	hPL	110	22	Below 10th centile		Mixed	Unknown		No
Steiner 1991	SGA	hPL	113	68	Below 10th centile		High	Unknown		No
Steiner 1991	SGA	Urinary E3	113	68	8 mg per 24 hours (30 w) to 12 mg/24 hours (40 w)		High	Unknown		No
Takeuchi 1985	SGA	EFW	210	39	Below 10th centile	Hadlock	Mixed	Unknown		No
Trudinger 1979	SGA	hPL	59	25	Below 10th centile		High	Unknown		No
Trudinger 1979	Stillbirth	hPL	59	1	Below 10th centile		High	Unknown		No
Turitz 2014	SGA	EFW	10642	1876	Below 10th centile	Hadlock	Mixed	Yes	Before 37 weeks	No
Valino 2016	SGA	EFW	3953	379	Below 10th centile	Unknown	Mixed	Yes		No
Valino 2016	SGA	PIGF	3953	379	Below 10th centile	Unknown	Mixed	Yes		No
Valino 2016	Stillbirth	PIGF	3953	1	Below 10th centile	Unknown	Mixed	Yes		No

Table 1. Characteristics of included studies table (Continued)

Voto 1988	SGA	UA	215	30	6 mg%		High	Unknown		No
Walker 2010	SGA	Placental grading	1238	104	Grade III		Mixed	Unknown	Before 37 weeks	No
Weerasinghe 1977	SGA	Urinary E3	327	45	-2 SD		High	Unknown		No
Weerasinghe 1977	Stillbirth	Urinary E3	327	3	-2 SD		High	Unknown		No
Weiner 2016	SGA	EFW	405	30	Below 10th centile	Hadlock	Mixed	No	37 weeks onwards	No
Westergaard 1984	SGA	hPL	392	28	Abnormal value		Mixed	Unknown	Before 37 weeks	No
Williams 2002	SGA	UA	456	87	450 µmol/L		High	No		No
Yassae 2003	SGA	UA	103	59	6 mg/dL		High	Unknown		No
Yassae 2003	Stillbirth	UA	103	12	6 mg/dL		High	Unknown		No
Ylikorkala 1973	Stillbirth	hPL	199	14	Below 2.5th centile		High	Unknown		No
Zhang 1990	SGA	hPL	121	38	4 µg/mL		High	No		No
Zhang 1990	Stillbirth	hPL	121	8	4 µg/mL		High	No		No

EFW: estimated fetal weight; **hPL:** human placental lactogen; **MoM:** multiple of the median; **PIGF:** Placental growth factor; **SD:** standard deviation; **SEFW:** sonographic estimated fetal weight; **SGA:** small-for-gestational age; **UA:** uric acid

Table 2. Indirect comparison of tests for assessment of small-for-gestational-age infants ≤tenth centile outcome

Ratio of diagnostic odds ratios (95% CI), P value		Estimated fetal weight	Human placental lactogen	Oestriol	Urinary oestriol	Placental growth factor	Uric acid
Studies;	DOR (95% CI)	21.3 (13.1 to 34.6)	4.78 (3.21 to 7.13)	4.00 (2.91 to 5.49)	3.59 (1.78 to 7.23)	2.73 (1.67 to 4.48)	2.36 (1.25 to 4.46)

Table 2. Indirect comparison of tests for assessment of small-for-gestational-age infants ≤tenth centile outcome (Continued)

	partici- pants (SGA cases)							
Estimated fetal weight	32; 51,702 (6169)	21.3 (13.1 to 34.6)						
Human placental lactogen	20; 3486 (624)	4.78 (3.21 to 7.13)	4.45 (2.38 to 8.25), P < 0.0001					
Oestriol	9; 2773 (373)	4.00 (2.91 to 5.49)	5.33 (2.98 to 9.52), P < 0.0001	1.20 (0.72 to 1.99), P = 0.48				
Urinary oestriol	9; 92,406 (7076)	3.59 (1.78 to 7.23)	5.93 (2.53 to 13.9), P < 0.0001	1.33 (0.59 to 2.98), P = 0.48	1.11 (0.52 to 2.40), P = 0.78			
Placental growth factor	7; 6405 (837)	2.73 (1.67 to 4.48)	7.79 (3.90 to 15.6), P < 0.0001	1.75 (0.93 to 3.30), P = 0.08	1.46 (0.81 to 2.63), P = 0.26	1.31 (0.56 to 3.09), P = 0.53		
Uric acid	8; 2884 (605)	2.36 (1.25 to 4.46)	9.02 (4.05 to 20.1), P < 0.0001	2.03 (0.96 to 4.29), P = 0.06	1.69 (0.83 to 3.44), P = 0.14	1.52 (0.59 to 3.92), P = 0.38	1.16 (0.52 to 2.59), P = 0.72	
Placental grading	12; 4940 (520)	2.34 (1.33 to 4.12)	9.09 (4.32 to 19.2), P < 0.0001	2.04 (1.02 to 4.08), P = 0.04	1.71 (0.89 to 3.26), P = 0.10	1.53 (0.62 to 3.77), P = 0.35	1.17 (0.55 to 2.47), P = 0.68	1.01 (0.43 to 2.36), P = 0.98

All available data were used for indirect comparison of the accuracy of the tests. The ratio of diagnostic odds ratios is the DOR of the test in the column divided by the DOR of the test in the row. If the ratio of DORs is greater than one, then the diagnostic accuracy of the test in the column is higher than that of the test in the row; if the ratio is less than one, the diagnostic accuracy of the test in the row is higher than that of the test in the column.

Table 3. Comparison of test accuracy at different levels of prevalence of small-for-gestational-age (birthweight $\leq$ tenth centile) infants

Prevalence (%)	Specificity	False positives	Test	Sensitivity (95% CI)	Missed SGA infants
10	0.74	234	EFW	0.88 (0.82 to 0.92)	12
			hPL	0.63 (0.53 to 0.71)	38
			Oestriol	0.58 (0.51 to 0.66)	42
			Urinary oestriol	0.56 (0.39 to 0.72)	45
			PIGF	0.49 (0.37 to 0.61)	52
			Uric acid	0.45 (0.31 to 0.61)	55
			Placental grading	0.45 (0.32 to 0.59)	55
19	0.74	211	EFW	0.88 (0.82 to 0.92)	23
			hPL	0.63 (0.53 to 0.71)	71
			Oestriol	0.58 (0.51 to 0.66)	80
			Urinary oestriol	0.56 (0.39 to 0.72)	84
			PIGF	0.49 (0.37 to 0.61)	97
			Uric acid	0.45 (0.31 to 0.61)	104
			Placental grading	0.45 (0.32 to 0.59)	105
35	0.74	169	EFW	0.88 (0.82 to 0.92)	42
			hPL	0.63 (0.53 to 0.71)	131
			Oestriol	0.58 (0.51 to 0.66)	146
			Urinary oestriol	0.56 (0.39 to 0.72)	155
			PIGF	0.49 (0.37 to 0.61)	179
			Uric acid	0.45 (0.31 to 0.61)	192
			Placental grading	0.45 (0.32 to 0.59)	193
10	0.88	108	EFW	0.74 (0.64 to 0.83)	26
			hPL	0.39 (0.30 to 0.49)	61
			Oestriol	0.35 (0.28 to 0.43)	65
			Urinary oestriol	0.33 (0.20 to 0.50)	68
			PIGF	0.24 (0.15 to 0.38)	76

Table 3. Comparison of test accuracy at different levels of prevalence of small-for-gestational-age (birthweight ≤ tenth centile) infants (Continued)

			Uric acid	0.27 (0.19 to 0.38)	73
			Placental grading	0.24 (0.15 to 0.36)	76
19	0.88	97	EFW	0.74 (0.64 to 0.83)	49
			hPL	0.39 (0.30 to 0.49)	116
			Oestriol	0.35 (0.28 to 0.43)	123
			Urinary oestriol	0.33 (0.20 to 0.50)	128
			PIGF	0.24 (0.15 to 0.38)	144
			Uric acid	0.27 (0.19 to 0.38)	139
			Placental grading	0.24 (0.15 to 0.36)	145
35	0.88	78	EFW	0.74 (0.64 to 0.83)	90
			hPL	0.39 (0.30 to 0.49)	212
			Oestriol	0.35 (0.28 to 0.43)	227
			Urinary oestriol	0.33 (0.20 to 0.50)	235
			PIGF	0.24 (0.15 to 0.38)	265
			Uric acid	0.27 (0.19 to 0.38)	255
			Placental grading	0.24 (0.15 to 0.36)	266
10	0.96	36	EFW	0.47 (0.35 to 0.59)	53
			hPL	0.17 (0.12 to 0.23)	84
			Oestriol	0.14 (0.11 to 0.19)	86
			Urinary oestriol	0.13 (0.07 to 0.23)	87
			PIGF	0.10 (0.07 to 0.16)	90
			Uric acid	0.09 (0.05 to 0.16)	92
			Placental grading	0.09 (0.05 to 0.15)	92
19	0.96	23	EFW	0.47 (0.35 to 0.59)	101
			hPL	0.17 (0.12 to 0.23)	159
			Oestriol	0.14 (0.11 to 0.19)	163
			Urinary oestriol	0.13 (0.07 to 0.23)	166
			PIGF	0.10 (0.07 to 0.16)	171

Table 3. Comparison of test accuracy at different levels of prevalence of small-for-gestational-age (birthweight $\leq$ tenth centile) infants (Continued)

35	0.96	26	Uric acid	0.09 (0.05 to 0.16)	173
			Placental grading	0.09 (0.05 to 0.15)	174
			EFW	0.47 (0.35 to 0.59)	186
			hPL	0.17 (0.12 to 0.23)	292
			Oestriol	0.14 (0.11 to 0.19)	301
			Urinary oestriol	0.13 (0.07 to 0.23)	305
			PIGF	0.10 (0.07 to 0.16)	315
			Uric acid	0.09 (0.05 to 0.16)	319
			Placental grading	0.09 (0.05 to 0.15)	319

EFW: estimated fetal weight; **hPL:** human placental lactogen; **PIGF:** placental growth factor.

The sensitivities were estimated from the SROC curves at quartiles of the observed specificity in the included studies. Using these sensitivities and specificities, along with quartiles of prevalence from the included studies, the numbers of missed SGA infants and false positives were calculated based on a hypothetical cohort of 1000 pregnant women predicted to have a small-for-gestational-age infant.

Table 4. Indirect comparison of biochemical tests for predicting stillbirth

Ratio of diagnostic odds ratios (95% CI),			Placental growth factor	Human placental factor	Urinary oestriol
P value					
	Studies; participants (stillbirths)	DOR (95% CI)	49.2 (12.7 to 191)	11.4 (4.29 to 30.2)	5.83 (4.91 to 6.92)
Placental growth factor	4; 5894 (16)	49.2 (12.7, 191)			
Human placental lactogen	6; 544 (36)	11.4 (4.29, 30.2)	4.32 (0.81 to 23.0), P = 0.08		
Urinary oestriol	7; 92,186 (651)	5.83 (4.91, 6.92)	8.44 (2.15 to 33.1), P = 0.004	1.95 (0.72 to 5.27), P = 0.17	
Uric acid	4; 2063 (37)	4.02 (0.95, 17.0)	12.2 (1.69 to 88.5), P = 0.016	2.83 (0.50 to 16.1), P = 0.22	1.45 (0.34 to 6.19), P = 0.60

All available data were used for the indirect comparison of the accuracy of the tests. The ratio of diagnostic odds ratios is the DOR of the test in the column divided by the DOR of the test in the row. If the ratio of DORs is greater than one, then the diagnostic accuracy of the test in the column is higher than that of the test in the row; if the ratio is less than one, the diagnostic accuracy of the test in the row is higher than that of the test in the column.

Table 5. Comparison of test accuracy at different levels of prevalence of stillbirth

Prevalence (%)	Specificity	False positives	Test	Sensitivity (95% CI)	Missed stillbirths
0.9	0.63	367	PIGF	0.97 (0.88 to 0.99)	1
			hPL	0.87 (0.72 to 0.95)	2
			Urinary oestriol	0.77 (0.74 to 0.80)	3
			Uric acid	0.70 (0.36 to 0.91)	3
1.7	0.63	364	PIGF	0.97 (0.88 to 0.99)	1
			hPL	0.87 (0.72 to 0.95)	3
			Urinary oestriol	0.77 (0.74 to 0.80)	4
			Uric acid	0.70 (0.36 to 0.91)	6
9.1	0.63	336	PIGF	0.97 (0.88 to 0.99)	4
			hPL	0.87 (0.72 to 0.95)	12
			Urinary oestriol	0.77 (0.74 to 0.80)	21
			Uric acid	0.70 (0.36 to 0.91)	28
0.9	0.78	218	PIGF	0.93 (0.78 to 0.98)	1
			hPL	0.76 (0.55 to 0.90)	3
			Urinary oestriol	0.62 (0.58 to 0.66)	4
			Uric acid	0.53 (0.21 to 0.83)	5
1.7	0.78	216	PIGF	0.93 (0.78 to 0.98)	2
			hPL	0.76 (0.55 to 0.90)	5
			Urinary oestriol	0.62 (0.58 to 0.66)	7
			Uric acid	0.53 (0.21 to 0.83)	8
9.1	0.78	200	PIGF	0.93 (0.78 to 0.98)	7
			hPL	0.76 (0.55 to 0.90)	22
			Urinary oestriol	0.62 (0.58 to 0.66)	35
			Uric acid	0.53 (0.21 to 0.83)	43
0.9	0.89	109	PIGF	0.86 (0.61 to 0.96)	2
			hPL	0.58 (0.35 to 0.79)	4

Table 5. Comparison of test accuracy at different levels of prevalence of stillbirth *(Continued)*

1.7	0.89	108	Urinary oestriol	0.42 (0.38 to 0.46)	6
			Uric acid	0.33 (0.11 to 0.68)	7
			PlGF	0.86 (0.61 to 0.96)	3
			hPL	0.58 (0.35 to 0.79)	8
			Urinary oestriol	0.42 (0.38 to 0.46)	10
			Uric acid	0.33 (0.11 to 0.68)	12
9.1	0.89	100	PlGF	0.86 (0.61 to 0.96)	13
			hPL	0.58 (0.35 to 0.79)	38
			Urinary oestriol	0.42 (0.38 to 0.46)	53
			Uric acid	0.33 (0.11 to 0.68)	61

hPL: human placental lactogen; **PlGF:** placental growth factor.

The sensitivities were estimated from the SROC curves at quartiles of the observed specificity in the included studies. Using these sensitivities and specificities, along with quartiles of prevalence from the included studies, the numbers of missed stillbirths and false positives were calculated based on a hypothetical cohort of 1000 pregnant women predicted to have a stillbirth.

APPENDICES

Appendix 1. Examples of placental function tests available for diagnostic use (compiled 29th March 2016)

Placental growth factor (PlGF)

Triage PlGF (Alere, San Diego) – point of care fluorescence immunoassay (<http://www.plgf.com/home/proposed-clinical-use-of-plgf/alere-triage-plgf.html>)

Elecsys™ Preeclampsia (sFlt-1 & PlGF) – automated immunoassay performed on Roche platform

(<http://www.cobas.com/home/product/clinical-and-immunochemistry-testing/elecsys-preeclampsia-assays-sFlt-1-PlGF.html>)

Oestriol (E3)

AutoDELFIA Unconjugated Estriol (Perkin Elmer) – automated fluorescence immunoassay performed on Perkin-Elmer platform. (<http://www.perkinelmer.co.uk/product/autodelfia-unconjugated-estriol-ue3-ki-b083-301>)

Beckman Coulter – automated immunoassay performed on Beckman Coulter platform (<https://www.beckmancoulter.com/wsrportal/bibliography?docname=DS14764A%20Access%20Unconjugated%20Estriol%20US%20Data%20Sheet.pdf>)

Elecsys™ Estradiol – automated immunoassay performed on Roche platform (http://www.cobas.com/content/dam/cobas_com/pdf/lists/parameter-list-swa.pdf)

Appendix 2. Search strategy

Database: Ovid MEDLINE(R) 1946 to October Week 3 2016 (October 26th 2016)

Search Strategy:

1 Placental insufficiency/

2 ((placenta\$ or f?etoplacental or uteroplacental) adj2 (insufficien\$ or fail\$ or function\$)).ti,ab.

3 fetal movement/

Biochemical tests of placental function versus ultrasound assessment of fetal size for stillbirth and small-for-gestational-age infants (Review)

214

- 4 fetal growth retardation/
- 5 ((reduc\$ or decline\$) adj2 f?etal movement).ti,ab.
- 6 (stillborn or stillbirth).ti,ab.
- 7 Stillbirth/
- 8 ((f?etal or intrauterine or intra-uterine) adj2 (growth or death\$ or loss\$)).ti,ab.
- 9 IUGR.ti,ab.
- 10 (small adj2 gestational age).ti,ab.
- 11 ((neonatal or perinatal or fetal or birth\$ or deliver\$) adj2 outcome\$).ti,ab.
- 12 f?etal move\$.ti,ab.
- 13 or/1-12
- 14 oestradiol.ti,ab.
- 15 estradiol.ti,ab.
- 16 exp Estradiol/
- 17 oestriol.ti,ab.
- 18 exp progesterone/
- 19 progesterone.ti,ab.
- 20 exp pregnenolone/
- 21 pregnenolone.ti,ab.
- 22 exp Chorionic Gonadotropin/
- 23 human chorionic gonadotrophin.ti,ab.
- 24 hCG.ti,ab.
- 25 placental lactogen/
- 26 hPL.ti,ab.
- 27 human placental lactogen.ti,ab.
- 28 human placental growth hormone.ti,ab.
- 29 placental protein 13.ti,ab.
- 30 placental growth factor.ti,ab.
- 31 plasma placental protein.ti,ab.
- 32 pregnancy specific glycoprotein\$.ti,ab.
- 33 Pregnancy-Specific beta 1-glycoproteins/
- 34 schwangerschaft protein 1.ti,ab.
- 35 pregnancy specific beta 1-glycoprotein.ti,ab.
- 36 exp ultrasonography, Prenatal/
- 37 (sonograph\$ or ultraso\$).ti,ab.
- 38 Grannum grading.ti,ab.

39 biomarkers/

40 biomarker\$.mp. or marker\$.ti,ab.

41 or/14-40

42 13 and 41

43 limit 42 to humans

Database: Ovid MEDLINE(R) In-Process & Other Non-Indexed Citations (October 25, 2016)

Search Strategy:

1 (placenta\$ or f?etoplacental or uteroplacental) adj2 (insufficien\$ or fail\$ or function\$)).ti,ab.

2 ((reduc\$ or decline\$) adj2 fe?tal movement).ti,ab.

3 (stillborn or stillbirth).ti,ab.

4 ((f?etal or intrauterine or intra-uterine) adj2 (growth or death\$ or loss\$)).ti,ab.

5 IUGR.ti,ab.

6 (small adj2 gestational age).ti,ab.

7 ((neonatal or perinatal or fetal or birth\$ or deliver\$) adj2 outcome\$).ti,ab.

8 f?etal move\$.ti,ab.

9 or/1-8

10 oestradiol.ti,ab.

11 estradiol.ti,ab.

12 oestriol.ti,ab.

13 progesterone.ti,ab.

14 pregnenolone.ti,ab.

15 human chorionic gonadotrophin.ti,ab.

16 hCG.ti,ab.

17 hPL.ti,ab.

18 human placental lactogen.ti,ab.

19 human placental growth hormone.ti,ab.

20 placental protein 13.ti,ab.

21 placental growth factor.ti,ab.

22 plasma placental protein.ti,ab.

23 pregnancy specific glycoprotein\$.ti,ab.

24 schwangerschaft protein 1.ti,ab.

25 pregnancy specific beta 1-glycoprotein.ti,ab.

26 (sonograph\$ or ultraso\$).ti,ab.

27 Grannum grading.ti,ab.

28 placental lactogen.ti,ab.

29 biomarker\$.ti,ab.

30 marker\$.ti,ab.

31 or/10-30

32 9 and 31

Database: Embase (Ovid) 1974 to week 4 October 2016

Search Strategy:

1 ((placenta\$ or fetoplacental or foetoplacental or uteroplacental) adj2 (insufficien\$ or fail\$ or function\$)).ti,ab.

2 ((reduc\$ or decline\$) adj2 (fetal or foetal) adj movement)).ti,ab.

3 (stillborn or stillbirth).ti,ab.

4 ((fetal or foetal or intrauterine or intra-uterine) adj2 (growth or death\$ or loss\$)).ti,ab.

5 IUGR.ti,ab.

6 (small adj2 gestational age).ti,ab.

7 ((neonatal or perinatal or fetal or foetal or birth\$ or deliver\$) adj2 outcome\$).ti,ab.

8 (fetal or foetal) adj move\$.ti,ab.

9 exp placenta insufficiency/

10 exp fetus movement/

11 exp intrauterine growth retardation/

12 exp stillbirth/

13 oestradiol.ti,ab.

14 estradiol.ti,ab.

15 exp estradiol/

16 exp estriol/

17 oestriol.ti,ab.

18 exp progesterone/

19 progesterone.ti,ab.

20 exp pregnenolone/

21 pregnenolone.ti,ab.

22 exp chorionic gonadotropin/

23 human chorionic gonadotropin.ti,ab.

24 hCG.ti,ab.

25 placental lactogen.ti,ab.

26 exp placenta lactogen/

27 hPL.ti,ab.

28 human placental growth hormone.ti,ab.

29 exp placenta protein/

30 placental protein 13.ti,ab.
31 placental growth factor.ti,ab.
32 plasma placental protein.ti,ab.
33 pregnancy specific glycoprotein\$.ti,ab.
34 exp pregnancy specific beta1 glycoprotein/
35 schwangerschaft protein 1.ti,ab.
36 pregnancy specific beta 1-glycoprotein.ti,ab.
37 exp fetus echography/
38 sonograph\$ or ultrason\$.ti,ab.
39 Grannum grading.ti,ab.
40 biological marker/
41 biomarker\$ or marker\$.ti,ab.
42 or/13-41
43 or/1-12
44 42 and 43
45 limit 44 to human

Database: Cochrane Library (Wiley) (CENTRAL) Issue 7 of 12 2016 (DARE, HTA,EED) Issue 2 of 4 2015, (CDSR) Issue 7 of 12 2016

Search strategy:

#1 MeSH descriptor: [Placental Insufficiency] explode all trees
#2 (placenta* or fetoplacental or foetoplacental or uteroplacental) near/2 (insufficienc* or fail* or function*)
#3 MeSH descriptor: [Fetal Movement] explode all trees
#4 MeSH descriptor: [Fetal Growth Retardation] explode all trees
#5 (reduc* or declin*) near/2 ("fetal move*") or ("foetal move*")
#6 stillborn or stillbirth
#7 MeSH descriptor: [Stillbirth] explode all trees
#8 (fetal or foetal or intrauterine or intra-uterine) near/2 (growth or death* or loss*)
#9 IUGR
#10 small near/2 (gestational next age)
#11 (neonatal or perinatal or fetal or birth* or deliver*) near/2 (outcome*)
#12 fetal next move*
#13 #1 or #2 or #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12
#14 oestradiol
#15 estradiol
#16 oestriol
#17 progesterone

- #18 MeSH descriptor: [Estradiol] explode all trees
- #19 MeSH descriptor: [Progesterone] explode all trees
- #20 pregnenolone
- #21 MeSH descriptor: [Pregnenolone] explode all trees
- #22 MeSH descriptor: [Chorionic Gonadotropin] explode all trees
- #23 "human chorionic gonadotrop*"
- #24 hCG
- #25 hPL
- #26 MeSH descriptor: [Placental Lactogen] explode all trees
- #27 "human placental lactogen"
- #28 "human placental growth hormone"
- #29 "placental protein 13"
- #30 "placental growth factor"
- #31 "plasma placental protein"
- #32 "pregnancy specific glycoprotein*"
- #33 MeSH descriptor: [Pregnancy-Specific beta 1-Glycoproteins] explode all trees
- #34 "schwangerschaft protein 1"
- #35 "pregnancy specific beta 1-glycoprotein"
- #36 MeSH descriptor: [Ultrasonography, Prenatal] explode all trees
- #37 sonograph* or ultraso*
- #38 "Grannum grading"
- #39 MeSH descriptor: [Biological Markers] explode all trees
- #40 biomarker* or marker*
- #41 #14 or #15 or #16 or #17 or #18 or #19 or #20 or #21 or #22 or #23 or #24 or #25 or #26 or #27 or #28 or #29 or #30 or #31 or #32 or #33 or #34 or #35 or #36 or #37 or #38 or #39 or #40
- #42 #13 and #41

Database: CINAHL (Ebsco) 1937 – present (28 October 2016)

Search strategy:

- S1 (placenta* or fetoplacental or foetoplacental or uteroplacental) N2 (insufficienc* or fail* or function*)
- S2 (MH"Placental Insufficiency")
- S3 (MH"Fetal Movement")
- S4 (MH"Fetal Growth Retardation")
- S5 (reduc* or decline) N2 (fetal move* or foetal move*)
- S6 stillborn or stillbirth
- S7 (MH"Perinatal Death")

S8 (fetal or foetal or intrauterine or intra-uterine) N2 (growth or death* or loss*)

S9 IUGR

S10 (small) N2 (gestational age)

S11 (neonatal or perinatal or fetal or birth* or deliver*) N2 (outcome*)

S12 fetal move* or foetal move*

S13 S1 or S2 or S3 or S4 or S5 or S6 or S7 or S8 or S9 or S10 or S11 or S12

S14 oestradiol or estradiol or oestriol or progesterone or pregnenolone

S15 (MH"Estradiol")

S16 (MH"Progesterone+")

S17 (MH "Gonadotropins, Chorionic")

S18 human chorionic gonadotrophin

S19 hCG

S20 placental lactogen

S21 human placental growth hormone

S22 placental protein 13

S23 placental growth factor

S24 plasma placental protein

S25 pregnancy specific glycoprotein*

S26 schwangerschaft protein 1

S27 pregnancy specific beta 1-glycoprotein*

S28 (MH"Ultrasonography, Prenatal")

S29 sonograph* or ultraso*

S30 Grannum grading

S31 S14 or S15 or S16 or S17 or S18 or S19 or S20 or S21 or S22 or S23 or S24 or S25 or S26 or S27 or S28 or S29 or S30

S32 S13 and S31

S33 S13 and S31 Limiters exclude MEDLINE records

Searches were also performed in ClinicalTrials.gov and WHO ICTRP; based on the nature of these databases full search strategies were not constructed, searches were performed using key terms such as: "stillbirth*" AND "predictors OR placenta*".

Appendix 3. QUADAS 2 tool for assessing methodological quality of included studies

Domain	Signalling question	Signalling question	Signalling question	Risk of bias	Concerns about applicability
Patient selection	Was a consecutive or random sample of patients enrolled?	Was a case-control design avoided?	Did the study avoid inappropriate exclusions?	Could the selection of patients have introduced bias?	Are there concerns that the included patients and setting do

(Continued)

				not match the review question?
	<p>Yes if participants were consecutively enrolled or if all eligible participants were enrolled or participants were randomly sampled.</p> <p>No if participants were selected from those eligible.</p> <p>Unclear if participant selection was not clear from the report.</p>	<p>Yes if a case control design was avoided.</p> <p>No if a case control design was used.</p> <p>Unclear if the study design could not be determined from the report.</p>	<p>Yes if the study avoided inappropriate exclusions (e.g. only excluded multiple pregnancy, congenital abnormalities).</p> <p>No if participants were excluded inappropriately (e.g. ethnicity, age, income).</p> <p>Unclear if appropriateness of exclusions could not be assessed from report.</p>	<p>Low risk if yes to all of the signalling questions.</p> <p>High or unclear risk if "no" or "unclear" was reported for at least one signalling question.</p> <p>Low concern if the sample of pregnant women represent the women indicated by the review question and if inappropriate exclusions were avoided.</p> <p>High concern if the sample of pregnant women are different from those indicated in the review question.</p> <p>Unclear concern if insufficient information was available.</p>
Index test – test of placental function	<p>Were the index test results interpreted without knowledge of the results of the reference standard?</p>	<p>If a threshold was used was it pre-specified?</p>	<p>Could the conduct or interpretation of the index test have introduced bias?</p>	<p>Are there concerns that the index test, its conduct or its interpretation differ from the review question?</p>
	<p>Yes if the result(s) of the test of placental function was interpreted without knowledge of the reference standard.</p> <p>No if the result(s) of the test of placental function was interpreted with knowledge of the reference standard.</p> <p>Unclear if this was not clear in the report.</p>	<p>Yes if the criteria for a positive result of the placental function test were pre-specified.</p> <p>No if the criteria for a positive result were not pre-specified or deviated from that specified.</p> <p>Unclear if this was not clear from the report.</p>	<p>Low risk if yes to all of the signalling questions.</p> <p>High or unclear risk if "no" or "unclear" was reported for at least one signalling question.</p>	<p>Low concern if the placental function test was performed as described in the review question (e.g. after 24 weeks of pregnancy to assess placental function).</p> <p>High concern if the placental function test was performed in a different way to that described in the review question.</p> <p>Unclear concern if insufficient information was available.</p>
Reference standard and target condition	<p>Is there reference standard likely to correctly classify the target condition?</p>	<p>Were the reference standard results interpreted without knowledge of the results of the index test?</p>	<p>Could the reference standard, its conduct or interpretation have introduced bias?</p>	<p>Are there concerns that the target condition as defined by the reference standard does not match the question?</p>

(Continued)

	<p>Yes if an acceptable reference standard was used (e.g. SGA = birthweight < 10th centile, Stillbirth = baby born with no signs of life after 24 weeks' gestation).</p> <p>No if pregnancy outcome was not classified by an acceptable reference standard (e.g. low birthweight < 2.5 kg).</p> <p>Unclear if this was not clear from the report.</p>	<p>Yes if pregnancy outcome (live or still-birth), and a diagnosis of a small for gestational age infant was made without the knowledge of results of the placental function test.</p> <p>No if pregnancy outcome and a diagnosis of a small for gestational age infant were made with the knowledge of the results of the placental function test.</p> <p>Unclear if this was not clear from the report.</p>	<p>Low risk if yes to all of the signalling questions.</p> <p>High or unclear risk if "no" or "unclear" was reported for at least one signalling question.</p>	<p>Low concern if acceptable reference standards were used and if the reference standard was interpreted without the knowledge of the placental function test.</p> <p>High concern if an acceptable reference standard was not used or the results were interpreted with knowledge of the result of the placental function test.</p> <p>Unclear concern if insufficient information was available.</p>
Flow and Timing	Was there an appropriate interval between the index test and reference standard?	Did all patients receive the same reference standard?	Were all patients included in the analysis?	Could the patient flow have introduced bias?
	<p>Yes If acquisition of the index test occurred prior to birth (reference standards both determined after birth).</p> <p>No if sample acquired after delivery of the infant (i.e. known reference standard).</p> <p>Unclear if this was not clear from the report.</p>	<p>Yes if all participants had the outcome of pregnancy and birthweight recorded.</p> <p>No if some participants do not have the outcome of pregnancy and birthweight recorded.</p> <p>Unclear if this was not clear from the report.</p>	<p>Yes if all participants recruited to the study were included in the final analysis.</p> <p>No if all participants were not included in the final analysis.</p> <p>Unclear if this was not clear from the report.</p>	<p>Low risk if yes to all of the signalling questions.</p> <p>High or unclear risk if "no" or "unclear" was reported for at least one signalling question.</p>

Appendix 4. Glossary

Aneuploidy: a condition where there are an abnormal number of chromosomes in a cell

Centile: percentile, below the tenth centile means in the bottom 10%

Ductus venosus: a fetal blood vessel that helps carry oxygenated blood to the heart

Echotexture: the appearance of human tissue when looked at using ultrasound

Efficacy: efficiency, the ability of a test to produce the desired result

False negative: a negative test result in an individual with the condition of interest

False positive: a positive test result in an individual without the condition of interest

Heterogeneity: variation, diversity

Fetal growth restriction (FGR: a condition where a fetus fails to attain its growth potential, i.e. is smaller than expected for its genetic potential.

hPL - human placental lactogen - a protein made by the trophoblast layer of the placenta.

Morphology: appearance, structure

Placental analyte: a substance produced by the placenta that can be measured and analysed

PLGF: placental growth factor- a protein made by the trophoblast layer of the placenta.

Small-for-gestational-age infant (SGA infant): the condition where the fetal weight or birthweight is beneath a specific threshold, generally considered to be the 10th centile.

SROC plot: summary receiver operator characteristic plot - a scatterplot of estimates of sensitivity and specificity from included studies.

Umbilical artery Doppler: a measurement of fetal blood flow through the umbilical artery using Doppler ultrasound.

Uterine artery Doppler: a measurement of maternal blood flow through the uterine artery using Doppler ultrasound.

Appendix 5. Risk of bias and applicability concerns summary for each study, by index test

[Figure 14](#), [Figure 15](#), [Figure 16](#), [Figure 17](#), [Figure 18](#), [Figure 19](#), [Figure 20](#)

Figure 14. Risk of bias and applicability concerns summary: review authors' judgements about each domain for each included study of human placental lactogen (hPL)

	Risk of Bias				Applicability Concerns		
	Patient Selection	Index Test: hPL	Reference Standard	Flow and Timing	Patient Selection	Index Test: hPL	Reference Standard
Altmann 1978	+	+	?	?	-	?	+
Chard 1985	+	?	?	+	+	+	-
Gohari 1978	+	+	?	+	-	?	+
Granat 1977	+	+	?	+	-	+	+
Howell 1985	+	+	+	+	+	+	+
Klebe 1990	+	?	?	+	?	+	-
Kunz 1976	+	+	?	+	-	-	+
Leader 1980	+	+	+	+	-	+	+
Lenstrup 1982	+	?	+	+	+	?	+
Lilford 1983	+	-	?	+	+	+	-
Marin 1979	+	?	+	?	+	+	+
Nice 2016	?	+	+	+	+	+	+
Nisbet 1982	-	+	+	?	+	+	+
Obiekwe 1983	+	?	?	+	+	+	-
Odendaal 1981	+	+	+	?	-	+	+
Sagen 1984	+	+	?	+	-	+	+
Siebert 1974	+	?	-	+	-	?	+
Spernal 1989	?	+	?	+	?	+	+
Steiner 1991	+	?	+	?	-	?	+
Trudinger 1979	+	+	?	+	-	+	+
Westergaard 1984	+	?	?	-	+	+	+
Ylikorkala 1973	+	+	?	+	-	+	+
Zhang 1990	+	+	?	+	-	+	-

High
 Unclear
 Low

Figure 15. Risk of bias and applicability concerns summary: review authors' judgements about each domain for each included study of placental grading.

	Risk of Bias				Applicability Concerns		
	Patient Selection	Index Test: Placental grading	Reference Standard	Flow and Timing	Patient Selection	Index Test: Placental grading	Reference Standard
Altmann 1978	+	+	?	?	-	+	+
Chen 2012	+	+	?	+	+	+	-
Chen 2012a	+	+	?	+	-	+	-
Chen 2015	+	+	+	+	-	+	+
Chitlange 1990	?	+	+	+	+	+	-
Estel 1989	+	?	?	?	-	+	+
Geerts 2016	+	+	+	+	-	+	+
Kazzi 1983a	+	+	?	-	-	+	+
McKenna 2005	+	+	?	+	-	+	+
Miller 1988	+	+	+	+	+	+	+
Montan 1986	+	+	+	+	+	+	?
Patterson 1983	-	+	?	+	-	+	+
Walker 2010	-	?	+	+	+	?	+

High
 Unclear
 Low

Figure 16. Risk of bias and applicability concerns summary: review authors' judgements about each domain for each included study of serum oestriol (E3).

	<u>Risk of Bias</u>				<u>Applicability Concerns</u>		
	Patient Selection	Index Test: E3	Reference Standard	Flow and Timing	Patient Selection	Index Test: E3	Reference Standard
Cedard 1979	+	+	?	+	-	+	+
Chard 1985	+	?	?	+	+	+	-
Gerhard 1986	+	+	+	+	+	+	+
Lenstrup 1982	+	?	+	+	+	?	+
Nielsen 1985	+	+	+	+	+	+	+
Nisbet 1982	-	+	+	?	+	+	+
Odendaal 1981	+	+	+	?	-	+	+
Palo 1987	+	+	+	+	-	+	+
Sagen 1984	+	+	?	+	-	+	+
Spernol 1989	?	+	?	+	?	+	+

High
Unclear
Low

Figure 17. Risk of bias and applicability concerns summary: review authors' judgements about each domain for each included study of estimated fetal weight (EFW).

	Risk of Bias				Applicability Concerns		
	Patient Selection	Index Test: EFW	Reference Standard	Flow and Timing	Patient Selection	Index Test: EFW	Reference Standard
Baird 2016	+	?	+	+	-	+	+
Barel 2016	+	+	+	+	+	+	+
Ben-Haroush 2007	+	+	?	?	-	+	+
Berkowitz 1988	+	+	?	+	-	+	+
Bikmetova 2013	+	?	?	?	?	+	?
Callec 2015	?	?	?	+	+	+	+
Chauhan 1999	+	+	?	+	-	+	+
Chauhan 1999a	+	+	?	+	-	+	+
Chauhan 2003	+	?	?	+	-	+	+
Chervenak 1984	+	+	?	+	-	+	?
Christensen 2015	+	?	+	+	-	+	-
Freire 2010	+	?	?	+	+	?	+
Gabbay-Benziv 2016	+	+	+	+	+	+	+
Geerts 2016	+	+	+	+	-	+	+
Griffin 2015	+	+	?	?	-	+	+
Gupta 2008	+	+	?	+	-	+	+
Hammad 2015	+	+	+	+	-	+	+
Hatfield 2010	+	-	+	+	-	-	+
Hendrix 2000	+	+	+	+	+	+	-
Laurin 1987	+	+	?	+	+	+	+
MacLeod 2013	+	+	?	+	+	+	-
Mahrar 1988	+	+	+	+	+	+	+
Ott 1984	+	+	+	+	+	+	+
Palo 1989	+	+	?	+	-	+	+
Roma 2015	+	+	?	+	+	+	+
Sekar 2016	+	+	?	+	+	+	+

Figure 17. (Continued)

Sekar 2016	+	+	?	+	+	+	+
Skovron 1991	+	+	?	+	+	+	+
Sovio 2015	+	+	+	?	+	+	+
Takeuchi 1985	?	+	?	?	+	+	+
Turitz 2014	+	+	?	+	+	+	+
Valino 2016	+	+	?	+	+	+	+
Weiner 2016	+	+	?	+	?	-	+

High
Unclear
Low

Figure 18. Risk of bias and applicability concerns summary: review authors' judgements about each domain for each included study of placental growth factor (PlGF).

		<u>Risk of Bias</u>				<u>Applicability Concerns</u>		
		Patient Selection	Index Test: PlGF	Reference Standard	Flow and Timing	Patient Selection	Index Test: PlGF	Reference Standard
	Benton 2016	+	+	+	+	-	+	-
	Chaiworapongsa 2013	+	-	?	+	+	?	+
	Griffin 2015	+	+	?	?	-	+	+
	Kienast 2016	+	-	?	+	+	?	+
	Molvarec 2013	?	+	+	+	-	+	+
	Nice 2016	?	+	+	+	+	+	+
	Shawkat 2015	+	+	+	+	-	+	+
	Valino 2016	+	+	?	+	+	?	+
		<div> High Unclear Low </div>						

Figure 19. Risk of bias and applicability concerns summary: review authors' judgements about each domain for each included study of uric acid.

	<u>Risk of Bias</u>				<u>Applicability Concerns</u>		
	Patient Selection	Index Test: Uric acid	Reference Standard	Flow and Timing	Patient Selection	Index Test: Uric acid	Reference Standard
Amini 2014	+	?	+	+	+	+	+
Bellomo 2011	+	?	?	?	-	+	+
Hawkins 2012	+	+	+	+	-	?	+
Jauniaux 1996	+	+	?	+	?	+	+
Odendaal 1997	+	+	?	+	-	+	+
Redman 1976	+	+	?	+	-	+	+
Voto 1988	+	?	?	+	-	+	+
Williams 2002	+	+	?	+	-	+	+
Yassaee 2003	+	?	?	?	-	-	?

High
Unclear
Low

Figure 20. Risk of bias and applicability concerns summary: review authors' judgements about each domain for each included study of urinary oestriol (UE3).

	<u>Risk of Bias</u>				<u>Applicability Concerns</u>		
	Patient Selection	Index Test: Urinary E3	Reference Standard	Flow and Timing	Patient Selection	Index Test: Urinary E3	Reference Standard
Beischer 1991	+	+	?	+	+	-	+
Campbell 1972	+	+	+	+	-	-	-
Chew 1976	+	+	+	+	-	-	+
Elliott 1970	+	+	+	+	-	-	+
Fliegner 1979	+	+	+	+	+	+	+
Kunz 1976	+	+	?	+	-	-	+
Oats 1979	+	+	?	+	+	-	+
Odendaal 1981	+	+	+	?	-	+	+
Steiner 1991	+	?	+	?	-	?	+
Weerasinghe 1977	+	+	?	+	-	-	-

High
Unclear
Low

Appendix 6. Within study comparisons of tests for identifying small-for-gestational age (birthweight $\leq$ 10th centile) infants

Study	Sensitivity (true positives/cases)		Difference (95% CI)	P value	Specificity (true negatives/non-cases)		Difference (95% CI)	P value
Estimated fetal weight (EFW) versus placental grading								
	EFW	Placental grad- ing			EFW	Placental grad- ing		
Geerts 2016	0.57 (34/60)	0.41 (21/51)	0.15 (-0.03 to 0.34)	P = 0.13	1.00 (150/150)	0.96 (132/137)	0.04 (0.01 to 0.07)	P = 0.02
Estimated fetal weight versus placental growth factor (PIGF)								
	EFW	PIGF			EFW	PIGF		
Valino 2016	(104/379)	(48/379)	0.15 (0.09 to 0.20)	P < 0.0001	(3519/3574)	(3424/3574)	0.03 (0.02 to 0.03)	P < 0.0001
Human placental lactogen (hPL) versus placental grading								
	hPL	Placental grad- ing			hPL	Placental grad- ing		
Altmann 1978	0.50 (3/6)	0.67 (4/6)	-0.17 (-0.72 to 0.38)	P = 1.00	0 (0/4)	0 (0/4)	0 (– to –)	–
Human placental lactogen versus placental growth factor								
	hPL	PIGF			hPL	PIGF		
Nice 2016	0.43 (10/23)	0.17(4/23)	0.26 (0.01 to 0.52)	P = 0.11	0.91 (49/54)	0.92 (49/53)	-0.02 (-0.12 to 0.09)	P = 1.0.
Human placental lactogen versus urinary oestriol (UE3)								
	hPL	UE3			hPL	UE3		
Kunz 1976	0.53 (8/15)	0.67 (10/15)	-0.13 (-0.48 to 0.21)	P = 0.71	0.76 (52/68)	0.82 (56/68)	-0.06 (-0.19 to 0.08)	P = 0.53
Odendaal 1981	0.81 (35/43)	0.21 (6/28)	0.60 (0.41 to 0.79)	P < 0.0001	0.53 (18/34)	0.94 (17/18)	-0.42 (-0.61 to -0.22)	P = 0.002
Steiner 1991	0.37 (25/68)	0.13 (9/68)	0.24 (0.10 to 0.38)	P = 0.003	0.62 (28/45)	0.62 (28/45)	0 (-0.20 to 0.20)	P = 1.0

(Continued)

Serum oestriol (E3) versus urinary oestriol

	E3	UE3			E3	UE3		
Odendaal 1981	0.43 (13/30)	0.21 (6/28)	0.22 (-0.01 to 0.45)	P = 0.10	0.87 (20/23)	0.94 (17/18)	-0.07 (-0.25 to 0.10)	P = 0.62

Differences in sensitivities and specificities between tests evaluated within each study are presented in the table. Five studies that evaluated hPL and oestriol were not included in this table as meta-analysis was performed (see Figure 10). The three studies of hPL versus urinary oestriol used different thresholds (see Figure 5).

Appendix 7. Studies with evaluations of both small-for-gestational age (birthweight $\leq$ 10th centile) infants and stillbirths

Estimates of sensitivity and specificity from studies that evaluated biochemical tests for both SGA infants and stillbirths (Figure 21)

Figure 21. Forest plot of studies with evaluations of both small-for-gestational-age (SGA birthweight $\leq$ tenth centile) infants and stillbirth. E3 = oestriol; hPL = human placental lactogen; PIGF = placental growth factor; UA = uric acid. Two studies (Altmann 1978; Chen 2012) evaluated placental grading for both SGA and stillbirth but not possible to include them on the plot. Studies are sorted by test and study identifier.

SGA data from studies with both SGA infants and stillbirths

Stillbirth data from studies with both SGA infants and stillbirths

WHAT'S NEW

Date	Event	Description
7 June 2019	Amended	Corrected typographical error in abstract background.

CONTRIBUTIONS OF AUTHORS

AEP Heazell (AEPH) and Clare Davenport (CD) conceived the idea for the systematic review. All authors contributed to the design of the review and writing the protocol. Susan Bayliss (SEB) undertook the literature searches. AEPH, Dexter Hayes (DH) and Melissa Whitworth (MKW) screened the titles and abstracts and extracted data from included studies. Yemisi Takwoingi (YT) performed the statistical analysis. All authors contributed to the final manuscript. AEPH is the guarantor for the review.

DECLARATIONS OF INTEREST

Susan Bayliss: none known.

Clare Davenport's employer (The University of Birmingham) received funding for her participation in this review as part of an NIHR clinical fellowship awarded to Alexander Heazell (the lead author and contact person).

Dexter Hayes: none known.

Yemisi Takwoingi's employer (The University of Birmingham) received funding for her participation in this review as part of an NIHR clinical fellowship awarded to Alexander Heazell (the lead author and contact person).

Melissa Whitworth was on the NICE guideline development group for Intrapartum Care in High Risk Pregnancy and travel to committee meetings was paid by NICE

Alexander Heazell has received research grants from Alere (UK) and Action Medical Research to investigate placental factors in maternal serum in women with reduced fetal movements. He is also a Supervisor for a Clinical Research Fellowship from Action Medical Research which incorporates projects to detect placental factors in maternal serum. In addition, he holds a Clinician Scientist Award from National Institute of Health Research (NIHR) (CS-2013-13-009) and this review is part of that programme of work. The views expressed here are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

SOURCES OF SUPPORT

Internal sources

- No sources of support supplied

External sources

- National Institute of Health Research, UK.

Alexander Heazell is funded by a Clinician Scientist Award from the National Institute of Health Research (CS-2013-009). This protocol presents independent research funded by the National Institute for Health Research (NIHR). The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

There are some minor differences between our published protocol ([Heazell 2016](#)) and this review.

- **Methods/investigation of heterogeneity:** in our protocol we stated "Where a sufficient number of studies assess the same index test, potential sources of heterogeneity will be separated into clinical (e.g. population studied, test type) and methodological (as appropriate) sources". In the review, we clarified this and edited the methods to "Where a sufficient number of studies assessed the same index test and there were at least four studies per subgroup of a categorical covariate, we performed meta-regression by adding the potential source of heterogeneity as a covariate to the HSROC model".
- **Statistical analysis and data synthesis**
 - * We clarified the use of the terms SGA (for SGA defined as birthweight $\leq$ 10th centile) and SGA3 for SGA defined as birthweight $<$ 3rd centile.
 - * We added "Before performing meta-analyses to compare test accuracy, we performed meta-analysis of each test separately for preliminary investigation of the shape of the SROC curve of each test and to explore if assuming common variances across tests for the random effects would be reasonable".
 - * We added "When there were adequate data, we also allowed the variance parameters for accuracy and threshold to depend on test type, i.e. differences in accuracy and threshold modelled as random effects"
 - * We also added "To quantify differences in accuracy between tests, we computed ratios of diagnostic odds ratios when SROC curves were symmetric or a common shape was assumed. Using the estimate statement within NLMIXED, we also estimated sensitivities along the SROC curves at fixed values of specificity that correspond to the median and interquartile range of specificities from the studies included in the comparative meta-analysis. We used these values along with the median and interquartile range of the

prevalence estimated from the studies to compute numbers of missed cases and false positives in a hypothetical cohort of 1000 pregnant women. We used these frequencies to illustrate the accuracy of the tests in absolute terms"

INDEX TERMS

Medical Subject Headings (MeSH)

*Infant, Small for Gestational Age; *Pregnancy Outcome; *Ultrasonography, Prenatal; Fetal Development [*physiology]; Fetal Growth Retardation; Perinatal Mortality; Placenta [*diagnostic imaging]; Stillbirth

MeSH check words

Female; Humans; Infant, Newborn; Pregnancy