Cochrane
Library

Cochrane Database of Systematic Reviews

Treatment of epilepsy for people with Alzheimer's disease (Review)

Liu J, Wang LN, Wu LY, Wang YP

Liu J, Wang LN, Wu LY, Wang YP.
Treatment of epilepsy for people with Alzheimer's disease.
Cochrane Database of Systematic Reviews 2018, Issue 12. Art. No.: CD011922.
DOI: [10.1002/14651858.CD011922.pub3](https://doi.org/10.1002/14651858.CD011922.pub3).

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	3
BACKGROUND	8
OBJECTIVES	8
METHODS	8
RESULTS	10
Figure 1.	11
Figure 2.	12
Figure 3.	13
DISCUSSION	14
AUTHORS' CONCLUSIONS	15
ACKNOWLEDGEMENTS	15
REFERENCES	16
CHARACTERISTICS OF STUDIES	17
DATA AND ANALYSES	20
Analysis 1.1. Comparison 1 Levetiracetam versus lamotrigine, Outcome 1 Proportion with seizure freedom.	20
Analysis 1.2. Comparison 1 Levetiracetam versus lamotrigine, Outcome 2 Reduction in seizure frequency of 50% or more.	20
Analysis 1.3. Comparison 1 Levetiracetam versus lamotrigine, Outcome 3 Proportion with adverse events.	20
Analysis 2.1. Comparison 2 Levetiracetam versus phenobarbital, Outcome 1 Proportion with seizure freedom.	21
Analysis 2.2. Comparison 2 Levetiracetam versus phenobarbital, Outcome 2 Reduction in seizure frequency of 50% or more. ..	21
Analysis 2.3. Comparison 2 Levetiracetam versus phenobarbital, Outcome 3 Proportion with adverse events.	22
Analysis 3.1. Comparison 3 Lamotrigine versus phenobarbital, Outcome 1 Proportion with seizure freedom.	22
Analysis 3.2. Comparison 3 Lamotrigine versus phenobarbital, Outcome 2 Reduction in seizure frequency of 50% or more.	22
Analysis 3.3. Comparison 3 Lamotrigine versus phenobarbital, Outcome 3 Proportion with adverse events.	23
APPENDICES	23
WHAT'S NEW	24
CONTRIBUTIONS OF AUTHORS	24
DECLARATIONS OF INTEREST	25
SOURCES OF SUPPORT	25
INDEX TERMS	25

[Intervention Review]

Treatment of epilepsy for people with Alzheimer's disease

Jia Liu¹, Lu-Ning Wang², Li-Yong Wu¹, Yu-Ping Wang¹¹Department of Neurology, Xuanwu Hospital, Capital Medical University, Beijing, China. ²Department of Geriatric Neurology, Chinese PLA General Hospital, Beijing, China**Contact address:** Jia Liu, Department of Neurology, Xuanwu Hospital, Capital Medical University, Changchun Street 45, Beijing, 100053, China. Jason_liu1984@163.com.**Editorial group:** Cochrane Epilepsy Group**Publication status and date:** New search for studies and content updated (no change to conclusions), published in Issue 12, 2018.**Citation:** Liu J, Wang LN, Wu LY, Wang YP. Treatment of epilepsy for people with Alzheimer's disease. *Cochrane Database of Systematic Reviews* 2018, Issue 12. Art. No.: CD011922. DOI: [10.1002/14651858.CD011922.pub3](https://doi.org/10.1002/14651858.CD011922.pub3).

Copyright © 2018 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Any type of seizure can be observed in Alzheimer's disease (AD). Antiepileptic drugs seem to prevent the recurrence of epileptic seizures in most people with AD. There are pharmacological and non-pharmacological treatments for epilepsy in people with AD. There are no current systematic reviews to evaluate the efficacy and tolerability of these treatments; this review aims to review those different modalities. This is an updated version of the original Cochrane Review published in Issue 11, 2016.

Objectives

To assess the efficacy and tolerability of pharmacological or non-pharmacological interventions for the treatment of epilepsy in people with AD (including sporadic AD and dominantly inherited AD).

Search methods

For the latest update, on 10 July 2018 we searched the Cochrane Register of Studies (CRS Web), which includes the Cochrane Epilepsy Group's Specialized Register and the Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE (Ovid 1946-), [ClinicalTrials.gov](https://www.clinicaltrials.gov), and the [WHO International Clinical Trials Registry Platform](https://www.who.int/clinicaltrialsregistryplatform) (ICTRP). In an effort to identify further published, unpublished and ongoing trials, we searched ongoing trials registers, reference lists and relevant conference proceedings, and contacted authors and pharmaceutical companies.

Selection criteria

We included randomized and quasi-randomized controlled trials investigating treatment for epilepsy in people with AD, with the outcomes of proportion of participants with seizure freedom or proportion of participants experiencing adverse events.

Data collection and analysis

Two review authors independently screened the titles and abstracts of identified records, selected studies for inclusion, extracted data, cross-checked the data for accuracy and assessed the methodological quality. We performed no meta-analyses due to the limited available data.

Main results

We included one randomized controlled trial on pharmacological interventions with 95 participants. No studies were found for non-pharmacological interventions. Concerning the proportion of participants with seizure freedom, no significant differences were found for the comparisons of levetiracetam (LEV) versus lamotrigine (LTG) (risk ratio (RR) 1.20, 95% confidence interval (CI) 0.53 to 2.71), LEV versus phenobarbital (PB) (RR 1.01, 95% CI 0.47 to 2.19), or LTG versus PB (RR 0.84, 95% CI 0.35 to 2.02). It seemed that LEV could improve cognition

and LTG could relieve depression, while PB and LTG could worsen cognition, and LEV and PB could worsen mood. Unclear risk of bias was found in allocation, blinding and selective reporting. We judged the quality of the evidence to be very low.

Authors' conclusions

This review does not provide sufficient evidence to support LEV, PB or LTG for the treatment of epilepsy in people with AD. Regarding efficacy and tolerability, no significant differences were found between LEV, PB and LTG. Large randomized controlled trials with a double-blind, parallel-group design are required to determine the efficacy and tolerability of treatment for epilepsy in people with AD.

PLAIN LANGUAGE SUMMARY

Treatment of epilepsy for people with Alzheimer's disease

Background

Alzheimer's disease is a risk factor for increased seizures in older people. Seizures of any type can be observed in Alzheimer's disease and are probably underestimated.

Study characteristics

We searched scientific databases for clinical trials comparing medication and non-medication-based treatments for epilepsy in people with Alzheimer's disease. We wanted to evaluate how well the treatment worked and if it had any side effects.

Key results

We included and analyzed one randomized controlled trial (a clinical study where people are randomly put into one or two or more treatment groups) with 95 participants. Concerning the proportion of participants with seizure freedom, no significant differences were found between the antiepileptic drugs (levetiracetam versus lamotrigine, levetiracetam versus phenobarbital, and lamotrigine versus phenobarbital). It seemed that levetiracetam could improve cognition (thinking) and lamotrigine could relieve depression; while phenobarbital and lamotrigine could worsen cognition, and levetiracetam and phenobarbital could worsen mood.

Quality of the evidence

The quality of the evidence from the study was very low and results should be interpreted with caution. Large randomized controlled trials, which have a double-blind parallel-group design, are required to determine how effective and well tolerated treatments are for epilepsy in people with Alzheimer's disease.

The evidence is current to July 2018.

SUMMARY OF FINDINGS

Summary of findings for the main comparison. Levetiracetam compared with lamotrigine for the treatment of epilepsy in people with Alzheimer's disease

Levetiracetam compared with lamotrigine for the treatment of epilepsy in people with Alzheimer's disease

Patient or population: people with AD and epileptic seizures

Settings: the community of Caltanissetta, Italy

Intervention: levetiracetam

Comparison: lamotrigine

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	lamotrigine	levetiracetam				
Proportion with seizure freedom	241 per 1000	289 per 1000 (128 to 653)	RR 1.20 (0.53 to 2.71)	67 (1 study)	⊕⊕⊕⊕ very low ¹	RR more than 1 favours levetiracetam
Proportion with adverse events	241 per 1000	158 per 1000 (60 to 419)	RR 0.65 (0.25 to 1.74)	67 (1 study)	⊕⊕⊕⊕ very low ¹	RR less than 1 favours levetiracetam
Reduction in seizure frequency of 50% or more	586 per 1000	711 per 1000 (492 to 1026)	RR 1.21 (0.84 to 1.75)	67 (1 study)	⊕⊕⊕⊕ very low ¹	RR more than 1 favours levetiracetam
Proportion with seizure freedom in dominantly inherited AD	Not reported	Not reported				
Change in cognition	See comment	See comment		67 (1 study)	⊕⊕⊕⊕ very low ¹	In the lamotrigine group, participants showed slight declines in MMSE and ADAS-Cog scores. In the levetiracetam group, MMSE scores reflected improvement by a mean of 0.23 points as compared with baseline. Similar improvement was observed in ADAS-Cog scores (−0.23).

Change in neuropsychiatric symptoms	See comment	See comment	67 (1 study)	⊕⊕⊕⊕ very low ¹	In the lamotrigine group, score change of -0.72 was recorded on the Cornell scale. In the levetiracetam group, mood score worsened with 0.20 points.
Improvement in quality of life	Not reported	Not reported			

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes.² The **corresponding risk** (and its 95% CI) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

AD: Alzheimer's disease; **ADAS-Cog:** Alzheimer's Disease Assessment Scale-Cognitive; **CI:** confidence interval; **MMSE:** Mini Mental State Examination; **RR:** risk ratio

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ Only one trial was included, with 67 randomized participants and potential methodological insufficiency.

² Assumed risk is calculated as the event rate in the control group per 1000 people (number of events divided by the number of participants receiving control treatment).

Summary of findings 2. Levetiracetam compared with phenobarbital for the treatment of epilepsy in people with Alzheimer's disease

Levetiracetam compared with phenobarbital for the treatment of epilepsy in people with Alzheimer's disease

Patient or population: patients with AD and epileptic seizures

Settings: the community of Caltanissetta, Italy

Intervention: levetiracetam

Comparison: phenobarbital

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	phenobarbital	levetiracetam				
Proportion with seizure freedom	286 per 1000	289 per 1000 (134 to 626)	RR 1.01 (0.47 to 2.19)	66 (1 study)	⊕⊕⊕⊕ very low ¹	RR more than 1 favours levetiracetam

Proportion with adverse events	357 per 1000	158 per 1000 (64 to 382)	RR 0.44 (0.18 to 1.07)	66 (1 study)	⊕⊕⊕⊕ very low ¹	RR less than 1 favours levetiracetam
Reduction in seizure frequency of 50% or more	643 per 1000	711 per 1000 (502 to 1003)	RR 1.11 (0.78 to 1.56)	66 (1 study)	⊕⊕⊕⊕ very low ¹	RR more than 1 favours levetiracetam
Proportion with seizure freedom in dominantly inherited AD	Not reported	Not reported				
Change in cognition	See comment	See comment		66 (1 study)	⊕⊕⊕⊕ very low ¹	In the phenobarbital group, significant worsening of cognitive performance was found with lower mean scores indicating aggravation of existing cognitive impairment at both 6 and 12 months post-randomization on MMSE and ADAS-Cog. In the levetiracetam group, MMSE scores reflected improvement by a mean of 0.23 points as compared with baseline. Similar improvement was observed in ADAS-Cog scores (-0.23).
Change in neuropsychiatric symptoms	See comment	See comment		66 (1 study)	⊕⊕⊕⊕ very low ¹	In the phenobarbital group, mood score worsened with 1.74 points. In levetiracetam group, mood score worsened with 0.20 points.
Improvement in quality of life	Not reported	Not reported				

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes.² The **corresponding risk** (and its 95% CI) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

AD: Alzheimer's disease; **ADAS-Cog:** Alzheimer's Disease Assessment Scale-Cognitive; **CI:** confidence interval; **MMSE:** Mini Mental State Examination; **RR:** risk ratio

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ Only one trial was included, with 66 randomized participants and potential methodological insufficiency.

² Assumed risk is calculated as the event rate in the control group per 1000 people (number of events divided by the number of participants receiving control treatment).

Summary of findings 3. Lamotrigine compared with phenobarbital for the treatment of epilepsy in people with Alzheimer's disease

Lamotrigine compared with phenobarbital for the treatment of epilepsy in people with Alzheimer's disease

Patient or population: patients with AD and epileptic seizures

Settings: the community of Caltanissetta, Italy

Intervention: lamotrigine

Comparison: phenobarbital

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	phenobarbital	lamotrigine				
Proportion with seizure freedom	286 per 1000	241 per 1000 (100 to 578)	RR 0.84 (0.35 to 2.02)	57 (1 study)	⊕⊕⊕⊕ very low ¹	RR less than 1 favours phenobarbital
Proportion with adverse events	357 per 1000	241 per 1000 (107 to 546)	RR 0.68 (0.30 to 1.53)	57 (1 study)	⊕⊕⊕⊕ very low ¹	RR less than 1 favours lamotrigine
Reduction in seizure frequency of 50% or more	643 per 1000	586 per 1000 (386 to 887)	RR 0.91 (0.60 to 1.38)	57 (1 study)	⊕⊕⊕⊕ very low ¹	RR less than 1 favours phenobarbital
Proportion with seizure freedom in dominantly inherited AD	Not reported	Not reported				
Change in cognition	See comment	See comment		57 (1 study)	⊕⊕⊕⊕ very low ¹	In the phenobarbital group, significant worsening of cognitive performance was found, with lower mean scores indicating aggravation of existing cognitive impairment at both 6 and 12 months post-randomization on MMSE and ADAS-Cog. In the lamotrigine group, participants showed slight declines in MMSE and ADAS-Cog scores.
Change in neuropsychiatric symptoms	See comment	See comment		57 (1 study)	⊕⊕⊕⊕ very low ¹	In the phenobarbital group, mood score worsened with 1.74 points. In the lamotrigine group, score change of -0.72 was recorded on the Cornell scale.

Improvement in quality of life	Not reported	Not reported
--------------------------------	--------------	--------------

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes.² The **corresponding risk** (and its 95% CI) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

AD: Alzheimer's disease; **ADAS-Cog:** Alzheimer's Disease Assessment Scale-Cognitive; **CI:** confidence interval; **MMSE:** Mini Mental State Examination; **RR:** risk ratio

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ Only one trial was included, with 57 randomized participants and potential methodological insufficiency.

² Assumed risk is calculated as the event rate in the control group per 1000 people (number of events divided by the number of participants receiving control treatment).

BACKGROUND

Description of the condition

Epilepsy is a common neurological disorder and becomes more frequent with age (Brodie 2009). Meanwhile, Alzheimer's disease (AD) is the most common neurodegenerative disease in older people, and is characterized by memory loss, cognitive decline, and behavioral disorders. Although epilepsy is not the predominant symptom in sporadic AD, it is more common in autosomal dominant AD (Wu 2012). It has been estimated that AD is a risk factor for increased seizures in older people (Pandis 2012). Approximately 10% to 22% of people with AD have at least one unprovoked seizure (Mendez 2003). Any type of seizure can be observed in AD (Rao 2009). The prevalence of epilepsy is probably underestimated (Tallis 2002), considering the unrecognised non-convulsive forms. Seizures can be seen even in the early stages of AD (Palop 2007), which suggests seizures may contribute to cognitive impairment.

Description of the intervention

Antiepileptic drugs (AEDs) are the current intervention for epilepsy treatment in people with AD. According to a previous study, the efficacy of AEDs in older people was proven to be better than that in the younger population (Mattson 1985). The first generation AEDs, such as valproic acid and benzodiazepines, can aggravate cognitive decline in people with AD (Fleisher 2011; Wu 2009). In contrast, the new generation AEDs, e.g. lamotrigine and gabapentin, seem to be well tolerated by older people (Rowan 2005; Saetre 2007). Drugs targeting beta amyloid may also be a rational choice for treatment of epilepsy in AD as beta amyloid accumulation can contribute to seizures, as confirmed by Down syndrome and AD mouse models (Puri 2001; Westmark 2008). Non-pharmacological interventions such as transcranial magnetic stimulation (TMS), deep brain stimulation and acupuncture are potentially beneficial for people with AD or people with epilepsy (Hsu 2015; Kimiskidis 2010; Laxpati 2014; McElroy-Cox 2009).

How the intervention might work

The common mechanisms between AD and epilepsy can probably be attributed to hippocampal sclerosis; that is, neuron loss and astrogliosis in cornu ammonis with sparing of granule cells in the dentate nucleus (Chin 2013). Furthermore, hippocampal sclerosis may be associated with dementia and neuropsychiatric symptoms in people with AD or epilepsy. In AD mouse models, a-beta amyloid and neurofibrillary tangle-related tau protein have been suggested to contribute to seizures (Roberson 2011; Westmark 2008). Some studies have demonstrated that seizures occur even earlier than amyloid pathology, which possibly causes the progressive neurodegeneration in AD (Minkeviciene 2009; Palop 2007). Thus, AEDs may reduce the progression of AD by controlling epilepsy, while drugs targeting beta amyloid may reduce seizures by preventing amyloid aggregation. With regard to the mechanisms of TMS, either excitatory or inhibitory responses can be generated in cortical tissues, which improve the symptoms of both AD and epilepsy.

Why it is important to do this review

At present, AED therapy has been tested in clinical trials and seems to prevent the recurrence of epileptic seizures in most people with AD (Rao 2009). Meanwhile, drugs targeting beta amyloid, such as bapineuzumab and solanezumab, have not been widely applied

in the treatment of AD. Therefore, their antiepileptic efficacy is still unknown. TMS treatment has been applied in AD and in epilepsy, but the efficacy of TMS for epilepsy in people with AD is rarely reported. In this review, we aim to evaluate the efficacy and tolerability of the treatment of epilepsy for people with AD. To our knowledge, no systematic review or meta-analysis on this topic exists.

OBJECTIVES

To assess the efficacy and tolerability of pharmacological or non-pharmacological interventions for the treatment of epilepsy in people with AD (including sporadic AD and dominantly inherited AD).

METHODS

Criteria for considering studies for this review

Types of studies

We included randomized and quasi-randomized controlled trials.

Types of participants

We included people diagnosed with Alzheimer's disease (AD) combined with epilepsy. We used the definitions of AD and epilepsy as provided in the original studies. We applied no limitations on gender or age.

Types of interventions

Experimental intervention: any pharmacological or non-pharmacological intervention for epilepsy, alone or combined with other treatment.

Control intervention: no treatment; placebo alone or combined with other treatment (concomitant interventions had to be the same in each group); different doses of the experimental intervention.

Types of outcome measures

The outcomes of all the participants initially randomized were collected and subjected to intention-to-treat (ITT) analysis. All the outcomes were measured at the endpoint, in comparison with the data from the baseline. We followed the endpoints as provided in the original publications, and merged the data at the same time point, if possible.

Primary outcomes

1. Proportion of participants with seizure freedom
2. Proportion of participants who experienced any adverse events (AEs)

Secondary outcomes

1. Proportion of participants with dominantly inherited AD with seizure freedom.
2. Reduction in seizure frequency of 50% or more.
3. Change in cognition, measured by Mini Mental State Examination (MMSE), Alzheimer's Disease Assessment Scale-Cognitive (ADAS-Cog), etc.

4. Change in neuropsychiatric symptoms, measured by Neuropsychiatric Inventory (NPI), Geriatric Depression Scale (GDS), etc.
5. Improvement in quality of life, measured by Quality of Life Scale, etc.

Search methods for identification of studies

Electronic searches

Searches were run for the original review on 1 February 2016. Subsequent searches were run on 10 July 2018. For the latest update we searched the following databases:

1. Cochrane Register of Studies (CRS Web, 10 July 2018), which includes the Cochrane Epilepsy Group's Specialized Register and the Cochrane Central Register of Controlled Trials (CENTRAL), using the search strategy set out in [Appendix 1](#);
2. MEDLINE (Ovid, 1946-), using the search strategy set out in [Appendix 2](#);
3. [ClinicalTrials.gov](#) (10 July 2018), using the search strategy set out in [Appendix 3](#);
4. [WHO International Clinical Trials Registry Platform](#) (ICTRP, 10 July 2018), using the search strategy set out in [Appendix 4](#).

Note: it is no longer necessary to search Embase, because randomized and quasi-randomized controlled trials from Embase are now included in CENTRAL.

Searching other resources

We also:

1. searched reference lists of reviews and retrieved articles for additional studies;
2. searched conference proceedings from the last three years for relevant studies, in World Congress of Neurology, International Conference of Alzheimer's Disease, Alzheimer's Association International Conference, and International Epilepsy Congress;
3. contacted researchers, pharmaceutical companies, and relevant trial authors to seek information about unpublished or incomplete trials.

We did not impose any language limitations for the search, and we attempted to obtain translations of articles where necessary.

Data collection and analysis

Selection of studies

Two review authors (JL, L-NW) independently evaluated titles and abstracts of identified trials to determine if they fulfilled the inclusion criteria. All potentially relevant studies were obtained in full text for further consideration. We listed the excluded studies and reported the reasons for exclusion. Any disagreements were resolved by discussion or by an independent party (L-YW) if necessary.

Data extraction and management

Two review authors (JL, L-NW) independently extracted eligible data from the published reports onto standardized forms, and cross-checked them for accuracy. Disagreements regarding data extraction were resolved by discussion between the review authors.

We used checklists to independently record details of the following:

1. study design;
2. total study duration;
3. methods of generating randomization schedule;
4. method of concealment of allocation;
5. blinding;
6. use of an ITT analysis (all participants initially randomized will be included in the analyses as allocated to groups);
7. AEs and dropouts for all reasons;
8. participants (country, number of participants, age, gender, inclusion and exclusion criteria);
9. comparison (details of the intervention in treatment and control groups, details of co intervention(s) in both groups, duration of treatment);
10. outcomes and time points of measures (number of participants in each group and outcome, regardless of compliance);
11. factors for heterogeneity (sample size, missing participants, confidence interval (CI) and P value in measurement, subgroup analyses).

Assessment of risk of bias in included studies

Two review authors (JL, L-NW) independently assessed the risk of bias for each study using the criteria outlined in the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011](#)). We resolved any disagreements by discussion or by involving an independent party (L-YW). We assessed the risk of bias according to the following domains.

1. Random sequence generation
2. Allocation concealment
3. Blinding of participants and personnel
4. Blinding of outcome assessment
5. Incomplete outcome data
6. Selective outcome reporting
7. Other bias

We assessed the risk of bias for each domain as high, low or unclear risk and provided information from the study report. We described sources of bias.

Measures of treatment effect

We expressed continuous data as mean differences (MDs) with 95% confidence intervals (CIs). Standardized mean difference (SMD) would have been used for the outcomes when they were measuring the same thing but in different ways. For all dichotomous outcomes we calculated risk ratios (RRs), again with 95% CIs. As it is possible that some trials (or groups within a trial) had no adverse events or no dropouts, we would have calculated risk differences (RDs) instead of RRs in these specific situations, again with 95% CIs. We used ITT analysis for all the outcomes. Different control groups were analysed separately.

Unit of analysis issues

We planned to deal with any unit of analysis issues according to the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011](#)).

Dealing with missing data

We planned to contact the authors of the studies for further details if any data were missing, and to establish the characteristics of unpublished trials through correspondence with the trial co-ordinator or principal investigator. According to the ITT principle, all randomized participants should be included.

Assessment of heterogeneity

We planned to use the standard χ^2 statistic and I^2 statistic to measure the heterogeneity (Higgins 2011), and make a judgement along with visual inspection of forest plots. For the χ^2 test, we would have rejected the hypothesis of tolerability if the P value was less than 0.10, and we would have interpreted an I^2 greater than 50% as representing significant heterogeneity. In this case, we would have tried to explore factors for heterogeneity.

Assessment of reporting biases

We planned to assess publication bias by funnel plot if we had found more than 10 studies. However, the review included only one study.

Data synthesis

If we found neither clinical nor statistical heterogeneity, we planned to pool results using a fixed-effect model. We would have analyzed different controls separately. In the case of statistical heterogeneity, we planned to pool the results using a random-effects model. For heterogeneity that could not be readily explained, we would not have pooled the data but only given a description of the results.

Subgroup analysis and investigation of heterogeneity

We planned to analyze subgroups of studies categorized according to the subtype of AD (sporadic AD and dominantly inherited AD), stage of AD (mild, moderate or severe AD), and the dosage and duration of interventions.

As a formal method of comparing subgroups, we planned to use the χ^2 test (to test for significant differences between subgroups of participants). For all statistical analyses, we used Review Manager 5 (RevMan 2014).

However, we did not perform subgroup analyses due to the limited available data.

Sensitivity analysis

We planned to use best-case and worst-case scenarios for taking into account missing data. We also planned to undertake the following sensitivity analyses to investigate unexplained heterogeneity and to test the robustness of results:

1. exclusion of cross-over trials from the analysis;
2. exclusion of studies at high risk of bias, with inadequate allocation concealment or lack of blinded outcome assessor;
3. comparison of fixed-effect versus random-effects models.

However, we were unable to perform any of these analyses due to the limited available data.

Summary of findings and quality of the evidence (GRADE)

We have presented three 'Summary of findings' tables, one for each comparison (Summary of findings for the main comparison; Summary of findings 2; Summary of findings 3). We reported all outcomes in the tables.

We determined the quality of the evidence using the GRADE approach (Schunemann 2011; GRADEPro 2004); and downgraded evidence in the presence of a high risk of bias in at least one study, indirectness of the evidence, unexplained heterogeneity or inconsistency, imprecision of results, and high probability of publication bias. We downgraded evidence by one level if we considered the limitation to be serious and by two levels if very serious.

RESULTS

Description of studies

Results of the search

After searching the electronic databases and excluding duplicates, we identified 247 references (Figure 1). After screening the titles and abstracts, we obtained the full papers of four studies and assessed them for eligibility. We excluded three studies, one because it had a non-randomized design (Campion 1995) and two because the participants were ineligible (Lovestone 2015; Musaeus 2017). The previous ongoing randomized controlled trial (RCT) (NCT01554683) was completed and released as Musaeus 2017, which we excluded as described. There are two ongoing RCTs (NCT02002819; NCT03489044).

Figure 1. Study flow diagram.

Included studies

One RCT with 95 participants met the inclusion criteria.

A total of 95 participants with Alzheimer's disease (AD) and epileptic seizures (41 males, 54 females) were enrolled. The participants had a mean age of 71.75 years (range: 65 to 82 years) and a mean education of 6.3 years (range: 5 to 17 years), and lived in the community. They were randomly assigned with an antiepileptic drug (AED) as monotherapy: 38 were administered levetiracetam (LEV), 28 phenobarbital (PB), and 29 lamotrigine (LTG). This study comprised a four-week dose adjustment period followed by a

12-month dose evaluation period. During the treatment periods, participants received LEV, PB, or LTG and were titrated to an effective dose. No participant was previously exposed to another AED. The initial target dosage of LEV was 500 mg/day, increased weekly by 500 mg/day; for PB it was 50 mg/day, increased weekly by 50 mg/day; and for LTG it was 25 mg/day, increased weekly by 25 mg/day. Thereafter, the dosage was individually adjusted. In summary, the mean daily dose of LEV monotherapy was 956 mg (range: 500 to 2000 mg/day); the mean daily dose of PB monotherapy was 90 mg/day (range: 50 to 100 mg/day); and the mean daily dose of LTG monotherapy was 57.5 mg/day (range: 25 to

100 mg/day). Further details of the included studies are provided in the [Characteristics of included studies](#).

Excluded studies

We excluded three studies after full-text evaluation ([Campion 1995](#); [Lovestone 2015](#); [Musaeus 2017](#)). The reasons for exclusion

were due to non-randomized design or ineligible participants (see [Characteristics of excluded studies](#)).

Risk of bias in included studies

The information regarding risk of bias is provided in [Figure 2](#) and [Figure 3](#).

Figure 2. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.

Figure 3. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

Allocation

The included study, [Cumbo 2010](#), did not describe the method used to generate allocation sequence and whether the methods of allocation were concealed or not. Therefore we regarded it as having an unclear risk of selection bias.

Blinding

The details of blinding were not mentioned in the included study. Thus, we evaluated it as having an unclear risk of both performance bias and detection bias.

Incomplete outcome data

Eighty-three participants (87%) completed the study (35 (92%) on LEV, 23 (82%) on PB, and 25 (86%) on LTG). The reasons for dropouts were given. Therefore, we assessed the study as having a low risk of attrition bias.

Selective reporting

We intended to use the table of 'Outcome Reporting Bias In Trials' (ORBIT) to evaluate selective outcome reporting ([Kirkham 2010](#)). However, we could not assess reporting bias as no pre-published protocols were available.

Other potential sources of bias

We found no other potential sources of bias. Insufficient numbers of trials were available for a funnel plot analysis.

Effects of interventions

See: [Summary of findings for the main comparison](#) Levetiracetam compared with lamotrigine for the treatment of epilepsy in people with Alzheimer's disease; [Summary of findings 2](#) Levetiracetam compared with phenobarbital for the treatment of epilepsy in people with Alzheimer's disease; [Summary of findings 3](#) Lamotrigine compared with phenobarbital for the treatment of epilepsy in people with Alzheimer's disease

See: [Summary of findings for the main comparison](#); [Summary of findings 2](#); [Summary of findings 3](#). An efficacy and tolerability analysis was conducted of all participants who had taken at least one dose, i.e. intention-to-treat (ITT) population. People who discontinued the drug for any reason were considered non-responders with 0% difference from baseline seizure frequency and were included in the efficacy results. There was no difference between groups in demographic or baseline characteristics. All the outcomes were measured at the (12-month) endpoint.

Primary outcomes

1. Proportion of participants with seizure freedom

At 12 months, 11 of 38 participants (29%) in the LEV group had become seizure free; seven of the 11 were seizure free from the start of therapy, and the other four became seizure free after two months (with a relatively high dosage, 2000 mg/day). In the PB group, eight of 28 participants (29%) were seizure free from the start of therapy at a dosage of 100 mg/day. In the LTG group, elimination of seizures was achieved by seven of 29 (24%) participants ([Analysis 1.1](#); [Analysis 2.1](#); [Analysis 3.1](#)).

2. Proportion of participants who experienced any adverse events (AEs)

In the LEV group, AEs were observed in six (16%) participants. These AEs included somnolence, asthenia, headache and dizziness. No participant withdrew due to AEs. In the PB group, 10 (36%) experienced AEs. Somnolence and asthenia were the most frequently reported AEs. PB was discontinued by four participants due to AEs. In the LTG group, seven (24%) participants reported AEs. No participant withdrew because of AEs. There was no statistically significant difference ([Analysis 1.3](#); [Analysis 2.3](#); [Analysis 3.3](#)).

Secondary outcomes

1. Proportion of participants with dominantly inherited AD with seizure freedom

No data reported.

2. Reduction in seizure frequency of 50% or more

In the LEV group, 27 of 38 (71%) had a greater than 50% reduction in seizure frequency at 12 months follow-up. In the PB group, 18 of 28 (64%) had a greater than 50% reduction in seizure frequency. In the LTG group, a greater than 50% reduction in seizure frequency was observed in 17 of 29 (59%) participants. There was no statistically significant difference ([Analysis 1.2](#); [Analysis 2.2](#); [Analysis 3.2](#)).

3. Change in cognition

In the LEV group, Mini Mental State Examination (MMSE) scores recorded at the endpoint reflected improvement by a mean of 0.23 points as compared with baseline. Similar improvement was observed in ADAS-Cog scores (−0.23). LEV was particularly associated with improved attention, short-term memory, and oral fluency. In the PB group, significant worsening of cognitive performance was found, with lower mean scores indicating aggravation of existing cognitive impairment at both six and 12 months post-randomization on MMSE and ADAS-Cog. Participants in the LTG group showed slight declines in MMSE and ADAS-Cog scores. Due to the lack of mean differences and standard deviations provided, qualitative analyses were unavailable.

4. Change in neuropsychiatric symptoms

Participants in the LTG group scored better on measures of mood. They exhibited progressive improvement on the Cornell scale for depression from six months onward. At the endpoint, a score change of −0.72 was recorded on the Cornell scale. Mood scores for the LEV group (0.20) and PB group (1.74) worsened. Due to the lack of standard deviations provided, qualitative analyses were unavailable.

5. Improvement in quality of life

No data reported.

DISCUSSION

Summary of main results

We included one randomized controlled trial (RCT) with 95 participants. We found methodological defects; details are provided in the [Characteristics of included studies](#) section. We were unable to carry out meta-analysis. There were no significant differences reported between levetiracetam (LEV), phenobarbital (PB) and lamotrigine (LTG) groups, in terms of participants with seizure freedom, reduction in seizure frequency of 50% or more, and adverse events. It seemed that LEV could improve cognition and LTG could relieve depression; while PB and LTG could worsen cognition, and LEV and PB could worsen mood.

Overall completeness and applicability of evidence

The one included RCT on pharmacological interventions reported no significant differences between LEV, PB and LTG in the outcomes of efficacy and safety. There were 95 participants randomized, and there were 12 dropouts at the endpoint. All the participants had concomitant cholinesterase inhibitor therapy for AD. Other concomitant medications, including diuretics, antihypertensives, lipid-reducing agents, and antidiabetic drugs, started prior to the baseline visit were allowed during the study. No studies were found for non-pharmacological interventions. Large, well-designed, parallel-group RCTs on pharmacological or non-pharmacological interventions are required.

Quality of the evidence

Only one RCT comprising 95 randomized participants was included. We identified methodological issues in the study. Therefore, we were uncertain about the results. The study did not provide details of allocation sequence generation, allocation concealment, or blinding of participants and personnel. Considering only one trial was included with a small sample size and potential methodological insufficiency, we judged the quality of the evidence to be very low.

Potential biases in the review process

The search for trials was rigorously performed based on the strategies in different electronic databases. We found one eligible RCT and two ongoing studies. To identify unpublished or incomplete trials, we also searched for protocols, but found no eligible studies. It is possible that certain unpublished trials were not identified. In addition, due to the inclusion of only one RCT, we could not assess publication bias using funnel plots.

Agreements and disagreements with other studies or reviews

To our knowledge, this is the first review to systematically evaluate the efficacy and tolerability of treatment of epilepsy in people with AD.

AUTHORS' CONCLUSIONS

Implications for practice

This review does not provide sufficient evidence to support levetiracetam (LEV), phenobarbital (PB), or lamotrigine (LTG) for the treatment of epilepsy in people with Alzheimer's disease (AD). Regarding efficacy and tolerability, no significant differences were found between LEV, PB and LTG.

Implications for research

Large randomized, double-blind, controlled, parallel-group clinical trials are required to determine the efficacy and tolerability of treatment of epilepsy in people with AD.

ACKNOWLEDGEMENTS

The authors acknowledge the help provided by the [Cochrane Epilepsy Group](#).

This review update was supported by the National Institute for Health Research, via Cochrane Infrastructure funding to the Epilepsy Group. The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the Systematic Reviews Programme, NIHR, NHS or the Department of Health.

REFERENCES

References to studies included in this review

Cumbo 2010 {published data only}

Cumbo E, Ligori LD. Levetiracetam, lamotrigine, and phenobarbital in patients with epileptic seizures and Alzheimer's disease. *Epilepsy & Behavior* 2010;**17**(4):461-6.

References to studies excluded from this review

Campion 1995 {published data only}

Campion D, Brice A, Hannequin D, Tardieu S, Dubois B, Calenda A, et al. A large pedigree with early-onset Alzheimer's disease: clinical, neuropathologic, and genetic characterization. *Neurology* 1995;**45**(1):80-5.

Lovestone 2015 {published data only}

Lovestone S, Boada M, Dubois B, Hüll M, Rinne JO, Huppertz HJ, et al. A phase II trial of tideglusib in Alzheimer's disease. *Journal of Alzheimer's Disease* 2015;**45**(1):75-88.

Musaeus 2017 {published data only}

Musaeus CS, Shafi MM, Santarnecchi E, Herman ST, Press DZ. Levetiracetam alters oscillatory connectivity in Alzheimer's disease. *Journal of Alzheimer's Disease* 2017;**58**(4):1065-76.

References to ongoing studies

NCT02002819 {unpublished data only}

Levetiracetam for Alzheimer's Disease-Associated Network Hyperexcitability.

NCT03489044 {unpublished data only}

An Investigation of Levetiracetam in Alzheimer's Disease.

Additional references

Brodie 2009

Brodie MJ, Elder AT, Kwan P. Epilepsy in later life. *Lancet Neurology* 2009;**8**(11):1019-30.

Chin 2013

Chin J, Scharfman HE. Shared cognitive and behavioral impairments in epilepsy and Alzheimer's disease and potential underlying mechanisms. *Epilepsy & Behavior* 2013;**26**(3):343-51.

Fleisher 2011

Fleisher AS, Truran D, Mai JT, Langbaum JB, Aisen PS, Cummings JL, et al. Chronic divalproex sodium use and brain atrophy in Alzheimer disease. *Neurology* 2011;**77**(13):1263-71.

GRADEPro 2004 [Computer program]

Brozek J, Oxman A, Schunemann H. GRADEPro Version 3.6 for Windows. GRADE Working Group, 2004.

Higgins 2011

Higgins JPT, Green S (editors). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [update

March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Hsu 2015

Hsu WY, Ku Y, Zanto TP, Gazzaley A. Effects of noninvasive brain stimulation on cognitive function in healthy aging and Alzheimer's disease: a systematic review and meta-analysis. *Neurobiology of Aging* 2015;**36**(8):2348-59.

Kimiskidis 2010

Kimiskidis VK. Transcranial magnetic stimulation for drug-resistant epilepsies: rationale and clinical experience. *European Neurology* 2010;**63**(4):205-10.

Kirkham 2010

Kirkham JJ, Dwan KM, Altman DG, Gamble C, Dodd S, Smyth R, et al. The impact of outcome reporting bias in randomised controlled trials on a cohort of systematic reviews. *BMJ* 2010;**340**:c365.

Laxpati 2014

Laxpati NG, Kasoff WS, Gross RE. Deep brain stimulation for the treatment of epilepsy: circuits, targets, and trials. *Neurotherapeutics* 2014;**11**(3):508-26.

Lefebvre 2011

Lefebvre C, Manheimer E, Glanville J. Chapter 6: Searching for studies. In: Higgins JPT, Green S (editors). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from <http://handbook.cochrane.org/>.

Mattson 1985

Mattson RH, Cramer JA, Collins JF, Smith DB, Delgado-Escueta AV, Browne TR, et al. Comparison of carbamazepine, phenobarbital, phenytoin, and primidone in partial and secondarily generalized tonic-clonic seizures. *The New England Journal of Medicine* 1985;**313**(3):145-51.

McElroy-Cox 2009

McElroy-Cox C. Alternative approaches to epilepsy treatment. *Current Neurology and Neuroscience Reports* 2009;**9**(4):313-8.

Mendez 2003

Mendez M, Lim G. Seizures in elderly patients with dementia: epidemiology and management. *Drugs & Aging* 2003;**20**(11):791-803.

Minkeviciene 2009

Minkeviciene R, Rheims S, Dobszay MB, Zilberter M, Hartikainen J, Fülöp L, et al. Amyloid beta-induced neuronal hyperexcitability triggers progressive epilepsy. *The Journal of Neuroscience* 2009;**29**(11):3453-62.

Palop 2007

Palop JJ, Chin J, Roberson ED, Wang J, Thwin MT, Bien-Ly N, et al. Aberrant excitatory neuronal activity and compensatory remodeling of inhibitory hippocampal circuits in mouse models of Alzheimer's disease. *Neuron* 2007;**55**(5):697-711.

Pandis 2012

Pandis D, Scarmeas N. Seizures in Alzheimer disease: clinical and epidemiological data. *Epilepsy Currents* 2012;**12**(5):184-7.

Puri 2001

Puri BK, Ho KW, Singh I. Age of seizure onset in adults with Down's syndrome. *International Journal of Clinical Practice* 2001;**55**(7):442-4.

Rao 2009

Rao SC, Dove G, Cascino GD, Petersen RC. Recurrent seizures in patients with dementia: frequency, seizure types, and treatment outcome. *Epilepsy & Behavior* 2009;**14**(1):118-20.

RevMan 2014 [Computer program]

Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Roberson 2011

Roberson ED, Halabisky B, Yoo JW, Yao J, Chin J, Yan F, et al. Amyloid- β /Fyn-induced synaptic, network, and cognitive impairments depend on tau levels in multiple mouse models of Alzheimer's disease. *The Journal of Neuroscience* 2011;**31**(2):700-11.

Rowan 2005

Rowan AJ, Ramsay RE, Collins JF, Pryor F, Boardman KD, Uthman BM, et al. New onset geriatric epilepsy: a randomized study of gabapentin, lamotrigine, and carbamazepine. *Neurology* 2005;**64**(11):1868-73.

Saetre 2007

Saetre E, Perucca E, Isojarvi J, Gjerstad L, LAM 40089 Study Group. An international multicenter randomized double-blind controlled trial of lamotrigine and sustained-release carbamazepine in the treatment of newly diagnosed epilepsy in the elderly. *Epilepsia* 2007;**48**(7):1292-302.

Schunemann 2011

Schunemann HJ, Oxman AD, Vist GE, Higgins JPT, Deeks JJ, Glasziou P, et al. Chapter 12: Interpreting results and drawing conclusions. In: Higgins JPT, Green S editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. Available from www.cochrane-handbook.org. The Cochrane Collaboration, 2011.

Tallis 2002

Tallis R, Boon P, Perucca E, Stephen L. Epilepsy in elderly people: management issues. *Epileptic Disorders* 2002;**4** Suppl 2:S33-9.

Westmark 2008

Westmark CJ, Westmark PR, Beard AM, Hildebrandt SM, Malter JS. Seizure susceptibility and mortality in mice that over-express amyloid precursor protein. *International Journal of Clinical and Experimental Pathology* 2008;**1**(2):157-68.

Wu 2009

Wu CS, Wang SC, Chang IS, Lin KM. The association between dementia and long-term use of benzodiazepine in the elderly: nested case-control study using claims data. *The American Journal of Geriatric Psychiatry* 2009;**17**(7):614-20.

Wu 2012

Wu L, Rosa-Neto P, Hsiung GY, Sadovnick AD, Masellis M, Black SE, et al. Early-onset familial Alzheimer's disease (EOFAD). *Canadian Journal of Neurological Sciences* 2012;**39**(4):436-45.

References to other published versions of this review

Liu 2015

Liu J, Wang LN. Treatment of epilepsy for people with Alzheimer's disease. *Cochrane Database of Systematic Reviews* 2015, Issue 10. [DOI: [10.1002/14651858.CD011922](https://doi.org/10.1002/14651858.CD011922)]

Liu 2016

Liu J, Wang LN. Treatment of epilepsy for people with Alzheimer's disease. *Cochrane Database of Systematic Reviews* 2016, Issue 11. [DOI: [10.1002/14651858.CD011922.pub2](https://doi.org/10.1002/14651858.CD011922.pub2)]

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Cumbo 2010

Methods	A prospective, randomized, three-arm parallel-group clinical trial.
Participants	<p>Inclusion criteria: mild to moderate disease (MMSE score: 10 to 26), age between 60 and 90 years, educational level $\geq$ 5 years, a diagnosis of partial epilepsy according to the International League Against Epilepsy diagnostic scheme, and a caregiver who could ensure compliance with treatment and provide the information required for psychometric assessments.</p> <p>Exclusion criteria: history of primary neurological or psychiatric disease other than AD, history of seizures experienced prior to the development of AD, drug or alcohol abuse, clinically significant or unstable medical or surgical disorders that could influence the outcome of the study, previous treatment for epilepsy, concomitant treatment with antidepressants or neuroleptics, use of investigational drugs, and refusal to give informed consent in writing.</p>

Cumbo 2010 (Continued)

A total of 95 people with AD and epileptic seizures (41 males, 54 females) with a mean age of 71.75 years (range: 65 to 82 years) and a mean education of 6.3 years (range: 5 to 17 years) living in the community were included in the study.

Participants were randomly assigned with an AED as monotherapy: 38 were administered LEV, 28 PB, and 29 LTG.

Interventions	<p>This study comprised a 4-week dose adjustment period followed by a 12-month dose evaluation period.</p> <p>During the treatment periods, participants received LEV, PB, or LTG and were titrated to an effective dose. No participant was previously exposed to another AED. The initial target dosage of LEV was 500 mg/day, increased weekly by 500 mg/day; for PB, it was 50 mg/day increased weekly by 50 mg/day; and for LTG, it was 25 mg/day increased weekly by 25 mg/day. Thereafter, the dosage was individually adjusted. The mean daily dose of LEV monotherapy was 956 mg (range: 500 to 2000 mg/day). The mean daily dose of PB monotherapy was 90mg/day (range: 50 to 100mg/day). The mean daily dose of LTG monotherapy was 57.5 mg/day (range: 25 to 100 mg/day).</p>
Outcomes	<p>Primary outcomes: percentage of participants achieving at least 50% and 100% (seizure-freedom) reduction in seizure frequency; adverse events.</p> <p>Secondary outcomes: changes in MMSE score; changes in ADAS-Cog score; changes in Cornell scale for depression.</p>
Notes	<p>All participants had concomitant cholinesterase inhibitor therapy for AD. Other concomitant medications, including diuretics, antihypertensives, lipid-reducing agents, and antidiabetic drugs, started prior to the baseline visit were allowed during the study.</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Each subject was randomly assigned to be treated with an AED as monotherapy: 38 were administered LEV, 28 PB, and 29 LTG. The details were not provided.
Allocation concealment (selection bias)	Unclear risk	The details were not provided.
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	The details were not provided.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	The details were not provided.
Incomplete outcome data (attrition bias) All outcomes	Low risk	83 (87%) subjects (35 on LEV, 23 on PB, 25 on LTG) completed the study. The reasons for dropouts were given.
Selective reporting (reporting bias)	Low risk	The outcomes reported in the methods and results were consistent; and all important outcomes expected were reported.
Other bias	High risk	Minimum necessary sample size was not calculated.

AD: Alzheimer's disease; ADAS-Cog: Alzheimer's Disease Assessment Scale-Cognitive; AED: antiepileptic drug; LEV: levetiracetam; LTG: lamotrigine; MMSE: Mini Mental State Examination; PB: phenobarbital

Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
Campion 1995	Not a randomized controlled trial.
Lovestone 2015	The participants were not eligible.
Musaey 2017	The participants were not eligible.

Characteristics of ongoing studies *[ordered by study ID]*

[NCT02002819](#)

Trial name or title	Levetiracetam for Alzheimer's Disease-Associated Network Hyperexcitability
Methods	Randomized cross-over assignment
Participants	Meets National Institute on Aging-Alzheimer's Association Workgroups criteria for probable AD dementia; age $\leq$ 80 years at time of screening; willing and able caregiver who has daily contact with the subject; MMSE score $\geq$ 18 and/or CDR $<$ 2 at the initial screening assessment
Interventions	Levetiracetam and placebo
Outcomes	Changes in executive function, epileptiform activity frequency, cognitive function and behavior and level of disability
Starting date	January 2018
Contact information	Keith A Vossel, MD, MSc. University of Minnesota - Clinical and Translational Science Institute
Notes	Estimated Study Completion Date: December 2019

[NCT03489044](#)

Trial name or title	An Investigation of Levetiracetam in Alzheimer's Disease
Methods	Double-blind, randomized, cross-over assignment
Participants	Meets the National Institute of Aging-Alzheimer's Association criteria for probable AD
Interventions	Levetiracetam and placebo
Outcomes	Hippocampal function, mood, quality of life and adverse events
Starting date	April 2018
Contact information	Arjune Sen, PhD, FRCPE. Nuffield Department of Clinical Neurosciences
Notes	Estimated Study Completion Date: December 2019

AD: Alzheimer's disease; CDR: Clinical Dementia Rating; MMSE: Mini Mental Status Exam

DATA AND ANALYSES

Comparison 1. Levetiracetam versus lamotrigine

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Proportion with seizure freedom	1	67	Risk Ratio (M-H, Random, 95% CI)	1.20 [0.53, 2.71]
2 Reduction in seizure frequency of 50% or more	1	67	Risk Ratio (M-H, Random, 95% CI)	1.21 [0.84, 1.75]
3 Proportion with adverse events	1	67	Risk Ratio (M-H, Random, 95% CI)	0.65 [0.25, 1.74]

Analysis 1.1. Comparison 1 Levetiracetam versus lamotrigine, Outcome 1 Proportion with seizure freedom.

Analysis 1.2. Comparison 1 Levetiracetam versus lamotrigine, Outcome 2 Reduction in seizure frequency of 50% or more.

Analysis 1.3. Comparison 1 Levetiracetam versus lamotrigine, Outcome 3 Proportion with adverse events.

Comparison 2. Levetiracetam versus phenobarbital

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Proportion with seizure freedom	1	66	Risk Ratio (M-H, Random, 95% CI)	1.01 [0.47, 2.19]
2 Reduction in seizure frequency of 50% or more	1	66	Risk Ratio (M-H, Random, 95% CI)	1.11 [0.78, 1.56]
3 Proportion with adverse events	1	66	Risk Ratio (M-H, Random, 95% CI)	0.44 [0.18, 1.07]

Analysis 2.1. Comparison 2 Levetiracetam versus phenobarbital, Outcome 1 Proportion with seizure freedom.

Analysis 2.2. Comparison 2 Levetiracetam versus phenobarbital, Outcome 2 Reduction in seizure frequency of 50% or more.

Analysis 2.3. Comparison 2 Levetiracetam versus phenobarbital, Outcome 3 Proportion with adverse events.

Comparison 3. Lamotrigine versus phenobarbital

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Proportion with seizure freedom	1	57	Risk Ratio (M-H, Random, 95% CI)	0.84 [0.35, 2.02]
2 Reduction in seizure frequency of 50% or more	1	57	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.60, 1.38]
3 Proportion with adverse events	1	57	Risk Ratio (M-H, Random, 95% CI)	0.68 [0.30, 1.53]

Analysis 3.1. Comparison 3 Lamotrigine versus phenobarbital, Outcome 1 Proportion with seizure freedom.

Analysis 3.2. Comparison 3 Lamotrigine versus phenobarbital, Outcome 2 Reduction in seizure frequency of 50% or more.

Analysis 3.3. Comparison 3 Lamotrigine versus phenobarbital, Outcome 3 Proportion with adverse events.

APPENDICES

Appendix 1. Cochrane Epilepsy Group Specialized Register

1. MESH DESCRIPTOR Alzheimer Disease EXPLODE ALL AND CENTRAL:TARGET
2. MESH DESCRIPTOR Dementia EXPLODE ALL AND CENTRAL:TARGET
3. (alzheimers* OR dement*):AB,KW,MC,MH,TI AND CENTRAL:TARGET
4. #1 OR #2 OR #3
5. MESH DESCRIPTOR Epilepsy EXPLODE ALL AND CENTRAL:TARGET
6. MESH DESCRIPTOR Seizures EXPLODE ALL AND CENTRAL:TARGET
7. (epileps* OR seizure* OR convuls*):AB,KW,MC,MH,TI AND CENTRAL:TARGET
8. #5 OR #6 OR #7 AND CENTRAL:TARGET
9. #4 AND #8
10. >01/02/2016:CRSINCENTRAL AND CENTRAL:TARGET
11. #9 AND #10

Appendix 2. MEDLINE (Ovid)

This strategy is based on the Cochrane Highly Sensitive Search Strategy for identifying randomized trials ([Lefebvre 2011](#)).

1. exp Alzheimer Disease/ or exp Dementia/
2. (alzheimers\$ or dement\$).tw.
3. 1 or 2
4. exp Epilepsy/
5. exp Seizures/

6. (epilep\$ or seizure\$ or convuls\$).tw.
7. 4 or 5 or 6
8. exp *Pre-Eclampsia/ or exp *Eclampsia/
9. 7 not 8
10. (randomized controlled trial or controlled clinical trial or pragmatic clinical trial).pt. or (randomi?ed or placebo or randomly).ab.
11. clinical trials as topic.sh.
12. trial.ti.
13. 10 or 11 or 12
14. exp animals/ not humans.sh.
15. 13 not 14
16. 3 and 9 and 15
17. limit 16 to ed=20160201-20180710
18. 16 not (1\$ or 2\$).ed.
19. 18 and (2016\$ or 2017\$ or 2018\$).dt.
20. 17 or 19
21. remove duplicates from 20

Appendix 3. ClinicalTrials.gov

Interventional Studies | Alzheimer Disease AND Epilepsy | First posted on or after 02/01/2016

Appendix 4. WHO International Clinical Trials Registry Platform (ICTRP)

Condition: Alzheimer AND Epilepsy

Recruitment status: All

Date of registration between 01/02/2016 and 10/07/2018

Phases: 2, 3, 4

WHAT'S NEW

Date	Event	Description
10 July 2018	New citation required but conclusions have not changed	Conclusions are unchanged.
10 July 2018	New search has been performed	Searches updated 10 July 2018; no new studies were identified.

CONTRIBUTIONS OF AUTHORS

JL and L-NW formulated the idea and developed the basis for the review.

L-YW served as the independent party in selection of studies and assessment of risk of bias.

The manuscript was completed by JL and L-NW, and revised by Y-PW.

JL will be in charge of updating the review.

Treatment of epilepsy for people with Alzheimer's disease (Review)

Copyright © 2018 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

DECLARATIONS OF INTEREST

JL: none known.

L-NW: none known.

Y-PW: none known.

L-YW: none known.

SOURCES OF SUPPORT

Internal sources

- No sources of support supplied

External sources

- National Institute for Health Research (NIHR), UK.

INDEX TERMS

Medical Subject Headings (MeSH)

Alzheimer Disease [*complications]; Anticonvulsants [administration & dosage] [*therapeutic use]; Cognition [drug effects]; Depression [drug therapy]; Epilepsy [*drug therapy]; Lamotrigine [administration & dosage] [*therapeutic use]; Levetiracetam [administration & dosage] [*therapeutic use]; Phenobarbital [administration & dosage] [*therapeutic use]; Secondary Prevention

MeSH check words

Aged; Aged, 80 and over; Female; Humans; Male