Published in final edited form as:

Org Lett. 2019 April 19; 21(8): 2941–2946. doi:10.1021/acs.orglett.9b01014.

Deaminative Reductive Cross-Electrophile Couplings of Alkylpyridinium Salts and Aryl Bromides

Jennie Liao[†], Corey H. Basch^{†,§}, Megan E. Hoerrner^{†,§}, Michael R. Talley[†], Brian P. Boscoe[‡], Joseph W. Tucker[‡], Michelle R. Garnsey^{‡,*}, and Mary P. Watson^{†,*}

[†]Department of Chemistry & Biochemistry, University of Delaware, Newark, Delaware 19716, United States

[‡]Medicine Design, Pfizer Worldwide Research and Development, Groton, Connecticut 06340, United States

Abstract

A nickel-catalyzed reductive cross-coupling of alkylpyridinium salts and aryl bromides has been developed using Mn as the reductant. Both primary and secondary alkylpyridinium salts can be used, and high functional group and heterocycle tolerance is observed, including for protic groups. Mechanistic studies indicate formation of an alkyl radical, and controlling its fate was key to the success of this reaction.

Graphical Abstract

Authors are required to submit a graphic entry for the Table of Contents (TOC) that, in conjunction with the manuscript title, should give the reader a representative idea of one of the following: A key structure, reaction, equation, concept, or theorem, etc., that is discussed in the manuscript. Consult the journal's Instructions for Authors for TOC graphic specifications.

The increasing prevalence of sp³-hybridized carbons in pharmaceuticals, along with their presence in natural products and other targets, has spurred exciting innovations in the development of reagents and methods for the incorporation of alkyl groups.^{1,2} Methods that

*Corresponding Authors M. G.: Michelle.Garnsey@pfizer.com. M.P.W.: mpwatson@udel.edu.

§These authors contributed equally.

The authors declare no competing financial interest.

Supporting Information. The Supporting Information is available free of charge on the ACS Publications website. Experimental details and data (PDF)

harness ubiquitous functional groups in cross-couplings provide opportunities to exploit previously untapped feedstocks and for late-stage functionalization.³ Our efforts focus on methods that enable alkyl primary amines to be transformed into alkyl electrophiles. Alkyl amines are prevalent in molecules ranging from simple starting materials to complex natural products and drug candidates.^{1b, 4} They are easy to prepare, stable, and broadly compatible with many functional groups, allowing them to be carried deep into a synthetic sequence particularly when protected.^{4c, 4d} From the perspective of medicinal discovery, alkyl amines are present in pharmaceutical libraries in vast numbers and offer considerable diversity. For example, Pfizer's internal chemical inventory has >47,000 alkyl primary amines (vs. <30,000 primary and secondary alkyl halides).

We identified that alkyl amines, including those with *unactivated* alkyl groups, can become effective reagents for cross-couplings via conversion to Katritzky pyridinium salts.^{5,6,7,5c, 5d, 8,9} These Katritzky pyridinium salts are easily prepared in a single step from the primary amine, are air- and moisture-stable, and are convenient to purify. Their formation is selective for primary amines, allowing other Lewis basic groups to be tolerated in the substrate. We demonstrated a nickel-catalyzed Suzuki–Miyaura arylation of alkylpyridinium salts to form C(sp³)–C(sp²) bonds, the first example of using an unactivated alkyl amine derivative in a cross-coupling.^{5a} Since our discovery, we and others have continued to develop this underappreciated class of reagents.^{5,10}

Although the use of aryl boronic acids presents control over the arylation regioselectivity, it also presents limitations. Due to the basic conditions required for boronic acid activation, pyridinium salts with β -electron-withdrawing groups undergo elimination, and other protic groups are also problematic. An excess (3.0 equiv) of the arylboronic acid is required.^{5a} Thirdly, aryl boronic acids must be prepared from aryl halides, adding steps and limiting opportunities for efficient parallel synthesis. Finally, the availability of aryl boronic acids is much more limited than aryl bromides. In contrast to only 6,200 (het)aryl boronic acids and esters, there are >56,000 (het)aryl bromides in Pfizer's inventory. Similar constraints limit the potential of any organometallic coupling partner. Thus, a method for direct use of aryl bromides would be of high value.

Despite the exquisite methods developed for reductive couplings of other alkyl electrophiles,^{3d, 3g, 11} it was unclear if the reduction of the pyridinium cation and potential side reactions of the alkyl radical intermediate could be controlled in the presence of a stoichiometric reductant. Overcoming these challenges, we report conditions for the efficient coupling of alkyl pyridinium salts with aryl bromides (Scheme 1). This method uses abundant and diverse aryl bromides, approximately equimolar amounts of alkyl pyridinium salt and aryl bromide, and the neutral conditions allow exceptional scope of pyridinium salts, particularly those with base-sensitive functional groups. During the preparation of this manuscript, Han and Rueping reported couplings of alkylpyridinium salts and aryl iodides.¹² These reports demonstrate feasibility of a reductive coupling. However, it is important to note that aryl iodides are less available than arylboronic acids and esters; for example, less than 300 (het)aryl iodides are in Pfizer's internal inventory, making these methods of limited utility for the incorporation of diverse and readily available aryl groups.

In an effort to optimize conditions for the incorporation of heteroaryls, we selected the reductive coupling of pyridinium salt **3a** and 3-bromoquinoline for optimization. To maximize the utility of this reaction for the use of precious alkyl amines and/or precious aryl bromides, we prioritized conditions that would allow approximately equimolar amounts of alkyl pyridinium salt and aryl halide to be used. Using NiBr₂·DME and 4,4'-di^tBuBipy, a promising 9% yield was observed with Zn as the reductant (Table 1, entry 1). However, dihydropyridine **5** was a major competing product, likely arising from addition of the alkyl radical intermediate to another equivalent of **3a**. Changing the reductant to Mn⁰ gave increased yield (entry 2). Further improvement was observed with the addition of LiCl or MgCl₂, by changing the solvent to NMP, and by reducing concentration (entries 3–5). These additives may accelerate reduction of the Ni^{II} intermediates or assist in activating the surface of the Mn⁰.¹³ Under these conditions, the equivalents of Mn⁰ could be reduced (entry 6). Hypothesizing that an electron-rich ligand was necessary to promote single-electron transfer (SET) to the pyridinium cation, we found that 4,4'-diOMeBipy provided an even higher yield of 74% (entry 7). Because the aryl halide is often the precious component, we investigated conditions with aryl bromide as the limiting reagent, and observed 85% isolated yield (1.2 equiv pyridinium **3a**, entry 8). Control experiments confirmed that NiCl₂·DME, ligand, and Mn⁰ are required (entries 9–11, 13) and that MgCl₂ still provides a beneficial effect (entry 12). With minimal precaution taken to exclude air and moisture (set up under air, no drying of glassware or solvents), 53% yield is observed (entry 14).

With these conditions in hand, we turned to the reductive coupling of secondary alkyl pyridinium salts. Disappointingly, only 19% yield of product **6** was observed, and dihydropyridine **7** was formed in 32% yield (Scheme 2). In the absence of Ni catalyst, product **7** is formed in 39% yield, indicating a non-catalyzed competitive pathway for the production of the isopropyl radical via direct reduction by Mn⁰. Hypothesizing that **7** forms from addition of the isopropyl radical to pyridinium cation **3b** instead of combination with a Ni(II) intermediate (see mechanism discussion below), we proposed that maintaining a low concentration of pyridinium cation vs Ni catalyst would prevent formation of **7** and give higher yield of **6**. Indeed, slow addition of a solution of **3b** and aryl bromide resulted in a much improved 84% yield.¹⁴

These optimized conditions proved general for a wide range of alkyl pyridinium salts and aryl bromides (Scheme 3). Primary and secondary alkyl pyridinium salts underwent coupling in good yield, as did a benzylic pyridinium salt (**15**). Even a quaternary carbon can be formed via cross-coupling of a tertiary alkyl pyridinium salt, albeit in 35% yield (**20**). Pyridinium salts with more bulky tertiary alkyls cannot be prepared due to hindrance of the 2,6-diphenyls. In addition to general exploration of functional group tolerance, we were particularly interested in substrates that failed under the basic conditions of our Suzuki–Miyaura arylation.^{5a} Excitingly, pyridinium salts with β-electron-withdrawing groups are effective substrates (**11**, **12**); elimination is not observed under these mild, neutral conditions. Protic functional groups (primary amide, indole, alcohol, and dihydropyridine) were also well tolerated (**16–18**, **23**, **27**). Successful pyridinium formation and arylation of a number of natural product and pharmaceutically relevant alkyl amines highlights another advantage of using alkyl amines as precursors for late-stage functionalization. For example,

pyridinium salts derived from amino acids (**21**, **22**) and amino alcohols (**23**) can be utilized. Amine intermediates used in the synthesis of the pharmaceuticals mosapride and Lipitor® can also be arylated (**24–26**).¹⁵ Notably, the arylation to form **25** was successful on gram scale, demonstrating the scalability of this chemistry. Finally, arylation of the pharmaceutical amlodipine provides a dramatic example of the functional groups tolerated under these mild reaction conditions (**27**).¹⁶

We also observed excellent scope with respect to the aryl bromide. Aryl boronate esters (**13**) can be incorporated into the products. The use of 2-pyridyl bromide enabled the challenging 2-pyridyl group to be installed (**14**), highlighting another advantage over using an aryl boronic acid.¹⁷ A range of other heteroaryl bromides was also effective, including quinolyl (**4**, **6**, **12**, **20**, **22**), 3-pyridyl (**8**), dibenzothiophenyl (**11**), benzofuranyl (**19**), and pyrimidyl (**21**). 3-Pyridyl iodide and chloride provided <10% yield.¹⁸ Aryl bromides with *ortho* substitution, electron-donating groups, and some complex heteroaryls provided low yields.¹⁹ Saturated heterocycles, including morpholines **9**, **24**, and **25**, piperazine **10**, piperidine **19**, pyrrolidine **22**, ketal **26**, and dihydropyridine **27**, were also tolerated. Additional functional groups compatible with this method include dioxalane (**9**, **14**), Boc-protected amines (**10**, **19**, **22**), esters (**11**, **16**, **21**, **26**, **27**), trifluoromethyls (**12**, **23**, **27**), nitriles (**15**, **26**), ethers (**15**, **27**), thioethers (**21**), isoxazoles (**25**), and aryl chlorides (**27**). However, these conditions fail for alkyl or alkynyl bromides.

We also investigated one-pot transformations of amine to arylalkane. Although simultaneous addition of pyridinium salt, arylbromide, catalyst, and other reagents failed, telescoping pyridinium formation and cross-coupling provided 50% yield (Scheme 4). This yield is lower than when pyridinium salt **3a** is isolated (75%), but this protocol may be advantageous in certain cases, such as parallel synthesis.

This reaction likely proceeds via a mechanism analogous to reductive cross-couplings of alkyl halides with aryl halides (Scheme 5A).²⁰ Oxidative addition of aryl bromide to a Ni⁰ intermediate generates an arynickel(II) bromide. Single-electron transfer (SET) from a Ni^I intermediate to the pyridinium salt generates a neutral pyridyl radical, which then fragments to give alkyl radical **28**. Alkyl radical **28** may also form via reduction by Mn⁰, particularly for secondary alkylpyridinium salts. Combination of radical **28** with an arynickel(II) intermediate delivers Ni^{III} species **29**. Reductive elimination then releases product **30** and regenerates a Ni^I intermediate. In support of the oxidative addition of the aryl bromide, 24% yield of product **4** is observed when tetrakis(dimethylamino)ethylene (TDAE) is used in place of Mn⁰, suggesting an arylmanganese intermediate is not required (Scheme 5B). The intermediacy of alkyl radical **28** is consistent with the observed opening of cyclopropylmethyl pyridinium **3s**, and the formation of TEMPO-trapped adduct **32** when TEMPO is added to the reaction. However, we cannot yet distinguish between a radical-chain bimetallic pathway and a radical-rebound pathway.²¹

In summary, we have developed a nickel-catalyzed reductive cross-coupling of alkyl pyridinium salts and aryl bromides. Given the abundance and diversity of alkyl amines and (het)aryl bromides available and the mild conditions, this method enables access to an exceptional variety of products, including those with protic and other base-sensitive

functional groups. The success of this method, and the ability to use primary, secondary, and one tertiary alkyl pyridinium salts, relied on controlling the relative concentration of pyridinium salt to shuttle the alkyl radical intermediate along the desired catalytic cycle.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

We thank NIH (R01 GM111820), Pfizer, Inc., and University of Delaware (UD) for University Graduate Fellowships (J. L., C.H.B.). We thank the Martin and Molander groups for helpful discussions. Data were acquired at UD on instruments obtained with assistance of NSF and NIH funding (NSF CHE0421224, CHE1229234, CHE0840401, and CHE1048367; NIH P20 GM104316, P20 GM103541, and S10 OD016267). We thank Lotus Separations, LLC, for assistance with SFC.

REFERENCES

- (a). Lovering F; Bikker J; Humblet C Escape from Flatland: Increasing Saturation as an Approach to Improving Clinical Success. *J. Med. Chem.* 2009, 52, 6752–6756; [PubMed: 19827778]
(b) McGrath NA; Brichacek M; Njardarson JT A Graphical Journey of Innovative Organic Architectures That Have Improved Our Lives. *J. Chem. Educ.* 2010, 87, 1348–1349.
- (a). Tasker SZ; Standley EA; Jamison TF Recent advances in homogeneous nickel catalysis. *Nature* 2014, 509, 299–309; [PubMed: 24828188] (b) Yan M; Lo JC; Edwards JT; Baran PS Radicals: Reactive Intermediates with Translational Potential. *J. Am. Chem. Soc.* 2016, 138, 12692–12714; [PubMed: 27631602] (c) Molander G; Milligan JA; Phelan JP; Badir SO Recent Advances in Alkyl Carbon-Carbon Bond Formation by Nickel/Photoredox Cross-Coupling. *Angew. Chem., Int. Ed.* 0, doi:10.1002/anie.201809431; (d) Zhou JS; Fu GC Cross-Couplings of Unactivated Secondary Alkyl Halides: Room-Temperature Nickel-Catalyzed Negishi Reactions of Alkyl Bromides and Iodides. *J. Am. Chem. Soc.* 2003, 125, 14726–14727. [PubMed: 14640646]
- (a). Johnston CP; Smith RT; Allmendinger S; MacMillan DW Metallaphotoredox-catalysed sp(3)-sp(3) cross-coupling of carboxylic acids with alkyl halides. *Nature* 2016, 536, 322–325; [PubMed: 27535536] (b) Gutiérrez-Bonet Á; Tellis JC; Matsui JK; Vara BA; Molander GA 1,4-Dihydropyridines as Alkyl Radical Precursors: Introducing the Aldehyde Feedstock to Nickel/Photoredox Dual Catalysis. *ACS Catal.* 2016, 6, 8004–8008; [PubMed: 27990318] (c) Qin T; Cornella J; Li C; Malins LR; Edwards JT; Kawamura S; Maxwell BD; Eastgate MD; Baran PS A General Alkyl-Alkyl Cross-Coupling Enabled by Redox-Active Esters and Alkylzinc Reagents. *Science* 2016, 352, 801–805; [PubMed: 27103669] (d) Huihui KM; Caputo JA; Melchor Z; Olivares AM; Spiewak AM; Johnson KA; DiBenedetto TA; Kim S; Ackerman LK; Weix DJ Decarboxylative Cross-Electrophile Coupling of N-Hydroxyphthalimide Esters with Aryl Iodides. *J. Am. Chem. Soc.* 2016, 138, 5016–5019; [PubMed: 27029833] (e) Cornella J; Edwards JT; Qin T; Kawamura S; Wang J; Pan CM; Gianatassio R; Schmidt M; Eastgate MD; Baran PS Practical Ni-Catalyzed Aryl-Alkyl Cross-Coupling of Secondary Redox-Active Esters. *J. Am. Chem. Soc.* 2016, 138, 2174–2177; [PubMed: 26835704] (f) Zhang X; MacMillan DW Alcohols as Latent Coupling Fragments for Metallaphotoredox Catalysis: sp³-sp² Cross-Coupling of Oxalates with Aryl Halides. *J. Am. Chem. Soc.* 2016, 138, 13862–13865; [PubMed: 27718570] (g) Molander GA; Traister KM; O'Neill BT Engaging Nonaromatic, Heterocyclic Tosylates in Reductive Cross-Coupling with Aryl and Heteroaryl Bromides. *J. Org. Chem.* 2015, 80, 2907–2911. [PubMed: 25711834]
- (a). Ruiz-Castillo P; Buchwald SL Applications of Palladium-Catalyzed C–N Cross-Coupling Reactions. *Chem. Rev.* 2016, 116, 12564–12649; [PubMed: 27689804] (b) Liu Y; Ge H Site-selective C–H arylation of primary aliphatic amines enabled by a catalytic transient directing group. *Nat. Chem.* 2017, 9, 26–32; (c) Lawrence SA, *Amines: Synthesis, Properties and Applications*. Cambridge University Press: New York, NY, 2004; (d) Nugent TC, *Chiral Amine Synthesis*. Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, 2010.

- 5 (a). Basch CH; Liao J; Xu J; Piane JJ; Watson MP Harnessing Alkyl Amines as Electrophiles for Nickel-Catalyzed Cross Couplings via C–N Bond Activation. *J. Am. Chem. Soc.* 2017, 139, 5313–5316; [PubMed: 28359153] (b) Plunkett S; Basch CH; Santana SO; Watson MP Harnessing Alkyl Pyridinium Salts as Electrophiles in De-aminative Alkyl-Alkyl Cross-Couplings. *J. Am. Chem. Soc.* 2019, DOI: 10.1021/jacs.1029b00111; (c) Liao J; Guan W; Boscoe BP; Tucker JW; Tomlin JW; Garnsey MR; Watson MP Transforming Benzylic Amines into Diarylmethanes: Cross-Couplings of Benzylic Pyridinium Salts via C–N Bond Activation. *Org. Lett.* 2018, 20, 3030–3033; [PubMed: 29745674] (d) Guan W; Liao J; Watson MP Vinylation of Benzylic Amines via C–N Bond Functionalization of Benzylic Pyridinium Salts. *Synthesis* 2018, 50, 3231–3237. [PubMed: 30174353]
- 6 (a). Bapat JB; Blade RJ; Boulton AJ; Epszajn J; Katrizky AR; Lewis J; Molina-Buendia P; Nie P-L; Ramsden CA Pyridines as Leaving Groups in Synthetic Transformations: Nucleophilic Displacements of Amino Groups, and Novel Preparations of Nitriles and Isocyanates. *Tetrahedron Lett.* 1976, 31, 2691–2694; (b) Katrizky AR; Marson CM Pyrylium Mediated Transformations of Primary Amino Groups into Other Functional Groups. *Angew. Chem., Int. Ed.* 1984, 23, 420–429; (c) Sowmiah S; Esperança JMSS; Rebelo LPN; Afonso CAM Pyridinium salts: from synthesis to reactivity and applications. *Org. Chem. Front.* 2018, 5, 453–493.
7. Ouyang K; Hao W; Zhang WX; Xi Z Transition-Metal-Catalyzed Cleavage of C–N Single Bonds. *Chem. Rev.* 2015, 115, 12045–12090. [PubMed: 26423200]
8. For electronically or strain-activated alkyl groups, see: Huang C-Y; Doyle AG Nickel-Catalyzed Negishi Alkylations of Styrenyl Aziridines. *J. Am. Chem. Soc.* 2012, 134, 9541–9544; [PubMed: 22414150] Li M-B; Wang Y; Tian S-K Regioselective and Stereospecific Cross-Coupling of Primary Allylic Amines with Boronic Acids and Boronates through Palladium-Catalyzed C–N Bond Cleavage. *Angew. Chem., Int. Ed.* 2012, 51, 2968–2971; Maity P; Shacklady-McAtee DM; Yap GPA; Sirianni ER; Watson MP Nickel-Catalyzed Cross Couplings of Benzylic Ammonium Salts and Boronic Acids: Stereospecific Formation of Diarylethanes via C–N Bond Activation. *J. Am. Chem. Soc.* 2013, 135, 280–285; [PubMed: 23268734] Jensen KL; Standley EA; Jamison TF Highly Regioselective Nickel-Catalyzed Cross-Coupling of N-Tosylaziridines and Alkylzinc Reagents. *J. Am. Chem. Soc.* 2014, 136, 11145–11152; [PubMed: 25055180] Shacklady-McAtee DM; Roberts KM; Basch CH; Song Y-G; Watson MP A general, simple catalyst for enantiospecific cross couplings of benzylic ammonium triflates and boronic acids: no phosphine ligand required. *Tetrahedron* 2014, 70, 4257–4263; [PubMed: 25364060] Zhang H; Hagihara S; Itami K Making Dimethylamino a Transformable Directing Group by Nickel-Catalyzed C–N Borylation. *Chem. Eur. J.* 2015, 21, 16796–16800; [PubMed: 26435307] Basch CH; Cobb KM; Watson MP Nickel-Catalyzed Borylation of Benzylic Ammonium Salts: Stereospecific Synthesis of Enantioenriched Benzylic Boronates. *Org. Lett.* 2016, 18, 136–139; [PubMed: 26679212] Hu J; Sun H; Cai W; Pu X; Zhang Y; Shi Z Nickel-Catalyzed Borylation of Aryl- and Benzyltrimethylammonium Salts via C–N Bond Cleavage. *J. Org. Chem.* 2016, 81, 14–24; [PubMed: 26628255] Moragas T; Gaydou M; Martin R Nickel-Catalyzed Carboxylation of Benzylic C–N Bonds with CO₂. *Angew. Chem., Int. Ed.* 2016, 55, 5053–5057; Yi Y-Q-Q; Yang W-C; Zhai D-D; Zhang X-Y; Li S-Q; Guan B-T Nickel-catalyzed C–N bond reduction of aromatic and benzylic quaternary ammonium triflates. *Chem. Commun.* 2016, 52, 10894–10897; Gui Y; Tian S-K Stereospecific Nucleophilic Substitution of Enantioenriched Tertiary Benzylic Amines via in Situ Activation with Benzyne. *Org. Lett.* 2017, 19, 1554–1557; [PubMed: 28333467] Guisán-Ceinos M; Martín-Heras V; Tortosa M Regio- and Stereospecific Copper-Catalyzed Substitution Reaction of Propargylic Ammonium Salts with Aryl Grignard Reagents. *J. Am. Chem. Soc.* 2017, 139, 8448–8451. [PubMed: 28613864]
9. For vinyl and aryl pyridinium salts, see: Buszek KR; Brown N N-Binylpyridinium and -ammonium Tetrafluoroborate Salts: New Electrophilic Coupling Partners for Pd(0)-Catalyzed Suzuki Cross-Coupling Reactions. *Org. Lett.* 2007, 9, 707–710; [PubMed: 17253706] Moser D; Duan Y; Wang F; Ma Y; O'Neill Matthew J; Cornella J Selective Functionalization of Aminoheterocycles by a Pyrylium Salt. *Angew. Chem., Int. Ed.* 2018, 57, 11035–11039.
- 10 (a). Klauk FJR; James MJ; Glorius F Deaminative Strategy for the Visible-Light-Mediated Generation of Alkyl Radicals. *Angew. Chem., Int. Ed.* 2017, 56, 12336–12339; (b) Ociepa M; Turkowska J; Gryko D Redox-Activated Amines in C(sp³)–C(sp) and C(sp³)–C(sp²) Bond Formation Enabled by Metal-Free Photoredox Catalysis. *ACS Catal.* 2018, 8, 11362–11367; (c) Wu J; He L; Noble A; Aggarwal VK Photoinduced Deaminative Borylation of Alkylamines. *J.*

Am. Chem. Soc. 2018, 140, 10700–10704; [PubMed: 30091912] (d)Sandfort F; Strieth-Kalthoff F; Klauck FJR; James MJ; Glorius F Deaminative Borylation of Aliphatic Amines Enabled by Visible Light Excitation of an Electron Donor–Acceptor Complex. *Chem. Eur. J.* 2018, 24, 17210–17214; [PubMed: 30290050] (e)Hu J; Wang G; Li S; Shi Z Selective C–N Borylation of Alkyl Amines Promoted by Lewis Base. *Angew. Chem., Int. Ed.* 2018, 57, 15227–15231; (f)Zhang M-M; Liu F Visible-light-mediated allylation of alkyl radicals with allylic sulfones via a deaminative strategy. *Organic Chemistry Frontiers* 2018, 5, 3443–3446; (g)For a recent three-component coupling of activated alkyl pyridinium salts, see: Klauck FJR; Yoon H; James MJ; Lautens M; Glorius F Visible-Light-Mediated Deaminative Three-Component Dicarbofunctionalization of Styrenes with Benzylic Radicals. *ACS Catal.* 2019, 9, 236–241; (h)Jiang X; Zhang MM; Xiong W; Lu LQ; Xiao WJ Deaminative (Carbonylative) Alkyl-Heck-type Reactions Enabled by Photocatalytic C–N Bond Activation. *Angew. Chem., Int. Ed.* 2019, 58, 2402–2406.

- 11 (a). Weix DJ Methods and Mechanisms for Cross-Electrophile Coupling of Csp² Halides with Alkyl Electrophiles. *Acc. Chem. Res.* 2015, 48, 1767–1775; [PubMed: 26011466] (b)Everson DA; Weix DJ Cross-Electrophile Coupling: Principles of Reactivity and Selectivity. *J. Org. Chem.* 2014, 79, 4793–4798; [PubMed: 24820397] (c)Everson DA; Shrestha R; Weix DJ Nickel-Catalyzed Reductive Cross-Coupling of Aryl Halides with Alkyl Halides. *J. Am. Chem. Soc.* 2010, 132, 920–921; [PubMed: 20047282] (d)Everson DA; Jones BA; Weix DJ Replacing Conventional Carbon Nucleophiles with Electrophiles: Nickel-Catalyzed Reductive Alkylation of Aryl Bromides and Chlorides. *J. Am. Chem. Soc.* 2012, 134, 6146–6159; [PubMed: 22463689] (e)Wang S; Qian Q; Gong H Nickel-Catalyzed Reductive Coupling of Aryl Halides with Secondary Alkyl Bromides and Allylic Acetate. *Org. Lett.* 2012, 14, 3352–3355; [PubMed: 22697415] (f)Molander GA; Traister KM; O'Neill BT Reductive Cross-Coupling of Nonaromatic, Heterocyclic Bromides with Aryl and Heteroaryl Bromides. *J. Org. Chem.* 2014, 79, 5771–5780; [PubMed: 24892751] (g)Wang X; Wang S; Xue W; Gong H Nickel-Catalyzed Reductive Coupling of Aryl Bromides with Tertiary Alkyl Halides. *J. Am. Chem. Soc.* 2015, 137, 11562–11565; [PubMed: 26325479] (h)Wang X; Ma G; Peng Y; Pitsch CE; Moll BJ; Ly TD; Wang X; Gong H Ni-Catalyzed Reductive Coupling of Electron-Rich Aryl Iodides with Tertiary Alkyl Halides. *J. Am. Chem. Soc.* 2018, 140, 14490–14497; [PubMed: 30296073] (i)Zhang P; Le CC; MacMillan DW Silyl Radical Activation of Alkyl Halides in Metallaphotoredox Catalysis: A Unique Pathway for Cross-Electrophile Coupling. *J. Am. Chem. Soc.* 2016, 138, 8084–8087; [PubMed: 27263662] (j)Amatore M; Gosmini C Direct Method for Carbon–Carbon Bond Formation: The Functional Group Tolerant Cobalt-Catalyzed Alkylation of Aryl Halides. *Chem. Eur. J.* 2010, 16, 5848–5852; [PubMed: 20379979] (k)Sun S-Z; Martin R Nickel-Catalyzed Umpolung Arylation of Ambiphilic α -Bromoalkyl Boronic Esters. *Angew. Chem., Int. Ed.* 2018, 57, 3622–3625.
- 12 (a). Han also reported reductive coupling with alkyl and alkynyl halides. See: Ni S; Li C; Mao Y; Pan Y; Han J Ni-Catalyzed Deamination Cross-Electrophile Coupling of Katritzky Salts with Halides via C–N Bond Activation. *ChemRxiv* 2019, DOI: 10.26434/chemrxiv.7638164.v7638161; (b)Yue H; Zhu C; Shen L; Geng Q; Hock KJ; Yuan T; Cavallo L; Rueping M Nickel-catalyzed C–N bond activation: activated primary amines as alkylating reagents in reductive cross-coupling. *Chem. Sci.* 2019, DOI: 10.1039/c1039sc00783k.
- 13 (a). Feng C; Cunningham DW; Easter QT; Blum SA Role of LiCl in Generating Soluble Organozinc Reagents. *J. Am. Chem. Soc.* 2016, 138, 11156–11159; [PubMed: 27547857] (b)Zhao C; Jia X; Wang X; Gong H Ni-Catalyzed Reductive Coupling of Alkyl Acids with Unactivated Tertiary Alkyl and Glycosyl Halides. *J. Am. Chem. Soc.* 2014, 136, 17645–17651. [PubMed: 25415424]
14. Aryl bromide was added via slow addition to prevent diaryl formation. See Supporting Information.
- 15 (a). Kato S; Morie T; Yoshida N Synthesis and Biological Activities of Metabolites of Mosapride, a New Gastroprokinetic Agent. *Chem. Pharm. Bull.* 1995, 43, 699–702; [PubMed: 7600620] (b)Roth BD Trans-6-[2-(3- or 4-carboxamido-substituted pyrrol-1-yl)alkyl]-4-hydroxypyran-2-one Inhibitors of Cholesterol Synthesis US 4,681,893, 1987.

16. Arrowsmith JE; Campbell SF; Cross PE; Stubbs JK; Burges RA; Gardiner DG; Blackburn KJ Long-acting dihydropyridine calcium antagonists. 1. 2-Alkoxymethyl derivatives incorporating basic substituents. *J. Med. Chem.* 1986, 29, 1696–1702. [PubMed: 2943898]
17. Everson DA; Buonomo JA; Weix DJ Nickel-Catalyzed Cross-Electrophile Coupling of 2-Chloropyridines with Alkyl Bromides. *Synlett* 2014, 25, 233–238. [PubMed: 24795502]
18. See Supporting Information.
19. For additional studies of heteroaryl bromides, see the Supporting Information.
20. Biswas S; Weix DJ Mechanism and Selectivity in Nickel-Catalyzed Cross-Electrophile Coupling of Aryl Halides with Alkyl Halides. *J. Am. Chem. Soc.* 2013, 135, 16192–16197. [PubMed: 23952217]
21. For further description of these pathways and related experiments, please see the Supporting Information.

- abundant and diverse aryl bromides
- avoids arylboronic acid (required in prior method)
- increased step efficiency
- neutral reaction conditions
- unprecedented functional group tolerance
- near equimolar ratio of coupling partners

Scheme 1.

Reductive coupling of alkyl pyridinium salts and aryl bromides

Scheme 2. Optimization of secondary alkyl pyridinium salt^a

^a Conditions: pyridinium salt **3b** (0.50 mmol), 3-bromoquinoline (1.1 equiv), NiCl₂·DME (10 mol %), 4,4'-diOMeBipy (12 mol %), Mn⁰ (2.0 equiv), MgCl₂ (1.0 equiv), NMP (0.17 M), 80 °C, 24 h, unless noted otherwise. Yields determined by ¹H NMR using 1,3,5-trimethoxybenzene as internal standard.

Scheme 3.
Substrate Scope^a

^a Conditions: pyridinium salt **3** (1.2 mmol), aryl bromide (1.0 mmol), NiCl₂-DME (10 mol %), 4,4'-dimethoxy-2,2'-bipyridine (12 mol %), Mn⁰ (2.0 equiv), MgCl₂ (1.0 equiv), NMP (0.17 M) 80 °C, 24 h. Average isolated yields (±5%) from duplicate experiments. ^b Slow addition of pyridinium salt **3** (1.0 mmol) and aryl bromide (1.1 mmol). ^c 4.0 mmol scale, single experiment.

Scheme 4.
Telescoped pyridinium formation and cross-coupling

A. Basic mechanistic proposal **B. Mechanistic experiments**

Scheme 5.
Mechanism and supporting experiments

Table 1.

Optimization of primary alkyl pyridinium salt^a

entry	ligand	additive	yield (%) ^b		
			4	5	Ar–Ar
1 ^{c,d}	4,4'-di ^t BuBipy	–	9	24	5
2 ^d	4,4'-di ^t BuBipy	–	26	25	4
3 ^d	4,4'-di ^t BuBipy	LiCl	54	14	4
4	4,4'-di ^t BuBipy	LiCl	60	12	10
5	4,4'-di ^t BuBipy	MgCl ₂	65	8	15
6 ^e	4,4'-di ^t BuBipy	MgCl ₂	67	6	17
7 ^e	4,4'-diOMeBipy	MgCl ₂	74	7	15
8 ^{e,f}	4,4'-diOMeBipy	MgCl ₂	85 ^g	n.d.	n.d.
9 ^{e,f,h}	4,4'-diOMeBipy	MgCl ₂	0	36	0
10 ^{e,f}	none	MgCl ₂	0	16	14
11 ^{e,f,h}	none	MgCl ₂	0	30	0
12 ^{e,f}	4,4'-diOMeBipy	–	50	24	12
13 ^{f,i}	4,4'-diOMeBipy	MgCl ₂	0	0	0
14 ^{e,f,j}	4,4'-diOMeBipy	MgCl ₂	53 ^g	n.d.	n.d.

^a Conditions: pyridinium salt **3a** (0.10 mmol), 3-bromoquinoline (1.1 equiv), NiCl₂·DME (10 mol %), ligand (12 mol %), Mn⁰ (3.0 equiv), additive (0 or 1.0 equiv), NMP (0.17 M), 80 °C, 24 h, unless noted otherwise.

^b Determined by ¹H NMR using 1,3,5-trimethoxybenzene as internal standard.

^c Zn⁰ instead of Mn⁰.

^d NiBr₂·DME (10 mol %), DMA (0.33 M).

^e Mn⁰ (2.0 equiv).

^f **3a** (1.2 equiv), 3-bromoquinoline (1.0 equiv).

^g 1.0 mmol scale. Isolated yield.

^h No NiCl₂·DME.

ⁱ No Mn⁰.

^jMinimal precautions to protect from air and moisture. n.d. = not determined.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript