Cochrane
Library

Cochrane Database of Systematic Reviews

Pro-coagulant haemostatic factors for the prevention and treatment of bleeding in people without haemophilia (Review)

Fabes J, Brunskill SJ, Curry N, Doree C, Stanworth SJ

Fabes J, Brunskill SJ, Curry N, Doree C, Stanworth SJ.

Pro-coagulant haemostatic factors for the prevention and treatment of bleeding in people without haemophilia.

Cochrane Database of Systematic Reviews 2018, Issue 12. Art. No.: CD010649.

DOI: [10.1002/14651858.CD010649.pub2](https://doi.org/10.1002/14651858.CD010649.pub2).

www.cochranelibrary.com

Pro-coagulant haemostatic factors for the prevention and treatment of bleeding in people without haemophilia
(Review)

Copyright © 2018 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

WILEY

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	3
SUMMARY OF FINDINGS	4
BACKGROUND	11
OBJECTIVES	12
METHODS	12
RESULTS	15
Figure 1.	16
Figure 2.	20
DISCUSSION	32
AUTHORS' CONCLUSIONS	34
ACKNOWLEDGEMENTS	34
REFERENCES	35
CHARACTERISTICS OF STUDIES	42
DATA AND ANALYSES	121
Analysis 1.1. Comparison 1 Fibrinogen: prophylactic trials: intervention vs inactive control, Outcome 1 All-cause mortality.	124
Analysis 1.2. Comparison 1 Fibrinogen: prophylactic trials: intervention vs inactive control, Outcome 2 Arterial thromboembolic events.	124
Analysis 1.3. Comparison 1 Fibrinogen: prophylactic trials: intervention vs inactive control, Outcome 3 Venous thromboembolic events.	125
Analysis 1.4. Comparison 1 Fibrinogen: prophylactic trials: intervention vs inactive control, Outcome 4 Mortality due to bleeding.	125
Analysis 1.5. Comparison 1 Fibrinogen: prophylactic trials: intervention vs inactive control, Outcome 5 RBC transfusion requirement.	126
Analysis 1.6. Comparison 1 Fibrinogen: prophylactic trials: intervention vs inactive control, Outcome 6 Receipt of a RBC transfusion.	126
Analysis 1.7. Comparison 1 Fibrinogen: prophylactic trials: intervention vs inactive control, Outcome 7 Blood loss.	127
Analysis 1.8. Comparison 1 Fibrinogen: prophylactic trials: intervention vs inactive control, Outcome 8 Allergic Adverse Events.	128
Analysis 2.1. Comparison 2 Fibrinogen: prophylactic trials: intervention vs haemostatically active control, Outcome 1 All-cause mortality.	130
Analysis 2.2. Comparison 2 Fibrinogen: prophylactic trials: intervention vs haemostatically active control, Outcome 2 Arterial thromboembolic events.	130
Analysis 2.3. Comparison 2 Fibrinogen: prophylactic trials: intervention vs haemostatically active control, Outcome 3 Venous thromboembolic events.	130
Analysis 2.4. Comparison 2 Fibrinogen: prophylactic trials: intervention vs haemostatically active control, Outcome 4 Mortality due to bleeding.	130
Analysis 2.5. Comparison 2 Fibrinogen: prophylactic trials: intervention vs haemostatically active control, Outcome 5 RBC transfusion requirement.	130
Analysis 2.6. Comparison 2 Fibrinogen: prophylactic trials: intervention vs haemostatically active control, Outcome 6 Receipt of a RBC transfusion.	131
Analysis 2.7. Comparison 2 Fibrinogen: prophylactic trials: intervention vs haemostatically active control, Outcome 7 Allergic Adverse Events.	131
Analysis 3.1. Comparison 3 Fibrinogen: therapeutic trials: intervention vs inactive control, Outcome 1 All-cause mortality.	133
Analysis 3.2. Comparison 3 Fibrinogen: therapeutic trials: intervention vs inactive control, Outcome 2 Arterial thromboembolic events.	134
Analysis 3.3. Comparison 3 Fibrinogen: therapeutic trials: intervention vs inactive control, Outcome 3 Venous thromboembolic events.	135
Analysis 3.4. Comparison 3 Fibrinogen: therapeutic trials: intervention vs inactive control, Outcome 4 Mortality due to bleeding.	135
Analysis 3.5. Comparison 3 Fibrinogen: therapeutic trials: intervention vs inactive control, Outcome 5 Receipt of a RBC transfusion.	136

Analysis 3.6. Comparison 3 Fibrinogen: therapeutic trials: intervention vs inactive control, Outcome 6 RBC transfusion requirement.	137
Analysis 3.7. Comparison 3 Fibrinogen: therapeutic trials: intervention vs inactive control, Outcome 7 Blood loss.	138
Analysis 3.8. Comparison 3 Fibrinogen: therapeutic trials: intervention vs inactive control, Outcome 8 Allergic Adverse Events.	138
Analysis 4.1. Comparison 4 Fibrinogen: therapeutic trials: intervention vs haemostatically active control, Outcome 1 All-cause mortality.	140
Analysis 4.2. Comparison 4 Fibrinogen: therapeutic trials: intervention vs haemostatically active control, Outcome 2 Arterial thromboembolic events.	140
Analysis 4.3. Comparison 4 Fibrinogen: therapeutic trials: intervention vs haemostatically active control, Outcome 3 Venous thromboembolic events.	140
Analysis 4.4. Comparison 4 Fibrinogen: therapeutic trials: intervention vs haemostatically active control, Outcome 4 Mortality due to bleeding.	141
Analysis 4.5. Comparison 4 Fibrinogen: therapeutic trials: intervention vs haemostatically active control, Outcome 5 RBC transfusion requirement.	141
Analysis 4.6. Comparison 4 Fibrinogen: therapeutic trials: intervention vs haemostatically active control, Outcome 6 Receipt of a RBC transfusion.	142
Analysis 4.7. Comparison 4 Fibrinogen: therapeutic trials: intervention vs haemostatically active control, Outcome 7 Allergic Adverse Events.	142
Analysis 5.1. Comparison 5 FXIII: prophylactic trials: intervention vs inactive control, Outcome 1 All-cause mortality.	144
Analysis 5.2. Comparison 5 FXIII: prophylactic trials: intervention vs inactive control, Outcome 2 Arterial thromboembolic events.	144
Analysis 5.3. Comparison 5 FXIII: prophylactic trials: intervention vs inactive control, Outcome 3 Venous thromboembolic events.	145
Analysis 5.4. Comparison 5 FXIII: prophylactic trials: intervention vs inactive control, Outcome 4 Mortality due to bleeding.	145
Analysis 5.5. Comparison 5 FXIII: prophylactic trials: intervention vs inactive control, Outcome 5 RBC transfusion requirement.	145
Analysis 5.6. Comparison 5 FXIII: prophylactic trials: intervention vs inactive control, Outcome 6 Receipt of a RBC transfusion.	146
Analysis 5.7. Comparison 5 FXIII: prophylactic trials: intervention vs inactive control, Outcome 7 Blood loss.	146
Analysis 5.8. Comparison 5 FXIII: prophylactic trials: intervention vs inactive control, Outcome 8 Allergic Adverse Events.	147
Analysis 6.1. Comparison 6 FXIII: therapeutic trials: intervention vs inactive control, Outcome 1 Allergic Adverse Events.	147
Analysis 7.1. Comparison 7 PCC: prophylactic trials: intervention vs inactive control, Outcome 1 All-cause mortality.	148
Analysis 7.2. Comparison 7 PCC: prophylactic trials: intervention vs inactive control, Outcome 2 Allergic Adverse Events.	148
ADDITIONAL TABLES	148
APPENDICES	165
CONTRIBUTIONS OF AUTHORS	171
DECLARATIONS OF INTEREST	171
SOURCES OF SUPPORT	171
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	171
INDEX TERMS	172

[Intervention Review]

Pro-coagulant haemostatic factors for the prevention and treatment of bleeding in people without haemophilia

Jez Fabes¹, Susan J Brunskill², Nicola Curry³, Carolyn Doree², Simon J Stanworth⁴

¹John Radcliffe Hospital, Oxford, UK. ²Systematic Review Initiative, NHS Blood and Transplant, Oxford, UK. ³Oxford Haemophilia & Thrombosis Centre, Churchill Hospital, Oxford, UK. ⁴National Institute for Health Research (NIHR) Oxford Biomedical Research Centre, Oxford University Hospitals NHS Foundation Trust and University of Oxford, Oxford, UK

Contact address: Susan J Brunskill, Systematic Review Initiative, NHS Blood and Transplant, Level 2, John Radcliffe Hospital, Headington, Oxford, Oxon, OX3 9BQ, UK. susan.brunskill@nhsbt.nhs.uk.

Editorial group: Cochrane Injuries Group.

Publication status and date: New, published in Issue 12, 2018.

Citation: Fabes J, Brunskill SJ, Curry N, Doree C, Stanworth SJ. Pro-coagulant haemostatic factors for the prevention and treatment of bleeding in people without haemophilia. *Cochrane Database of Systematic Reviews* 2018, Issue 12. Art. No.: CD010649. DOI: [10.1002/14651858.CD010649.pub2](https://doi.org/10.1002/14651858.CD010649.pub2).

Copyright © 2018 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Some hospital patients may be at risk of or may present with major bleeding. Abnormalities of clotting (coagulation) are often recorded in these people, and the traditional management has been with transfusions of blood components, either to prevent bleeding (prophylactic) or to treat bleeding (therapeutic). There is growing interest in the use of targeted therapies with specific pro-coagulant haemostatic (causing bleeding to stop and to keep blood within a damaged blood vessel) factor concentrates in place of plasma.

Objectives

To assess the effects and safety of pro-coagulant haemostatic factors and factor concentrates in the prevention and treatment of bleeding in people without haemophilia.

Search methods

We searched for randomised controlled trials (RCTs) in the Cochrane Central Register of Controlled Trials (2018, issue 3), MEDLINE (from 1948), Embase (from 1974), CINAHL (from 1938), PubMed (publications in process to 18 April 2018), PROSPERO, Transfusion Evidence Library (from 1950), LILACS (from 1980), IndMED (from 1985), KoreaMed (from 1934), Web of Science Conference Proceedings Citation Index (from 1990) and ongoing trial databases to 18 April 2018.

Selection criteria

We included RCTs that compared intravenous administration of a pro-coagulant haemostatic factor concentrate, either with placebo, current best or standard treatment, or another pro-coagulant haemostatic factor concentrate for prevention or treatment of bleeding. There was no restriction on the types of participants. We excluded studies of desmopressin, tranexamic acid and aminocaproic acid and use of pro-coagulant haemostatic factors for vitamin K over-anticoagulation.

Data collection and analysis

We followed standard Cochrane methodological procedures.

Main results

We identified 31 RCTs with 2392 participants and 22 ongoing trials. There were 13 therapeutic RCTs that randomised 1057 participants (range from 20 to 249 participants) and 18 prophylactic trials that randomised 1335 participants (range 20 to 479 participants). The pro-

coagulant haemostatic factor concentrate was fibrinogen in 23 trials, Factor XIII in seven trials and pro-thrombin complex concentrates (PCC) in one trial.

Seventeen trials had industrial funding or support, eight studies either did not declare their funding or were unclear about their source of funding and six studies declared non-industrial funding sources.

Certainty in the evidence and included study bias

Our certainty in the evidence, using GRADE criteria, ranged from very low to high across all outcomes. We assessed most outcomes as being of low certainty. Risks of bias were a concern in many of the RCTs; randomisation methodology was unclear in 15 RCTs, with allocation concealment unclear in 14 RCTs and at high risk of bias in five RCTs. The blinding status of outcome assessors was unclear in 13 RCTs and at high risk of bias in five RCTs, although most outcomes in these trials were objective and not prone to observer bias. Study personnel were often unblinded or insufficient information was available to assess their level of blinding (five RCTs were at unclear risk and seven at high risk of bias).

Primary outcomes

All-cause mortality was reported by 21 RCTs, arterial thromboembolic events by 22 RCTs, and venous thromboembolic events by 21 RCTs.

Fibrinogen concentrate: prophylactic trials with inactive comparator (nine RCTs)

The trials had heterogeneous clinical settings and outcome time points, so we did not pool the data. Compared to placebo, there was no evidence that prophylactic fibrinogen concentrate reduced all-cause mortality (4 RCTs; 248 participants). Compared to inactive comparators there was low- to moderate-quality evidence that prophylactic fibrinogen concentrate did not increase the risk of arterial or venous thromboembolic complications (7 RCTs; 398 participants).

Fibrinogen concentrate: prophylactic trials with active comparator (two RCTs)

There was no mortality or incidence of thromboembolic events in these two RCTs (with 57 participants).

Fibrinogen concentrate: therapeutic trials with inactive comparator (eight RCTs)

The trials had heterogeneous surgical settings and outcome time points, so we pooled data for subgroups only. Compared to an inactive comparator, there was no evidence (quality ranging from low to high) that fibrinogen concentrate reduced all-cause mortality in actively bleeding participants (7 RCTs; 724 participants). Compared to inactive comparators there was no evidence that the use of fibrinogen concentrate in active bleeding increased arterial (7 RCTs; 607 participants) or venous (6 RCTs; 562 participants) thromboembolic events.

Fibrinogen concentrate: therapeutic trials with active comparator (four RCTs)

We did not pool the outcome data, as they were not measured at comparable time points. Compared to other active pro-coagulant agents, there was no evidence (very low to moderate quality) that fibrinogen concentrate reduced all-cause mortality in actively bleeding participants (4 RCTs; 220 participants). There was no evidence that fibrinogen concentrate increased the risk of arterial (3 RCTs; 126 participants) or venous (4 RCTs; 220 participants) thromboembolic events.

FactorXIII: Prophylactic trials with inactive comparator (six trials)

The trials were heterogeneous in their surgical settings and time points for outcome analysis, so we pooled data for subgroups only. Compared to an inactive comparator, there was no evidence that prophylactic Factor XIII reduced all-cause mortality (5 RCTs; 414 participants). There was no evidence (very low to low quality) of a difference in the arterial or venous event rate between Factor XIII and inactive comparators (4 trials; 354 participants).

FactorXIII: therapeutic trials with inactive comparator (one trial)

There was no mortality or incidence of thromboembolic events in this trial.

Prothrombin complex concentrate (PCC): prophylactic trials with inactive comparator (one trial)

There was no evidence (moderate quality) that PCC reduced all-cause mortality (1 trial; 78 participants). No thromboembolic complications were reported in this trial.

Authors' conclusions

The paucity of good-quality comparable evidence precludes the drawing of conclusions for clinical practice. Further research is required to determine the risk-to-benefit ratio of these interventions. The sample sizes of future RCTs would need to be greatly increased to detect a reduction in mortality or thromboembolic events between treatment arms. To improve consistency in outcome reporting, the development of core outcome sets is essential and may help address a number of the limitations identified in this review.

PLAIN LANGUAGE SUMMARY

The effectiveness of drugs to help blood clot in the prevention and treatment of bleeding in people without haemophilia

Review question

Do medicines that help in the prevention and treatment of bleeding reduce the risk of death, the risk of experiencing a blood clot and reduce the amount of blood lost in people who do not suffer from haemophilia (a bleeding disorder), who are bleeding, or at risk of bleeding?

Background

Coagulopathy, defined as a failure of normal blood clotting, is common in severe illness, trauma and major surgery. This can make bleeding worse and may cause death. Medicines are available that are blood clotting factors that can be given into the vein to treat coagulopathy and bleeding. However, we lack enough information about how effective and safe they are.

This review combines all the data available on these medicines to assess their effectiveness and safety.

Study characteristics

We searched the medical literature up to 18 April 2018 for randomised controlled trials (RCTs), as these provide the most reliable evidence. We identified 31 relevant RCTs that contained results from 2392 participants and compared these medicines either against placebo (inactive treatment) or another medicine or blood product. The RCTs focused on three types of factors that may improve blood clotting: fibrinogen (a clotting factor that increases clot strength), clotting factor XIII (important for holding clots together) and prothrombin complex concentrate (a combination of four precursor clotting factors).

The trials either gave these medicines before bleeding occurred (prophylactically) or to treat existing bleeding (therapeutically). Most trials focused on surgery, especially heart surgery, trauma, and bleeding after childbirth.

Seventeen of the RCTs we identified had industrial support, eight had unclear funding sources and six declared non-industrial funding.

Key results

No medicine had any effect on the risk of dying, regardless of clinical setting or how the medicine was used. However, our certainty in the results is low and this finding may change in the future when new studies are published.

No medicine increased the risk of harmful clots in the veins or arteries, but our certainty in these findings is low.

Prophylactic fibrinogen reduced bleeding after heart and orthopaedic surgery compared to placebo. Prophylactic fibrinogen, compared to placebo, almost halved the need for blood transfusion following heart surgery, and reduced the need by three-quarters in other surgery. Fibrinogen reduced the need for blood transfusion when used to treat bleeding.

Prophylactic factor XIII reduced bleeding after heart surgery.

Sample sizes in future randomised trials need to be greatly increased in order to show any differences in overall survival and death due to bleeding.

Certainty of the evidence

Our certainty in the evidence is low, but future research may change the findings of this review.

Conclusion

The low certainty of the evidence makes it difficult to draw conclusions about how well these medicines work and whether they should be used in current health care. Further research is required in larger-scale RCTs to determine the benefit and costs of these treatments and whether this outweighs their risks.

SUMMARY OF FINDINGS

Summary of findings for the main comparison. FIBRINOGEN: Therapeutic trials: intervention compared to inactive Control for the prevention and treatment of bleeding in participants without haemophilia

FIBRINOGEN: Therapeutic trials: intervention compared to inactive control for the prevention and treatment of bleeding in participants without haemophilia

Patient or population: the prevention and treatment of bleeding in patients without haemophilia

Setting: hospital

Intervention: FIBRINOGEN in therapeutic trials

Comparison: Inactive control

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	N° of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with inactive control	Risk with fibrinogen in therapeutic trials				
All-cause mortality	Study population		RR 1.46 (0.71 to 2.99)	97 (2 RCTs)	⊕⊕⊕⊖ MODERATE ^a	-
Up to 28 days following admission	184 per 1000	268 per 1000 (130 to 549)				
All-cause mortality	Study population		RR 5.00 (0.25 to 102.00)	120 (1 RCT)	⊕⊕⊖⊖ LOW ^{a,b}	-
In-hospital mortality up to 30 days post-operatively	0 per 1000	0 per 1000 (0 to 0)				
All-cause mortality	Study population		-	294 (2 RCTs)	⊕⊕⊖⊖ LOW ^{a,b}	No events
Up to 6 weeks post-natally	see comment	see comment				
All-cause mortality	Study population		RR 0.23 (0.05 to 1.01)	213 (2 RCTs)	⊕⊕⊕⊕ HIGH	-
Up to 46 days post-operatively	85 per 1000	20 per 1000 (4 to 86)				

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR:** Risk ratio

GRADE Working Group grades of evidence

High certainty: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

^aWe downgraded by one level as neither study was designed to measure mortality.

^bWe downgraded by one level, due to low event rate.

Summary of findings 2. FIBRINOGEN: Therapeutic trials: intervention compared to haemostatically active control for the prevention and treatment of bleeding in participants without haemophilia

FIBRINOGEN: Therapeutic trials: intervention compared to haemostatically active control for the prevention and treatment of bleeding in participants without haemophilia

Patient or population: the prevention and treatment of bleeding in patients without haemophilia

Setting: hospital

Intervention: FIBRINOGEN in therapeutic trials

Comparison: Haemostatically active control

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	N° of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with haemostatically active control	Risk with FIBRINOGEN in therapeutic trials				
All-cause mortality	Study population		not estimable	63 (1 RCT)	⊕⊕⊕⊖ MODERATE ^a	No events
Up to 7 days post-operatively or hospital discharge	0 per 1000	0 per 1000 (0 to 0)				
All-cause mortality	Study population		not estimable	20 (1 RCT)	⊕⊕⊖⊖ LOW ^{a,b}	No events
Up to 28 days post-operatively	0 per 1000	0 per 1000 (0 to 0)				
All-cause mortality	Study population		RR 0.95 (0.06 to 14.30)	43 (1 RCT)	⊕⊕⊖⊖ LOW ^{a,c}	-
Up to 30 days post-operatively	48 per 1000	45 per 1000 (3 to 681)				
All-cause mortality	Study population		RR 2.20 (0.45 to 10.78)	94 (1 RCT)	⊕⊖⊖⊖ VERY LOW ^{a,c,d}	-
Up to 30 days post-operatively	45 per 1000	100 per 1000				

(20 to 490)

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR:** Risk ratio

GRADE Working Group grades of evidence

High certainty: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

^aWe downgraded by one level due to low event rate.

^bWe downgraded by one level due to concerns about risks of bias both in the way that allocation of treatment was concealed at randomisation and whether blinding to treatment could be maintained for the duration of the trial in this "open-label trial where adequacy of surgical field haemostasis reported as a subjective outcome".

^cWe downgraded by one level as the study did not measure mortality as an outcome.

^dWe downgraded by one level due to concerns about risks of bias in how blinding to treatment intervention could be maintained for the duration of the trial in this "open-label trial where adequacy of surgical field haemostasis reported as a subjective outcome", and the non-reporting of all outcomes that the investigators said they would measure (selective outcome reporting).

Summary of findings 3. FIBRINOGEN: Prophylactic trials: intervention compared to inactive control for the prevention and treatment of bleeding in participants without haemophilia

FIBRINOGEN: Prophylactic trials: intervention compared to inactive control for the prevention and treatment of bleeding in participants without haemophilia

Patient or population: the prevention and treatment of bleeding in patients without haemophilia

Setting: hospital

Intervention: FIBRINOGEN: prophylactic trials

Comparison: inactive control

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	N° of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with inactive control	Risk with FIBRINOGEN in prophylactic trials				
Venous thromboembolic events	Study population		-	122 (2 RCTs)	⊕⊕⊕⊕ LOW ^{a,b}	No events in one trial
Up to 30 days post-operatively	see comment	see comment				
Arterial thrombotic events	Study population		not estimable	80 (2 RCTs)	⊕⊕⊕⊕ MODERATE ^a	-

Up to 4 days post-operatively	25 per 1000	0 per 1000 (0 to 0)				
Arterial thrombotic events	Study population		RR 0.33 (0.01 to 8.02)	116 (1 RCT)	⊕⊕⊕⊖ LOW ^{a,c}	-
Up to 30 days post-operatively	17 per 1000	6 per 1000 (0 to 138)				
Arterial thrombotic events - up to 30 days post-operatively	Study population		not estimable	122 (2 RCTs)	⊕⊕⊕⊖ LOW ^{a,b}	-
	51 per 1000	0 per 1000 (0 to 0)				
Venous thrombotic events	Study population		not estimable	80 (2 RCTs)	⊕⊕⊕⊖ MODERATE ^a	-
Up to 4 days post-operatively	0 per 1000	0 per 1000 (0 to 0)				

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR:** Risk ratio

GRADE Working Group grades of evidence

High certainty: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

^aWe downgraded by one level due to low event rate.

^bWe downgraded by one level due to concerns about risks of bias in the non-reporting of all outcomes that the investigators said they would measure (selective outcome reporting).

^cWe downgraded by one level due to concerns about risks of bias in the way that allocation of the treatment was concealed at the time of randomisation.

Summary of findings 4. FIBRINOGEN: Prophylactic trials: intervention compared to haemostatically active control for the prevention and treatment of bleeding in participants without haemophilia

FIBRINOGEN: Prophylactic trials: intervention compared to haemostatically active control for the prevention and treatment of bleeding in participants without haemophilia

Patient or population: the prevention and treatment of bleeding in patients without haemophilia

Setting: hospital

Intervention: FIBRINOGEN: prophylactic trials
Comparison: Haemostatically active control

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	N° of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with haemostatically active control	Risk with FIBRINOGEN in prophylactic trials				
Arterial thromboembolic events	Study population		not estimable	49 (1 RCT)	⊕⊕⊕⊕ LOW ^{a,b}	-
	0 per 1000	0 per 1000 (0 to 0)				
Arterial thromboembolic events - Duration of hospital stay (whole population)	Study population		not estimable	49 (1 RCT)	⊕⊕⊕⊕ LOW ^{a,b}	-
	0 per 1000	0 per 1000 (0 to 0)				
Venous thromboembolic events	Study population		not estimable	49 (1 RCT)	⊕⊕⊕⊕ LOW ^{a,b}	-
	0 per 1000	0 per 1000 (0 to 0)				
Venous thromboembolic events Duration of hospital stay (whole population)	Study population		not estimable	49 (1 RCT)	⊕⊕⊕⊕ LOW ^{a,b}	-
	0 per 1000	0 per 1000 (0 to 0)				

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR:** Risk ratio

GRADE Working Group grades of evidence

High certainty: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

^aWe downgraded by one level as the thromboembolic events (arterial or venous) in these participants given fibrinogen prophylactically were not an outcome of interest.

^bWe downgraded by one level due to the small number of participants available for this outcome.

Summary of findings 5. FXIII: Prophylactic trials: intervention compared to inactive control for the prevention and treatment of bleeding in participants without haemophilia

FXIII: Prophylactic trials: intervention compared to inactive control for the prevention and treatment of bleeding in participants without haemophilia

Patient or population: the prevention and treatment of bleeding in patients without haemophilia

Setting: hospital

Intervention: FXIII: prophylactic trials

Comparison: inactive control

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	N° of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with inactive control	Risk with FXIII in prophylactic trials				
Arterial thromboembolic events	Study population		RR 0.67 (0.28 to 1.62)	282 (2 RCTs)	⊕⊕⊕⊖ LOW ^{a,b}	-
Up to 5 to 7 weeks post-operatively	81 per 1000	54 per 1000 (23 to 131)				
Arterial thromboembolic events	Study population		RR 3.00 (0.14 to 66.53)	22 (1 RCT)	⊕⊕⊕⊖ LOW ^{b,c}	-
Up to 30 days post-operatively	0 per 1000	0 per 1000 (0 to 0)				
Arterial thromboembolic events	Study population		RR 0.20 (0.01 to 3.97)	50 (1 RCT)	⊕⊕⊕⊖ VERY LOW ^{b,c,d}	-
Post-operatively	80 per 1000	16 per 1000 (1 to 318)				
Venous thromboembolic events - Up to 5 to 7 weeks post-operatively	Study population		RR 0.93 (0.06 to 14.67)	282 (2 RCTs)	⊕⊕⊕⊖ LOW ^{a,b}	-
	7 per 1000	7 per 1000 (0 to 108)				
Venous thromboembolic events Up to 30 days post-operatively	Study population		RR 3.00 (0.14 to 66.53)	22 (1 RCT)	⊕⊕⊕⊖ LOW ^{b,c}	-
	0 per 1000	0 per 1000 (0 to 0)				
Venous thromboembolic events	Study population		not estimable	50 (1 RCT)	⊕⊕⊕⊖ VERY LOW ^{b,c,d}	-
Post-operatively	0 per 1000	0 per 1000				

(0 to 0)

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR:** Risk ratio

GRADE Working Group grades of evidence

High certainty: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

^aWe downgraded by one level due to concerns about attrition, reporting and selection bias across these trials.

^bWe downgraded by one level as thromboembolic events (arterial or venous) in these participants given factor XIII prophylactically were not an outcome of interest.

^cWe downgraded by one level due to low event rates.

^dWe downgraded by one level due to risks of bias concerns in how concealment of treatment intervention was attempted (unsealed envelopes in the patients notes) and the subsequent impact this had on the level of blinding to treatment intervention that could be maintained for the duration of the trial.

BACKGROUND

A term which we use frequently throughout this review is 'haemostatic'. Haemostatic means to cause bleeding to stop, keeping blood within a damaged blood vessel or tissue.

Description of the condition

Coagulopathy (a dysfunctional clotting response) is common in critically-ill people, both in those with and without bleeding, because of the nature of their illness. Coagulopathy is typically defined by abnormalities of standard coagulation tests performed in haematology laboratories, most commonly defined by prolongation of the prothrombin time or the activated partial thromboplastin time. These are *in vitro* tests of the time taken for a plasma sample to start to form a fibrin clot. Notably, the prothrombin time or the activated partial thromboplastin time or both are poor predictors of bleeding and there are no clear definitions of a threshold that defines a clinically significant prolongation of either test (Tripodi 2009).

In a recent, multi-centre, prospective, observational trial in adult critical care, which evaluated just under 2000 intensive-treatment-unit (ITU) admissions, one-third of patients without active bleeding were shown to develop coagulopathy during their ITU admission (Walsh 2009).

The most common current treatment for people with coagulopathy remains the transfusion of plasma. Plasma is the liquid component of blood in which different cellular elements and macromolecules are suspended. Plasma contains pro-coagulant haemostatic proteins which are a key component of the haemostatic system. These pro-coagulant haemostatic proteins, known as blood clotting factors, interact with each other, with platelets and tissues to form stable blood clots. Around one-third of people with coagulopathy received transfusions of plasma (termed fresh frozen plasma (FFP) in the UK), with around half of all FFP transfusions given to reduce a perceived risk of bleeding or prior to invasive procedures (Stanworth 2011; Walsh 2009).

However, there is uncertainty around the effectiveness of plasma transfusion in clinical practice, and the risk-benefit profile of FFP remains unknown (Yang 2012). The recognised risks associated with its use include fluid overload, transfusion-related acute lung injury, along with the generic risks associated with transfusion of allogeneic blood products (SHOT 2011). A national audit of 5000 FFP transfusion episodes to consecutive patients from 196 hospitals across England indicated little efficacy for plasma transfusion, with very small median change in duration of prothrombin time following FFP transfusion (Stanworth 2010).

Description of the intervention

Given concerns about the effectiveness of plasma transfusion, there is increasing interest in the targeted use of specific or purified intravenous pro-coagulant haemostatic factors. Many different intravenous pro-coagulant haemostatic factors exist, including those derived from plasma as concentrates, or those developed as specific recombinant factors. The advantages of plasma-derived coagulation factor concentrates include greater standardisation (whether at the stage of production or during manufacture) and reduced infective and compatibility complications when compared with blood transfusions through the application of techniques such as pasteurisation, nanofiltration and adsorption

to remove viruses. The advantages of recombinant-derived factors include potentially unlimited supply, with reduced reliance on the availability of blood donors, as well as negligible infective risk. Finally, both recombinant and plasma-derived intravenous concentrates offer the potential advantage of providing specific pro-coagulant haemostatic factors more rapidly and at higher concentrations than is possible with the use of traditional plasma transfusion.

Some of the plasma-derived coagulation factor concentrates contain multiple pro-coagulant haemostatic factors. Prothrombin complex concentrates are plasma-derived coagulation factor concentrates that contain three or four (vitamin K-dependent) factors at high concentration.

Two observational cohort studies have examined the use of prothrombin complex concentrates as an adjunct to fibrinogen concentrate in the management of bleeding following trauma. Whilst neither of these studies demonstrated reduced mortality in those receiving prothrombin complex concentrates, one of the studies demonstrated an association between receiving prothrombin complex concentrates and a reduced transfusion requirement (Görlinger 2011). Safety data on the use of prothrombin complex concentrates are limited. In particular, rates of thrombosis, both arterial and venous, and disseminated intravascular coagulation need to be fully assessed (Grottke 2011).

In contrast to concentrates containing multiple factors, single pro-coagulant haemostatic factors allow the treatment of individual clotting factor deficiencies to be specifically targeted. A specific concentrate of the key pro-coagulant haemostatic factor fibrinogen (or factor I) is available, although currently only licensed for use in many countries for people with inherited disorders of fibrinogen. The observation that normal-to-high levels of fibrinogen (in the range 1.5 to 4.0 g/L) prior to the onset of major obstetric haemorrhage are associated with improved outcomes (Charbit 2007; de Lloyd 2011), has led clinicians to consider targeted therapy with concentrates of fibrinogen in this cohort of women, rather than the more traditional use of FFP.

How the intervention might work

Specific coagulation factor concentrates are presumed to stop or prevent bleeding by providing a source of pro-coagulant haemostatic factor(s) which, alongside contributions from platelets and the endothelium, interact to produce a stronger fibrin clot. The rationale for their use is based on either restoring or increasing levels of pro-coagulant haemostatic factors, or activating clotting systems by infusing recombinant activated complexes to initiate coagulation.

It is hypothesised that pro-coagulant haemostatic factors administered intravenously might work more effectively than plasma (either FFP or cryoprecipitate) to deliver a more concentrated source of factors which will raise factor levels more effectively and predictably using a standardised product. Given that the current processes for thawing plasma would not apply to pro-coagulant haemostatic factors, it is also likely that these agents would be more rapidly available for use in emergency situations.

However, as for all pro-coagulant haemostatic agents, adverse events are recognised. In particular, use of these agents might be expected to be associated with a greater rate of thromboembolic

events. This has been clearly demonstrated in meta-analyses of RCTs of recombinant factor VIIa in people without haemophilia (Levi 2010; Simpson 2012).

Why it is important to do this review

In a short period there has been a rapid shift in the clinician's approach to the management of coagulopathy and bleeding, with increasing interest in the use of specific intravenous pro-coagulant haemostatic factors instead of the current standard plasma transfusion. Because of the real uncertainties about the effectiveness and safety of FFP, wider use of the pro-coagulant haemostatic factors and evaluation in clinical trials are anticipated, not least to understand their benefits given their increased cost in comparison to plasma transfusion. There is increasing understanding of the complex changes that occur in people with acquired major bleeding, particularly after trauma and parturition, and it is clear that many clotting factor levels change during active bleeding.

This review explores both the use of single clotting factor concentrate therapy (at the time of publication these were fibrinogen, factor XIII and a prothrombin complex concentrate composite) but we also anticipate that clotting factor concentrates will be evaluated in combination to treat the complexity of clotting changes that occurs in major haemorrhage. Lessons should also be learned from the story of rFVIIa, which was 'marketed' as a universal haemostatic agent in people without haemophilia but was then found, as new randomised trials were published, to have uncertain benefit with harm (increased arterial thromboembolic events; Levi 2010; Simpson 2012), emphasising the importance of a systematic review to identify and appraise the RCT literature as it evolves. This Cochrane Review builds on the experience gained from the rFVIIa Cochrane Review (Simpson 2012), and comprehensively identifies and critically appraises all the RCT evidence assessing the use of intravenous pro-coagulant haemostatic factors.

This systematic review differs from two other 'clotting factor concentrate' Cochrane Reviews (Wikkelsø 2013; Johansen 2015). Our review did not assess the use of pro-coagulant haemostatic factors and factor concentrates for the reversal of vitamin K antagonist treatment in bleeding and non-bleeding people, which is the focus of Johansen 2015. Our review does not include trials where the intervention was recombinant factor VIIa, antifibrinolytic therapy or desmopressin, as is the case in Wikkelsø 2013. Our review includes trials in which the participant group were neonates, unlike Wikkelsø 2013 which excluded them.

This systematic review indicates where high-quality evidence is lacking, so informing healthcare professionals where more evidence is required before these expensive drugs are widely accepted into clinical practice.

OBJECTIVES

To assess the effects and safety of pro-coagulant haemostatic factors and factor concentrates in the prevention and treatment of bleeding in people without haemophilia.

METHODS

Criteria for considering studies for this review

Types of studies

We included only randomised controlled trials (RCTs), regardless of language, date or place of publication. We included open-label and unblinded studies.

Types of participants

We included adults and children (with no age restriction) either at risk of bleeding or already bleeding due to any cause. We excluded people with known inherited bleeding disorders (e.g. haemophilia) unless any data were separately available for the participants without the inherited condition. We excluded studies of healthy volunteers.

Types of interventions

We assessed separately whether a pro-coagulant haemostatic factor concentrate was administered as a treatment or in a prophylactic manner for the following comparisons:

- a pro-coagulant haemostatic factor concentrate (either alone or in combination with another pro-coagulant haemostatic factor concentrate) versus either placebo or current best/standard treatment (e.g. plasma/FFP).
- one pro-coagulant haemostatic factor concentrate (either alone or in combination with another pro-coagulant haemostatic factor concentrate) versus another different pro-coagulant haemostatic factor concentrate (either alone or in combination with another pro-coagulant haemostatic factor concentrate).

Each of the above combinations could be given in any dose and for any duration of treatment.

We included RCTs of the administration of prothrombin complex concentrates if used for conditions other than reversal of vitamin K antagonist therapy.

We only included studies where pro-coagulant haemostatic factors were administered intravenously.

We excluded:

- pro-coagulant haemostatic factors administered topically, e.g. thrombin or fibrin glue, as they are the subject of a previous Cochrane Review (Carless 2009).
- RCTs of desmopressin, tranexamic acid and aminocaproic acid, due to their different mechanisms of action from other pro-coagulant haemostatic agents. We also excluded RCTs in which a pro-coagulant haemostatic agent was used exclusively to reverse warfarin (or a similar agent), but after the review had been started.
- Studies exclusively administering rFVIIa, as they have been the subject of a previous Cochrane Review (Simpson 2012). However, studies in which rFVIIa was a co-intervention with other intravenous pro-coagulant haemostatic factors (i.e. in separate comparator arms or co-administration of rFVIIa and another factor/reagent) were eligible for inclusion.

We are aware that studies of fibrinogen concentrate in bleeding have been the focus of a previous Cochrane Review (Wikkelsø 2013),

but as this is a rapidly expanding field with a significant number of new studies published since the [Wikkelsso 2015](#) review, and as we have presented the analysis from an alternative perspective, we have included the fibrinogen concentrate studies in this review,

Types of outcome measures

We presented outcome data from all identified RCTs by prophylactic and therapeutic indications for use. In prophylactic use, the agent was given to prevent bleeding which may be anticipated prior to or during a surgical operation; in therapeutic use, the agent was given to treat bleeding which is established and ongoing.

There was variation in the time points at which each outcome was measured, due to the nature of the participants and conditions across the trials; we took this into account when reporting and analysing the outcome data for the included trials. Most of the trials reported outcomes acutely postpartum and in the 24 hours following any given intervention or procedure, when compared to baseline measures for outcomes such as blood loss and transfusion requirement. These formed our main time point comparators for these outcomes. In addition, where outcome data were reported for other time points we have also included these data. For mortality, arterial and venous thromboembolic events, bleeding and adverse events, data were presented at different time points in each trial, making it difficult to generate a cohesive picture.

Primary outcomes

- Mortality (all causes) (principal primary outcome for therapeutic intervention usage)
- The number of participants suffering arterial and/or venous thromboembolic events (principal primary outcome for prophylactic intervention usage)

Secondary outcomes

- Mortality due to bleeding
- Red blood cell transfusion requirement (both volume of transfusion and the receipt of a transfusion)
- Blood loss
- Allergic reaction
- Change to laboratory measures of coagulation, both standard laboratory tests and global measurements of coagulation (such as thromboelastography (TEG) or rotational thromboelastometry (ROTEM), or thrombin generation)

Search methods for identification of studies

The Systematic Review Initiative's Information Specialist (CD) formulated the search strategies used, in collaboration with the Cochrane Injuries Group.

Electronic searches

We searched for RCTs in the following electronic databases between the dates listed below. There were no restrictions by language or publication status. All searches were undertaken on 18 April 2018 and the search strategies for all sources are detailed in [Appendix 1](#).

- Cochrane Central Register of Controlled Studies (CENTRAL) (the Cochrane Library 2018, Issue 3) on 18 April 2018;

- MEDLINE (OvidSP, 1948 to 18 April 2018);
- Embase (OvidSP, 1974 to 18 April 2018);
- CINAHL (EBSCOHOST, 1938 to 18 April 2018);
- PubMed, in process and epublications ahead of print only, on 18 April 2018;
- PROSPERO on 18 April 2018;
- Transfusion Evidence Library (www.transfusionevidencelibrary.com, 1950 to 18 April 2018);
- LILACS (from 1980 to 18 April 2018);
- IndMed (from 1985 to 18 April 2018);
- KoreaMed (from 1934 to 18 April 2018);
- Web of Science Conference Proceedings Citation Index - Science (CPCI-S, from 1990 to 18 April 2018).

We also searched for ongoing trials in the following databases. We undertook all searches on 18 April 2018 and the search strategies are detailed in [Appendix 1](#).

- ClinicalTrials.gov (www.clinicaltrials.gov);
- EU Clinical Trials Register (EUDRACT, www.clinicaltrialsregister.eu);
- WHO International Clinical Trials Registry Platform (ICTRP, www.who.int/ictpr)
- ISRCTN Register (www.isrctn.com/)

We combined searches in MEDLINE, Embase and CINAHL with adaptations of the Cochrane RCT search filter, as detailed in Chapter 6 of the *Cochrane Handbook for Systematic Reviews of interventions* ([Higgins 2011](#)).

Searching other resources

We augmented database searching by:

- Handsearching of reference lists; we checked references of all identified trials and relevant review articles for further literature. We limited these searches to 'first generation' reference lists;
- Personal contacts; we contacted authors of relevant trials, trial groups and experts worldwide who are known to be active in the field for unpublished material or further information on ongoing trials.

Data collection and analysis

Selection of studies

One author (CD) initially screened all search results for relevance against the eligibility criteria and discarded all those that were clearly irrelevant. Thereafter, three review authors (JF, NC, YL) independently screened all the remaining results (titles, abstracts and full text) for relevance against the full eligibility criteria. We retrieved full-text papers for all references for which a decision of eligibility was not made from the title and abstract alone. We resolved differences of opinion through discussion and consensus, where necessary, with recourse to a fourth review author (SB). Trials that did not meet our eligibility criteria are detailed in the [Characteristics of excluded studies](#) table.

Data extraction and management

Five review authors (JF, NC, SB, YL, GS) independently extracted data onto standardised forms. Two of the review authors (JF, GS) piloted these forms on two included RCTs and made changes

that were agreed upon as needed. Throughout the data extraction process, we resolved disagreements by consensus, with recourse to a third review author (SS) as needed. The review authors were not blinded to the names of study authors, institutions, journals or the outcomes of the trials. We extracted data on the setting of the trial, the methods and statistical assumptions made, the characteristics of participants and interventions and details of the outcomes measured and the timing of the outcome assessments.

Where studies describe red blood cell transfusion in units (Jeppsson 2016; Karlsson 2011; Lance 2012; Najafi 2014; Sadeghi 2014), we converted these to millilitres to permit comparison with other studies reporting millilitres of red blood cells transfused. We have used a conversion ratio of one unit of red blood cells equivalent to 300 millilitres.

Assessment of risk of bias in included studies

Three review authors (JF, SB, NC) independently assessed all included trials for possible risks of bias, as described in Chapter 8 of the *Cochrane Handbook* (Higgins 2011). We classified trials as being at low, high or unclear risk of bias. To assess risks of bias, the authors included the following questions in the 'Risk of bias' table for each included trial.

- Was the allocation sequence adequately generated?
- Was allocation adequately concealed?
- Was knowledge of the allocated intervention adequately prevented (i.e. blinded) throughout the trial for participants, trial personnel and outcome assessors?
- Were incomplete outcome data adequately addressed for the main outcome?
- Are reports of the trial free of selective outcome reporting?
- Was the trial apparently free of other problems that could put it at risk of bias?

We anticipated that knowledge of the allocated intervention would be difficult to blind from clinicians and participants. However we felt that it would be possible to blind knowledge of the allocated intervention from outcome assessors. We bore these considerations in mind when making an assessment of the risk of bias for the adequacy of blinding of participants, trial personnel and outcome assessors.

Measures of treatment effect

We present dichotomous outcomes as relative risks (RRs) with 95% confidence intervals (CIs). We present continuous outcomes as mean differences (MDs) with 95% confidence intervals. Where data are not presented in this way, but we were able to make adjustments from the data given, we did this for the following studies:

- [Godje 2006](#); we present the data in tabular form without quoting numeric values. We extracted these data quantitatively using digital measurement of the relevant table data and axes.

A number of studies reported outcome data for blood loss and transfusion requirement using median values with interquartile ranges (IQRs) (12 and 17 studies, respectively); we have commented on these values in the text and tabulated them in [Table 1](#) and [Table 2](#).

Unit of analysis issues

In the included trials the participant is the unit of analysis.

We did not encounter unit-of-analysis issues. However, if in any future updates of this review we identify any cluster-randomised trials we will treat these in accordance with the advice given in the *Cochrane Handbook* (Higgins 2011).

Dealing with missing data

We contacted study authors by email to request missing data or to clarify data that were presented. The list of authors we contacted and the information obtained is outlined below.

- [Collins 2017](#) provided additional data about mortality.
- [Curry 2018](#) provided additional data about arterial and venous thromboembolic adverse events and the number of participants exposed to a red blood cell transfusion.
- [Haas 2015](#) (C) and [Haas 2015](#) (S) provided additional data about the volume of red blood cell units transfused, the number of participants exposed to a red blood cell transfusion, mortality, bleeding mortality and arteriovenous thromboembolic events.
- [Karkouti 2013](#) provided additional data on adverse events.
- [Korte 2009](#) provided additional information on the number of participants in each study arm.
- [Lance 2012](#) provided additional data about outcome time points, mortality, adverse events and red blood cell transfusion.
- [Ranucci 2015](#) provided additional information about the number of participants exposed to a red blood cell transfusion.
- [Sabate 2016](#) provided additional data on the incidence of bleeding mortality, arterial and venous thromboembolic events and the volume of red blood cell units transfused.
- [Innerhofer 2017](#), [Jeppsson 2016](#), [Karlsson 2011](#) and [Wikkelsø 2015](#) were unable to provide any additional data that we could include in this review.
- [Bilecen 2017](#); [Cui 2010](#); [Galas 2014](#); [Levy 2009](#); [Najafi 2014](#); [Nascimento 2016](#); [Pournajafian 2015](#); [Rahe-Meyer 2013](#); [Rahe-Meyer 2016](#); [Sadeghi 2014](#); [Soleimani 2016](#); and [Tanaka 2014](#) provided no response to our communication.
- The contact email addresses of [Bregenzer 1999](#); [Fenger Erikson 2009](#); [Godje 2006](#) generated an automatic failure-to-deliver email communication and no alternative contact details were available.
- There were no relevant or usable contact details provided in [Rasche 1982](#); [Shirahata 1990](#); [Turner 1981](#), and no alternative contact details were available.

Method of analysis

Of the 31 included trials, 11 declared an intention-to-treat (ITT) analysis, 13 outlined per protocol analysis methods and seven studies did not identify their method of analysis. Due to this, and the paucity of data that could be usefully harmonised on participant cohort, intervention and outcome time point we have included per protocol data.

Assessment of heterogeneity

We assessed clinical heterogeneity based on the data extracted about the characteristics of the included trials. We have discussed possible explanations for the observed heterogeneity within the review.

Where we considered trials to be clinically homogeneous, we conducted meta-analysis and assessed statistical heterogeneity of treatment effects between trials by using a χ^2 test with a significance level at $P < 0.1$. We used the I^2 index to quantify the amount of possible heterogeneity ($I^2 > 30\%$ as moderate heterogeneity and $I^2 > 75\%$ as considerable heterogeneity). When we found heterogeneity, we explored its potential causes with prespecified subgroup and sensitivity analyses.

Assessment of reporting biases

We made every effort to identify unpublished trials relevant to this review. We accept that funnel-plot asymmetry, of which publication bias is one cause, is difficult to detect with the small number of trials (i.e. less than 10) often encountered in systematic reviews. In future updates, we will assess publication bias using funnel plots if we include 10 or more trials in any of our predefined comparison subgroups.

Data synthesis

We undertook meta-analysis where trials were sufficiently homogeneous in terms of clinical setting and trial design and there were sufficient data of a suitable type. We used the Cochrane statistical software Review Manager 5 (Review Manager 2014). We used a fixed-effect model for combining data in the first instance. Where we identified heterogeneity in a fixed-effect meta-analysis, we noted this and repeated the analysis using a random-effects model. We did not remove any study from any analysis due to a high risk of bias. We grouped outcome data into the outcomes listed earlier in the [Types of outcome measures](#) section.

As well as quantitative synthesis, the overall interpretation of the data incorporated insights from a qualitative summary. We based our conclusions on patterns of results identified across clearly-tabulated results of included trials, as well as summary measures. We considered both the direction and magnitude of any effect. We document individual trial results in the 'Additional Tables' section of the review.

For meta-analysis, we log-transformed blood loss outcome data and conducted the analysis using the transformed values. A meta-analysis (using arithmetic means) of the differences in means using the transformed blood loss data corresponds to a meta-analysis (using geometric means) of the ratio of means in the original scale.

Subgroup analysis and investigation of heterogeneity

We performed one subgroup analysis due to heterogeneity, using the first of our predefined subgroups.

- Children versus adults (because different definitions of coagulopathy apply). We defined adults as being 18 years or older at the time of recruitment.
- Trials of participants with brain/neurological injuries. For example, acute intracranial injury versus no acute intracranial injury (in view of the added effect of central nervous system injury on coagulopathy and therefore the possibly different efficacy of interventions in this participant group).

We did not use the second of our predefined subgroups as these data were sparse in the included studies.

Sensitivity analysis

We assessed the robustness of our findings by performing sensitivity analyses for the trials where 1) allocation concealment was adequate, and 2) in which 25% of participants or fewer were lost in the one meta-analysis with noted heterogeneity.

'Summary of findings' tables

We created 'Summary of findings' tables for each intervention, following guidance in the *Cochrane Handbook* (Higgins 2011). We assessed the quality of the evidence for each outcome using GRADE (Schünemann 2011). The outcomes included in the 'Summary of findings' tables are:

- All-cause mortality for trials in which the indication of use for the intervention was therapeutic;
- Arterial and venous thromboembolic complications for trials in which the indication of use for the intervention was prophylactic.

RESULTS

Description of studies

Results of the search

The search strategy identified 4781 references with an additional 379 ongoing studies; these were reduced to 2770 references plus 319 ongoing studies once duplicates had been removed. Of these, 2681 references and 288 ongoing studies were ruled out for being clearly irrelevant to the scope of this review. We screened the remaining 89 references and 31 ongoing trials for eligibility. We included 31 studies (from 61 references) and identified 22 ongoing trials (from 25 references); we excluded 14 trials (from 28 references) and six ongoing studies (from six references). Full details are reported in the PRISMA flow diagram (Figure 1).

Figure 1. Trial flow diagram.

Figure 1. (Continued)

We found four non-English publications. Two papers required translation from German to assess eligibility for inclusion (Rasche 1982; Scherer 1994). Following translation, we deemed one paper ineligible (Scherer 1994), whilst we rated the second trial as eligible for inclusion (Rasche 1982). A third paper was translated from Portuguese using Google translate and we included it in the review (Sadeghi 2014). The fourth paper was translated from Persian by a volunteer translator found through the Cochrane TaskExchange (Pournajafian 2015), and was also included.

One reference reported data from two distinct surgical groups: scoliosis surgery and craniostylosis surgery (Haas 2015). We report these data as two separate studies throughout the review: Haas 2015 (S) and Haas 2015 (C).

The 31 included studies randomised a total of 2392 participants. There were 13 therapeutic trials that randomized 1057 participants (ranging from 20 to 249) and 18 prophylactic trials that randomised a total of 1335 participants (ranging from 20 to 479). Two trials had three arms (Godje 2006; Karkouti 2013), and one trial had five arms (Levy 2009). All three were prophylactic trials.

Included studies

We give further details about the included studies in the [Characteristics of included studies](#) table and [Table 3](#); [Table 4](#); [Table 5](#); [Table 6](#); [Table 7](#); [Table 8](#) and [Table 9](#).

Setting

Three trials were multicentre, multinational trials (Karkouti 2013; Levy 2009; Rahe-Meyer 2016) and six were multicentre trials, based in a single country: Germany (Bregenzer 1999), the UK (Collins 2017; Curry 2018), Spain (Sabate 2016), Sweden (Jeppsson 2016) and Denmark (Wikkelsø 2015). The remaining 22 trials were single-centre trials. Four trials were conducted in Iran (Najafi 2014; Sadeghi 2014; Pournajafian 2015; Soleimani 2016), three trials were conducted in Germany (Godje 2006; Rahe-Meyer 2013; Rasche 1982), three trials were conducted in Switzerland (Haas 2015 (S); Haas 2015 (C); Korte 2009) and the Netherlands (Lance 2012; Bilecen 2017), whilst one trial was each conducted in Brazil

(Galas 2014), Austria (Innerhofer 2017), Canada (Nascimento 2016), China (Cui 2010), Denmark (Fenger Erikson 2009), Great Britain (Turner 1981), Italy (Ranucci 2015), Japan (Shirahata 1990), Sweden (Karlsson 2011), and the USA (Tanaka 2014).

The clinical setting varied across the trials: 22 were in an elective surgical setting (Cui 2010; Fenger Erikson 2009; Galas 2014; Godje 2006; Haas 2015 (S) and (C); Karkouti 2013; Karlsson 2011; Korte 2009; Lance 2012; Levy 2009; Najafi 2014; Rahe-Meyer 2013; Ranucci 2015; Sadeghi 2014; Tanaka 2014; Bilecen 2017; Jeppsson 2016; Pournajafian 2015; Rahe-Meyer 2016; Sabate 2016; Soleimani 2016), five in an urgent medical setting (Wikkelsø 2015; Collins 2017; Curry 2018; Innerhofer 2017; Nascimento 2016), and four in a non-urgent medical setting (Bregenzer 1999; Rasche 1982; Shirahata 1990; Turner 1981). Definitions of elective and urgent have been taken from the trial manuscript.

Participants

Three trials were undertaken in infants (Haas 2015 (C); Shirahata 1990; Turner 1981), and three in paediatric populations (Cui 2010; Galas 2014; Haas 2015 (S)); the remaining 25 trials enrolled adult participants. We considered and analysed the infant trials separately for all outcomes other than allergic adverse events, where they have been grouped with the adult studies. We have included the paediatric population trials in analyses alongside the trials with adult populations, as we consider the populations in the three paediatric trials (Cui 2010, mean age 38.4 months; Haas 2015 (S), mean age of 12 years and Galas 2014, participants under 15 years of age) to be physiologically comparable to the populations in the trials recruiting adults.

Interventions

Prophylactic trials

Eighteen trials addressed prophylactic use of pro-coagulant factors, 16 in comparison to inactive agents (Cui 2010; Fenger Erikson 2009; Godje 2006; Karkouti 2013; Karlsson 2011; Korte 2009; Levy 2009; Najafi 2014; Pournajafian 2015; Ranucci 2015; Rasche 1982; Sabate 2016; Sadeghi 2014; Shirahata 1990; Soleimani 2016;

Turner 1981), and two in comparison to an active comparator (Haas 2015 (C); Haas 2015 (S)).

Inactive comparator trials

Of the 16 inactive comparator trials, seven recruited participants undergoing cardiac surgery (Cui 2010; Godje 2006; Karkouti 2013; Karlsson 2011; Levy 2009; Ranucci 2015; Sadeghi 2014). The other trials focused on elective cystectomy (Fenger Erikson 2009), gastrointestinal cancer (Korte 2009), acute leukaemia (Rasche 1982), spinal fusion (Pournajafian 2015), liver transplantation (Sabate 2016), trans-urethral prostatic resection (Soleimani 2016), and elective hip replacement arthroplasty (Najafi 2014). Both neonatal population trials used inactive comparators and focused on babies at risk of intracranial haemorrhage (Shirahata 1990; Turner 1981).

Six of these prophylactic trials used factor XIII as the pro-coagulant haemostatic intervention (Godje 2006; Karkouti 2013; Korte 2009; Levy 2009; Rasche 1982; Shirahata 1990), and nine used fibrinogen (Cui 2010; Fenger Erikson 2009; Karlsson 2011; Najafi 2014; Pournajafian 2015; Ranucci 2015; Sabate 2016; Sadeghi 2014; Soleimani 2016). Turner 1981 used a range of agents, including a prothrombin complex concentrate supplied by the Scottish National Blood Transfusion Service Protein Fractionation Centre: fibrinogen, factor VIII and platelets.

The inactive comparator agents were haemostatically inactive placebo products in 10 trials (Fenger Erikson 2009; Karkouti 2013; Korte 2009; Levy 2009; Najafi 2014; Pournajafian 2015; Ranucci 2015; Sabate 2016; Sadeghi 2014; Soleimani 2016), or nothing other than normal care (Cui 2010; Godje 2006; Karlsson 2011; Rasche 1982; Shirahata 1990; Turner 1981).

Active comparator trials

The participants in the two trials with an active comparator were neonates undergoing craniosynostosis or paediatric patients undergoing scoliosis surgery: both compared two doses of fibrinogen concentrate (Haas 2015 (C); Haas 2015 (S)).

Therapeutic trials

Thirteen trials addressed therapeutic use of pro-coagulant factors: nine in comparison to inactive agents (Bilecen 2017; Bregenzner 1999; Collins 2017; Curry 2018; Jeppsson 2016; Nascimento 2016; Rahe-Meyer 2013; Rahe-Meyer 2016; Wikkelsø 2015), and four compared to an active comparator (Galas 2014; Innerhofer 2017; Lance 2012; Tanaka 2014).

Inactive comparator trials

The inactive comparator trials focused on cardiac surgery (Bilecen 2017; Jeppsson 2016; Rahe-Meyer 2013; Rahe-Meyer 2016), ulcerative colitis (Bregenzner 1999), trauma (Curry 2018; Nascimento 2016), and postpartum haemorrhage (Collins 2017; Wikkelsø 2015). The pro-coagulant haemostatic interventions were fibrinogen in eight studies (Bilecen 2017; Collins 2017; Curry 2018; Jeppsson 2016; Nascimento 2016; Rahe-Meyer 2013; Rahe-Meyer 2016; Wikkelsø 2015), and factor XIII (Bregenzner 1999). The comparators in all nine trials were haemostatically inactive placebo products.

Active comparator trials

The active comparator trials addressed cardiac surgery (Galas 2014; Tanaka 2014), trauma (Innerhofer 2017), and major elective surgery

(Lance 2012). The intervention in all these trials was fibrinogen compared with FFP (Innerhofer 2017), cryoprecipitate (Galas 2014), platelets (Tanaka 2014), and as a partial replacement for FFP (Lance 2012).

Co-interventions

Most trials documented the use of blood products (red blood cells and platelets) as well as transfusion of clotting products (FFP, prothrombin complex concentrate, cryoprecipitate) to treat bleeding or clotting derangement during the course of the trial.

Non-blood product co-interventions were reported by 26 trials. One trial reported routine administration of Vitamin K to all participants (Turner 1981), 11 trials reported administration of protamine (Bilecen 2017; Cui 2010; Godje 2006; Jeppsson 2016; Karkouti 2013; Karlsson 2011; Lance 2012; Levy 2009; Rahe-Meyer 2013; Rahe-Meyer 2016; Ranucci 2015), 14 trials reported the use of an antifibrinolytic agent (Bilecen 2017; Bregenzner 1999; Collins 2017; Curry 2018; Haas 2015 (C); Haas 2015 (S); Innerhofer 2017; Karkouti 2013; Nascimento 2016; Rahe-Meyer 2013; Rahe-Meyer 2016; Ranucci 2015; Sabate 2016; Wikkelsø 2015) and two trials documented the use of post-intervention low-molecular weight heparin (Karlsson 2011; Korte 2009). Five trials did not report details of co-interventions administered or outline additional transfusion policies (Galas 2014; Najafi 2014; Sadeghi 2014; Shirahata 1990; Soleimani 2016).

Primary outcomes of included trials

Twelve trials reported transfusion requirement, or the avoidance of transfusion, as a primary outcome (Collins 2017; Haas 2015 (C); Haas 2015 (S); Karkouti 2013; Najafi 2014; Rahe-Meyer 2013; Rahe-Meyer 2016; Ranucci 2015; Sabate 2016; Sadeghi 2014; Tanaka 2014; Wikkelsø 2015). Four trials reported blood loss as the primary outcome (Bilecen 2017; Galas 2014; Jeppsson 2016; Soleimani 2016), and two trials each had primary outcomes of adverse events (Karlsson 2011; Levy 2009), or clot firmness following surgery (Fenger Erikson 2009; Korte 2009). Other primary outcomes were time to chest wall closure (Cui 2010), the incidence of multiorgan failure (Innerhofer 2017), mortality (Turner 1981), time to cessation of bleeding (Bregenzner 1999), clotting time and clot formation time (Lance 2012) and bleeding complications (Rasche 1982). Two trials were feasibility studies (Curry 2018; Nascimento 2016), and three studies did not report their primary outcome (Godje 2006; Pournajafian 2015; Shirahata 1990).

Funding sources

Seventeen trials received support from a commercial company for their study: CSL Behring (Bilecen 2017; Collins 2017; Curry 2018; Fenger Erikson 2009; Galas 2014; Haas 2015; Karlsson 2011; Lance 2012; Nascimento 2016; Rahe-Meyer 2013; Rahe-Meyer 2016; Ranucci 2015; Tanaka 2014), Novo Nordisk (Karkouti 2013; Korte 2009; Levy 2009), and Centeon Pharma GmbH (Bregenzner 1999). Eight of these trials also received support from an academic institution or charity (Bregenzner 1999; Collins 2017; Fenger Erikson 2009; Galas 2014; Karlsson 2011; Korte 2009; Nascimento 2016; Ranucci 2015).

Four studies were solely supported by grants from an academic institution or a charity (Jeppsson 2016; Sabate 2016; Sadeghi 2014; Wikkelsø 2015). Eight trials did not report receipt of any funding (Cui 2010; Godje 2006; Innerhofer 2017; Najafi 2014; Rasche 1982;

[Shirahata 1990](#); [Soleimani 2016](#); [Turner 1981](#)), whilst in one trial it was unclear whether or not funding had been received to support the trial ([Pournajafian 2015](#)).

Excluded studies

For the full list of excluded trials see [Characteristics of excluded studies](#).

We excluded six trials as the participants were ineligible: they were on warfarin and received the pro-coagulant haemostatic factor for anticoagulation reversal ([Boulis 1999](#); [Demeyere 2010](#); [Goldstein 2015](#); [Papakitsos 2014](#); [Sarode 2013](#); [van Aart 2006](#)). We excluded one trial after translation from German and review of the full text for not being a randomised controlled trial ([Scherer 1994](#)). One study was excluded on review of the full text as it was a review of the literature ([Maegle 2015](#)). We excluded five studies as the study

interventions did not meet the inclusion criteria for this review ([Curry 2015](#); [Garrigue 2017](#); [Masoumi 2016](#); [Miyata 2014](#); [Qiu 2016](#)). One study was excluded as it was terminated early for feasibility reasons after recruiting only a single participant to each study arm ([TOP-CLOT Trial](#)).

We identified 22 trials as ongoing studies. Full details of these trials are available in the [Characteristics of ongoing studies](#) table.

Risk of bias in included studies

Please refer to [Figure 2](#) for visual representations of the assessments of risk of bias across all trials and for each item in the included trials. See the 'Risk of bias' section in the [Characteristics of included studies](#) for more detailed information about the biases identified within the individual trials.

Figure 2. Risk of bias summary: review authors' judgements about each risk of bias item for each included trial. Thirty-one trials are included in this review (Haas 2015 reports two trials).

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants (performance bias)	Blinding of study personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Bilecen 2017	+	+	+	+	+	?	+	+
Bregenzer 1999	?	?	+	+	+	-	-	?
Collins 2017	+	+	+	+	?	+	+	+
Cui 2010	?	?	+	-	+	?	+	?
Curry 2018	+	+	+	+	?	+	?	?
Fenger Erikson 2009	?	?	+	+	+	+	+	+
Galas 2014	+	+	+	?	+	+	+	+
Godje 2006	?	-	+	-	-	+	+	?
Haas 2015	?	+	+	-	?	+	+	?
Innerhofer 2017	+	+	+	-	-	+	-	+
Jeppsson 2016	?	+	+	+	?	+	+	+
Karkouti 2013	+	?	+	-	+	+	+	-
Karlsson 2011	?	?	+	+	+	?	+	+
Korte 2009	+	+	+	+	?	?	+	?
Lance 2012	?	-	+	?	+	+	?	+
Lewy 2009	+	+	+	+	?	-	-	-
Najafi 2014	+	?	?	?	?	+	-	?
Nascimento 2016	+	+	+	+	?	+	+	?
Pournajafian 2015	?	?	+	?	?	-	?	?
Rahe-Meyer 2013	+	+	+	+	+	+	+	-
Rahe-Meyer 2016	?	?	+	+	?	+	+	-
Rosinski 2015	+	-	+	+	?	+	+	?

Figure 2. (Continued)

Rahe-Meyer 2016	+	+	+	+	+	+	+
Ranucci 2015	+	+	+	+	+	+	+
Rasche 1982	+	+	+	+	+	+	+
Sabate 2016	+	+	+	+	+	+	+
Sadeghi 2014	+	+	+	+	+	+	+
Shirahata 1990	+	+	+	+	+	+	+
Soleimani 2016	+	+	+	+	+	+	+
Tanaka 2014	+	+	+	+	+	+	+
Turner 1981	+	+	+	+	+	+	+
Wikkelsso 2015	+	+	+	+	+	+	+

Allocation

Random sequence generation

Prophylactic trials

We considered nine trials to be at low risk of bias, as randomisation was done centrally by a Voice response system (Karkouti 2013; Levy 2009), or was randomly generated (Korte 2009; Najafi 2014; Ranucci 2015; Sabate 2016; Sadeghi 2014; Soleimani 2016; Turner 1981). We judged nine trials to be at unclear risk of bias, as the method of randomisation was not adequately reported (Cui 2010; Fenger Erikson 2009; Godje 2006; Haas 2015 (C); Haas 2015 (S); Karlsson 2011; Pournajafian 2015; Rasche 1982; Shirahata 1990).

Therapeutic trials

We considered seven trials to be at low risk of bias, as randomisation was done centrally (Curry 2018; Rahe-Meyer 2013; Wikkelsso 2015), or was randomly generated (Bilecen 2017; Galas 2014; Innerhofer 2017; Nascimento 2016). We judged six trials to be at unclear risk of bias, as the method of randomisation was not adequately reported (Bregenzer 1999; Collins 2017; Jeppsson 2016; Lance 2012; Rahe-Meyer 2016; Tanaka 2014).

Concealment of treatment allocation

Prophylactic trials

We considered five trials to be at low risk of bias for allocation concealment, as assignment was done centrally (Levy 2009; Sabate 2016) or was randomly generated (Haas 2015 (C); Haas 2015 (S); Korte 2009). We considered two trials to be at high risk of bias due to incorrect methods being used to conceal treatment allocation (Godje 2006; Ranucci 2015). We judged 11 trials to be at unclear risk of bias because no description of allocation concealment was provided (Cui 2010; Fenger Erikson 2009; Karkouti 2013; Karlsson 2011; Najafi 2014; Pournajafian 2015; Rasche 1982; Sadeghi 2014; Shirahata 1990; Soleimani 2016; Turner 1981).

Therapeutic trials

We considered seven trials to be at low risk of bias for allocation concealment, as assignment was done centrally (Rahe-Meyer 2013)

or was randomly generated (Bilecen 2017; Curry 2018; Galas 2014; Innerhofer 2017; Jeppsson 2016; Nascimento 2016). We considered three trials to be at high risk of bias, due to incorrect methods being used to conceal treatment allocation (Lance 2012; Tanaka 2014; Wikkelsso 2015). We judged three trials to be at unclear risk of bias, because no description of allocation concealment was provided (Bregenzer 1999; Collins 2017; Rahe-Meyer 2016).

Blinding

Blinding of participants and personnel (performance bias)

Prophylactic intervention

We considered 16 trials to be at low risk of **participant** performance bias as, i) participants were under anaesthesia when their intervention was administered (Cui 2010; Fenger Erikson 2009; Godje 2006; Haas 2015 (C); Haas 2015 (S) Karkouti 2013; Karlsson 2011; Korte 2009; Levy 2009; Pournajafian 2015; Ranucci 2015; Sabate 2016), ii) were too young to appreciate the nature of their intervention (Shirahata 1990; Turner 1981), iii) were unaware of the intervention content (Soleimani 2016), or iv) the plastic container vials used were covered and details of the contents not visible (Sadeghi 2014). We considered two trials, Najafi 2014 and Rasche 1982 to be at unclear risk of performance bias as insufficient information was provided to permit an alternative judgement. We did not consider any trials to be at high risk of participant performance bias.

We considered nine trials to be at low risk of **personnel** performance bias as: i) staff independent from the trial team prepared the interventions (Fenger Erikson 2009; Korte 2009; Sabate 2016; Soleimani 2016); ii) interventions were provided in visually identical containers (Levy 2009); iii) the plastic container vials used were covered and details of the contents were not visible (Sadeghi 2014); iv) clinical staff were blinded to treatment allocation (Karlsson 2011; Ranucci 2015); and v) outcome was not likely to be influenced by lack of blinding (Shirahata 1990). We considered six trials to be at high risk of personnel performance bias due to: i) the availability of the allocation concealment device in the participant's notes (Godje 2006); ii) the apparent non-concealment

of the notably visual difference between the intervention and control (Turner 1981); iii) personnel being unblinded to treatment allocation (Haas 2015 (C); Haas 2015 (S)); and iv) a lack of information on blinding where knowledge of treatment allocation could introduce performance bias (Cui 2010; Karkouti 2013). We considered three trials to be at unclear risk of personnel performance bias, as insufficient information was provided to permit an alternative judgement (Najafi 2014; Pournajafian 2015; Rasche 1982).

Therapeutic trials

We considered all 13 trials to be at low risk of **participant** performance bias because: i) intervention and placebo were identical in composition (Bregenzer 1999; Collins 2017; Curry 2018; Nascimento 2016; Wikkelsø 2015); ii) participants were under anaesthesia when their intervention was administered with the trial personnel and clinicians blinded to treatment allocation (Bilecen 2017; Galas 2014; Jeppsson 2016; Lance 2012; Rahe-Meyer 2013; Rahe-Meyer 2016; Tanaka 2014); and iii) participants had a low level of consciousness or were unconscious at the time of the study intervention, although the clinicians administering the intervention were not blinded to its nature (Innerhofer 2017).

We considered nine trials to be at low risk of **personnel** performance bias because the placebo was identical in composition to the intervention (Bilecen 2017; Bregenzer 1999; Collins 2017; Curry 2018; Jeppsson 2016; Nascimento 2016; Rahe-Meyer 2013; Rahe-Meyer 2016; Wikkelsø 2015). We considered two trials to be at high risk of personnel performance bias as these were open-label trial designs (Innerhofer 2017; Tanaka 2014), and we considered two trials to be at an unclear risk of performance bias, as insufficient information was provided to permit an alternative judgement (Galas 2014; Lance 2012).

Blinding of outcome assessors (detection bias)

Prophylactic intervention

We considered six trials to be at low risk of detection bias, as outcome assessments were: i) made by trial personnel blinded to treatment allocation (Cui 2010; Fenger Erikson 2009; Karkouti 2013; Karlsson 2011); or ii) knowledge of treatment allocation would not affect assessment of objective outcomes (Rasche 1982; Shirahata 1990). We considered two trials to be at high risk of detection bias due to lack of appropriate allocation concealment (Godje 2006; Turner 1981), as well as notable functional and visual differences between the intervention and control in Turner 1981. We considered 10 trials to be at unclear risk of detection bias, as insufficient information was provided to permit an alternative judgement (Haas 2015 (C); Haas 2015 (S); Korte 2009; Levy 2009; Najafi 2014; Pournajafian 2015; Ranucci 2015; Sabate 2016; Sadeghi 2014; Soleimani 2016).

Therapeutic trials

We considered six trials to be at low risk of detection bias as outcomes were: i) measured by participants blinded to their treatment allocation and by an independent reference laboratory (Bregenzer 1999); ii) trial personnel were blinded to treatment allocation (Bilecen 2017; Galas 2014; Rahe-Meyer 2013; Wikkelsø 2015); and iii) knowledge of allocation would not influence assessment of the trial's objective outcomes (Lance 2012). We considered two trials to be at high risk of detection bias as these were open-label trial designs (Innerhofer 2017; Tanaka 2014). We

considered five studies to be at unclear risk of detection bias, as insufficient information was provided to permit a judgement of high or low risk of detection bias (Collins 2017; Curry 2018; Jeppsson 2016; Nascimento 2016; Rahe-Meyer 2016).

Incomplete outcome data

Prophylactic intervention

We considered 11 trials to be at low risk of attrition bias, with all participants accounted for within the study flow diagram and no withdrawals (Fenger Erikson 2009; Godje 2006; Haas 2015 (C); Haas 2015 (S); Karkouti 2013; Najafi 2014; Ranucci 2015; Sadeghi 2014; Turner 1981), or where participants were lost to follow-up this was evenly spread across trial arms (Sabate 2016; Soleimani 2016). We considered two trials to be at high risk of attrition bias because no details were reported about the intervention groups of participants excluded from analysis (Levy 2009) and four participants were excluded at unclear phases of the study, but following recruitment and may have been excluded following randomisation and treatment administration (Pournajafian 2015). We considered five trials to be at unclear risk of attrition bias because: i) the issue of participant dropout and post-randomisation exclusion was not addressed (Cui 2010; Karlsson 2011; Rasche 1982; Shirahata 1990); and ii) the number of participants randomised to each treatment arm or analysed by outcome was not reported (Korte 2009).

Therapeutic trials

We considered 11 trials to be at low risk of attrition bias, as all participants were accounted for within the study flow diagram and there were no withdrawals (Collins 2017; Curry 2018; Galas 2014; Innerhofer 2017; Jeppsson 2016; Lance 2012; Nascimento 2016; Rahe-Meyer 2013; Rahe-Meyer 2016; Tanaka 2014; Wikkelsø 2015). We considered one trial to be at high risk of attrition bias, because a third of all participants were not available for the intention-to-treat analyses (Bregenzer 1999). We considered one trial to be at unclear risk of attrition bias, because we determined that the missing data forms from participants in both study arms were unlikely to have an impact on outcome; as the data were balanced across study groups, this presented an unclear risk of bias (Bilecen 2017).

Selective reporting

Prophylactic trials

We considered 14 trials to be at low risk of reporting bias, because all the outcomes were prespecified by the authors and provided in the results (Cui 2010; Fenger Erikson 2009; Godje 2006; Haas 2015 (C); Haas 2015 (S); Karkouti 2013; Karlsson 2011; Korte 2009; Ranucci 2015; Rasche 1982; Sabate 2016; Sadeghi 2014; Soleimani 2016; Turner 1981). We considered two trials to be at high risk of reporting bias, as the trials did not report data for all predefined outcomes (Levy 2009; Najafi 2014). We considered two trials to be at unclear risk of reporting bias, as outcomes were not predefined (Pournajafian 2015; Shirahata 1990).

Therapeutic trials

We considered nine trials to be at low risk of reporting bias, because all the outcomes were prespecified by the authors and covered in the results (Bilecen 2017; Collins 2017; Galas 2014; Jeppsson 2016; Nascimento 2016; Rahe-Meyer 2013; Rahe-Meyer 2016; Tanaka 2014; Wikkelsø 2015).

We considered two trials to be at high risk of reporting bias (Bregenza 1999; Innerhofer 2017). In Bregenza 1999, due to the early discontinuation of the study, data were not reported for a number of prespecified outcomes, including haemoglobin and haematocrit levels. In Innerhofer 2017, the published manuscript did not report findings from all the outcomes they purported to measure. We considered two trials to be at unclear risk of reporting bias, as there was no trial protocol and no clarity about the outcomes measured in the trial (Lance 2012). In Curry 2018, the study protocol outlined the use of ROTEM data, but none were reported.

Other potential sources of bias

Prophylactic trials

We determined four studies to be at low risk of other potential sources of bias in study design; Fenger Erikson 2009 appeared free of all concerns, while Haas 2015 (C), Haas 2015 (S) and Karlsson 2011 declared industrial funding and support, but reported no conflicts of interest or funder control over data or publication. We considered one trial (Karkouti 2013) to be at high risk of bias due to variations in transfusion practice, despite declared protocols and the fact that the company producing the product and sponsoring the trial oversaw the trial operations, audited all data, and conducted the statistical analyses. There is also unclear ownership of trial data, although the publication states full author access.

We determined that 13 trials were at an unclear risk of other bias. In seven of these trials, the reasoning was: i) participants enrolled in Cui 2010 appear to have been randomised prior to application of exclusion criteria with more participants excluded from the intervention arm of the study, although the allocation and exclusion numbers were not stated; ii) Godje 2006 did not declare sources of funding or a statement of author interests; iii) Korte 2009 declared industry funding with an industrial employee on the byline, but did not provide any indication of roles or independence of publication or data ownership; iv) Levy 2009 declared industrial funding of the study and three authors, including the senior author, were industrial employees. However there was no statement of data ownership, study independence or industrial control of data, manuscript content or publication; v) Ranucci 2015 declared full author access to data and decision to publish, independent of industrial funding, but ownership of the data was unclear; vi) Sabate 2016 provided a power analysis identifying that 66 participants were required in each study group, hence insufficient participants (99) were recruited for the study's primary outcome conclusions; and vii) the source of funding for Turner 1981 was unclear. In the other six trials there was insufficient information from which to determine the risk of any other forms of bias (Najafi 2014; Pournajafian 2015; Rasche 1982; Sadeghi 2014; Shirahata 1990; Soleimani 2016).

Therapeutic trials

We deemed six therapeutic trials to be at low risk of other sources of bias. Five trials (Bilecen 2017; Galas 2014; Innerhofer 2017; Jeppsson 2016; Lance 2012) appeared free of all concerns, whilst the sixth (Collins 2017) declared industrial support for the study but provided detailed information about involvement and low industrial influence.

We considered two trials to be at high risk of bias, due to a declaration of funding by CS Behring and having industrial employees as authors without providing any information about data ownership or control, study, manuscript and publication independence (Rahe-Meyer 2013; Rahe-Meyer 2016). Five trials did not provide sufficient information for us to determine risk of other potential sources of bias (Bregenza 1999; Curry 2018; Nascimento 2016; Tanaka 2014; Wikkelsso 2015).

Effects of interventions

See: **Summary of findings for the main comparison FIBRINOGEN: Therapeutic trials: intervention compared to inactive Control for the prevention and treatment of bleeding in participants without haemophilia**; **Summary of findings 2 FIBRINOGEN: Therapeutic trials: intervention compared to haemostatically active control for the prevention and treatment of bleeding in participants without haemophilia**; **Summary of findings 3 FIBRINOGEN: Prophylactic trials: intervention compared to inactive control for the prevention and treatment of bleeding in participants without haemophilia**; **Summary of findings 4 FIBRINOGEN: Prophylactic trials: intervention compared to haemostatically active control for the prevention and treatment of bleeding in participants without haemophilia**; **Summary of findings 5 FXIII: Prophylactic trials: intervention compared to inactive control for the prevention and treatment of bleeding in participants without haemophilia**

We present the outcome data in the following way.

- by outcome.
- by type of intervention.
- by whether the intervention was used in a prophylactic or therapeutic setting.
- by whether the intervention was being compared with a haemostatically inactive comparator (for example, placebo or saline) or a haemostatically active comparator (whether a different dose of the same intervention or a different haemostatically active intervention).

However, the paucity of quantifiable data that can be combined in a meta-analysis is such that we cannot provide quantitative analysis for most comparisons.

We analysed the three trials enrolling neonates separately from the other included trials (as previously described) (Haas 2015 (C); Shirahata 1990; Turner 1981). Due to the small volumes of the placebo agents and their inactive nature in two trials (Shirahata 1990; Turner 1981), we have compared all trials in this group as if their comparators are equivalent.

Three trials assessed the prophylactic use of factor XIII against an inactive comparator with multiple factor XIII dosing arms (Godje 2006; Karkouti 2013; Levy 2009). While we appreciate it is not possible to generate an exact comparison across these trials, to facilitate some meaningful analysis we have included the data for the 2500 unit (Godje 2006) and 35 unit/kg arms (Karkouti 2013; Levy 2009), as these should correlate for a 'typical' 70 kg person. Each trial included only a single control arm and we have included them in full.

In Fenger Erikson 2009, the average participant weight was 77 kg, and we have used this value to calculate total intervention doses given.

In Korte 2009, the breakdown of the 22 randomised participants was not reported by intervention group, but based on graphical data in the report we have been able to identify that 11 participants were included in each study arm.

Primary outcome: mortality (all-cause)

Fibrinogen concentrate: prophylactic trials with inactive comparator

(Nine trials: Cui 2010; Fenger Erikson 2009; Karlsson 2011; Najafi 2014; Pournajafian 2015; Ranucci 2015; Sabate 2016; Sadeghi 2014; Soleimani 2016)

Four trials comparing fibrinogen concentrate versus inactive comparators with 248 participants reported data on all-cause mortality (Fenger Erikson 2009; Karlsson 2011; Ranucci 2015; Sabate 2016), but we did not pool data due to differences in the timing of the outcome measurement.

There was no evidence of a difference in the number of deaths between the fibrinogen and the inactive comparator arms: risk ratio (RR) 0.33, 95% confidence interval (CI) 0.04 to 3.11; in 116 participants up to 30 days post-complex cardiac surgery (Ranucci 2015) and RR 0.31, 95% CI 0.03 to 2.83; in 92 participants prior to hospital discharge after liver transplantation (Sabate 2016). Fenger Erikson 2009 and Karlsson 2011 reported no events at up to four days after radical cystectomy amongst 20 participants and up to seven days after coronary artery bypass graft surgery amongst 20 participants, respectively (Analysis 1.1).

Fibrinogen concentrate: prophylactic trials with active comparator

(Two participant groups reported in 1 trial: Haas 2015 (C); Haas 2015 (S))

No mortality was reported by either Haas 2015 (C) or Haas 2015 (S), with 49 participants in total when comparing two different thresholds for administering prophylactic fibrinogen concentrate in complex surgery (Analysis 2.1).

Fibrinogen concentrate: therapeutic trials with inactive comparator

(Eight trials: Bilecen 2017; Collins 2017; Curry 2018; Jeppsson 2016; Nascimento 2016; Rahe-Meyer 2013; Rahe-Meyer 2016; Wikkelsø 2015)

All-cause mortality data from 724 participants was reported by seven trials comparing fibrinogen concentrate with an inactive agent in the management of active bleeding (Bilecen 2017; Collins 2017; Curry 2018; Nascimento 2016; Rahe-Meyer 2013; Rahe-Meyer 2016; Wikkelsø 2015). Differences in the timing of the outcome measure and clinical setting of the trial prevented us from pooling data across the seven trials (Analysis 3.1).

Bilecen 2017 did not demonstrate any evidence of a difference in in-hospital mortality within 30 days of cardiac surgery between the fibrinogen and inactive comparator intervention arms: RR 5.00, 95% CI 0.25 to 102.00; 120 participants; low-quality evidence (Summary of findings for the main comparison).

Pooled data at 46 days following aortic valve surgery (Rahe-Meyer 2013) and complex cardiovascular surgery (Rahe-Meyer 2016) did not demonstrate any evidence of a difference in

the number of deaths between the fibrinogen and inactive comparator intervention arms: RR 0.23, 95% CI 0.05 to 1.01; 213 participants; high-quality evidence (Summary of findings for the main comparison).

Two trials investigating the role of fibrinogen concentrate in the management of trauma haemorrhage (Curry 2018; Nascimento 2016) did not demonstrate any evidence of a difference in the number of deaths between the fibrinogen and inactive comparator intervention arms at 28 days following admission: RR 1.46, 95% CI 0.71 to 2.99; 97 participants; moderate-quality evidence (Summary of findings for the main comparison).

In the management of postpartum haemorrhage no deaths were reported or recorded in two trials six weeks postnatally (Collins 2017; 50 participants and Wikkelsø 2015; 244 participants).

Fibrinogen concentrate: therapeutic trials with active comparator

(Four trials: Galas 2014; Innerhofer 2017; Lance 2012; Tanaka 2014)

Four studies comparing fibrinogen concentrate to haemostatically active comparators reported on all-cause mortality in 220 participants at different time points (Galas 2014; Innerhofer 2017; Lance 2012; Tanaka 2014). No studies reported comparable time points, hence we did not pool data in a meta-analysis across the trials (Analysis 4.1).

Following major elective surgery, there was no evidence of a difference in the number of deaths between fibrinogen and FFP at post-operative day 30 (Lance 2012: RR 0.95, 95% CI 0.06 to 14.30; 43 participants, low-quality evidence (Summary of findings 2)) or 30 days after major trauma (Innerhofer 2017: RR 2.20, 95% CI 0.45 to 10.78; 94 participants, very low-quality evidence (Summary of findings 2)).

Following cardiac surgery there were no recorded deaths in either the fibrinogen or cryoprecipitate arms at post-operative day seven or hospital discharge (Galas 2014: 63 participants) or when fibrinogen concentrate was compared to platelet infusion at 28 days post-operatively (Tanaka 2014: 20 participants).

Factor XIII (FXIII): prophylactic trials with inactive comparator

(Six trials: Godje 2006; Karkouti 2013; Korte 2009; Levy 2009; Rasche 1982; Shirahata 1990)

All-cause mortality was reported for 414 participants in five trials comparing prophylactic factor XIII with a haemostatically inactive control in adult populations (Godje 2006; Karkouti 2013; Korte 2009; Levy 2009; Rasche 1982) (Analysis 5.1).

Following coronary artery bypass graft surgery there was no evidence of a difference in the number of deaths between the factor XIII and placebo intervention arms at five to seven weeks post-operatively in Karkouti 2013: RR 2.78, 95% CI 0.11 to 67.73; 266 participants. Levy 2009 reported no events in 16 participants who had undergone coronary artery bypass graft surgery. Godje 2006 recorded no deaths prior to hospital discharge in 50 participants following cardiac surgery.

In non-cardiac scenarios there was no evidence of a difference in all-cause mortality between factor XIII and haemostatically inactive control groups at an unspecified time point in acute

leukaemia patients in [Rasche 1982](#): RR 1.07, 95% CI 0.49 to 2.32; 60 participants; or at a median follow-up of 340 days from elective gastrointestinal surgery in [Korte 2009](#): RR 1.00, 0.26 to 3.91; 22 participants.

Factor XIII (FXIII): therapeutic trials with inactive comparator

(one trial: [Bregenzler 1999](#))

[Bregenzler 1999](#) did not report on all-cause mortality.

Prothrombin complex concentrate (PCC): prophylactic trials with inactive comparator

(one trial: [Turner 1981](#))

In [Turner 1981](#) there was no evidence of a difference in the number of deaths amongst pre-term neonates (time point not stated) when PCC was compared to no treatment: RR 1.05, 95% CI 0.71 to 1.53; 78 participants ([Analysis 7.1](#)).

Primary outcome: arterial and venous thromboembolic events (VTE)

Fibrinogen concentrate: prophylactic trials with inactive comparator

(Nine trials: [Cui 2010](#); [Fenger Erikson 2009](#); [Karlsson 2011](#); [Najafi 2014](#); [Pournajafian 2015](#); [Ranucci 2015](#); [Sabate 2016](#); [Sadeghi 2014](#); [Soleimani 2016](#))

Seven trials reported the incidence of arterial events in 398 participants where fibrinogen concentrate was given prophylactically in comparison to an inactive control ([Fenger Erikson 2009](#); [Karlsson 2011](#); [Najafi 2014](#); [Ranucci 2015](#); [Sabate 2016](#); [Sadeghi 2014](#); [Soleimani 2016](#)) ([Analysis 1.2](#); [Analysis 1.3](#)). We did not combine data across all studies due to differences in the time point of measurement and the clinical settings of the trials.

Arterial events

In participants undergoing cardiac surgery, at four days post-operatively [Sadeghi 2014](#) reported no arterial events in either study arm while [Karlsson 2011](#) reported a single event in each arm (60 and 20 participants, respectively). At 30 days following cardiac surgery, there was no evidence of a difference in the number of arterial events in [Ranucci 2015](#): RR 0.33, 95% CI 0.01 to 8.02; 116 participants; low-quality evidence ([Summary of findings 3](#)).

In studies that assessed fibrinogen concentrate in elective surgery, no arterial events were reported up to three days following radical cystectomy in [Fenger Erikson 2009](#) (20 participants), up to one week after trans-urethral prostatic resection in [Soleimani 2016](#) (60 participants) or at 30 days following hip replacement surgery in [Najafi 2014](#) (30 participants). In [Sabate 2016](#), there was no evidence of a difference in arterial event incidence between the intervention arms 30 days after liver transplantation: RR 0.31, 95% CI 0.03 to 2.83; 92 participants, low-quality evidence ([Summary of findings 3](#)).

Venous events

Following cardiac surgery, there was no evidence of a difference in the incidence of venous thromboembolic complications at four days between the intervention arms in [Karlsson 2011](#): RR 3.00, 95% CI 0.14 to 65.90; 20 participants, moderate-quality evidence ([Summary of findings 3](#)).

No venous events were reported at four days following cardiac surgery in 60 participants in [Sadeghi 2014](#) or amongst 116 participants at 30 days following cardiac surgery in [Ranucci 2015](#). Following elective surgery [Fenger Erikson 2009](#) (20 participants), [Soleimani 2016](#) (60 participants) and [Najafi 2014](#) (30 participants) did not report any venous events at three, seven and 30 days, respectively.

In [Sabate 2016](#) there was no evidence of a difference in the incidence of venous thromboembolic complications between the intervention arms 30 days following liver transplantation: RR 0.31, 95% CI 0.01 to 7.32; 92 participants, low-quality evidence ([Summary of findings 3](#)).

Fibrinogen concentrate: prophylactic trials with active comparator

(Two trials reported in 1 study: [Haas 2015](#) (C); [Haas 2015](#) (S))

[Haas 2015](#) (C) and [Haas 2015](#) (S) reported no arterial or venous thromboembolic events in either fibrinogen concentrate supplementation arms (total 49 participants) during the entire participant hospital stay.

Fibrinogen concentrate: therapeutic trials with inactive comparator

(Eight trials: [Bilecen 2017](#); [Collins 2017](#); [Curry 2018](#); [Jeppsson 2016](#); [Nascimento 2016](#); [Rahe-Meyer 2013](#); [Rahe-Meyer 2016](#); [Wikkelso 2015](#))

Six trials reported both arterial and venous thromboembolic events at a single time point each in a total of 562 participants ([Bilecen 2017](#); [Collins 2017](#); [Curry 2018](#); [Jeppsson 2016](#); [Rahe-Meyer 2013](#); [Wikkelso 2015](#)). Additionally, adverse event data from [Nascimento 2016](#) (45 participants) could be used to calculate the per capita event rate for arterial events but not venous events ([Analysis 3.2](#); [Analysis 3.3](#)).

Arterial events

Neither [Collins 2017](#) nor [Wikkelso 2015](#) found any arterial events at six weeks following postpartum haemorrhage in 50 and 244 participants, respectively. There were no reported arterial events in [Nascimento 2016](#) at 28 days following admission for trauma (45 participants). [Curry 2018](#) found no evidence of a difference between the fibrinogen and control arms in the arterial events rate at 28 days following admission for trauma: RR 0.47, 95% CI 0.05 to 4.82; 39 participants.

In the cardiac surgery studies, there was no evidence of any differences in the number of arterial events at post-operative day 30 ([Bilecen 2017](#); RR 2.33, 95% CI 0.63 to 8.60; 120 participants), day 45 ([Rahe-Meyer 2013](#): RR 1.10, 95% CI 0.07 to 16.85; 61 participants) and during hospital stay ([Jeppsson 2016](#): RR 0.33, 95% CI 0.01 to 7.80; 48 participants).

Venous events

There were no recorded venous thromboembolic events following cardiac surgery in [Bilecen 2017](#) (120 participants) or [Jeppsson 2016](#) (48 participants), or at six weeks following postpartum haemorrhage in [Wikkelso 2015](#) (244 participants). In the three trials recording venous thromboembolic events there was no evidence of a difference in the event rate between the fibrinogen and inactive comparator intervention arms: following cardiac surgery in [Rahe-](#)

Meyer 2013: RR 0.37, 95% CI 0.02 to 8.66; 61 participants; at six weeks following postpartum haemorrhage in Collins 2017: RR 0.92, 95% CI 0.06 to 13.95, 50 participants; and 28 days after an admission with major trauma in Curry 2018: RR 4.76, 95% CI 0.24 to 93.19; 39 participants.

Fibrinogen concentrate: therapeutic trials with active comparator

(Four trials: Galas 2014; Innerhofer 2017; Lance 2012; Tanaka 2014)

Three studies reported on both arterial and venous complications in 126 participants receiving therapeutic fibrinogen concentrate or active comparator (Galas 2014; Lance 2012; Tanaka 2014). Additionally Innerhofer 2017 reported on the incidence of venous thromboembolic complications in a further 94 participants. We did not pool data collectively due to differences in the timing of the outcome measurement (Analysis 4.2; Analysis 4.3), the active comparator and the clinical setting of the trials.

Arterial events

There was no evidence of a difference in arterial event incidence between the fibrinogen and active comparator arms in Galas 2014 seven days after cardiac surgery: RR 0.44, 95% CI 0.09 to 2.10; 63 participants; or in Tanaka 2014 28 days after cardiac surgery: RR 0.33, 95% CI 0.02 to 7.32; 20 participants; or in Lance 2012 30 days after major elective surgery: RR 2.87, 95% CI 0.12 to 66.75; 43 participants.

Venous events

No venous events were reported in either cardiac surgery trial (Galas 2014; Tanaka 2014).

There was no evidence of a difference in venous thromboembolic event incidence between the fibrinogen and active comparator arm in Innerhofer 2017 30 days after trauma admission: RR 0.63, 95% CI 0.21 to 1.87; 94 participants; or following major surgery in Lance 2012: RR 3.00, 95% CI 0.12 to 77.83; 43 participants.

Factor XIII (FXIII): prophylactic trials with inactive comparator

(six trials: Godje 2006; Karkouti 2013; Korte 2009; Levy 2009; Rasche 1982; Shirahata 1990)

Four trials of factor XIII prophylaxis reported on thromboembolic complications in 338 participants in cardiac surgery (Godje 2006; Karkouti 2013; Levy 2009) and surgery for gastrointestinal cancer (Korte 2009) (Analysis 5.2; Analysis 5.3)

Arterial events

No arterial events were reported at up to seven weeks following cardiac surgery in Levy 2009 (16 participants).

There was no evidence of a difference in arterial thromboembolic events in Karkouti 2013 up to seven weeks following cardiac surgery: RR 0.67, 95% CI 0.28 to 1.62; 266 participants; low-quality evidence; or in Godje 2006 at an unspecified post-operative time point: RR 0.20, 95% CI 0.01 to 3.97; 50 participants; very low-quality evidence; or Korte 2009 following gastrointestinal surgery: RR 3.00, 95% CI 0.14 to 66.53; 22 participants; low-quality evidence.

Venous events

No venous events were reported at up to seven weeks following cardiac surgery in Levy 2009 (16 participants) or Godje 2006 (50 participants).

There was no evidence of a difference in venous thromboembolic events in Karkouti 2013 up to seven weeks following cardiac surgery: RR 0.93, 95% CI 0.06 to 14.67; 266 participants; low-quality evidence; or Korte 2009 following gastrointestinal surgery: RR 3.00, 95% CI 0.14 to 66.53; 22 participants; low-quality evidence.

Factor XIII (FXIII): therapeutic trials with inactive comparator

(one trial: Bregenzner 1999)

Bregenzner 1999 did not report the incidence of arterial or venous thromboembolic complications.

Prothrombin complex concentrate (PCC): prophylactic trials with inactive comparator

(One trial; Turner 1981)

Turner 1981 did not report the incidence of arterial or venous thromboembolic complications.

Secondary outcome: mortality due to bleeding

Fibrinogen concentrate: prophylactic trials with inactive comparator

(Nine trials: Cui 2010; Fenger Erikson 2009; Karlsson 2011; Najafi 2014; Pournajafian 2015; Ranucci 2015; Sabate 2016; Sadeghi 2014; Soleimani 2016)

Fenger Erikson 2009, Karlsson 2011; Ranucci 2015; and Sabate 2016 reported on this outcome in 248 participants. There was only a single death due to bleeding reported in the fibrinogen arm of Sabate 2016: RR 2.76, 95% CI 0.12 to 65.92; 92 participants (Analysis 1.4).

Fibrinogen concentrate: prophylactic trials with active comparator

(Two participant groups reported in one trial: Haas 2015 (C); Haas 2015 (S))

No deaths due to bleeding were reported in Haas 2015 (C) or Haas 2015 (S) from 49 participants in either fibrinogen supplementation arm (Analysis 2.4).

Fibrinogen concentrate: therapeutic trials with inactive comparator

(Eight trials: Bilecen 2017; Collins 2017; Curry 2018; Jeppsson 2016; Nascimento 2016; Rahe-Meyer 2013; Rahe-Meyer 2016; Wikkelso 2015)

Five studies reported data on death due to haemorrhage with a total of 539 participants. Rahe-Meyer 2013 reported no deaths due to bleeding occurred following cardiac surgery in 152 participants up to post-operative day 46 (Analysis 3.4).

Following postpartum haemorrhage, no deaths due to bleeding were reported at six weeks postnatally in either Collins 2017 or Wikkelso 2015 in 50 and 244 participants, respectively.

Mortality due to bleeding following administration of therapeutic fibrinogen at 28 days following trauma admission was reported and pooled for [Curry 2018](#) and [Nascimento 2016](#): there was no evidence of any difference in the number of deaths between the intervention arms: RR 2.45, 95% CI 0.38 to 15.76; 93 participants.

Fibrinogen concentrate: therapeutic trials with active comparator

(Four trials: [Galas 2014](#); [Innerhofer 2017](#); [Lance 2012](#); [Tanaka 2014](#)).

Four trials with 220 participants reported no incidents of mortality due to bleeding to post-cardiac surgery day seven ([Galas 2014](#): 63 participants) and day 28 ([Tanaka 2014](#): 20 participants; [Lance 2012](#): 43 participants), or at 30 days following trauma ([Innerhofer 2017](#): 94 participants) ([Analysis 4.4](#)).

Factor XIII (FXIII): prophylactic trials with inactive comparator

(Six trials: [Godje 2006](#); [Karkouti 2013](#); [Korte 2009](#); [Levy 2009](#); [Rasche 1982](#); [Shirahata 1990](#))

Four trials with 392 participants reported data for this outcome ([Godje 2006](#); [Karkouti 2013](#); [Levy 2009](#); [Rasche 1982](#)) ([Analysis 5.4](#)).

[Rasche 1982](#) reported mortality due to bleeding at a median of 32 days following the study intervention: there was no evidence of a difference in mortality due to bleeding in people with acute leukaemia given factor XIII infusion compared to no intervention: RR 0.80, 95% CI 0.20 to 3.28; 60 participants.

[Godje 2006](#); [Karkouti 2013](#); [Levy 2009](#) reported on this outcome following cardiac surgery and reported that no deaths due to bleeding occurred (total 332 participants).

Factor XIII (FXIII): therapeutic trials with inactive comparator

(One trial: [Bregenzner 1999](#))

[Bregenzner 1999](#) did not report on this outcome.

Prothrombin complex concentrate (PCC): prophylactic trials with inactive comparator

(One trial: [Turner 1981](#)).

[Turner 1981](#) did not report on this outcome.

Secondary outcome: red blood cell (RBC) transfusion requirement

See [Table 1](#) for individual study data for this outcome, which was reported in a number of ways (means, median, total) across the included studies.

Fibrinogen concentrate: prophylactic trials with inactive comparator

(Nine trials: [Cui 2010](#); [Fenger Erikson 2009](#); [Karlsson 2011](#); [Najafi 2014](#); [Pournajafian 2015](#); [Ranucci 2015](#); [Sabate 2016](#); [Sadeghi 2014](#); [Soleimani 2016](#))

Mean red blood cell volume required per participant

[Karlsson 2011](#), [Najafi 2014](#) and [Sadeghi 2014](#) (110 participants in total) reported these data as mean red blood cell volume required per participant, but pooling of data was not possible due to the lack of common time points. Following cardiac surgery, fibrinogen

concentrate reduced the red blood cell transfusion requirement at 12 hours post-operatively ([Karlsson 2011](#); mean difference (MD) -150.00, 95% CI -205.72 to -94.28 mL; 20 participants; although this effect was not reproduced intra-operatively (MD -90.00, 95% CI -281.13 to 101.13 mL; 60 participants). There was no evidence of a difference in red blood cell requirement between the intervention arms at 24 hours post-cardiac surgery in [Sadeghi 2014](#) (MD -150.00, 95% CI -401.53 to 101.53 mL; 60 participants) or in the perioperative period following hip replacement surgery ([Najafi 2014](#); MD -78.00, 95% CI -317.29 to 161.29 mL; 30 participants) ([Analysis 1.5](#)).

Receipt of a red blood cell transfusion

Nine studies (with 470 participants) reported the number of participants receiving a red cell transfusion ([Cui 2010](#); [Fenger Erikson 2009](#); [Karlsson 2011](#); [Najafi 2014](#); [Pournajafian 2015](#); [Ranucci 2015](#); [Sabate 2016](#); [Sadeghi 2014](#); [Soleimani 2016](#)).

Following cardiac surgery three studies provided a common time point at 24 hours post-operatively and we pooled the data ([Cui 2010](#); [Ranucci 2015](#); [Sadeghi 2014](#)). The data showed a reduction in the number of participants receiving a RBC transfusion with the use of prophylactic fibrinogen in cardiac surgery with moderate heterogeneity: RR 0.53, 95% CI 0.37 to 0.76; $P < 0.001$; 207 participants; $I^2 = 31\%$ ([Analysis 1.6](#)). This association was also present when we used a random-effects model. We noted no change to the effect estimate, but an increase in heterogeneity (analysis not shown) when we restricted the analysis to:

- trials including adult participants only: RR 0.53, 95% CI 0.36 to 0.78, $I^2 = 65\%$; 176 participants) ([Ranucci 2015](#); [Sadeghi 2014](#));
- trials where the number of participants who were not included in the analysis was unclear: RR 0.53, 95% CI 0.36 to 0.78, $I^2 = 65\%$; 176 participants) ([Ranucci 2015](#); [Sadeghi 2014](#)).

We noted a change in the effect estimate in that there was no difference in blood loss between the intervention arms and a reduction in heterogeneity when we restricted the analysis to trials with a low or uncertain risk of bias: RR 0.67, 95% CI 0.44 to 1.01, $I^2 = 0\%$; 91 participants) ([Cui 2010](#); [Sadeghi 2014](#)).

We also found evidence for a reduction in the number of participants receiving a transfusion between the fibrinogen and inactive comparator arm at 48 hours post-operatively in [Ranucci 2015](#): RR 0.59, 95% CI 0.38 to 0.92; 116 participants. Conversely, [Karlsson 2011](#) did not find any evidence for a reduction in the number of participants requiring a red blood cell transfusion at 12 hours following coronary artery bypass grafting: RR 0.33, 95% CI 0.04 to 2.69; 20 participants.

We pooled data from the elective, non-cardiac surgery trials ([Fenger Erikson 2009](#); [Soleimani 2016](#)), and found evidence of a reduction in the number of participants requiring a red blood cell transfusion following prophylactic fibrinogen at 48 hours post-operatively: RR 0.23, 95% CI 0.07 to 0.68; $P = 0.008$; 80 participants; $I^2 = 0\%$; [Analysis 1.6](#)). In the trial of participants undergoing spinal fusion surgery ([Pournajafian 2015](#)) and liver transplantation ([Sabate 2016](#)) there was no evidence of a difference in the number of participants requiring a RBC transfusion between the intervention arms at 24 hours following surgery: RR 1.01, 95% CI 0.76 to 1.33; 92 participants; and RR 0.07, 95% CI 0.00 to 1.22; 41 participants, respectively).

Four studies reported data at a range of different time points, as medians with IQRs (Cui 2010; Fenger Erikson 2009; Ranucci 2015; Sabate 2016). See Table 1 for these values.

Fibrinogen concentrate: prophylactic trials with active comparator

(Two participant groups reported in one trial: Haas 2015 (C); Haas 2015 (S))

Haas 2015 (C) and Haas 2015 (S) reported the mean and median quantity of red cells transfused (in 49 participants). This trial compared receipt of fibrinogen concentrate at two predefined intraoperative fibrinogen concentrations: early substitution and conventional substitution. In both surgical settings, participants in the conventional substitution arm received a greater quantity of red cells within the first 24 hours following surgery than participants in the early substitution arm (see Table 1), although there was no evidence of a statistically significant difference between the intervention arms in either trial (Haas 2015 (C): MD -8.70 (95% CI -21.71 to 4.31) mL/kg; 30 participants; and Haas 2015(S): MD -3.50, 95% CI -9.99 to 2.99 mL/kg; 19 participants; (Analysis 2.5).

All 30 participants in the craniosynostosis trial (Haas 2015 (C)) received a RBC transfusion within 24 hours of surgery. There was no evidence of a difference in the number of participants requiring a transfusion at 24 hours following surgery in the scoliosis trial (Haas 2015(S)): RR 0.60, 95% CI 0.25 to 1.46; 19 participants).

Fibrinogen concentrate: therapeutic trials with inactive comparator

(Eight trials: Bilecen 2017; Collins 2017; Curry 2018; Jeppsson 2016; Nascimento 2016; Rahe-Meyer 2013; Rahe-Meyer 2016; Wikkelso 2015)

Six trials including 568 participants reported data for RBC transfusion requirement (Bilecen 2017; Collins 2017; Curry 2018; Jeppsson 2016; Rahe-Meyer 2013; Wikkelso 2015). All six trials reported the number of participants who received a RBC transfusion, although differences in the measurement time points and the clinical setting of the trials prevented us from pooling the data (Analysis 3.5).

Bilecen 2017 and Rahe-Meyer 2013 reported the incidence of receipt of a RBC transfusion within 24 hours of delivery of the study drug in the management of bleeding during cardiac surgery. There was evidence of a reduction in receipt of a RBC transfusion for the participants who received fibrinogen in comparison with the placebo group: RR 0.54, 95% CI 0.39 to 0.75; $P < 0.001$; 178 participants; $I^2 = 0\%$; Analysis 3.5). The investigators also reported the median number of red blood cell units transfused (within 24 hours of delivery of the study drug), which was less in the fibrinogen group than in the placebo group (see Table 1). This finding was not repeated at the later time point reported by Jeppsson 2016, who found no evidence of a difference in the number of participants requiring a RBC transfusion during their entire hospital stay: RR 1.00, 95% CI 0.38 to 2.66; 48 participants.

Wikkelso 2015 found no evidence of a difference in the receipt of a RBC transfusion in 244 women with a postpartum haemorrhage who received fibrinogen or placebo (saline) within four hours (RR 0.39, 95% CI 0.13 to 1.22), within 24 hours (RR 0.72, 95% CI 0.38 to 1.38), and within seven days (RR 0.95, 95% CI 0.58 to 1.54) of receipt

of the study drug. The investigators also reported the number of women who received a transfusion of four units or more (time point unspecified), with no evidence of a difference between the fibrinogen and the placebo (saline) group: RR 2.74, 95% CI 0.71 to 10.57; 244 participants. Similarly Collins 2017 found no evidence of a difference in the number of participants receiving a red blood cell transfusion between the fibrinogen concentrate and placebo arms before hospital discharge for postpartum haemorrhage: RR 0.96, 95% CI 0.55 to 1.69; 55 participants.

Curry 2018 found no difference in the incidence of RBC transfusion following admission for trauma at three hours (RR 1.05, 95% CI 0.86 to 1.29; 46 participants), at six hours (RR 1.05, 95% CI 0.84 to 1.30; 43 participants), or 24 hours (RR 1.11, 95% CI 0.91 to 1.36; 41 participants).

Only Jeppsson 2016 reported the volume of red blood cells transfused; during the hospital stay following cardiac surgery there was no evidence of a difference between participants in whom bleeding was managed with fibrinogen concentrate and those who received placebo: MD -210.00, 95% CI -608.81 to 188.81 mL; 48 participants (Analysis 3.6).

Median values are reported in Table 1. These data were presented in seven studies (Bilecen 2017; Collins 2017; Curry 2018; Jeppsson 2016; Nascimento 2016; Rahe-Meyer 2013; Rahe-Meyer 2016).

Fibrinogen concentrate: therapeutic trials with active comparator

(Four trials: Galas 2014; Innerhofer 2017; Lance 2012; Tanaka 2014)

All four trials (total of 220 participants) reported data for RBC transfusion requirement. Lance 2012 reported the volume of RBC transfusion required following major elective surgery when fibrinogen concentrate was used as a partial replacement for FFP in the management of bleeding, but found no evidence of a difference between the intervention arms (MD -120.00, 95% CI -546.93 to 306.93 mL; 43 participants (Analysis 4.5).

Three trials (total of 177 participants) reported the number of participants receiving a RBC transfusion (Galas 2014; Innerhofer 2017; Tanaka 2014), but due to differences in the haemostatically active control used and the time points reported we did not undertake any meta-analyses. Following cardiac surgery Tanaka 2014 reported no evidence of a difference in red cell exposure by 24 hours post-operatively between the study's fibrinogen and platelet arms: RR 1.00, 95% CI 0.75 to 1.34; 20 participants. Innerhofer 2017 reported no evidence of a difference between fibrinogen and FFP for red cell exposure 24 hours after a trauma admission: RR 1.04, 95% CI 0.90 to 1.21; 94 participants, and Galas 2014 found no difference at post-operative day seven between fibrinogen and cryoprecipitate: RR 0.86, 95% CI 0.72 to 1.02; 63 participants (Analysis 4.6).

Median values are reported in Table 1 for Innerhofer 2017.

Factor XIII (FXIII): prophylactic trials with inactive comparator

(Six trials: Godje 2006; Karkouti 2013; Korte 2009; Levy 2009; Rasche 1982; Shirahata 1990)

Five trials (472 participants) reported data for RBC transfusion requirement (Godje 2006; Karkouti 2013; Korte 2009; Levy 2009; Rasche 1982).

One study (Godje 2006) reported the volume of red blood cells transfused and found no evidence of a difference between the intervention arms: -MD 0.70, 95% CI -1.63 to 0.23; 50 participants; Analysis 5.5). Karkouti 2013 reported data for the number of participants who received a RBC transfusion following elective surgery; there was no evidence of a difference between factor XIII and placebo: RR 0.97, 95% CI 0.70 to 1.35; 266 participants (Analysis 5.6). Levy 2009 stated that this outcome was measured in their trial, but did not report their findings.

Data were reported as medians with IQRs only for Karkouti 2013, Korte 2009 and Rasche 1982. In Korte 2009, the surgical participants in the haemostatically inactive control (placebo) arm received more RBC units than participants in the haemostatically active arm. In Rasche 1982, people with acute leukaemia in the haemostatically active arm required more RBC units than those in the haemostatically inactive control (placebo) arm. Median values are reported in Table 1.

Factor XIII (FXIII): therapeutic trials with inactive comparator

(One trial: Bregenzler 1999)

Bregenzler 1999 did not report on this outcome.

Prothrombin complex concentrate (PCC): prophylactic trials with inactive comparator

(One trial: Turner 1981)

Turner 1981 did not report on this outcome.

Blood loss

See Table 2 for individual study data for this outcome, which was reported in a number of ways (means, median, total) across the included studies.

Fibrinogen concentrate: prophylactic trials with inactive comparator

(Nine trials: Cui 2010; Fenger Erikson 2009; Karlsson 2011; Najafi 2014; Pournajafian 2015; Ranucci 2015; Sabate 2016; Sadeghi 2014; Soleimani 2016)

Post-operative bleeding was reported by eight trials with 378 participants (Cui 2010; Fenger Erikson 2009; Karlsson 2011; Najafi 2014; Pournajafian 2015; Ranucci 2015; Sadeghi 2014; Soleimani 2016) (Analysis 1.7).

In three trials, there was evidence of a difference in blood loss between the treatment arms with more blood being lost in the inactive comparator group than in the group that received fibrinogen concentrate (Analysis 1.7). We pooled data for the two cardiac surgery trials which reported blood loss at 12 hours post-operatively (Karlsson 2011; Sadeghi 2014): Ratio of Geometric Means 0.68, 95% CI 0.60 to 0.76; $P < 0.001$; $I^2 = 0\%$; 180 participants. Pournajafian 2015 reported evidence of a 21% reduction in bleeding at 24 hours following spinal fusion surgery for prophylactic fibrinogen concentration in comparison to an inactive comparator: Ratio of Geometric Means 0.77, 95% CI 0.66 to 0.89; $P < 0.001$; 41 participants.

There was no evidence of a difference between the intervention arms:

- in intra-operative (Ratio of Geometric Means 0.90, 95% CI 0.64 to 1.27; 30 participants) or 24-hour blood loss during hip surgery (Ratio of Geometric Means 0.99, 95% CI 0.80 to 1.24; 30 participants) in Najafi 2014;
- in intra-operative (Ratio of Geometric Means 1.02, 95% CI 0.75 to 1.37; 60 participants); or 48-hour blood loss following prostatic resection (Ratio of Geometric Means 0.85, 95% CI 0.57 to 1.27; 60 participants) in Soleimani 2016;
- in blood loss at one hour following cardiac surgery (Ratio of Geometric Means 0.75, 95% CI 0.52 to 1.09; 31 participants) in Cui 2010;
- in blood loss during and up to 48 hours after radical cystectomy (Ratio of Geometric Means 0.94, 95% CI 0.67 to 1.33; 20 participants) in Fenger Erikson 2009.

Ranucci 2015 reported the median quantity of blood lost from chest drain in the first 12 hours following elective cardiac surgery in 116 participants: there was less blood loss in participants receiving fibrinogen compared to the placebo control (saline); see Table 2.

Fibrinogen concentrate: prophylactic trials with active comparator

(Two participant groups reported in one trial; Haas 2015 (C); Haas 2015 (S))

Haas 2015 (C) and Haas 2015 (S) reported the median calculated total blood loss at 24 hours following surgery. This trial compared receipt of fibrinogen concentrate at two predefined intraoperative fibrinogen concentrations: early substitution and conventional substitution. In both surgical settings, participants in the conventional substitution arm had a greater total blood loss 24 hours post-operatively than those in the early substitution arm. Median values are reported in Table 2.

Fibrinogen concentrate: therapeutic trials with inactive comparator

(Eight trials: Bilecen 2017; Collins 2017; Curry 2018; Jeppsson 2016; Nascimento 2016; Rahe-Meyer 2013; Rahe-Meyer 2016; Wikkelsø 2015)

Five trials with 614 participants reported data for blood loss post-intervention.

The only trial that reported mean blood loss following use of fibrinogen concentrate as a treatment for bleeding (Jeppsson 2016) found no evidence of a difference in bleeding volume during surgery (Ratio of Geometric Mean 0.87, 95% CI 0.68 to 1.11; 48 participants) or at 12 hours post-operatively (Ratio of Geometric Mean 0.86, 95% CI 0.67 to 1.11; 48 participants), although blood loss was lower in the fibrinogen arm at all time points (Analysis 3.7).

Median blood loss was reported in five trials (Bilecen 2017; Collins 2017; Jeppsson 2016; Rahe-Meyer 2016; Wikkelsø 2015). Bilecen 2017 provided median blood loss data at seven time points following cardiac surgery up to 24 hours post-operatively. Participants who received fibrinogen demonstrated lower bleeding volumes at all time points. Collins 2017 and Wikkelsø 2015 provided median haemorrhage volumes following postpartum haemorrhage. In the fibrinogen intervention arm, Collins 2017 found there was less blood loss at 24 hours following study intervention, while in Wikkelsø 2015 there was no difference at up to six weeks postpartum. Jeppsson 2016 and Rahe-Meyer 2016

reported median blood loss after cardiac surgery. [Jeppsson 2016](#) found comparable bleeding volumes during surgery with lower volumes at 12 hours post-operatively in the fibrinogen arm. [Rahe-Meyer 2016](#) demonstrated lower blood loss at 6, 12 and 24 hours post-operatively in the active intervention arm.

Median values are reported in [Table 2](#).

Fibrinogen concentrate: therapeutic trials with active comparator

(Four trials: [Galas 2014](#); [Innerhofer 2017](#); [Lance 2012](#); [Tanaka 2014](#))

Two trials with 80 participants reported median data for blood loss post-intervention ([Galas 2014](#); [Tanaka 2014](#)). [Tanaka 2014](#) reported median blood loss as chest drainage to 12 hours post-cardiac surgery, and [Galas 2014](#) reported median blood loss at 48 hours following cardiac surgery. In both trials those participants who received the fibrinogen concentrate lost less blood than participants who received the respective haemostatically active control: cryoprecipitate in [Galas 2014](#), and platelets in [Tanaka 2014](#). Median values are reported in [Table 2](#).

Factor XIII (FXIII): prophylactic trials with inactive comparator

(Six trials: [Godje 2006](#); [Karkouti 2013](#); [Korte 2009](#); [Levy 2009](#); [Rasche 1982](#); [Shirahata 1990](#))

Only two trials (total of 66 participants) reported mean blood loss following prophylactic administration of factor XIII, both in cardiac surgery ([Godje 2006](#); [Levy 2009](#)). Blood loss was reported at a range of time points; at all time points blood loss was lower in the haemostatically active arm. We pooled data for two common time points, eight and 24 hours post-operatively. There was evidence of reduced blood loss in participants who received factor XIII when compared to an inactive comparator at eight hours (Ratio of Geometric Mean 0.79, 95% CI 0.65 to 0.97; $P = 0.02$; $I^2 = 0\%$; 66 participants) but this did not persist at 24 hours post-operatively (Ratio of Geometric Mean 0.85, 95% CI 0.72 to 1.00; 66 participants) ([Analysis 5.7](#)).

[Godje 2006](#) reported data at three further time points, with evidence of a reduction in blood loss for participants who had received factor XIII, in comparison to placebo at 12 hours (Ratio of Geometric Mean 0.78, 95% CI 0.63 to 0.97; $P = 0.02$; 50 participants), at 36 hours (Ratio of Geometric Mean 0.82, 95% CI 0.68 to 0.98; $P = 0.03$; 50 participants), and at 48 hours (Ratio of Geometric Mean 0.73, 95% CI 0.62 to 0.86; $P < 0.001$; 50 participants) post-operatively. [Levy 2009](#) reported blood loss data at the time of drain removal and observed that there was no evidence of a difference in blood loss between the intervention arms at those time points (Ratio of Geometric Mean 0.68, 95% CI 0.32 to 1.47; 16 participants).

[Korte 2009](#) reported median blood loss at the completion of surgery for gastrointestinal cancer and noted greater blood loss in the Factor XIII arm when compared to placebo, see [Table 2](#) for median values.

Factor XIII (FXIII): therapeutic trials with inactive comparator

(One trial: [Bregenzner 1999](#))

[Bregenzner 1999](#) did not report on blood loss post-intervention.

Prothrombin complex concentrate (PCC): prophylactic trials with inactive comparator

(One trial: [Turner 1981](#))

[Turner 1981](#) did not report on blood loss post-intervention.

Secondary outcome: allergic adverse reactions

Fibrinogen concentrate: prophylactic trials with inactive comparator

(Nine trials: [Cui 2010](#); [Fenger Erikson 2009](#); [Karlsson 2011](#); [Najafi 2014](#); [Pournajafian 2015](#); [Ranucci 2015](#); [Sabate 2016](#); [Sadeghi 2014](#); [Soleimani 2016](#))

One cardiac surgery trial ([Ranucci 2015](#)), and two non-cardiac surgery trials ([Fenger Erikson 2009](#); [Soleimani 2016](#)), with 196 participants in total, provided data on the incidence of allergic or hypersensitivity events following fibrinogen concentrate administration. No adverse events were reported at any time point.

Fibrinogen concentrate: prophylactic trials with active comparator

(Two participant groups reported in one trial: [Haas 2015](#) (C); [Haas 2015](#) (S))

No allergic adverse events to fibrinogen concentrate were reported in either arm of [Haas 2015](#) (C) or [Haas 2015](#) (S) in a total of 49 participants up to 24 hours post-operatively.

Fibrinogen concentrate: therapeutic trials with inactive comparator

(Eight trials: [Bilecen 2017](#); [Collins 2017](#); [Curry 2018](#); [Jeppsson 2016](#); [Nascimento 2016](#); [Rahe-Meyer 2013](#); [Rahe-Meyer 2016](#); [Wikkelsø 2015](#))

Four trials reported data on hypersensitivity reactions to fibrinogen concentrate or placebo in 470 participants. A single allergic event was recorded within 24 hours of placebo administration in the setting of postpartum haemorrhage ([Wikkelsø 2015](#): RR 0.33 (95% CI 0.01 to 8.06); 244 participants). No allergic reactions were reported at any time point following cardiac surgery ([Bilecen 2017](#); [Rahe-Meyer 2013](#)), or trauma ([Nascimento 2016](#)).

Fibrinogen concentrate: therapeutic trials with active comparator

(Four trials: [Galas 2014](#); [Innerhofer 2017](#); [Lance 2012](#); [Tanaka 2014](#))

Two trials with 106 participants reported data on allergic adverse reactions and found no hypersensitivity reactions to fibrinogen concentrate or the active comparators cryoprecipitate ([Galas 2014](#)), and FFP ([Lance 2012](#)).

Factor XIII (FXIII): prophylactic trials with inactive comparator

(Six trials: [Godje 2006](#); [Karkouti 2013](#); [Korte 2009](#); [Levy 2009](#); [Rasche 1982](#); [Shirahata 1990](#))

Five trials ([Godje 2006](#); [Korte 2009](#); [Levy 2009](#); [Rasche 1982](#); [Shirahata 1990](#)) reporting on 206 participants reported no incidents of allergy to factor XIII, although the time points were unclear in all of these studies.

Factor XIII (FXIII): therapeutic trials with inactive comparator

(One trial: [Bregenzler 1999](#))

[Bregenzler 1999](#) found no evidence of hypersensitivity to factor XIII in 28 participants.

Prothrombin complex concentrate (PCC): prophylactic trials with inactive comparator

(One trial: [Turner 1981](#))

[Turner 1981](#) reported no allergic reactions during administration of PCC prophylaxis in 78 participants.

Change to laboratory measures of coagulation, both standard laboratory tests and global measurements of coagulation

Due to the volume and heterogeneity of time points, definitions, reporting formats and tests performed, we did not undertake statistical analyses of these data. Instead we report the number of studies which reported on this outcome and the measures that they used to assess the outcome.

Fibrinogen concentrate: prophylactic trials with inactive comparator

(Nine trials: [Cui 2010](#); [Fenger Erikson 2009](#); [Karlsson 2011](#); [Najafi 2014](#); [Pournajafian 2015](#); [Ranucci 2015](#); [Sabate 2016](#); [Sadeghi 2014](#); [Soleimani 2016](#))

Eight trials (410 participants) reported measures of coagulation ([Cui 2010](#); [Fenger Erikson 2009](#); [Karlsson 2011](#); [Najafi 2014](#); [Pournajafian 2015](#); [Ranucci 2015](#); [Sabate 2016](#); [Sadeghi 2014](#)). All studies reported standard laboratory measures (most commonly Clauss fibrinogen levels) except [Cui 2010](#), where only TEG results were reported. Three studies reported viscoelastic tests and standard laboratory measures ([Fenger Erikson 2009](#); [Ranucci 2015](#); [Sabate 2016](#)), and only one study reported additional global testing (thrombin generation: [Fenger Erikson 2009](#)). All studies took samples at baseline and five reported a coagulation outcome at 24 hours ([Karlsson 2011](#); [Najafi 2014](#); [Pournajafian 2015](#); [Sabate 2016](#); [Sadeghi 2014](#)). There was marked heterogeneity across the studies in timing and frequency of testing.

Fibrinogen concentrate: prophylactic trials with active comparator

(Two participant groups reported in one trial: [Haas 2015](#) (C); [Haas 2015](#) (S))

These two trials analysed standard coagulation tests (activated partial thromboplastin time, international normalised ratio (INR), Clauss fibrinogen and factor XIII) and ROTEM at baseline alone ([Haas 2015](#) (C); [Haas 2015](#) (S)).

Fibrinogen concentrate: therapeutic trials with inactive comparator

(Eight trials: [Bilecen 2017](#); [Collins 2017](#); [Curry 2018](#); [Jeppsson 2016](#); [Nascimento 2016](#); [Rahe-Meyer 2013](#); [Rahe-Meyer 2016](#); [Wikkelso 2015](#))

All eight trials (with 759 participants) reported standard clotting tests, including Clauss fibrinogen, with all studies reporting results from baseline. Three studies ([Collins 2017](#); [Rahe-Meyer 2013](#); [Rahe-Meyer 2016](#)) reported additional ROTEM measures, with all three

using a FIBTEM test (a ready-to-use ROTEMsystem reagent for use with citrated whole blood, and assesses the firmness of the fibrin clot). The reporting of post-intervention coagulation tests varied, particularly with timing of tests. Between two and 10 tests were performed between two hours and day 11 post-intervention. The commonest reported time was 24 hours, which was conducted in six studies ([Bilecen 2017](#); [Collins 2017](#); [Jeppsson 2016](#); [Nascimento 2016](#); [Rahe-Meyer 2016](#); [Wikkelso 2015](#)).

Fibrinogen concentrate: therapeutic trials with active comparator

(Four trials: [Galas 2014](#); [Innerhofer 2017](#); [Lance 2012](#); [Tanaka 2014](#))

All four trials (with 220 participants) reported standard clotting assays and viscoelastic measures (all used ROTEM). Three studies reported coagulation factor levels, which varied between trials: factors II, VII, IX, X and XIII ([Galas 2014](#)); factors II, VIII, IX, X, anti-thrombin ([Lance 2012](#)); and factor XIII and anti-thrombin ([Innerhofer 2017](#)). All studies reported baseline results and three of the four reported 24-hour results ([Galas 2014](#); [Innerhofer 2017](#); [Tanaka 2014](#)).

Factor XIII (FXIII): prophylactic trials with inactive comparator

(Six trials: [Godje 2006](#); [Karkouti 2013](#); [Korte 2009](#); [Levy 2009](#); [Rasche 1982](#); [Shirahata 1990](#))

Five trials reported FXIII levels at baseline and at one or more subsequent time points ([Godje 2006](#); [Karkouti 2013](#); [Korte 2009](#); [Levy 2009](#); [Rasche 1982](#)), which varied widely between 30 minutes post-intervention ([Karkouti 2013](#); [Levy 2009](#)) to once a week testing ([Rasche 1982](#)). One trial, with 58 neonates, reported only one set of coagulation tests (factor XIII, Clauss fibrinogen, plasminogen, anti-thrombin, proteins C and S, alpha-2 anti-plasmin, plasmin-anti-plasmin and fibrinogen degradation products), taken within the first week of birth ([Shirahata 1990](#)). Standard clotting tests (activated partial thromboplastin time, prothrombin time and Clauss fibrinogen) were reported in two studies ([Godje 2006](#); [Rasche 1982](#)), and two studies reported Clauss fibrinogen only ([Korte 2009](#); [Shirahata 1990](#)). Only one study reported ROTEM measures ([Korte 2009](#)).

Factor XIII (FXIII): therapeutic trials with inactive comparator

(One trial: [Bregenzler 1999](#))

[Bregenzler 1999](#) reported FXIII measures at nine time points, including baseline and up to day 42 post-therapy in 20 participants. No other coagulation parameter was reported.

Prothrombin complex concentrate (PCC): prophylactic trials with inactive comparator

(One trial: [Turner 1981](#))

Standard clotting assays (activated partial thromboplastin time, prothrombin time, Clauss fibrinogen) were reported at two time points, baseline and post-intervention, in the 78 neonates in this trial ([Turner 1981](#)).

DISCUSSION

Summary of main results

We identified 31 randomised controlled trials which involved a total of 2392 participants, and compared the use of intravenous pro-coagulant haemostatic factors with either placebo or another haemostatically active agent.

Unlike inherited bleeding conditions, where a person has a deficiency in one coagulation factor alone, acquired coagulopathy is more complex, often involving multiple deficiencies of coagulant factors, and which also evolves with time. For example, in trauma haemorrhage, when trauma coagulopathy is present it leads initially to a reduction in pro-coagulant factors (i.e. including factors V, XIII and fibrinogen); an increase in fibrinolytic factors (i.e. tissue plasminogen activator (t-PA) with low plasminogen activator inhibitor type 1 (PAI-1) and alpha 2-antiplasmin (A2-AP)) as well as an alteration of anticoagulant factors (i.e. high activated protein C levels). These initial changes can be associated with a bleeding phenotype, although at later stages can shift towards a pro-thrombotic state. This has implications for the management of acquired bleeding, as it suggests that single pro-haemostatic therapies may have less effect. This review therefore was aimed at understanding the trial data to inform efficacy and safety of multiple pro-coagulants, or in combination.

Our review identified data on three different pro-haemostatic agents as interventions: fibrinogen (23 trials), Factor XIII (7 trials) and PCC (1 trial). While the earlier studies focused on factor XIII, there has been an increase in the investigation of fibrinogen concentrate since 2010, as demonstrated by fibrinogen concentrate being the focus of most of the trials published in the last five years. Eighteen trials explored the use of these pro-haemostatic agents as prophylaxis for bleeding and 13 trials explored their therapeutic use. Fibrinogen was the focus of 12 of the 13 therapy-orientated trials whilst six of the factor XIII trials tested used it as a prophylactic agent.

Summary of primary outcomes

The primary outcomes were all-cause mortality (reported by 21 trials), arterial thromboembolic events (reported by 22 trials) and venous thromboembolic events (reported by 21 trials). The results of meta-analyses did not demonstrate an impact of pro-haemostatic agents on mortality or the incidence of arterial or venous thromboembolic events. The overall quality of this evidence ranged from very low to high ([Summary of findings for the main comparison](#); [Summary of findings 2](#); [Summary of findings 3](#); [Summary of findings 4](#); [Summary of findings 5](#)).

Summary of secondary outcomes

Data were available for all of our secondary outcomes: mortality due to bleeding, red blood cell transfusion requirement, blood loss and allergic reaction. We also reported on changes to laboratory measures of coagulation, both standard laboratory tests and global measurements of coagulation, but meta-analysis of these data were limited by heterogeneity in the assays and reported results.

Overall there was no evidence of a decrease in mortality due to bleeding in any of the 10 trials that contributed data for this outcome. Fibrinogen concentrate was associated with a reduction in the individual patient risk of being exposed to a red blood cell

transfusion when compared to inactive comparators. This finding was evident in both cardiac and non-cardiac surgery settings and in the prophylactic and therapeutic settings ([Analysis 1.6](#), [Analysis 3.5](#)). Additionally, prophylactic fibrinogen concentrate was found to reduce post-operative bleeding in two studies of cardiac surgery and one trial in orthopaedic surgery, when compared to a haemostatically inactive comparator ([Analysis 1.7](#)). Overall, the prophylactic use of fibrinogen concentrate almost halves the risk of a person requiring a blood transfusion following cardiac surgery and reduces it by 75% in non-cardiac surgery compared to placebo ([Analysis 1.6](#)). Prophylactic factor XIII demonstrated efficacy at reducing the volume of post-operative bleeding in cardiac surgery compared to placebo ([Analysis 5.7](#)).

All studies that reported blood loss following surgery found greater blood loss in the haemostatically inactive arm (no use of pro-haemostatic agents), irrespective of differences in the clinical setting, the intervention or how and when blood loss was measured. Across all trials only a single intervention-related allergic adverse event was recorded in an inactive-control study arm participant.

Overall the data evaluating coagulation test results were highly variable, with data presenting a wide range of standard as well as global tests (viscoelastic and thrombin generation). There was no consensus on the timing or frequency of results which were used to evaluate the 'laboratory efficacy' of the intervention in each clinical trial. Of more importance is the overwhelming lack of any attempt by trialists to combine laboratory data with clinically meaningful endpoints; for example, if a pro-haemostatic drug is being given with the aim of preventing or treating bleeding, laboratory data would be best presented in combination with clinical endpoints such as 'cessation of bleeding' or 'time to haemostasis'.

Overall completeness and applicability of evidence

Completeness

The interpretation of the findings of our review are limited by the individual trial data that contributed to meta-analyses. Specifically, meta-analysis of the data across trials was not possible for many outcomes, due to a lack of common time points or inadequate or inconsistent definitions of time points. We were only able to report data on our primary outcomes of mortality and thromboembolic events in 21 out of 31 trials.

The identified studies recruited participants across a broad range of clinical settings in which pro-haemostatic agents are likely to be used. These include trauma, postpartum haemorrhage and cardiac surgery, where clinical outcomes might be modified significantly by optimal use of pro-haemostatic agents. However, other clinical areas of bleeding risk, such as gastro-intestinal bleeding, were poorly represented in the included trials.

Applicability

Applicability of the review findings is limited. None of the included studies was powered to detect harm or adverse events, and there is little documentation of adverse events to the 90-day time point that is recommended for hospital-associated thromboembolic complications. The poor quality of the evidence means that the results for outcomes assessed in this review cannot be applied in other clinical settings (See 'Summary of findings' tables).

Quality of the evidence

We used GRADE to summarise the quality of evidence for the two primary outcomes: mortality and thromboembolic events in the 'Summary of findings' tables. We focused on mortality for the studies assessing the use of pro-coagulant haemostatic agents as a treatment for bleeding and thromboembolic events to assess the prophylactic use of pro-coagulant haemostatic agents. The overall quality of the evidence ranged from very low to high, with most trial outcomes being rated as low quality. We considered no trial to be at low risk of bias in all domains, but we downgraded half the outcomes by one level for risk of bias. Domains with high risk of bias included allocation concealment, blinding of study personnel and outcome assessors, incomplete outcome data and selective reporting. The small cohorts and rare mortality and thrombotic events introduced risks of imprecision. Lastly, the trials in this review represented most of the clinical areas in which bleeding is observed, but not all clinical areas were represented in each of the intervention comparisons. Moreover the trials did not set out to explore the outcomes of interest to this review, and this introduced inconsistency. Lastly, we did not downgrade any outcome due to inconsistency, perhaps as a consequence of GRADE assessments being based on outcome data for one or two studies due to diversity in the clinical setting and the timing of outcome measurements, which limited our ability to pool data across studies (as has been highlighted above).

Potential biases in the review process

The searches performed for this review were extensive and we have tried to identify all trials eligible for inclusion, regardless of their publication status or language of publication. We have also attempted to obtain missing data from all the included studies to provide a complete picture of the primary and secondary review outcomes, and were able to acquire this from eight authors. The review authors performed screening and data extraction in duplicate to minimise bias and were blinded to each other's results.

Included study bias

Determining the risks of bias of these studies highlighted a range of methodological issues. A significant number of studies were unclear in their randomisation methodology (15 studies) and allocation concealment (14 at unclear risk and five at high risk of bias). Any impairment of the randomisation process is known to generate an overestimation of the clinical efficacy of any intervention (Schulz 1995). Similarly the blinding status of outcome assessors was often not addressed (13 at unclear risk and five at high risk of bias), although most outcomes in these trials were objective and therefore not prone to observer bias. More important, however, is the frequency with which study personnel were either unblinded or where insufficient information was available to assess their level of blinding (five at unclear risk and seven at high risk of bias). This is concerning for outcomes such as transfusion, especially where no algorithm or threshold for transfusion was applied, where decisions are likely to be prone to bias.

Limitations of this review

Most of the studies we identified that contributed to this review were small-scale, with a mean of 78 participants (range 20 to 479). This review is clearly limited to the data published and provided by the trial authors. A surprising number of studies did not provide mortality data as an outcome (10 studies). Most outcomes of

interest in this review are objective and have well-defined criteria, although the definitions of thromboembolic events, allergy and blood loss provided were less robust.

Sample sizes in future randomized trials powered on clinical outcomes of mortality (or even preferably mortality due to bleeding) would need to be greatly increased. Analogous examples for power calculations would be CRASH2 (CRASH-2 2010), WOMAN (WOMAN 2017) and HALT IT (Roberts 2014, Brenner 2018), which recruited between 15,000 and 20,000 participants each.

We have grouped the wide range of study clinical settings into the broad primary categories of therapy or prophylaxis. However, we recognise heterogeneity within these settings and at further levels. For example, trials within the cardiac group may include more routine coronary artery bypass graft surgery (Karlsson 2011; Sadeghi 2014), or complex aortic surgery (Rahe-Meyer 2016; Ranucci 2015), presumably with different levels of bleeding risk.

The study interventions, doses and administration regimens varied. However, several of these agents act through common pathways of coagulation (factor XIII and fibrinogen, for example, principally act to increase clot strength and reduce the susceptibility of clots to fibrinolysis), and in some recent trials these interventions were used in combination (Innerhofer 2017). Fibrinogen concentrate doses ranged from one gram (Pournajafian 2015; Sadeghi 2014) to six grams (Curry 2018; Nascimento 2016), and factor XIII therapy ranged from a single dose (Godje 2006; Karkouti 2013) to nine daily doses (Bregenzer 1999). In this review we have not differentiated between the doses given, as at present there are no robust data to support any one dose or dose calculation as providing clinical efficacy in a given setting. However, the more recent fibrinogen concentrate papers have implemented fibrinogen-dosing protocols based on fibrinogen levels or viscoelastic markers of clot strength (Bilecen 2017; Collins 2017; Cui 2010; Haas 2015; Rahe-Meyer 2013; Ranucci 2015; Sabate 2016). While this approach might eventually permit individualised approaches to product dosing, it increases the difficulty of comparing these studies. Additionally it is likely that some studies reported here administered haemostatic agent doses below that required for clinical efficacy (Pournajafian 2015; Sadeghi 2014), with the potential of false negative findings.

The published data on markers and indices of coagulation status were very heterogeneous and it is difficult to offer a useful interpretation of these data.

Agreements and disagreements with other studies or reviews

There are no other published reviews of this scope on factor XIII for comparison. The paucity of data for the use of pro-thrombin complex concentrates may have precluded systematic reviews of randomised trials, although there are reviews of observational studies (Zeng 2017).

Clinical findings and conclusions

The findings of this review are consistent with other systematic reviews of the use of fibrinogen concentrate. An analysis of one prospective and seven retrospective studies in trauma found no impact on overall mortality and recorded a broad range of recorded mortality time points and heterogeneity in the reported outcomes (Mengoli 2017). Similarly, analysis of blood loss and transfusion was impaired by variation in the use of continuous and discrete

data presentation. Another systematic review that included four studies (two RCTs and two non-randomised controlled studies) concluded that fibrinogen concentrate in the perioperative and major haemorrhage settings was safe and reduced bleeding and the transfusion of blood components (Franchini 2012). A Cochrane Review of fibrinogen concentrate in bleeding patients was published in 2013 and contains six RCTs that compared fibrinogen to placebo or usual care in adults with bleeding (Wikkelso 2013). We have included these RCTs in this review (Cui 2010; Fenger Erikson 2009; Galas 2014; Karlsson 2011; Lance 2012; Rahe-Meyer 2013). In keeping with our conclusions, the authors found that fibrinogen concentrate reduced transfusion requirement, but there were insufficient data to draw conclusions about the impact on mortality or other clinical efficacy.

Risk of bias and quality of included studies

The three reviews highlighted above reached comparable conclusions to this review about the poor quality of evidence and frequently indeterminate or high risks of bias in the included studies.

AUTHORS' CONCLUSIONS

Implications for practice

The inadequate quality of evidence in most of the studies included in this review means that we cannot draw conclusions for clinical practice or the use of these interventions outside controlled trials. Further research is required in larger-scale randomised controlled trials to determine the risk-to-benefit ratio of these interventions. We anticipate that data from the identified ongoing trials, some of which are larger studies, will provide much higher-quality evidence around clinical outcomes and efficacy.

At present we lack a clear picture of the cost effectiveness of these interventions, due to a lack of health economic data. It is likely that many pro-haemostatic agents are more costly than blood components, including plasma. Most studies co-administer the pro-coagulant agents with a broad range of other haemostatic interventions as part of the clinical setting or routine management of haemorrhage. Additionally, the progression towards viscoelastic testing and algorithmic clotting factor approaches to transfusion means that these agents are often administered in combination. However, at present we lack the data required to assess how these interventions interact or work together.

Implications for research

The main implication for this review is to direct the need for further research. Whilst at one level the numbers of trials represents a body of data to inform clinical practice, the limitations preclude the drawing of robust conclusions without additional studies. There is considerable current clinical interest in the use of these agents (we identified 22 ongoing trials), which is likely to expand in the coming years. Sample sizes in future randomised trials powered for clinical outcomes of mortality (or even preferably mortality due to bleeding) would need to be greatly increased. Analogous examples for power calculations would be CRASH-2 (CRASH-2 2010), WOMAN (WOMAN 2017) and HALT IT (Brenner 2018; Roberts 2014), which recruited between 15,000 and 20,000 participants each.

To improve consistency in outcome reporting, the use of core outcome sets (COS) are increasingly advocated by both the CONSORT and SPIRIT statements (Moher 2010; Chan 2013), and may help address a number of the limitations in reporting identified in this review. A COS is defined as a “minimum (number of outcomes) that should be measured and reported in all clinical trials of a specific condition” (Gargon 2017). The Core Outcome Measures in Effectiveness Trials (COMET) database has assisted in driving international collaboration amongst clinical researchers across a plethora of clinical conditions, such as critical care and rheumatological diseases, and it may be timely to look at this methodology for outcomes of bleeding. In particular, consideration of how to combine clinically meaningful endpoints (such as achievement of haemostasis) with reliable laboratory data must be a focus for future studies, in order to improve the quality of the available evidence for laboratory thresholds to guide clinicians about when to administer haemostatic therapy. Future updates of this review may need to be split by interventions used as treatment versus prophylaxis.

Researchers may also consider the lessons from previous research in the management of coagulopathy and bleeding. Firstly, the extensive trial programmes undertaken to evaluate recombinant factor VIIa consisted initially of poor-quality studies with small participant numbers in a heterogeneous range of clinical settings which demonstrated variable degrees of clinical efficacy that overestimated the benefit while concealing the underlying harm. It is essential that this process is not repeated. Secondly, there were two large studies of tranexamic acid (TXA) that demonstrated a positive impact on death from bleeding. CRASH-2 was a study of just over 20,000 participants that showed that the administration of TXA reduced overall mortality and death from bleeding at a statistically significant level (CRASH-2 2010). Subsequent analysis of the data, in combination with data from over 20,000 participants in the WOMAN study of TXA in postpartum haemorrhage (WOMAN 2017), found that administration of an early dose of TXA improved outcomes. The multicentre and multi-national design, the pragmatic approach and large participant numbers of these studies permitted statistically robust and clinically useful conclusions to be drawn about practice in a real-world setting.

ACKNOWLEDGEMENTS

We would like to thank Drs Graham Barker, Gemma Simons, Yulia Lin and Stuart McKechnie for their help with the development and writing of the protocol for this review.

Drs Gemma Simons (GS) and Yulia Lin (YL) were involved with the screening (YL), data extraction (GS, YL), and risk of bias assessment (GS) of the studies identified at the first search, and we thank them for their valuable help with these activities.

We would like to acknowledge the assistance and expertise of Tahereh Ziafazel in the translation of Pournajafian 2015.

This project was supported by the UK National Institute for Health Research, through Cochrane Infrastructure funding to the Cochrane Injuries Group. The views and opinions expressed are those of the authors and do not necessarily reflect those of the Systematic Reviews Programme, NIHR, NHS or the Department of Health.

REFERENCES

References to studies included in this review

Bilecen 2017 {published data only}

Bilecen S, de Groot JAH, Kalkman CJ, Spanjersberg AJ, Bruinsma GJBB, Moons KGM, et al. Effect of Fibrinogen Concentrate on Intraoperative Blood Loss Among Patients With Intraoperative Bleeding During High-Risk Cardiac Surgery. *JAMA* 2017;**317**(7):738-47.

Bregenzner 1999 {published data only}

Bregenzner N, Carsar I, Andus T, Hämling J, Malchow H, Schreiber S, et al. Lack of clinical efficacy of additional factor XIII treatment in patients with steroid refractory colitis. *Zeitschrift für Gastroenterologie* 1999;**37**(10):99-104.

Collins 2017 {published data only}

Bruynseels D, Dick J, Elton C, Mallaiah S, Collis RE. International Journal of Obstetric Anesthesia. 2016; Vol. 26:S62016.

* Collins PW, Cannings-John R, Bruynseels D, Mallaiah S, Dick J, Elton C, et al. Viscoelastometric-guided early fibrinogen concentrate replacement during postpartum haemorrhage: OBS2, a double-blind randomized controlled trial. *British Journal of Anaesthesia* 2017;**119**(3):411-21.

Weeks A, Collis R, Mallaiah S, Elton C, Dick J, Canning-John R, et al. Fibrinogen concentrate versus placebo for treatment of postpartum haemorrhage: A multicentre, prospective, double blind randomised control pilot study (OBS2). *BJOG: An International Journal of Obstetrics and Gynaecology*. 2016; Vol. 123:177.

Cui 2010 {published data only}

Cui Y, Hei F, Long C, Feng Z, Zhao J, Yan F, et al. Perioperative monitoring of thromboelastograph on blood protection and recovery for severely cyanotic patients undergoing complex cardiac surgery. *Artificial Organs* 2010;**34**(11):955-60.

Curry 2018 {unpublished data only}

A multi-centre, randomised, double blind, placebo-controlled trial evaluating the effects of early administration of fibrinogen concentrate in adults with major traumatic haemorrhage. ISRCTN registry ISRCTN67540073 (accessed 05.02.2018).

* Curry N, Foley C, Wong H, Mora A, Curnow E, Zarankaite A, et al. Early fibrinogen concentrate therapy for major haemorrhage in trauma (E-FIT 1). Results from a UK multi-centre, randomised, double blind, placebo-controlled pilot trial and comparisons with two fibrinogen supplementation trauma trials. The primary author, Dr Nicola Curry.

Fenger Erikson 2009 {published data only}

Fenger-Eriksen C, Jensen TM, Kristensen B, Jensen KM, Tonnesen E, Ingerslev J, et al. Fibrinogen concentrate corrects dilutional coagulopathy: a prospective, randomized placebo controlled clinical trial. *Blood* 2008;**112**(111):Abstract Number 1227.

* Fenger-Eriksen C, Jensen TM, Kristensen BS, Jensen KM, Tonnesen E, Ingerslev J, et al. Fibrinogen substitution improves

whole blood clot firmness after dilution with hydroxyethyl starch in bleeding patients undergoing radical cystectomy: a randomized, placebo-controlled clinical trial. *Journal of Thrombosis & Haemostasis* 2009;**7**(5):795-802.

Galas 2014 {published data only}

Galas F, Hajjar L, Sorensen B, Almeida J, Sundin M, Guimaraes V, et al. Randomized comparison of fibrinogen concentrate versus cryoprecipitate for bleeding control in paediatric cardiac surgery (FICCS study). *Critical Care* 2012;**16**(Suppl 1):S156-7. [PUBMED: PMC3363856]

* Galas FRBG, de Almedia JP, Fukushima JT, Vincent JL, Osawa EA, Zeferino S, et al. Hemostatic effects of fibrinogen concentrate compared with cryoprecipitate in children after cardiac surgery: A randomized pilot trial. *The Journal of Thoracic and Cardiovascular Surgery* 2014;**148**(4):1647-55.

Godje 2006 {published data only}

Godje O, Gallmeier U, Calatzis A, Schelian M, Friedl R, Grunewald M, et al. Reduction of blood loss and blood transfusions after cardiac surgery by application of plasma coagulation factor XIII. *Annals of Hematology*. 2002. 2002; Vol. 81 (Suppl 1):A22.

* Godje O, Gallmeier U, Schelian M, Grunewald M, Mair H. Coagulation factor XIII reduces postoperative bleeding after coronary surgery with extracorporeal circulation. *Thoracic & Cardiovascular Surgeon* 2006;**54**(1):26-33.

Haas 2015 {published data only}

Fibrinogen for treatment of paediatric dilutional coagulopathy FibPaed study. ClinicalTrials.gov NCT 01487837.

* Haas T, Spielmann N, Restin T, Seifert B, Henze G, Obwegeser J, et al. Higher fibrinogen concentrations for reduction of transfusion requirements during major paediatric surgery: a prospective randomised controlled trial. *British Journal of Anaesthesia* 2015;**115**(2):234-43.

Haas T, Spielmann N, Cushing M. Dosing of fibrinogen concentrate during major paediatric surgery-calculating the increase in clot firmness. *Transfusion medicine. Conference: 18th annual NATA symposium on patient blood management, haemostasis and thrombosis*. 2017; Vol. 27:63.

Haas T, Spielmann N, Cushing M, Schmugge M, Weiss M. Influence of factor XIII levels on clot firmness during major paediatric surgery. *Transfusion Medicine*. 2015; Vol. 25:45.

Innerhofer 2017 {published data only}

Correction to Lancet Haematol 2017; 4: e258-71. The Lancet Haematology 2017; Vol. 4, issue 6:e257.

* Innerhofer P, Fries D, Mittermayr M, Innerhofer N, von Langen D, Hell T, et al. Reversal of trauma-induced coagulopathy using first-line coagulation factor concentrates or fresh frozen plasma (RETIC): a single-centre, parallel-group, open-label, randomised trial. *Lancet Haematology* 2017;**4**(6):e258-71.

Jeppsson 2016 {published data only}

Jeppsson A, Walden K, Roman-Emanuel C, Thimour-Bergstrom L, Karlsson M. Preoperative supplementation with fibrinogen concentrate in cardiac surgery: A randomized controlled study. *British Journal of Anaesthesia* 2016;**116**(2):208-14.

Karkouti 2013 {published data only}

Karkouti K, von Heymann C, Jespersen CM, Korte W, Levy JH, Ranucci M, et al. Efficacy and safety of recombinant factor XIII on reducing blood transfusions in cardiac surgery: A randomized, placebo-controlled, multicenter clinical trial. *The Journal of Thoracic and Cardiovascular Surgery* 2013;**146**(4):927-39. [PUBMED: 23820174]

Karlsson 2011 {published data only}

Baghaei F, Karlsson M, Ternstrom L, Hyllner M, Flinck A, Skrtic S, et al. Prophylactic fibrinogen infusion reduces bleeding after coronary artery bypass surgery: A prospective randomized pilot study. *Journal of Thrombosis and Haemostasis* 2009;**7**(S2):697.

Karlsson M, Ternstrom L, Hyllner M, Baghaei F, Flinck A, Skrtic S, et al. Prophylactic fibrinogen infusion reduces bleeding after coronary surgery. A prospective randomised pilot study. *Thrombosis & Haemostasis* 2009;**102**(1):137-44.

* Karlsson M, Ternstrom L, Hyllner M, Baghaei F, Skrtic S, Jeppsson A. Prophylactic fibrinogen infusion in cardiac surgery patients: effects on biomarkers of coagulation, fibrinolysis, and platelet function. *Clinical & Applied Thrombosis/Hemostasis* 2011;**17**(4):396-404.

Korte 2009 {published data only}

Korte WC, Szadkowski C, Gahler A, Gabi K, Kownacki E, Eder M, et al. Factor XIII substitution in surgical cancer patients at high risk for intraoperative bleeding. *Anesthesiology* 2009;**110**(2):239-45.

Lance 2012 {published data only}

Lance MD, Ninivaggi M, Hamulyak K, van Pampus EC, ten Cate H, Heemskerk JW. Early transfusion with FFP or FFP plus fibrinogen concentrate in massive hemorrhage: A randomised intervention trial. *Transfusion and Haemostasis* 2005;**100**(5):1416-21.

Lance MD, Ninivaggi M, Marcus MAE, Hamulyak K, van Pampus ECM, ten Cate H, et al. Partial pro-hemostatic effect of transfusion of fibrinogen concentrate replacing fresh frozen plasma in massive hemorrhage. *Journal of Thrombosis and Haemostasis* 2011;**9**(Suppl 2):S229.

* Lance MD, Ninivaggi M, Schols SEM, Feijge MAH, Oehrl SK, Kuiper GJ, et al. Perioperative dilutional coagulopathy treated with fresh frozen plasma and fibrinogen concentrate: a prospective randomized intervention trial. *Vox Sanguinis* 2012;**103**:25-34.

Levy 2009 {published data only}

Safety of rFXIII in patients following first time myocardial revascularisation requiring cardiopulmonary bypass. ClinicalTrials.gov; reference number NCT02239146.

* Levy JH, Gill R, Nussmeier NA, Olsen PS, Andersen HF, Booth FV, et al. Repletion of factor XIII following cardiopulmonary bypass using a recombinant A-subunit homodimer. A preliminary report. *Thrombosis & Haemostasis* 2009;**102**(4):765-71.

Najafi 2014 {published data only}

Effect of prophylactic administration of fibrinogen concentrate on volume of transfused packed RBC during and after total hip arthroplasty in hospital during 2011-12. IRCT trial number 201012264784N2.

* Najafi A, Moharari RS, Orandi AA, Etezadi F, Sanatkar M, Khajavi MR, et al. Prophylactic administration of fibrinogen concentrate in perioperative period of total hip arthroplasty: a randomized clinical trials study. *Acta Medica Iranica* 2014;**52**(11):804-10.

Nascimento 2016 {published data only}

Nascimento B, Callum J, Tien H, Peg H, Rizoli S, Karanicolas P, et al. Fibrinogen in the initial resuscitation of severe trauma (FiIRST): a randomized feasibility trial. *British Journal of Anaesthesia* 2016;**117**(6):775-82.

Pournajafian 2015 {published data only}

Pournajafian AL, Ghodraty MR, Moheeni M, Ghamari AA, Sadeghi F. Reduction of intraoperative bleeding with fibrinogen administration in posterior spinal fusion surgery. *Journal of Isfahan Medical School* 2015;**33**(347):1370-9.

Rahe-Meyer 2013 {published data only}

Haemocompletan®P during aortic replacement. www.ClinicalTrials.gov, trial identifier: NCT00701142 [accessed 05 January 2016].

Rahe-Meyer N, Hanke A, Schmidt DS, Hagl C, Pichlmaier M. Fibrinogen concentrate reduces intraoperative bleeding when used as first-line hemostatic therapy during major aortic replacement surgery: results from a randomized, placebo-controlled trial. *Journal of Thoracic and Cardiovascular Surgery* 2013;**145**(3 Suppl.):S178-85. [PUBMED: 23410777]

Rahe-Meyer N, Solomon C, Hanke A, Schmidt DS, Hagl C, Pichlmaier M. Haemostatic therapy with fibrinogen concentrate for controlling major bleeding during complex cardiovascular surgery: results of a randomised, placebo-controlled trial. *British Journal of Anaesthesia*. 2012; Vol. 108 (S2):ii75-104.

* Rahe-Meyer N, Solomon C, Hanke A, Schmidt DS, Knoerzer D, Hochleitner G, et al. Effects of Fibrinogen Concentrate as First-line Therapy during Major Aortic Replacement Surgery. *Anesthesiology* 2013;**118**:40-50. [PUBMED: 23249928]

Solomon C, Hagl C, Rahe-Meyer N. Time course of haemostatic effects of fibrinogen concentrate administration in aortic surgery. *British Journal of Anaesthesia* 2013;**110**(6):947-56.

Solomon C, Rahe-Meyer N. Fibrinogen concentrate as first-line therapy in aortic surgery reduces transfusion requirements in patients with platelet counts over or under 100X10⁹/L. *Blood Transfusion* 2015;**13**:248-54.

Solomon C, Schochl H, Hagl C, Rahe-Meyer N. Fibrinogen concentrate as first line haemostatic therapy during aortic replacement surgery. *International Society of thrombosis and Haemostasis*. 2012; Vol. 10:e7.

Solomon C, Schochl H, Hanke A, Hagl C, Rahe-Meyer N. Fibrinogen concentrate causes a short lived increase in plasma fibrinogen and fibrin based clot quality when used as a first line haemostatic therapy during aortic replacement surgery: a randomized trial. *British Journal of Anaesthesia*. 2012; Vol. 108 (S2):ii276-7.

Rahe-Meyer 2016 {published data only}

Rahe-Meyer N, Levy JH, Mazer CD, Schramko A, Klein AA, Brat R, et al. Randomized evaluation of fibrinogen vs placebo in complex cardiovascular surgery (REPLACE): a double blind phase III study of haemostatic therapy. *British Journal of Anaesthesia* 2016;**117**(1):41-51.

Ranucci 2015 {published data only}

The ZERo PLASma Trial (ZEPLAST): Avoidance of Fresh Frozen Plasma in Cardiac Surgery. www.ClinicalTrials.gov, registry number NCT01471730 (accessed 05.02.2018).

Ranucci M, Baryshnikova E. Fibrinogen supplementation after cardiac surgery: insights from the Zero-Plasma trial (ZEPLAST). *British Journal of Anaesthesia* 2016;**116**(5):618-23.

* Ranucci M, Baryshnikova E, Crapelli GB, Rahe-Meyer N, Menicanti L, Frigola A. Randomized, double-blinded, placebo-controlled trial of fibrinogen concentrate supplementation after complex cardiac surgery. *Journal of the American Heart Association* 2015;**4**:e002066.

Rasche 1982 {published data only}

Rasche H, Haghou F, Gaus W, Dietrich M, Hoelzer D, Pflieger H, et al. [Blood clotting factor XIII substitution in acute leukaemia: result of a randomized and controlled study]. [German]. [Blutgerinnungsfaktor-XII-Substitution bei akuter Leukämie]. *Deutsche Medizinische Wochenschrift* 1982;**107**(49):1482-6.

Sabate 2016 {published data only}

Sabate A, Gutierrez R, Beltra J, Mellado P, Blasi A, Acosta F, et al. Impact of preemptive fibrinogen concentrate on transfusion requirements in liver transplantation: a multicenter, randomized, double-blind, placebo-controlled trial. *American Journal of Transplantation* 2016;**16**:2421-9.

Sadeghi 2014 {published data only}

Effect of prophylactic fibrinogen concentrate infusion on postoperative bleeding in coronary artery bypass graft surgery: a randomised, double blind clinical trial. IRCT trial registration number 201011085140N1.

* Sadeghi M, Atefyekta R, Azimaraghi O, Marashi SM, Aghajani Y, Ghadimi F, et al. A randomized, double blind trial of prophylactic fibrinogen to reduce bleeding in cardiac surgery [Estudo randômico e duplo-cego de profilaxia com fibrinogênio para reduzir o sangramento em cirurgia cardíaca]. *Revista Brasileira de Anestesiologia* 2014;**64**(4):253-7.

Shirahata 1990 {published data only}

Nakamura T, Ariyoshi N, Shirahata A. Effect of prophylactic administration of factor XIII concentrate on intraventricular haemorrhage in premature infants. *Thrombosis and Haemostasis* 2009;**62**(1989):287.

* Shirahata A, Nakamura T, Shimono M, Kaneko M, Tanaka S. Blood coagulating findings and the efficacy of Factor XIII concentrate in premature infants with intracranial haemorrhages. *Thrombosis Research* 1990;**57**:755-63.

Soleimani 2016 {published data only}

Masoumi N, Soleimani M, Lashay A, Nooraie N. Fibrinogen concentrate effect on perioperative bleeding in transurethral resection of the prostate: a double blind placebo-controlled and randomized study. *BJU International*. 2017; Vol. 119:44.

* Soleimani M, Masoumi N, Nooraie N, Lashay A, Safarienejad MR. The effect of fibrinogen concentrate on perioperative bleeding in transurethral resection of the prostate: a double-blind placebo-controlled and randomized study. *Journal of Thrombosis and Haemostasis* 2017;**15**:255-62.

Tanaka 2014 {published data only}

RiaSTAP vs. Conventional Transfusion in Patients Having Heart Valve Surgery (RiaCT). ClinicalTrials.gov Identifier: NCT01283321; terminated 2015 (insufficient funding to complete total projected enrolment; accessed 05.02.2018).

* Tanaka AK, Egan K, Szlam F, Ogawa S, Roback JD, Sreeram G, et al. Transfusion and hematologic variables after fibrinogen or platelet transfusion in valve replacement surgery: preliminary data of purified lyophilized human fibrinogen concentrate versus conventional transfusion. *Transfusion* 2014;**54**(1):109-18. [PUBMED: 23718572]

Turner 1981 {published data only}

Turner T, Prowse CV, Prescott RJ, Cash JD. A clinical trial on the early detection and correlation of haemostatic defects in selected high-risk neonates. *British Journal of Haematology* 1981;**47**:65-75. [PUBMED: 7002204]

Wikkelso 2015 {published data only}

Fibrinogen Concentrate as Initial Treatment for Postpartum Haemorrhage: A Randomised Clinically Controlled Trial (FIB-PPH). www.ClinicalTrials.gov, registry number NCT01359878 (accessed 06 January 2016).

* Wikkelssoe AJ, Edwards HM, Afshari A, Stensballe J, Langhoff-Roos J, Albrechtsen C, et al. Pre-emptive treatment with fibrinogen concentrate for postpartum haemorrhage: randomised controlled trial. *British Journal of Anaesthesia* 2015;**114**(4):623-33.

Wikkelssoe AJ, Edwards HM, Afshari A, Stensballe J, Langhoff-Roos J, Albrechtsen C, et al. Fibrinogen concentrate as initial treatment for postpartum haemorrhage: FIB-PPH study. *Transfusion Medicine*. 2015; Vol. 25 (S1):7-72.

Wikkelssoe AJ, Afshari A, Stensballe J, Langhoff-Roos J, Albrechtsen C, Ekelund K, et al. The FIB-PPH trial: fibrinogen concentrate as initial treatment for postpartum haemorrhage:

study protocol for a randomised controlled trial. *Trials* 2012;**13**(110):1-8.

References to studies excluded from this review

ACTRN12615001293550p 2015 {published data only}

To determine the feasibility of early administration of Fibrinogen concentrate in trauma patients with ongoing haemorrhagic shock requiring activation of the massive transfusion protocol. ANZCTR registry identifier ACTRN12615001293550p (accessed 05.02.2018).

Boulis 1999 {published data only}

Boulis NM, Bobek MP, Schmaier A, Hoff JT. Use of factor IX complex in warfarin-related intracranial haemorrhage. *Neurosurgery* 1999;**45**(5):1113-8.

Curry 2015 {published data only}

Curry N, Rourke C, Davenport R, Beer S, Pankhurst L, Deary A, et al. Early cryoprecipitate for major haemorrhage in trauma: a randomised controlled feasibility trial. *British Journal of Anaesthesia* 2015;**115**(1):76-83.

Demeyere 2010 {published data only}

Demeyere R, Arnout J, Strengers P, Gasthuisberg UZ. Prothrombin complex concentrate versus fresh frozen plasma in patients on oral anticoagulant therapy undergoing cardiac surgery: a randomized study. *Critical Care* 2006;**10**(Suppl 1):233.

* Demeyere R, Gillardin S, Arnout J, Strengers PF. Comparison of fresh frozen plasma and prothrombin complex concentrate for the reversal of oral anticoagulants in patients undergoing cardiopulmonary bypass surgery: a randomized study. *Vox Sanguinis* 2010;**99**(3):251-60.

Garrigue 2017 {published data only}

Garrigue D, Godier A, Glacet A, Labreuche J, Kipnis E, Paris C, et al. French lyophilized plasma versus fresh frozen plasma for the initial management of trauma-induced coagulopathy: a randomized open-label trial. *Journal of Thrombosis & Haemostasis* 2018;**16**(3):481-9.

Goldstein 2015 {published data only}

An open label, randomized, multicentre, phase IIIb to assess the safety, efficacy and tolerance of BERIPLEX P/N (Kcentra) compared with plasma for rapid reversal of coagulopathy induced by vitamin K antagonists in subjects requiring an urgent surgical procedure (BE1116_3003). *ClinicalTrials.gov* NCT00803101.

* Goldstein JN, Refaai MA, Milling TJ Jr, Lewis B, Goldberg-Alberts R, Hug BA, et al. Four-factor prothrombin complex concentrate versus plasma for rapid vitamin K antagonist reversal in patients needing urgent surgical or invasive interventions: a phase 3b, open-label, non-inferiority, randomised trial. *The Lancet* 2015;**385**:2077-87.

Milling TJ Jr, Refaai MA, Sarode R, Lewis B, Mangione A, Durn BL, et al. Safety of 4-factor prothrombin complex concentrate (4F-PCC) versus plasma for vitamin K antagonist reversal: an

integrated data analysis of two phase III clinical trials. *American Journal of Haematology*. 2014; Vol. 89 (6):E26.

Refaai MA, Goldstein JN, Lee ML, Milling TJ Jr, Sarode R. Increased risk of volume overload with plasma compared to four-factor prothrombin complex concentrate for the emergency reversal of vitamin K antagonist therapy. *Academic Emergency Medicine* 2014;**21**(5):S243.

Refaai MA, Goldstein JN, Milling TJ Jr, Lewis B, Goldberg-Alberts R, Hug BA, et al. Randomized phase IIIb study of rapid vitamin K antagonist reversal in patients requiring an urgent surgical procedure: four-factor prothrombin complex concentrate is superior to plasma. *Blood*. 2013; Vol. 122:21.

Masoumi 2016 {published data only}

Masoumi SG, Bigdelian H, Mousavian SM, Mirzaei A, Abedi H, Ziaiefard M. Comparing the efficacy of synthetic fibrinogen plus albumin to fresh frozen plasma in the prevention of coagulation disorders in pediatric heart surgery. *Isfahan University of Medical Sciences* 2016;**119**:44.

Miyata 2014 {published data only}

Miyata S, Sasaki H, Ueda Y, Takamoto S, Usui A, Nishiwaki K, et al. Liberal versus restrictive transfusion strategy of platelet concentrate and cryoprecipitate in thoracic aortic surgery: A multicenter randomized trial. Abstracts of the HAA 2014 Annual Scientific Meeting. 2014:407.

NCT01706159 {published data only}

A Multicenter, Randomised, Double-blind, Placebo-controlled, Multiple-dose Trial With rFXIII Administered to Subjects With Mild to Moderate Active Ulcerative Colitis. *ClinicalTrials.gov* Identifier: NCT01706159 (accessed 05.02.2018).

NCT02542306 {published data only}

Notes for Safety and Efficacy of Fibrinogen Concentrate in Aortic Arch Surgery Involving Moderate Hypothermic Circulatory Arrest. *ClinicalTrials.gov* Identifier: NCT02542306 (accessed 20.4.2018).

NCT02736812 {published data only}

Interest of Pre-hospital Administration of Lyophilized Plasma to Prevent or Treat Coagulopathy Associated With Post-traumatic Hemorrhagic Shock (PREHO-PLYO Study). *ClinicalTrials.gov* Identifier: NCT02736812 (accessed 05.08.2018).

Papakitsos 2014 {published data only}

Papakitsos G, Kapsali A, Papakitsou T, Roimba A. Comparison of two dose-regimens of prothrombin complex concentrate in urgent anticoagulation reversal: A prospective randomised study. *Haematologica*. 2014; Vol. 99:267.

Qiu 2016 {published data only}

Qiu M, Zhang X, Cai H, Xu Z, Lin H. The impact of hemocoagulase for improvement of coagulation and reduction of bleeding in fracture-related hip hemiarthroplasty geriatric patients: a prospective, single-blinded, randomized, controlled study. *Injury* 2016;**48**(4):914-9.

Sarode 2013 {published data only}

Lang E. 4F-PCC was noninferior to plasma in patients with acute major bleeding who needed urgent VKA reversal. American College of Physicians [Journal Club] 2014; Vol. 160, issue 8:JC6.

Milling TJ Jr, Refaai MA, Goldstein JN, Goldberg-Alberts R, Schneider A, Hug B, et al. Thrombotic events after vitamin K antagonist reversal with 4-factor prothrombin complex concentrate or plasma. *Annals of Emergency Medicine*. 2013; Vol. 62 (4s):S10.

Milling TJ Jr, Refaai MA, Sarode R, Lewis B, Mangione A, Durn BL, et al. Safety of 4-factor prothrombin complex concentrate (4F-PCC) versus plasma for vitamin K antagonist reversal: an integrated data analysis of two phase III clinical trials. *American Journal of Haematology*. 2014; Vol. 89 (6):E26.

Refaai MA, Goldstein JN, Lee ML, Milling TJ Jr, Sarode R. Increased risk of volume overload with plasma compared to four-factor prothrombin complex concentrate for the emergency reversal of vitamin K antagonist therapy. *Academic Emergency Medicine*. 2014; Vol. 21:5 (Suppl 1).

Refaai MA, Goldstein JN, Milling TJ, Foehl H, Hug B, Sarode R. Correction of INR and coagulation factor levels in a randomized clinical trial of a four-factor prothrombin complex concentrate (PCC-4) versus plasma for urgent vitamin K antagonist reversal. *Annals of Emergency Medicine*. 2012; Vol. 60 (4s):S10.

Sarode R, Milling T, Refaai MA, Mangione A, Schneider A, Goldstein JN. Randomized Phase IIb study comparing the safety and efficacy of four-factor prothrombin complex concentrate with plasma in subjects receiving Vitamin K antagonists with major bleeding. *American Journal of Haematology*. 2012; Vol. 87 (S1):S146.

* Sarode R, Milling TJ, Refaai MA, Mangione A, Schneider A, Durn BL, et al. Efficacy and safety of a 4-factor prothrombin complex concentrate in patients on vitamin K antagonists presenting with major bleeding: a randomized, plasma-controlled, phase IIb study. *Circulation* 2013;**128**:1234-43. [PUBMED: 23935011]

Scherer 1994 {published data only}

Scherer R, Gille A, Erhard J, Paar D, Kox WJ. [The effect of substitution with AT III- and PPSB-concentrates in patients with terminal liver insufficiency]. [German] [Substitutionseffekt von AT III- und PPSB-Konzentraten bei Patienten mit terminaler Leberinsuffizienz]. *Anaesthesist* 1994;**43**(3):178-82.

TOP-CLOT Trial {published data only}

Trial of RiaSTAP Versus Cryoprecipitate to Lower Operative Transfusions (TOP-CLOT). www.ClinicalTrials.gov, registry number (accessed 05.02.2018); study terminated (feasibility issues prevent completion of recruitment).

UMIN000022194 {published data only}

Invasive intervention study on the safety of fibrinogen concentrate used in obstetric hemorrhage. UMIN-CTR register: UMIN000022194 (accessed 05.02.2018).

UMIN000022666 {published data only}

Application of fibrinogen concentrate (Fibrinogen HT) to uncontrollable massive bleeding. UMIN-CTR registry: UMIN000022666 (accessed 05.02.2018).

van Aart 2006 {published data only}

van Aart L, Eijkhout H, Kamphuis JS, Dam M, Strengers PFW, Ploeger B. The use of prothrombin complex investigated [Gebruik van protrombinecomplex onderzocht [Dutch]. *Pharmaceutisch weekblad* 2001;**136**(3):84-7.

* van Aart L, Eijkhout HW, Kamphuis JS, Dam M, Schattenkerk ME, Schouten TJ, et al. Individualized dosing regimen for prothrombin complex concentrate more effective than standard treatment in the reversal of oral anticoagulant therapy: An open, prospective randomized controlled trial. *Thrombosis Research* 2006;**118**:313-20. [PUBMED: 16182346]

References to studies awaiting assessment
Siemens 2017 {published data only}

Fibrinogen concentrate supplementation in the management of bleeding during paediatric cardiopulmonary bypass: a phase 1B/2A, open label dose escalation study. EudraCT Number: 2013-003532-68 (accessed 05.04.2018).

* Siemens K, Hunt BJ, Harris J, Nyman A, Perkins J, Murdoch IA, et al. Fibrinogen in paediatric cardiac surgery: the FIB-CON trial. *Transfusion Medicine*. 2017; Vol. 27:19.

References to ongoing studies
ANZCTR12613001279718 {published data only}

Prothrombinex-VF vs Fresh Frozen Plasma for the treatment of bleeding post cardiopulmonary bypass. Australian New Zealand Clinical Trials Registry 12613001279718 (accessed 13 January 2016).

ANZCTR12614000691640 {published data only}

The administration of fibrinogen concentrate (a component of blood involved in clotting process) to be given to those women who, after delivery of babies, develop severe bleeding to establish if this drug will prevent further blood loss or reduce the volume of blood loss. Australia New Zealand Clinical Trial Registry, number: ACTRN12614000691640 (accessed 13 January 2016).

EudraCT: 2010-022923-31 {published data only}

A multicenter double-blind, placebo controlled, randomized, pilot trial to assess the efficacy of pre-hospital administration of Fibrinogen Concentrate (FGTW) in trauma patients, presumed to bleed (FI in TIC). EudraCT Identifier: 2010-022923-31 (accessed 20.4.18).

EudraCT: 2013-003532-68 {published data only}

Fibrinogen concentrate supplementation in the management of bleeding during paediatric cardiopulmonary bypass: a phase 1B/2A, open label dose escalation study. EudraCT Identifier: 2013-003532-68 (accessed 13 January 2016).

EudraCT: 2013-004484-31 {published data only}

Rivaroxaban and PCC: Prothrombin Complex Concentrate in patients with bleeding complications related to Rivaroxaban [Eine prospektive klinische Pilotstudie zur Untersuchung der Wirksamkeit von Prothrombin Komplex Konzentraten (Beriplex®) bei Patienten mit Blutungskomplikationen unter der Einnahme von Rivaroxaban (Xarelto®)]. EudraCT Identifier: 2013-004484-31 (accessed 13 January 2016).

NCT00994045 {published data only}

Coagulopathy During Surgery for the Repair of Extent 4 Thoraco-Abdominal Aortic Aneurysms - Feasibility Study of the Use of Fibrinogen Concentrate by Infusion in Place of Fresh Frozen Plasma. ClinicalTrials.gov Identifier: NCT00994045 (accessed 05.02.2018).

NCT01539057 {published data only}

The Efficacy of the Administration of Fibrinogen in Liver Transplantation (FibstudLT). ClinicalTrials.gov Identifier: NCT01539057 (accessed 13 January 2016).

NCT02155725 {published data only}

Study on the Efficacy and Safety of a Therapeutic Strategy of PPH Comparing Early Administration of Human Fibrinogen vs Placebo in Patients Treated With IV Prostaglandins Following Vaginal Delivery. ClinicalTrials.gov Identifier: NCT02155725 (accessed 05.08.2018).

* Ducloy-Bouthors A, Mignon A, Huissoud C, Grouin J, Mercier FJ. Fibrinogen concentrate as a treatment for postpartum haemorrhage-induced coagulopathy: A study protocol for a randomised multicentre controlled trial. The fibrinogen in haemorrhage of DELivery (FIDEL) trial. *Anaesthesia Critical Care and Pain Medicine* 2016;**35**:293-8.

NCT02203968 {published data only}

Fibrinogen in the Initial Resuscitation of Severe Trauma (FiIRST). ClinicalTrials.gov Identifier: NCT02203968 (accessed 13 January 2016).

NCT02253082 {published data only}

Vasculopathic Injury and Plasma as Endothelial Rescue - OCTAplus Trial (VIPER-OCTA). ClinicalTrials.gov Identifier: NCT02253082 (accessed 13 January 2016).

NCT02299947 {published data only}

Bloodtransfusions After Aortic Surgery (FIBTEG). ClinicalTrials.gov Identifier: NCT02299947 (accessed 13 January 2016).

NCT02344069 {published data only}

Effect of Immediate, Pre-emptive Fibrinogen Concentrate in Patients With Trauma Haemorrhage Needing Haemostatic Resuscitation - a Randomized, Controlled, Double-blinded Investigator-initiated Pilot Trial. ClinicalTrials.gov Identifier: NCT02344069 (accessed 05.08.2018).

* Stensballe J, Sorensen AM, Steinmetz J, Henriksen HH, Lange T, Larsen CF, et al. Fibrinogen in trauma haemorrhage. *Transfusion medicine*. 2017; Vol. 27:19.

NCT02429453 {published data only}

FFP Versus PCC in Traumatic Intracranial Hemorrhage. ClinicalTrials.gov Identifier: NCT02429453 (accessed 13 January 2016).

NCT02528708 {published data only}

A Program to Evaluate Riastap® and FIBTEM® for the Early Control and Treatment of Postpartum Hemorrhage (PERFECT PPH). ClinicalTrials.gov Identifier: NCT02528708 (accessed 05.08.2018) currently suspended pending review to commence.

NCT02557672 {published data only}

Prothrombin Complex Concentrate (PCC) Compared to Fresh Frozen Plasma (FFP) for Post-cardiopulmonary Bypass Coagulopathy and Bleeding, a Prospective Randomized Trial at Large US Medical Center. ClinicalTrials.gov Identifier: NCT02557672 (accessed 05.02.2018).

NCT02745041 {published data only}

Fibrinogen Concentrate vs Cryoprecipitate in Traumatic Haemorrhage: A Pilot Randomised Controlled Trial. ClinicalTrials.gov Identifier: NCT02745041 (accessed 05.08.2018).

* Winearls J, Wulschleger M, Wake E, Hurn C, Furry J, Ryan G, et al. Fibrinogen Early In Severe Trauma study (FEISTY): study protocol for a randomised controlled trial. *Trials* 2017;**18**:241.

NCT02777424 {published data only}

Prothrombin Complex Concentrate Versus Fresh Frozen Plasma to Correct Coagulation Disorders in Adult Neurosurgical Patients. ClinicalTrials.gov Identifier: NCT02777424 (accessed 05.08.2018).

NCT02822599 {published data only}

The Role of Human Fibrinogen Concentrate (RiaSTAP) in Decreasing Blood Loss and the Need for Component Blood Therapy in Infants Undergoing Cardiopulmonary Bypass. ClinicalTrials.gov Identifier: NCT02822599 (accessed 05.02.2018).

NCT03014700 {published data only}

Repurposing of Fibrinogen Concentrate as a Cost-Effective and Safe Hemostatic Agent in Infants Undergoing Cardiac Surgery on Cardiopulmonary Bypass. ClinicalTrials.gov Identifier: NCT03014700 (accessed 05.02.2018).

NCT03037424 {published data only}

Prospective, Multi-center, Randomized, Active-control, Non-inferiority Study Comparing Fibrinogen Concentrate With Cryoprecipitate for the Treatment of Acquired Hypofibrinogenemia in Bleeding Adult Cardiac Surgical Patients. ClinicalTrials.gov Identifier: NCT03037424 (accessed 05.08.2018).

NCT03183479 {published data only}

The Effects of Fibrinogen Concentrate Infusion on Perioperative Blood Loss and Allogeneic Blood Conservation in Patients Undergoing Scoliosis Surgery. ClinicalTrials.gov Identifier: NCT03183479 (accessed 05.02.2018).

NCT03218722 {published data only}

Early Administration of Prothrombin Concentrate Complex in Patients With Acute Hemorrhage Following Severe Trauma (PROCOAG). ClinicalTrials.gov Identifier: NCT03218722 (accessed 20.4.18).

Additional references

Brenner 2018

Brenner A, Arribas M, Cuzick J, Jairath V, Stanworth S, Ker K, et al. Outcome measures in clinical trials of treatments for acute severe haemorrhage. *Trials* 2018;**19**(1):533-9.

Carless 2009

Carless PA, Henry DA, Anthony DM. Fibrin sealant use for minimising peri-operative allogeneic blood transfusion. *Cochrane Database of Systematic Reviews* 2009, Issue 3. [DOI: [10.1002/14651858.CD004171](https://doi.org/10.1002/14651858.CD004171); PUBMED: 12804501]

Chan 2013

Chan AW, Tetzlaff JM, Gøtzsche PC, Altman DG, Mann H, Berlin JA, et al. SPIRIT 2013 explanation and elaboration: guidance for protocols of clinical trials. *BMJ* 2013;**8**(346):e7586.

Charbit 2007

Charbit B, Mandelbrot L, Samain E, Baron G, Haddaoui B, Keita H, et al. The decrease of fibrinogen is an early predictor of the severity of postpartum hemorrhage. *Journal of Thrombosis and Haemostasis* 2007;**5**(2):266-73. [PUBMED: 17087729]

CRASH-2 2010

CRASH-2 Trial Collaborators. Effects of tranexamic acid on death, vascular occlusive events, and blood transfusion in trauma patients with significant haemorrhage (CRASH-2): a randomised, placebo-controlled trial. *Lancet* 2010;**376**:23-32. [DOI: [10.1016/S0140-6736\(10\)60835-5](https://doi.org/10.1016/S0140-6736(10)60835-5)]

de Lloyd 2011

de Lloyd L, Bovington R, Kaye A, Collis RE, Rayment R, Sanders J, et al. Standard haemostatic tests following major obstetric haemorrhage. *International Journal of Obstetric Anaesthesia* 2011;**20**(2):135-41. [PUBMED: 21439811]

Dentali 2011

Dentali F, Marchesi C, Pierfranceschi MG, Crowther M, Garcia D, Hylek E, et al. Safety of prothrombin complex concentrates for rapid anticoagulation reversal of vitamin K antagonists: a meta-analysis. *Thrombosis and Haemostasis* 2011;**106**(3):429-38.

Franchini 2012

Franchini M, Lippi G. Fibrinogen replacement therapy: a critical review of the literature. *Blood Transfusion* 2012;**10**(1):23-7.

Gargon 2017

Gargon E, Williamson PR, Young B. Improving core outcome set development: qualitative interviews with developers provided pointers to inform guidance. *Journal of Clinical Epidemiology* 2017;**86**:140-52.

Grottke 2011

Grottke O, Braunschweig T, Spronk HM, Esch S, Rieg AD, van Oerle R, et al. Increasing concentrations of prothrombin complex concentrate induce disseminated intravascular coagulation in a pig model of coagulopathy with blunt liver injury. *Blood* 2011;**118**(7):1943-51. [PUBMED: 21670472]

Görlinger 2011

Görlinger K, Dirkmann D, Hanke AA, Kamler M, Kottenberg E, Thielmann M, et al. First-line therapy with coagulation factor concentrates combined with point-of-care coagulation testing is associated with decreased allogeneic blood transfusion in cardiovascular surgery: a retrospective, single-center cohort study. *Anesthesiology* 2011;**115**(6):1179-91.

Haas 2012

Haas T, Korte W, Spielmann N. Perioperative course of FXIII in children undergoing major surgery. *Paediatric Anaesthesiology* 2012;**22**:641-6.

Higgins 2011

Higgins JPT, Altman DG, Sterne JAC (editors). Chapter 8: Assessing risk of bias in included studies. Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). Available from www.cochrane-handbook.org. The Cochrane Collaboration, 2011.

Johansen 2015

Johansen M, Wikkeso A, Lunde J, Wetterslev J, Afshari A. Prothrombin complex concentrate for reversal of vitamin K antagonist treatment in bleeding and non-bleeding patients. *Cochrane Database of Systematic Reviews* 2015, Issue 7. [DOI: [10.1002/14651858.CD010555](https://doi.org/10.1002/14651858.CD010555)]

Kearney 1989

Kearney RA, Rosales JK, Howes WJ. Craniosynostosis: an assessment of blood loss and transfusion practices. *Canadian Journal of Anaesthesia* 1989;**36**:473-7.

Kozek-Langenecker 2013

Kozek-Langenecker SA, Afshari A, Albaladejo P. Management of severe perioperative bleeding guidelines from the European Society of Anaesthesiology. *European Journal of Anaesthesiology* 2013;**30**:270-382.

Levi 2010

Levi M, Levy JH, Andersen HF, Truloff D. Safety of recombinant activated factor VII in randomized clinical trials. *New England Journal of Medicine* 2010;**363**(19):1791-800. [PUBMED: 21047223]

Mengoli 2017

Mengoli C, Franchini M, Marano G, Pupella S, Vaglio S, Marietta M, et al. The use of fibrinogen concentrate for the management of trauma-related bleeding: a systematic review and meta-analysis. *Blood Transfusion* 2017;**15**(4):318-24.

Moher 2010

Moher D, Hopewell S, Schulz KF, Montori V, Gotzsche PC, Devereaux PJ, et al. CONSORT 2010 Explanation and

Elaboration: updated guidelines for reporting parallel group randomised trials. *BMJ* 2010;**340**:c869.

Review Manager 2014 [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Roberts 2014

Roberts I, Coats T, Edwards P, Gilmore I, Jairath V, Ker K, et al. HALT-IT - tranexamic acid for the treatment of gastrointestinal bleeding: study protocol for a randomised controlled trial. *Trials* 2014;**15**:450-63.

Schulz 1995

Schulz KF, Chalmers I, Hayes RJ, Altman DG. Empirical evidence of bias. Dimensions of methodological quality associated with estimates of treatment effects in controlled trials. *JAMA* 1995;**273**(5):408-12.

Schünemann 2011

Schünemann HJ, Oxman AD, Vist GE, Higgins JPT, Deeks JJ, Glasziou P, et al. Chapter 12: Interpreting results and drawing conclusions. In: Higgins JPT, Green S editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011.

SHOT 2011

Serious Hazards of Transfusion (SHOT). Annual SHOT Report 2011. <http://www.shotuk.org/shot-reports/shot-annual-report-summary-2011> (accessed 1/4/2013).

Simpson 2012

Simpson E, Lin Y, Stanworth S, Birchall J, Doree C, Hyde C. Recombinant factor VIIa for the prevention and treatment of bleeding in patients without haemophilia. *Cochrane Database of Systematic Reviews* 2012, Issue 3. [DOI: [10.1002/14651858.CD005011.pub4](https://doi.org/10.1002/14651858.CD005011.pub4); PUBMED: 21328270]

Stanworth 2010

Stanworth SJ, Morris TP, Gaarder C, Goslings JC, Maegele M, Cohen MJ, et al. Reappraising the concept of massive transfusion in trauma. *Critical Care (London, England)* 2010;**14**(6):R239. [PUBMED: 21192812]

Stanworth 2011

Stanworth SJ, Grant-Casey J, Lowe D, Laffan M, New H, Murphy MF, et al. The use of fresh-frozen plasma in England:

high levels of inappropriate use in adults and children. *Transfusion* 2011;**51**(1):62-70. [PUBMED: 20804532]

Tripodi 2009

Tripodi A, Chantarangkul V, Mannucci PM. Acquired coagulation disorders: revisited using global coagulation/anticoagulation testing. *British Journal of Haematology* 2009;**147**(1):77-82. [PUBMED: 19659548]

Volis 2012

Voils SA, Baird B. Systematic review: 3-factor versus 4-factor prothrombin complex concentrate for warfarin reversal: does it matter?. *Thrombosis Research* 2012;**130**(6):833-40.

Walsh 2009

Walsh CE, Valentino LA. Factor VIII prophylaxis for adult patients with severe haemophilia A: results of a US survey of attitudes and practices. *Haemophilia* 2009;**15**(5):1014-21. [PUBMED: 19493018]

Wikkelsø 2013

Wikkelsø A, Lunde J, Johansen M, Stensballe J, Wetterslev J, Møller AM, et al. Fibrinogen concentrate in bleeding patients. *Cochrane Database of Systematic Reviews* 2013, Issue 8. [DOI: [10.1002/14651858.CD008864](https://doi.org/10.1002/14651858.CD008864)]

WOMAN 2017

WOMAN Trial Collaborators. Effect of early tranexamic acid administration on mortality, hysterectomy, and other morbidities in women with post-partum haemorrhage (WOMAN): an international, randomised, double-blind, placebo-controlled trial. *Lancet* 2017;**389**:10084:2105-16. [DOI: [10.1016/S0140-6736\(17\)30638-4](https://doi.org/10.1016/S0140-6736(17)30638-4)]

Yang 2012

Yang L, Stanworth S, Hopewell S, Doree C, Murphy M. Is fresh-frozen plasma clinically effective? An update of a systematic review of randomized controlled trials. *Transfusion* 2012;**52**(8):1673-86. [PUBMED: 22257164]

Zeng 2017

Zeng L, Choonara I, Zhang L, Li Y, Shi J. Effectiveness of prothrombin complex concentrate (PCC) in neonates and infants with bleeding or risk of bleeding: a systematic review and meta-analysis. *European Journal of Pediatrics* 2017;**176**(5):581-9.

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Bilecen 2017

Methods

Randomised controlled, single-centre trial.

Randomly assigned at: At the end of cardiopulmonary bypass, when operative haemostasis had been achieved an assessment was made after 5 minutes if a patient had an activated clotting time < 140secs; temp > 36 celsius; pH > 7.3; Hb > 8.5 or Hct > 25%. Gauzes were weighed and if less than 60 mL this was defined as a dry surgical field; if > 250 mL surgical bleeding was defined and further surgery and

Bilecen 2017 (Continued)

haemostasis was required. If the swabs contained 60 - 250 mL blood then 'intraoperative bleeding' was defined and these patients received fibrinogen concentrate or placebo.

Calculated sample size: Power analysis

Participants	<p>Condition: Patients undergoing elective high-risk cardiac surgery.</p> <p>Inclusion criteria: Older than 18 years and had undergone elective high-risk cardiac surgery, defined as combined coronary artery bypass graft surgery and valve repair or replacement surgery, the replacement of multiple valves, aortic root reconstruction, or reconstruction of the ascending aorta or aortic arch.</p> <p>Exclusion criteria: Proof or suspicion of a congenital or acquired coagulation disorder, any stroke or myocardial infarction within 2 months preceding surgery, venous or arterial thrombosis, clopidogrel use in the 5 days preceding surgery, glycoprotein IIb/IIIa receptor antagonist use in the 2 days preceding surgery, or INR > 1.4 in patients who used anti-coagulants</p> <p>Number of participants enrolled: 203</p> <p>Number of participants randomly assigned to the intervention/placebo: 60/60</p> <p>Number of participants who received the treatment/placebo and were included in the outcome analyses: 60/60</p>
Interventions	<p>Intervention: Fibrinogen concentrate (dose depended on fibrinogen level at end of surgery and calculated according to participant's weight)</p> <p>Comparator: Placebo (albumin in 0.9% sodium chloride solution)</p> <p>Additional treatments received whilst in the trial: Antifibrinolytic prophylaxis, post-operative tranexamic acid, post-operative PCC. Transfusion based on standard operating procedure for hospital intensive care unit.: if Hb < 8.5g/dL, 1 unit red blood cells were administered; if a participant lost more than 1 litre of blood, 2 pools FFP were administered; if a participant lost 2+ litres of blood, 4 FFP units were administered; if a participant had a platelet count < 80 × 10⁹/L, 1 unit of platelet concentrate (5 donors) was administered</p> <p>Following on from the ICU protocol, participants experiencing a blood loss > 400 mL in 1 hour or exceeding 300 mL per hour for 2 hours or exceeding 200 mL per hour for 3 hours while in the ICU received 4 units of red blood cells, 4 units FFP, and 1 unit of platelet concentrate (5 donors) via transfusion, and the attending surgeon was informed. Participants whose platelet counts were lower than 75 × 10⁹/L received 1 unit of platelet concentrate (5 donors) via transfusion</p>
Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> Intra-operative blood loss measured between intervention (i.e. infusion of study medication after completion of cardiopulmonary bypass) and closure of the chest when the surgery ended <p>Secondary:</p> <ul style="list-style-type: none"> Measured blood loss 1, 3, 6, 12, and 24 hours after the intervention The proportion of participants who had received a transfusion The number of units of RBC, FFP, platelet concentrate, and any blood product units used between when the intervention started and when the chest was closed and for the 24-hour period after the intervention The amount of procoagulants and antifibrinolytics (tranexamic acid, desmopressin, PCC, recombinant factor VIIa and fibrinogen concentrate) given during the perioperative and post-operative periods Adverse events: 30-day mortality, myocardial infarction, stroke, transient Ischaemic attack, renal failure, venous thromboembolism, allergic reactions to fibrinogen concentrate, wound infections, reoperative thoracotomy <p>Duration of follow-up: Until chest closure for primary outcome; up to 30 days for other outcomes.</p>

Bilecen 2017 (Continued)

Notes

Population nationality: Dutch

Correspondence: Dr. Süleyman Bilecen (s.bilecen@umcutrecht.nl).

Clinical trial registration number: [Clinicaltrials.gov](https://clinicaltrials.gov) Identifier: NCT01124981 and EudraCT No: 2009-018086-12

Sources of funding: CSL Behring sponsored this study and donated the bottles of study medication

Dates the trial was conducted: February 2011 to December 2014

Declarations of interest (among primary researchers): All authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest and none were reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Web-based randomisation protocol using an unstratified fixed block size of 4. The order of blocks was also randomised
Allocation concealment (selection bias)	Low risk	Web-based randomisation protocol
Blinding of participants (performance bias)	Low risk	Under general anaesthesia
Blinding of study personnel (performance bias)	Low risk	Syringes of intervention visibly undifferentiable. Trial interventions made up outside the operating theatre, study drug made up in a room where only a certain number of research unblinded personnel could go, in amber colour syringes. Fibrinogen levels not released for first 24 hours
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Syringes of intervention visibly undifferentiable. Trial interventions made up outside the operating theatre, study drug made up in a room where only a certain number of research unblinded personnel could go, in amber colour syringes. Fibrinogen levels not released for first 24 hours
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Data forms missing from participants in both study arms. Unlikely to have an impact on outcome and missing data balanced across study groups.
Selective reporting (reporting bias)	Low risk	All prespecified outcomes were reported in the results. Study protocol available
Other bias	Low risk	One participant given post-operative fibrinogen, however this does not represent a protocol violation

Bregenzer 1999

Methods

Randomised controlled, multicentre trial

Randomly assigned at: Not stated

Calculated sample size: Power analysis

Participants

Condition: ulcerative colitis and macroscopically visible severe intestinal bleeding

Bregenzner 1999 (Continued)

Inclusion criteria: Confirmed flare of ulcerative colitis not responding to steroids for at least 21 days, clinical activity index ≥ 8

Exclusion criteria: Use of antithrombotic drugs affecting platelet function, coumarin derivatives or therapeutic heparinisation

Number of participants randomly assigned to the intervention: 17/11

Number of participants included in the outcome analyses: 11/9

Interventions	<p>Intervention: Continuation of basic steroid treatment with addition of factor XIII for iv injection (at 3750 units on Day 0 and 1250 units on days 1 through 9)</p> <p>Comparator: Continuation of basic steroid treatment with placebo (freeze-dried product for iv injection of the same composition as the intervention, but not enriched with FXIII)</p> <p>Additional treatments received whilst in the trial:</p> <p>Basic steroid treatment for the treatment of ulcerative colitis</p> <p>Number of participants enrolled: Planned for 75 but the trial was prematurely stopped due to observation of no difference between the treatment arms at the first interim analysis.</p>
Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> Time to cessation of macroscopically visible intestinal bleeding within 14 days of starting treatment <p>Secondary:</p> <ul style="list-style-type: none"> Clinical scores of ulcerative colitis severity (time points not stated) Endoscopic/histological findings (time points not stated) Factor XIIIa and XIIIb levels (time points not stated) <p>Duration of follow-up: Up to day 42</p>
Notes	<p>Population nationality: German</p> <p>Correspondence: Dr N Bregenzner, Department of Internal Medicine, University of Regensburg, D93042 Regensburg, Germany</p> <p>Clinical trial registration number: not stated</p> <p>Sources of funding: trial supported by Centeon Pharma GmbH</p> <p>Dates the trial was conducted: October 1994 to January 1997</p> <p>Declarations of interest (among primary researchers): not stated</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information was reported from which to make an assessment of this bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information was reported from which to make an assessment of this bias
Blinding of participants (performance bias)	Low risk	The study reports that "the blinding of patients and the investigators was carried out by labelling", and that the placebo was a "freeze dried product for I.V. injection of the same composition as the FXIII concentrate", so it is unlikely that participants would be able to guess their treatment allocation

Bregenz 1999 (Continued)

Blinding of study personnel (performance bias)	Low risk	The study reports that "the blinding of patients and the investigators was carried out by labelling", and that the placebo was a "freeze dried product for I.V. injection of the same composition as the FXIII concentrate"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	<p>Primary outcome: assessed daily by the participants using a study checklist</p> <p>Secondary outcomes: clinically assessed, although unclear by whom</p> <p>All outcomes were combined to calculate FXIIa and FXIIb levels in each participant. The study reports that the "FXIII analyses were performed by an independent reference laboratory and results were not made known to the treating physicians"</p> <p>Overall judged to be at low risk of detection bias.</p>
Incomplete outcome data (attrition bias) All outcomes	High risk	<p>Quote: "The first planned blinded interim analysis of the primary and secondary efficacy criteria for 28 patients ... showed no differences between groups... the trial was prematurely stopped".</p> <p>This analysis was intention-to-treat (with assumptions made for participants not available for the analysis (N = 8)*). The planned per protocol analysis was not possible due to the high number of participants that had to be excluded from the per protocol analysis (N = 16); a large number (not defined) due to major protocol violations.</p> <p>*The reason for the unavailability of data for analyses was aggravation of ulcerative colitis (N = 2 in placebo group; N = 4 in the factor XIII group); loss to follow-up (N = 1 in the factor XIII group); withdrawal of consent (N = 1 in the factor XIII group).</p>
Selective reporting (reporting bias)	High risk	No trial protocol is available and outcome information was not reported for a number of prespecified outcomes, including haemoglobin and haematocrit levels
Other bias	Unclear risk	Insufficient information was reported from which to make an assessment of this bias

Collins 2017

Methods	<p>Multicentre, randomised, double-blind, placebo-controlled trial set in teaching hospital obstetric units</p> <p>Randomly assigned at: Following delivery of the baby and with ongoing bleeding</p> <p>Calculated sample size: Power analysis</p>
Participants	<p>Condition: Postpartum haemorrhage</p> <p>Inclusion criteria: Women aged 18 years and above and ≥ 24 weeks gestation who had ongoing major postpartum haemorrhage (1000 – 1500 mL blood loss)</p> <p>Exclusion criteria: Declined transfusion, placenta accreta diagnosed antenatally, undergone an invasive procedure to control bleeding or if there was clinical suspicion of amniotic fluid embolus. Detailed inclusion and exclusion criteria have been published elsewhere</p> <p>Number of participants enrolled: 57</p> <p>Number of participants randomly assigned to the intervention/control: 29/28</p>

Collins 2017 (Continued)

Number of participants who received the treatment/control and were included in the outcome analyses: 28/27

Interventions	<p>Intervention: Fibrinogen concentrate (variable dose with aim to increase FIBTEM A5 to > 22 mm in the fibrinogen arm)</p> <p>Comparator: 0.9% sodium chloride solution</p> <p>Additional treatments received whilst in the trial: At randomisation 1 g of tranexamic acid was infused if not already given and 4 units of FFP ordered that were given if bleeding was ongoing. Other blood and blood products as per local standard operating procedures</p>
Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> The number of allogeneic blood products (RBC, FFP, cryoprecipitate, platelets) infused after study medication until hospital discharge <p>Secondary:</p> <ul style="list-style-type: none"> Measured blood loss Numbers of units of RBCs, FFP, cryoprecipitate, and platelet transfusion within the first 24 hours Cell salvage volume Change in FIBTEM and Clauss fibrinogen Invasive procedures Initiation of breastfeeding Safety endpoints were thrombosis to 6 weeks, level 2 and 3 admission and length of hospital stay <p>Duration of follow-up: Primary outcome until hospital discharge, other outcomes until 6 weeks post-intervention</p>
Notes	<p>Population nationality: British</p> <p>Correspondence: peter.collins@wales.nhs.uk</p> <p>Clinical trial registration number: ISRCTN46295339</p> <p>Sources of funding: Supported by an investigator-led grant from CSL Behring. The trial's sponsor and funder had no role in study design; collection, analysis and interpretation of data; writing the report; or the decision to submit the report for publication. The trial funder reviewed the manuscript before submission and no changes were made to the manuscript following this review</p> <p>Dates the trial was conducted: 29 June 2013 and 26 November 2015</p> <p>Declarations of interest (among primary researchers): P.W.C. and R.E.C. research support from TEM International and Haemonectics. Funding for lectures from CSL Behring. A.D.W. is the inventor of the PPH butterfly, a device to facilitate uterine compression for the treatment of postpartum haemorrhage. The patent is held by his employer, the University of Liverpool, but A.D.W. would receive royalty payments for any sales. All other authors: no interest declared</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated random numbers
Allocation concealment (selection bias)	Low risk	Consecutive blinded numbered study packs

Collins 2017 (Continued)

Blinding of participants (performance bias)	Low risk	Study medication syringes were blacked out to prevent identification
Blinding of study personnel (performance bias)	Low risk	Blinding by identical intervention boxes, black syringes and technique for drawing up the trial interventions
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Blinded to allocation, although potential for outcome assessors to access fibrinogen levels and potentially determine allocation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Complete outcome data available. 1 participant dropped out in each study arm, with reasons documented
Selective reporting (reporting bias)	Low risk	Study protocol available
Other bias	Low risk	Industrial support for study but well detailed involvement and low influence

Cui 2010

Methods	Randomised controlled trial, single centre Randomly assigned at: Not stated Calculated sample size: Not stated
Participants	Condition: Paediatric cardiac surgery Inclusion criteria: Haematocrit > 54% before operation; the diagnosis was transposition of the great arteries or double outlet right ventricle, which was given by 1 designated ultrasound doctor; the operation that the patients underwent was arterial switch operation or double roots transplantation Exclusion criteria: history of blood disease; anticoagulation treatment before surgery; medication that affects haemostasis (such as prostaglandin E1); difficult sternal closure because of anatomic or surgical reasons. Number of participants enrolled: 40 Number of participants randomly assigned to the intervention: 20/20 Number of participants who received the treatment and were included in the outcome analyses: 17/14
Interventions	Intervention: Fibrinogen administration (0.5 - 1 g) combined with traditional transfusion, guided by TEG (no further details provided) Comparator: Traditional transfusion guided by clinical experience Additional treatments received whilst in the trial: None stated
Outcomes	Primary: <ul style="list-style-type: none">Time to chest wall closure Secondary: <ul style="list-style-type: none">Total FFP volume used in operation and in ICU during the first 24 hoursTotal RBC units used in operation and in ICU during the first 24 hoursTotal platelet units used at closure time and in ICU during the first 24 hours

Cui 2010 (Continued)

- Amount of chest drainage at 1, 6 and 24 hours
- Mechanical ventilator time
- ICU stay
- Hospitalisation time

Duration of follow-up: first 24 hours in the ICU

Notes

Population nationality: Chinese

Correspondence: jinpingfw@hotmail.com

Clinical trial registration number: Not stated

Sources of funding: Not stated

Dates the trial was conducted: September 2007 to April 2009

Declarations of interest (among primary researchers): Not stated

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information presented to make an assessment of this bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information presented to make an assessment of this bias
Blinding of participants (performance bias)	Low risk	No information is reported on whether blinding was undertaken in this trial, but participants were under general anaesthesia when the intervention was administered and therefore bias due to knowledge of allocation concealment is not relevant
Blinding of study personnel (performance bias)	High risk	No information is reported on whether blinding was undertaken in this trial, but as some of the outcomes measured in this trial (e.g. ventilation time, time to discharge and FFP usage) are open to bias if treatment allocation were known, then we rate this at high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Low risk	No information is reported on whether blinding was undertaken in this trial, but as the outcomes were objective and measured during the hospital stay we felt this domain was not subject to performance bias, and hence rated it at low risk of bias
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Following randomisation 3 participants in the intervention group and 6 in the comparator group were excluded from the trial for the following reasons: history of blood disease, anticoagulation treatment before surgery, medication that affects haemostasis, difficult sternal closure for anatomical or surgical reasons (reasons not attributed to intervention group or the timing of exclusion in relation to receipt or not of intervention). It is also unclear whether the number of participants included in the outcome analyses were those randomised or those available for outcome analysis: we have assumed the latter for the purposes of this review
Selective reporting (reporting bias)	Low risk	All outcomes outlined in the Methods have been reported on in the Results of the study
Other bias	Unclear risk	Case-mix likely to be quite heterogeneous and ages quite different (although not significantly so, but value of the standard deviation suggests a broad spread of ages in each intervention group). Additionally participants appear

Cui 2010 (Continued)

to have been randomised prior to application of exclusion criteria with more patients excluded from the intervention arm of the study (although the allocation and exclusion numbers are not stated)

Curry 2018

Methods	<p>Multi-centre, double-blind, placebo-controlled trial, conducted in 5 UK major trauma centres</p> <p>Randomly assigned at: On arrival in A&E department</p> <p>Calculated sample size: Power analysis</p>
Participants	<p>Condition: Acute trauma</p> <p>Inclusion criteria: Adults (judged to be 16 years or older); were actively bleeding and in haemorrhagic shock; and required activation of the major haemorrhage protocol or had already received a transfusion of emergency red blood cells</p> <p>Exclusion criteria: Patient transferred from another hospital; the trauma team leader deemed the injury incompatible with life; more than 3 hours had elapsed from time of injury (taken as time of 999 call, if unknown); pregnant women and severe isolated or unsalvageable head injury</p> <p>Number of participants randomly assigned to the intervention/control: 24/24; 9 patients subsequently did not receive the study intervention as excluded as per protocol; final allocation 20/19</p> <p>Number of participants who received the treatment/control and were included in the outcome analyses: 20/19</p>
Interventions	<p>Intervention: Fibrinogen concentrate, 6 g</p> <p>Comparator: 0.9% sodium chloride solution</p> <p>Additional treatments received whilst in the trial: Major haemorrhage protocol therapy, as per hospital protocol</p> <p>Number of participants enrolled: 48.</p>
Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> Feasibility of delivering fibrinogen therapy within 45 minutes to adult trauma patients Proportion of participants whose fibrinogen levels were maintained $\geq 2\text{g/L}$ during active haemorrhage <p>Secondary:</p> <ul style="list-style-type: none"> Mortality at 3, 6 and 24 hours, and 28 days from admission Transfusion requirements, in numbers of units, at 3, 6 and 24 hours Duration of organ support In-patient stay, including ICU/HDU Quality of life A haemorrhage assessment tool, completed at 3 hours from admission, was piloted in this study, aiming to determine whether this tool may provide a useful clinical measure of haemostasis Safety was measured by symptomatic thrombotic events; arterial (e.g. myocardial infarction, stroke) and venous (e.g. pulmonary embolism, deep venous thrombosis) during hospital stay Laboratory measures included Clauss fibrinogen at 2 hours after admission and at day 7 from admission <p>Duration of follow-up: 28 days post-admission $\pm$ 4 days</p>
Notes	<p>Population nationality: British.</p>

Curry 2018 (Continued)

Correspondence: nicola.curry@ouh.nhs.uk

Clinical trial registration number: ISRCTN67540073

Sources of funding: CSLBehring - financial support only and had no role in initiation of the trial, its methods or process and no control over data or publication

Dates the trial was conducted: December 2015 to November 2016

Declarations of interest (among primary researchers): Not stated

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Centralised computerised randomisation system
Allocation concealment (selection bias)	Low risk	Sequentially-numbered allocation boxes
Blinding of participants (performance bias)	Low risk	Enrolled by waiver, generally intubated and hence unaware of allocation
Blinding of study personnel (performance bias)	Low risk	Blinding method previously validated in another RCT and published; blinded intervention containers, prevention of study personnel seeing intervention reconstitution and blacked-out syringes
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Blacked-out syringes were used, so outcome assessors were blinded to treatment allocation. However, outcome assessors would have had access to the blood results which may identify whether fibrinogen concentrate was given, hence we rate the risk being as unclear
Incomplete outcome data (attrition bias) All outcomes	Low risk	Outcome data complete
Selective reporting (reporting bias)	Unclear risk	Protocol outlined use of ROTEM data but no ROTEM data reported
Other bias	Unclear risk	The funder, CSL Behring, provided financial support only and had no role in initiation of the trial, its methods or process and no control over data or publication.

Fenger Erikson 2009

Methods	<p>Randomised controlled, single-centre trial</p> <p>Randomly assigned at: a dilution level (blood loss having been substituted 1:1 with HES 130/0.4) of 30%</p> <p>Calculated sample size: Power calculation</p>
Participants	<p>Condition: Radical cystectomy</p> <p>Inclusion criteria: Older than 17 years of age, undergoing radical cystectomy for suffering localised bladder cancer (T₁₋₂N₀M₀)</p>

Fenger Erikson 2009 (Continued)

Exclusion criteria: Coagulation disorders, defined as abnormal values of platelet count, prothrombin time, activated partial thromboplastin time, fibrinogen, antithrombin, or D-dimer; treatment with oral vitamin K antagonists; intake of non-steroid anti-inflammatory drugs within 2 days prior to surgery; renal or hepatic dysfunction; ischaemic heart disease; pregnancy; and known hypersensitivity to hydroxyethyl starch

Number of participants enrolled: 27

Number of participants randomly assigned to the intervention: 10/11

Number of participants included in the outcome analyses: 10/10

Interventions	<p>Intervention: Fibrinogen concentrate at 45 mg/kg, administered intra-operatively</p> <p>Comparator: Intra-operative placebo (placebo = isotonic saline, administered intra-operatively at a dose of 2.25 mL/kg)</p> <p>Additional treatments received whilst in the trial: Ringer's lactate solution (4 mL/kg/hour) and hydroxyethyl starch 130/0.4</p>
Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> Whole blood maximum clot firmness as determined by thromboelastometry <p>Secondary:</p> <ul style="list-style-type: none"> Other thromboelastometric variables Platelet function Thrombin generation Bleeding and requirement for peri- and post-operative blood product transfusion <p>Duration of follow-up: up to 48 hours post-operatively</p>
Notes	<p>Population nationality: Danish</p> <p>Correspondence: b.sorensen@kcl.ac.uk</p> <p>Clinical trial registration number: Eudra CT number 2007-00026-44</p> <p>Sources of funding: Unrestricted research grant from CSL Behring, the University of Aarhus Research Foundation and the A. P. Møller and Hustru Chastine Mc-Kinney Møllers Foundation</p> <p>Dates the trial was conducted: June 2007 to March 2008</p> <p>Declarations of interest (among primary researchers): No stated conflicts of interest</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information was reported from which to make an assessment of this bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information was reported from which to make an assessment of this bias
Blinding of participants (performance bias)	Low risk	Participants under general anaesthesia when the intervention was administered
Blinding of study personnel (performance bias)	Low risk	The study reports that the "trial medication was prepared and administered by a second person thus the trial staff were blinded to treatment" and that

Fenger Erikson 2009 (Continued)

		"allocation concealment was sustained until all recruitment, data collection, transfusion requirement data and laboratory analyses were complete":
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The study reports that the "trial medication was prepared and administered by a second person thus the trial staff were blinded to treatment" and that "allocation concealment was sustained until all recruitment, data collection, transfusion requirement data and laboratory analyses were complete":
Incomplete outcome data (attrition bias) All outcomes	Low risk	1 participant randomised to placebo did not receive placebo due to an interruption in their operation: this participant was not included in the final outcome analyses. All other participants and their outcome data were reported by the study
Selective reporting (reporting bias)	Low risk	All prespecified outcomes were reported in the Results
Other bias	Low risk	The study appears to be free of other sources of bias

Galas 2014

Methods	Randomised controlled, single-centre trial Randomly assigned at: post-surgery and after heparin neutralisation provided patients met 2 inclusion criteria: diffuse bleeding from capillary beds at wound surfaces requiring haemostatic therapy and plasma fibrinogen < 1 g/dl Calculated sample size: None stated
Participants	Condition: Paediatric cardiac surgery Inclusion criteria: Age under 15 years, undergoing cardiac surgery cardiopulmonary bypass, intra-operative bleeding and hypofibrinogenaemia Exclusion criteria: Previous coagulopathy (clinical history or INR > 1.5), low platelet count (< 100,000), product allergy, urgent procedures, active infection Number of participants enrolled: 688 patients were screened: 63 were randomised (274 were either not bleeding or did not meet laboratory target for inclusion; 351 met an exclusion criterion) Number of participants randomly assigned to the intervention: 30/33 Number of participants included in the outcome analyses: 30/33
Interventions	Single dose of either intervention given at time of intra-operative bleeding Intervention: Fibrinogen concentrate (60 mg/kg body weight, Haemocomplettan®) Comparator: Cryoprecipitate (10 mL/kg body weight). Additional treatments received whilst in the trial: None stated
Outcomes	Primary: <ul style="list-style-type: none">Post-operative blood losses during 48 hours after surgery Secondary: <ul style="list-style-type: none">Percentage of participants exposed to allogenic blood products (RBCs, FFP, platelet concentrate and cryoprecipitate) during the intraoperative periodDuration of mechanical ventilation during ICU stayVasopressor requirement during ICU stay

Galas 2014 (Continued)

- Incidence of acute myocardial infarction, stroke, acute kidney injury requiring dialysis, septic shock, re-operation, peripheral artery occlusion, deep vein thrombosis, pulmonary embolism up to hospital discharge
- Death up to post-operative day 7 or hospital discharge
- Length of ICU stay;
- Length of hospital stay

Duration of follow-up: Up to post-operative day 7 or hospital discharge

Notes

Population nationality: Brazilian
Correspondence: ludhmila@usp.br

Clinical trial registration number: NCT01187225

Sources of funding: CSL Behring (Sao Paulo, Brazil) provided the trial drug and the testing of clotting factors and thromboelastometry; University of Sao Paulo, Brazil

Dates the trial was conducted: August 2010 to November 2011

Declarations of interest (among primary researchers): "Nothing to disclose with regard to commercial support"

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random-number table
Allocation concealment (selection bias)	Low risk	Opaque envelopes opened sequentially
Blinding of participants (performance bias)	Low risk	Participants were under anaesthesia when the intervention was administered
Blinding of study personnel (performance bias)	Unclear risk	Insufficient information was reported from which to make an assessment of this bias
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The study reports that outcome assessors were unaware of trial group assignments. No further details given
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	The trial protocol is available on www.clinicaltrials.gov ; all outcomes prespecified in the protocol have been reported on in the study in the prespecified way
Other bias	Low risk	The study appears to be free of other sources of bias

Godje 2006

Methods

Randomised controlled, single-centre trial

Randomly assigned at: Not stated

Godje 2006 (Continued)

Calculated sample size: Power calculation

Participants	<p>Condition: Cardiac surgery</p> <p>Inclusion criteria: Age 18 - 80 yrs, bodyweight 60 - 90 kg, scheduled coronary artery bypass graft surgery with left internal thoracic use</p> <p>Exclusion criteria: No intake of aspirin or other anti-coagulants for 5 days prior to procedure, normal standard coagulation laboratory data, no allergy to human plasma products</p> <p>Number of participants enrolled: Not stated</p> <p>Number of participants randomly assigned to the intervention: 25/25/25</p> <p>Number of participants included in the outcome analyses: 25/25/25 (but for bleeding and transfusion requirement outcomes, the numbers analysed were: 24/25/23)</p>
Interventions	<p>Intervention 1: Infusion of 1250 units of factor XIII immediately after administration of protamine (protamine was administered during the scheduled coronary artery bypass graft surgery)</p> <p>Intervention 2: Infusion of 2500 units of factor XIII immediately after administration of protamine (protamine was administered during the scheduled coronary artery bypass graft surgery)</p> <p>Comparator: Placebo infusion, standard surgical and post-operative care</p> <p>Additional treatments received whilst in the trial: "routine standard care"</p>
Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> Not stated <p>Secondary:</p> <ul style="list-style-type: none"> Standard haematology cell counts at 6,12, 24, 36, 48 hours and 5 days after administration of trial drug, Factor XIII levels at 6, 12, 24 hours and 5 days after administration of trial drug Drain volumes up to 6, 12, 36 and 48 hours after administration of the trial drug Administration of red blood cells, FFP and platelets <p>Duration of follow-up: Up to trial day 5</p>
Notes	<p>Population nationality: German</p> <p>Correspondence: goedje@pulsion.com</p> <p>Clinical trial registration number: None</p> <p>Sources of funding: Not stated.</p> <p>Dates the trial was conducted: Not stated</p> <p>Declarations of interest (among primary researchers): Not stated</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information was reported from which to make an assessment of this bias
Allocation concealment (selection bias)	High risk	The study reports that sealed envelopes were used to allocate treatment groups, but no further information on the envelopes is provided (e.g. whether they were sequentially numbered or opaque or both)

Godje 2006 (Continued)

Blinding of participants (performance bias)	Low risk	Participants under general anaesthesia when the intervention was administered
Blinding of study personnel (performance bias)	High risk	The study reports that "the trial drug was prepared in blackened syringes immediately prior to the operation by a person not involved in the trial or in the treatment of patients..." and that "due to the blinded design of the trial, neither the treating medical staff nor the trial nurses were informed about the type of trial medication, however this information was available within a sealed envelope in the records of the patients at any time". The fact that (unsecured) envelopes were clearly accessible to anyone interested makes this a high risk of performance bias
Blinding of outcome assessment (detection bias) All outcomes	High risk	The study does not report who undertook outcome assessment; treatment allocation was placed in unsecured envelopes within the participant's notes
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "No patients were lost during the follow-up and all patients left the hospital alive". 3 participants: (2 placebo and 1 who had received factor XIII 1250 IE) required a re-operation due to hints of early by-pass occlusion. All participants were retrospectively excluded from the analyses of bleeding and transfusion requirement outcomes. In 1 participant (received factor XII 12500 IE), exclusion seems to be during the post-operative period when they underwent a re-thoracotomy due to high drain loss. In the other 2 patients, their bleeding and transfusion requirement data were excluded from the analysis at the time of revision, but the study does not report the timing of the revision surgery. The reason for the exclusion of these 3 participants' data is because they were deemed incompatible with the other participants in the trial. All participants were included in the outcome reporting for the other trial outcomes
Selective reporting (reporting bias)	Low risk	All prespecified outcomes were reported, as prespecified, in the Results. No trial protocol
Other bias	Unclear risk	No declaration of source of funding or statement of author interests

Haas 2015

Methods	Single-centre, single-blinded, parallel randomised controlled trial Randomly assigned at: Not stated. Calculated sample size: A power of 80% and a 2-sided type I error of 0.05 based on a meaningful reduction for RBC transfusion revealed a number of 18 per group: for a better estimation of clinical relevance, 30 participants were scheduled for each group
Participants	Condition: elective craniosynostosis or scoliosis surgery Inclusion criteria: Male and female patients aged between 6 months and 17 years who were to have elective craniosynostosis or scoliosis surgery; received parental consent Exclusion criteria: Any diagnosed pre-existing congenital or acquired coagulation disorder; a medical history consistent with increased bleeding tendency; ongoing anticoagulation therapy; drug intake that could cause bleeding; clinical signs or diagnosis of acute thromboembolism; participation in another clinical trial; pregnant or lactating women Number of participants enrolled: 57: 31 who were undergoing craniosynostosis surgery; 26 who were undergoing scoliosis surgery

Haas 2015 (Continued)

Number of participants randomly assigned to the intervention: craniostomosis surgery = 17/14; scoliosis surgery = 10/9.

Number of participants included in the outcome analyses: craniostomosis surgery = 17/13; scoliosis surgery = 10/9 (but 2/5 of those undergoing scoliosis surgery did not receive the allocated treatment, although they were included in the analyses)

Interventions

Intervention: Fibrinogen concentrate (dose of 30 mg/kg, pump controlled iv administration over 10 minutes) if FIBTEM mean clot firmness < 13 mm: early substitution

Comparator: Fibrinogen concentrate (dose of 30 mg/kg) if FIBTEM mean clot firmness < 8 mm: conventional

Treatment was administered at any intra-operative measurement independent of the presence of bleeding. The trigger of < 8 mm in the control group was established based on the recommendation from a European guideline (Kozek-Langenecker 2013), and the concentration of < 13 mm represented the baseline concentration for this population (Haas 2012)

FIBTEM was also performed 10 minutes after the end of fibrinogen administration and fibrinogen concentrate was repeatedly administered if threshold FIBTEM concentrations were not reached

FIBTEM is a ready-to-use ROTEM system reagent for use with citrated whole blood. It assess the firmness of the fibrin clot

Additional treatments received whilst in the trial: All patients received antifibrinolytics prophylaxis by administration of tranexamic acid at an initial dose of 15 mg/kg followed by continuous infusion of 1.5 mg/kg/hour until 3 hours after transfer to the paediatric ICU

Outcomes
Primary:

- Total volume of transfused red blood cells per kg bodyweight (including transfused salvaged autologous blood) within 24 hours after surgery

Secondary:

- Requirements for all other transfused blood products within 24 hours after surgery
- Intraoperative blood loss based on the estimated total blood volume (calculated as described by Kearney 1989)
- Length of stay in paediatric ICU at 14 days after surgery
- Discharge status at 14 days after surgery
- Occurrence of bleeding that required revision surgery up to 14 days after surgery
- Treatment-related adverse events occurring within 14 days following surgery

Duration of follow-up: To day 14 following surgery

Notes

Population nationality: Swiss

Correspondence: thorsten.haas@kispi.uzh.ch

Clinical trial registration number: NCT01487837, prospectively registered (registered on 5 December 2011; trial started in January 2012)

Sources of funding: The work was supported by departmental funding for scientific member of staff by CSL Behring AG.

Dates the trial was conducted: February 2012 to September 2014

Declarations of interest (among primary researchers): The contact author has received speaker fees and travel support from the trial funders and 4 other drug companies. No other authors report a conflict of interest

Risk of bias

Haas 2015 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information about the sequence generation to permit a judgement. All that is stated is that "a randomisation list was used"
Allocation concealment (selection bias)	Low risk	<p>A telephone randomisation sequence was used</p> <p>At the beginning of the trial, a randomisation list, stratified by procedure was prepared by a statistician and sent to dedicated, trained staff members in the hospital pharmacy. After the start of anaesthesia for a consented trial participant, the investigator retrieved the randomisation group by calling the pharmacy</p> <p>Participants were randomised to 1 of the treatment groups after the start of anaesthesia and stratified by type of surgery</p>
Blinding of participants (performance bias)	Low risk	Participants under general anaesthesia when the intervention was administered
Blinding of study personnel (performance bias)	High risk	The study reports that "intraoperative staff and trial team members were not blinded to the randomisation. To mitigate this limitation, we used a strictly-enforced transfusion algorithm and blinded post-operative caregivers" (pg 8). "The staff in the PICU unit were blinded to the randomisation group"
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information was provided to permit a judgement
Incomplete outcome data (attrition bias) All outcomes	Low risk	No obvious missing data: 1 participant undergoing craniostomy surgery, randomised to the comparator arm, was excluded from the analysis as following randomisation and after an unexpected change in surgical procedure, the procedure and the expected blood loss was considered not comparable by the surgeon, but data for all other participants are available
Selective reporting (reporting bias)	Low risk	<p>The trial protocol is available as a trial registration document</p> <p>All outcomes detailed of interest on the trial registration document have been reported on in the study</p>
Other bias	Unclear risk	Member of author group on industrial salary with role in study design and data collection, declared no conflict of interest. No other concerns re. sources of bias.

Innerhofer 2017

Methods	<p>Single-centre, parallel-group, open-label, randomised trial</p> <p>Randomly assigned at: At admission to Emergency Department</p> <p>Calculated sample size: Power analysis</p>
Participants	<p>Condition: Adult major trauma</p> <p>Inclusion criteria: Adults (aged 18 – 80 years) admitted to the trauma centre with trauma exhibiting an Injury Severity Score > 15 and clinical signs or risk of substantial haemorrhage were screened for trauma-induced coagulopathy, which was defined as abnormally low fibrin polymerisation as measured with bedside ROTEM using the FIBTEM assay (10 min value of fibrinogen polymerisation (FibA10) < 9 mm) or prolonged initiation of coagulation in the extrinsically activated ROTEM (ExTEM) assay (coagulation time of ExTEM assay (ExCT) > 90s).</p>

Pro-coagulant haemostatic factors for the prevention and treatment of bleeding in people without haemophilia (Review)

Innerhofer 2017 (Continued)

Exclusion criteria: Injuries that were judged incompatible with survival, cardiopulmonary resuscitation on the scene, isolated brain injury, burn injury, avalanche injury, and pre-hospital coagulation therapy other than tranexamic acid. Additional exclusion criteria on www.clinicaltrials.gov: delayed (> 6 hours after trauma) admission to the Emergency Department; known use of oral anticoagulants, or platelet aggregation inhibitors within 5 days before injury; known history of severe allergic reaction to plasma products; known history of congenital haemostasis disturbance, IgA or Protein C deficiency; a history of thromboembolic events or heparin-induced thrombocytopenia type 2 within the last year; a body weight < 45 kg and > 150 kg; known to be pregnant; Jehovah's Witness; known participation in another clinical trial; known refusal of a participation in this clinical trial

Number of participants enrolled: 292 in total

Number of participants randomly assigned to the intervention: 52/48

Number of participants who received the treatment and were included in the outcome analyses: 50/44

Interventions	<p>Intervention: Fibrinogen concentrate and clotting factors (50 mg/kg fibrinogen)</p> <p>Comparator: FFP (15 mL/kg)</p> <p>Additional treatments received whilst in the trial: Cross-over therapy as rescue therapy; in FFP study arm participants received fibrinogen or PCC, or both. Also transfusion of RBC units and other blood products</p>
Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> • Occurrence of multiple organ failure during ICU stay as assessed by the daily SOFA score <p>Secondary:</p> <ul style="list-style-type: none"> • Length of ICU stay and hospital stay • Need for, and duration of haemofiltration • Ventilator-free days • Occurrence of sepsis, infection, venous thromboembolism, pulmonary embolism and in-hospital mortality at 24 hours and by day 30 • Numbers of transfusions of RBC units and platelet concentrates during the first 24 hours • Plasmatic and viscoelastic tests • Clinically relevant time intervals (time until randomisation, time until start of first study drug administration, and time until first normalised ROTEM (together with bleeding score of 0 or 1)) <p>Post-hoc;</p> <ul style="list-style-type: none"> • Mean SOFA score adjusted for injury severity score up to the first 7 days at the ICU • Analysis for the primary endpoint adjusted for injury severity score and brain injury status <p>Duration of follow-up: Duration of ICU stay for primary outcome; other outcomes throughout hospital stay to day 30</p>
Notes	<p>Population nationality: Austrian.</p> <p>Correspondence: petra.innerhofer@tirol-kliniken.at</p> <p>Clinical trial registration number: NCT01545635</p> <p>Sources of funding: None declared, departmental support</p> <p>Dates the trial was conducted: 3 March 2012 to 20 February 2016</p> <p>Declarations of interest (among primary researchers): PI has received personal fees from Baxter GmbH, CSL Behring GmbH, Fresenius Kabi GmbH Austria, Bayer GmbH Austria, and LFB, and non-financial support from TEM International, outside the submitted work. DF has received grants and personal fees from CSL Behring, LFB, and BBraun, outside the submitted work. MM has received personal fees</p>

Innerhofer 2017 (Continued)

from CSL Behring GmbH, outside the submitted work. IHL has received personal fees and non-financial support from Fresenius Austria, and non-financial support from MSD and Pfizer, outside the submitted work. SS has received personal fees from Fresenius Kabi GmbH Austria, outside the submitted work. All other authors declare no competing interests

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated random sequence
Allocation concealment (selection bias)	Low risk	Opaque envelopes, sequentially-opened, stratified for traumatic brain injury
Blinding of participants (performance bias)	Low risk	Unconscious or low conscious level at time of intervention
Blinding of study personnel (performance bias)	High risk	Unblinded, open-label
Blinding of outcome assessment (detection bias) All outcomes	High risk	For venous thromboembolism diagnosis outcome assessors were blinded to therapy. However for all other outcomes the research team were not blinded to therapy and made clinical decisions for input of management of participants
Incomplete outcome data (attrition bias) All outcomes	Low risk	Participant dropouts in each arm declared and explained and distributed across study arms
Selective reporting (reporting bias)	High risk	The trial initially declared a plan to report coagulopathy changes as an outcome, but data for this outcome were not reported in the published trial results
Other bias	Low risk	No other identifiable sources of bias

Jeppsson 2016

Methods	Randomised controlled, multicentre trial Randomly assigned at: Not stated. Study intervention given pre-operatively Calculated sample size: Power analysis
Participants	Condition: Elective cardiac surgery Inclusion criteria: Accepted for elective coronary artery bypass graft surgery and had a plasma fibrinogen concentration of ≤ 3.8 g/litre Exclusion criteria: Known liver or kidney disease, bleeding disorder, or ongoing treatment with a P2Y12-inhibitor Number of participants enrolled: 52 Number of participants randomly assigned to the intervention/placebo: 26/26 Number of participants who received the treatment/placebo and were included in the outcome analyses: 24/24

Jeppsson 2016 (Continued)

Interventions	<p>Intervention: Fibrinogen concentrate, 2 g</p> <p>Comparator: 0.9% sodium chloride solution</p> <p>Additional treatments received whilst in the trial: Blood and blood product infusion, general anaesthesia 300 units per kg of heparin and 1 mg protamine per 100 units of heparin</p>	
Outcomes	<p>Primary:</p> <ul style="list-style-type: none">• Reduction of mediastinal blood loss into the drain at 12 hours post-operatively <p>Secondary:</p> <ul style="list-style-type: none">• Estimated intraoperative bleeding volume• Total perioperative bleeding volume• Total number of blood components given• The proportion of transfused participants• Haemoglobin levels at 2 hours and 24 hours after cardiopulmonary bypass• Fibrinogen level, platelet count, prothrombin time and activated partial thromboplastin time before during and after cardiopulmonary bypass <p>Duration of follow-up: Primary outcome: 12 hours post-operatively; secondary outcomes up to duration of hospital stay</p>	
Notes	<p>Population nationality: Swedish</p> <p>Correspondence: anders.jepsson@vgregion.se</p> <p>Clinical trial registration number: NCT00968045</p> <p>Sources of funding: Swedish Heart-Lung Foundation, Sahlgrenska hospital and Gothenburg Medical Society</p> <p>Dates the trial was conducted: April 2009 to February 2012</p> <p>Declarations of interest (among primary researchers): A.J. has received honoraria from CSL Behring for lectures and advisory work, and research support for other studies than the present one. M.K. has received an honorarium from CSL Behring for a lecture</p>	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not stated
Allocation concealment (selection bias)	Low risk	Identical/masked infusion bottles; only independent research technician aware of allocation (not involved further in study) - randomisation with closed envelopes in pharmacy
Blinding of participants (performance bias)	Low risk	Under general anaesthesia
Blinding of study personnel (performance bias)	Low risk	Unaware of allocation but did have access to blood results, so may be able to work out by fibrinogen levels
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Unaware of allocation but did have access to blood results, so may be able to work out by fibrinogen levels

Jeppsson 2016 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	Incomplete outcome data but balanced participant losses across study arms
Selective reporting (reporting bias)	Low risk	All prespecified outcomes have been reported in the Results
Other bias	Low risk	Study appears clear of other sources of bias

Karkouti 2013

Methods	<p>Randomised controlled, multicentre trial</p> <p>Randomly assigned at: not stated</p> <p>Calculated sample size: Calculated for 70% power for 15% greater transfusion avoidance with FXIII, significance of 0.05</p>
Participants	<p>Condition: non-urgent cardiac surgery</p> <p>Inclusion criteria: aged 18 - 80 year undergoing non-urgent cardiac artery bypass graft surgery requiring cardio-pulmonary bypass, with moderate risk of transfusion</p> <p>Exclusion criteria: coagulation disorders; liver dysfunction (aspartate aminotransferase or alanine transaminase $\geq 2 \times$ upper limit); platelets under 100; autoimmune disease; creatinine ≥ 175 or dialysis; haematocrit $< 27\%$ during cardiopulmonary bypass; thromboembolic events (excluding cardiac); stroke; ST elevation myocardial infarction within 7 days; left ventricular ejection fraction $< 30\%$</p> <p>Number of participants enrolled: 579 (100 not randomised: 78 due to inclusion/exclusion criteria; 22 for other reasons)</p> <p>Number of participants randomly assigned to the intervention: 164/162/153</p> <p>Number of participants included in the outcome analyses: 143/138/128</p>
Interventions	<p>Intervention 1: single intravenous dose of recombinant factor XIII at 17.5 iU/kg before the induction of anaesthesia</p> <p>Intervention 2: single intravenous dose of recombinant factor XIII at 35 iU/kg before the induction of anaesthesia</p> <p>Comparator: single dose of an unspecified placebo before the induction of anaesthesia</p> <p>Additional treatments received whilst in the trial: Tranexamic acid and protamine during cardiopulmonary bypass.</p>
Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> Percentage of participants avoiding any allogeneic transfusions within 7 days or discharge home <p>Secondary:</p> <ul style="list-style-type: none"> Units of blood products transfused within 7 days or discharge home Percentage of participants avoiding any RBC transfusion and massive RBC transfusion (5 or more units) within 7 days or discharge home Incidence of reoperation until the end of the trial (= 7 weeks after trial drug administration) <p>Duration of follow-up: up to 7 weeks after trial drug administration</p>
Notes	<p>Population nationality: Multinational (13 European hospitals, 5 Canadian hospitals, 7 North American hospitals, 4 Japanese hospitals, 2 Israeli hospitals)</p> <p>Correspondence: keyvan.karlouti@uhn.ca</p>

Karkouti 2013 (Continued)

Clinical trial registration number: NCT00914589 prospectively registered. (Registered on 3 June 2009; trial started in July 2009)

Sources of funding: Novo Nordisk

Dates the trial was conducted: 27 July 2009 through to 23 February 2011

Declarations of interest (among primary researchers): 6 of the 8 authors have declared receiving some level of commercial support; 1 of these authors is employed by the company sponsoring the trial.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	The trial reports "randomised (1:1:1) using an interactive voice/Web response system".
Allocation concealment (selection bias)	Unclear risk	Insufficient information was reported from which to make an assessment of this bias
Blinding of participants (performance bias)	Low risk	Participants under general anaesthesia when the intervention was administered
Blinding of study personnel (performance bias)	High risk	The study does not outline how intervention selection was performed or who was aware of this selection, such that transfusion outcomes are potentially open to bias
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Outcome assessors were reported to be blinded to the intervention received by the participant
Incomplete outcome data (attrition bias) All outcomes	Low risk	Significant dropout and exclusion rates but all cases accounted for and documented. 479 participants were randomised, but only 409 participants received their allocated intervention: all due to withdrawals pre-dosing. All 409 participants were included in the outcome analyses although only 356 participants completed the trial: the reason for all exclusions and non-completion of the trial are documented in the study report
Selective reporting (reporting bias)	Low risk	All prespecified outcomes were reported in the Results: trial protocol registered online
Other bias	High risk	Variations in transfusion practice despite protocols. The company producing the product and sponsoring the trial oversaw the trial operations, audited all data, and conducted the statistical analyses. Unclear ownership of data, although state full author access.

Karlsson 2011

Methods	Single-centre randomised controlled trial Randomly assigned at: on arrival in the operating theatre immediately before surgery Calculated sample size: Not stated
Participants	Condition: elective cardiac surgery Inclusion criteria: elective coronary artery bypass graft, preoperative plasma fibrinogen < 3.8 g/L

Karlsson 2011 (Continued)

Exclusion criteria: kidney or liver disease, bleeding disorders, surgical source of bleeding at acute re-exploration

Number of participants enrolled: 20

Number of participants randomly assigned to the intervention: 10/10

Number of participants included in the outcome analyses: 10/10

Interventions	<p>Intervention: Preoperative infusion of fibrinogen concentrate (2 g) at 5 minutes following baseline measurements</p> <p>Control: No intervention</p> <p>Additional treatments received whilst in the trial: Heparin (300 units/kg body weight) to maintain an activated clotting time of > 480 seconds during cardiopulmonary bypass; protamine (1 mg protamine/100 units heparin) for heparin reversal, RBC, aspirin</p>
Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> Clinical adverse events, including graft occlusion on CT up to 3 to 4 days following surgery Markers of coagulation, including ROTEM, fibrinogen levels <p>Secondary:</p> <ul style="list-style-type: none"> Post-operative bleeding 24 hours after surgery Blood transfusions within 24 hours following surgery Haemoglobin concentration 24 hours after surgery Thromboelastometry <p>Duration of follow-up: to 3 to 4 days following surgery</p>
Notes	<p>Population nationality: Swedish</p> <p>Correspondence: martin.karlsson@hjl.gu.se</p> <p>Clinical trial registration number: None reported, assumed not registered</p> <p>Sources of funding: Supported in part by</p> <ul style="list-style-type: none"> the Swedish Heart & Lung Foundation, CSL Behring, Marburg, Germany Sahlgrenska University Hospital, Gothenburg, Sweden <p>Dates the trial was conducted: September to December 2006</p> <p>Declarations of interest (among primary researchers): None stated</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Although the study reports that "random assignment was conducted using unmarked envelopes each containing a card indicating treatment with fibrinogen or control", no information as to whether the cards/envelopes were shuffled was provided
Allocation concealment (selection bias)	Unclear risk	There is insufficient information to fully determine the risk of bias for this domain. See above for what is reported by the study

Karlsson 2011 (Continued)

Blinding of participants (performance bias)	Low risk	Participants under general anaesthesia when the intervention was administered
Blinding of study personnel (performance bias)	Low risk	The trial reports that only the trial co-ordinator was informed about allocation assignment. None of the operating staff, surgeon, anaesthesiologist nor the staff at the ICU were informed of trial assignment Whilst there is potential for blinding to have been broken during infusion of fibrinogen product in this study such that, as no transfusion trigger was allocated for red blood cell transfusion, this outcome could be at risk of clinician bias
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The study reports that only the trial co-ordinator was informed about treatment allocation. The outcomes are all clinical and objective. No indication is provided as to who reported these but as they occurred within a short time following surgery and all clinical staff were blinded to treatment allocation, we felt that outcome assessment would not likely be influenced by lack of blinding
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	The number of participants included in outcome analysis was not reported for all outcomes (with the exception of TEG and post-operative cardiac CT scan) We have assumed for the analysis that as the outcomes were measured in the immediate aftermath of surgery and all 20 participants were discharged from clinic at 7 days that all were included in outcome analyses
Selective reporting (reporting bias)	Low risk	The trial protocol is available: all prespecified outcomes outlined in the protocol were reported in the Results
Other bias	Low risk	Study support included industrial funding but sponsors had no involvement in the study or publication.

Korte 2009

Methods	Randomised controlled, single-centre trial Randomly assigned at: not stated Calculated sample size: Power calculation
Participants	Condition: elective surgery for gastrointestinal cancer Inclusion criteria: elective surgery for gastrointestinal cancer, 18 yrs or older, preoperative fibrin monomer > 3µg/L, American Society of Anaesthesiology physical status classification score of 2 or higher Exclusion criteria: congenital bleeding disorder, intolerance to planned transfusion triggers, contraindications to the use of plasma proteins, history of cerebrovascular or cardiovascular events, symptomatic peripheral artery disease, deep venous thrombosis or pulmonary embolism within the last 5 yrs, BMI > 35kg/m ² , and pregnancy Number of participants enrolled: 25 Number of participants randomly assigned to the intervention: not clear, 22 overall as there were 3 early dropouts due to rescheduling of surgery Number of participants included in the outcome analyses: not clear, the study reports evaluating 22 participants (pg 241)

Korte 2009 (Continued)

Interventions	<p>Intervention: Factor XIII (30 U/kg) infused 15 minutes after the beginning of surgery</p> <p>Comparator: Placebo: albumin infusion.</p> <p>Additional treatments received whilst in the trial: Heparin starting the day before surgery as per thromboprophylaxis throughout hospitalisation guidelines</p>	
Outcomes	<p>Primary:</p> <ul style="list-style-type: none">• Clot firmness over 195 minutes of surgery <p>Secondary:</p> <ul style="list-style-type: none">• Fibrinogen and prothrombin fragments F1 + F2 levels at 195 minutes of surgery• Blood loss upon completion of surgery• Blood product support upon completion of surgery <p>Duration of follow-up: not specifically stated</p>	
Notes	<p>Population nationality: Swiss</p> <p>Correspondence: wolfgang.korte@ikch.ch</p> <p>Clinical trial registration number: Not registered</p> <p>Sources of funding:</p> <ul style="list-style-type: none">• Institute for Clinical Chemistry and Haematology, the Institute for Anesthesiology, Kantonsspital St. Gallen, St. Gallen, Switzerland• CSL Behring, Bern, Switzerland; Dade Behring, Marburg, Germany; Pentapharm, Basel, Switzerland• Novo Nordisk, Zurich, Switzerland <p>Dates the trial was conducted: None stated</p> <p>Declarations of interest (among primary researchers): One author (WK) has received support from commercial drug companies and one author (JD) is an employee of a sponsor company of the trial, CSL Behring, Bern</p>	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Participants were randomised using a predefined computer-based block-wise randomisation plan
Allocation concealment (selection bias)	Low risk	Randomisation was performed by the Pharmacy Department and was kept blinded to anyone outside the Pharmacy during the trial period
Blinding of participants (performance bias)	Low risk	No details are given as to the blinding of the participants, but the participants were anaesthetised during the administration of the intervention or placebo and special care was taken to make the trial intervention (factor 13 substitution) and the placebo (albumin) indistinguishable
Blinding of study personnel (performance bias)	Low risk	Low risk was assigned as the study reports that the trial medication was prepared by the pharmacy and directly shipped to the anaesthesiologist in the operating room ready to be used. In addition, special care was taken to make the trial intervention (factor 13 substitution) and the placebo (albumin) indistinguishable
Blinding of outcome assessment (detection bias)	Unclear risk	No details are given as to the blinding of the outcome assessors or who these people would be

Korte 2009 (Continued)

All outcomes

Incomplete outcome data (attrition bias) All outcomes	Unclear risk	The study does not report the number of participants randomised to each treatment arm and in some cases the number of participants analysed per outcome. 22 of the 25 enrolled participants were evaluated. The study does report the number of protocol violations (1 per treatment arm, but seemingly evaluated) and the number who dropped out before receiving their randomised intervention (3 due to rescheduling of surgery at short notice and these participants were not treated or followed up within the trial)
Selective reporting (reporting bias)	Low risk	All outcomes prespecified in the Methods were reported in the Results
Other bias	Unclear risk	Interim analysis but no evidence this has led to an overestimation of the impact of intervention. Industry sponsored and industrial employee on author list but no indication of roles or independence of publication or data ownership.

Lance 2012

Methods	Randomised controlled trial, single-centre Randomly assigned at: massive bleeding during or after surgery (massive bleeding was defined as prolonged blood loss of > 150 mL/hour or > 1.5 mL/kg/20 min or acute blood loss of > 700 mL at once) Calculated sample size: None stated
Participants	Conditions: elective surgery (cardiovascular, abdominal or spinal column) Inclusion criteria: Admitted for cardiovascular, major abdominal or orthopaedic surgery, expected to last > 120 mins Exclusion criteria: Age under 18 yrs, active HIV infection, known coagulation abnormalities, deep hypothermia with circulatory arrest or preoperative need of transfusion Number of participants enrolled: 307 (255 did not meet the pre-defined inclusion criteria of a massive haemorrhage) Number of participants randomly assigned to the intervention: 52 (but 9 did not require treatment), so numbers = 22/21 Number of participants included in the outcome analyses: 22/21 for all, but only 16/16 for the whole blood ROTEM analysis (primary outcome analysis)
Interventions	Intervention: Two units of FFP plus 2g of fibrinogen (Haemocomplettan P 20 g/L) transfused as a consequence of massive bleeding during or after surgery Comparator: 4 units of FFP transfused as a consequence of massive bleeding during or after surgery. Additional treatments received whilst in the trial: heparin in 25 participants who received cardiovascular surgery
Outcomes	Primary: <ul style="list-style-type: none">• Pre- and post-intervention, ROTEM analysis, including clotting time and clot formation time Secondary: <ul style="list-style-type: none">• Fibrinogen levels• Other coagulation factor levels

Lance 2012 (Continued)

- Haematological cell counts

Duration of follow-up: Not stated

Notes

Population nationality: Dutch

Correspondence: marcus.lance@mumc.nl

Clinical trial registration number: NL.23565.068.08

Sources of funding: CSL-Behring, Breda (Netherlands), Marburg (Germany)

Dates the trial was conducted: Not stated

Declarations of interest (among primary researchers): 2 authors (SS and JH) hold unrestricted grants with the funding sources. All other authors had no conflict of interest to disclose

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information was reported from which to make an assessment of this bias
Allocation concealment (selection bias)	High risk	Closed envelope method: no further details are reported
Blinding of participants (performance bias)	Low risk	No details are given as to the blinding of the participants but participants were under general anaesthesia when the intervention was administered
Blinding of study personnel (performance bias)	Unclear risk	Insufficient information was reported from which to make an assessment of this bias. It is unclear whether the anaesthetist (administering the intervention) has control over other components of care, or is involved in trial, which could impact on transfusion decisions and outcome assessment
Blinding of outcome assessment (detection bias) All outcomes	Low risk	No details are given as to the blinding of the outcome assessors or who these people would be, but as the outcomes were all objective, we felt that outcome assessment would not be influenced by lack of blinding
Incomplete outcome data (attrition bias) All outcomes	Low risk	All the participants randomised in this trial were included in the outcome analyses, with the exception of one of the primary analyses: ROTEM of whole blood where only 74% (N = 16/16) participants were included. No details were given as to the reason for this, but ROTEM analyses of platelet-free plasma and platelet-rich plasma were also undertaken and full data on these primary outcome analyses have been reported
Selective reporting (reporting bias)	Unclear risk	No trial protocol and it is not entirely clear what were the outcomes of interest in this trial. In addition, the study does not address the finding of a deep vein thrombosis in 1 participant, or consider whether reported adverse events could be associated with the interventions
Other bias	Low risk	The study appears to be free of other sources of bias

Levy 2009

Methods

Prospective, randomised, multi-centre, dose-escalation trial.

Levy 2009 (Continued)

Randomly assigned prior to anaesthesia

Calculated sample size: Power calculation

Participants	<p>Conditions: Elective cardiac surgery</p> <p>Inclusion criteria: Aged 35 – 75 years, scheduled for elective first-time myocardial revascularisation requiring cardiopulmonary bypass</p> <p>Exclusion criteria: Pre-operative anaemia (haemoglobin < 11.8 g/dl (women) or < 13.5 g/dl (men)), nephropathy (creatinine > 1.4 mg/dl), acute myocardial infarction within 24 hours prior to surgery, or a history of non-cardiac vascular diseases or extra-myocardial thromboembolic events. Patients treated with clopidogrel, heparin or glycoprotein IIb/IIIa inhibitors. Intraoperative exclusion criteria were suspected or confirmed intra-operative MI, treatment with anti-fibrinolytics (aprotinin, epsilon-aminocaproic acid or tranexamic acid), treatment with recombinant, activated factor VII and severe haemodynamic instability</p> <p>Number of participants enrolled: not stated</p> <p>Number of participants randomly assigned to the intervention: 43; N in intervention 1 = 10; intervention 2 = 9; intervention 3 = 8; intervention 4 = 8; comparator = 8.</p> <p>Number of participants included in the outcome analyses: as randomised for the primary outcome</p>
Interventions	<p>After termination of cardiopulmonary bypass:</p> <p>Intervention 1: Infusion of a single dose of recombinant FXIII-A2 at 11.9 units/kg, given as a slow iv injection within 10 to 20 minutes after the first protamine dose</p> <p>Intervention 2: Infusion of a single dose of recombinant FXIII-A2 at 25 units/kg, given as a slow iv injection within 10 to 20 minutes after the first protamine dose</p> <p>Intervention 3: Infusion of a single dose of recombinant FXIII-A2 at 35 units/kg, given as a slow iv injection within 10 to 20 minutes after the first protamine dose;</p> <p>Intervention 4: Infusion of a single dose of recombinant FXIII-A2 at 50 units/kg, given as a slow iv injection within 10 to 20 minutes after the first protamine dose</p> <p>Comparator: Infusion of a single dose of variable doses of 'placebo' compound (no further details given), given as a slow iv injection within 10 to 20 minutes after the first protamine dose</p> <p>Additional treatments received whilst in the trial: protamine</p>
Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> Incidence and severity of adverse events from trial drug administration through to follow-up at 5 to 7 weeks <p>Secondary:</p> <ul style="list-style-type: none"> Immunogenicity (antibody formation against recombinant FXIII-A2 and yeast) at the follow-up visit (5 to 7 weeks) FXIII activity measurements and concentration of free FXIII-B subunit Chest tube drainage volume Requirement for allogeneic blood products (red blood cells, FFP, platelets, cryoprecipitate, fibrinogen concentrate) The percentage of subjects avoiding allogeneic blood product within 7 days of surgery or until discharge (whichever occurred first) <p>Duration of follow-up: 5 - 7 weeks after receipt of intervention</p>
Notes	<p>Population nationality: Mixed (9 centres in the USA, 1 centre each from Denmark and Germany)</p>

Levy 2009 (Continued)

Correspondence: jlevy01@emory.edu

Clinical trial registration number: none stated

Sources of funding: Novo Nordisk.

Dates the trial was conducted: Not stated

Declarations of interest (among primary researchers): 3 authors (JL, RG, NN) serve as consultants for the funding source; 3 authors (HA, FB, CJ) are employees of the funding source

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Participants were randomised prior to anaesthesia using an Interactive Voice Response System
Allocation concealment (selection bias)	Low risk	Participants were randomised prior to anaesthesia using an Interactive Voice Response System and the interventions were supplied in identical containers as a lyophilised powder for reconstitution
Blinding of participants (performance bias)	Low risk	No details are given as to the blinding of the participants, but as the participants were anaesthetised during the administration of the intervention and were not involved in outcome assessment
Blinding of study personnel (performance bias)	Low risk	The study reports that the interventions were provided in visually identical containers
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details are given as to the blinding of the outcome assessors or who these people would be
Incomplete outcome data (attrition bias) All outcomes	High risk	All patients (N = 43) were included in the safety evaluation; 38 participants in a per protocol analysis and 35 in the pharmacokinetic analysis. The number and reason for participant exclusion from analysis was reported but not the details of the intervention group to which they were randomised, thus it is not possible to state whether the missing outcome data was balanced across the groups
Selective reporting (reporting bias)	High risk	1 predefined outcome, the percentage of participants who avoided receiving allogeneic red blood cells within 7 days of surgery or until discharge, was not reported in the Results All other predefined outcomes were reported in the Results
Other bias	High risk	Industrial sponsor of study. Three authors including senior author are employees of sponsor. No statement of data ownership, study independence or industrial control of data, manuscript content or publication.

Najafi 2014

Methods Prospective, randomised, single-centre trial

Randomly assigned at: Not stated

Calculated sample size: Yes

Participants **Conditions:** Total hip replacement arthroplasty

Najafi 2014 (Continued)

Inclusion criteria: All candidates for total hip replacement arthroplasty; ASA physical status class I or II; metabolic equivalent of the task >4; haemoglobin level > 10 g/dL.

Exclusion criteria: No consent to stay in project at any time; ASA class > 2; metabolic equivalent of the task < 4; pregnancy; any history of sensitivity to blood products; history of myocardial infarction; deep vein thrombosis; pulmonary emboli; cerebrovascular accident and known haematologic disorder

Number of participants enrolled: not stated

Number of participants randomly assigned to the intervention: 15/15

Number of participants included in the outcome analyses: 15/15

Interventions	<p>Intervention: Fibrinogen concentrate 30 mg/kg infused after induction of general anaesthesia. The fibrinogen concentrate was dissolved in 100 mL and infused within 10 minutes of being dissolved in distilled water</p> <p>Comparator: Normal saline administered in equal volume as fibrinogen</p> <p>Additional treatments received whilst in the trial: None stated</p>
Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> • Volume of RBC transfused during and 24 hours after total hip arthroplasty surgery • Volume of bleeding during and 24 hours after total hip arthroplasty surgery • Haemoglobin concentration level 1 hr from start of surgery and at 24 hours after surgery; • Complications of transfusion during and 24 hours after surgery. <p>Secondary:</p> <ul style="list-style-type: none"> • Volume of transfused other blood products during and after surgery • Thromboembolic events during hospitalisation and after discharge up to 2 weeks (according to fibrinogen half life) • Transfusion-related long-term complication after surgery • Cardiovascular complications due to bleeding during and after surgery <p>Duration of follow-up: Not stated</p>
Notes	<p>Population nationality: Iranian</p> <p>Correspondence: horandi@sina.tums.ac.ir</p> <p>Clinical trial registration number: Iranian Registry of Clinical Trials registry number IRC-T201012264784N2. Retrospectively registered while recruiting. Registered on 11 June 2012</p> <p>Sources of funding: None stated</p> <p>Dates the trial was conducted: August 2011 to February 2013</p> <p>Declarations of interest (among primary researchers): None stated</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Table of random numbers
Allocation concealment (selection bias)	Unclear risk	Insufficient information was reported from which to make an assessment of this bias

Najafi 2014 (Continued)

Blinding of participants (performance bias)	Unclear risk	Insufficient information was reported from which to make an assessment of this bias
Blinding of study personnel (performance bias)	Unclear risk	Insufficient information was reported from which to make an assessment of this bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information was reported from which to make an assessment of this bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	There were no missing data; all randomised participants were included in the outcome analyses
Selective reporting (reporting bias)	High risk	Trial registration form is available (www.irct.ir) but data were not reported for 2 predefined outcomes: 'volume of transfused other blood products during and after surgery' and 'transfusion related long term complication long term after surgery'
Other bias	Unclear risk	Insufficient information was reported from which to make an assessment of this bias

Nascimento 2016

Methods	<p>Prospective, randomised, single-centre trial</p> <p>Randomly assigned at: Within 30 mins of hospital arrival</p> <p>Calculated sample size: Feasibility study</p>
Participants	<p>Condition: Severe adult trauma</p> <p>Inclusion criteria: Adult (age > 18 yrs) severe trauma (blunt or penetrating) patients were eligible if they were: assessed by the trauma team at the research institution and identified as being at risk for significant haemorrhage as evidenced by: systolic arterial pressure < 100 mmHg and requiring uncrossmatched RBC transfusion at any time from injury until 30 mins after hospital arrival</p> <p>Exclusion criteria: Receipt of any blood or blood products before admission to trauma centre; presented more than 6 hours after injury; estimated body weight < 50 kg; known or suspected pregnancy; catastrophic brain injury (defined as any of: Glasgow Coma Scale of 3 as a result of brain injury; need for immediate neurosurgery, focal signs such as anisocoria or imaging evidence of intracranial bleeding with mass effect, transcranial gunshot wound, or open skull fracture with exposure/loss of brain tissue); non-haemorrhagic shock (i.e. obstructive (cardiac tamponade, tension pneumothorax and massive pulmonary emboli, neurogenic, cardiogenic, or septic); underlying hereditary or acquired coagulopathy; known or suspected use of anticoagulant medications such as warfarin, low-molecular weight heparin, and direct thrombin and factor Xa inhibitors; or were moribund and predicted to expire in a few hours</p> <p>Number of participants enrolled: 50</p> <p>Number of participants randomly assigned to the intervention/placebo: 21/24</p> <p>Number of participants who received the treatment/placebo and were included in the outcome analyses: 20/24</p>
Interventions	<p>Intervention: Fibrinogen concentrate, 6 g</p> <p>Comparator: 0.9% normal saline solution</p>

Nascimento 2016 (Continued)

Additional treatments received whilst in the trial: Blood, other clotting and haemostatic factors as required.

Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> Feasibility of delivery of intervention within 60 mins of admission <p>Secondary:</p> <ul style="list-style-type: none"> 28-day all-cause mortality Rates of symptomatic thromboembolic complications (defined by the evidence of deep venous thrombosis, or myocardium infarction, or cerebral vascular accident, or pulmonary embolism, or arterial thrombosis at any time during hospital stay) Rates of asymptomatic deep vein thrombosis (evidenced by leg Doppler performed at day 7 of hospital stay) Incidence of acute lung injury/acute respiratory distress syndrome (defined by the Berlin Classification of acute lung injury) during hospitalisation Allergic reactions to the SI infusion Feasibility: duration of infusion, wastage of trial intervention, missed participants, randomisation errors Fibrinogen levels <p>Duration of follow-up: Primary outcome at 1 hour after arrival in hospital; secondary outcomes up to 28 days post-admission</p>
Notes	<p>Population nationality: Canadian</p> <p>Correspondence: Barto.Nascimento@sunnybrook.ca</p> <p>Clinical trial registration number: NCT02203968.</p> <p>Sources of funding: Supported by grants from</p> <ul style="list-style-type: none"> Canadian Forces Health Services, Defence Research and Development Canada CSL Behring <p>Dates the trial was conducted: October 2014 to November 2015</p> <p>Declarations of interest (among primary researchers): B.N. received speaker honoraria and a research grant from CSL Behring, the manufacturer of RiaSTAPTM. S.R. received unrestricted educational grants for a Consensus Conference on ROTEMTM from CSL Behring and TEM International. H.T. was part of the Canadian Armed Forces during the time this trial was conducted. H.P. is a scientist at the Defence Research and Development Canada and Canadian Forces Health Services</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	A computerised random-number generator was used to generate sequences of random numbers, blocks of 4
Allocation concealment (selection bias)	Low risk	Allocation was concealed with sealed opaque envelopes in the blood bank
Blinding of participants (performance bias)	Low risk	Identical infusion bags and equipment covered with amber bag to ensure blinding
Blinding of study personnel (performance bias)	Low risk	Identical infusion bags and equipment covered with amber bag to ensure blinding

Nascimento 2016 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Unaware of allocation but did possibly know fibrinogen levels
Incomplete outcome data (attrition bias) All outcomes	Low risk	Unequal loss of participants from each group but major outcomes presented in ITT format and dropouts prior to intervention and due to case-specific events
Selective reporting (reporting bias)	Low risk	Study protocol with objectives listed is available
Other bias	Unclear risk	Insufficient information was reported from which to make an assessment of this bias

Pournajafian 2015

Methods	Prospective, randomised, single-centre trial Randomly assigned at: Not stated Calculated sample size: Power analysis
Participants	Condition: Posterior spinal fusion surgery Inclusion criteria: 18 - 60 years old, BMI < 30kg/m ² and ASA Class I or II Exclusion criteria: History of posterior spinal fusion surgery; cardiovascular, respiratory, cerebrovascular, liver or kidney diseases; coagulopathy; diabetes; use of beta and calcium channel blockers, digoxin, tricyclic antidepressants, anticoagulants or clonidine; alcohol or drug abuse; the presence of a tumour in the surgical area; expansion of surgical area to more than 'three parts'; any type of excessive and abnormal bleeding required coagulant agents. In addition, when the procedure lasted more than 4 hours, these patients were excluded from the study Number of participants enrolled: 41 Number of participants randomly assigned to the intervention: 21/20 Number of participants who received the treatment and were included in the outcome analyses: 21/20 Note that 45 participants were initially enrolled but with 4 subsequent exclusions for unclear reasons. It is also unclear when these people were excluded, as some exclusion criteria (i.e. duration of surgery) could be applied following randomisation and administration of the study intervention.
Interventions	Intervention: Fibrinogen 1 g Comparator: Placebo, unclear nature Additional treatments received whilst in the trial: Unclear
Outcomes	Primary: Unclear Secondary: Unclear Duration of follow-up: Unclear
Notes	Population nationality: Iranian. Correspondence: pournajafian.ar@iums.ac.ir Clinical trial registration number: Not stated

Pournajafian 2015 (Continued)

Sources of funding: Unclear

Dates the trial was conducted: 2014 to 2015

Declarations of interest (among primary researchers): Unclear

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not stated
Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of participants (performance bias)	Low risk	Intervention given under general anaesthesia
Blinding of study personnel (performance bias)	Unclear risk	Not stated
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	State that outcome assessors were blinded to participant allocation, but transfusion thresholds not declared and blood transfusion likely to have been determined by clinicians for whom blinding status is not clear
Incomplete outcome data (attrition bias) All outcomes	High risk	4 participants excluded at unclear phases of the study, but following recruitment. Potentially excluded following randomisation and treatment administration
Selective reporting (reporting bias)	Unclear risk	No study protocol available. No declared outcomes
Other bias	Unclear risk	Insufficient information available to make an assessment of this source of bias

Rahe-Meyer 2013

Methods	Randomised controlled, single-centre trial Randomly assigned before surgery Calculated sample size: To provide a reduction in transfusion requirement of 4.5 units of RBCs in the intervention arm with 80% power and 5% sensitivity
Participants	Conditions: Elective aortic valve replacement surgery Inclusion criteria: aged 18 or above; aortic valve replacement surgery involving cardiopulmonary bypass; all aortic valve operations with aortic root/ascending aortic replacement (thoracic aortic with/without aortic arch replacement) and thoracoabdominal replacements Exclusion criteria: previous surgery at the same site, congenital or acquired clotting disorder, myocardial infarction or stroke within 2 months of surgery, use of aspirin, clopidogrel or vitamin K antagonist within 2 - 5 days of surgery Number of participants enrolled: 80 Number of participants randomly assigned to the intervention: 38/42 Number of participants included in the outcome analysed: 29/32
Interventions	Trial treatment was administered within 5 minutes of a clinically relevant bleeding event

Rahe-Meyer 2013 (Continued)

Intervention: Fibrinogen concentrate with dose determined by FIBTEM test. Median (IQR range) dose: 8 g (6 to 9 g)

Comparator: Normal (0.9%) saline. Median (IQR range) dose: 400 mL (300 to 450 mL)

Additional treatments received whilst in the trial: All participants received tranexamic acid during surgery Post-operative bleeding managed with platelet or FFP transfusions (either trial arm)

Outcomes	<p>Primary:</p> <ul style="list-style-type: none">• Total number of units of allogeneic blood components given during first 24 hours post-operatively <p>Secondary:</p> <ul style="list-style-type: none">• Number of units of each blood components given• Proportion of participants receiving no blood products• Number of days not in ICU or hospital within 45 days of operation• Mortality at 45 days after surgery;• Treatment-emergent adverse events within 10 days of treatment• Serious adverse events within 45 days of treatment <p>Duration of follow-up: day 45</p>	
Notes	<p>Population nationality: German</p> <p>Correspondence: niels.rahe-meyer@franziskus.de</p> <p>Clinical trial registration number: Prospectively registered (16 June 2008): NCT00701142</p> <p>Sources of funding: Sponsored by CSL Behring</p> <p>Dates the trial was conducted: June 2008 through to April 2010</p> <p>Declarations of interest (among primary researchers): 3 authors (DS, DK and GH) were employed by the funding company</p>	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation numbers were assigned sequentially, in a 1:1 ratio, with a block size of 4, stratified by type of aortic surgery
Allocation concealment (selection bias)	Low risk	Centrally allocated by contract research organisation
Blinding of participants (performance bias)	Low risk	Intervention while under general anaesthesia. No subjective outcomes
Blinding of study personnel (performance bias)	Low risk	Intervention product made up centrally by unblinded pharmacists. Point-of-care lab technicians were also unblinded The investigators, operating room and ICU staff were blinded to treatment allocation and did not have access to FIBTEM or fibrinogen concentration results after trial interventions were administered. Opaque syringes that were visually identical
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The investigators, operating room and ICU staff were blinded to treatment allocation and did not have access to FIBTEM or fibrinogen concentration results after trial interventions were administered
Incomplete outcome data (attrition bias)	Low risk	24% of participants in both arms (9 participants randomised to receive fibrinogen and 10 participants randomised to placebo) did not receive their allocated

Rahe-Meyer 2013 (Continued)

All outcomes		ed intervention: this is fully detailed and similar reasons for this by treatment group were given in the study report.
		All participants who received their allocated intervention were included in the efficacy and safety analyses of outcomes
Selective reporting (reporting bias)	Low risk	The trial registration form is available and all outcomes detailed on the form have been reported in the Results section
Other bias	High risk	Three study authors were employees of the funding company during the study. No statement of data ownership or publication decision making and responsibility.

Rahe-Meyer 2016

Methods	Randomised, controlled multi-centre international study Randomly assigned at: At the start of surgery Calculated sample size: Power analysis
Participants	Condition: Complex cardiovascular surgery Inclusion criteria: Age 18 and above. first 5 mins bleeding mass of 60 – 250 g; body temperature $\geq 35^{\circ}\text{C}$; activated clotting time within 25% of the baseline value; and blood pH > 7.3 . The 5-min bleeding mass was defined as weight gained by surgical swabs after 5 min of packing of the surgical site; it was initially measured after surgical haemostasis was completed after separation from cardiopulmonary bypass and heparin reversal. Elective open surgical procedure on any part of the aorta requiring cardiopulmonary bypass, with or without other cardiac procedures Exclusion criteria: Re-operative aortic surgery at the same site as the original procedure; operation for infection; and any known coagulation disorder. Prior use of any haemostatic therapy during the surgical procedure; and any perception that study participation could threaten the patient's safety Number of participants enrolled: 519 participants were enrolled; 152 treated as per inclusion criteria Number of participants randomly assigned to the intervention: 78/74 Number of participants who received the treatment and were included in the outcome analyses: 78/74
Interventions	Intervention: Fibrinogen concentrate Comparator: 0.9% saline solution Additional treatments received whilst in the trial: Red blood cell units if Hb < 7 or if end organ ischaemia, acute blood loss or patient-specific reason, antifibrinolytics could be given according to each site SOP. Heparin and protamine as per cardiac bypass
Outcomes	Primary: <ul style="list-style-type: none">The number of units of allogeneic blood products (FFP, platelets, and RBCs) administered during the 24 hours after administration of study medication Secondary: <ul style="list-style-type: none">Number of units of individual allogeneic blood productsTotal avoidance of transfusionsNeed for re-operationMortalityPlasma fibrinogen (central laboratory, Clauss assay)

Rahe-Meyer 2016 (Continued)

- MCF (FIBTEM assay performed at each site using a ROTEM device)
- Incidence of treatment-emergent adverse events throughout the 46-day observation period
- Changes in vital signs and laboratory values and ECG assessments
- Viral transmission by enzyme-linked immunosorbent assays and polymerase chain reaction tests

Duration of follow-up: Primary outcome at 24 hours after study intervention; other outcomes up to 46 days post-operatively

Notes

Population nationality: Austria, Brazil, Canada, Czech republic, Copenhagen, Finland, Germany, Italy, Japan, Poland, UK.

Correspondence: niels.rahe-meyer@franziskus.de

Clinical trial registration number: NCT01475669; EnduraCT: 2011-002685-20

Sources of funding: CSL Behring

Dates the trial was conducted: 23 January 2012 to 11 September 2014.

Declarations of interest (among primary researchers):

- N.R.-M. serves on steering committees and advisory boards and is Principal Investigator for CSL Behring and Merck. J.H.L. serves on steering committees for Boehringer Ingelheim, CSL Behring, Grifols, Instrumentation Laboratories, and Janssen.
- C.D.M. has received research support and honoraria from CSL Behring, Medicines Company, and Fresenius Kabi.
- A.S. has received honoraria from CSL Behring, Grifols, B. Braun, and Fresenius Kabi in addition to educational grant funding from Orion.
- A.A.K. has received educational grant funding from CSL Behring, Brightwake, and Pharamacosmos.
- R.B. and Y.O. have no interest to declare.
- Y.U. serves on steering committees for CSL Behring in Japan and has received honoraria from CSL Behring for consultation.
- D.S.S. and R.R. are full-time employees and shareholders of CSL Behring.
- R.G. serves on steering committees for CSL Behring, serves on advisory boards for Octapharma, and has been paid to speak on behalf of CSL Behring, Octapharma, and Roche

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	By interactive voice system although method not defined
Allocation concealment (selection bias)	Unclear risk	Study medication was prepared by a pharmacist or blood bank but concealment of allocation not stated
Blinding of participants (performance bias)	Low risk	Under general anaesthesia
Blinding of study personnel (performance bias)	Low risk	Investigator blinding was maintained by providing placebo at an equivalent volume to the fibrinogen concentrate dose and by using identical opaque syringes
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	It is unclear whether fibrinogen levels were available to staff
Incomplete outcome data (attrition bias)	Low risk	No loss to follow-up. 10 participant dropouts during study, equally distributed across the study arms

Rahe-Meyer 2016 (Continued)

All outcomes

Selective reporting (reporting bias)	Low risk	All prespecified outcomes have been reported in the Results
Other bias	High risk	Two study authors were employees of the funding company during the study. No statement of data ownership or publication decision making and responsibility.

Ranucci 2015

Methods	Randomised controlled, single-centre trial. Randomly assigned before surgery Calculated sample size: based on the primary endpoint and calculated by reference to own hospital's transfusion data for the past 5 years
Participants	Conditions: Elective, complex, cardiac surgery Inclusion criteria: Age > 18 years, planned complex cardiac surgery operation with expected duration of 90 minutes or longer and presence of 1 of the following risk factors: age > 65 years, non-elective surgery, serum creatinine level > 1.36 g/dl, redo surgery Exclusion criteria: Participation in another randomised controlled trial, refusal to participate, emergency surgery, known congenital or acquired coagulopathy, baseline antithrombin levels < 80%, known auto-immune disorder, pregnancy, major antiplatelet agents not discontinued within 5 days of surgery, pre-operative haematocrit value < 35% and a body surface area < 1.7m ² Number of participants enrolled: 119 Number of participants randomly assigned to the intervention: 58/58 Number of participants included in the outcome analysed: 58/58
Interventions	Intervention: Fibrinogen concentrate with dose determined by the mean clot firmness test on the FIBTEM) Comparator: Normal (0.9%) saline Additional treatments received whilst in the trial: Intraoperative dose of tranexamic acid and protamine
Outcomes	Primary: <ul style="list-style-type: none">Avoidance of allogeneic blood products (RBCs, FFP and platelets) transfusion during hospital stay up to 30 days Secondary: <ul style="list-style-type: none">Reduction in allogeneic blood products (RBCs, FFP and platelets) transfusions to 30 daysMassive blood transfusion within the first post-operatively 24 hoursNumber of participants experiencing blood transfusion of 7 or more RBC units within the first post-operative 24 hoursBleeding (chest drain output) within the first post-operative 12 hoursOperative and 30-day mortalityThromboembolic complications in-hospital or within 30 post-operative days Duration of follow-up: day 30

Ranucci 2015 (Continued)

Notes

Population nationality: Italian

Correspondence: cardioanesthesia@virgilio.it

Clinical trial registration number: Prospectively registered (7 November 2011): NCT01471730

Sources of funding: Funded by

- local research funds of the IRCCS* Policlinico San Donato
- trial drugs and the ROTEM reagents were provided by CSL Behring free of charge

Dates the trial was conducted: 30 November 2011 through to 10 December 2014

Declarations of interest (among primary researchers): 2 authors (DK and GH) were employed by the funding company.

* IRCCS translates as an "institution of in-patient care in a scientific nature", it is a hospital in which activities of clinical and translational research are carried out alongside the organization and management of health services, and the provision of highly specialized hospital care. Such centres have administrative autonomy and a different legal standing to non-IRCCS institutions

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	State computer-generation of random numbers
Allocation concealment (selection bias)	High risk	The trial used "sealed, opaque numbered envelopes" specially prepared for the trial, but they did not state whether the envelopes were sequentially-numbered, hence a high risk of bias grading
Blinding of participants (performance bias)	Low risk	Participants were anaesthetised when they received their trial treatment
Blinding of study personnel (performance bias)	Low risk	The study reports that an unblinded biologist prepared the intervention and ran the ROTEM tests: the anaesthetists, surgical staff, and clinical staff in the ICU all blinded to treatment allocation
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	There was no direct comment in paper about outcome assessors, although the database staff were blinded to treatment allocation
Incomplete outcome data (attrition bias) All outcomes	Low risk	The study reports no participants were lost to follow-up
Selective reporting (reporting bias)	Low risk	Trial protocol is available on ClinicalTrials.gov and all listed outcomes are reported in the study
Other bias	Unclear risk	Two study authors disclose personal fees from the funding company but declare these as received outside this study. The study declares full author access to data and decision to publish, although it is unclear as to the ownership of the data.

Rasche 1982

Methods

Randomised controlled, single-centre trial

Rasche 1982 (Continued)

Randomly assigned within 5 days of hospital admission

Calculated sample size: Not stated

Participants	<p>Conditions: Acute leukaemia</p> <p>Inclusion criteria: Acute leukaemia patients aged 16 - 70 yrs with a first or relapse presentation of the primary illness</p> <p>Exclusion criteria: Evidence of cerebral mass haemorrhage at admission, platelets $\leq 50,000/\mu\text{L}$, factor XIII $\leq 60\%$ of normal activity</p> <p>Number of participants enrolled: 179</p> <p>Number of participants randomly assigned to the intervention: 29/31</p> <p>Number of participants included in the outcome analyses: 29/31</p>
Interventions	<p>Intervention: Factor XIII concentrate (1000 units) on day 1 of treatment and a subsequent daily dose of 500 units in addition to normal care protocols</p> <p>Comparator: Normal care protocols</p> <p>Additional treatments received whilst in the trial: Normal care protocols</p>
Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> Bleeding complications <p>Secondary:</p> <ul style="list-style-type: none"> Maximum body temperature Platelet and red blood cell transfusion requirement Weekly full blood counts <p>Duration of follow-up: Not stated</p>
Notes	<p>Population nationality: German</p> <p>Correspondence: none stated</p> <p>Clinical trial registration number: None stated</p> <p>Sources of funding: None stated</p> <p>Dates the trial was conducted: 15 February 1978 to 31 July 1980</p> <p>Declarations of interest (among primary researchers): None stated</p> <p>The paper was translated from German for inclusion in this review</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information was reported from which to make an assessment of this bias (paper translated from German for this review)
Allocation concealment (selection bias)	Unclear risk	Insufficient information was reported from which to make an assessment of this bias (paper translated from German for this review).

Rasche 1982 (Continued)

Blinding of participants (performance bias)	Unclear risk	No details were provided as to whether anyone involved in the trial was blinded to treatment allocation
Blinding of study personnel (performance bias)	Unclear risk	No details were provided as to whether anyone involved in the trial was blinded to treatment allocation
Blinding of outcome assessment (detection bias) All outcomes	Low risk	No details were provided as to whether anyone involved in the trial was blinded to treatment allocation, but whether or not outcome assessors were blinded to treatment allocation is unlikely to have an impact on the reported outcomes (laboratory data or disease course)
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	It would seem from the study tables that all randomised participants were included in the analyses of outcomes, but it is difficult to confirm this from the text
Selective reporting (reporting bias)	Low risk	All prespecified outcomes have been reported in the Results
Other bias	Unclear risk	Sponsored by intervention product manufacturer. Insufficient information available to determine risk of bias.

Sabate 2016

Methods	Randomised, multicentre, double-blind, placebo-controlled trial Randomly assigned at: Immediately prior to surgery Calculated sample size: Power analysis
Participants	Condition: Scheduled for liver transplantation surgery Inclusion criteria: Adults aged 18 – 80 years who were scheduled for liver transplantation Exclusion criteria: History of allergic reaction to fibrinogen concentrate, known history of thromboembolic events in the last 30 days, known or suspected pregnancy, previous randomisation in same trial, known presence of congenital bleeding disorder, and aspirin or warfarin therapy. Excluded if indication for transplantation was familial polyneuropathy and living donors; acute liver failure, biliary cirrhosis and sclerosing cholangitis, and non-heart-beating donors. Patients with a plasma fibrinogen concentration > 2.9g/L in the 24 hours prior to surgery Number of participants enrolled: 99 Number of participants randomly assigned to the intervention/control: 51/48 Number of participants who received the treatment/control and were included in the outcome analyses: 48/44
Interventions	Intervention: Fibrinogen concentrate (dose calculated to reach a target plasma concentration of 2.9g/L fibrinogen) Comparator: 0.9% sodium chloride solution Additional treatments received whilst in the trial: Crystalloids to maintain blood volume, colloids were sometimes used; bicarbonate, iv calcium. Blood transfusion according to SOP: Hb > 80; platelets > 50, FFP 2 units if bleeding not corrected by RBC units and platelets; tranexamic acid (500 mg) if lysis > 15% at 60 mins on ROTEM
Outcomes	Primary:

Sabate 2016 (Continued)

- Percentage of participants requiring transfusion of RBC units s during the LT procedure

Secondary:

- Percentages of participants requiring FFP, fibrinogen concentrate and platelets during surgery and within 24 hours within each group
- Number of units of RBC units s, FFP and platelets transfused during and within 24 hours of surgery
- Grams of fibrinogen administered during and within 24 hours of surgery
- Number of patients and grams of tranexamic acid administered during surgery;
- Mortality during the operative period until hospital discharge
- Liver graft survival
- Thrombotic complications of any type and cause during the hospital stay or 30 days after surgery

Duration of follow-up: Primary outcome at end of surgery; other outcomes up to hospital discharge

Notes

Population nationality: Spanish

Correspondence: asabatep@bellvitgehospital.cat and asabatep@ub.edu

Clinical trial registration number: EudraCT, 2010-024584 and NCT01539057

Sources of funding: This study was funded by the Spanish Department of Health through the Institute of Heath Carlos III Grant

Dates the trial was conducted: 8 August 2012, to 25 February 2014

Declarations of interest (among primary researchers): None declared

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	The randomisation sequence was created using a computer-generated random list, which was then stratified according to whether the baseline haemoglobin concentration was < 95 or ≥ 95g/L and by centre (1:1 ratio, in blocks of multiples of 2 U)
Allocation concealment (selection bias)	Low risk	Sealed consecutively-numbered kits made up centrally and distributed
Blinding of participants (performance bias)	Low risk	Participants under general anaesthesia
Blinding of study personnel (performance bias)	Low risk	Only independent nurse who prepared intervention aware of allocation; no other involvement in study. Specific administration kits so treating teams were blinded
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Fibrinogen levels were unblinded during transplant procedure so potential for staff to determine which participant received the fibrinogen concentrate
Incomplete outcome data (attrition bias) All outcomes	Low risk	Although the case record form was lost in 7 participants, these losses were equally distributed across study arms and the participants did not receive the trial intervention
Selective reporting (reporting bias)	Low risk	All prespecified outcomes have been reported in the Results

Sabate 2016 (Continued)

Other bias	Unclear risk	Power analysis identified that 66 participants were required in each group, hence insufficient participants recruited
------------	--------------	---

Sadeghi 2014

Methods	Randomised controlled, single-centre trial Randomly assigned at: Not stated Calculated sample size: Not stated
Participants	Conditions: Elective revascularisation cardiac surgery Inclusion criteria: Scheduled to undergo first elective coronary artery bypass surgery Exclusion criteria: Haematologic or liver disease, previously diagnosed, uncontrolled diabetes or dependence on insulin, pregnancy, unstable angina, high serum creatinine > 130 mol/L, left ventricular ejection fraction < 35%, serum fibrinogen > 3.5 g/L Number of participants enrolled: not stated Number of participants randomly assigned to the intervention: 30/30 Number of participants included in the outcome analyses: 30/30
Interventions	Intervention: Fibrinogen, 1 g dissolved in 50 mL normal saline for 15 minutes, administered 30 minutes prior to anaesthesia induction Comparator: 1 g saline, administered 30 minutes prior to anaesthesia induction Additional treatments received whilst in the trial: None stated
Outcomes	Primary <ul style="list-style-type: none"> Volume of post-operative haemorrhage 0, 12 and 24 hours after surgery Secondary <ul style="list-style-type: none"> Mean number of red cell infusions in the group during the 1st 24 hours after surgery Blood lost within the first 12 hours after surgery Mean prothrombin time 24 hours after surgery Haemoglobin concentration, platelet count, partial thromboplastin time, prothrombin time and plasma fibrinogen concentrations, 24 hours after surgery Thrombotic adverse events during first 72 hours after surgery Duration of follow-up: to 72 hours after surgery
Notes	Population nationality: Iranian Correspondence: ali.movafeghi@gmail.com Clinical trial registration number: Retrospectively randomised (07 June 2011) on the Iranian Registry of Clinical Trials, registry number IRCT201011085140N1 Sources of funding: Tehran University of Medical Sciences Dates the trial was conducted: 25 January 2009 to 29 June 2010 Declarations of interest (among primary researchers): The authors declared no conflicts of interest.

Sadeghi 2014 (Continued)

Study translated from Portuguese into English using Google translate 16 July 2015

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	A computer-generated list was used
Allocation concealment (selection bias)	Unclear risk	Insufficient information was reported from which to make an assessment of this bias (paper translated from Portuguese for this review)
Blinding of participants (performance bias)	Low risk	The study reports that all the plastic container vials in the trial were covered by tissue paper and the intervention or placebo were not visible
Blinding of study personnel (performance bias)	Low risk	Anaesthetists not involved in trial made up drugs. The study reports that "all necessary medicines for the trial were prepared by a non-anaesthetist" and that all the plastic container vials in the trial were covered by tissue paper and the intervention or placebo were not visible
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information was reported from which to make an assessment of this bias (paper translated from Portuguese for this review)
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing data: study reports that all participants were included in the analysis
Selective reporting (reporting bias)	Low risk	Trial registration details are available and the outcomes listed on the form are reported on in the study
Other bias	Unclear risk	Insufficient information was reported from which to make an assessment of this bias (paper translated from Portuguese for this review)

Shirahata 1990

Methods	Randomised controlled, single-centre trial Randomly assigned on Admission to neonatal ICU Calculated sample size: Not stated
Participants	Conditions: Prematurity of birth Inclusion criteria: Either gestational age under 37 weeks or birth weight under 2 kg if artificially ventilated; or gestational age under 33 weeks or birth weight under 1.5 kg if not ventilated Exclusion criteria: Treatment with phenobarbital, pancuronium and ethamsylate; anomalies of the central nervous system, death within 6 hours of delivery Number of participants enrolled: not stated Number of participants randomly assigned to the intervention: 30/28 Number of participants included in the outcome analyses: 30/28
Interventions	Intervention: FXIII at a dose of 70 to 100 units administered slowly for at least 10 minutes intravenously within 6 hours of delivery Comparator: "non-treated group"

Shirahata 1990 (Continued)

Additional treatments received whilst in the trial: None stated

Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> Not specifically stated. However the objective of the trial was to see if administration of a FXIII concentrate could prevent severe intraventricular haemorrhage <p>Secondary:</p> <ul style="list-style-type: none"> Not specifically stated, the study lists a number of observation items <p>Duration of follow-up: up to 7 days after delivery</p>
Notes	<p>Population nationality: Japanese</p> <p>Correspondence: A Shirahata MD, Department of Paediatrics, University of Occupational and Environmental Health, Japan, 1-1 Iseigaoka, Nishi-ku, Yahata, Kitakyushu, Japan</p> <p>Clinical trial registration number: not registered</p> <p>Sources of funding: Not stated.</p> <p>Dates the trial was conducted: January 1986 to October 1987</p> <p>Declarations of interest (among primary researchers): None reported</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information was reported from which to make an assessment of this bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information was reported from which to make an assessment of this bias
Blinding of participants (performance bias)	Low risk	No information on blinding was reported. Participants were neonates and as treatment (intervention or no treatment) was administered by a clinician within 6 hours of delivery we do not believe parents' knowledge of allocation would elevate the risk of performance bias
Blinding of study personnel (performance bias)	Low risk	No information on blinding was reported but we felt that the outcome and the outcome measurement are not likely to be influenced by lack of blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	No information on blinding was reported but we felt that the outcome and the outcome measurement are not likely to be influenced by lack of blinding
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	The trial does not address issue of loss to follow-up or dropouts
Selective reporting (reporting bias)	Unclear risk	No outcomes formally stated: Insufficient information was reported from which to make an assessment of this bias
Other bias	Unclear risk	Insufficient information was reported from which to make an assessment of this bias.

Soleimani 2016

Methods	<p>Randomised controlled, single-centre trial.</p> <p>Randomly assigned at: Pre-operative visit</p> <p>Calculated sample size: Power analysis</p>
Participants	<p>Condition: Clinical and laboratory evidence of benign prostatic hyperplasia</p> <p>Inclusion criteria: Benign prostatic hyperplasia due to undergo transurethral resection of the prostate</p> <p>Exclusion criteria: Previous history of prostate cancer or prostatectomy; taking finasteride or any anti-coagulant agent; having a known blood abnormality or coagulopathy; a history of thromboembolism; chronic liver disease; chronic kidney disease with a serum creatinine level of > 2mg/dL; and a preoperative plasma fibrinogen level of ≥ 400mg/dL</p> <p>Number of participants enrolled: 72</p> <p>Number of participants randomly assigned to the intervention/control: 36/36</p> <p>Number of participants who received the treatment/control and were included in the outcome analyses: 31/29</p>
Interventions	<p>Intervention: Fibrinogen 2 g 15 - 30 mins pre-operatively</p> <p>Comparator: 0.9% sodium chloride solution</p> <p>Additional treatments received whilst in the trial: Transurethral resection of the prostate surgery including use of irrigating fluid</p>
Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> Amount of bleeding during and after surgery <p>Secondary:</p> <ul style="list-style-type: none"> Amount of blood transfusion required Volume of irrigating fluid used Operative time Safety <p>Duration of follow-up: Primary outcome up to 12 hours post-operatively, other outcomes up to 1 week post-operatively</p>
Notes	<p>Population nationality: Iranian</p> <p>Correspondence: info@safarinejad.com</p> <p>Clinical trial registration number: IRCT201305097457N13</p> <p>Sources of funding: None declared</p> <p>Dates the trial was conducted: August 2014 to September 2015</p> <p>Declarations of interest (among primary researchers): The authors state that they have no conflict of interest</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Sequence generated by associate anaesthesiologist during the preoperative visit by assigning random numbers from a random-number table

Soleimani 2016 (Continued)

Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of participants (performance bias)	Low risk	Participants not informed of nature of intervention, given while awake pre-operatively. Intervention given by an anesthesiology technician who was not involved further in the study
Blinding of study personnel (performance bias)	Low risk	Participants not informed of nature of intervention, given while awake pre-operatively. Intervention given by an anaesthesiology technician who was not involved further in the study
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not stated; the study does not identify whether assessors would have access to fibrinogen level results
Incomplete outcome data (attrition bias) All outcomes	Low risk	Some loss to follow-up and loss of participant data although this is distributed equally across study groups
Selective reporting (reporting bias)	Low risk	All prespecified outcomes have been reported in the Results
Other bias	Unclear risk	Insufficient information was reported from which to make an assessment of this bias

Tanaka 2014

Methods	Randomised controlled, single-centre trial Randomly assigned at: not stated Calculated sample size: None stated - interim analysis
Participants	Conditions: Elective valve replacement or repair surgery Inclusion criteria: Elective cardiopulmonary bypass procedures via median sternotomy for a single valve replacement or repair, combined mitral and aortic valve replacements, or combined valve replacement and cardiac bypass graft surgery. If haemostatic condition of surgical field either moderate bleeding (controllable with applied pressure) or severe (multiple diffuse bleeding sites) then randomly assigned to trial intervention Exclusion criteria: Emergency surgery, redo valve procedure, age (< 18 or > 85 years), any known congenital or acquired bleeding disorders, dialysis-dependent renal failure, liver dysfunction (more than 2-fold increase in liver enzymes), pregnancy or nursing, and intake of clopidogrel or coumadin within 5 days of surgery Number of participants enrolled: 51 Number of participants randomly assigned to the intervention: 10/10 Number of participants included in the outcome analyses: 10/10
Interventions	Intervention: Single dose (4 g) of fibrinogen (RiaSTAP®) given within 30 minutes of intervention decision Comparator: 1 unit of apheresis platelets (median volume 230 mL) within 30 minutes of intervention decision Additional treatments received whilst in the trial: Additional transfusions (after trial intervention) in the presence of bleeding
Outcomes	Primary:

Tanaka 2014 (Continued)

- Haemostatic condition in the surgical field after receipt of the trial intervention
- Haemostatic blood product use over the first 24 hours
- Percentage of participants with thromboembolic events through to Day 28 visit (up to 6 - 8 weeks post-operatively)
- Mortality through to Day 28 visit (up to 6 - 8 weeks post-operatively)

Secondary:

- 12-hour volumes of chest tube drainage
- Re-exploration within 24 hours of surgery
- Troponin levels at 24 hours after surgery
- Length of ICU stay (hours)
- Length of hospital stay (days)

Duration of follow-up: Up to 6 to 8 weeks post-operatively

Notes

Population nationality: American.
Correspondence: tanakak@upmc.edu

Clinical trial registration number: Prospectively registered (24 January 2011): NCT01283321

Sources of funding: Funding from CSL Behring; single author received a consulting fee from ROTEM, Inc

Dates the trial was conducted: January 2011 to to May 2013

Declarations of interest (among primary researchers): The contact author (KT) has received a consulting fee from the company supplying analytic equipment. No other author declared any conflicts of interest

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Trial states randomised but provides no further information
Allocation concealment (selection bias)	High risk	Closed-envelope method. As no further details given, judged to be at high risk of bias
Blinding of participants (performance bias)	Low risk	Participants were under general anaesthesia at time of intervention
Blinding of study personnel (performance bias)	High risk	Open-label trial with adequacy of surgical field haemostasis reported as a subjective outcome
Blinding of outcome assessment (detection bias) All outcomes	High risk	Open-label trial with adequacy of surgical field haemostasis reported as a subjective outcome
Incomplete outcome data (attrition bias) All outcomes	Low risk	Complete follow-up on all participants
Selective reporting (reporting bias)	Low risk	All stated outcomes reported appropriately. Protocol registered at www.clinicaltrials.gov
Other bias	Unclear risk	This is an interim analysis of this study and does not include a power calculation or declaration of intended numbers of participants, hence it is unclear

Tanaka 2014 (Continued)

whether the participant numbers are sufficient to determine the efficacy of the interventions

Turner 1981

Methods	Randomised controlled, single-centre trial Randomly assigned at: Not stated Calculated sample size: Not stated
Participants	<p>Conditions: Prematurity of birth</p> <p>Inclusion criteria: Infants were screened for inclusion if they:</p> <ol style="list-style-type: none"> weighed < 1500 g and were < 34 weeks gestation were < 37 weeks gestation, weighing between 1500 - 2000 g, and with idiopathic respiratory distress syndrome needing umbilical arterial catheterisation for ABG; were mature infants with severe intrapartum asphyxia + an Apgar of < 5 at 5 minutes after birth <p>To proceed to randomisation, they had to show abnormal haemostasis (defined as: prothrombin time > 28 secs, partial thromboplastin time > 100 secs; fibrinogen < 1 g/L; platelet count < 100) within 48 hours of birth</p> <p>Exclusion criteria: Not stated</p> <p>Number of participants enrolled: 244 infants across the three groups above</p> <p>Number of participants randomly assigned to the intervention: 39/39</p> <p>Number of participants included in the outcome analyses: 39/39</p>
Interventions	<p>Intervention: Prothrombin complex concentrate (PCC)* at a dose of 1 mL/kg for prolonged prothrombin time (if from groups 1 and 2 above) or at a standard dose of 2 mL/kg (if from group 3 above)</p> <p>*PCC was from Scottish National Blood Transfusion Service Protein Fractionation Centre containing approximately 25 times the normal plasma concentration of coagulation factors XI, VII, IX and X and 10 u/mL heparin. Low fibrinogen treated with cryoprecipitate, single-donor units containing approximately 200 mg of fibrinogen and 80 units of factor VIII. Thrombocytopenia treated with platelets, single-donor units containing about 0.7×10^{11} platelets. Dosing until coagulation screen demonstrated sustained therapy response</p> <p>Comparator: Usual care protocols: dosing until coagulation screen demonstrated sustained therapy response</p> <p>Additional treatments received whilst in the trial: All participants in this trial received Vitamin K supplementation (dose not stated). Heparin and clotting factors given for disseminated intravascular coagulation. Red blood cell transfusion as required for all participants</p>
Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> Mortality (all-cause) incidence of intraventricular haemorrhage (post-mortem) <p>Secondary:</p> <ul style="list-style-type: none"> Correction of haemostatic defects <p>Duration of follow-up: at least 12 (but up to 36) months from birth</p>
Notes	<p>Population nationality: British</p> <p>Correspondence: No longer available.</p>

Turner 1981 (Continued)

Clinical trial registration number: not registered.

Sources of funding: Not stated.

Dates the trial was conducted: Not stated.

Declarations of interest (among primary researchers): Not stated.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Closed-envelope system arranged by an impartial observer (statistician author on the paper)
Allocation concealment (selection bias)	Unclear risk	Closed-envelope system arranged by an impartial observer (statistician author on the paper)
Blinding of participants (performance bias)	Low risk	Neonatal participants
Blinding of study personnel (performance bias)	High risk	No comments were made in the study report. However, we know that the intervention is notably visually different from the control, so this would probably mean trial personnel would be aware of treatment received.
Blinding of outcome assessment (detection bias) All outcomes	High risk	"Closed envelope system arranged by an impartial observer". However we know that the intervention is notably visually and functionally different from the control and requires more tests involved for response, this would probably mean outcome assessors would be aware of treatment received.
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants were included in the Results.
Selective reporting (reporting bias)	Low risk	The study reports the results of all outcomes predefined in the Methods section.
Other bias	Unclear risk	Source of funding unclear, no declaration of author interests.

Wikkelso 2015

Methods	Parallel randomised controlled, multicentre trial Randomly assigned at: Not stated Calculated sample size: Sample size of 245 patients to demonstrate a reduction in RR of 33% of the proportion of participants receiving postpartum transfusion (with a power of 80% difference, a 2-sided type 1 error of 0.05 and up to 15% dropout)
Participants	Conditions: Postpartum haemorrhage Inclusion criteria: Aged > 18 years with postpartum haemorrhage defined as bleeding from the uterus or birth canal or both, within 24 hours postpartum; caesarean section with an estimated perioperative blood loss > 1 litre or vaginal delivery with either estimated blood loss > 0.5 litre and intended manual removal of placenta or estimated blood loss > 1 litre and intended manual exploration of the uterus because of continuous bleeding after delivery of placenta Blood loss up to 0.5 litre was assessed by visual inspection of the absorbent delivery pad and thereafter determined by weighing of drapes and pads as described in Danish national guidelines

Wikkelso 2015 (Continued)

Exclusion criteria: Known inherited coagulation deficiencies; antenatal thrombotic treatment;

pre-pregnancy weight < 45 kg; refusal to receive a blood transfusion

Number of participants enrolled: 1967

Number of participants randomly assigned to the intervention: 124/125

Number of participants included in the outcome analyses: 123/121 women respectively included in the ITT analysis; 120/119 respectively in the per-protocol analysis

Interventions	<p>Intervention: Fixed dose of 2 g of fibrinogen concentrate dispensed using a syringe pump infusion over 20 minutes in 100 mL sterile water (mean (standard deviation) dose = 26 (SD 4.2) mg/kg)</p> <p>This intervention was applied in a pragmatic early pre-emptive treatment regimen giving a fixed dose to all women, irrespective of body weight and not guided by measurement of fibrinogen concentration or other haemostatic measures</p> <p>Comparator: 100 mL of isotonic saline (mean (SD) dose = 1.2 (0.2) mg/kg).</p> <p>Additional treatments received whilst in the trial: All participants could receive additional interventions, including tranexamic acid, hydroxyethyl starch and iv fluids at the discretion of their attending physician and supported by regional guidelines</p>	
Outcomes	<p>Primary:</p> <ul style="list-style-type: none">• Red blood cell transfusion during a 6-week follow-up period postpartum <p>Secondary:</p> <ul style="list-style-type: none">• Severe postpartum haemorrhage, during hospital stay or until 6 weeks post-intervention• Estimated blood loss, during hospital stay or until 6 weeks post-intervention• Total amount of blood transfused, during hospital stay or until 6 weeks post-intervention• Recurrence of bleeding, up to 6 weeks post-intervention• Haemoglobin level < 5.8 g/dL, during hospital stay or until 6 weeks post-intervention• Side-effects including thromboembolic complications, up to 6 weeks post-intervention• Composite outcome of "severe postpartum haemorrhage" defined by Charbit 2007 as a decrease in haemoglobin of > 40 g/dL, transfusion of > 4 units of red blood cells, haemostatic intervention (angiographic embolisation, surgical arterial ligation or hysterectomy) or maternal death <p>Duration of follow-up: up to 6 weeks post-intervention</p>	
Notes	<p>Population nationality: Danish</p> <p>Correspondence: wikkelse@gmail.com</p> <p>Clinical trial registration number: Prospectively registered (20 May 2011): NCT01359878</p> <p>Sources of funding: The Danish Council for Independent Research Medical Sciences Grant</p> <p>Dates the trial was conducted: 30 May 2011 to 11 July 2013</p> <p>Declarations of interest (among primary researchers): None declared</p>	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random sequence generation occurred through computer-generation with a fixed block size that was stratified by centre (trial protocol)

Wikkelso 2015 (Continued)

Allocation concealment (selection bias)	High risk	Although the authors report that allocation concealment was secured through the use of "sealed opaque numbered envelopes", for this domain to be rated at of low risk of bias, the envelopes have to be sequentially numbered
Blinding of participants (performance bias)	Low risk	<p>The intervention and placebo were both yellow-coloured and both provided in 50 mL opaque syringes (the placebo was dyed to near-match the colour of the fibrinogen concentrate).</p> <p>The study reports that all included in the trial were blinded to treatment allocation.</p> <p>The success of blinding to treatment allocation was measured by asking during the follow-up period if "they had any hint of the allocated treatment and if so why". The full results are reported as supplementary data. Only 1 patient had discovered what treatment she had received due to deliberate unblinding for an adverse event: all other women were unaware of their treatment allocation</p>
Blinding of study personnel (performance bias)	Low risk	<p>The intervention and placebo were both yellow-coloured and both provided in 50 mL opaque syringes (the placebo was dyed to near-match the colour of the fibrinogen concentrate).</p> <p>The study reports that all included in the trial (trial investigators and care providers) were blinded to treatment allocation with anaesthetic personnel not involved in patient care randomising the patient and dispensing trial medication.</p> <p>The success of blinding to treatment allocation was measured by asking in the operating theatre following trial intervention administration if "they had any hint of the allocated treatment and if so why". The full results are reported as supplementary data. 94% (220) of the anaesthetists who evaluated blinding had no idea of which treatment was provided</p>
Blinding of outcome assessment (detection bias) All outcomes	Low risk	<p>The intervention and placebo were both yellow-coloured and both provided in 50 mL opaque syringes (the placebo was dyed to near-match the colour of the fibrinogen concentrate).</p> <p>The study reports that all included in the trial were blinded to treatment allocation</p>
Incomplete outcome data (attrition bias) All outcomes	Low risk	<p>The study reports both the ITT and the per-protocol analyses.</p> <p>All participants (N = 123 of 124 fibrinogen group; N = 121 of 125 placebo group) who were fully and properly consented were included in the ITT analysis for all outcomes reported. 2 participants (1 from each group) received their trial medication, but were incorrectly consented and not included in any ITT outcome analyses. The reason for the lack of full and properly attained consent are documented in the supplementary data for this trial and are similar between the groups</p> <p>The secondary outcome 'decrease in haemoglobin > 40g litre⁻¹' was missing in 148 (61%) cases in the ITT analysis and there is no indication of the distribution of these cases between the intervention and control arms of the study</p> <p>Transexamic acid, that might enhance the action of the fibrinogen intervention, was given in a comparable proportion of participants in each study arm prior to the study intervention (intervention 33%, placebo 30%) and after the study intervention (intervention 7%, placebo 6%). No significant difference in the rate between groups was reported (P > 0.05)</p>

Wikkelsø 2015 (Continued)

Selective reporting (reporting bias)	Low risk	The trial protocol is available and all of the trial's prespecified outcomes (primary and secondary) have been reported in the prespecified way
Other bias	Unclear risk	ITT analysis included all randomised participants except the 5 (1 fibrinogen and 4 placebo arm) for whom full standing consent was not received. 2 of these 5 participants (1 from each arm) received the intervention as allocated ITT data have been published in this review. Per-protocol analysis data were reported in the supplementary data that omitted data for participants meeting the exclusion criteria or who did not receive the full intervention dose. These women were analysed as treated. Outcomes were comparable to the ITT analysis with no significant differences found in primary or secondary outcomes

A&E = Accident and Emergency department; ASA = American Society of Anesthesiologists; BMI = body mass index; CT = computed tomography; ECG = echocardiogram; FFP = fresh frozen plasma; FIBTEM = a ready-to-use ROTEM® system reagent for use with citrated whole blood and assesses the clot firmness of the fibrin clot; Hb = haemoglobin; HDU = high dependency unit; ICU = intensive care unit; IE = intrinsic efficacy (The property of a drug that determines the amount of biological effect produced per unit of drug-receptor complex formed); IgA = immunoglobulin A; INR = international normalised ratio iv = intravenous; PCC = prothrombin complex concentrate; RBC = red blood cell; RCT = randomised controlled trial; ROTEM = rotational thromboelastometry; SOFA score = Sepsis-Related Organ Failure Assessment score; TEG = thromboelastogram.

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
ACTRN12615001293550p 2015	This study was not randomised
Boulis 1999	The patients were a specific group, on warfarin (as a prevention for arterial or venous thromboembolic disease) who received the pro-coagulant haemostatic factor for anticoagulation reversal
Curry 2015	The study intervention was cryoprecipitate rather than an injectable haemostatic clotting factor
Demeyere 2010	The patients were a specific group, on warfarin (as a prevention for arterial or venous thromboembolic disease) who received the pro-coagulant haemostatic factor for anticoagulation reversal
Garrigue 2017	The study intervention was lyophilised plasma rather than an injectable haemostatic clotting factor
Goldstein 2015	The patients were a specific group, on warfarin (as a prevention for arterial or venous thromboembolic disease) who received the pro-coagulant haemostatic factor for anticoagulation reversal
Masoumi 2016	The intervention in this study is not appropriate for this review
Miyata 2014	The intervention in this study is not appropriate for this review
NCT01706159	The study population are not actively bleeding
NCT02542306	The study design is not randomised
NCT02736812	The intervention in this study (plasma) is not appropriate for this review
Papakitsos 2014	The patients were a specific group, on warfarin (as a prevention for arterial or venous thromboembolic disease) who received the pro-coagulant haemostatic factor for anticoagulation reversal
Qiu 2016	The intervention in this study is not appropriate for this review

Study	Reason for exclusion
Sarode 2013	The patients were a specific group, on warfarin (as a prevention for arterial or venous thromboembolic disease) who received the pro-coagulant haemostatic factor for anticoagulation reversal
Scherer 1994	Not a randomised controlled trial
TOP-CLOT Trial	The study was terminated after randomisation of two participants due to feasibility issues
UMIN000022194	The study design is not randomised
UMIN000022666	The study design is not randomised
van Aart 2006	The patients were a specific group, on warfarin (as a prevention for arterial or venous thromboembolic disease) who received the pro-coagulant haemostatic Factor for anticoagulation reversal

Characteristics of studies awaiting assessment *[ordered by study ID]*

[Siemens 2017](#)

Methods	Randomised controlled, single-centre trial; phase 1B/2A, open-label dose escalation study Randomly assigned at: following ROTEM performed 1 hour pre-end bypass Calculated sample size: Not stated
Participants	<p>Conditions: Neonates and infants (below 12 kg) who are at risk of mediastinal bleeding following cardiopulmonary bypass surgery for congenital heart disease</p> <p>Inclusion criteria:</p> <p>Congenital heart disease requiring non-emergency* surgery on cardiopulmonary bypass; age range: > 36 weeks corrected gestation; Weight 2.5 – 12 kg; Informed consent to participate</p> <p>*Non-emergency is defined as surgery that can be delayed > 24 hours following diagnosis of congenital heart disease</p> <p>Exclusion criteria:</p> <p>Known pre-existing inherited coagulopathy; known pre-existing inherited thrombophilia; recent, acute (within previous 2 weeks) thrombosis in a major vessel or thrombotic-related major complications (as defined in sections 2.43 and 7.3); administration of antiplatelet agents (e.g. aspirin) < 48 hrs prior to surgery; known hypersensitivity / allergy to study drug or similar products; history of anaphylaxis; enrolment in another clinical trial in the previous 3 months; parent/guardian unable to provide informed consent (this can include insufficient understanding of the trial, as judged by the clinician taking consent); major co-morbidity likely to increase risk of mortality from surgical procedure; significant renal/liver impairment within 2 days of planned surgery (creatinine > 2x ULN, ALT > 2x ULN)</p>
Interventions	<p>Participants were randomised to receive fibrinogen concentrate: placebo (2:1 ratio) if FIBTEM mean clot firmness ≤ 6 mm (physiological 7 – 24 mm)</p> <p>If FIBTEM mean clot firmness ≥ 7 mm participants entered the monitoring cohort</p> <p>Individualised fibrinogen concentrate dose calculation included participant weight, bypass circuit volume, haematocrit and intraoperative FIBTEM mean clot firmness</p> <p>Desired mean clot firmness was 8 – 13 mm and fibrinogen levels 1.5 – 2.5 g/L</p> <p>Comparator: Solution for injection</p>

Siemens 2017 (Continued)

Additional treatments received whilst in the trial: Not stated

Outcomes

Objective: To determine the dose of intraoperative human fibrinogen concentrate required to achieve physiological levels of fibrin polymerisation of 8 to 13 mm as measured by the ROTEM measure of fibrin-based clotting: FIBTEM mean clot firmness (equating to plasma fibrinogen concentrations of 1.5 to 2.5 g/L), immediately prior to separation from cardiopulmonary bypass in neonates and children < 12 kg

(a) To provide preliminary efficacy and safety data following human fibrinogen concentrate administration

(b) To document ROTEM profiles intra- and post-operatively

Primary Outcome:

- Fibrinogen concentration and FIBTEM mean clot firmness measured within 5 minutes of completion of fibrinogen concentrate administration

Secondary Outcomes:

Efficacy

- Mediastinal drain losses in first 24 hours after PICU admission
- Requirement for delayed sternal closure due to clinical bleeding/tamponade
- Requirement for ancillary blood transfusions in first 24 hours after PICU admission
- Use of intra- and post-operative ancillary clotting products (as per transfusion algorithm)
- Fibrinogen levels and ROTEM variables T4 – T6

Safety (incidence of major thrombotic event/thromboembolic-associated complications)

- Surgical shunt occlusion
- Imaging evidence of intracardiac/major vessel thrombosis up to discharge or day 30 post-operative
- Stroke from sinovenous thrombosis or arterial ischaemia of cardioembolic origin
- Pulmonary embolism
- Superior vena cava syndrome
- Thrombotic vessel obstruction requiring active treatment with thrombolysis/mechanical intervention (open thrombectomy, cardiac catheter)
- Cardiopulmonary arrest associated with thrombosis
- Allergic/hypersensitivity reaction to study drug

All safety analyses will be up to hospital discharge or day 30 post-operative (whichever comes first)

The remaining endpoints will be evaluated up to 24 hours after PICU admission with fibrinogen levels and ROTEM variables evaluated at time points T4 - T6

T4: at PICU admission

T5: 4 hrs after PICU admission

T6: 24 hrs after PICU admission

Duration of follow-up: Up to day 30 post-operatively

Notes

Population nationality: British

Siemens 2017 (Continued)

Correspondence: Dr. Shane Tibby, shane.tibby@gstt.nhs.uk

Clinical trial registration number: Endura-CT Number: 2013-003532-68.

Sources of funding: CSL Behring Ltd

Dates the trial was conducted: Unclear

Sponsor: Guy's and St Thomas NHS Foundation Trust.

Interim data from this study have been published in abstract form only. We do not have the full data and hence this trial is awaiting assessment

FIBTEM = a ready-to-use ROTEM® system reagent for use with citrated whole blood and assesses the clot firmness of the fibrin clot, PICU = Paediatric Intensive Care Unit, ROTEM = rotational thromboelastometry,

Characteristics of ongoing studies [ordered by study ID]

ANZCTR12613001279718

Trial name or title	Prothrombinex-VF versus fresh frozen plasma for the treatment of bleeding post-cardiopulmonary bypass
Methods	Randomised controlled trial
Participants	<p>Patients undergoing cardiac surgery requiring cardiopulmonary bypass and determined to have non-surgical bleeding post-heparin reversal. Planned sample size = 200</p> <p><i>Exclusion criteria;</i></p> <ul style="list-style-type: none"> • Patients with pre-existing bleeding diatheses • Patients receiving Prothrombinex or FFP less than 5 days before cardiopulmonary bypass • Patients receiving heart or lung transplant • Patients having implantation of ventricular assist devices
Interventions	Human 3 factor prothrombin complex concentrate (Prothrombinex VF) 25 IU/kg I.V. bolus for the first-line treatment of non-surgical bleeding post-cardiopulmonary bypass versus FFP 15 mL/kg bolus iv as first-line treatment for non-surgical bleeding post-cardiopulmonary bypass
Outcomes	<p>Primary:</p> <ul style="list-style-type: none"> • The composite end point of those participants receiving either allogeneic red blood cells or platelets after 15 mL/kg FFP or Prothrombinex 25IU/kg, up to 24 hours post-operation • The total cost of treatment following randomisation, which will include, drugs, procedures, imaging, length of stay in ICU and hospital. We will also include best estimation of blood product costs during hospital stay <p>Secondary:</p> <ul style="list-style-type: none"> • Incidence of thrombotic complications defined by Classification of Hospital Acquired Diagnoses (CHADx) groups: 5.2 pulmonary embolism and 5.8 venous thrombosis/embolism (not progressing to a pulmonary embolism): assessment will be clinical and based on discharge coding • Respiratory indices (PaO₂/FiO₂ ratio: the ratio of arterial oxygen partial pressure to fractional inspired oxygen) at 6 and 12 hours post-operation • Overall mortality at 30 days • Length of stay in ICU and hospital • Chest drain output at 6 and 24 hours • ICU arrival haemoglobin • Total surgical red cell loss (Mercuriali formula) at post-operative day 5

ANZCTR12613001279718 (Continued)

Starting date	13 January 2014
Contact information	Dr Warren Pavey, Department of Anaesthesia and Pain Medicine, Royal Perth Hospital, Wellington Street, Perth WA 6001, Australia warren.pavey@health.wa.gov.au
Notes	Funding received from Western Australia State Health Research Advisory Committee and Royal Perth Hospital Date registered on trial website: 20 November 2013

ANZCTR12614000691640

Trial name or title	The administration of fibrinogen concentrate (a component of blood involved in clotting process) to be given to those women who, after delivery of babies, develop severe bleeding, to establish if this drug will prevent further blood loss or reduce the volume of blood loss
Methods	Randomised controlled trial
Participants	Women after delivery with a postpartum blood loss of more than 1000 mL and with signs of ongoing bleeding Planned sample size =245 <i>Inclusion criteria;</i> <ul style="list-style-type: none"> • Women with estimated blood loss greater than 1000 mL in the immediate postpartum period (< 4 hours postpartum) • Women with persistent postpartum haemorrhage not responding to first-line uterotonic therapy (i.e. syntocinon infusion), and manual uterine compression in whom further haemostatic intervention is being considered <i>Exclusion criteria;</i> <ul style="list-style-type: none"> • Women with coagulopathic disorder secondary to pre-existing liver disease • Women with confirmed venous thromboembolism occurring in the last 4 weeks of pregnancy • Women with confirmed allergy to fibrinogen concentrate • Woman who refuse blood products
Interventions	A single dose of fibrinogen concentrate 4 g given intravenously upfront versus placebo (saline, given intravenously)
Outcomes	Primary: <ul style="list-style-type: none"> • Efficacy: 25% reduction in total measured blood volume loss at 24 hours. To compare volume blood loss measured we will correlate volume loss by calculating starting haemoglobin, to post-bleed haemoglobin and include fluid and blood and products administered Secondary: <ul style="list-style-type: none"> • Incidence of moderate postpartum haemorrhage (blood loss between 1000 and 2000 mL) • Incidence of severe postpartum haemorrhage (blood loss > 2000 mL) • Total blood product usage over 24 hours • Activation of massive transfusion protocol within 24 hours • Need for haemostatic intervention (balloon tamponade, uterine artery embolisation, surgical arterial ligation, hysterectomy) within 24 hours • Anaemia with Hb < 80 g/L at 24 hours post-delivery at 24 hours • Admission to ICU/ department of critical care medicine within 6 months

ANZCTR12614000691640 (Continued)

- Maternal mortality at 6 months
- Infection rates (wound infection, endometritis, pneumonia) at 3 months
- Breast feeding rates at hospital discharge and at 6 weeks
- Quality of life indicators as per WHO Disability Assessment Schedule 2.0/post-traumatic stress disorder score at 6 weeks
- Anaemia at 6 weeks postpartum (Hb < 100 g/L)

Starting date	(Anticipated) 01 December 2014
Contact information	Dr Joreline van der Westhuizen, Anaesthetic Department Auckland City Hospital jayv@adhb.govt.nz
Notes	Funding source: Auckland City Hospital Date registered on trial website: 1 July 2014

EudraCT: 2010-022923-31

Trial name or title	A multicenter double-blind, placebo controlled, randomized, pilot trial to assess the efficacy of pre-hospital administration of fibrinogen concentrate in trauma patients, presumed to bleed (FI in TIC)
Methods	Randomised controlled trial
Participants	<p>Trauma patients with major bleeding or occult bleeding with the need for volume replacement therapy</p> <p>Intended recruitment of 60 adult participants</p> <p><i>Inclusion</i></p> <ul style="list-style-type: none"> • Trauma patient • ≥ 18 years of age, either sex • Major bleeding or occult bleeding • Need for volume replacement therapy • Patient who will be admitted to one of the participating hospitals <p><i>Exclusion</i></p> <ul style="list-style-type: none"> • Solely penetrating trauma • Solely head injury • Ongoing severe haemodynamic instability refractory to therapy (vasopressor, volume) • Inevitably fatal injury as evaluated by emergency physician • Need for CPR on the scene • Deep hypothermia (< 30 °C) • Pregnant women • Known recent history of thromboembolic events within the last 6 months • Known to be on anticoagulant therapy • Known refusal of participation in this clinical trial
Interventions	Pre-hospital fibrinogen concentrate vs placebo
Outcomes	<p>Primary Outcome</p> <p>Change from baseline (before fibrinogen administration) to hospital admission change in FIBTEM maximum clot firmness</p>

EudraCT: 2010-022923-31 (Continued)

Secondary Outcomes

Other ROTEM parameters and coagulation tests
Incidence of thrombosis after 7 days assessed by duplex sonography

Starting date	Unclear
Contact information	Dr. Fries, pamela.schech@i-med.ac.at
Notes	Sponsor: Medical University Innsbruck Funding from LFB Biomedicaments, France and United States Army

EudraCT: 2013-003532-68

Trial name or title	Fibrinogen concentrate supplementation in the management of bleeding during paediatric cardiopulmonary bypass: a phase 1B/2A, open label dose escalation trial
Methods	Randomised controlled open trial
Participants	<p>Neonates and infants (to 23 months old), < 12 kg, who are at risk of mediastinal bleeding following cardiopulmonary bypass surgery for congenital heart disease. Planned sample size = 90</p> <p><i>Inclusion criteria;</i></p> <ul style="list-style-type: none"> • Congenital heart disease requiring non-emergency* surgery on cardiopulmonary bypass • Age range: > 36 weeks corrected gestation • Weight 2.5 – 12 kg • Informed consent to participate <p><i>Exclusion criteria;</i></p> <ul style="list-style-type: none"> • Known pre-existing inherited coagulopathy • Known pre-existing inherited thrombophilia • Recent, acute (within previous 2 weeks) thrombosis in a major vessel or thrombotic-related major complications • Administration of antiplatelet agents (e.g. aspirin) < 48 hrs prior to surgery • Known hypersensitivity/allergy to trial drug or similar products • History of anaphylaxis • Enrolment in another clinical trial in the previous 3 months • Parent/guardian unable to provide informed consent (this can include insufficient understanding of the trial, as judged by the clinician taking consent) • Major co-morbidity likely to increase risk of mortality from surgical procedure • Significant renal/liver impairment within 2 days of planned surgery (creatinine > 2x ULN, alanine transaminase > 2x ULN)
Interventions	<p>Patients will be screened while on cardiopulmonary bypass (approximately 1 hour before end of operation) using Rotational Thromboelastometry: FIBTEM maximum clot firmness</p> <p>If maximum clot firmness < 7 mm, participants will be randomised to fibrinogen concentrate or placebo in a 2:1 ratio.</p> <p>Dose will be tailored to participant based upon measured FIBTEM maximum clot firmness and desired target range.</p> <p>Fibrinogen concentrate will be administered while on cardiopulmonary bypass</p>

EudraCT: 2013-003532-68 (Continued)

Intervention: Human fibrinogen concentrate (RiaSTAP®)

Control: placebo (intravenous solution for injection)

Outcomes	<p>Primary outcomes</p> <ol style="list-style-type: none"> 1. Fibrinogen concentration (measured using the Clauss method) 2. Fibrin polymerisation (measured by rotational thromboelastometry FIBTEM maximum clot firmness) achieved within 5 minutes of completion of study drug <p>Secondary outcome</p> <p><i>Efficacy</i></p> <ul style="list-style-type: none"> • Mediastinal drain losses in first 24 hours after PICU admission • Requirement for delayed sternal closure due to clinical bleeding/tamponade • Requirement for ancillary blood transfusions in first 24 hours after PICU admission • Use of intra- and post-operative ancillary clotting products (as per transfusion algorithm) • Fibrinogen levels and ROTEM variables at times T4 – T6 (see below for details). <p><i>Safety</i></p> <ul style="list-style-type: none"> • Incidence of major thrombotic event/thromboembolic-associated complications • Surgical shunt occlusion • Imaging evidence of intracardiac/major vessel thrombosis up to discharge or day 30 post-operative • Stroke from sinovenous thrombosis or arterial ischaemia of cardioembolic origin • Pulmonary embolism • Superior vena cava syndrome • Thrombotic vessel obstruction requiring active treatment with: thrombolysis/mechanical intervention (open thrombectomy, cardiac catheter) • Cardiopulmonary arrest associated with thrombosis • Allergic/hypersensitivity reaction to trial drug <p>Time point(s) of evaluation of the end points</p> <p>All safety analyses will be up to hospital discharge or day 30 post-operative (whichever comes first). All efficacy endpoints will be evaluated up to 24 hours after PICU admission with fibrinogen levels and ROTEM variables evaluated at time points T4 - T6. T4: at PICU admission T5: 4 hours after PICU admission T6: 24 hours after PICU admission</p>
Starting date	11 March 2014
Contact information	Dr Shane Tibby, Guy's and St Thomas' NHS Foundation Trust. shane.tibby@gstt.nhs.uk
Notes	Investigator network involved with the trial: Medicines For Children Research Network (MCRN) Trial sponsor: CSL Behring Ltd Study completed (March 2016) and awaiting publication

EudraCT: 2013-004484-31

Trial name or title	Rivaroxaban and PCC: Prothrombin Complex Concentrate in patients with bleeding complications related to Rivaroxaban
Methods	Not stated
Participants	<p>Patients with bleeding complications. Target sample size = 20</p> <p><i>Inclusion criteria;</i></p> <ul style="list-style-type: none"> • Patient at the obvious age of ≥ 18 years of either sex • Patients receiving rivaroxaban • Patients with significant bleeding with the need of reversal and/or patients needing acute reversal of rivaroxaban anticoagulation effects <p><i>Exclusion criteria;</i></p> <ul style="list-style-type: none"> • Additional treatment with further procoagulant therapy including recombinant activated factor seven, FEIBA or other coagulation factor concentrates (except fibrinogen concentrate, desmopressin and tranexamic acid) • Risk for thromboembolic events higher than the bleeding risk as anticipated by the physician • Pregnant or nursing women • Patient with suspected or confirmed sepsis • Patient with known recent history of thromboembolic events within the last 3 months • Patients who disagree to participate in the trial (for emergency patients: patients with known refusal of a participation in this clinical trial) • Known active participation in another clinical trial • Any condition, including the presence of laboratory abnormalities, which would place the subject at unacceptable risk if he/she would participate in the trial or confound in the ability to interpret data from the trial • Any serious medical condition, laboratory abnormality, or psychiatric illness that would prevent the subject from signing the informed consent form
Interventions	Rivaroxaban (tablet) versus Prothrombin Complex Concentrate
Outcomes	<p>Primary</p> <ul style="list-style-type: none"> • Difference in thrombin generation <p>Secondary</p> <ul style="list-style-type: none"> • Alterations of following parameters from and correlation with blood level of rivaroxaban (Xarelto®) and thrombin generation: <ul style="list-style-type: none"> - Thrombin generation - Single factor profiles - Standard coagulation tests outcome (prothrombin time, activated partial thromboplastin time, fibrinogen, antithrombin time, etc.) - ROTEM® results (FIBTEM, ExTEM, InTEM, HepTEM) - Further biological parameters - Differences between sample groups
Starting date	10 January 2014
Contact information	Projektmanagement, Medizinische Universität Innsbruck / Univ.-Klinik für Allg. u.l Chirurg. Intensivmedizin, Innsbruck, Austria

EudraCT: 2013-004484-31 (Continued)

bettina.schenk@i-med.ac.at

Notes	Trial sponsor: CSL Behring Ltd
-------	--------------------------------

NCT00994045

Trial name or title	Coagulopathy during surgery for the repair of extent 4 thoraco-abdominal aortic aneurysms - feasibility study of the use of fibrinogen concentrate by infusion in place of fresh frozen plasma
Methods	Randomised single-blinded controlled trial
Participants	<p>20 participants enrolled</p> <p><i>Inclusion Criteria</i></p> <ul style="list-style-type: none"> • Undergoing elective thoracoabdominal aneurysm repair. • Over 18 years of age. • Able to give written informed consent. <p><i>Exclusion Criteria</i></p> <ul style="list-style-type: none"> • Previous aortic surgery (re-do surgery) • Emergency surgery • Pregnancy • Females of child-bearing age (less than 45 years) not using medically approved method of contraception • Congenital or acquired coagulopathy • Known allergy to study drug
Interventions	<p>Active Comparator: Fresh Frozen Plasma</p> <p>Active Comparator: Fibrinogen concentrate</p>
Outcomes	<p>Primary Outcome</p> <ul style="list-style-type: none"> • Pattern of coagulation disturbance in the conventional treatment (FFP) and fibrinogen concentrate groups (time frame: intraoperatively and up to 24 hours post-operatively). <p>Secondary Outcomes</p> <ul style="list-style-type: none"> • Proportion of participants in the fibrinogen group in whom FFP transfusion is required during surgery • Units of FFP transfused, during surgery and up to 24 hours after surgery. • Units of platelets and allogeneic red cells transfused, during surgery and up to 24 hours after surgery • Intra-operative blood loss
Starting date	June 2010
Contact information	Principal Investigator: Akastair Nimmo, NHS Lothian
Notes	<p>Study Sponsor: University of Edinburgh</p> <p>Collaborators: NHS Lothian, CSL Behring</p> <p>Note: This study is completed but unpublished. Additional data have not been sought</p> <p>Note: This study is randomised and controlled but only participant blinded</p>

NCT00994045 (Continued)

Note: There has been a change in primary and secondary outcomes for this study, prior to participant recruitment

NCT01539057

Trial name or title	The efficacy of the administration of fibrinogen in liver transplantation (FibstudLT)
Methods	Multicentre, randomised, double-blind, placebo-controlled trial
Participants	<p>Patients undergoing liver transplantation. Planned sample size = 132</p> <p><i>Inclusion Criteria:</i></p> <ul style="list-style-type: none"> • Patients candidates for liver transplantation • Patients with a value of pre-transplant plasma fibrinogen less than 2.9 g / L. <p><i>Exclusion Criteria;</i></p> <ul style="list-style-type: none"> • Patients with a value of fibrinogen in the 24 hours prior to the intervention than 2.9 g/L • Known history of thromboembolic events in 30 days • Known or suspected pregnancy • Previous randomisation in this trial • Known or suspected allergy to trial products or related products • Known presence of congenital bleeding disorder. Patients treated with aspirin, warfarin • The following indications for transplantation: familial polyneuropathy, acute liver failure, biliary cirrhosis and sclerosing cholangitis, Budd-Chiari syndrome • Heart beating donors and living donor • Patient reluctant to participate in the trial
Interventions	Intravenous fibrinogen (administered until an expected plasmatic value of 2.9 g/L is achieved) versus placebo (saline of the same dose in volume as the intravenous fibrinogen)
Outcomes	<p>Primary</p> <p>Percentage of participants requiring transfusion of red blood cell units during the procedure</p> <p>Secondary</p> <ul style="list-style-type: none"> • Percentage of participants requiring blood products other than red cell concentrates during the procedure. Number of red blood cell units transfused during surgery. Number of units of fresh frozen plasma transfused during surgery. Number of platelet units transfused during surgery. Grams of fibrinogen administered during surgery • Operative outcome, measured at 4 weeks: operative mortality, liver graft survival, thrombotic complications of all types and causes • Liver transplantation outcome, measured at 1 year • Follow-up of graft survival and participant mortality 1 year after liver transplantation
Starting date	July 2012
Contact information	Antoni Sabate, Head of Anesthesiology, Hospital Universitari de Bellvitge, Spain
Notes	<p>The trial is open but not recruiting participants (as verified in December 2014; website accessed 13 January 2016)</p> <p>Sponsors and collaborators: Hospital Universitari de Bellvitge and the Spanish Clinical Research Network - CAIBER</p>

NCT02155725

Trial name or title	Fibrinogen in Haemorrhage of Delivery (FIDEL)
Methods	Randomised, double-blind, multicentre, placebo-controlled trial
Participants	<p>Women with a postpartum haemorrhage following vaginal delivery. Planned sample size: 434</p> <p><i>Inclusion Criteria;</i></p> <ul style="list-style-type: none"> • Signed and dated informed consent form • Vaginal delivery • postpartum haemorrhage requiring iv administration of prostaglandins • At least one available result of Hb level during the third trimester of pregnancy • Women 18 years and older • Covered by healthcare insurance in accordance with local requirements <p><i>Exclusion Criteria;</i></p> <ul style="list-style-type: none"> • Caesarean section • Haemostatic intervention (as ligation, embolisation or hysterectomy) already decided at the time of inclusion • Known placenta praevia or accreta • Hb level < 10g/dl during the third trimester of pregnancy • History of venous or arterial thromboembolic event • Known inherited bleeding or thrombotic disorders • Treatment with low-molecular-weight heparin within 24 hours prior to the inclusion • Treatment with acetylsalicylic acid within 5 days prior to the inclusion • Treatment with vitamin K antagonists within 7 days prior to the inclusion • Administration of fibrinogen concentrate within 48 hours prior to the inclusion • Administration of FFP, platelets units or pro haemostatic drugs, tranexamic acid and recombinant FVIIa or prothrombin complex concentrates within 48 hours prior to the inclusion • Administration of red blood cells within 3 months prior to the inclusion • Participation in another interventional clinical trial within 30 days prior to the inclusion • Previous inclusion/enrolment in the present clinical trial • Known history of hypersensitivity or other severe reaction to any component of Clottafact® or placebo • Minors, majors under guardianship, persons staying in health or social institutes and people deprived of their freedom • Known drug or alcohol abuse • Patients whose use of concomitant medication may interfere with the interpretation of data • Any other current significant medical condition that might interfere with treatment evaluation according to the investigator's judgement • Patients who are unlikely to survive through the treatment period and evaluation • Patients transferred from another service
Interventions	Human fibrinogen concentrate (2 vials (200 mL)/3g intravenous) versus placebo (2 vials (200 mL))
Outcomes	<p>Primary</p> <ul style="list-style-type: none"> • Percentage of participants losing at least 4 g/dl of Hb, and/or requiring the transfusion of at least 2 units of red blood cells within the 48 hours following the administration of the intervention <p>Secondary</p> <ul style="list-style-type: none"> • Evolution of haemorrhage at 48 hours

NCT02155725 (Continued)

- Percentage of participants losing at least 4 g/dL and 3 g/dL of Hb by the Hb reference level at 48 hours
- Percentage of participants requiring at least 1 haemostatic intervention at 48 hours
- Maternal morbi-mortality at up to 6 ± 2 weeks after delivery
- Length of stay in resuscitation or in ICU or both
- Change in plasma fibrinogen level between H0 and Day 2, after early administration
- Number of participants with at least 1 adverse event or serious adverse event or both
- Evolution of haemorrhage at 48 hours: Percentage of participants losing at least 4 g/dL by Hb level measured at inclusion
- Evolution of haemorrhage at 48 hours : Percentage of participants requiring the transfusion of at least 2 units of red blood cells
- Evolution of haemorrhage at 48 hours: Number of units of transfused blood products
- Evolution of haemorrhage at 48 hours: total blood loss
- Maternal morbi-mortality at each follow-up assessment (up to 6 ± 2 weeks after delivery): SOFA score in women transferred to resuscitation or ICU or both
- Maternal morbi-mortality at each follow-up assessment (up to 6 ± 2 weeks after delivery): death during the trial

Starting date	April 2014
Contact information	Dr Anne-Sophie Ducloy-Bouthors, Maternité Jeanne de Flandre - 59037 Lille, France. anne-sophie.ducloy@chru-lille.fr
Notes	Sponsors and collaborators: Laboratoire Français de Fractionnement et de Biotechnologies

NCT02203968

Trial name or title	Fibrinogen in the Initial Resuscitation of Severe Trauma (FiIRST)
Methods	Randomised, double-blind feasibility trial
Participants	Planned sample size = 50 <i>Inclusion criteria;</i> <ul style="list-style-type: none"> • Injured trauma (penetrating or blunt) patients who are at risk of significant bleeding, defined as: <ul style="list-style-type: none"> * Systolic blood pressure ≤ 100mmHg at any time from the injury scene until 30 mins after hospital admission; AND * Red blood cell transfusion has been ordered by the trauma team leader (or delegate) <i>Exclusion criteria;</i> <ul style="list-style-type: none"> • Patients in shock which the aetiology is purely not related to bleeding: <ul style="list-style-type: none"> * Cardiogenic (myocardial or valvular dysfunction); * Distributive (septic, anaphylactic, acute adrenal insufficiency and neurogenic) * Obstructive (cardiac tamponade, tension pneumothorax and massive pulmonary emboli) • Severe head injury, defined as any of the following: <ul style="list-style-type: none"> * Glasgow coma scale of 3 due to severe TBI, * TBI with clear indication of immediate neurosurgical intervention based on clinical findings (mechanism of trauma associated with focal signs such as anisocoria with fixed pupil) or on CT results (bleeding causing mass effect); * Unsalvageable head injury such as through-through gunshot wound to the head, open skull fracture with exposure/loss of brain tissue; as per the trauma team or neurosurgery initial clinical assessment or as per initial CT of the head findings

NCT02203968 (Continued)

- Known complete or incomplete spinal cord injury
- Known hereditary or acquired coagulopathies unrelated to the trauma resuscitation (e.g. known hepatic dysfunction)
- Use of anticoagulant medications such as warfarin, low-molecular-weight heparin, and direct thrombin and factor Xa inhibitors
- Moribund with evidence of unsalvageable injuries and withdrawal of care, as per the trauma team
- Received blood products prior to admission
- Patients with estimated body weight under 50 Kg
- Patients with known or suspected pregnancy
- Patients arriving more than 6 hours after injury

Interventions

Fibrinogen concentrate (300 mL of RiaSTAP® infused) versus placebo (normal saline will be administered intravenously as a single 300 mL rapid infusion within 1 hour of hospital admission)

Outcomes
Primary

- Feasibility endpoint: proportion of participants receiving trial intervention within first hour of hospital admission

Secondary

- Other feasibility endpoints and physiologic endpoints at 1 year
 - * Proportion of participants receiving trial intervention prior to any blood transfusion
 - * Times to randomisation, issue, and start of infusion
 - * Duration of infusion
 - * Wastage of trial intervention
 - * Proportion of participants with blood tests performed for all time points
 - * Proportion of missed participants (eligible but not randomised)
 - * Randomisation errors
- Physiologic endpoints:
 - * Plasma fibrinogen levels
 - * Rotational Thromboelastometry/Thromboelastography parameters (functional fibrinogen, FIBTEM, FIBTEM plus)
 - * Plasmin anti-plasmin complex
 - * Tissue plasminogen activator
 - * Plasminogen activator inhibitor-1
 - * D-dimer levels
 - * International normalized ratio
 - * Partial thromboplastin time
 - * Complete blood count
 - * Endogenous thrombin potential
 - * Factor XIII; factor V; and activated protein C will be obtained on admission, and at 3, 11, 23 and 47 hours following the start time of trial infusion

NCT02203968 (Continued)

- Clinical Parameters at 1 year
 - * Rates of symptomatic thromboembolic complications in both trial arms defined by the evidence of any of the following any time during hospital stay:
 - * Deep venous thrombosis; myocardium infarction; cerebral vascular accident; pulmonary embolism; and arterial thrombosis
 - * Rates of asymptomatic deep venous thrombosis by leg doppler at day 7 of hospital stay
 - * Mortality by exsanguination
 - * All-cause mortality at 28 days (mainly due to exsanguination; mainly neurological/due to traumatic brain injury/withdrawal of care; or mainly due to multiple organ failure/sepsis)
 - * Incidence of acute lung injury/acute respiratory distress syndrome before day 28
 - * Multiple organ failure
 - * Total amount of crystalloids used in 24 hours
 - * Units of blood and blood products transfused in 24 hours of hospitalisation

Starting date	October 2014
Contact information	Dr Barto Nascimento, Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada barto.nascimento@sunnybrook.ca
Notes	Trial sponsored by Sunnybrook Health Sciences Centre

NCT02253082

Trial name or title	Vasculopathic Injury and Plasma as Endothelial Rescue - OCTApLas Trial (VIPER-OCTA)
Methods	Single-centre randomised, single-blinded, controlled, investigator-initiated pilot trial
Participants	<p>Patients undergoing emergency surgery for thoracic aortic dissections. Planned sample size = 42</p> <p><i>Inclusion criteria</i></p> <ul style="list-style-type: none"> • Patient eligible for emergency surgery on cardiopulmonary bypass pump for a thoracic aortic dissections AND • Age > 18 years AND • Consent obtainable from patient or by proxy (independent physicians or next of kin or both) <p><i>Exclusion criteria</i></p> <ul style="list-style-type: none"> • Documented refusal of blood transfusion OR • FFP transfusion before randomisation OR • Aortic dissection due to trauma OR • Treatment with Glycoprotein IIb/IIIa inhibitors < 24 hours from screening OR • Withdrawal from active therapy OR • Expected to die < 24 hours OR • Previously within 30 days included in a randomised trial, if known at the time of enrolment • Known IgA deficiency with documented antibodies against IgA • Known hypersensitivity to OctaplasLG®: the active substance, any of the excipients (sodium citrate dihydrate, sodium dihydrogenphosphate dihydrate or glycine) or residues from the manufacturing process (Tri (N-Butyl) Phosphate (TNBP) and Octoxynol (Triton X-100)) • Known severe deficiencies of protein S • Pregnancy (non-pregnancy confirmed by patient being post-menopausal or having a negative serum-hCG)

NCT02253082 (Continued)

Interventions	OctaplasLG® OctaplasLG® versus fresh frozen plasma
Outcomes	<p>Primary</p> <ul style="list-style-type: none"> Plasma levels of endothelial markers (Syndecan-1, soluble thrombomodulin, sE-selectin, sVE-cadherin) at 24 hours after arrival in ICU for post-operative care, as compared to baseline <p>Secondary</p> <ul style="list-style-type: none"> Plasma levels of endothelial markers (Syndecan-1, soluble thrombomodulin, sE-selectin, sVE-cadherin) at 48 hours post-operatively, as compared to baseline Acute Kidney Injury according to RIFLE Criteria in the first 7 post-operative days Renal replacement therapy SOFA worst score during ICU stay 30-day and 90-day mortality P-CRP, IL-6, P-Catecholamines at 24 hours and 48 hours Length of stay in ICU and hospital Severe adverse reactions <p>Tertiary</p> <ul style="list-style-type: none"> TRALI TACO
Starting date	November 2014
Contact information	Dr Jakob Stensballe, Rigshospitalet, Denmark stensballe@rh.dk
Notes	Also registered on EudraCT with number 2014-000452-28

NCT02299947

Trial name or title	Blood transfusions after aortic surgery (FIBTEG)
Methods	Randomised, double blind trial
Participants	<p>Patients undergoing surgery for ascending aortic aneurysm. Planned sample size = 82.</p> <p><i>Inclusion Criteria;</i></p> <ul style="list-style-type: none"> Elective surgery for thoracic aneurysm Aged > 18 years <p><i>Exclusion Criteria;</i></p> <ul style="list-style-type: none"> Prior thrombosis or myocardial infarction Congenital coagulation disorder Use of anti-coagulants prior to surgery Prior thoracic surgery Pregnancy Pre-operative fibrinogen concentration < 1 g/L
Interventions	Fibrinogen (Haemocomplettan P) versus placebo (saline solution)

NCT02299947 (Continued)

Outcomes

Primary

- Number of transfused blood products 24 hours after surgery

Secondary

- Blood loss 24 hours after surgery
- Re-operation within 30 day after surgery
- Mortality at 30 days

Starting date	August 2014
Contact information	Eline Vlot, St Antonius Hospital, Utrecht, The Netherlands e.vlot@antoniusziekenhuis.nl
Notes	Sponsors and collaborators: St. Antonius Hospital, Utrecht, The Netherlands

NCT02344069

Trial name or title	Pilot Randomized trial of Fibrinogen in Trauma Haemorrhage (PRooF-iTH)
Methods	Single-centre, randomised (1:1, active:placebo), placebo-controlled, double-blinded, investigator-initiated phase II trial
Participants	Expected enrolment - 40 patients <i>Inclusion criteria</i> <ul style="list-style-type: none"> • Trauma patient received directly from the scene of the accident • Age ≥18 years • Initiated order of transfusion of at least 1 blood component within the first hour of arrival • Predicted to need transfusion package therapy during the initial resuscitation (first 2 hours) • Consent obtainable from patient or scientific guardians (independent physicians or next of kin or both) <i>Exclusion criteria</i> <ul style="list-style-type: none"> • Duration of > 2 hours from time of accident to arrival at trauma centre • Known anticoagulant treatment (vitamin K antagonist, dabigatran, rivaroxaban, apixaban) • Severe isolated TBI • Moribund patient with devastating injuries and expected to die within 1 hour of admission • Withdrawal from active therapy • Previously, within 30 days, included in a randomised trial, if known at the time of enrolment • Known body weight < 55 kg • Any blood product prior to inclusion
Interventions	Active intervention: intravenous injection of fibrinogen concentrate (RiaSTAP®); 60 – 70 mg/kg ideally within 1 hour of arrival prior to blood products Control: 0.9% saline solution
Outcomes	Primary Outcome <ul style="list-style-type: none"> • Change in thrombelastograph TEG® functional fibrinogen maximum amplitude (MA) in millimetres at 15 mins after the intervention

NCT02344069 (Continued)

Secondary Outcomes

- TEG® Functional Fibrinogen maximum amplitude in millimetres at 2, 6, 24 and 72 hours post-intervention
- TEG® maximum amplitude in millimetres at 15 mins, 2, 6, 24 and 72 hours post-intervention
- Transfusion requirements (red blood cells units or FFP or platelets) at 2, 6, 24, 72 hours and in total at day 30
- Total use of haemostatic therapy (i.e. use of coagulation factor concentrates and tranexamic acid) in the first 24 and 72 hours, omitted from this is active treatment (intervention)
- Time to intervention or placebo
- Time to FFP and platelet transfusion
- Percentage of participants receiving intervention or placebo < 1 hour of arrival
- Time to surgical control of bleeding as noted by the surgeon
- Severe adverse reactions at day 30, defined as symptomatic thromboembolism at day 30 and anaphylaxis at day 30
- 24-hour and 30-day mortality

Starting date	February 2015
Contact information	Pär I Johansson, Section for Transfusion Medicine, Capitol Region Blood Bank, Copenhagen, Denmark per.johansson@regionh.dk
Notes	Trial sponsor: CSL Behring Ltd

NCT02429453

Trial name or title	FFP versus PCC in traumatic intracranial hemorrhage
Methods	Randomised, open-label trial
Participants	<p>Patients with a traumatic intracranial haemorrhage. Planned sample size = 30</p> <p><i>Inclusion Criteria;</i></p> <ul style="list-style-type: none"> • Coumadin use • INR of 2.0 or higher on arrival at the trial centre • Evidence on cranial imaging of subdural haematoma, epidural haematoma, cerebral contusion, traumatic subarachnoid haemorrhage, or traumatic intraparenchymal haemorrhage <p><i>Exclusion Criteria;</i></p> <ul style="list-style-type: none"> • Unable to obtain consent • Estimated survival < 24 hours • Hypersensitivity to 4 factor PCC • Concomitant use of novel vitamin K antagonists • Religious/social prohibition to receiving blood products • Need for emergent, non-neurosurgical operative intervention • Mechanical heart valves
Interventions	FFP (a single dose based on INR per the following regimen: 2 U for INR of 2 - 2.5; 3 U for INR of 2.5 - 3; 4 U for INR of 3 - 3.5; 5 U for INR of 3.5 - 4; 6 U for INR of 4+) versus 4 factor PCC (a single dose of 4 factor PCC per the following dosing regimen: 25 U/kg for INR of 2 - 4; 35 U/kg for INR of 4 - 6; 50 U/kg

NCT02429453 (Continued)

for INR of 6+; maximum dosing weight of 100 kg, participants may be dispensed $\pm$ 10% of ordered dose)

Outcomes	<p>Primary</p> <ul style="list-style-type: none"> Rapid reversal of warfarin as measured by INR drawn at 30 minutes after transfusion <p>Secondary</p> <ul style="list-style-type: none"> Radiographic expansion of traumatic intracerebral haemorrhage as measured by CT scan within 24 hours of presentation Timing of reversal of warfarin as measured by INR drawn at 3 hours, 8 hours and 24 hours after transfusion Thromboelastography response as measured by results of ROTEM analysis at 30 minutes and 24 hours after transfusion Absolute INR reversal as measured by INR drawn 24 hours after transfusion Need for operative intervention as measured by need for neurosurgical procedure during the hospitalisation Estimated blood loss during any neurosurgical procedure Further transfusion needs as measured by number of units of blood/platelet/plasma products transfused during the hospitalisation In-hospital mortality Total hospital cost 30-day outcome as measured by the Glasgow outcome score Complications as measured by development of deep vein thrombosis, pulmonary embolism, myocardial infarction, stroke, unanticipated intubation, heart failure, or need for aggressive diuresis during the hospitalisation
Starting date	April 2015
Contact information	Dr Jian Guan, University of Utah Medical Center, Salt Lake City, USA jian.guan@hsc.utah.edu
Notes	Sponsors and collaborators: University of Utah

NCT02528708

Trial name or title	A Program to Evaluate RiaSTAP® and FIBTEM® for the Early Control and Treatment of Postpartum Hemorrhage (PERFECT PPH)
Methods	Prospective, single-centre, randomised, placebo-controlled, double-blind clinical trial
Participants	<p>Estimated enrolment: 40 participants</p> <p><i>Inclusion Criteria</i></p> <ul style="list-style-type: none"> Informed consent from participant Age $\geq$ 18 years and $<$ 50 years Primary postpartum haemorrhage defined as bleeding from uterus or the birth canal or both within 24 hours postpartum Vaginal delivery or Caesarean delivery (irrespective of aetiology of PPH, such as accreta), with estimated blood loss $>$ 1000 mL and ongoing bleeding notwithstanding standard treatment measures (volume replacement, uterine massage, uterotonic agents) FIBTEM®- A10 $<$ 18 mm (corresponding to a maximum clot firmness value of $<$ 20 mm and to a plasma fibrinogen level approximately $<$ 3 g/L)

NCT02528708 (Continued)

	<p>Exclusion Criteria:</p> <ul style="list-style-type: none"> Refusal to give written informed consent Refusal to receive blood transfusion Known inherited deficiencies of coagulation Personal history of thrombosis Either pre-pregnancy or ante-partum antithrombotic treatment due to increased risk of thrombosis Administration of platelets, FFP or cryotherapy prior to study drug
Interventions	<p>Intervention: fibrinogen concentrate (RiaSTAP®)</p> <ul style="list-style-type: none"> The dose of fibrinogen concentrate needed to achieve target plasma fibrinogen levels will be calculated using a formula that accounts for the baseline FIBTEM® - A10 value and the participant's body weight assessed at hospital admission. In general, a 70 kg participant requires a fibrinogen dose of approximately 0.5 g to increase the MCF by approximately 1mm <p>Placebo: 0.9% saline solution</p>
Outcomes	<p>Primary Outcome</p> <ul style="list-style-type: none"> Maximum clot firmness (via FIBTEM A10) at 15 minutes <p>Secondary Outcomes</p> <ul style="list-style-type: none"> Not stated.
Starting date	Not stated.
Contact information	Michael J Paidas, Yale School of Medicine
Notes	<p>Estimated primary completion date: January 2020</p> <p>Study Sponsor: Yale University</p> <p>Collaborator: CSL Behring</p> <p>Note: Study is currently suspended having never begun; it is pending review to be reopened</p>

NCT02557672

Trial name or title	Prothrombin complex concentrate (PCC) compared to fresh frozen plasma (FFP) for post-cardiopulmonary bypass coagulopathy and bleeding, a prospective randomized trial at large US Medical Center.
Methods	Randomised controlled trial, open-label
Participants	<p>Estimated enrolment: 100 participants, by invitation only</p> <p><i>Inclusion Criteria</i></p> <ul style="list-style-type: none"> 18 years of age or more Elective cardiac surgical procedure using cardiopulmonary bypass Evidence of excessive microvascular bleeding in the surgical field as determined by the surgical team in addition to a prothrombin time > 16.6 secs or INR > 1.6 secs <p><i>Exclusion Criteria</i></p> <ul style="list-style-type: none"> Inability to grant informed consent or comply with study procedure

NCT02557672 (Continued)

- History of hypercoagulable condition (e.g. factor V Leiden, AT-3 deficiency, prothrombin gene mutation, anti-phospholipid antibody syndrome, etc.) or previous unprovoked thromboembolic complications
- Coagulopathic conditions such as factor deficiencies, factor inhibitors, heparin-induced thrombocytopenia, or use of intravenous anticoagulants other than heparin at the time of cardiovascular surgery
- Thromboembolic event within past 3 months
- Received oral therapy with clopidogrel, prasugrel, rivaroxaban or dabigatran within the past 5 days
- Patients taking chronic warfarin therapy who have not discontinued treatment and demonstrated an INR < 1.3 prior to surgery
- Fibrinogen level < 150 mg/dL on initial post-cardiopulmonary bypass results
- Antithrombin 3 level < 80% control (preoperative)
- Emergency open heart surgery
- Cardiopulmonary bypass time is expected to be < 30 minutes
- Age < 18 years
- Pregnant

Interventions

Active comparator: Fresh frozen plasma, 10 - 15 mL/kg (rounded to nearest unit)

Experimental intervention: Prothrombin complex concentrate (human, Kcentra) 15 units/kg

After cardiopulmonary bypass participants will receive protamine at dose 0.01 mg/unit of heparin given with target activated clotting time within 10% of baseline value. After protamine administration the activated clotting time, complete blood count, prothrombin time, INR, activated partial thromboplastin time and fibrinogen will be collected by pre-existing arterial access. If the activated clotting time is > 10% above the baseline additional protamine will be given at the anaesthetist's discretion. Evaluation and determination of excessive microvascular

bleeding in the surgical field will occur 10 minutes after return of activated clotting time to within 10% of baseline. Participants with clinical evidence of excessive microvascular bleeding in the surgical field as determined by the surgical team, along with a prothrombin time > 16.6 secs/INR > 1.6 secs will receive the student intervention or comparator

Outcomes
Primary Outcomes

- Blood loss at 24 hours; blood loss as collected by chest tube drains 24 hours after surgery
- Blood product transfusion at 24 hours; total quantity of all blood products transfused 24 hours after surgery

Secondary Outcomes

- None stated

Starting date

August 2016

Contact information

Gregory Nuttal, Mayo Clinic

Notes

Study sponsor: Mayo Clinic

Collaborators: CSL Behring

Estimated completion date: July 2018

Note: This study is open-label

NCT02745041

Trial name or title	Fibrinogen Early In Severe Trauma study (FEISTY)
Methods	Multicentre randomised controlled trial
Participants	<p>Adult trauma patients judged by treating clinicians to have clinically significant haemorrhage or potential for significant transfusion requirements with an Assessment of Blood Consumption Score ≥ 2</p> <p>Estimated enrolment of 100 adult trauma patients</p> <p><i>Inclusion criteria</i></p> <ul style="list-style-type: none"> Adult (≥ 18 years) affected by trauma Judged to have significant haemorrhage or predicted to require significant transfusion with Assessment of Blood Consumption Score ≥ 2 or by treating clinician judgement <p><i>Exclusion criteria</i></p> <ul style="list-style-type: none"> Injury judged incompatible with survival Previous fibrinogen administration this admission Pretrauma centre dedicated fibrinogen replacement Known objection to blood products Known coagulation disorder Pregnancy Enrolment in competing study
Interventions	Fibrinogen supplementation comprising either protocol-directed fibrinogen concentrate or cryoprecipitate therapy
Outcomes	<p>Primary Outcome</p> <ul style="list-style-type: none"> The time interval in minutes from FIBTEM analysis and clinical scenario suggesting that fibrinogen supplementation is required to the completion of the first administration of fibrinogen supplementation (fibrinogen concentrate or cryoprecipitate). <p>Secondary Outcomes</p> <ul style="list-style-type: none"> Effects of fibrinogen supplementation (fibrinogen concentrate or cryoprecipitate) on plasma fibrinogen Time to randomisation Time course of fibrinogen levels as measured by FIBTEM and Clauss Fibrinogen Transfusion requirements Clinically relevant thromboembolic complications Mortality at several time points including hospital discharge and 90 days
Starting date	2017
Contact information	James Winearls, james.winearls@health.qld.gov.au
Notes	Note: clinicians will be unblinded to intervention

NCT02777424

Trial name or title	Prothrombin complex concentrate versus fresh frozen plasma to correct coagulation disorders in adult neurosurgical patients
---------------------	---

NCT02777424 (Continued)

Methods	Prospective, randomised, multicentre study
Participants	<p>Estimated enrolment: 46 participants</p> <p><i>Inclusion Criteria</i></p> <ul style="list-style-type: none"> • Patient with spontaneous intracranial haemorrhage or traumatic intracranial haemorrhage or patient requiring neurological surgery • Coagulation disorder defined by prothrombin time < 60% <p><i>Exclusion Criteria</i></p> <ul style="list-style-type: none"> • Concomitant use with oral anticoagulant drugs • Acquired deficiency of coagulation factors whose treatment is established • Hypersensitivity to PCC • History of thrombocytopenia induced by heparin • Disseminated intravascular coagulation • Extracranial active bleeding • Hypersensitivity to vitamin K
Interventions	<p>Active comparator: Prothrombin Complex Concentrate (25iU/kg factor IX)</p> <p>Active comparator: Fresh Frozen Plasma (15 mL/kg)</p>
Outcomes	<p>Primary Outcome</p> <ul style="list-style-type: none"> • Proportion of participants with correction of prothrombin time (prothrombin time > 60%, at end of treatment administration, an average of 1 hour) <p>Secondary Outcomes</p> <ul style="list-style-type: none"> • Not stated
Starting date	January 2016
Contact information	Laurence Salomon, lsalomon@for.paris
Notes	Note: Open-label study, estimated completion date July 2017

NCT02822599

Trial name or title	The role of human fibrinogen concentrate (RiaSTAP®) in decreasing the need for component blood therapy in infants undergoing cardiopulmonary bypass
Methods	Randomised, controlled, double-blind trial
Participants	<p>Estimated enrolment of 30 participants</p> <p><i>Inclusion Criteria</i></p> <ul style="list-style-type: none"> • Neonatal and infant cardiac patients presenting for open-heart surgery at Nicklaus Children's Hospital. <p><i>Exclusion Criteria</i></p> <ul style="list-style-type: none"> • Patients above 1 year of age • Patients known to have had an anaphylactic or severe reaction to the study drug or its components

NCT02822599 (Continued)

	<ul style="list-style-type: none"> • ROTEM results at the time of rewarming showing a FIBTEM maximum clot firmness > 15 mm
Interventions	<p>Active comparator: RiaSTAP® fibrinogen concentrate</p> <ul style="list-style-type: none"> • Group will receive an infusion of RiaSTAP® fibrinogen concentrate after termination of cardiopulmonary bypass at a dose of 70 mg/kg after randomisation to this group <p>Placebo comparator: Saline</p> <ul style="list-style-type: none"> • Group 2 will receive a placebo consisting of normal saline 0.9% after randomisation to this group
Outcomes	<p>Primary Outcome</p> <ul style="list-style-type: none"> • post-operative bleeding volume and transfusion requirement after surgery 24 hours after surgery <p>Secondary Outcomes</p> <ul style="list-style-type: none"> • Adverse events 24 hours after surgery • Requirement for ventilator support • Length of stay in the ICU • Re-exploration for bleeding • Length of stay in the hospital
Starting date	January 2017
Contact information	Jenny Esteves jenny.esteves@mch.com
Notes	<p>Estimated completion date: April 2019</p> <p>Study Sponsor: Nicklaus Children's Hospital, Miami</p>

NCT03014700

Trial name or title	Repurposing of fibrinogen concentrate as a cost-effective and safe hemostatic agent in infants undergoing cardiac surgery on cardiopulmonary bypass
Methods	Randomised, single-blind, controlled trial
Participants	<p>Planned sample size: 30 participants</p> <p><i>Inclusion Criteria</i></p> <ul style="list-style-type: none"> • Neonates of at least 32 weeks of gestational age and infants up to 12 months of age with the diagnosis of congenital heart disease, requiring open-heart surgery with cardiopulmonary bypass <p><i>Exclusion Criteria</i></p> <ul style="list-style-type: none"> • Pre-existing coagulopathy, including unexplained bleeding or history of clotting
Interventions	<p>Active comparator: Cryoprecipitate</p> <p>Active comparator: Fibrinogen concentrate</p>
Outcomes	<p>Primary Outcome</p> <ul style="list-style-type: none"> • Intraoperative allogenic donor transfusions (from administration of the drug during surgery to 7 days post-operatively) <p>Secondary Outcomes (approximately 3 - 14 days of hospitalisation)</p>

NCT03014700 (Continued)

- Coagulation laboratory values
- Transfusion requirements
- Ventilator times
- ICU and hospital length of stay
- Haematocrit, fibrinogen level and thromboelastogram values

Starting date	9 January 2017
Contact information	Glyn D Williams: jumbo@stanford.edu
Notes	Estimated study completion date: June 2018 Study sponsor: Stanford University Collaborators: Emory University

NCT03037424

Trial name or title	Prospective, multi-center, randomized, active-control, non-inferiority study comparing fibrinogen concentrate with cryoprecipitate for the treatment of acquired hypofibrinogenemia in bleeding adult cardiac surgical patients
Methods	Prospective, multi-centre, randomised, active-control, single-blinded, non-inferiority phase 3 trial Up to 12 Canadian hospitals will participate
Participants	Estimated Enrolment: 1200 participants <i>Inclusion Criteria</i> <ul style="list-style-type: none"> • Patients undergoing cardiac surgery with cardiopulmonary bypass in whom fibrinogen supplementation is ordered in accordance with accepted clinical standards (significant haemorrhage and known or presumed hypofibrinogenaemia) <i>Exclusion Criteria</i> <ul style="list-style-type: none"> • Receipt of fibrinogen-containing products, including concentrate or cryoprecipitate within 24 hours before surgery (to exclude patients with significant blood loss before surgery) • History of severe allergic reaction to cryoprecipitate or fibrinogen concentrate • Refusal of fibrinogen concentrate or cryoprecipitate due to religious or other reasons • Fibrinogen level known to be > 3.0 g/L within 30 minutes of trial intervention administration (to eliminate the risk of raising patients' fibrinogen levels to > 4.0 g/L with supplementation, which is the upper limit of the normal range) • Known pregnancy
Interventions	Active comparator: Fibrinogen concentrate Octafibrin Active comparator: Cryoprecipitate
Outcomes	Primary Outcome <ul style="list-style-type: none"> • The number of red blood cells, pooled or apheresis platelets and plasma administered during the first 24 hours after termination of cardiopulmonary bypass Secondary Outcomes <ul style="list-style-type: none"> • None declared

NCT03037424 (Continued)

- Safety outcomes will be measured for the first 28 days after surgery

Starting date	22 February 2017
Contact information	Keyvan Karkouti, keyvan.karkouti@uhn.ca
Notes	<p>Estimated primary completion: 30 January 2019</p> <p>One interim analysis will be conducted after 600 participants have been treated to determine whether the study should be terminated for safety reasons, demonstrated non-inferiority or futility reasons</p> <p>Study Sponsor: University Health Network, Toronto</p>

NCT03183479

Trial name or title	The effects of fibrinogen concentrate infusion on blood loss and allogeneic blood conservation in scoliosis surgery
Methods	Prospective, randomised, double-blind, placebo-controlled trial
Participants	<p>Patients older than 12 with adolescent idiopathic scoliosis planned for elective posterior scoliosis correction surgery</p> <p>Planned sample size: 104 participants</p> <p><i>Inclusion Criteria</i></p> <ul style="list-style-type: none"> • Patients diagnosed as adolescent idiopathic scoliosis • Planned for elective posterior scoliosis correction surgery at Peking Union Medical College Hospital <p><i>Exclusion Criteria</i></p> <ul style="list-style-type: none"> • Preoperative anaemia • Preoperative congenital or acquired coagulopathy • Ongoing anticoagulation therapy or other drug therapy that could cause bleeding • Clinical signs or diagnosis of acute thromboembolism • Emergency surgery • Redo surgery
Interventions	Fibrinogen concentrate 30 mg/kg versus placebo (normal saline solution)
Outcomes	<p>Primary Outcome</p> <ul style="list-style-type: none"> • Perioperative blood loss (during hospital stay up to 30 days) <p>Secondary Outcomes</p> <ul style="list-style-type: none"> • Intraoperative blood loss (from skin incision until wound closure, up to 12 hours) • Perioperative allogeneic red blood cell transfusion (hospital stay up to 30 days) • Perioperative plasma transfusion (hospital stay up to 30 days) • Perioperative platelets transfusion (hospital stay up to 30 days)
Starting date	17 May 2017
Contact information	Yuguang Huang, MD at garybeijingpumch@163.com

NCT03183479 (Continued)

Notes

Estimated study completion date: September 2018

NCT03218722

Trial name or title	Early administration of prothrombin complex concentrate in patients with acute hemorrhage following severe trauma (PROCOAG)
Methods	Randomised controlled trial
Participants	<p>Inclusion Criteria:</p> <ul style="list-style-type: none"> • Primary admission for severe trauma • Out-of-hospital transfusion or red blood cell unit transfusion within the first hour following hospital admission • Clinical prediction of massive transfusion or Assessment of Blood Consumption score ≥ 2 • Informed consent signed by a relative or emergency procedure • Affiliation to a social security system <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> • Cardiac arrest before randomisation • Secondary transfer from another hospital (a technical stop is accepted) • Post-traumatic lesions out of therapeutic resources with death expected in the hour following hospital admission • Anti-coagulation treatment (vitamin K antagonist or novel oral anticoagulant) • Pregnancy • Hypersensitivity to active substances or one of the excipients of KANOKAD® • Patient treated with an experimental medicine within the last 30 days • Decision to place limitations on therapy before randomisation • Patient protected by article L1121-6 and L1121-7 of the French Public health code and patient subject to a legal protection order
Interventions	<p>Pro-Thrombin complex concentrate (25 IU/kg factor IX) versus placebo (normal saline solution)</p> <p>Both study arms will receive conventional trauma management strategy</p>
Outcomes	<p>Primary</p> <ul style="list-style-type: none"> • Blood products transfused in the first 24 hours <p>Secondary</p> <ul style="list-style-type: none"> • Red blood cell units transfused in the first 24 hours following hospital admission. This outcome is measured in number of bags administered • FFP transfused in the first 24 hours following hospital admission. This outcome is measured in number of bags administered • Platelets transfused in the first 24 hours following hospital admission. This outcome is measured in number of bags administered • Time to achieve prothrombin ratio < 1.5 within the first 24 hours • Time to haemostasis within the first 24 hours following admission. Hemostasis is defined as bleeding control in the surgical field or resolution of contrast blush after embolisation during interventional radiology • Thrombo-embolic events during ICU stay (an average of 21 days) • Mortality at 24 hours and Day 28) • ICU-free days during hospital stay (an average of 28 days). Number of in-hospital days outside ICU

NCT03218722 (Continued)

- Ventilator-free days during ICU stay (an average of 21 days). Number of days without mechanical ventilation
- Hospital-free days (Time Frame: Within the first 28 days)Number of days outside hospital
- Glasgow Outcome Scale Extended at day 28
- Hospitalisation status at day 28
- Cost of the strategy at day 8 and day 28

Starting date	29 December 2017
Contact information	Pierre Bouzat, pbouzat@chu-grenoble.fr Anais Adolle, aadolle@chu-grenoble.fr
Notes	Study Sponsor: University Hospital, Grenoble, France

ASA = American Society of Anesthesiologists; CT = computed tomography; EXTEM test = a test which mildly activates haemostasis via the physiological activator tissue factor; FEIBA = Factor Eight Inhibitor Bypassing Activity; FFP = fresh frozen plasma; FIBTEM = a ready-to-use ROTEM® system reagent for use with citrated whole blood and assesses the clot firmness of the fibrin clot; Hb = haemoglobin; HepTEM test = an assay which represents an INTEM assay performed in the presence of heparinase, a heparin- (or LMWH-) degrading enzyme; ICU = intensive care unit; IgA = immunoglobulin A; INR = international normalised ratio; INTEM test = mildly activates the contact phase of haemostasis iv = intravenous; PCC = prothrombin complex concentrate; PICU = paediatric intensive care unit; RIFLE criteria = used to define the spectrum of progressive kidney injury seen in acute kidney injury; ROTEM® = rotational thromboelastometry; SOFA score = Sepsis-Related Organ Failure Assessment score; TACO = transfusion-associated circulatory overload; TBI = traumatic brain injury; TEG = thromboelastography; TRALI = transfusion-related acute lung injury; ULN = upper limit of normal; U = units.

DATA AND ANALYSES

Comparison 1. Fibrinogen: prophylactic trials: intervention vs inactive control

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 All-cause mortality	4		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
1.1 Up to 4 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.2 Up to 7 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.3 Up to 30 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.4 Prior to hospital discharge	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2 Arterial thromboembolic events	7		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
2.1 Up to 3 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.2 Up to 4 days post-operatively	2		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.3 Up to 7 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.4 Up to 30 days post-operatively in patients who have had cardiac surgery	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.5 Up to 30 days post-operatively in patients who had undergone elective surgery	2		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3 Venous thromboembolic events	7		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
3.1 Up to 3 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3.2 Up to day 4 post-operatively	2		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3.3 Up to 7 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3.4 Up to 30 days post-operatively in patients who have had cardiac surgery	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3.5 Up to 30 days post-operatively in participants who had undergone elective surgery	2		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
4 Mortality due to bleeding	4		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
4.1 Up to 3 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
4.2 Up to 7 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
4.3 Up to 30 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
4.4 Prior to hospital discharge	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
5 RBC transfusion requirement	3		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
5.1 Intra-operatively	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
5.2 Up to 12 hours post-operatively	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
5.3 Up to 24 hours post-operatively	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
5.4 In peri-operative period (unclear time-point)	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
6 Receipt of a RBC transfusion	8		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
6.1 Intra-operatively	1	92	Risk Ratio (M-H, Fixed, 95% CI)	1.42 [0.75, 2.68]
6.2 Up to 12 hours post-operatively	1	20	Risk Ratio (M-H, Fixed, 95% CI)	0.33 [0.04, 2.69]
6.3 Up to 24 hours post-operatively	3	207	Risk Ratio (M-H, Fixed, 95% CI)	0.53 [0.37, 0.76]
6.4 Up to 24 hours post-operatively	1	41	Risk Ratio (M-H, Fixed, 95% CI)	0.07 [0.00, 1.22]
6.5 During surgery and up to 24 hours post-operatively	1	92	Risk Ratio (M-H, Fixed, 95% CI)	1.01 [0.76, 1.33]
6.6 Up to 48 hours post-operatively in patients who have had cardiac surgery	1	116	Risk Ratio (M-H, Fixed, 95% CI)	0.59 [0.38, 0.92]
6.7 Up to 48 hours post-operatively in patients who have undergone elective, non-cardiac surgery	2	80	Risk Ratio (M-H, Fixed, 95% CI)	0.23 [0.07, 0.68]
7 Blood loss	7		Ratio of Geometric Means (Fixed, 95% CI)	Subtotals only
7.1 Intraoperative blood loss (mL)	2		Ratio of Geometric Means (Fixed, 95% CI)	0.96 [0.77, 1.21]
7.2 Chest tube drainage at 1 hour post operatively (ml/kg/hr)	1		Ratio of Geometric Means (Fixed, 95% CI)	0.75 [0.52, 1.09]
7.3 Chest tube drainage at 6 hours post-operatively (ml/kg/hr)	1		Ratio of Geometric Means (Fixed, 95% CI)	0.93 [0.76, 1.14]
7.4 Chest tube drainage in first 12 hours post-operatively (mL)	2		Ratio of Geometric Means (Fixed, 95% CI)	0.68 [0.60, 0.76]
7.5 Bleeding up to 24 hours post-operatively (mL)	2		Ratio of Geometric Means (Fixed, 95% CI)	0.83 [0.73, 0.94]
7.6 Bleeding at up to 48 hours post-operatively (mL)	1		Ratio of Geometric Means (Fixed, 95% CI)	0.85 [0.57, 1.27]
7.7 Bleeding during surgery and up to 48 hours post-operatively (mL)	1		Ratio of Geometric Means (Fixed, 95% CI)	0.94 [0.67, 1.33]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
8 Allergic Adverse Events	3		Odds Ratio (M-H, Fixed, 95% CI)	Totals not selected
8.1 Up to 2 days post-operatively	1		Odds Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
8.2 Up to 3 days post-operatively	1		Odds Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
8.3 Up to 30 days post-operatively	1		Odds Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Analysis 1.1. Comparison 1 Fibrinogen: prophylactic trials: intervention vs inactive control, Outcome 1 All-cause mortality.

Study or subgroup	Experimental n/N	Control n/N	Risk Ratio M-H, Fixed, 95% CI	Risk Ratio M-H, Fixed, 95% CI
1.1.1 Up to 4 days post-operatively				
Fenger Erikson 2009	0/10	0/10		Not estimable
1.1.2 Up to 7 days post-operatively				
Karlsson 2011	0/10	0/10		Not estimable
1.1.3 Up to 30 days post-operatively				
Ranucci 2015	1/58	3/58		0.33[0.04,3.11]
1.1.4 Prior to hospital discharge				
Sabate 2016	1/48	3/44		0.31[0.03,2.83]

Analysis 1.2. Comparison 1 Fibrinogen: prophylactic trials: intervention vs inactive control, Outcome 2 Arterial thromboembolic events.

Study or subgroup	Experimental n/N	Control n/N	Risk Ratio M-H, Fixed, 95% CI	Risk Ratio M-H, Fixed, 95% CI
1.2.1 Up to 3 days post-operatively				
Fenger Erikson 2009	0/10	0/10		Not estimable
1.2.2 Up to 4 days post-operatively				
Karlsson 2011	1/10	1/10		1[0.07,13.87]
Sadeghi 2014	0/30	0/30		Not estimable
1.2.3 Up to 7 days post-operatively				
Soleimani 2016	0/31	0/29		Not estimable
1.2.4 Up to 30 days post-operatively in patients who have had cardiac surgery

Analysis 1.3. Comparison 1 Fibrinogen: prophylactic trials: intervention vs inactive control, Outcome 3 Venous thromboembolic events.

Analysis 1.4. Comparison 1 Fibrinogen: prophylactic trials: intervention vs inactive control, Outcome 4 Mortality due to bleeding.

Analysis 1.5. Comparison 1 Fibrinogen: prophylactic trials: intervention vs inactive control, Outcome 5 RBC transfusion requirement.

Analysis 1.6. Comparison 1 Fibrinogen: prophylactic trials: intervention vs inactive control, Outcome 6 Receipt of a RBC transfusion.

Analysis 1.7. Comparison 1 Fibrinogen: prophylactic trials: intervention vs inactive control, Outcome 7 Blood loss.

Analysis 1.8. Comparison 1 Fibrinogen: prophylactic trials: intervention vs inactive control, Outcome 8 Allergic Adverse Events.

Study or subgroup	Experimental n/N	Control n/N	Odds Ratio M-H, Fixed, 95% CI	Odds Ratio M-H, Fixed, 95% CI
1.8.1 Up to 2 days post-operatively				
Soleimani 2016	0/31	0/29		Not estimable
1.8.2 Up to 3 days post-operatively				
Fenger Erikson 2009	0/10	0/10		Not estimable
1.8.3 Up to 30 days post-operatively				
Ranucci 2015	0/58	0/58		Not estimable

Favours fibrinogen 0.01 0.1 1 10 100 Favours inactive control

Comparison 2. Fibrinogen: prophylactic trials: intervention vs haemostatically active control

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 All-cause mortality	1		Odds Ratio (M-H, Fixed, 95% CI)	Totals not selected
1.1 Duration of hospital stay (whole population)	1		Odds Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2 Arterial thromboembolic events	1		Odds Ratio (M-H, Fixed, 95% CI)	Totals not selected
2.1 Duration of hospital stay (whole population)	1		Odds Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3 Venous thromboembolic events	1		Odds Ratio (M-H, Fixed, 95% CI)	Totals not selected
3.1 Duration of hospital stay (whole population)	1		Odds Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
4 Mortality due to bleeding	1		Odds Ratio (M-H, Fixed, 95% CI)	Totals not selected
4.1 Duration of hospital stay (whole population)	1		Odds Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
5 RBC transfusion requirement	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
5.1 Surgical time plus 24 hours post-operatively - Craniosynostosis (ml/kg)	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
5.2 Surgical time plus 24 hours post-operatively - Scoliosis (ml/kg)	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
6 Receipt of a RBC transfusion	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
6.1 Surgical time plus 24 hours post-operatively - Craniosynostosis	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
6.2 Surgical time plus 24 hours post-operatively - Scoliosis	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
7 Allergic Adverse Events	1		Odds Ratio (M-H, Fixed, 95% CI)	Totals not selected
7.1 Up to 24 hours post-operatively (whole population)	1		Odds Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Analysis 2.1. Comparison 2 Fibrinogen: prophylactic trials: intervention vs haemostatically active control, Outcome 1 All-cause mortality.

Study or subgroup	Experimental n/N	Control n/N	Odds Ratio M-H, Fixed, 95% CI	Odds Ratio M-H, Fixed, 95% CI
2.1.1 Duration of hospital stay (whole population)				
Haas 2015	0/27	0/22		Not estimable
			Favours fibrinogen 0.01 0.1 1 10 100	Favours active control

Analysis 2.2. Comparison 2 Fibrinogen: prophylactic trials: intervention vs haemostatically active control, Outcome 2 Arterial thromboembolic events.

Study or subgroup	Experimental n/N	Control n/N	Odds Ratio M-H, Fixed, 95% CI	Odds Ratio M-H, Fixed, 95% CI
2.2.1 Duration of hospital stay (whole population)				
Haas 2015	0/27	0/22		Not estimable
			Favours fibrinogen 0.01 0.1 1 10 100	Favours active control

Analysis 2.3. Comparison 2 Fibrinogen: prophylactic trials: intervention vs haemostatically active control, Outcome 3 Venous thromboembolic events.

Study or subgroup	Experimental n/N	Control n/N	Odds Ratio M-H, Fixed, 95% CI	Odds Ratio M-H, Fixed, 95% CI
2.3.1 Duration of hospital stay (whole population)				
Haas 2015	0/27	0/22		Not estimable
			Favours fibrinogen 0.01 0.1 1 10 100	Favours active control

Analysis 2.4. Comparison 2 Fibrinogen: prophylactic trials: intervention vs haemostatically active control, Outcome 4 Mortality due to bleeding.

Study or subgroup	Experimental n/N	Control n/N	Odds Ratio M-H, Fixed, 95% CI	Odds Ratio M-H, Fixed, 95% CI
2.4.1 Duration of hospital stay (whole population)				
Haas 2015	0/27	0/22		Not estimable
			Favours fibrinogen 0.01 0.1 1 10 100	Favours active control

Analysis 2.5. Comparison 2 Fibrinogen: prophylactic trials: intervention vs haemostatically active control, Outcome 5 RBC transfusion requirement.

Study or subgroup	Experimental		Control		Mean Difference Fixed, 95% CI	Mean Difference Fixed, 95% CI
	N	Mean(SD)	N	Mean(SD)		
2.5.1 Surgical time plus 24 hours post-operatively - Craniostomosis (ml/kg)						
Haas 2015	17	33.1 (17.9)	13	41.8 (18.1)		-8.7[-21.71,4.31]
					Favours fibrinogen -100 -50 0 50 100	Favours active control

Analysis 2.6. Comparison 2 Fibrinogen: prophylactic trials: intervention vs haemostatically active control, Outcome 6 Receipt of a RBC transfusion.

Analysis 2.7. Comparison 2 Fibrinogen: prophylactic trials: intervention vs haemostatically active control, Outcome 7 Allergic Adverse Events.

Comparison 3. Fibrinogen: therapeutic trials: intervention vs inactive control

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 All-cause mortality	7		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Up to 28 days following admission	2	97	Risk Ratio (M-H, Fixed, 95% CI)	1.46 [0.71, 2.99]
1.2 In-hospital mortality up to 30 days post-operatively	1	120	Risk Ratio (M-H, Fixed, 95% CI)	5.0 [0.25, 102.00]
1.3 Up to 6 weeks postnatally	2	294	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.4 Up to 46 days post-operatively	2	213	Risk Ratio (M-H, Fixed, 95% CI)	0.23 [0.05, 1.01]
2 Arterial thromboembolic events	7		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
2.1 During hospital stay	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.2 Up to 28 days following admission	2		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.3 Up to 30 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.4 Up to 45 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.5 Up to 6 weeks postnatally	2		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3 Venous thromboembolic events	6		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
3.1 During hospital stay	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3.2 Up to 28 days following admission	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3.3 Up to 30 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3.4 Up to 45 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3.5 Up to 6 weeks postnatally	2		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
4 Mortality due to bleeding	5		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
4.1 Up to 28 days following admission	2	93	Risk Ratio (M-H, Fixed, 95% CI)	2.45 [0.38, 15.76]
4.2 Up to 6 weeks postnatally	2	294	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
4.3 Up to 46 days post-operatively	1	152	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
5 Receipt of a RBC transfusion	6		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
5.1 Intraoperative: Between study medication and chest closure	1	117	Risk Ratio (M-H, Fixed, 95% CI)	0.15 [0.01, 2.75]
5.2 Up to 3 hours following admission	1	46	Risk Ratio (M-H, Fixed, 95% CI)	1.05 [0.86, 1.29]
5.3 Within 4 hours of receipt of the study drug	1	244	Risk Ratio (M-H, Fixed, 95% CI)	0.39 [0.13, 1.22]
5.4 Up to 6 hours following admission	1	43	Risk Ratio (M-H, Fixed, 95% CI)	1.05 [0.84, 1.30]
5.5 Up to 24 hours following admission	1	41	Risk Ratio (M-H, Fixed, 95% CI)	1.11 [0.91, 1.36]
5.6 Within 24 hours of receipt of study drug in patients who have had cardiac surgery	2	178	Risk Ratio (M-H, Fixed, 95% CI)	0.54 [0.39, 0.75]
5.7 Within 24 hours of receipt of study drug in women who have ex-	1	244	Risk Ratio (M-H, Fixed, 95% CI)	0.72 [0.38, 1.38]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
perienced a post-partum haemorrhage				
5.8 Within 7 days of receipt of the study drug	1	244	Risk Ratio (M-H, Fixed, 95% CI)	0.95 [0.58, 1.54]
5.9 From end of intervention to hospital discharge	1	55	Risk Ratio (M-H, Fixed, 95% CI)	0.96 [0.55, 1.69]
5.10 During hospital stay	1	48	Risk Ratio (M-H, Fixed, 95% CI)	1.0 [0.38, 2.66]
6 RBC transfusion requirement	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
6.1 During hospital stay	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
7 Blood loss	1		Ratio of Geometric Mean (Fixed, 95% CI)	Totals not selected
7.1 Intra-operative	1		Ratio of Geometric Mean (Fixed, 95% CI)	0.0 [0.0, 0.0]
7.2 Post-operative (within 12 hours)	1		Ratio of Geometric Mean (Fixed, 95% CI)	0.0 [0.0, 0.0]
7.3 Intra- and post-operative (within 12 hours)	1		Ratio of Geometric Mean (Fixed, 95% CI)	0.0 [0.0, 0.0]
8 Allergic Adverse Events	4		Risk Difference (M-H, Fixed, 95% CI)	Totals not selected
8.1 Up to 24 hours from study medication	1		Risk Difference (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
8.2 Up to 10 days from study medication	1		Risk Difference (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
8.3 Up to 28 days following admission	1		Risk Difference (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
8.4 Up to 30 days from study medication	1		Risk Difference (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Analysis 3.1. Comparison 3 Fibrinogen: therapeutic trials: intervention vs inactive control, Outcome 1 All-cause mortality.

Study or subgroup	Experimental n/N	Control n/N	Risk Ratio M-H, Fixed, 95% CI	Weight	Risk Ratio M-H, Fixed, 95% CI
3.1.1 Up to 28 days following admission					
Curry 2018	10/24	7/24		78.13%	1.43[0.65,3.13]

Analysis 3.2. Comparison 3 Fibrinogen: therapeutic trials: intervention vs inactive control, Outcome 2 Arterial thromboembolic events.

Study or subgroup	Experimental n/N	Control n/N	Risk Ratio M-H, Fixed, 95% CI	Risk Ratio M-H, Fixed, 95% CI
Collins 2017	0/26	0/24		Not estimable
Wikkelso 2015	0/123	0/121		Not estimable
Favours fibrinogen 0.01 0.1 1 10 100 Favours inactive control				

Analysis 3.3. Comparison 3 Fibrinogen: therapeutic trials: intervention vs inactive control, Outcome 3 Venous thromboembolic events.

Study or subgroup	Experimental n/N	Control n/N	Risk Ratio M-H, Fixed, 95% CI	Risk Ratio M-H, Fixed, 95% CI
3.3.1 During hospital stay				
Jeppsson 2016	0/24	0/24		Not estimable
3.3.2 Up to 28 days following admission				
Curry 2018	2/20	0/19		4.76[0.24,93.19]
3.3.3 Up to 30 days post-operatively				
Bilecen 2017	0/60	0/60		Not estimable
3.3.4 Up to 45 days post-operatively				
Rahe-Meyer 2013	0/29	1/32		0.37[0.02,8.66]
3.3.5 Up to 6 weeks postnatally				
Collins 2017	1/26	1/24		0.92[0.06,13.95]
Wikkelso 2015	0/123	0/121		Not estimable
Favours fibrinogen 0.002 0.1 1 10 500 Favours inactive control				

Analysis 3.4. Comparison 3 Fibrinogen: therapeutic trials: intervention vs inactive control, Outcome 4 Mortality due to bleeding.

Study or subgroup	Experimental n/N	Control n/N	Risk Ratio M-H, Fixed, 95% CI	Weight	Risk Ratio M-H, Fixed, 95% CI
3.4.1 Up to 28 days following admission					
Curry 2018	2/24	1/24		68.12%	2[0.19,20.61]
Nascimento 2016	1/21	0/24		31.88%	3.41[0.15,79.47]
Subtotal (95% CI)	45	48		100%	2.45[0.38,15.76]
Total events: 3 (Experimental), 1 (Control)					
Heterogeneity: Tau ² =0; Chi ² =0.07, df=1(P=0.79); I ² =0%					
Test for overall effect: Z=0.94(P=0.35)					
3.4.2 Up to 6 weeks postnatally					
Collins 2017	0/26	0/24			Not estimable
Wikkelso 2015	0/123	0/121			Not estimable
Subtotal (95% CI)	149	145			Not estimable
Total events: 0 (Experimental), 0 (Control)					
Heterogeneity: Not applicable					
Test for overall effect: Not applicable					
Favours fibrinogen 0.002 0.1 1 10 500 Favours inactive control					

Study or subgroup	Experimental n/N	Control n/N	Risk Ratio M-H, Fixed, 95% CI	Weight	Risk Ratio M-H, Fixed, 95% CI
3.4.3 Up to 46 days post-operatively					
Rahe-Meyer 2016	0/78	0/74			Not estimable
Subtotal (95% CI)	78	74			Not estimable
Total events: 0 (Experimental), 0 (Control)					
Heterogeneity: Not applicable					
Test for overall effect: Not applicable					
Test for subgroup differences: Not applicable					
<div> <div>Favours fibrinogen</div> <div>0.002 0.1 1 10 500</div> <div>Favours inactive control</div> </div>					

Analysis 3.5. Comparison 3 Fibrinogen: therapeutic trials: intervention vs inactive control, Outcome 5 Receipt of a RBC transfusion.

Study or subgroup	Active n/N	Inactive n/N	Risk Ratio M-H, Fixed, 95% CI	Weight	Risk Ratio M-H, Fixed, 95% CI
3.5.1 Intraoperative: Between study medication and chest closure					
Bilecen 2017	0/58	3/59		100%	0.15[0.01,2.75]
Subtotal (95% CI)	58	59		100%	0.15[0.01,2.75]
Total events: 0 (Active), 3 (Inactive)					
Heterogeneity: Not applicable					
Test for overall effect: Z=1.29(P=0.2)					
3.5.2 Up to 3 hours following admission					
Curry 2018	21/23	20/23		100%	1.05[0.86,1.29]
Subtotal (95% CI)	23	23		100%	1.05[0.86,1.29]
Total events: 21 (Active), 20 (Inactive)					
Heterogeneity: Not applicable					
Test for overall effect: Z=0.47(P=0.64)					
3.5.3 Within 4 hours of receipt of the study drug					
Wikkelso 2015	4/123	10/121		100%	0.39[0.13,1.22]
Subtotal (95% CI)	123	121		100%	0.39[0.13,1.22]
Total events: 4 (Active), 10 (Inactive)					
Heterogeneity: Not applicable					
Test for overall effect: Z=1.61(P=0.11)					
3.5.4 Up to 6 hours following admission					
Curry 2018	19/21	19/22		100%	1.05[0.84,1.3]
Subtotal (95% CI)	21	22		100%	1.05[0.84,1.3]
Total events: 19 (Active), 19 (Inactive)					
Heterogeneity: Not applicable					
Test for overall effect: Z=0.42(P=0.67)					
3.5.5 Up to 24 hours following admission					
Curry 2018	19/20	18/21		100%	1.11[0.91,1.36]
Subtotal (95% CI)	20	21		100%	1.11[0.91,1.36]
Total events: 19 (Active), 18 (Inactive)					
Heterogeneity: Not applicable					
Test for overall effect: Z=1(P=0.32)					
<div> <div>Favours fibrinogen</div> <div>0.01 0.1 1 10 100</div> <div>Favours inactive control</div> </div>

Analysis 3.6. Comparison 3 Fibrinogen: therapeutic trials: intervention vs inactive control, Outcome 6 RBC transfusion requirement.

Analysis 3.7. Comparison 3 Fibrinogen: therapeutic trials: intervention vs inactive control, Outcome 7 Blood loss.

Analysis 3.8. Comparison 3 Fibrinogen: therapeutic trials: intervention vs inactive control, Outcome 8 Allergic Adverse Events.

Comparison 4. Fibrinogen: therapeutic trials: intervention vs haemostatically active control

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
1 All-cause mortality	4		Risk Ratio (M-H, Fixed, 95% CI)	Totals not select- ed
1.1 Up to 7 days post-operatively or hospital discharge	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.2 Up to 28 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.3 Up to 30 days post-operatively in patients who have undergone major elective surgery	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.4 Up to 30 days post-operatively in trauma patients	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2 Arterial thromboembolic events	3		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
2.1 Up to 7 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.2 Up to 28 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.3 Up to 30 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3 Venous thromboembolic events	4		Odds Ratio (M-H, Fixed, 95% CI)	Totals not selected
3.1 Up to 7 days post-operatively	1		Odds Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3.2 Up to 28 days post-operatively	1		Odds Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3.3 Up to 30 days post-operatively	1		Odds Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3.4 Up to day 30 following trauma	1		Odds Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
4 Mortality due to bleeding	4		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
4.1 Up to 7 days post-operatively	1	63	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
4.2 Up to 30 days post-operatively	2	63	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
4.3 Up to day 30 following trauma	1	94	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
5 RBC transfusion requirement	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
5.1 Up to 30 days post-operatively	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
6 Receipt of a RBC transfusion	3		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
6.1 Up to 24 hours post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
6.2 Up to 24 hours following trauma	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
6.3 Up to 7 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
7 Allergic Adverse Events	2		Odds Ratio (M-H, Fixed, 95% CI)	Totals not selected
7.1 Up to 7 days post-operatively	1		Odds Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
7.2 Up to 30 days post-operatively	1		Odds Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Analysis 4.1. Comparison 4 Fibrinogen: therapeutic trials: intervention vs haemostatically active control, Outcome 1 All-cause mortality.

Analysis 4.2. Comparison 4 Fibrinogen: therapeutic trials: intervention vs haemostatically active control, Outcome 2 Arterial thromboembolic events.

Analysis 4.3. Comparison 4 Fibrinogen: therapeutic trials: intervention vs haemostatically active control, Outcome 3 Venous thromboembolic events.

Analysis 4.4. Comparison 4 Fibrinogen: therapeutic trials: intervention vs haemostatically active control, Outcome 4 Mortality due to bleeding.

Analysis 4.5. Comparison 4 Fibrinogen: therapeutic trials: intervention vs haemostatically active control, Outcome 5 RBC transfusion requirement.

Analysis 4.6. Comparison 4 Fibrinogen: therapeutic trials: intervention vs haemostatically active control, Outcome 6 Receipt of a RBC transfusion.

Analysis 4.7. Comparison 4 Fibrinogen: therapeutic trials: intervention vs haemostatically active control, Outcome 7 Allergic Adverse Events.

Comparison 5. FXIII: prophylactic trials: intervention vs inactive control

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 All-cause mortality	5		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
1.1 Until hospital discharge	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.2 Up to 5 to 7 weeks post-operatively	2		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.3 Up to 340 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.4 Time point not stated	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2 Arterial thromboembolic events	4		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
2.1 Up to 5 to 7 weeks post-operatively	2		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.2 Up to 30 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.3 Post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3 Venous thromboembolic events	4		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
3.1 Up to 5 to 7 weeks post-operatively	2		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3.2 Up to 30 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3.3 Post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
4 Mortality due to bleeding	4		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
4.1 Up to a median of 32 days from study intervention	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
4.2 Until hospital discharge	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
4.3 Up to 5 to 7 weeks post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
4.4 Up to 340 days post-operatively	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
5 RBC transfusion requirement	1		Mean Difference (IV, Fixed, 95% CI)	Totals not selected
5.1 From the point of study medication	1		Mean Difference (IV, Fixed, 95% CI)	0.0 [0.0, 0.0]
6 Receipt of a RBC transfusion	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
6.1 Up to 7 days post-operatively (or discharge)	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
7 Blood loss	2		Ratio of Geometric Mean (Fixed, 95% CI)	Subtotals only
7.1 Chest tube drainage at 8 hours post-operatively (mL)	2		Ratio of Geometric Mean (Fixed, 95% CI)	0.79 [0.65, 0.97]
7.2 Chest drain volume at 12 hours post-operatively (mL)	1		Ratio of Geometric Mean (Fixed, 95% CI)	0.78 [0.63, 0.97]
7.3 Chest tube drainage at 24 hours post-operatively (mL)	2		Ratio of Geometric Mean (Fixed, 95% CI)	0.85 [0.72, 1.00]
7.4 Chest drain volume at 36 hours post-operatively (mL)	1		Ratio of Geometric Mean (Fixed, 95% CI)	0.82 [0.68, 0.98]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
7.5 Chest drain volume at 48 hours post-operatively (mL)	1		Ratio of Geometric Mean (Fixed, 95% CI)	0.73 [0.62, 0.86]
7.6 Chest drainage at drain removal (mL)	1		Ratio of Geometric Mean (Fixed, 95% CI)	0.68 [0.32, 1.47]
8 Allergic Adverse Events	5		Odds Ratio (M-H, Fixed, 95% CI)	Subtotals only
8.1 Unclear timepoint	5	206	Odds Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Analysis 5.1. Comparison 5 FXIII: prophylactic trials: intervention vs inactive control, Outcome 1 All-cause mortality.

Study or subgroup	Experimental n/N	Control n/N	Risk Ratio M-H, Fixed, 95% CI	Risk Ratio M-H, Fixed, 95% CI
5.1.1 Until hospital discharge				
Godje 2006	0/25	0/25		Not estimable
5.1.2 Up to 5 to 7 weeks post-operatively				
Karkouti 2013	1/138	0/128		2.78[0.11,67.73]
Levy 2009	0/8	0/8		Not estimable
5.1.3 Up to 340 days post-operatively				
Korte 2009	3/11	3/11		1[0.26,3.91]
5.1.4 Time point not stated				
Rasche 1982	9/29	9/31		1.07[0.49,2.32]
<div style="display: flex; justify-content: space-between; align-items: center;"> Favours FXIII 0.005 0.1 1 10 200 Favours inactive control </div>				

Analysis 5.2. Comparison 5 FXIII: prophylactic trials: intervention vs inactive control, Outcome 2 Arterial thromboembolic events.

Study or subgroup	Experimental n/N	Control n/N	Risk Ratio M-H, Fixed, 95% CI	Risk Ratio M-H, Fixed, 95% CI
5.2.1 Up to 5 to 7 weeks post-operatively				
Karkouti 2013	8/138	11/128		0.67[0.28,1.62]
Levy 2009	0/8	0/8		Not estimable
5.2.2 Up to 30 days post-operatively				
Korte 2009	1/11	0/11		3[0.14,66.53]
5.2.3 Post-operatively				
Godje 2006	0/25	2/25		0.2[0.01,3.97]
<div style="display: flex; justify-content: space-between; align-items: center;"> Favours FXIII 0.002 0.1 1 10 500 Favours inactive control </div>				

Analysis 5.3. Comparison 5 FXIII: prophylactic trials: intervention vs inactive control, Outcome 3 Venous thromboembolic events.

Analysis 5.4. Comparison 5 FXIII: prophylactic trials: intervention vs inactive control, Outcome 4 Mortality due to bleeding.

Analysis 5.5. Comparison 5 FXIII: prophylactic trials: intervention vs inactive control, Outcome 5 RBC transfusion requirement.

Analysis 5.6. Comparison 5 FXIII: prophylactic trials: intervention vs inactive control, Outcome 6 Receipt of a RBC transfusion.

Study or subgroup	Experimental n/N	Control n/N	Risk Ratio M-H, Fixed, 95% CI	Risk Ratio M-H, Fixed, 95% CI
5.6.1 Up to 7 days post-operatively (or discharge)				
Karkouti 2013	47/138	45/128		0.97[0.7,1.35]
			Favours FXIII	Favours inactive control

Analysis 5.7. Comparison 5 FXIII: prophylactic trials: intervention vs inactive control, Outcome 7 Blood loss.

Study or subgroup	Experi- mental N	Control N	log[Ratio of Geometric Mean] (SE)	Ratio of Geometric Mean IV, Fixed, 95% CI	Weight	Ratio of Geo- metric Mean IV, Fixed, 95% CI
5.7.1 Chest tube drainage at 8 hours post-operatively (mL)						
Godje 2006	0	0	-0.2 (0.115)		77.71%	0.79[0.63,0.99]
Levy 2009	0	0	-0.2 (0.215)		22.29%	0.8[0.52,1.22]
Subtotal (95% CI)					100%	0.79[0.65,0.97]
Heterogeneity: Tau ² =0; Chi ² =0, df=1(P=0.97); I ² =0%						
Test for overall effect: Z=2.28(P=0.02)						
5.7.2 Chest drain volume at 12 hours post-operatively (mL)						
Godje 2006	0	0	-0.2 (0.111)		100%	0.78[0.63,0.97]
Subtotal (95% CI)					100%	0.78[0.63,0.97]
Heterogeneity: Tau ² =0; Chi ² =0, df=0(P<0.0001); I ² =100%						
Test for overall effect: Z=2.25(P=0.02)						
5.7.3 Chest tube drainage at 24 hours post-operatively (mL)						
Godje 2006	0	0	-0.2 (0.089)		93.41%	0.86[0.72,1.02]
Levy 2009	0	0	-0.3 (0.337)		6.59%	0.74[0.38,1.43]
Subtotal (95% CI)					100%	0.85[0.72,1.02]
Heterogeneity: Tau ² =0; Chi ² =0.18, df=1(P=0.67); I ² =0%						
Test for overall effect: Z=1.92(P=0.06)						
5.7.4 Chest drain volume at 36 hours post-operatively (mL)						
Godje 2006	0	0	-0.2 (0.091)		100%	0.82[0.68,0.98]
Subtotal (95% CI)					100%	0.82[0.68,0.98]
Heterogeneity: Not applicable						
Test for overall effect: Z=2.21(P=0.03)						
5.7.5 Chest drain volume at 48 hours post-operatively (mL)						
Godje 2006	0	0	-0.3 (0.081)		100%	0.73[0.62,0.86]
Subtotal (95% CI)					100%	0.73[0.62,0.86]
Heterogeneity: Not applicable						
Test for overall effect: Z=3.87(P=0)						
5.7.6 Chest drainage at drain removal (mL)						
Levy 2009	0	0	-0.4 (0.391)		100%	0.68[0.32,1.47]
Subtotal (95% CI)					100%	0.68[0.32,1.47]
Heterogeneity: Not applicable						
Test for overall effect: Z=0.97(P=0.33)						

Study or subgroup	Experimental N	Control N	log[Ratio of Geometric Mean] (SE)	Ratio of Geometric Mean IV, Fixed, 95% CI	Weight	Ratio of Geo- metric Mean IV, Fixed, 95% CI
Test for subgroup differences: $\chi^2=1.85$, $df=1$ ($P=0.87$), $I^2=0\%$						
				0.05 0.2 1 5 20		Favours FXIII Favours inactive control

Analysis 5.8. Comparison 5 FXIII: prophylactic trials: intervention vs inactive control, Outcome 8 Allergic Adverse Events.

Study or subgroup	Experimental n/N	Control n/N	Odds Ratio M-H, Fixed, 95% CI	Weight	Odds Ratio M-H, Fixed, 95% CI
5.8.1 Unclear timepoint					
Godje 2006	0/25	0/25			Not estimable
Korte 2009	0/11	0/11			Not estimable
Levy 2009	0/8	0/8			Not estimable
Rasche 1982	0/29	0/31			Not estimable
Shirahata 1990	0/30	0/28			Not estimable
Subtotal (95% CI)	103	103			Not estimable
Total events: 0 (Experimental), 0 (Control)					
Heterogeneity: Not applicable					
Test for overall effect: Not applicable					
			0.01 0.1 1 10 100		Favours FXIII Favours inactive control

Comparison 6. FXIII: therapeutic trials: intervention vs inactive control

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Allergic Adverse Events	1		Odds Ratio (M-H, Fixed, 95% CI)	Totals not selected
1.1 Timepoint not stated	1		Odds Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Analysis 6.1. Comparison 6 FXIII: therapeutic trials: intervention vs inactive control, Outcome 1 Allergic Adverse Events.

Study or subgroup	Experimental n/N	Control n/N	Odds Ratio M-H, Fixed, 95% CI	Weight	Odds Ratio M-H, Fixed, 95% CI
6.1.1 Timepoint not stated					
Bregenzer 1999	0/17	0/11			Not estimable
			0.01 0.1 1 10 100		Favours FXIII Favours inactive control

Comparison 7. PCC: prophylactic trials: intervention vs inactive control

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 All-cause mortality	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
1.1 Time point not stated	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2 Allergic Adverse Events	1		Odds Ratio (M-H, Fixed, 95% CI)	Totals not selected
2.1 Timepoint not stated	1		Odds Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Analysis 7.1. Comparison 7 PCC: prophylactic trials: intervention vs inactive control, Outcome 1 All-cause mortality.

Study or subgroup	Experimental n/N	Control n/N	Risk Ratio M-H, Fixed, 95% CI	Risk Ratio M-H, Fixed, 95% CI
7.1.1 Time point not stated				
Turner 1981	23/39	22/39		1.05[0.71,1.53]
			Favours PCC	Favours inactive control

Analysis 7.2. Comparison 7 PCC: prophylactic trials: intervention vs inactive control, Outcome 2 Allergic Adverse Events.

Study or subgroup	Experimental n/N	Control n/N	Odds Ratio M-H, Fixed, 95% CI	Odds Ratio M-H, Fixed, 95% CI
7.2.1 Timepoint not stated				
Turner 1981	0/39	0/39		Not estimable
			Favours PCC	Favours inactive control

ADDITIONAL TABLES

Table 1. Red blood cell transfusion requirement

Trial	Number included in analyses	Clinical condition	Transfusion Protocol?	When was transfusion requirement measured	How were findings reported	Findings: Intervention/Comparator
Fibrinogen concentrate: prophylactic trials with inactive comparator						
Cui 2010	17/14	Surgery (E) : cardiac	Not stated	During ICU stay	Median (IQR) number of red blood cell units transfused	0 (0 - 0)/0 (0 - 1.25)
					Median (IQR) total red blood cell usage	0 (0 - 1.5)/1.5 (0 - 2.3)

Table 1. Red blood cell transfusion requirement (Continued)

Fenger Erikson 2009	10/10	Surgery (E): radical cystectomy	Not stated	During surgery	Median (IQR) number of red blood cell units transfused	2 (0 - 5) / 2.5 (0 - 6)
				48 hours post-operatively		0 (0 - 2)/1.5 (0 - 2)
				Total transfused		3.5 (0 - 5)/4.0 (0 - 6)
				Within 48 hours post-operatively	Total number (%) patients requiring a red blood cell transfusion	2 (20%)/8 (80%)
Karlsson 2011	10/10	Surgery (E) : cardiac	Yes	Within 12 hours post-operatively	Mean (SD) number of red blood cell units transfused	0.2 (0.13)/0.7 (0.27)
Najafi 2014	15/15	Surgery (E): total hip replacement arthroplasty	Yes	"Perioperative period"	Mean (SD) "sum of transfused packed cells"	0.8 (1.01)/1.06 (1.2)
Pournajafian 2015	21/20	Surgery (E): posterior spinal fusion	Not stated	Up to 24 hours post-operatively	Number (%) patients transfused	0 (0%)/6 (30%)
Ranucci 2015	58/58	Surgery (E): cardiac	Yes	Duration of hospital stay until 30 days	Total number (%) of patients receiving a red blood cell transfusion	19 (33%)/32 (55%)
					Median (IQR) number of units transfused	0 (0 - 1)/1 (0 - 2)
				Within 24 hours of surgery	Mean (SD) mL of red blood cells infused for transfused patients only	136 (178)/356 (488)
				From 24 to 48 hours after surgery	Mean (SD) mL of red blood cells infused for transfused patients only	420 (760)/218 (552)
Sabate 2016	48/44	Surgery (E): liver transplantation	Yes	During surgery	Median (IQR) number of units transfused	0 (0 - 1)/0 (0 - 0.5)
					Total number (%) of patients receiving a red blood cell transfusion	17 (52.9%)/11 (42.7%)
				During and up to 24 hours after surgery	Median (IQR) number of units transfused	2 (0 - 6)/3 (0 - 6)
					Total number (%) of patients receiving a red blood cell transfusion	33 (68.8%)/30 (68.2%)

Table 1. Red blood cell transfusion requirement (Continued)

Sadeghi 2014	30/30	Surgery (E) : cardiac	Yes	Within first 24 hours	Mean (SD) number of red blood cell infusions	1.5 (1.8)/2 (1.5)
					Total number (%) of patients re- ceiving a red blood cell transfu- sion	15 (50)/21 (70)
Soleimani 2016	31/29	Surgery (E): transurethral resec- tion of the prostate.	Not stated	post-operatively	Total number (%) of patients re- ceiving a red blood cell transfu- sion	1 (3.2%)/5 (17.2%)
Fibrinogen concentrate: prophylactic trials with active comparator						
Haas 2015	10/9	Surgery (E): scoliosis	Yes	Over 24 hours from the start of surgery	Median (IQR) mL/kg of red blood cells transfused	0 (0 - 15.3)/18.4 (0 - 23.8)
					Mean (SD) mL/kg of red blood cells transfused	3.7 (6.2)/7.2 (8.0)
					Total number (%) of patients re- ceiving a red blood cell transfu- sion	4 (40%)/6 (67%)
	17/ 13	Surgery (E): craniosyn- ostosis	Yes	Over 24 hours from the start of surgery	Median (IQR) mL/kg of red blood cells transfused	28.2 (21.2- 49.9)/55.5 (27.5 - 61.8)
					Mean (SD) mL/kg of red blood cells transfused	33.1 (17.9)/41.8 (18.1)
					Total number (%) of patients re- ceiving a red blood cell transfu- sion	17 (100%)/13 (100%)
Fibrinogen concentrate: therapeutic trials with inactive comparator						
Bilecen 2017	58/59	Surgery (E): cardiac	Yes	Between study intervention and chest closure	Total number (%) of patients re- ceiving a red blood cell transfu- sion	0 (0%)/3 (5%)
				Between study intervention and 24 hours	Total number (%) of patients re- ceiving a red blood cell transfu- sion	10 (17%)/20 (33%)
					Median (IQR) number of units transfused	0 (0 - 1)/0 (0 - 4)
Collins 2017	28/27	postpartum haemor- rhage	Not stated	From study in- tervention to dis- charge	Median (IQR) number of units transfused	1 (0 - 2)/1 (0 - 2)
					Total number (%) of patients who received a red blood cell transfu- sion	13 (46.6%)/13 (48.1%)

Table 1. Red blood cell transfusion requirement (Continued)

					Total number of red blood cell transfusions	37/38
Curry 2018	24/24	Trauma	Yes	3hrs following admission	Median (IQR) number of units transfused	4 (2 - 6)/2 (2 - 6)
					Total number (%) of patients receiving a red blood cell transfusion	21 (91.3%)/20 (87.0%)
				6hrs following admission	Median (IQR) number of units transfused	3 (2 - 6)/2 (2 - 6)
					Total number (%) of patients receiving a red blood cell transfusion	19 (90.5%)/19 (86.4%)
				24hrs following admission	Median (IQR) number of units transfused	4 (2 - 8)/2 (2 - 5)
					Total number (%) of patients receiving a red blood cell transfusion	19 (95.0%)/18 (85.7%)
Jeppsson 2016	24/24	Surgery (E): cardiac	Yes	During hospital stay	Mean (SD) units of red blood cells transfused	0.63 (1.17)/1.33 (3.11)
					Median (IQR) number of units transfused	0 (0 - 1)/0 (0 - 1)
Nascimben-to 2016	21/24	Trauma	Yes	At 24 hours	Median (IQR) number of units transfused	3 (2 - 5)/3 (2 - 4)
Rahe-Meyer 2013	29/32	Surgery (E): aortic valve replacement	Yes	24 hours after receipt of study medication	Total number (%) of patients receiving a red blood cell transfusion	16 (55)/32 (100)
					Median (IQR) number of units transfused	0 (0 - 3)/2 (2-5)
Rahe-Meyer 2016	78/74	Surgery (E): complex cardiac	Yes	24 hours after receipt of study medication	Median (IQR) number of units transfused	1 (0 - 3)/0 (0 - 2)
Wikkelso 2015	123/121	postpartum haemorrhage	Yes	Need for a red blood cell transfusion up to 4 hours post study drug	Total number (%) of patients receiving a red blood cell transfusion	4 (3)/10 (8)
				Need for a red blood cell transfusion up to 24 hours post study drug		14 (11)/19 (16)
				Need for a red blood cell trans-		25 (20)/26 (21)

Table 1. Red blood cell transfusion requirement (Continued)

				fusion up to 7 days post study drug		
				"During the 6 week period postpartum"		25 (20)/26 (21)
				Time-point not stated	Total number (%) of patients who received a transfusion of > 4 units of red blood cells.	8 (7)/3 (3)
Fibrinogen concentrate: therapeutic trials with active comparator						
Galas 2014	30/33	Surgery (E): cardiac	Yes	post-operative day 7	Total number (%) of patients receiving a red blood cell transfusion	25 (83)/32 (97)
Tanaka 2014	10/10	Surgery (E): valve replacement or repair	Yes	At 24 hours post-operatively	Total number (%) of patients receiving a red blood cell transfusion	9 (90)/9 (90)
Lance 2012	22/21	Surgery (E): cardiovascular, abdominal or spinal column	Yes	Following study intervention	Mean (SD) number of red blood cell units transfused	4.98 (3.01)/5.38 (2.38)
Innerhofer 2017	50/44	Trauma	Yes	At 24 hours after injury	Total number (%) of patients receiving a red blood cell transfusion	25 (90%)/39 (89%)
					Median (IQR) number of red blood cell units transfused	4 (2 - 7)/6 (4 - 11)
Factor XIII (FXIII): prophylactic trials with inactive comparator						
Godje 2006	25/25	Surgery (E) : cardiac	Yes	At 24 hours post-operatively	Mean (SD) number of red blood cell units transfused	2.6 (1.875)/3.3 (1.397)
Karkouti 2013	138/128	Surgery (E) : cardiac	Not stated	Until post-operative day 7 or hospital discharge (whichever occurred first)	Median (IQR) number of red blood cell units transfused	0 (0-1)/0 (0-1)
					Total number (%) of patients receiving a red blood cell transfusion	46 (33)/43 (34)
Korte 2009	22 in total	Surgery (E): gastrointestinal cancer	Yes	Unclear time point	Median (IQR) mL of red blood cells transfused	150 (0-700)/200 (0-800)
Levy 2009	8/8	Surgery (E) : cardiac	Not stated	Until post-operative day 7 or hospital discharge (whichever occurred first)	Total number of patients requiring allogeneic blood products	No data was reported

Table 1. Red blood cell transfusion requirement (Continued)

					Percentage of patients avoiding a red blood cell transfusion	No data was reported
Rasche 1982	29/31	Acute leukaemia	Yes	Not stated	Median (IQR) number of red blood cell units transfused	8 (0 - 34)/6 (0 - 18)
Factor XIII (FXIII): therapeutic trials with inactive comparator						
PCC: prophylactic trials with inactive comparator						

(E) = elective surgery; ICU = intensive care unit, IQR = inter quartile range; SD = standard deviation.

Table 2. Blood loss

Trial	Number included in analyses intervention/comparator	Clinical condition	Transfusion Protocol?	How the outcome was measured	Findings: intervention	Findings: Comparator
Fibrinogen concentrate: prophylactic trials with inactive comparator						
Cui 2010	17/14	Surgery (E) : cardiac	Yes	Mean (SD) quantity (mL/kg/h) of chest drainage at 1 h post-operatively	2.9 (2.0) mL/kg/h	3.5 (1.6) mL/kg/h
				Mean (SD) quantity (mL/kg/h) of chest drainage at 6 h post-operatively	1.3 (0.1) mL/kg/h	1.5 (0.6) mL/kg/h
				Median (IQR) quantity (mL/kg/h) of chest drainage at 24 h post-operatively	0.6 (0.4 to 0.8) mL/kg/h	0.7 (0.6 to 0.9) mL/kg/h
Fenger Erikson 2009	10/10	Surgery (E): radical cystectomy	Not stated	Mean (SD) quantity (mL) blood loss at time of intervention	2682 (962) mL	2933 (1320) mL
Karlsson 2011	10/10	Surgery (E) : cardiac	Yes	Mean (SD): "post-operative bleeding"	565 (150) mL/12 h	830 (268) mL/12 h
Najafi 2014	15/15	Surgery (E): total hip replacement arthroplasty	Yes	Mean (SD) quantity (mL) of intraoperative blood loss	744.6 (337) mL	870 (498) mL
				Mean (SD) quantity (mL) of post-operative bleeding to 24 hours post surgery	231.6 (67) mL	236.6 (81) mL
Pournajafian 2015	21/20	Surgery (E): posterior spinal fusion	Not stated	Mean (SD) mL blood loss up to 24 hours post-operatively	533.3 (157.9) mL	679 (130) mL
Ranucci 2015	58/58	Surgery (E): cardiac	Yes	Median (IQR) quantity (mL) lost from chest drain in first 12 hours	300 (200 to 400) mL/12 h	355 (250 to 600) mL/12 h

Table 2. Blood loss (Continued)

Sadeghi 2014	30/30	Surgery (E) : cardiac	Yes	Mean (SD) quantity (mL) of blood lost in chest drain in first 12 h post-operatively	477 (143) mL	703 (179) mL
Soleimani 2016	31/29	Surgery (E): transurethral resec- tion of the prostate.	Not stated	Mean (SD) quantity (mL) of blood lost intraoperatively	521 (290) mL	557 (411) mL
				Mean (SD) quantity (mL) of blood lost post-operatively	291 (270) mL	341 (314) mL
Fibrinogen concentrate: prophylactic trials with active comparator						
Haas 2015	10/9	Surgery (E): scoliosis	Yes	Median (IQR) "calculated total blood loss (%)" by 24 h post-operatively	36.5 (14.9 to 54.3) %	51.0 (38.5 to 69.2) %
	17/ 13	Surgery (E): craniosyn- ostosis	Yes	Median (IQR) "calculated total blood loss (%)" by 24 h post-operatively	89.7 (77.7 to 112.9) %	156.9 (110..5 to 187.3) %
Fibrinogen concentrate: therapeutic trials with inactive comparator						
Bilecen 2017	58/59	Surgery (E): cardiac	Yes	Median (IQR) mL blood loss from study intervention to chest closure	50 (29 - 100) mL	70 (33 - 145) mL
				Median (IQR) mL blood loss from ICU arrival to 1hr	70 (35 - 130) mL	90 (46 - 149) mL
				Median (IQR) mL blood loss in ICU from 1hr to 3h	80 (50 - 156) mL	110 (40 - 220) mL
				Median (IQR) mL blood loss in ICU from 3h to 6h	100 (54 - 169) mL	110 (60 - 208) mL
				Median (IQR) mL blood loss in ICU from 6h to 12h	110 (80 - 160) mL	125 (83 - 224) mL
				Median (IQR) mL blood loss in ICU from 12h to 24h	130 (80 - 180) mL	160 (90 - 270) mL
				Median (IQR) cumulative mL blood loss in 24 hours	570 (390 - 730) mL	690 (400 - 1090) mL
Collins 2017	28/27	postpartum haemor- rhage	Not stated	Median (IQR) mL blood loss within 24h of study intervention	225 (100 - 341) mL	300 (60 - 800) mL
Jeppsson 2016	24/24	Surgery (E): cardiac	Yes	Mean (SD) mL intraoperative blood loss	306 (110) mL	375 (202) mL
				Median (IQR) mL intraoperative blood loss	300 (200- 400) mL	300 (200-500) mL
				Mean (SD) mL post-operative blood loss (within 12h)	796 (523) mL	897 (553) mL
				Median (IQR) mL post-operative blood loss (within 12h)	650 (500- 835) mL	730 (543-980) mL

Table 2. Blood loss (Continued)

				Mean (SD) mL total blood loss (within 12h)	1103 (518) mL	1272 (588) mL
				Median (IQR) mL total blood loss (within 12h)	913 (815–1230) mL	1185 (930–1398) mL
Rahe-Meyer 2016	78/74	Surgery (E): complex cardiac	Yes	Median (IQR) mL chest tube drainage volume at 6h	260.0 (155.0–410.0) mL	297.5 (200.0–455.0) mL
				Median (IQR) mL chest tube drainage volume at 12h	405.0 (245.0–600.0) mL	447.5 (320.0–700.0) mL
				Median (IQR) mL chest tube drainage volume at 24h	590.0 (400.5–839.5) mL	682.5 (530.0–1050.0) mL
Wikkelso 2015	123/121	postpartum haemor-rhage	Yes	Median (IQR) quantity (mL) lost to 6 weeks postpartum	1700 (1500 to 2000) mL	1700 (1400 to 2000) mL
Fibrinogen concentrate: therapeutic trials with active comparator						
Galas 2014	30/33	Surgery (E): cardiac	Yes	Median (IQR) quantity (mL) chest drainage from intraoperative to 48 h post-operatively	320 (157 - 750) mL	410 (215 - 510) mL
Tanaka 2014	10/10	Surgery (E): valve re-placement or repair	Yes	Median (IQR) quantity (mL) chest tube drainage at 12 h post-operatively	925 (500 to 1693) mL	1315 (653 to 2965) mL
Factor XIII (FXIII): prophylactic trials with inactive comparator						
Godje 2006	25/25	Surgery (E) : cardiac	Yes	Mean (SD) drain volume (ml) at 6 h post-operatively [§]	231 (113) mL	278 (98) mL
				Mean (SD) drain volume (ml) at 12 h post-operatively [§]	371 (170) mL	458 (160) mL
				Mean (SD) drain volume (ml) at 24 h post-operatively [§]	649 (227) mL	747 (222) mL
				Mean (SD) drain volume (ml) at 36 h post-operatively [§]	798 (273) mL	969 (309) mL
				Mean (SD) drain volume (ml) at 48h post-operatively [§]	958 (376) mL	1231 (175) mL
Korte 2009	22 in total	Surgery (E): gastroin-testinal cancer	Yes	Median (IQR) quantity (ml) blood loss at completion of surgery	750 (400 to 1000) mL	1050 (700 to 1800) mL
Levy 2009	8/8	Surgery (E) : cardiac	Not stated	Mean (SD) quantity (mL) of chest drainage at 8 h post-operatively	358 (93.5) mL	505 (301) mL
				Mean (SD) quantity (mL) of chest drainage at 24h post-operatively	903 (738) mL	1155 (804) mL

Table 2. Blood loss (Continued)

	Mean (SD) quantity (mL) of chest drainage at CT removal	1007 (912) mL	1489 (1395) mL
Factor XIII (FXIII): therapeutic trials with inactive comparator			
PCC: prophylactic trials with inactive comparator			

[^] "Calculated total blood loss (%) was based on the estimated total blood volume (calculated as described by [Kearney 1989](#)) " [Haas 2015](#)

^{*} "The visual assessment of the surgical field was performed by the senior surgical staff as follows: 0 = excellent haemostasis (dry field), 1 = mild bleeding (oozing), 2 = moderate bleeding (controllable with applied pressure) and 3 = severe bleeding (multiple diffuse bleeding sites)" [Tanaka 2014](#)

^{\$} Findings are reported graphically, not numerically.

(E) = elective surgery; h = hours; ICU = intensive care unit, IQR = inter quartile range; mL = millilitres; SD = standard deviation.

Table 3. Fibrinogen concentrate: prophylactic trials with inactive comparator

Trial	Num- ber ran- domised (N = inter- vention/N = compara- tor)	Number in- cluded in analyses (N = inter- vention/N = compara- tor)	Intervention	Comparator	Co-inter- ventions	Primary Outcomes	Clinical Con- dition	Population Age
Cui 2010	40 (20/20)	31 (17/14)	Fibrinogen concentrate (0.5-1g) combined with tradi- tional transfusion, guided by TEG	NONE: Tradi- tional transfu- sion guided by clinical experi- ence.	Protamine, FFP, red blood cells, platelets.	Time to chest wall closure.	Surgery (E) : cardiac	Paediatric population
Fenger Erik- son 2009	21 (10/11)	20 (10/10)	Fibrinogen concentrate (45mg/kg), administered in- tra-operatively	PLACEBO: iso- tonic saline, ad- ministered in- tra-operatively at a dose of 2.25 mL/kg).	Intra-opera- tive FFP, red blood cells.	Whole blood maxi- mum clot firmness as determined by thromboelastome- try.	Surgery (E): radical Cys- tectomy	Adult popu- lation
Karlsson 2011	20 (10/10)	20 (10/10)	Preoperative infusion of fib- rinogen concentrate (2g) at 5 minutes following baseline measurements.	NONE	Heparin, protamine, red blood cells, as- pirin.	Clinical adverse events, including graft occlusion on CT up to 3 to 4d post- operatively.	Surgery (E) : cardiac	Adult popu- lation
Najafi 2014	30 (15/15)	30 (15/15)	Fibrinogen concentrate (30 mg/kg) infused after induction of general anaesthesia	PLACEBO: nor- mal saline ad- ministered in equal volume as fibrinogen.	None stat- ed.	Volume of red blood cells transfused dur- ing and 24h post-op- eratively.	Surgery (E): total hip re- placement arthroplasty	Adult popu- lation
Pournajafi- an 2015	41 (21/20)	41 (21/20)	Fibrinogen concentrate (1g) infused after induction of gen- eral anaesthesia.	PLACEBO: nor- mal saline.	None stat- ed.	Not stated.	Surgery (E): posterior spinal fusion.	Adult popu- lation
Ranucci 2015	116 (58/58)	116 (58/58)	Fibrinogen concentrate with dose determined by the mean clot firmness test on the FIBTEM.	PLACEBO: Nor- mal (0.9%) saline.	TXA dur- ing surgery, protamine.	Avoidance of allo- geneic blood prod- ucts transfusion dur-	Surgery (E): cardiac	Adult popu- lation

Table 3. Fibrinogen concentrate: prophylactic trials with inactive comparator (Continued)

						ing hospital stay up to 30d.		
Sabate 2016	99 (51/48)	92 (48/44)	Fibrinogen concentrate with dose calculated to reach a target plasma concentration of 2.9 g/L.	PLACEBO: Normal (0.9%) saline.	Red blood cells, platelets, FFP, tranexamic acid.	The percentage of patients requiring transfusion of red blood cell units during the liver transplant procedure.	Surgery (E): liver transplantation.	Adult population
Sadeghi 2014	60 (30/30)	60 (30/30)	Fibrinogen (1g) administered 30 minutes prior to anaesthesia induction.	PLACEBO: saline administered 30 minutes prior to anaesthesia induction	None stated.	Volume of post-operative haemorrhage 0, 12 and 24h post-operatively.	Surgery (E): cardiac	Adult population
Soleimani 2016	72 (36/36)	60 (31/29)	Fibrinogen concentrate (2g) administered 15-30 minutes before the start of surgery.	PLACEBO: 50ml normal saline.	None stated.	Bleeding volume during and after surgery.	Surgery (E): transurethral resection of the prostate.	Adult population

* For the outcomes bleeding and transfusion requirements, 1 patient from the Factor XIII, 1250 IE arm and 2 patients from the placebo arm were excluded from the analysis due to re-operations following hints of early bypass occlusions. All patients were included in the analysis of the other outcomes.

d = days; (E) = elective surgery; FIBTEM = is a ready-to-use ROTEM® system reagent for use with citrated whole blood and assesses the clot firmness of the fibrin clot, h = hours; PCC = prothrombin complex concentrate; mins = minutes; wks = weeks.

Table 4. Fibrinogen concentrate: prophylactic trials with active comparator

Trial	Number randomised (N = intervention/ N = comparator)	Number included in analyses (N = intervention/ N = comparator)	Intervention	Comparator	Co-interventions	Primary Outcomes	Clinical Condition	Population Age
Haas 2015	Craniosynostosis surgery = 36 (17/14).	Craniosynostosis surgery = 36 (17/13).	Fibrinogen concentrate if FIBTEM MCF under 13mm (early substitution group). Total dose administered per subject in	Fibrinogen concentrate if FIBTEM MCF under 8 mm (conventional). Total dose administered per subject in	Tranexamic acid	Total volume of transfused red blood cells per kg bodyweight within 24h	Surgery (E): craniosynostosis or scoliosis	Paediatric population

Table 4. Fibrinogen concentrate: prophylactic trials with active comparator (Continued)

Scoliosis surgery = 26 (10/9)	Scoliosis surgery = 26 (10/9)	cranosynostosis surgery; median of 90mg kg ⁻¹ (IQR, 75 to 120 mg kg ⁻¹); Scoliosis surgery = median of 60mg kg ⁻¹ (IQR, 30 to 68 mg kg ⁻¹).	cranosynostosis surgery = median of 90mg kg ⁻¹ (IQR, 60 to 90 mg kg ⁻¹); Scoliosis surgery = median of 30mg kg ⁻¹ (IQR, 30 to 60 mg kg ⁻¹).	post-operatively
----------------------------------	----------------------------------	--	---	------------------

CPB = cardiopulmonary bypass; (E) = elective surgery; FFP = fresh frozen plasma; FIBTEM = is a ready-to-use ROTEM® system reagent for use with citrated whole blood and assesses the clot firmness of the fibrin clot; h = hours; INR = International Normalised Ratio; IQR = inter quartile range; PCC = prothrombin complex concentrate; (U) = urgent surgery/treatment.

Table 5. Fibrinogen concentrate: therapeutic trials with inactive comparator

Trial	Number randomised (N = intervention/ N = comparator)	Number included in analyses (N = intervention/ N = comparator)	Intervention	Comparator	Co-interventions	Primary Outcomes	Clinical Condition	Population Age
Bilecen 2017	120 (60/60)	115 (58/57)	Fibrinogen concentrate , dose calculated to provide a target plasma fibrinogen concentration of 2.5 g/L.	PLACEBO : 2g of albumin diluted with 50ml 0.9% sodium chloride solution.	Red blood cells, platelets, FFP, tranexamic acid.	The intraoperative blood loss measured between intervention and closure of the chest.	Surgery (E): cardiac	Adult population
Collins 2017	57 (29/28)	55 (28/27)	Fibrinogen concentrate , dose determined by FIBTEM to elevate A5 result above 22mm.	PLACEBO : 50ml of normal saline.	Red blood cells, platelets, FFP, tranexamic acid.	The number of allogeneic blood products (red blood cell, FFP, cryoprecipitate, platelets) infused after study medication until hospital discharge.	postpartum haemorrhage	Adult population
Curry 2018	48 (24/24)	48 (24/24)	Fibrinogen concentrate 6g infusion.	PLACEBO : 300ml of normal saline.	Major haemorrhage therapy as per hospital protocol: red blood cells, platelets,	Feasibility of delivery of FgC therapy within 45 minutes to adult trauma patients and the proportion of participants whose fibrinogen levels were	Trauma	Adult population

Table 5. Fibrinogen concentrate: therapeutic trials with inactive comparator (Continued)

					FFP, tranexamic acid, cryoprecipitate.	maintained ≥ 2 g/L during active haemorrhage.		
Jeppsson 2016	52 (26/26)	48 (24/24)	Fibrinogen (2g) as an infusion over 5 minutes, immediately before surgery.	PLACEBO: 100ml of normal saline.	300 units kg ⁻¹ of heparin, 1 mg protamine per 100 units of heparin. Red blood cells, platelets, FFP, additional fibrinogen.	Mediastinal drain loss during the first 12 hours post-operatively.	Surgery (E): cardiac	Adult population
Nascimento 2016	50 (25/25)	45 (21/24)	Fibrinogen concentrate , 6g as a 3 minute infusion.	PLACEBO: 300ml of normal saline as a 3 minute infusion.	Red blood cells, tranexamic acid, FFP, platelets and cryoprecipitate.	The feasibility of providing the study intervention within 1 hour of hospital admission.	Trauma	Adult population
Rahe-Meyer 2013	80 (38 /42)	71 (29/32)	Fibrinogen concentrate , dose determined by FIBTEM test: median (IQR range) dose: 8g (6 to 9g).	PLACEBO: normal (0.9%) saline. Median (IQR range) dose: 400ml (300 to 450ml).	TXA during surgery, platelets or FFP.	Total number of units of allogeneic blood components given during first 24h post-operatively	Surgery (E): aortic valve replacement	Adult population
Rahe-Meyer 2016	152 (78/74)	152 (78/74)	Fibrinogen concentrate , dose determined by FIBTEM testing to generate an MCF of 22mm.	PLACEBO: 0.9% sodium chloride solution.	Red blood cells, platelets, FFP, tranexamic acid.	The number of units of allogeneic blood products (FFP, platelets and red blood cells) administered during the 24 hours after administration of study medication.	Surgery (E): complex cardiac	Adult population
Wikkelso 2015	249 (124 /125)	249 (PPA: 123 /121) (ITT: 120 /119)	Fixed dose of 2g of fibrinogen concentrate dispensed using a syringe pump infusion over 20 minutes	PLACEBO: 100ml of isotonic saline (mean (standard deviation) dose =1.2 (0.2) mg/kg).	TXA, hydroxyethyl starch, intravenous fluids.	Red blood cell transfusion during a 6 wk follow-up period postpartum	postpartum haemorrhage	Adult population

d = days; (E) = elective surgery; FFP = fresh frozen plasma; FIBTEM = is a ready-to-use ROTEM® system reagent for use with citrated whole blood and assesses the clot firmness of the fibrin clot; h = hours; INR = International Normalised Ratio, IQR = inter quartile range; mins = minutes; PCC = prothrombin complex concentrate; TXA = tranexamic acid; (U) = urgent surgery/ treatment; wks = weeks.

Table 6. Fibrinogen concentrate: therapeutic trials with active comparator

Trial	Num- ber ran- domised (N = inter- vention/ N = compara- tor)	Number in- cluded in analyses (N = inter- vention/ N = compara- tor)	Intervention	Comparator	Co-inter- ventions	Primary Outcomes	Clinical Condition	Population Age
Galas 2014	63 (30/33)	63 (30/33)	Single dose of Fibrinogen concentrate (60mg/kg body weight, Haemocomplettan®) given at time of intra-operative bleeding.	Single dose of Cryoprecipitate (10ml/kg body weight) given at time of intra-operative bleeding.	Not Stated	post-operative blood losses during 48h after surgery	Surgery (E): cardiac	Paediatric population
Lance 2012	43 (22/21)	43 (22/21) (but only 16/16 for primary outcome analysis).	FFP (2 units) + Fibrinogen (2g)	FFP (4 units)	Protamine, red blood cells	Pre- and post-transfusion ROTEM analysis	Surgery (E): cardiovascular, abdominal or spinal column	Adult population
Tanaka 2014	20 (10/10)	20 (10/10)	A single dose (4g) of Fibrinogen (RiaSTAP®) given within 30 minutes of intervention decision.	One unit of apheresis platelets (median volume 230 ml) within 30 minutes of intervention decision.	Red blood cells, platelets, plasma, cryoprecipitate and fibrinogen if triggers met	i) Haemostatic condition in the surgical field post-intervention; ii) Haemostatic blood product use over the first 24h; iii) Percentage of patients with thromboembolic events at 6-8 weeks post-operatively;	Surgery (E): valve replacement or repair	Adult population

Table 6. Fibrinogen concentrate: therapeutic trials with active comparator (Continued)

Innerhofer 2017	100	94	Fibrinogen concentrate (50 mg/kg of bodyweight) for abnormal ROTEM fibrin polymerisation (FibA10 < 9 mm) or four-Factor PCC (20 IU/kg of bodyweight) for delayed initial thrombin formation (ExCT > 90s or prothrombin time index < 35%). FXIII concentrate (20 IU/kg of bodyweight) was administered with each second fibrinogen dose.	FFP (15 ml/kg bodyweight).	Tranexamic acid 20mg/kg, FXIII concentrate (20 IU/kg of bodyweight),	iv) Mortality at 6-8 weeks post-operatively Occurrence of multiple organ failure during ICU stay as assessed by the daily SOFA score.	Trauma	Adult population
	(52/48)	(50/44)						

* 20 of the 47 patients had a 2nd dose of 20ml of PCC: in 3 because of continued bleeding and in 17 due to INR target not being met. 3 of the 47 patients received a 3rd dose of PCC due to an unmet INR level.

^ 3 of the 46 patients had a 2nd dose of 20 ml PCC: 1 due to a continued bleed and 2 did not reach the target INR.

d = days; (E) = elective surgery; FFP = fresh frozen plasma; h = hours; INR = International Normalised Ratio; mins = minutes; PCC = prothrombin complex concentrate; ROTEM® = rotational thromboelastometry; TXA = tranexamic acid; (U) = urgent surgery/ treatment; wks = weeks.

Table 7. Factor XIII (FXIII): prophylactic trials with inactive comparator

Trial	Number randomised (N = intervention/ N = comparator)	Number included in analyses (N = intervention/ N = comparator)	Intervention	Comparator	Co-interventions	Primary Outcomes	Clinical Condition	Population Age
Godje 2006	75 (25/25/ 25)	75 (25/25/25)*	Infusion of FactorXIII1 1250 units or2) 2500 units*immediately after administration of protamine (protamine was administered during the scheduled coronary artery bypass graft surgery)* This is the group that was included in this review.	NONE: standard surgical and post-operative care	Protamine for heparin reversal FFP, red blood cells, platelets	Not stated	Surgery (E) : cardiac	Adult population
Karkouti 2013	479 (164/162/153)	409 (143/138/128)	A single intravenous dose of FactorXIII 1) 17.5 IU/kg or2) 35 IU/kg*before the induction of anaesthesia.* This is the group that was included in this review.	PLACEBO: single dose of an unspecified	Protamine, antifibrinolytics during cardiopul-	Percentage of patients avoiding any allogeneic	Surgery (E) : cardiac	Adult population

Table 7. Factor XIII (FXIII): prophylactic trials with inactive comparator (Continued)

				placebo before the induction of anaesthesia.	monary bypass, blood products	transfusions within 7d or discharge home.		
Korte 2009	25 (no further details stated)	22 (no further details stated)	FactorXIII (30 U/kg) infused 15 minutes after the beginning of surgery.	PLACEBO: albumin infusion.	Low molecular weight heparin, FFP, red blood cells, platelets	Clot firmness over 195 mins of surgery	Surgery (E): gastrointestinal cancer	Adult population
Levy 2009	43 (10/9/8/8)	43 (10/9/8/8)	A single infusion of FactorXIII at 4 doses:1) 11.9 units/kg;2) 25 units/kg;3) 35 units/kg*;4) 50 units/kg.* This is the group that was included in this review.	PLACEBO: no further details given.	Protamine for heparin reversal FFP, red blood cells, platelets Cryoprecipitate, Fibrinogen	Incidence and severity of adverse events from intervention administration to follow-up at 5 to 7 wks	Surgery (E) : cardiac	Adult population
Rasche 1982	60 (29/31)	60 (29/31)	FactorXIII (1000 units) on day 1 of treatment and a subsequent daily dose of 500 units plus normal care protocols.	NONE: normal care products	Red blood cells, platelets, whole blood	Bleeding complications	Acute leukaemia	Adult population
Shirahata 1990	58 (30/28)	58 (30/28)	FactorXIII (dose of 70 to 100 units) administered slowly for at least 10 minutes intravenously within 6 hours of delivery.	NONE	Not stated	Not stated	Prematurity	Infant population

Table 8. Factor XIII (FXIII): therapeutic trials with inactive comparator

Trial	Number randomised (N = intervention/ N = comparator)	Number included in analyses (N = intervention/ N = comparator)	Intervention	Comparator	Co-interventions	Primary Outcomes	Clinical Condition	Population Age
Bregenzer 1999	28 (17/11)	20 (11/9)	FactorXIII as an i.v. injection (at 3750 units on Day 0 and 1250 units on days 1 through 9) as an	PLACEBO: continuation of ba-	TXA,	Time to cessation of macroscopically visible intestinal bleed-	Ulcerative colitis	Adult population

Table 8. Factor XIII (FXIII): therapeutic trials with inactive comparator *(Continued)*

addition to basic steroid treatment. sic steroid treatment platelets or FFP post-intervention ing within 14d of starting treatment.

Table 9. PCC: prophylactic trials with inactive comparator

Trial	Number randomised (N = intervention/ N = comparator)	Number included in analyses (N = intervention/ N = comparator)	Intervention	Comparator	Co-interventions	Primary Outcomes	Clinical Condition	Population Age
Turner 1981	78 (39/39)	78 (39/39)	PCC (dose of 1ml/kg or 2ml/kg)	NONE: normal care protocols.	Vitamin K, heparin and clotting Factors for disseminated intravascular coagulation, red blood cells	i) Mortality (all-cause) ii) Incidence of intraventricular haemorrhage (post-mortem)	Prematurity	Infant population

APPENDICES

Appendix 1. Search strategies 2018

CENTRAL & DARE (The Cochrane Library)

#1 MeSH descriptor: [Factor XIII] explode all trees

#2 MeSH descriptor: [Fibrinogen] this term only

#3 MeSH descriptor: [Factor IX] explode all trees

#4 MeSH descriptor: [Prothrombin] this term only

#5 (coagulation Factor* or Factor concentrate* or clotting Factor* or recombinant Factor* or plasma-derived concentrate*):ti

#6 ((procoagulant* or pro-coagulant* or prohaemostatic or prohemostatic* or pro-haemostatic* or pro-hemostatic*) near/2 (Factor* or agent*))

#7 fibrinogen concentrate* or Riastap or Riastat or Haemocomplettan* or Clottagen* or Fibroraas

#8 prothrombin complex* or prothrombin concentrate* or Beriplex* or Ocplex* or Octaplex* or Kcentra* or Cofact or Prothrombinex* or "Proplex-T" or Prothroraa* or Haemosolvex* or Prothromplex* or "HT Defix" or Facnyne* or "UMAN Complex" or Kaskadil* or Confidex* or PPSB or Proplex* or ProthoRAAS* or Protromplex* or "Pushu Laishi"

#9 Factor XIII concentrate* or rFXIII or Fibrogammin or Corifact* or recombinant Factor XIII

#10 Factor ix concentrate* or Bebulin* or Profilnine*

#11 #1 or #2 or #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10

#12 MeSH descriptor: [Afibrinogenemia] this term only

#13 MeSH descriptor: [Factor XIII Deficiency] this term only

#14 hypofibrinogen?em* or afibrinogen?em* or dysfibrinogen?em* OR (trauma* or polytrauma*):ti

#15 (Factor xiii or Factor i or Factor ix) near/3 deficien*

#16 MeSH descriptor: [Hemorrhage] explode all trees

#17 (haemorrhag* or hemorrhag* or bleed* or bloodloss* or (blood near/2 los*))

#18 #12 or #13 or #14 or #15 or #16 or #17

#19 #11 and #18

#20 MeSH descriptor: [Factor XIII] explode all trees and with qualifiers: [Administration & dosage - AD, Adverse effects - AE, Therapeutic use - TU]

#21 MeSH descriptor: [Fibrinogen] explode all trees and with qualifiers: [Administration & dosage - AD, Adverse effects - AE, Therapeutic use - TU]

#22 MeSH descriptor: [Factor IX] explode all trees and with qualifiers: [Administration & dosage - AD, Adverse effects - AE, Therapeutic use - TU]

#23 MeSH descriptor: [Prothrombin] this term only and with qualifiers: [Administration & dosage - AD, Adverse effects - AE, Therapeutic use - TU]

#24 #19 or #20 or #21 or #22 or #23

#25 ((haemophilia* or hemophilia*) not (non or without)):ti

#26 #24 not #25

MEDLINE (Ovid)

1. Afibrinogenemia/ or Factor XIII Deficiency/
2. (hypofibrinogen?em* or afibrinogen?em* or dysfibrinogen?em*).tw.
3. ((Factor xiii or Factor I or Factor ix) adj3 deficien*).tw.
4. exp Hemorrhage/
5. (haemorrhag* or hemorrhag* or bleed* or bloodloss* or (blood adj3 los*)).tw.
6. (trauma* or polytrauma*).ti.
7. or/1-6
8. exp Factor XIII/
9. exp Factor IX/
10. *Fibrinogen/
11. Prothrombin/
12. (coagulation Factor* or Factor concentrate* or clotting Factor* or recombinant Factor* or plasma-derived concentrate*).ti.
13. ((procoagulant* or pro-coagulant* or prohaemostatic or prohemostatic* or pro-haemostatic* or pro-hemostatic*) adj2 (Factor* or agent*)).tw.
14. (fibrinogen concentrate* or Riastap or Riastat or Haemocomplettan* or Clottagen* or Fibroraas*).tw.
15. (prothrombin complex* or prothrombin concentrate* or Beriplex* or Ocplex* or Octaplex* or Kcentra* or Cofact or Prothrombinex* or "Proplex-T" or Prothroras* or Haemosolvex* or Prothromplex* or "HT Defix" or Facnyne* or "UMAN Complex" or Kaskadil* or Confidex* or PPSB or Proplex* or ProthoRAAS* or Protromplex* or "Pushu Laishi").tw.
16. (Factor ix concentrate* or Bebulin* or Profilnine*).tw.
17. (Factor XIII concentrate* or rFXIII or Fibrogammin or Corifact* or recombinant Factor XIII).tw.
18. or/8-17
19. 7 and 18
20. exp Factor XIII/ad, ae, th, tu
21. Fibrinogen/ad, ae, th, tu
22. exp Factor IX/ad, ae, th, tu
23. Prothrombin/ad, ae, th, tu
24. or/19-23
25. ((haemophilia* or hemophilia*) not (non or without)).ti.
26. 24 not 25

EMBASE (Ovid)

1. blood clotting Factor deficiency/ or exp blood clotting Factor 13 deficiency/ or exp fibrinogen defect/ or exp prothrombin deficiency/
2. ((Factor xiii or Factor i or Factor ix) adj3 deficien*).tw.
3. (hypofibrinogen?em* or afibrinogen?em* or dysfibrinogen?em*).ti,ab.
4. exp Bleeding/
5. (haemorrhag* or hemorrhag* or bleed* or bloodloss* or (blood adj3 los*)).tw.
6. Multiple Trauma/

7. (trauma* or polytrauma*).ti.
8. or/1-7
9. Blood Clotting Factor 13 concentrate/ or blood clotting Factor 13a/ or blood clotting Factor 13b/
10. blood clotting Factor 9 complex/ or blood clotting Factor 9 concentrate/ or blood clotting Factor 9a/ or recombinant blood clotting Factor 9/
11. *Fibrinogen/
12. Prothrombin Complex/
13. (coagulation Factor* or Factor concentrate* or clotting Factor* or recombinant Factor* or plasma-derived concentrate*).ti.
14. ((procoagulant* or pro-coagulant* or prohaemostatic or prohemostatic* or pro-haemostatic* or pro-hemostatic*) adj (Factor* or agent*)).tw.
15. (fibrinogen concentrate* or Riastap or Riastat or Haemocomplettan* or Clottagen* or Fibroraas*).tw.
16. (prothrombin complex* or prothrombin concentrate* or Beriplex* or Ocplex* or Octaplex* or Kcentra* or Cofact or Prothrombinex* or "Proplex-T" or Prothroras* or Haemosolvex* or Prothromplex* or "HT Defix" or Facnyne* or "UMAN Complex" or Kaskadil* or Confidex* or PPSB or Proplex* or ProthoRAAS* or Protromplex* or "Pushu Laishi").tw.
17. (Factor ix concentrate* or rFIX or Bebulin* or Profilnine* or recombinant Factor ix).tw.
18. (Factor XIII concentrate* or rFXIII or Corifact* or Fibrogammin* or recombinant Factor XIII).tw.
19. or/9-18
20. 8 and 19
21. fibrinogen concentrate/
22. blood clotting Factor 13 concentrate/ae, ct, ad, cb, cm, cr, dv, do, it, dt, to, im, iv, pr, pe, pk, pd
23. blood clotting Factor 9 concentrate/ae, ct, ad, cb, cm, cr, dv, do, it, dt, to, im, iv, pr, pe, pk, pd
24. fibrinogen/ae, ct, ad, cb, cm, cr, dv, do, it, dt, to, ia, ce, dl, im, io, ip, pl, iv, vi, pa, pr, pe, pk, pd, rp, sc, th, tp
25. prothrombin complex/ae, ct, ad, cb, cm, cr, dv, do, it, dt, to, iv, po, pa, pr, pe, pk, pd
26. or/20-25
27. ((haemophilia* or hemophilia*) not (non or without)).ti.
28. 26 not 27

CINAHL (EBSCOHOST)

- S1 (MH "Afibrinogenemia")
- S2 hypofibrinogen?em* or afibrinogen?em* or dysfibrinogen?em*
- S3 TX (Factor xiii or Factor i or Factor ix) N3 deficient*
- S4 (MH "Hemorrhage+")
- S5 TI (haemorrhag* or hemorrhag* or bleed* or bloodloss* or (blood N3 los*)) OR AB (haemorrhag* or hemorrhag* or bleed* or bloodloss* or (blood N3 los*))
- S6 S1 OR S2 OR S3 OR S4 OR S5
- S7 (MH "Fibrinogen")
- S8 (MH "Thromboplastin")
- S9 TI coagulation Factor* or Factor concentrate* or clotting Factor* or recombinant Factor* or plasma-derived concentrate*

S10 TI (((procoagulant* or pro-coagulant* or prohaemostatic or prohemostatic* or pro-haemostatic* or pro-hemostatic*) N2 (Factor* or agent*))) OR AB (((procoagulant* or pro-coagulant* or prohaemostatic or prohemostatic* or pro-haemostatic* or pro-hemostatic*) N2 (Factor* or agent*)))

S11 TX Factor XIII concentrate* or rFXIII or Fibrogammin or Corifact* or recombinant Factor XIII

S12 TX Factor ix concentrate* or Bebulin* or Profilnine*

S13 TX prothrombin complex* or prothrombin concentrate* or Beriplex* or Ocplex* or Octaplex* or Kcentra* or Cofact or Prothrombinex* or "Proplex-T" or Prothroras* or Haemosolvex* or Prothromplex* or "HT Defix" or Facnyne* or "UMAN Complex" or Kaskadil* or Confidex* or PPSB or Proplex* or ProthoRAAS* or Protromplex* or "Pushu Laishi"

S14 TX fibrinogen concentrate* or Riastap or Riastat or Haemocomplettan* or Clottagen* or Fibroraas*

S15 S7 OR S8 OR S9 OR S10 OR S11 OR S12 OR S13 OR S14

S16 S6 AND S15

S17 (MH "Fibrinogen/AD/AE/TU")

S18 S16 OR S17

S19 TI ((haemophilia* or hemophilia*) not (non or without))

S20 S18 NOT S19

PUBMED

1. haemorrhag*[TIAB] OR hemorrhag*[TIAB] OR bleed*[TIAB] OR bloodloss*[TIAB] OR blood los*[TIAB] OR "loss of blood" [TIAB]

2. hypofibrinogenaem*[TIAB] OR hypofibrinogenem*[TIAB] OR afibrinogenem*[TIAB] OR afibrinogenaem*[TIAB] OR dysfibrinogenem*[TIAB] OR dysfibrinogenaem*[TIAB]

3. Factor xiii deficien*[TI] OR Factor i deficien*[TI] OR Factor ix deficien*[TI] OR trauma*[TI] OR polytrauma*[TI]

4. #1 OR #2 OR #3

5. coagulation Factor*[TI] OR Factor concentrate*[TI] OR clotting Factor*[TI] OR recombinant Factor*[TI] OR plasma-derived concentrate*[TI]

6. procoagulant*[TIAB] OR pro-coagulant*[TIAB] OR prohaemostatic[TIAB] OR prohemostatic*[TIAB] OR pro-haemostatic*[TIAB] OR pro-hemostatic*[TIAB]

7. fibrinogen concentrate*[TIAB] OR Haemocomplettan*[TIAB] OR Clottagen*[TIAB] OR Fibroraas*[TIAB] OR Riastap*[TIAB] OR Riastat*[TIAB]

8. prothrombin complex*[TIAB] OR prothrombin concentrate*[TIAB] OR Beriplex*[TIAB] OR Ocplex*[TIAB] OR Octaplex*[TIAB] OR Kcentra*[TIAB] OR Cofact*[TIAB] OR Prothrombinex*[TIAB] OR Proplex-T*[TIAB] OR Prothroras*[TIAB] OR Haemosolvex*[TIAB] OR Prothromplex*[TIAB] OR "HT Defix"[TIAB] OR Facnyne*[TIAB] OR PPSB*[TIAB] OR UMAN Complex*[TIAB] OR Kaskadil*[TIAB] OR Confidex*[TIAB] OR Proplex*[TIAB] OR ProthoRAAS*[TIAB] OR Protromplex*[TIAB] OR "Pushu Laishi"[TIAB]

9. Factor IX concentrate*[TIAB] OR rFIX[TIAB] OR Bebulin*[TIAB] OR Profilnine*[TIAB] OR recombinant Factor IX[TIAB]

10. Factor XIII concentrate*[TIAB] OR rFXIII[TIAB] OR Corifact*[TIAB] OR Fibrogammin[TIAB] OR recombinant Factor XIII[TIAB]

11. #5 OR #6 OR #7 OR #8 OR #9 OR #10

12. #4 AND #11

13. ((haemophili*[TI] OR hemophili*[TI]) NOT (non[TI] OR without[TI]))

14. #12 NOT #13

15. publisher[SB] NOT pubstatusnihms

16. #14 AND #15

17. (random* OR blind* OR "control group" OR placebo* OR controlled OR groups OR trial* OR "systematic review" OR "meta-analysis" OR metaanalysis OR "literature search" OR medline OR cochrane OR embase) AND (publisher[sb] OR inprocess[sb] OR pubmednotmedline[sb])

18. #16 AND #17

TRANFUSION EVIDENCE LIBRARY

(haemorrhage OR hemorrhage OR bleed OR bleeding OR bloodloss OR hypofibrinogenaemia OR hypofibrinogenemia OR afibrinogenemia OR afibrinogenaemia OR dysfibrinogenemia OR dysfibrinogenaemia OR Factor xiii deficiency OR Factor i deficiency OR Factor ix deficiency OR trauma OR polytrauma OR traumatic) AND (coagulation Factor OR Factor concentrate OR clotting Factor OR recombinant Factor OR plasma-derived concentrate OR procoagulant OR pro-coagulant OR prohaemostatic OR prohemostatic OR pro-haemostatic OR pro-hemostatic OR coagulation Factors OR Factor concentrates OR clotting Factors OR recombinant Factors OR plasma-derived concentrates OR procoagulants OR pro-coagulants OR prohaemostatics OR prohemostatics OR pro-haemostatics OR pro-hemostatics OR fibrinogen concentrate OR fibrinogen concentrates OR Haemocomplettan OR Clottagen OR Fibroraas OR Riastap OR Riastat OR prothrombin complex OR prothrombin concentrate OR prothrombin concentrates OR Beriplex OR Ocplex OR Octaplex OR Kcentra OR Cofact OR Prothrombinex OR Proplex OR Prothraas OR Haemosolvex OR Prothromplex OR HT Defix OR Facnyne OR PPSB OR UMAN Complex OR Kaskadil OR Confidex OR Prothraas OR Protromplex OR Pushu Laishi OR Factor XIII concentrate OR rFXIII OR Fibrogammin OR recombinant Factor IX) OR Factor ix concentrate OR rFIX OR Bebulin OR Profilnine OR recombinant Factor IX)

LILACS

(tw:(("coagulation Factor" OR "coagulation Factors" OR "Factor concentrate" OR "Factor concentrates" OR "clotting Factor" OR "clotting Factors" OR "recombinant Factor" OR "recombinant Factors" OR "plasma-derived concentrate" OR "plasma-derived concentrates" OR procoagulant OR procoagulants OR pro-coagulant OR pro-coagulants OR prohaemostatic OR prohaemostatics OR prohemostatic OR prohemostatics OR pro-haemostatic OR pro-haemostatics OR pro-hemostatic OR pro-hemostatics OR "fibrinogen concentrate" OR "fibrinogen concentrates" OR haemocomplettan OR clottagen OR fibroraas OR riastat OR riastap OR "prothrombin complex" OR "prothrombin concentrate" OR "prothrombin concentrates" OR beriplex OR ocplex OR octaplex OR kcentra OR cofact OR prothrombinex OR proplex OR prothraas OR haemosolvex OR prothromplex OR facnyne OR PPSB OR UMAN OR kaskadil OR confidex OR prothraas OR protromplex OR "Pushu Laishi" OR "Factor XIII concentrate" OR rfxiii OR fibrogammin OR "recombinant Factor XIII" OR profilnine OR "Factor ix concentrate" OR rfix OR bebulin OR "recombinant Factor IX")) AND db:("LILACS") AND type_of_study:("clinical_trials"))

INDMED

1. (haemorrhage OR hemorrhage OR bleed OR bleeding OR bloodloss OR blood loss OR hypofibrinogenaemia OR hypofibrinogenemia OR afibrinogenemia OR afibrinogenaemia OR dysfibrinogenemia OR dysfibrinogenaemia OR trauma OR traumatic)

2. (Factor OR Factors OR plasma-derived concentrate OR procoagulant OR pro-coagulant OR prohaemostatic OR prohemostatic OR pro-haemostatic OR pro-hemostatic OR fibrinogen concentrate OR Haemocomplettan OR Clottagen OR Fibroraas OR Riastat OR Riastap OR prothrombin complex OR prothrombin concentrate OR Beriplex OR Ocplex OR Octaplex OR Kcentra OR Cofact OR Prothrombinex OR Proplex OR Prothraas OR Haemosolvex OR Prothromplex OR HT Defix OR Facnyne OR UMAN Complex OR Kaskadil OR Confidex OR PPSB OR Proplex OR Prothraas OR Protromplex OR Pushu Laishi OR Bebulin OR Profilnine OR Factor XIII concentrate OR rFXIII OR Fibrogammin OR Corifact)

3. (randomized OR randomised OR randomly OR blind OR blinded OR trial OR control group OR controlled OR groups)

4. 1 AND 2 AND 3

KOREAMED

"Randomized Controlled Trial" [PT] AND ("Hemorrhage" [MH] OR "Factor XIII Deficiency" [MH] OR "Afibrinogenemia" [MH] OR hemorrhag* [ALL] OR bleed* [ALL] OR "Prothrombin" [MH] OR prothrombin [ALL] OR concentrate*[TI] OR "Fibrinogen" [MH] OR fibrinogen [ALL] OR "Factor XIII" [MH] OR trauma* [ALL] OR Factor* [ALL])

OR

"Randomized Controlled Trial" [PT] AND (procoagulant* [ALL] OR prohaemostatic* [ALL] OR prohemostatic* [ALL] OR Haemocomplettan [ALL] OR Clottagen [ALL] OR Fibroraas [ALL] OR Beriplex [ALL] OR Octaplex [ALL] OR Cofact [ALL] OR Prothrombinex [ALL] OR Proplex [ALL] OR Prothraas [ALL] OR Haemosolvex [ALL] OR Profilnine [ALL] OR Prothromplex [ALL] OR Bebulin [ALL] OR Facnyne [ALL] OR Kaskadil [ALL] OR rFXIII [ALL] OR Fibrogammin [ALL] OR Riastap [ALL] OR Riastat [ALL] OR Ocplex [ALL] OR Corifact [ALL] OR Confidex [ALL])

WEB OF SCIENCE – CPCI-S

#1 TI=(haemorrhag* OR hemorrhag* OR bleed* OR bloodloss OR blood loss OR hypofibrinogenaemi* OR hypofibrinogenemi* OR afibrinogenemi* OR afibrinogenaemi* OR dysfibrinogenemi* OR dysfibrinogenaemi* OR Factor xiii deficien* OR Factor I deficien* OR Factor ix deficien* OR trauma* OR polytrauma*)

#2 TI=(concentrate* OR procoagulant* OR pro-coagulant* OR prohaemostatic OR prohemostatic* OR pro-haemostatic* OR pro-hemostatic* OR "coagulation Factor" OR "clotting Factor" OR "recombinant Factor" OR Haemocomplettan* OR Clottagen OR Fibroraas OR Riastat OR Riastap OR "prothrombin complex" OR Beriplex OR Ocplex OR Octaplex OR Kcentra OR Cofact OR Prothrombinex OR Proplex OR Prothraas OR Haemosolvex OR Prothromplex OR Defix OR Facnyne OR UMAN OR Kaskadil OR Confidex OR PPSB OR Proplex OR ProthoRAAS OR Protromplex OR Pushu Laishi OR Bebulin OR Profilnine OR rFXIII OR Fibrogammin OR Corifact)

#3 TS=(randomized OR randomised OR trial)

#4 #1 AND #2 AND #3

ONGOING TRIALS

ClinicalTrials.gov

Study Type: Interventional Studies

Condition: hemorrhage OR trauma

Interventions/Title: coagulation Factor OR recombinant Factor OR procoagulant OR prohemostatic OR fibrinogen concentrate OR prothrombin complex OR Factor IX concentrate OR Factor XIII concentrate OR Haemocomplettan* OR Clottagen OR Fibroraas OR Riastat OR Riastap OR Beriplex OR Ocplex OR Octaplex OR Kcentra OR Cofact OR Prothrombinex OR Proplex OR Prothraas OR Haemosolvex OR Prothromplex OR Defix OR Facnyne OR UMAN OR Kaskadil OR Confidex OR PPSB OR ProthoRAAS OR Protromplex OR Pushu Laishi OR Bebulin OR Profilnine OR rFIX OR rFXIII OR Fibrogammin OR Corifact

WHO ICTRP

Title/Intervention: procoagulant OR prohemostatic OR prohaemostatic OR Factor concentrate OR prothrombin complex OR prothrombin concentrate OR fibrinogen concentrate OR coagulation Factor OR clotting Factor OR plasma-derived concentrate OR Haemocomplettan OR Clottagen OR Fibroraas OR Riastat OR Riastap OR Beriplex OR Ocplex OR Octaplex OR Kcentra OR Cofact OR Prothrombinex OR Proplex OR Prothraas OR Haemosolvex OR Prothromplex OR HT Defix OR Facnyne OR UMAN Complex OR Kaskadil OR Confidex OR PPSB OR Proplex OR ProthoRAAS OR Protromplex OR Pushu Laishi OR rFIX OR Factor IX concentrate OR Bebulin OR Profilnine OR Factor XIII concentrate OR rFXIII OR Fibrogammin OR Corifact

Condition: hemorrhage OR haemorrhage OR haemorrhaging OR hemorrhaging OR bleed OR bleeding OR trauma OR deficiency

Recruitment Status: ALL

EUDRACT

((hemorrhage OR trauma) AND (coagulation Factor OR Factor concentrate OR clotting Factor OR recombinant Factor OR plasma-derived concentrate OR procoagulant OR pro-coagulant OR prohaemostatic OR prohemostatic OR pro-haemostatic OR pro-hemostatic OR fibrinogen OR Haemocomplettan OR Clottagen OR Fibroraas OR Riastat OR Riastap OR prothrombin complex OR prothrombin concentrate OR Beriplex OR Ocplex OR Octaplex OR Kcentra OR Cofact OR Prothrombinex OR Proplex OR Prothraas OR Haemosolvex OR Prothromplex OR HT Defix OR Facnyne OR UMAN Complex OR Kaskadil OR Confidex OR PPSB OR Proplex OR ProthoRAAS OR Protromplex OR Pushu Laishi OR rFIX OR Factor IX concentrate OR Bebulin OR Profilnine OR Factor XIII concentrate OR rFXIII OR Fibrogammin OR Corifact))

ISRCTN

((hemorrhage OR haemorrhage OR haemorrhaging OR hemorrhaging OR bleed OR bleeding OR trauma) AND (coagulation Factor OR Factor concentrate OR clotting Factor OR recombinant Factor OR plasma-derived concentrate OR procoagulant))- 7

((hemorrhage OR haemorrhage OR haemorrhaging OR hemorrhaging OR bleed OR bleeding OR trauma) AND (procoagulant OR pro-coagulant OR prohaemostatic OR prohemostatic OR pro-haemostatic OR pro-hemostatic OR fibrinogen concentrate OR Riastap OR Riastat OR Haemocomplettan OR Clottagen OR Fibroraas))- 8

((hemorrhage OR haemorrhage OR haemorrhaging OR hemorrhaging OR bleed OR bleeding OR trauma) AND (prothrombin complex OR prothrombin concentrate OR Beriplex OR Ocplex OR Octaplex OR Kcentra OR Cofact OR Prothrombinex OR Proplex OR Prothraas OR Haemosolvex OR Prothromplex OR Protromplex))- 5

((hemorrhage OR haemorrhage OR haemorrhaging OR hemorrhaging OR bleed OR bleeding OR trauma) AND (rFIX OR Factor IX concentrate OR Bebulin OR Profilnine OR Facnyne OR UMAN OR Kaskadil OR Confidex OR PPSB OR Fibrogammin OR Corifact OR Factor XIII concentrate OR rFXIII))- 0

CONTRIBUTIONS OF AUTHORS

Jez Fabes did most of the data extraction, 'Risk of bias' assessments and data analysis for this review, and led on the writing of the final review text.

Susan Brunskill is a methodological advisor for this review and has contributed to the data extraction, 'Risk of bias' assessments, and data analysis, and wrote some of the review text and provided project support throughout the review process.

Nikki Curry is a content advisor for this review and has undertaken screening, data extraction and 'Risk of bias' assessments for many of the included studies.

Carolyn Doree is an information specialist and has written and run all the search strategies for this review, and did an early screening of the search results.

Simon Stanworth is a content advisor for this review and has contributed to the review in this capacity.

DECLARATIONS OF INTEREST

Nicola Curry and Simon Stanworth were the chief investigators for the E-FIT 1 trial which has been included in this review as [Curry 2018](#).

Jez Fabes: None known.

Susan Brunskill: None known.

Carolyn Doree: None known.

SOURCES OF SUPPORT

Internal sources

- NHS Blood & Transplant, Research and Development, UK.

External sources

- No sources of support supplied

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

The differences between the protocol and the review are:

- **Eligible interventions:** In the protocol we stated that we would include RCTs in which pro-coagulant haemostatic agents were used as indications for warfarin (or a similar agent) over-anticoagulation reversal. However, when extracting data from six identified studies in which a pro-coagulant haemostatic was used exclusively to reverse warfarin (or a similar agent) ([Boulis 1999](#); [Demeyere 2010](#); [Goldstein 2015](#); [Papakitsos 2014](#); [Sarode 2013](#); [van Aart 2006](#)), it became clear that clinically the participants in these studies were distinct from participants in the other included studies. In participants already receiving warfarin (or a similar agent) the baseline risk of arterial or venous thromboembolic events is higher than in non-anticoagulated participants, due to the condition for which they have been receiving anti-coagulation therapy (i.e. warfarin or a similar agent). The use of pro-coagulant haemostatic factors in warfarin reversal has been explored in at least two published systematic reviews ([Dentali 2011](#); [Volis 2012](#)).
- **Subgrouping of the data:** In the protocol we said we would differentiate between trials that gave the intervention prophylactically and those which gave the intervention therapeutically (i.e. to treat active bleeding). We also said that we would differentiate between studies comparing a haemostatically active intervention with a haemostatically inactive intervention and studies comparing two haemostatically active interventions. However during data analysis and reporting it became clear that before differentiating between prophylactic and therapeutic interventions and haemostatically active or inactive comparators, we needed to group the studies by type of intervention: fibrinogen concentrate, factor XIII and prothrombin complex concentrate (PCC). We made this decision to provide the most clinically relevant results and comparisons. We determined that comparisons that contained mixed interventions (i.e. fibrinogen concentrate and factor XIII) would provide limited clinical utility.
- **Wording of the primary outcome:** We updated our primary outcome of incidence of arterial and/or venous thromboembolic events to "the number of participants suffering arterial or venous thromboembolic events, or both" to highlight the individual patient risk of suffering a thromboembolic adverse event rather than the study population event incidence rate. We felt this provided more clinical value when comparing intervention risks, and also permitted a patient-centred discussion were this data to be used in future risk discussions with patients; describing the risk to the individual patient is more intuitive and appropriate.

INDEX TERMS**Medical Subject Headings (MeSH)**

Blood Coagulation Factors [*therapeutic use]; Cause of Death; Factor XIII [*therapeutic use]; Fibrinogen [*therapeutic use]; Hemorrhage [mortality] [prevention & control] [*therapy]; Hemostatics [*therapeutic use]; Randomized Controlled Trials as Topic

MeSH check words

Humans