Published in final edited form as:

J Proteome Res. 2019 February 01; 18(2): 753–758. doi:10.1021/acs.jproteome.8b00703.

Tools for 3D Interactome Visualization

Andrew Keller¹, Juan D. Chavez¹, Jimmy K. Eng¹, Zorian Thornton¹, and James E. Bruce^{1,*}

¹Department of Genome Sciences, University of Washington, Seattle, WA 98195 USA

Abstract

In cells, intra- and inter-molecular interactions of proteins confer function and the dynamic modulation of this interactome is critical to meet changing needs required to support life. Cross-linking and mass spectrometry (XL-MS) enable detection of both intra- and inter-molecular protein interactions in organelles, cells, tissues and organs. Quantitative XL-MS enables detection of interactome changes in cells due to environmental, phenotypic, pharmacological, or genetic perturbations. We have developed new informatics capabilities, the first to enable three-dimensional visualization of multiple quantitative interactome datasets, acquired over time or with varied perturbation levels, to reveal relevant dynamic interactome changes. These new tools are integrated within release 3.0 of our online cross-linked peptide database and analysis tool suite XLinkDB. With the recent rapid expansion in XL-MS for protein structural studies and the extension to quantitative XL-MS measurements, 3D interactome visualization tools are of critical need.

Graphical Abstract

*Corresponding Author: jimbruce@uw.edu. Tel: 206-543-0220.

SUPPORTING INFORMATION

The following supporting information is available free of charge at ACS website <http://pubs.acs.org>

Keywords

Crosslinking; mass spectrometry; protein interaction networks; protein complexes; quantitative proteomics; dynamic interactome; crosslink database; heatmap; interactome movies; interaction surfaces

---

INTRODUCTION

The use of quantitative crosslinking to compare protein interactions among samples is rapidly gaining popularity and importance^{1–3}. Cleavable chemical crosslinkers, in particular, allow for the identification by mass spectrometry of crosslinks *in situ* in cells, tissues, and whole organisms^{4–7}. By providing distance constraints on protein residues, crosslinks help elucidate the structures of proteins and protein complexes^{8–10}. Quantitative crosslink data can be derived from samples isotopically labeled light or heavy, either at the level of the crosslinked peptides using technologies such as SILAC^{3, 11}, or at the level of the crosslinker¹². In either case, the light and heavy samples are combined together prior to XL-MS. Identified light and heavy crosslinks are then quantified according to the peak areas of their parent ions, as acquired in MS¹, to yield a ratio of abundance in the light sample (experimental condition) versus the heavy sample (reference condition). Quantitative cross-linked peptide information can be combined with protein abundance level measurements to delineate conformational and interaction changes due to posttranslational modifications, or protein interactor- or co-factor-induced allosteric changes, rather than changes in protein abundance.

A unique opportunity for new insight on biological function is presented by quantitative interactome measurements during perturbation. For example, visualization of interactome changes in samples treated with increasing concentrations of drugs, or samples crosslinked longitudinally during an environmental perturbation can reveal functional conformational and protein interaction changes not evident in any other large-scale data. Combining the XL-MS data of multiple samples enables one to pool their crosslink identifications which can then be quantified separately in each sample. Analyzing the data in this manner facilitates the researcher's ability to compare quantitative behavior among samples to find trends of interest. This is especially true in cases where samples have different experimental conditions, but a common reference condition. Comparison of crosslink levels among the samples is then possible by comparing their abundance ratios with respect to that of the common reference condition. This gives the opportunity to visualize the interactions of related samples as a third dimension, in a 3D interactome, to help understand how the interactions change over time or with respect to varied perturbation levels.

We have previously reported XLinkDB^{13–15}, a database that offers unique informatic tools for storing and visualizing protein interaction topology data. More recently, we have developed additional quantitative interactome analysis informatics tools to support visualization of 3 dimensional interactome datasets. These tools are integrated into XLinkDB 3.0 which stores many crosslink characteristics, including the distance in the context of protein structures, and the distance in known protein-protein interaction (PPI) databases. Support for quantitation now enables storage for one or more related samples, of

a crosslink log2ratio, standard error (among contributing replicates), and computed p-value reflecting the likelihood that an observed log ratio significantly deviates from zero. Additionally, protein abundance level measurements of crosslinked proteins can also be integrated within XLinkDB 3.0. One can view and explore crosslink data in a browser, as either a table with heatmap, or as a PPI network in Cytoscape¹⁶.

MATERIALS AND METHODS

Crosslink Data Upload to XLinkDB

Quantitative crosslink data in tab delimited file format can be uploaded to XLinkDB at <http://xlinkdb.gs.washington.edu/xlinkdb/>. The file must contain a header line with the following column names:

1. PeptideA
2. ProteinA (with optional,nsp1,wt1,ProteinA2,nsp2,wt2,.....)
3. CrosslinkPositionA
4. PeptideB
5. ProteinB (with optional,nsp1,wt1,ProteinB2,nsp2,wt2,.....)
6. CrosslinkPositionB

Optional additional crosslink columns include:

7. Confidence
8. NumberIDs

Additional columns for quantitation can be added with names, for each quantified sample, as:

9. Exp_Ref_Type_log2ratio
10. Exp_Ref_Type_log2stdev
11. Exp_Ref_Type_pvalue
12. Exp_Ref_Type_chromatogram

where 'Exp', 'Ref', 'Type' are the experiment condition, reference condition, and quantitation type ('silac', 'd0/d8'), respectively. They must in combination be unique for each included sample ratio. The Exp_Ref_Type_log2stdev, Exp_Ref_Type_pvalue, and Exp_Ref_Type_chromatogram columns are optional.

Values stored in columns of each non-header entry of the file are:

1. First peptide sequence, without modifications
2. Protein1,nsp1,wt1,Protein2,nsp2,wt2..... where Protein1, Protein2,... are the ranked list of protein Uniprot IDs corresponding to the first peptide, with their numbers of sibling peptides (nsp) and weight (wt) apportioned by

ProteinProphet¹⁷, when available. Only a single Protein is required. Other values are optional.

3. Position in sequence of first peptide, starting with 0 for the N-terminal residue
4. Second peptide sequence, without modifications
5. Protein1,nsp1,wt1,Protein2,nsp2,wt2..... where Protein1, Protein2,... are the ranked list of protein Uniprot IDs corresponding to the second peptide, with their numbers of sibling peptides (nsp) and weight (wt) apportioned by ProteinProphet, when available. Only a single Protein is required. Other values are optional.
6. Position in sequence of second peptide, starting with 0 for the N-terminal residue

Additional optional columns:

7. Probability that the non-redundant crosslink is correctly identified in the data set, a value between 0 and 1
8. Number of identifications of the non-redundant crosslink in the dataset

Additional columns for quantitation:

9. Mean log2 ratio of experiment to reference condition, among contributing replicates
10. Standard deviation of log2ratios among contributing replicates
11. P-value assessing likelihood of observing mean log2 ratio by chance
12. Chromatogram, enabling one to view the light and heavy parent raw data traces upon clicking a log2ratio value in the table or Cytoscape colored network edge. They must be extracted out of the raw data for the experiment and reference parent peaks ranging over the time of quantitation. As many as four chromatograms can be imported, separated by commas and written as an extended single line. The format is given below, and requires square bracket delimiters as illustrated:

```
[[silac,ms1_exp,m/z,Rawfilename,NormalizationFactor],
[2580.8,0],[2581.4,0],[2582.0,0],[2588.8,281176],[2589.9,195922],.....,
[2536.7,0]],
[[silac,ms1_ref,m/z,Rawfilename, NormalizationFactor],
[2580.8,0],[2581.4,0],[2582.0,0],[2588.8,190996],[2589.9,236953],.....,
[2609.6,0]]
```

where ms1_exp and ms1_ref indicate the extracted ms1 data for the experimental and reference conditions, respectively. One must specify m/z values of the light and heavy parent peaks, Rawfilename, the name of the raw data file, and NormalizationFactor, the amount added to each log2ratio to center the distribution (0 when no normalization was applied). This is useful to display along with the chromatogram so researchers know how it resulted in the recorded

log2ratio. ‘d0/d8’ can be substituted for ‘silac’ when appropriate. The following brackets each contain pairs of retention time in seconds, and the integrated peak intensity within ppm tolerance.

Note that the first, third, fourth, and sixth columns must be unique in the uploaded file, since they together define a unique crosslink. For the protein columns, when ProteinProphet results are known, one can include its apportionment of the number of sibling peptides and weight to the proteins, as indicated above, following the protein Uniprot ID.

Quantitative Proteome Data Upload to XLinkDB

Quantitative proteome data associated with a network on XLinkDB can be uploaded in tab delimited file format at <http://xlinkdb.gs.washington.edu/xlinkdb/>. The file must contain a header line with the following column names:

1. Protein
2. Exp_Ref_Type_log2ratio
3. Exp_Ref_Type_log2stdev
4. Exp_Ref_Type_pvalue

Values are as described above for crosslink data, but with only a single protein in the first column, matching names of ProteinA or ProteinB in the crosslink data upload. The Type is ‘shotgun’. Only the first 2 columns are required. Note that the first column value must be unique in the uploaded file.

Interaction Surfaces between Two Proteins

The interaction surfaces are computed by finding all amino acids in the structure or modeled docking PDB files with the two interacting proteins, that are within a user specified Ca-Ca Euclidean distance from any residue of the other protein. The surface residues of each protein are passed to the NGL Viewer with coloring instructions, along with crosslinks.

Rotating View Html Page

PNG files were downloaded from the XLinkDB Cytoscape PPI network colored, in succession, by multiple samples treated with increasing concentrations of 17-AAG, and assembled into an HTML page using the JQuery code available at <https://github.com/mcnitt/jquery-infinite-rotator>. The HTML page was loaded into a browser and saved as a screenshot movie file with QuickTime.

RESULTS AND DISCUSSION

Table View

Crosslinks uploaded to XLinkDB can be viewed in table format with one row per non-redundant crosslink. Information displayed for each crosslink, in columns, includes the released crosslinked peptide sequences and crosslink sites, proteins, distance in context of available protein structures and models, numbers of corresponding protein pairs, and numbers of identifications. With this information available for each identified crosslink,

researchers can efficiently explore the data. One can sort by any column values, or filter the table to view only crosslinks with specific features, such as corresponding peptide or proteins, or numbers of identifications. Users have the option to display any or all sample quantitation information associated with the dataset, including the log₂ratio, standard error, and p-value. Figure 1 shows selected rows of the table on XLinkDB displaying log₂ratio values in several samples as a heatmap. By sorting columns, one can identify crosslinks exhibiting similar abundance trends in the samples.

Charts summarizing some features of the displayed crosslink data, including cross-linked distance in the context of protein structures and models, as well as quantitation log₂ratios, are included at the bottom of the table. The distances are useful indicators of crosslinks that are not consistent with known protein structures, but possibly with alternative conformations. In cases where more than a single ratio is displayed in the table, plots of the multiple log₂ratios for each crosslink are plotted as trend lines (Figure 2A), which when clicked, go to a table with the single entry of that crosslink. When only a single ratio is displayed in the table, the number of crosslinks having log₂ratio values within various bounds are displayed as a clickable bar graph (Figure 2B), with links going to a table containing the contributing crosslinks with ratios within its bounds. These features help one to explore the dataset to identify and assess crosslinks with quantitation values of interest.

A primary challenge involved in analysis and interpretation of large-scale interactome data arises from the difficulties encountered in handling raw mass spectrometry and chromatographic data, protein structures and models, as well as offering seamless integration with other existing PPI database information. Within XLinkDB 3.0, users can make raw data available during the upload and if so, one can click on a log₂ratio value in the table to immediately view the chromatographic profiles of contributing light and heavy parent peaks. This provides crucial assessment of the quality of the quantitative measurement (Figure 3).

Hyperlinks were also added in XLinkDB 3.0 to several other column values in the table, each leading to pertinent additional information. For example, clicking on the entry in the table column entitled “Network Distance” for cases other than “Intra” presents the paths through the network connecting the two crosslinked proteins found in other existing PPI databases. Clicking on the entry under the table column entitled “Num Prot Pairs” displays all proteins corresponding to each peptide, and when available, their apportioned weights, as computed by ProteinProphet, indicating their relative likelihood as the origin of the crosslinked peptide. Clicking the “NGL Viewer” button displays the crosslink within the PDB structure in NGL (Rose and Hildebrand, 2015). In the case of inter-protein crosslinks with an available multimeric structure or docked model, the user has the option of additionally coloring the interaction surface of both proteins at a specified distance (Figure 4). This can help illustrate and validate how the two interacting proteins may recognize and fit with one another. Surfaces defined in this manner can potentially be used to design drugs that bind to proteins to disrupt specific interactions.

Data displayed in the table can be downloaded as a tab delimited file. Quantified data in this format can easily be subjected to clustering analysis to identify common trends among

crosslinks. Data can also be imported to Cytoscape to view the PPI network as a graphical display.

Interaction Network View

It is often useful to view crosslink data as a PPI network with crosslinks organized according to their corresponding proteins. In XLinkDB, one can view a network in a browser with Cytoscape, where nodes represent crosslink sites in proteins, and edges represent non-redundant crosslinks. Nodes corresponding to different sites in the same protein are normally grouped in circles according to Uniprot ID, though any layout is supported. The interaction network clearly indicates which proteins interact with one another. One can explore the network, zooming in or out, and centering the view near any region. Users have the option to save the current view so that when they return to the network at a later time, it will revert to that same position and zoom level.

Exploring the network is facilitated by options to color edges according to many crosslink attributes, such as the distance in the context of protein structures, distance in known PPI networks, confidence of identification, and numbers of corresponding protein pairs. When coloring by the various attributes, one can click on the edge to obtain the same pertinent information displayed upon clicking the value in the table view, as described above.

For networks with quantitative information, edges can be colored by any available sample log2ratio (Figure 5), as well as p-value. When coloring by log2ratio for which raw data is available, one can click on an edge to view the chromatograms of contributing light and heavy parent peaks. One can easily view and compare ratios corresponding to different contributing samples for crosslinks of interest by saving a view near the crosslinks and selecting edge coloring for each ratio, in succession. When edges are colored by log2ratio, the edge is solid if there is only a single corresponding protein pair possible, but dashed if there are two or more. This can occur when two or more proteins share indistinguishable peptide sequences and is used to alert users in that case that the quantitation may not apply literally to the two displayed most likely corresponding proteins. When viewing the edges colored by the number of corresponding protein pairs, one can click on the edge to view all proteins corresponding to each peptide, and when available, their apportioned likelihoods, as computed by ProteinProphet. Proteome level quantitation information in datasets can be displayed as node colors of the same hue scale used to color network edges according to crosslink-level quantitation.

For each of the crosslink attributes by which edge coloring is possible, one can also apply filtering according to a specified range of values. For example, one can select to view only crosslinks with low p-values with respect to some ratios, and high p-values with respect to others. Or one can specify to view only crosslinks with a single possible corresponding protein pair. One can also explicitly filter for a list of specified proteins, whereby only nodes of those proteins, and edges connecting them, are shown (Figure 5). One can obtain a table view of the network data currently displayed by clicking the 'Generate/Download Network Table' button. This is especially useful when one has applied filtering criteria to view the network, since those criteria are applied to the resulting table view.

Viewing the 3D Interactome

In cases in which a network has quantitation with respect to a number of related samples, one can use the XLinkDB Cytoscape network to construct a movie of images of the quantified PPI networks that comprise a 3D interactome. By coloring edges according to each sample log₂ratio in succession and downloading for each, a PNG file, one can easily assemble the network images into a rotating movie (see Materials and Methods). Viewing the movie as a 3D interactome, it is possible to experience the dynamic changes in quantified samples. For example, if the quantified samples are obtained over the time course of a perturbation, observing the movie of time-successive images of the network with edges colored by quantitation gives a clear indication of the protein conformational changes occurring over time. Movies of the network images with quantitation corresponding to increased doses of a drug applied to a sample, in a similar manner, help to illustrate the changes in conformation induced by exposure to the drug. Changes observed among subsets of edges are strong indicators of alternative protein conformations.

As an example, a set of light SILAC labeled human HeLa samples were treated with concentrations of the Hsp90 inhibitor, 17-N-allylamino-17-demethoxygeldanamycin (17-AAG) ranging from 0nM (control) to 1000nM¹⁹. Following treatment, cells were mixed with equal amounts of heavy labeled cells treated with no drug, crosslinked, subjected to ReACT XL-MS²⁰, and quantified with MassChroQ¹⁴. The data were uploaded to XLinkDB where the PPI network was viewed in Cytoscape, with crosslink edges colored in series by log₂ratio of samples exposed to increasing doses of 17-AAG. Images of heat shock protein HS90A and HS90B components of this network were downloaded as PNG files, embedded in an HTML rotator, and captured as a movie with QuickTime (Figure S1). Viewing the movie as a 3D interactome, one can see drug-induced increased levels of HS90A homo-dimerization at residue 443, and HS90A-HS90B hetero-dimerization at residues 443 and 435, respectively. In contrast, no increase in analogous HS90B homo-dimerization at residue 435 was observed. The majority of changes occur with as little as 100nM 17-AAG treatment, though some additional changes in observed levels of crosslinks occur with higher doses of the drug. Exploring the 3D interactome, researchers can efficiently identify subsets of crosslinks with similar quantitation changes that could specify alternative drug-induced conformations, and which could be used to guide structure modeling.

CONCLUSIONS

Interactome data as acquired with XL-MS offer a new window through which to observe both protein conformational and PPI information in complex samples such as functional organelles, live cells, and tissues. We have previously developed a suite of informatics tools embodied within web-based XLinkDB that enable researchers to visualize large cross-linked datasets, resulting interaction networks, as well as links on protein and protein interaction structures and models. Quantitative XL-MS studies are only beginning to reveal conformational and interactome dynamics but hold great potential for providing unique molecular-level insight on biological function. However, these datasets require further advanced informatics tools to enable multi-dimensional interactome visualization. To facilitate such exploration, we developed and implemented new tools to store and view

quantitative crosslink datasets in XLinkDB in its 3.0 release, available online at <http://xlinkdb.gs.washington.edu/xlinkdb/>. These new capabilities allow researchers to visualize interactome dynamics in conjunction with proteome-level quantitation with respect to virtually any perturbation. Crosslinks of interest can easily be assessed for the quality of their quantitation, as well as other features such as consistency with known protein structures. In the future, we intend to enable upload of XL-MS data analyzed with XLinkProphet²¹, and store the mass spectra of the released peptides, upon which the crosslink identifications are based.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

We thank members of the Bruce Lab for helpful discussions and suggestions. This work was supported by grants 5R01-GM086688, 2R01-HL110349, 5U19AI107775, 1R01HL142628, NSF DBI 659680, and in part by the UW Proteome Resource UWPR95794UWPR.

REFERENCES

1. Chavez JD; Eng JK; Schweppe DK; Cilia M; Rivera K; Zhong X; Wu X; Allen T; Khurgel M; Kumar A; Lampropoulos A; Larsson M; Maity S; Morozov Y; Pathmasiri W; Perez-Neut M; Pineyro-Ruiz C; Polina E; Post S; Rider M; Tokmina-Roszyk D; Tyson K; Vieira Parrine Sant'Ana D; Bruce JE, A General Method for Targeted Quantitative Cross-Linking Mass Spectrometry. *PLoS One* 2016, 11, (12), e0167547. [PubMed: 27997545]
2. Yu C; Kandur W; Kao A; Rychnovsky S; Huang L, Developing new isotope-coded mass spectrometry-cleavable cross-linkers for elucidating protein structures. *Anal Chem* 2014, 86, (4), 2099–106. [PubMed: 24471733]
3. Chavez JD; Schweppe DK; Eng JK; Zheng C; Taipale A; Zhang Y; Takara K; Bruce JE, Quantitative interactome analysis reveals a chemoresistant edgotype. *Nat Commun* 2015, 6, 7928. [PubMed: 26235782]
4. Zhang H; Tang X; Munske GR; Tolic N; Anderson GA; Bruce JE, Identification of protein-protein interactions and topologies in living cells with chemical cross-linking and mass spectrometry. *Mol Cell Proteomics* 2009, 8, (3), 409–20. [PubMed: 18936057]
5. Kao A; Chiu CL; Vellucci D; Yang Y; Patel VR; Guan S; Randall A; Baldi P; Rychnovsky SD; Huang L, Development of a novel cross-linking strategy for fast and accurate identification of cross-linked peptides of protein complexes. *Mol Cell Proteomics* 2011, 10, (1), M110 002212.
6. Kandur WV; Kao A; Vellucci D; Huang L; Rychnovsky SD, Design of CID-cleavable protein cross-linkers: identical mass modifications for simpler sequence analysis. *Org Biomol Chem* 2015, 13, (38), 9793–807. [PubMed: 26269432]
7. Tang X; Munske GR; Siems WF; Bruce JE, Mass spectrometry identifiable cross-linking strategy for studying protein-protein interactions. *Anal Chem* 2005, 77, (1), 311–8. [PubMed: 15623310]
8. Holding AN, XL-MS: Protein cross-linking coupled with mass spectrometry. *Methods* 2015, 89, 54–63. [PubMed: 26079926]
9. Leitner A; Faini M; Stengel F; Aebersold R, Crosslinking and Mass Spectrometry: An Integrated Technology to Understand the Structure and Function of Molecular Machines. *Trends Biochem Sci* 2016, 41, (1), 20–32. [PubMed: 26654279]
10. Orban-Nemeth Z; Beveridge R; Hollenstein DM; Rampler E; Stranzl T; Hudecz O; Doblmann J; Schlogelhofer P; Mechtler K, Structural prediction of protein models using distance restraints derived from cross-linking mass spectrometry data. *Nat Protoc* 2018, 13, (3), 478–494. [PubMed: 29419816]

11. Ong SE; Blagoev B; Kratchmarova I; Kristensen DB; Steen H; Pandey A; Mann M, Stable isotope labeling by amino acids in cell culture, SILAC, as a simple and accurate approach to expression proteomics. *Mol Cell Proteomics* 2002, 1, (5), 376–86. [PubMed: 12118079]
12. Rampler E; Stranzl T; Orban-Nemeth Z; Hollenstein DM; Hudecz O; Schlogelhofer P; Mechtler K, Comprehensive Cross-Linking Mass Spectrometry Reveals Parallel Orientation and Flexible Conformations of Plant HOP2-MND1. *J Proteome Res* 2015, 14, (12), 5048–62. [PubMed: 26535604]
13. Schweppe DK; Zheng C; Chavez JD; Navare AT; Wu X; Eng JK; Bruce JE, XLinkDB 2.0: integrated, large-scale structural analysis of protein crosslinking data. *Bioinformatics* 2016, 32, (17), 2716–8. [PubMed: 27153666]
14. Valot B; Langella O; Nano E; Zivy M, MassChroQ: a versatile tool for mass spectrometry quantification. *Proteomics* 2011, 11, (17), 3572–7. [PubMed: 21751374]
15. Zheng C; Weisbrod CR; Chavez JD; Eng JK; Sharma V; Wu X; Bruce JE, XLink-DB: database and software tools for storing and visualizing protein interaction topology data. *J Proteome Res* 2013, 12, (4), 1989–95. [PubMed: 23413830]
16. Shannon P; Markiel A; Ozier O; Baliga NS; Wang JT; Ramage D; Amin N; Schwikowski B; Ideker T, Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* 2003, 13, (11), 2498–504. [PubMed: 14597658]
17. Nesvizhskii AI; Keller A; Kolker E; Aebersold R, A statistical model for identifying proteins by tandem mass spectrometry. *Anal Chem* 2003, 75, (17), 4646–58. [PubMed: 14632076]
18. Li H; Leung K; Wong M, idock: A multithreaded virtual screening tool for flexible ligand docking. 2012, 77–84.
19. Chavez JD; Schweppe DK; Eng JK; Bruce JE, In Vivo Conformational Dynamics of Hsp90 and Its Interactors. *Cell Chem Biol* 2016, 23, (6), 716–26. [PubMed: 27341434]
20. Weisbrod CR; Chavez JD; Eng JK; Yang L; Zheng C; Bruce JE, In vivo protein interaction network identified with a novel real-time cross-linked peptide identification strategy. *J Proteome Res* 2013, 12, (4), 1569–79. [PubMed: 23413883]
21. Keller A; Chavez JD; Bruce JE, Increased Sensitivity with Automated Validation of XL-MS Cleavable Peptide Crosslinks. *Bioinformatics* 2018, (bty720).

Peptide A	Protein A	Residue Number A	Peptide B	Protein B	Residue Number B	Network Distance	XL Distance (Å)	Number ReACT IDs	Confidence	Num Prot Pairs	log2ratio_1_Control	log2ratio_2_100nmAAG	log2ratio_3_250nmAAG	log2ratio_4_500nmAAG	log2ratio_5_1000nmAAG
FYEQFSKNIK	HS90A_HUMAN	443	FYEQFSKNIK	HS90A_HUMAN	443	1	36.47	35	1	1	-0.17	3.3	2.74	3.34	3.17
ILKVR	HS90A_HUMAN	410	FYEQFSKNIK	HS90A_HUMAN	443	Intra	24.78	14	1	3	0.45	1.76	1.73	1.94	1.86
IMKAGALR	HS90A_HUMAN	615	FYEQFSKNIK	HS90A_HUMAN	443	Intra	20.83	104	1	2	0.06	0.55	0.94	1.26	1.1
FYEAFSKNLK	HS90B_HUMAN	435	FYEQFSKNIK	HS90A_HUMAN	443	1	####	74	1	1	-0.06	0.98	0.94	1.34	1.26
APFDLFENK	HS90B_HUMAN	347	FYEAFSKNLK	HS90B_HUMAN	435	Intra	17.75	57	1	1	0.45	-0.03	0.37	0.01	0.74

Figure 1.
Table view illustrating crosslink abundance ratios of various samples with respect to a common control as a sortable heatmap.

Figure 2.
(A) Line chart showing crosslink quantitation log2ratio value trends in various samples. (B)
Bar chart showing numbers of crosslinks with various log2ratio values in a single sample.

Figure 3. Example chromatographic traces of light and heavy parent masses in two replicate files, displayed upon clicking their crosslink log₂ratio value in the table. The indicated normalization multiplier was applied to the peak area ratios to achieve a log₂ratio distribution centered at zero.

Figure 4.

NGL Viewer showing identified inter-protein crosslinks spanning mouse mitochondrial proteins NADH-ubiquinone oxidoreductase (NDUS1) and NADH dehydrogenase flavoprotein 1 (NDUV1), with the interaction surfaces of each protein displayed at 10 Å in a model dimeric structure generated by iDock¹⁸.

Figure 5.
Cytoscape interaction network filtered for human heat shock proteins HS90A and HS90B, with edges colored by log2ratio in the selected sample, treatment of HeLa cells with 1000nM 17-N-allylamino-17-demethoxygeldanamycin (17-AAG), an Hsp90 inhibitor, versus control.