Published in final edited form as:

Psychol Med. 2019 June ; 49(8): 1392–1399. doi:10.1017/S0033291719000047.

The structure of genetic and environmental influences on normative personality, abnormal personality traits, and personality disorder symptoms

Kenneth S. Kendler^{1,2,3}, Steven H. Aggen^{1,2}, Nathan Gillespie^{1,2}, Robert F. Krueger⁴, Nikolai Czajkowski^{5,6}, Eivind Ystrom^{5,6}, and T. Reichborn-Kjennerud^{5,7}

¹Virginia Institute for Psychiatric and Behavioral Genetics, Richmond, VA, USA

²Departments of Psychiatry, Virginia Commonwealth University, Richmond, VA, USA

³Human and Molecular Genetics, Virginia Commonwealth University, Richmond, VA, USA

⁴Department of Psychology, University of Minnesota, Minneapolis, MN, USA

⁵Division of Mental Health, Norwegian Institute of Public Health, Oslo Norway

⁶Department of Psychology, University of Oslo, Oslo Norway

⁷Institute of Clinical Medicine, University of Oslo, Oslo Norway

Abstract

Background.—Can the structure of genetic and environmental influences on normative personality traits (NPTs), abnormal personality traits (APT), and DSM-IV criteria for personality disorders (PD) fit a high or low congruence model positing, respectively, close or more limited etiologic continuity?

Method.—Exploratory factor analysis was applied to transformed correlation matrices from Cholesky twin decompositions obtained in OpenMx. In 2801 adult twins from the Norwegian Institute of Public Health Twin Panel, NPTs and APTs were assessed by self-report using the Big Five Inventory (BFI) and PID-5-Norwegian Brief Form (PID-5-NBF), respectively. PDs were assessed at interview using the Structured Interview for DSM-IV Personality (SIDP-IV).

Results.—The best model yielded three genetic and three unique environmental factors. Genetic factors were dominated, respectively, by (i) high loadings on nearly all PDs and NPT/APT neuroticism and compulsivity, (ii) negative loadings on NPT agreeableness/conscientiousness and positive loadings on APT/PD measures of antisocial traits, and (iii) negative loadings on NPT extraversion and histrionic PD, and positive loadings on APT detachment and schizoid/avoidant PD. Unique environmental factors were dominated, by (i) high loadings on all PDs, (ii) high

Author for correspondence: Kenneth S. Kendler, kenneth.kendler@vcuhealth.org.

Conflict of interest. None.

Ethical standards. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. Written informed consent was obtained from all participants. Ethical approval for both waves was granted by The Norwegian Data Inspectorate and the Regional Ethical Committee.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291719000047>

loadings on all APT dimensions and NPT neuroticism, and (iii) negative loadings on NPT extraversion and positive loadings on NPT detachment/avoidant PD.

Conclusions.—Two genetic and one environmental common factor were consistent with a high congruence model while one genetic and two environmental factors were more supportive of a low congruence model. The relationship between genetic and environmental influences on personality assessed by NPTs, APTs, and PDs is complex and does not fit easily into a low or high congruence model.

Keywords

Genetic modeling; personality; personality disorder; twins

The Five Factor Model, including dimensions of Extraversion, Agreeableness, Conscientiousness, Neuroticism, and Openness to Experience (John and Srivastava, 1999), has been widely accepted as a description of normative personality. These normative personality traits (NPTs), typically assessed by self-report questionnaires, are moderately heritable in twin studies (McGue, 2002; Livesley and Jang, 2008; South *et al.*, 2015). Assessments for maladaptive/abnormal personality traits (APTs), based on the same Five Factor Model, have been developed (Clark, 1993; Clark *et al.*, 1994; Livesley and Jackson, 2009; Krueger *et al.*, 2012), studied in twin populations, and also found to be heritable (Livesley *et al.*, 1993; Jang *et al.*, 1996; South *et al.*, 2016). The DSM criteria for personality disorders (PDs) as first proposed in DSM-III (American Psychiatric Association, 1980), modestly modified in DSM-III-R (American Psychiatric Association, 1987) and DSM-IV (American Psychiatric Association, 2000), and now included in DSM-5 (American Psychiatric Association, 2013), have also been examined in twin samples – most often as ‘criteria counts’ – and shown to be heritable (Livesley *et al.*, 1993; Torgersen *et al.*, 2000; Jang and Vernon, 2001; Kendler *et al.*, 2008; Livesley and Jang, 2008; Reichborn-Kjennerud, 2008).

What is the nature of the etiologic relationship between NPTs, APTs, and PDs? Some studies suggest that DSM PDs, while not equivalent to NPYs, are nonetheless well represented by the Five Factor Model (Costa and Widiger, 2002; Saulsman and Page, 2004; Samuel and Widiger, 2008) suggesting that the traits of normal and abnormal personality exist on a single normally distributed continuum with NPTs, APTs, and PDs located at increasingly extreme locations (Livesley, 2007). Others contend that a four-factor structure is more appropriate as openness has little relationship with the DSM PDs (Dyce and O’Connor, 1998; Livesley *et al.*, 1998; Austin and Deary, 2000; Markon *et al.*, 2005).

Since the seminal publication by Robins and Guze (1970), familial/genetic factors have played a central role in the evaluation of psychiatric diagnostic classifications. A few studies have investigated common genetic influences on normal and abnormal personality with variable results (Jang and Livesley, 1999; Wright *et al.*, 2017).

In Norwegian population-based twins, we obtained measures of NPTs, APTs, and PDs. Three prior publications from our group on this sample have focused on the relationship between two of these three dimensions: NPTs and APTs (Kendler *et al.*, 2017), APTs and

PDs (Reichborn-Kjennerud *et al.*, 2017), and NPTs and PDs (Czajkowski *et al.*, 2018). In this report, for the first time, we conduct a computationally demanding multivariate twin analysis of all three of these personality domains with the specific goal of determining the degree of cross-domain coherence in their genetic and environmental determinants.

We seek to evaluate two hypotheses. A *high congruence model* predicts four or five genetic and environmental factors each anchored by one of the big-five dimensions of normal personality and reflected in parallel measures of APTs and PDs. Such results would reflect a close relationship between the genetic and environmental influences on NPTs, APTs, and PDs. A *low congruence model*, by contrast, predicts factors that largely index these three distinct domains of personality and which would therefore reflect limited etiologic continuity between NPTs, APTs, and PDs.

Methods

Our sample came from the Norwegian Institute of Public Health (NIPH) Twin Panel (Harris *et al.*, 2006). Twins born 1967–1979 ($N = 15\,370$) were identified through the Norwegian National Medical Birth Registry. Repeated contact with them began in 1992. Data used here came from an interview (1999–2004, wave 1) that assessed DSM-IV Axis I and Axis II disorders, and a self-report follow-up questionnaire (2010–2011, wave 2) including items assessing normative and maladaptive personality. Of the 3221 twin pairs eligible for wave 1, 1391 complete pairs (43.2%) and 19 single twins (0.6% pairwise) participated (63% female; mean age = 28.2 years, range = 19–36). Of the 2801 twins eligible for wave 2, 2393 twins (86.8%, 1063 twin pairs and 267 single twins) participated (64% female, mean age = 37.8 years, range = 30–44). Wave 1 interviews were administered by advanced psychology students or psychiatric nurses who received standardized training and supervision. Written informed consent was obtained from all participants. Ethical approval was granted by The Norwegian Data Inspectorate and the Regional Ethical Committee.

Measures

In the wave 2 self-report twin questionnaire, normative personality domains were assessed by the 44-item Big Five Inventory (BFI) (John *et al.*, 1991). A single common factor model was fitted to the five-point Likert-type item response option sets to test the unidimensionality of each BFI constructs. Based on satisfactory single-factor model fits, item aggregate sum score variables were constructed for each BFI construct. Six maladaptive personality domain constructs (Negative Emotion, Detachment, Antagonism, Disinhibition, Psychoticism, and Compulsivity) were assessed using a shortened 36-item version of the PID-5: the PID-5-Norwegian Brief Form (PID-5-NBF) (Krueger *et al.*, 2012). These items used a four-point Likert-type response format. Item integer scores were reverse coded and anchored at zero so that higher values indicated greater levels of maladaptive behavior. Similar psychometric modeling was conducted to verify the unidimensionality of each of the PID-5-NBF constructs. The resulting sum score aggregate variables were often highly positively skewed. Log transformations were applied to each sum score incremented by one prior to twin modeling.

All criteria for lifetime DSM-IV Axis II PDs were assessed using a Norwegian version of the Structured Interview for DSM-IV Personality (SIDP-IV) at wave 1 (Pfohl *et al.*, 1995). As detailed previously, inter-rater reliability of these assessments was high with a mean intra-class correlation of the endorsed criteria across the PDs equal to +0.89 (Kendler *et al.*, 2008).

Six of these 10 core PDs were also reassessed at a wave 2 interview 10 years later. The SIDP-IV used non-pejorative questions organized into topical sections rather than by individual PDs to facilitate the flow of the interview. The SIDP-IV interview was conducted after the Composite International Diagnostic Interview (CIDI) to aid interviewers in distinguishing between persistent long-term patterns of behavior and temporary states due to existing Axis I disorders. A 5-year rule was used to ensure the persistence of each of the PD criteria as representing the expression of a more trait-like personality pathology. Each symptom criterion was scored on an ordered four-point scale (0 = absent, 1 = subthreshold, 2 = present, and 3 = strongly present). Due to the low prevalence of full PD diagnoses, a dimensional approach was adopted scaling PD liability using a count of the positively endorsed criterion and a model for ordered thresholds. First, each PD criterion was dichotomized based on a sub-threshold cut-off (0 = absent *v.* 1 = sub-threshold or greater). These binary criteria were then summed for each respective PD to form aggregate variables. Due to strong positive skewness of these sum scores, each count variable was reorganized into a four-category (three threshold) ordinal scale (0 = 0 criteria, 1 = 1–2 criteria, 2 = 3–4 criteria, and 3 = 4 or more criteria) for each PD. We have previously examined the validity of this rescaling approach by testing whether a multiple threshold model is consistent with an assumption of bivariate normality to determine whether these categorized counts of endorsed criteria reflected ordered differences in severity on a single, normally distributed continuum of liability. This assumption was empirically supported for all 10 PDs examined.

Twin modeling

A Cholesky decomposition was specified for the full set of five NPTs, six APTs, and 10 PDs. Due to the size of this decomposition, the Cholesky was fit to a two-group MZ/DZ twin data structure with sexes combined. Model optimization was carried out using a Full Information Maximum Likelihood (FIML) raw data method as implemented in the R 3.1.2 (R Development Core Team, 2014) OpenMx package (Boker *et al.*, 2011; Neale *et al.*, 2016). To fit a Cholesky decomposition to twin data with so many variables (42 total), variables of different scale types were specified to facilitate optimization. The five BFI sum score variables were symmetrically distributed and treated as quasicontinuous. The six log-transformed PID-5-NBF variables were also treated as quasi-continuous. The 10 rescaled PD symptom count variables were treated as ordinal. However, rather than estimating all three thresholds per PD, a specification using fixed and free thresholds was implemented. With three thresholds per PD, the first threshold is fixed to 0, the second to 1, and the third estimated. This parameterization allows for a mean and variance to be estimated for each ordinal PD variable (Mehta *et al.*, 2004). This approach reduces the computational demands of threshold integration and allows all the observed variables to be modeled as continuously distributed using a FIML approach. Additionally, FIML makes use of all the available data providing more robust inference against missing data.

The main Cholesky decompositions included all A (additive genetic) and E (unshared environment-error) paths. Next, the Cholesky non-zero lower diagonal A and E path estimates were converted into covariance matrices and then rescaled to 21×21 correlation matrices. These additive genetic and unique environmental correlation matrices were then submitted to exploratory factor analyses (EFA) in which different numbers of factors were extracted using an oblique Geomin rotation in the R Psych Package (Revelle, 2015). The number of factors to be extracted was guided by Scree plots and content analyses (Horn, 1965). The minimum residual factor extraction method in the R psych package was used for all EFA solutions, since in some cases, the correlation matrices derived from the Cholesky path estimates resulted in non-positive definite conditions. This required adjustments to the Eigen values of the original derived correlation matrix to obtain a rotated EFA solution.

Due to the large number of freely estimated parameters in these Cholesky decompositions (755 Cholesky paths plus MZ and DZ means and thresholds parameterized as previously described for the full ACE model and 524 for the AE model) and the amount of computational time required to carry out FIML optimization, it became apparent that relying on a conventional single estimate of global fit to evaluate models was not practical. An alternative strategy was adopted where the degree of consistency of the end results of the EFA rotated solutions would be compared across a series of separate Cholesky optimizations obtained using different sets of reasonably bounded random starting values. Run times on a Unix multi-cluster server using parallel processing across multiple cores varied for the full ACE and reduced AE Cholesky decomposition but were in general much longer than typical optimization problems (e.g. 10 or fewer variables). Model fit was evaluated by the Akaike (AIC) and Bayesian information criteria (BIC) (Akaike, 1987; Raftery, 1993).

Results

Model fitting

We ran 10 models each with different random start values. Two models did not converge and were not further considered. Results of the remaining eight models, which took a mean (S.D.) of 18.9 (1.8) days to converge, are seen in Table 1. The fifth run produced the best fit by both indices and therefore became our best-fit model. Runs 1 and 6 produced the second and third best-fit models.

Genetic factors

Examination of the Scree plot provided evidence for three or four genetic factors. While the three-factor solution was readily interpretable, the four-factor solution was less satisfactory (online Supplementary Appendix Table S1) in that the second factor from the three-factor solution was divided into two poorly defined and less interpretable dimensions. We therefore focused on the three-factor solution, the loadings of which are shown in Table 2 and illustrated in Fig. 1a–c.

The first factor had salient loadings on eight of the 10 DSM-IV PDs (all but schizoid and antisocial), N from the BFI, and negative emotionality and compulsivity from the PID-5-NBF. We labeled this a *General Personality Disorder/Neuroticism* common genetic factor.

Factor 2 had strong negative loadings on agreeableness and conscientiousness from the BFI, and strong positive loadings on antagonism and detachment from the PID-5-NBF and antisocial PD. It also had moderate positive loadings on disinhibition and psychoticism from the PID-5-NBF. We labeled this an *Antisocial/Antagonism* common genetic factor.

Factor 3 had substantial negative loadings on extraversion and openness from the BFI and histrionic PD, and positive loadings on detachment from the PID-5-NBF and schizoid and avoidant PD. This latent dimension was called a *Schizoid-Introversion* common genetic factor.

Correlations between the genetic factors was strongest between factors 1 and 2 (+0.57), modest between factors 1 and 3 (−0.27), and estimated at zero between factors 2 and 3.

Environmental factors

A Scree plot for the exploratory oblique factor analyses of the environmental correlation matrix suggested a four-factor solution. However, because the fourth factor had only a single loading >0.40 (online Supplementary Appendix Table S2), we focused on the more easily interpretable three-factor solution presented in Table 3 and Fig. 2.

The first factor had strong positive loadings $\geq +0.47$ on all 10 DSM-IV PDs ranging with quite small loadings on all BFI and PID-5-NBF scales ($<|0.05|$). This was clearly a *General Personality Disorder* common environmental factor.

Factor 2 had strong loadings on all six of the PID-5-NBF scales with a moderately strong loading on the BFI N scale. Loadings on this factor for all the PDs and other BFI scales were small. This factor was best characterized as a *Pathological Personality Trait/Neuroticism* common environmental factor.

Factor 3 had substantial negative loadings on two BFI scales (extraversion and openness) and positive loadings on BFI neuroticism scale, the Detachment scale from the PID-5-NBF and avoidant PD. This is perhaps best termed an *Introversion-Avoidant* common environmental factor.

Inter-factor correlations for these environmental common factors were all modest: factors 1 and 2 +0.25, factors 1 and 3 +0.10, and factors 2 and 3 +0.19. The congruency coefficients between the genetic and environmental factors are presented in Table 3 (Lorenzo-Seva and ten Berge, 2006).

Discussion

We sought to clarify the underlying structure of genetic and environmental risk factors for individual PDs and specific NPT and APT dimensions, positing two heuristic hypotheses. The high congruence model predicted four or five genetic and environmental common factors each anchored in a distinct ‘big five’ dimension and each loading on a corresponding set of APTs and PDs. The low congruence model posited modest etiologic continuity between normative and disordered personality, predicting that the common genetic and

environmental factors would be largely ‘level-specific’ in their influence – which separate etiological factors reflecting normative personality, maladaptive personality, and PDs.

Our results were complex and did not cleanly fit the predictions of either hypothesis. Rather, our findings provided support for some features of each. The first genetic factor resembled a ‘level-specific’ PD factor loading strongly on eight of the 10 PDs. However, it was also ‘anchored’ by the NPT of N and the related APT traits of negative emotionality and compulsivity. The second genetic common factor was clearly cross-level but was anchored by not one but two big five factors: A and C. The third factor most closely resembled that predicted by the high congruence model with clinically sensible cross-level loadings anchored by the single NPT of extraversion.

By contrast, the first two of our three environmental factors were consistent with the low congruence model with, respectively, strong level-specific loadings on all 10 PDs and all six PID-F factors (plus N). The third environmental factor, however, was as predicted by a high-congruence model anchored by the BFI dimension of extraversion with appropriate cross loadings on detachment from the PID-5-NBF and avoidant PD.

An overall interpretation of our findings also needs to consider the inter-factor correlations, the only prominent one of which was between the first and second genetic factors (+0.57). Of note, eight of the 11 variables with prominent loadings on the first genetic factor are PDs, and four out of seven of the prominent loadings on the second factor come from the PID-5-NBF. Thus, this correlation is likely driven by genetic factors that broadly predispose both to most PDs and high scores on most PID-5-NBF scales. This result suggests an aggregate association between genetic risk factors for PDs and APTs but not necessarily the strong genetic correlations between clinically similar individual PDs and APT dimensions predicted by the high congruence model.

The structure of genetic factors found in the current study resembles that in our previous examination of these 10 DSM-IV PDs (Kendler *et al.*, 2008). The best-fitting model included three genetic factors. The first had substantial loadings on most PDs reflecting a general vulnerability to PD pathology. The second had strong loadings on antisocial and borderline PD, whereas the third had substantial loadings only on schizoid and avoidant PD.

Our results also resemble several prior three-factor models of personality. Our genetic factors closely resemble the three factors proposed in the Eysenck Personality Questionnaire (Eysenck and Eysenck, 1975) with factor 1 reflecting neuroticism, factor 2 ‘psychoticism’ (low agreeableness, antagonism, antisocial traits, and PID-5-NBF psychoticism), and factor 3 extraversion. Our findings are also similar to those presented as part of a hierarchical meta-analytic study of normal and abnormal personality where the factors Negative Emotionality, Disinhibition, and Positive emotionality were related to (1) Neuroticism, (2) Agreeableness (reversed) and Conscientiousness (reversed), and (3) Extraversion and Openness (Markon *et al.*, 2005).

A controversial topic during the DSM-5 deliberations was the placement of the Alternative DSM-5 model for PDs in section III (American Psychiatric Association, 2013; Zachar *et al.*, 2016). This alternative model was predicated on a high coherence hypothesis – that strong

and specific relationships exist between individual PDs and specific NPT and APT dimensions. The PID-5-NBF was explicitly developed to measure key dimensions of this alternative model (Krueger *et al.*, 2012; South *et al.*, 2016). Our results provide evidence both for and against the alternative DSM-5 model.

Our second and third genetic factors indicate that for the antisocial PD and the cluster of schizoid, histrionic, and avoidant PDs, genetic risk factors were well reflected in our measures of normative and APTs. This is consistent with the high coherence hypothesis and supportive of the alternative DSM model. But such relationships were not found for other important PDs including borderline, narcissistic, obsessive–compulsive, and schizotypal. Results for the first genetic factor were more ambiguous. On the one hand, it could be viewed as largely a ‘PD-specific’ factor as would be predicted by our low-coherence model, hence considered evidence against the alternative DSM model. On the other hand, it could be seen as a version of the ‘General Criteria for PD’ in the alternative model which, in our data, also reflects the normative and APTs of neuroticism and negative emotionality.

Unexpectedly, our environmental risk factors were considerably less supportive of the alternative model than our genetic factors. Our findings suggest that, with the likely exception of extraversion and associated APTs and PDs, most environmental experiences that impact on personality are not shared between PDs and the expected specific APT or NPT dimensions.

In aggregate, our results present evidence against the hypothesis that the full pattern of DSM-IV PDs can be captured by the standard APT and/or NPT dimensions. Whether this picture would differ if facet-level data were available for APT and NPT will need to be addressed by future studies. Our own results do, however, show significant areas of overlap of PDs, APTs, and NPT dimensions that were considerably stronger when reflecting genetic than environmental risk processes. Further empirical and psychometric research will be needed to bring these assessment methods into a closer consilience.

Our results can be profitably viewed in the context of our prior analyses with this sample where we showed that, *in aggregate*, the APT dimensions of the PID-5-NBF accounted for considerably more of the genetic variance of individual PDs (Reichborn-Kjennerud *et al.*, 2017) than the NPT dimensions of the BFI (Czajkowski *et al.*, 2018). These results suggest at a global level an expected pattern that is a closer genetic relationship between PDs and APTs than between PDs and NPTs. We observe this in our own findings of inter-factor genetic correlations. However, in this paper, we are asking a more stringent question, and one more relevant to evaluating the alternative DSM model. Instead of asking whether all the APT or NPT dimensions can predict genetic liability to an individual PD, we inquire about the genetic relationships between individual PDs and specific NPT and APT dimensions. It is to this question that we see support both for and against the high coherence alternative DSM model.

Limitations

These results should be interpreted in the context of eight potentially important methodological limitations. First, results derive from a Norwegian population-based twin

sample and may not generalize to other populations. Second, the instruments used to assess NPTs and APTs were both short and lacked facet level traits. Greater communality might have been found between PDs and NPT and APT had they been assessed with longer instruments. Third, PDs were assessed at the wave 1 interview and NPTs and APTs at wave 2. Could the first genetic or environmental common factors loading strongly on the PDs in part reflect occasion-specific effects? This is inconsistent with prior evidence of very high genetic correlations for PDs measured at waves 1 and 2 (Kendler *et al.*, 2015; Reichborn-Kjennerud *et al.*, 2015). To test this formally, we fitted AE models to all scales from the BFI and PID-5-NBF and the six PDs that were assessed at wave 2 and then repeated these analyses using the scores from the same PDs from the wave 1 assessment. We readily identified a PD-dominant genetic and environmental common factor in these analyses (see online Supplementary Appendix Tables S3–S6). The mean loadings on the six PDs in PD-dominant genetic common factor did not significantly differ in the analyses containing the wave 1 (0.57 ± 0.13) v. wave 2 data (0.49 ± 0.11) ($t = 0.40$, $df = 10$, N.S.). Findings were similar for the environmental common factors: 0.58 ± 0.04 v. 0.56 ± 0.07 ($t = 0.21$, $df = 10$, N.S.). The results suggest that our PD-dominant genetic and environmental first common factors were not the result of the time difference in assessment of the DSM-IV PDs and the BFI and PID-5-NBF.

Fourth, given the complexity of our modeling, it was not feasible to explicitly examine sex differences. When fitted to simpler models, prior twin analyses in this sample have consistently been unable to detect sex effects on PDs (Kendler *et al.*, 2006; Reichborn-Kjennerud *et al.*, 2007; Torgersen *et al.*, 2008). Fifth, attrition occurred in this twin sample from the original birth registry through our wave 2 assessments. Detailed analyses of this attrition show that cooperation was strongly predicted by female sex, monozygosity, older age, and education, but by neither psychiatric symptoms nor drug use (Tambs *et al.*, 2009). While attrition bias in our results is possible, the full-information maximum likelihood methods used here are typically robust against large biases due to missingness. Sixth, given the very long run time for our AE models and the lack of substantial shared environmental effects for our PD, BFI and PID-5-NBF measures in simpler analyses (Kendler *et al.*, 2006; 2017; Reichborn-Kjennerud *et al.*, 2007; Torgersen *et al.*, 2008; South *et al.*, 2016), we did not extensively explore twin models containing such ‘C’ effects. We did however run two such ‘ACE’ models (mean run time 33 days). Both fit considerably worse than our best-fit AE models by both AIC and BIC.

Seventh, given the rarity of full syndromal PDs in the general population, we were unable to study PDs defined by diagnostic thresholds. Instead, we took the commonly used approach of analyzing a quantitative index of the number of endorsed criteria. While not ideal, population-based twin samples would have to be very large to contain sufficient numbers of individuals meet full PD criteria to support these kinds of analyses.

Finally, because of the very large number of parameters in our Cholesky decompositions, we could not rely on conventional methods of sequentially fitting and comparing increasingly more complex models. More skepticism is therefore warranted about our findings. However, we examined the genetic and environmental factor structures from the two other best-fit models (runs 1 and 6). As seen in the online Supplementary Appendix Tables S7 and S8,

while these results differed in some modest details from those of the best-fit model, the overall pattern was very similar, providing added confidence in our findings.

Conclusion

Three major approaches have arisen in the fields of psychiatry and clinical psychology toward the assessment of personality. The inter-relationship between them has been subject of controversy. In this report we provide, for the first time to our knowledge, information about the relationship between the genetic and environmental determinants of these three major approaches. The results are nuanced and provide limited support for both the low and high congruence models. In aggregate, they suggest caution regarding simple overarching theoretical claims about the etiological relationships between NPTs, APTs, and PDs.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments.

Financial support. This project was supported by the Research Council of Norway grant 196148/V50 and NIH grant RO1DA037558. Previous collection and analysis of twin data from this project was in part supported by grant MH-068643 from the National Institutes of Health and grants from the Norwegian Research Council, the Norwegian Foundation for Health and Rehabilitation, the Norwegian Council for Mental Health, the Borderline Foundation, and the European Commission. These funding agencies played no role in the design and conduct of the study, its collection, management, analysis, and interpretation of the data or in the preparation, review, or approval of the manuscript.

References

- Akaike H (1987) Factor analysis and AIC. *Psychometrika* 52, 317–332.
- American Psychiatric Association (1980) Diagnostic and Statistical Manual of Mental Disorders, 3rd Edn. Washington, DC: American Psychiatric Association.
- American Psychiatric Association (1987) Diagnostic and Statistical Manual of Mental Disorders, Revised 3rd Edn. Washington, DC: American Psychiatric Association.
- American Psychiatric Association (2000) Diagnostic and Statistical Manual of Mental Disorders, 4th Edn. Revised, Fourth-Revised ed. Washington, DC: American Psychiatric Association.
- American Psychiatric Association (2013) Diagnostic and Statistical Manual of Mental Disorders, 5th Edn. *DSM-5*. Washington, DC: American Psychiatric Association.
- Austin EJ and Deary IJ (2000) The ‘four As’: a common framework for normal and abnormal personality? *Personality and Individual Differences* 28, 977–995.
- Boker S, Neale M, Maes H, Wilde M, Spiegel M, Brick T, Spies J, Estabrook R, Kenny S, Bates T, Mehta P and Fox J (2011) Openmx: an open source extended structural equation modeling framework. *Psychometrika* 76, 306–317. [PubMed: 23258944]
- Clark LA (1993) Schedule for Nonadaptive and Adaptive Personality (SNAP): Manual for Administration, Scoring, and Interpretation. Minneapolis: University of Minnesota Press.
- Clark LA, Vorhies L and McEwen JL (1994) Personality disorder symptomatology from the five-factor model perspective In Costa PT and Widiger TA (eds), *Personality Disorders and the Five-Factor Model of Personality*. Washington, DC: American Psychological Association, 125–147.
- Costa PT and Widiger TA (2002) *Personality Disorders and the Five-Factor Model of Personality*, 2nd Edn. Washington, DC: American Psychological Association.
- Czajkowski N, Aggen SH, Krueger RF, Kendler KS, Neale MC, Knudsen GP, Gillespie N, Røysamb E, Tambs K and Reichborn-Kjennerud T (2018) A twin study of normative personality and DSM-IV

- personality disorder criterion counts; evidence for separate genetic influences. *American Journal of Psychiatry* 175, 649–656. [PubMed: 29558815]
- Dyce JA and O'Connor BP (1998) Personality disorders and the five-factor model: a test of facet-level predictions. *Journal of Personality Disorders* 12, 31–45. [PubMed: 9573518]
- Eysenck HJ and Eysenck SBG (1975) *Manual of the Eysenck Personality Questionnaire*. London: Hodder and Stoughton.
- Harris JR, Magnus P and Tambs K (2006) The Norwegian Institute of Public Health twin program of research: an update. *Twin Research and Human Genetics* 9, 858–864. [PubMed: 17254421]
- Horn JL (1965) A rationale and test for the number of factors in factor analysis. *Psychometrika* 30, 179–185. [PubMed: 14306381]
- Jang KL and Livesley WJ (1999) Why do measures of normal and disordered personality correlate? A study of genetic comorbidity. *Journal of Personality Disorders* 13, 10–17. [PubMed: 10228923]
- Jang K and Vernon PA (2001) Genetics In Livesley WJ (ed.), *Handbook of Personality Disorders: Theory, Research, and Treatment*. New York: Guilford Press, pp. 177–195.
- Jang KL, Livesley WJ, Vernon PA and Jackson DN (1996) Heritability of personality disorder traits: a twin study. *Acta Psychiatrica Scandinavica* 94, 438–444. [PubMed: 9020996]
- John OP and Srivastava S (1999) The big-five trait taxonomy: history, measurement, and theoretical perspectives In Pervin LA and John OP (eds), *Handbook of Personality: Theory and Research*, 2nd Edn. New York: Guilford Press, pp. 102–139.
- John OP, Donahue EM and Kentle RL (1991) *The Big Five Inventory-Versions 4a and 54* Berkeley CA: University of California Berkeley, Institute of Personality and Social Research.
- Kendler KS, Czajkowski N, Tambs K, Torgersen S, Aggen SH, Neale C and Reichborn-Kjennerud T (2006) Dimensional representations of DSM-IV Cluster A personality disorders in a population-based sample of Norwegian twins: a multivariate study. *Psychological Medicine* 36, 1583–1591. [PubMed: 16893481]
- Kendler KS, Aggen SH, Czajkowski N, Roysamb E, Tambs K, Torgersen S, Neale MC and Reichborn-Kjennerud T (2008) The structure of genetic and environmental risk factors for DSM-IV personality disorders: a multivariate twin study. *Archives of General Psychiatry* 65, 1438–1446. [PubMed: 19047531]
- Kendler KS, Aggen SH, Neale MC, Knudsen GP, Krueger RF, Tambs K, Czajkowski N, Ystrom E, Orstavik RE and Reichborn-Kjennerud T (2015) A longitudinal twin study of cluster A personality disorders. *Psychological Medicine* 45, 1531–1538. [PubMed: 25394477]
- Kendler KS, Aggen SH, Gillespie N, Neale MC, Knudsen GP, Krueger RF, Czajkowski N, Ystrom E and Reichborn-Kjennerud T (2017) The genetic and environmental sources of resemblance between normative personality and personality disorder traits. *Journal of Personality Disorders* 31, 193–207. [PubMed: 27322578]
- Krueger RF, Derringer J, Markon KE, Watson D and Skodol AE (2012) Initial construction of a maladaptive personality trait model and inventory for DSM-5. *Psychological Medicine* 42, 1879–1890. [PubMed: 22153017]
- Livesley WJ (2007) A framework for integrating dimensional and categorical classifications of personality disorder. *Journal of Personality Disorders* 21, 199–224. [PubMed: 17492921]
- Livesley WJ and Jackson DN (2009) *Dimensional Assessment of Personality Pathology-Basic Questionnaire (DAPP-BQ) Technical Manual*. Port Huron, MI: Sigma Assessment Systems, Inc.
- Livesley WJ and Jang KL (2008) The behavioral genetics of personality disorder. *Annual Reviews of Clinical Psychology* 4, 247–274.
- Livesley WJ, Jang KL, Jackson DN and Vernon PA (1993) Genetic and environmental contributions to dimensions of personality disorder. *American Journal of Psychiatry* 150, 1826–1831. [PubMed: 8238637]
- Livesley WJ, Jang KL and Vernon PA (1998) Phenotypic and genetic structure of traits delineating personality disorder. *Archives of General Psychiatry* 55, 941–948. [PubMed: 9783566]
- Lorenzo-Seva U and ten Berge JMF (2006) Tucker's congruence coefficient as a meaningful index of factor similarity. *Methodology* 2, 57–64.

- Markon KE, Krueger RF and Watson D (2005) Delineating the structure of normal and abnormal personality: an integrative hierarchical approach. *Journal of Personality and Social Psychology* 88, 139–157. [PubMed: 15631580]
- McGue M (2002) Genetics of personality In Rimoin R, Pyeritz R and Korf B (eds.), *Emery & Rimoin's Principles and Practice of Medical Genetics*, 4th Edn. vol. 3 London: Churchill Livingstone, pp. 2791–2800.
- Mehta PD, Neale MC and Flay BR (2004) Squeezing interval change from ordinal panel data: latent growth curves with ordinal outcomes. *Psychological Methods* 9, 301–333. [PubMed: 15355151]
- Neale MC, Hunter MD, Pritikin JN, Zahery M, Brick TR, Kirkpatrick RM, Estabrook R, Bates TC, Maes HH and Boker SM (2016) *Openmx 2.0: extended structural equation and statistical modeling*. *Psychometrika* 81, 535–549. [PubMed: 25622929]
- Pfohl B, Blum N and Zimmerman M (1995) *Structured Interview for DSM-IV Personality (SIDP-IV)*. Iowa City: University of Iowa, Department of Psychiatry.
- Raftery AE (1993) Bayesian model selection in structural equation models In Bollen KA and Long JS (eds.), *Testing Structural Equation Models*, 1st Edn. Newbury Park, CA: Sage, pp. 163–180.
- R Development Core Team (2014) *The R Project for Statistical Computing; R: A language and environment for statistical computing*. [3.1.2]. Vienna, Austria, R Foundation for Statistical Computing Computer Program.
- Reichborn-Kjennerud T (2008) Genetics of personality disorders. *Psychiatric Clinics of North America* 31, 421–440. [PubMed: 18638644]
- Reichborn-Kjennerud T, Czajkowski N, Neale MC, Orstavik RE, Torgersen S, Tambs K, Roysamb E, Harris JR and Kendler KS (2007) Genetic and environmental influences on dimensional representations of DSM-IV cluster C personality disorders: a population-based multivariate twin study. *Psychological Medicine* 37, 645–653. [PubMed: 17134532]
- Reichborn-Kjennerud T, Czajkowski N, Ystrom E, Orstavik R, Aggen SH, Tambs K, Torgersen S, Neale MC, Roysamb E, Krueger RF, Knudsen GP and Kendler KS (2015) A longitudinal twin study of borderline and antisocial personality disorder traits in early to middle adulthood. *Psychological Medicine* 45, 3121–3131. [PubMed: 26050739]
- Reichborn-Kjennerud T, Krueger RF, Ystrom E, Torvik FA, Rosenstrom TH, Aggen SH, South SC, Neale MC, Knudsen GP, Kendler KS and Czajkowski NO (2017) Do DSM-5 Section II personality disorders and Section III personality trait domains reflect the same genetic and environmental risk factors? *Psychological Medicine* 47, 2205–2215. [PubMed: 28414014]
- Revelle W (2015) *Psych: Procedures for Psychological, Psychometric, and Personality Research*. [1.5.6]. Northwestern University. Computer Program.
- Robins E and Guze SB (1970) Establishment of diagnostic validity in psychiatric illness: its application to schizophrenia. *American Journal of Psychiatry* 126, 983–987. [PubMed: 5409569]
- Samuel DB and Widiger TA (2008) A meta-analytic review of the relationships between the five-factor model and DSM-IV-TR personality disorders: a facet level analysis. *Clinical Psychology Review* 28, 1326–1342. [PubMed: 18708274]
- Saulsman LM and Page AC (2004) The five-factor model and personality disorder empirical literature: a meta-analytic review. *Clinical Psychology Review* 23, 1055–1085. [PubMed: 14729423]
- South SC, Reichborn-Kjennerud T, Eaton NR and Krueger RF (2015) Genetics of personality In Mikulincer M, Shaver PR, Cooper ML and Larsen RJ (eds.), *APA handbooks in psychology. APA handbook of personality and social psychology*, Vol. 4. Personality processes and individual differences. Washington, DC, US: American Psychological Association, pp. 31–60.
- South SC, Krueger RF, Knudsen GP, Ystrom E, Czajkowski N, Aggen SH, Neale MC, Gillespie NA, Kendler KS and Reichborn-Kjennerud T (2016) A population based twin study of DSM-5 maladaptive personality domains. *Personality Disorders* 8, 366–375. [PubMed: 27797545]
- Tambs K, Ronning T, Prescott CA, Kendler KS, Reichborn-Kjennerud T, Torgersen S and Harris JR (2009) The Norwegian Institute of Public Health twin study of mental health: examining recruitment and attrition bias. *Twin Research and Human Genetics* 12, 158–168. [PubMed: 19335186]
- Torgersen S, Lygren S, Oien PA, Skre I, Onstad S, Edvardsen J, Tambs K and Kringlen E (2000) A twin study of personality disorders. *Comprehensive Psychiatry* 41, 416–425. [PubMed: 11086146]

- Torgersen S, Czajkowski N, Jacobson K, Reichborn-Kjennerud T, Roysamb E, Neale MC and Kendler KS (2008) Dimensional representations of DSM-IV cluster B personality disorders in a population-based sample of Norwegian twins: a multivariate study. *Psychological Medicine* 38, 1617–1625. [PubMed: 18275631]
- Wright ZE, Pahlen S and Krueger RF (2017) Genetic and environmental influences on Diagnostic and Statistical Manual of Mental Disorders-Fifth Edition (DSM-5) maladaptive personality traits and their connections with normative personality traits. *Journal of Abnormal Psychology* 126, 416–428. [PubMed: 28368150]
- Zachar P, Krueger RF and Kendler KS (2016) Personality disorder in DSM-5: an oral history. *Psychological Medicine* 46, 1–10. [PubMed: 26482368]

Fig. 1.

(a) Parameter estimates for genetic common factor 1 from the best-fit model (run 5) from a three-factor oblique rotation of the genetic correlation matrix. Normative personality traits are in blue, abnormal personality traits are in red, and personality disorders are in yellow. Abbreviations and full names for the variables are as follows: EXTR, extraversion; AGRE, agreeableness; CONS, conscientiousness; NEUR, neuroticism; OPEN, openness to experience; NEM, negative emotionality; DET, detachment; ANT, antagonism; DIS, disinhibition; COM, compulsivity; PSYC, psychoticism; PARA, paranoid personality

disorder; SCHI, schizoid personality disorder; SCHAT, schizotypal personality disorder; ANTI, antisocial personality disorder; BORD, borderline personality disorder; HIST, histrionic personality disorder; NARC, narcissistic personality disorder; AVOI, avoidant personality disorder; DEPE, dependent personality disorder; OBCO, obsessive–compulsive personality disorder. (b) Parameter estimates for genetic common factor 2 from a three-factor oblique rotation of the genetic correlation matrix. Normative personality traits are in blue, abnormal personality traits in red, and personality disorders are in yellow. (c) Parameter estimates for genetic common factor 3 from a three-factor oblique rotation of the genetic correlation matrix. Normative personality traits are in blue, abnormal personality traits are in red, and personality disorders are in yellow.

Fig. 2. Parameter estimates for environmental common factors from the best-fit model (run 5) from a three-factor oblique rotation of the environmental correlation matrix. Normative personality traits are in blue, abnormal personality traits are in pink, and personality disorders are in yellow. For abbreviations, see legend to Fig. 1a.

Table 1.

Model fit statistics and run times for models run with random start values

Model ^a	-2lnL	AIC	BIC	Run time (days)
Run 1	80 881	-24 453	-301 026	16 days
Run 2	81 029	-24 305	-300 878	19 days
Run 3	83 223	-22 111	-298 683	21 days
Run 4	81 290	-24 044	-300 616	19 days
Run 5	80 877	-24 457	-301 029	16 days
Run 6	80 955	-24 379	-300 951	21 days
Run 7	81 044	-24 290	-300 863	20 days
Run 8	81 261	-24 073	-300 646	19 days

^aFive hundred and twenty-four estimated parameters, 1410 observations.**Bolded** – best-fit model by both AIC and BIC.

Table 2.

Loadings on genetic factors for dimensions of normative personality, personality disorder traits, and personality disorders from the best-fit model

Personality scale	Loadings by factor ^a		
	1	2	3
Normative personality			
Extraversion	−0.18	−0.13	−0.81
Agreeableness	0.14	−0.86	−0.17
Conscientiousness	−0.18	−0.59	−0.10
Neuroticism	0.69	−0.08	0.23
Openness	0.05	0.29	−0.55
Personality disorder traits			
Negative emotionality	0.61	0.28	0.28
Detachment	0.05	0.64	0.68
Antagonism	−0.04	0.89	−0.07
Disinhibition	0.44	0.45	−0.02
Compulsivity	0.50	0.24	0.12
Psychoticism	0.37	0.57	0.01
Personality disorders			
Paranoid	0.89	−0.02	−0.12
Schizotypal	0.85	−0.19	0.01
Schizoid	0.38	0.24	0.48
Antisocial	0.15	0.60	−0.35
Borderline	0.87	0.09	−0.34
Histrionic	0.74	0.03	−0.66
Narcissistic	0.52	0.37	−0.39
Avoidant	0.64	0.00	0.53
Dependent	1.00	−0.42	0.05
Obsessive–compulsive	0.33	0.26	−0.09

Personality scale	Loadings by factor^a		
	1	2	3
Inter-factor correlations			
Factor 1	–	+0.57	–0.27
Factor 2	–	–	0.00
Factor 3	–	–	–

^aLoading bolded if it was the highest for that dimension or, if not highest, an absolute value of >0.45.

Table 3.

Loadings on environmental factors for dimensions of normative personality, personality disorder traits, and personality disorders from the best-fit model

Personality scale	Loadings by factor		
	1	2	3
Normative personality			
Extraversion	−0.01	−0.08	−0.83
Agreeableness	0.00	−0.28	−0.25
Conscientiousness	−0.01	−0.22	−0.28
Neuroticism	0.03	0.43	0.39
Openness	0.03	0.06	−0.36
Personality disorder traits			
Negative emotionality	−0.03	0.74	0.19
Detachment	0.02	0.41	0.33
Antagonism	0.04	0.47	−0.23
Disinhibition	0.00	0.57	−0.09
Compulsivity	−0.02	0.53	−0.03
Psychoticism	0.04	0.60	0.04
Personality disorders			
Paranoid	0.68	−0.07	0.07
Schizotypal	0.70	−0.06	0.14
Schizoid	0.47	−0.11	0.05
Antisocial	0.47	0.12	−0.10
Borderline	0.63	0.06	−0.01
Histrionic	0.58	0.07	−0.19
Narcissistic	0.62	0.01	−0.14
Avoidant	0.45	−0.01	0.31
Dependent	0.54	0.02	0.11
Obsessive–compulsive	0.54	0.07	−0.11
Inter-factor correlations			

Personality scale	Loadings by factor		
	1	2	3
Factor 1	–	+0.25	+0.10
Factor 2	–	–	+0.19
Factor 3	–	–	–
Congruency coefficients* between genetic and environmental factors			
Genetic factor 1	+0.83	+0.40	+0.24
Genetic factor 2	+0.15	+0.69	+0.07
Genetic factor 3	–0.20	+0.21	+0.81