Published in final edited form as:

Nat Ecol Evol. 2019 May ; 3(5): 765–771. doi:10.1038/s41559-019-0871-9.

Ancient Genomes Indicate Population Replacement in Early Neolithic Britain

Selina Brace^{1,*}, Yoan Diekmann^{2,*}, Thomas J. Booth^{1,*}, Lucy van Dorp³, Zuzana Faltyskova², Nadin Rohland⁴, Swapan Mallick^{3,5,6}, Iñigo Olalde⁴, Matthew Ferry^{4,6}, Megan Michel^{4,6}, Jonas Oppenheimer^{4,6}, Nasreen Broomandkhoshbacht^{4,6}, Kristin Stewardson^{4,6}, Rui Martiniano⁷, Susan Walsh⁸, Manfred Kayser⁹, Sophy Charlton^{1,12}, Garrett Hellenthal³, Ian Armit¹⁰, Rick Schulting¹¹, Oliver E. Craig¹², Alison Sheridan¹³, Mike Parker Pearson¹⁴, Chris Stringer¹, David Reich^{4,5,6,#}, Mark G. Thomas^{2,3,#,∞}, and Ian Barnes^{1,#,∞}

¹Department of Earth Sciences, Natural History Museum, London SW7 5BD, UK ²Research Department of Genetics, Evolution and Environment, University College London, London WC1E 6BT, UK ³UCL Genetics Institute (UGI), University College London, London WC1E 6BT, UK ⁴Department of Genetics, Harvard Medical School, Boston, Massachusetts 02115, USA ⁵Broad Institute of MIT and Harvard, Cambridge, Massachusetts 02142, USA ⁶Howard Hughes Medical Institute, Harvard Medical School, Boston, Massachusetts 02115, USA ⁷Department of Genetics, University of Cambridge, 20 Downing Place, Cambridge, CB2 3DY, UK ⁸Department of Biology, Indiana University-Purdue University Indianapolis, 723 W. Michigan Street, Indianapolis, IN 46202, USA ⁹Department of Genetic Identification, Erasmus MC University Medical Centre Rotterdam, Rotterdam, The Netherlands ¹⁰School of Archaeological and Forensic Sciences, University of Bradford, Richmond Building, Bradford, BD7 1AZ, UK ¹¹Institute of Archaeology, University of Oxford, 36 Beaumont St., Oxford, OX1 2PG, UK ¹²Bioarch, Environment Building, University of York, Wentworth Way, York, YO10 5DD, UK ¹³National Museums Scotland, Chambers Street, Edinburgh, EH1 1JF, UK ¹⁴Institute of Archaeology, University College London, 31-34 Gordon Square, London, WC1H 0PY, UK

Abstract

The roles of migration, admixture and acculturation in the European transition to farming have been debated for over 100 years. Genome-wide ancient DNA studies indicate predominantly Aegean ancestry for continental Neolithic farmers, but also variable admixture with local

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use: http://www.nature.com/authors/editorial_policies/license.html#terms

∞ corresponding authors.

* these authors contributed equally

these authors co-supervised the work

Data availability—BAM files (a file per library, before re-aligning around InDels; see Supplementary Table 1) have been deposited at ENA under study accession PRJEB31249.

Competing Interests Statement

We confirm that none of the authors have a competing financial and/or non-financial interest in relation to the work described.

Author Contributions I.B. and M.G.T conceived the project, and Y.D., S.B., Z.F., O.C. and T.B. contributed to the project design. S.B., Y.D., T.B., L.v.D., N.R., S.M., I.O., M.F., M.M., J.O., N.B., K.S., R.M., S.C. and S.W. generated and analysed data. I.B., M.G.T., Y.D., S.B., T.B., M.K., S.W., G.H., I.A., R.S., O.C., A.S., M.P.P., C.S. and D.R. contributed to the sampling strategy and the interpretation of results. I.B., M.G.T., Y.D., S.B. and T.B. wrote the paper, with contributions from all other authors.

Mesolithic hunter-gatherers. Neolithic cultures first appear in Britain ca. 4000 BCE, a millennium after they appear in adjacent areas of continental Europe. The pattern and process of this delayed British Neolithic transition remains unclear. We assembled genome-wide data from six Mesolithic and 67 Neolithic individuals found in Britain, dating from 8500-2500 BCE. Our analyses reveal persistent genetic affinities between Mesolithic British and Western European hunter-gatherers. We find overwhelming support for agriculture being introduced to Britain by incoming continental farmers, with small, geographically-structured levels of hunter-gatherer ancestry. Unlike other European Neolithic populations, we detect no resurgence of hunter-gatherer ancestry at any time during the Neolithic in Britain. Genetic affinities with Iberian Neolithic individuals indicate that British Neolithic people were mostly descended from Aegean farmers who followed the Mediterranean route of dispersal. We also infer considerable variation in pigmentation levels in Europe by ca. 6000 BCE.

Introduction

The transition to farming marks one of the most important ecological shifts in human evolution. The processes by which this transition occurred have been a matter of intense debate for over a century^{1–3}, although across continental Europe ancient DNA studies indicate a predominant role for expanding Neolithic farmer populations of mostly Aegean ancestry (Aegean Neolithic Farmers; ANF)^{4–15}. ANF-derived populations dispersed throughout Europe via two major routes; one along the Mediterranean and the other through Central and into Northern Europe^{7, 11}. Both dispersing populations introgressed repeatedly with local Mesolithic foragers, which gradually increased their proportion of European Mesolithic ancestry^{7, 13–15}.

The nature of the Neolithic transition in Britain remains unclear because of the millennium-long delay in its appearance after the establishment of farming in adjacent regions of continental Europe^{1–3}, and the lack of genome-wide data from British Mesolithic hunter-gatherers. Whilst there is universal agreement amongst archaeologists that there was a dramatic change in material culture in Britain around 4000 BCE, there are divergent views regarding the extent to which this change was influenced by cultural or demographic processes^{1–3}. The British Isles lie farthest from the Aegean origin^{4–15} of the migrating farmers that influenced the development of the Neolithic across Europe, are geographically isolated from continental Europe by large bodies of water, and had maritime climates which differ from the majority of mainland Europe; all factors which may alter the nature of the adoption of farming. The relationship between British and continental European Mesolithic populations is also of interest as Britain geographically abuts two genetically-distinct but contemporaneous populations: Western European and Scandinavian Mesolithic hunter-gatherers (WHGs & SHGs), and could have potentially harboured ancestry from earlier (~19,000 to 15,000 BCE) Magdalenian Palaeolithic hunter-gatherer populations^{16–18}.

Results

Here, we report the first whole genome data from six Mesolithic (including ‘Cheddar Man’ from Gough’s Cave, Somerset) and 16 Neolithic British individuals, and combine it with

data from 51 previously published Neolithic British individuals¹² to characterise the Mesolithic to Neolithic transition in Britain (Figure 1, Supplementary Figure S16). Our Mesolithic samples date from 8750-8459 cal. BCE (Early Mesolithic Aveline's Hole, Somerset, England) to 4256-3803 cal. BCE (Late Mesolithic Cnoc Coig, Oronsay, western Scotland). Our Neolithic samples date from 3951-3780 cal. BCE (Early Neolithic McArthur Cave, western Scotland) to 2570-2347 cal. BCE (Late Neolithic Isbister, Orkney, Scotland). We combined data generated in two different ways. For 35 individuals, we generated new whole genome shotgun sequencing data, including the first full genomes from the British Mesolithic (at 2.3x) and Neolithic (at 10.7x) individuals. For all samples we enriched next generation sequencing libraries for approximately 1.24 million single nucleotide polymorphisms (SNPs) (median coverage 0.88x). When available we merged data obtained from both methods and identify the most likely allele at each locus (see Material and Methods). These were combined with ancient genomic data from 67 previously reported individuals^{4–7, 9–12, 14, 16–22} (see Supplementary Table S1) and modern genomic data from diverse global populations²³.

All British Mesolithic individuals cluster with Western and Scandinavian hunter-gatherers in a principal components analysis (Figure 2). By contrast, all directly-dated individuals who post-date 4000 BCE, and undated individuals associated with Neolithic monuments, cluster tightly near Iberian and Central European Middle Neolithic individuals. By examining the degree of allele sharing of British Mesolithic individuals to various European hunter-gatherer individuals/groups (SHG, EHG and El Mirón, see Supplementary Figures S1-S4), we were able to attribute them confidently to the WHG group. Comparison of British Mesolithic individuals to different Mesolithic WHGs (Berry au Bac - France, Ranchot⁸⁸ - France, Loschbour - Luxembourg, La Braña - Spain, KO1 - Hungary; Supplementary Figures S5-S6, S11-S14) indicates that all most closely resemble Loschbour. When we compared the remaining British Mesolithic genomes to Loschbour and Cheddar Man (our highest-coverage British Mesolithic sample; ~2.3X), we found no significant excess of shared drift for either individual, indicating that Loschbour, Ranchot and the British Mesolithic samples do not form separate clusters (Supplementary Figure S7).

To investigate the proportions of Aegean farmer-related ancestry in the British samples we modelled them as mixtures of ANFs and European WHGs using the *qpAdm* method, which studies ensembles of f_4 statistics (Figure 3, Supplementary Figure S8)²⁴. The genomes of all British Mesolithic individuals can be explained almost entirely by WHG ancestry, the remainder (<7.3%) likely stemming from poorly matching portions of the genome. Most of the ancestry in all British Neolithic individuals could be attributed to ANFs (>56%, ~74% on average), indicating a substantial shift in ancestry with the transition to farming. To investigate the proximate source of ANF ancestry in British Neolithic individuals, we examined affinities with Early Neolithic individuals from Iberia and Central Europe. We compare Early over Middle Neolithic individuals as the latter are contemporary with the British Early Neolithic, making them an unlikely direct source. For all British Neolithic individuals considered we inferred more shared drift with Early Neolithic Iberians (Figure 4A, Supplementary Figure S9). However, these f_4 statistic-based inferences may be sensitive to levels of WHG admixture, such that the similarity in WHG admixture proportions in Early Neolithic Iberian and British samples, but lower estimates in Central European Early

Neolithic individuals, is driving the inference of an Iberian rather than Central European source for Early British farmers. To examine this possibility in more detail we performed a more powerful haplotype-based analysis.

Using a chromosome painting approach²⁵ we obtained patterns of haplotype matching between our high coverage British Neolithic sample and a global modern reference panel (Supplementary Materials Section 7). We found similar patterns of donor haplotype matching in the British Neolithic genome to those inferred for other high coverage Neolithic genomes from Ireland and Iberia. These were more similar than the same profiles obtained for high coverage Neolithic genomes from Central Europe (Figure 5a). Inferred ancestry coefficients (see Materials and Methods) further support this connection between the British, Irish and Iberian Neolithic⁶ and are consistent with the same ancestral populations bringing the Neolithic to Britain and Ireland (Figure 5b,c, Supplementary Table S8). Additional modelling using global modern populations²⁶ as ancestry surrogates suggests this population is best represented today by components found in French and Spanish peoples (Figure 5c, Supplementary Table S9).

In order to test for a potential second ANF ancestry stream from Central Europe, we explicitly modelled WHG and Early Neolithic populations in *qpGraph* (see Supplementary Fig. S23 and Supplementary Table S10). The results suggest that the limited Central European Neolithic admixture we find in British Neolithic populations is regionally structured, with populations from England showing the highest levels of admixture, followed by populations from Scotland. We infer no Central European admixture in Neolithic farmers from Wales. However, we caution that the model fits are poor and so these inferences should be considered preliminary.

We inferred some significant geographic structure in WHG admixture proportions among the British Early Neolithic individuals (see Supplementary Table S4 for statistical comparison of inferred WHG admixture proportions); those from Wales retain the lowest levels of WHG admixture, followed by those from South-West and Central England. Neolithic individuals from South-East England and Scotland show significantly higher WHG admixture proportions. These proportions remain stable from the Early into the Middle/Late Neolithic. To infer levels of WHG introgression occurring between Iberian Early Neolithic populations and early British farmers, we estimated admixture proportions using *qpAdm*²⁴. We detected little excess (~10%) WHG ancestry beyond that already present in Iberian Early Neolithic individuals, supporting little or no additional admixture with British hunter-gatherers, particularly in Wales, South-West and Central England (Figures 3 and 4B; Supplementary Table S4). This result appears to be slightly at odds with the f_4 results presented in Supplementary Figure S7, which indicate that some British Neolithic samples share genetic affinities with Cheddar Man over Loschbour, although it is difficult to say in these cases whether this is due to genuine substantial admixture with British WHGs or with other WHGs in northern Europe. We regressed individual WHG ancestry proportions in British Neolithic farmers (shown in Supplementary Figure S8) against latitude and longitude and found a significant positive southwest to northeast cline (Supplementary Figure S15).

To further explore WHG introgression in Britain we applied ALDER27 to pairs of Early Neolithic regional samples to estimate the timing of WHG/ANF admixture events (Supplementary Table S3). Only Early Neolithic farmers from western Scotland show evidence of WHG introgression within 10 generations. Two individuals from Raschoille Cave had estimated introgression events occurring 4.0 ± 3.4 generations before they lived, which is sufficiently recent in their past that it likely occurred in Britain. The elevated levels of WHG ancestry we see in Neolithic samples from South East England are older, and therefore probably a result of farmer-forager interactions in mainland Europe. Chronological modelling (using OxCal 4.328) of available direct Early Neolithic radiocarbon data from individuals showing ANF ancestry suggests that continental farmers arrive in Britain by 3975-3722 cal. BCE (95% confidence), 481 years after to 27 years before (95% confidence) the death of our latest Mesolithic individual showing no ANF ancestry (Supplementary Materials Section 6). Our model suggests that continental farmers arrive marginally earlier in the west (although see the discussion in the Supplementary Materials), and rapidly disperse into other regions of Britain (including the Orkney Islands). The latest regional appearance of ANF ancestry is in Central England and occurs 59 to 386 years (95% confidence) after it first appears in Britain.

To explore variation in pigmentation of European populations, we predicted pigmentation in higher-coverage Mesolithic and Neolithic Europeans using HRISplex-S29. We infer that Cheddar Man mostly likely had blue/green eyes, dark brown possibly black hair, and dark or dark to black skin, while our highest coverage early Neolithic individual had brown eyes, black possibly dark brown hair, and dark to intermediate skin (see Supplementary Materials Section 3). Together with the pigmentation prediction outcomes we obtained for Loschbour and La Braña, these results imply that different pigmentation levels coexisted in Europe by around ca.6000 BCE.

Discussion

The six British Mesolithic genomes examined here are typical of WHGs, indicating that this population spread to the furthest northwestern point of early Holocene Europe after moving from southeastern Europe, or further east, from approximately 12,000 BCE¹⁷. This genetic similarity among British and European Mesolithic individuals spans a period in Britain (ca. 8,500-4,000 BCE) that includes the cultural transition to the Late Mesolithic and the separation of Britain from continental Europe. Our analyses indicate that the appearance of Neolithic practices and domesticates in Britain ca. 4,000 BCE was mediated overwhelmingly by immigration of farmers from continental Europe^{1–2}, and strongly reject the hypothesised adoption of farming by indigenous hunter-gatherers as the main process³. British farmers were substantially descended from Iberian Neolithic-related populations whose ancestors had expanded along a Mediterranean route^{6, 11}, although with a minority portion of their ancestry from populations who took the Danubian route¹². The affinities we find between Neolithic individuals from the British Isles and modern individuals from France are consistent with populations sharing ancestry with Neolithic groups in Iberia moving into northern France via the Atlantic seaboard and/or southern France, mixing to a limited degree with Neolithic populations from Central Europe before travelling across the Channel^{1–2, 30}.

One explanation for the British Neolithic cline in WHG ancestry is that a single population moved across Britain from a western entry point, and progressively admixed with local hunter-gatherers. This scenario is consistent with the western distribution of megalithic cultures along the Atlantic seaboard³¹, and is supported by the radiocarbon evidence suggesting a marginally earlier date for the arrival of ANF ancestry in the west of Britain¹. However, the lack of evidence for substantive WHG introgression into British Neolithic populations – outside of western Scotland – favours this cline reflecting multiple source populations with variable proportions of WHG admixture having entered different parts of Britain. This interpretation is consistent with archaeological evidence for regional British Neolithic cultures showing links to varied parts of mainland Europe² and our qpGraph analysis indicating geographically-structured Neolithic Central European admixture. Overall, the regional variation in ancestry of British Neolithic populations likely reflects both differing degrees of admixture between farmers and local foragers (e.g. western Scotland), and multiple continental source populations carrying variable WHG and Neolithic Central European ancestry.

Evidence for only low levels of WHG introgression among British Early Neolithic people is striking given the extensive and complex admixture processes inferred for continental Neolithic populations^{7, 13–15, 32–33}. Low levels of admixture between these two groups on the wave front of farming advance in continental Europe have been attributed to the maintenance of cultural and reproductive boundaries for several centuries after initial contact, before more extensively mixing³². Similarly, isotopic and genetic data from the west coast of Scotland suggest the potential coexistence of genetically distinct hunter-fisher-gatherers and farmers, albeit for a maximum of a few centuries³⁴. However, there is no evidence for a resurgence of WHG ancestry in the British Neolithic, consistent with limited evidence for Mesolithic cultural artefacts in Britain beyond 4000 BC^{1–2}, and with a major dietary shift from marine to terrestrial resources at this time (see Supplementary Materials Section 5)³⁵.

Conclusion

In contrast to other European regions, the transition to farming in Britain occurred with little introgression from resident foragers – either during initial colonization, or throughout the Neolithic. This may reflect low Late Mesolithic population density in Britain and/or an introduction of farming by populations who had mastered the technologies needed to thrive in northern and western continental Europe during the previous two millennia^{1–2}.

Materials and Methods

Ancient DNA Extraction and Sequencing

DNA extractions and library preparations for all samples with newly reported data were conducted in a dedicated ancient DNA laboratory (NHM, London). We used approximately 25mg of finely drilled bone powder and followed the DNA extraction protocol described in Dabney *et al.*³⁶ but replaced the Zymo-Spin V column binding apparatus with a high pure extender assembly from the High Pure Viral Nucleic Acid Large Volume Kit (Roche). Library preparations followed the partial uracil–DNA–glycosylase treatment described in

Rohland *et al.*³⁷ and a modified version of the Meyer & Kircher³⁸ protocol. Library modifications: the initial DNA fragmentation step was not required; all clean-up steps used MinElute PCR purification kits (Qiagen). The index PCR step included double indexing³⁹, the polymerase AmpliTaq Gold and the addition of 0.4mg/mL BSA. The index PCR was set for 20 cycles with three PCR reactions conducted per library. Libraries were screened for DNA preservation on an Illumina NextSeq platform, with paired-ends reads. Promising libraries were further enriched at the NHM using in-solution hybridisation capture enrichment kits (Mybait-3) from MYcroarray. The baits were designed to cover ca. 20K SNP's (5,139 functional and 15,002 neutral SNP's) at 4x tiling. The capture protocol followed the manufacturer's instructions in the Mybait manual v3. Captured libraries were sequenced on an Illumina NextSeq platform (NHM) using paired-ends reads. Newly reported data from 36 of these libraries was also obtained at the dedicated ancient DNA lab in Harvard Medical School by enriching in solution for approximately 1.24 million targeted SNPs. We sequenced these libraries on an Illumina NextSeq500 instrument, iteratively sequencing more until we estimated that the additional number of targeted SNPs hit per newly generated sequence was less than 1 per 100.

Bioinformatics

All sequence reads underwent adapter and low-quality base trimming, and overlapping reads pairs were collapsed with AdapterRemoval⁴⁰. Non-collapsed reads and those of length less than 30bp were discarded, and the remaining aligned against the hs37d5 human reference genome with BWA⁴¹. Mapped reads with a mapping quality of at least 30 were merged per individual and re-aligned around InDels with GATK⁴². Resulting BAM files were split by flowcell and lane, and empirical ATLAS⁴³ post mortem damage patterns estimated per individual per lane for lanes with at least 5.5 million reads, otherwise per individual per flowcell. ATLAS BQSR base quality score recalibration tables were generated per lane for lanes with at least 5.5 million reads, otherwise per flowcell. We generated recalibrated BAM files per individual with ATLAS recalBAM, and used these to estimate mitochondrial contamination and determine mitochondrial and Y-chromosome haplogroups with ContamMix⁴⁴, Yleaf⁴⁵ and Phy-Mer⁴⁶. We considered mitochondrial contamination to be tolerable if 0.98 was included in the confidence intervals. Haploid genotypes were called with ATLAS "allelePresence" with theta fixed at 0.001, determining the most likely base at a given position. Heterozygosity estimates, shown in Supplementary Figure S10, were computed with ATLAS "estimate Theta" and a default window size of 1Mbp, excluding windows that overlap with telo- or centromeres.

PCA

Principal component analysis was performed with LASER⁴⁷ following the approach described previously⁹. After generating a reference space of modern Western Eurasian individuals¹⁰, we projected the BAM files of ancient reference individuals (see Supplementary Table S1 for references) and the British individuals presented here into the reference space via Procrustes analysis implemented in LASER.

f-statistics

The f -statistics presented here, *i.e.* outgroup f_3 , f_4 , $qpAdm$, and $qpGraph$ were computed with $qpPop$, $qpDstat$ in f_4 mode, $qpAdm$, and $qpGraph$ from the ADMIXTOOLS24 package with default parameters on the positions defined by the *HOIII* set of SNPs7. Ancient individuals analysed here are listed in Supplementary Table S1, including the explanation of all population labels used (WHG, SHG, etc.). Modern reference individuals were first published in Mallick *et al.*23. All $qpAdm$ runs used the set of outgroups Han, Karitiana, Mbuti, Onge, Papuan, Mota, Ust'-Ishim, MA1, El Mirón, GoyetQ116-1.

ALDER

We used ALDER27 to estimate the dates of admixture between WHG and ANF. All combinations we tested are listed in Supplementary Table S3, which consisted of the pairs or groups of individuals specified in the first column and WHG and ANF (individuals constituting WHG and ANF are given in Supplementary Table S1).

Chronological Modelling

We used OxCal 4.328 to produce chronological models of the arrival and spread of ANF ancestry into Britain (Supplementary Materials Section 6). We only used Early Neolithic (4000-3500 BC) radiocarbon dates that had been obtained from material or individuals where there was palaeogenetic data indicating ANF ancestry. We divided these samples into five regional populations: Western Britain, Central England, Eastern England, Western Scotland and the Orkney Isles. Dates associated with each region were grouped as Phases (Supplementary Figure S19). We used the Boundary function to produce probability distributions for the arrival of ANF ancestry in Britain as a whole, and for each region. We used the Difference function to produce probability distributions for the time between the death of the latest individual with wholly WHG ancestry and the arrival of populations with ANF ancestry, as well as between the arrival of ANF ancestry in Britain as a whole and the different regions of Britain.

Haplotype-based analyses

We used CHROMOPAINTER25 to summarize DNA patterns in our ancient individuals, including other high coverage publicly available ancient genomes from relevant cultures and time periods, to infer the proportion of DNA for which ancient individuals most closely matched to individuals from a global panel of modern donor groups48–50. This panel included many population samples from across West Eurasia, as well as 35 labelled groups from within the British Isles. We generated matching profiles when considering SNPs independently (allele sharing) and also when considering the correlations between neighbouring SNPs (haplotype sharing). To do so we first merged high quality diploid calls for our selected high coverage ancient genomes and jointly phased the resultant dataset of 159,287 SNPs using SHAPEITv251. We performed additional mixture modelling on our generated allele and haplotype sharing profiles implemented in SOURCEFIND26 to form target groups as mixtures of the DNA sharing profiles of other included groups. We performed two sets of analyses: i) using all modern groups (or a subset of) to model the ancestry of ancient individuals, and ii) using different sets of ancient individuals, plus the

modern Yoruba and Han, to model the ancestry of modern world-wide groups. Further details are provided in Supplementary Materials Section 7.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We are grateful to the five anonymous reviewers whose constructive comments helped to improve this article considerably. The authors would like to thank the Longleat Estate, Tom Lord at Lower Winskill Farm, Barry Chandler at Torquay Museum, Andrew Chamberlain at the University of Manchester, Linda Wilson and Graham Mullan at the University of Bristol Spelaeological Society, Elizabeth Walker, Adam Gwilt and Jody Deacon at the National Museum of Wales, Andy Maxted at Brighton Museum, Marta Lahr at the Duckworth Laboratory, Barry Lane at Wells Museum, Martin Smith at Bournemouth University, David Rice at the Museum of Gloucester and Rob Kruszynski at the Natural History Museum for providing access to samples. In addition, Y.D. wishes to thank Jens Blöcher, Amelie Scheu, Christian Sell and Joachim Burger for discussions on the bioinformatic pipeline, and Vivian Link for help with ATLAS. M.G.T. and I.B. were supported by a Wellcome Trust Investigator Award (project 100713/Z/12/Z). L.v.D acknowledges financial support from the Newton Trust grant MR/P007597/1. R.M. was supported by an EMBO Long-Term Fellowship (ALTF 133-2017). D.R. was supported by NIH grant GM100233, by NSF HOMINID BCS-1032255, and by an Allen Discovery Center of the Paul Allen Foundation, and is a Howard Hughes Medical Institute investigator. C.S. is supported by the Calleva Foundation and the Human Origins Research Fund. S.W. was supported by the U.S National Institute of Justice grant 2014-DN-BX-K031.

References

1. Collard M, Edinborough K, Shennan S, Thomas MG. Radiocarbon evidence indicates that migrants introduced farming to Britain. *J Arch Sci.* 2010; 37:866–870.
2. Sheridan, JA. The Neolithisation of Britain and Ireland. *Landscapes in Transition.* Finlayson, B, Warren, G, editors. Oxbow; Oxford: 2010. 89–105.
3. Thomas, J. *The Birth of Neolithic Britain: An interpretive account.* Oxford University Press; Oxford: 2013.
4. Skoglund P, et al. Genomic Diversity and Admixture Differs for Stone-Age Scandinavian Foragers and Farmers. *Science.* 2014; 344:747–750. [PubMed: 24762536]
5. Gamba C, et al. Genome flux and stasis in a five millennium transect of European prehistory. *Nat Commun.* 2014; 5
6. Cassidy LM, et al. Neolithic and Bronze Age migration to Ireland and establishment of the insular Atlantic genome. *Proc Natl Acad Sci USA.* 2015; 113:368–373. [PubMed: 26712024]
7. Haak W, et al. Massive migration from the steppe was a source for Indo-European languages in Europe. *Nature.* 2015; 522:207–211. [PubMed: 25731166]
8. Broushaki F, et al. Early Neolithic genomes from the eastern Fertile Crescent. *Science.* 2016; 353:499–503. [PubMed: 27417496]
9. Hofmanová Z, et al. Early farmers from across Europe directly descended from Neolithic Aegeans. *Proc Natl Acad Sci USA.* 2016; 113:6886–6891. [PubMed: 27274049]
10. Lazaridis I, et al. Genomic insights into the origin of farming in the ancient Near East. *Nature.* 2016; 536:419–424. [PubMed: 27459054]
11. Olalde I, et al. A Common Genetic Origin for Early Farmers from Mediterranean Cardial and Central European LBK Cultures. *Mol Biol Evol.* 2015; 32:3132–3142. [PubMed: 26337550]
12. Olalde I, et al. The Beaker Phenomenon And The Genomic Transformation Of Northwest Europe. *Nature.* 2018; 555:190–196. [PubMed: 29466337]
13. González-Forbes G, et al. Paleogenomic Evidence for Multi-generational Mixing between Neolithic Farmers and Mesolithic Hunter-Gatherers in the Lower Danube Basin. *Curr Biol.* 2017; 27:1801–1810. [PubMed: 28552360]
14. Lipson M, et al. Parallel palaeogenomic transects reveal complex genetic history of early European farmers. *Nature.* 2017; 551:368–372. [PubMed: 29144465]

15. Mathieson I, et al. The genomic history of southeastern Europe. *Nature*. 2018; 555(7695):197–203. [PubMed: 29466330]
16. Günther T, et al. Genomics of Mesolithic Scandinavia reveal colonization routes and high-latitude adaptation. *PLoS Biol*. 2018; 16:e2003703. [PubMed: 29315301]
17. Fu Q, et al. The genetic history of Ice Age Europe. *Nature*. 2016; 534:200–205. [PubMed: 27135931]
18. Olalde I, et al. Derived immune and ancestral pigmentation alleles in a 7,000-year-old Mesolithic European. *Nature*. 2014; 507:225–228. [PubMed: 24463515]
19. Mathieson I, et al. Genome-wide patterns of selection in 230 ancient Eurasians. *Nature*. 2015; 528:499–503. [PubMed: 26595274]
20. Kiliç GM, et al. The Demographic Development of the First Farmers in Anatolia. *Curr Biol*. 2016; 26:2659–2666. [PubMed: 27498567]
21. Jones ER, et al. The Neolithic Transition in the Baltic Was Not Driven by Admixture with Early European Farmers. *Curr Biol*. 2017; 27:576–582. [PubMed: 28162894]
22. Lazaridis I, et al. Ancient human genomes suggest three ancestral populations for present-day Europeans. *Nature*. 2014; 513:409–413. [PubMed: 25230663]
23. Mallick S, et al. The Simons Genome Diversity Project: 300 genomes from 142 diverse populations. *Nature*. 2016; 538:201–206. [PubMed: 27654912]
24. Patterson N, et al. Ancient Admixture in Human History. *Genetics*. 2012; 192:1065–1093. [PubMed: 22960212]
25. Lawson DJ, et al. Inference of population structure using dense haplotype data. *PLoS Genet*. 2012; 8(1):e1002453. [PubMed: 22291602]
26. Chacon-Duque JC, et al. Latin Americans show wide-spread Converso ancestry and the imprint of local Native ancestry on physical appearance. *Nature Comms*. 2018; 9
27. Loh P-R, Lipson M, Patterson N, et al. Inferring admixture histories of human populations using linkage disequilibrium. *Genetics*. 2013; 193:1233–1254. [PubMed: 23410830]
28. Bronk Ramsey C. Bayesian analysis of radiocarbon dates. *Radiocarbon*. 2009; 51(1):337–360.
29. Chaitanya L, et al. The HRISplex-S system for eye, hair and skin colour prediction from DNA: Introduction and forensic developmental validation. *Forensic Sci Int Genet*. 2018; 35:123–135. [PubMed: 29753263]
30. Whittle, AWR, Healy, F, Bayliss, A, Allen, MJ. *Gathering time: dating the early Neolithic enclosures of southern Britain and Ireland*. Oxford: Oxbow Books; 2011.
31. Scarre C. *The Early Neolithic of western France and megalithic origins in Atlantic Europe*. Oxford Journal of Archaeology. 1992; 11(2):121–154.
32. Bollongino R, et al. 2000 Years of Parallel Societies in Stone Age Central Europe. *Science*. 2013; 342:479–481. [PubMed: 24114781]
33. Fraser M, et al. New insights on cultural dualism and population structure in the Middle Neolithic Funnel Beaker culture on the island of Gotland. *J Arch Sci Rep*. 2018; 17:325–334.
34. Charlton S, et al. Finding Britain's last hunter-gatherers: A new biomolecular approach to 'unidentifiable' bone fragments utilising bone collagen. *J Arch Sci*. 2016; 73:55–61.
35. Schulting, RJ, Bori, D. A tale of two processes of Neolithisation: southeast Europe and Britain/Ireland. *Neolithic Europe: essays in honour of Professor Alasdair Whittle*. Bickle, P, Cummings, V, Hofmann, D, Pollard, J, editors. Oxford: Oxbow Books; 2017. 82–104.
36. Dabney J, et al. Complete mitochondrial genome sequence of a Middle Pleistocene cave bear reconstructed from ultrashort DNA fragments. *Proc Natl Acad Sci USA*. 2013; 110:15758–15763. [PubMed: 24019490]
37. Rohland N, Harney E, Mallick S, Nordenfelt S, Reich D. Partial uracil-DNA-glycosylase treatment for screening of ancient DNA. *Philos Trans Royal Soc B*. 2014; 370
38. Meyer M, Kircher M. *Illumina Sequencing Library Preparation for Highly Multiplexed Target Capture and Sequencing*. Cold Spring Harb Protoc. 2010
39. Kircher M, Sawyer S, Meyer M. Double indexing overcomes inaccuracies in multiplex sequencing on the Illumina platform. *Nucleic Acids Res*. 2011; 40

40. Schubert M, Lindgreen S, Orlando L. Adapter Removal v2: rapid adapter trimming, identification, and read merging. *BMC Res Notes*. 2016; 9
41. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*. 2009; 25:1754–1760. [PubMed: 19451168]
42. Mckenna A, et al. The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res*. 2010; 20:1297–1303. [PubMed: 20644199]
43. Link V, et al. ATLAS: Analysis Tools for Low-depth and Ancient Samples. *Biorxiv*. 2017; doi: 10.1101/105346
44. Fu Q, et al. A Revised Timescale for Human Evolution Based on Ancient Mitochondrial Genomes. *Curr Biol*. 2013; 23:553–559. [PubMed: 23523248]
45. Navarro-Gomez D, et al. Phy-Mer: a novel alignment-free and reference-independent mitochondrial haplogroup classifier. *Bioinformatics*. 2014; 31:1310–1312. [PubMed: 25505086]
46. Ralf A, Montiel González D, Zhong K, Kayser M. Yleaf: software for human Y-chromosomal haplogroup inference from next generation sequencing data. *Mol Biol Evol*. 2018; 35:1291–1294. [PubMed: 29518227]
47. Wang C, Zhan X, Liang L, Abecasis GR, Lin X. Improved Ancestry Estimation for both Genotyping and Sequencing Data using Projection Procrustes Analysis and Genotype Imputation. *Am J Hum Genet*. 2015; 96:926–937. [PubMed: 26027497]
48. Hellenthal G, et al. A genetic atlas of human admixture history. *Science*. 2014; 343(6172):747–751. [PubMed: 24531965]
49. Busby GB, et al. The role of recent admixture in forming the contemporary West Eurasian genomic landscape. *Current Biology*. 2015; 25(19):2518–2526. [PubMed: 26387712]
50. Leslie S, et al. The fine-scale genetic structure of the British population. *Nature*. 2015; 519(7543):309. [PubMed: 25788095]
51. Delaneau O, et al. A linear complexity phasing method for thousands of genomes. *Nat Methods*. 2012; 9(2):179–81.

Figure 1. Map of sample locations.

Geographical locations of British samples analysed here. Numbers indicate the number of samples from a given location.

Figure 2. PCA of modern and ancient West-Eurians.

British and additional ancient samples are projected onto the reference space computed on present-day West-Eurasian populations. See Material and Methods for computational details and Supplementary Table 1 for information on the samples.

Abbreviations: European (Eur.), Pleistocene (Plei.), hunter-gatherer (H.-G.), British Isle (Brit.-I.), Middle Neolithic (M.-Neo.)

Figure 3. WHG and ANF ancestry components of British and Central European Neolithic populations.

The relative WHG and ANF ancestry in Early and Middle Neolithic British and continental European populations quantified by *qpAdm*. Percentages indicate error estimates computed by block jackknifing with a block size of 5 cM²⁴. See Material and Methods for computational details and Supplementary Table 1 for the lists of samples grouped into WHG and the different Neolithic populations.

Abbreviations: Neolithic (Neo.), Early-, Middle-, and Late Neolithic (EN, MN, LN), South East (SouthE), South West (SouthW)

Figure 4. Affinities of British and continental Neolithic populations.

(A) We compute f_4 symmetry testing statistics for different British EN, MN, and LN and continental MN populations comparing shared drift with Central European EN and Iberian EN populations. A positive Z-score above 2 corresponds to a significant affinity to the Iberian EN over Central European EN population. **(B)** *Quantifying excess WHG ancestry in British EN compared to the Iberian EN population.* We compute *qpAdm* estimates of WHG and Anatolian and Iberian ANF populations in EN samples from Wales, England, and Scotland. See Supplementary Table 1 for the lists of samples grouped into WHG and the

different Neolithic populations. Percentages and bars indicate error estimates computed by block jackknifing with a block size of 5 cM²⁴.

Figure 5. Patterns of haplotype sharing across high coverage aDNA samples.

(A) Hierarchical clustering of total-variation-distance (TVD) between CHROMOPAINTER inferred haplotype sharing profiles of seven high coverage Neolithic individuals when compared to a global modern reference panel. (B) Inferred ancestry proportions (SOURCEFIND inferred mixing coefficients) of high coverage ancient genomes, coloured as per the legend and outer pie ring colour, relative to a panel of ancient genomes, plus modern Yoruba and Han (as given in the legend at top). Raw proportions and standard errors are provided in Supplementary Table S8. (C) Inferred ancestry proportions of five high coverage Neolithic individuals (triangles coloured as in (B)) relative to a global modern reference panel. The size of the blue circle provides the majority inferred contributions, as given in the scale at bottom right, with all possible modern contributors shown with a black dot. Raw proportions and standard errors are provided in Supplementary Table S9.