Published in final edited form as:

Urology. 2019 January ; 123: 280–286. doi:10.1016/j.urology.2018.04.045.

Complications of Delivery Among Mothers with Spina Bifida

Courtney L. Shepard, MD, MS¹, Phyllis L. Yan, MA², Stephanie J. Kielb, MD³, Daniela A. Wittmann, PhD, LMSW⁴, Elisabeth H. Quint, MD⁵, Kate H. Kraft, MD⁶, and John M. Hollingsworth, MD, MS²

¹Divisions of Pediatric Urology and Health Services Research, Department of Urology and University of Michigan Institute for Healthcare Policy and Innovation, Ann Arbor

²Divisions of Endourology and Health Services Research, Department of Urology and University of Michigan Institute for Health Policy and Innovation, Ann Arbor

³Division of Female Pelvic Medicine and Reconstructive Surgery, Department of Urology, Northwestern Medicine, Chicago

⁴Division of Health Services Research, Department of Urology, University of Michigan, Ann Arbor

⁵Division of Pediatric and Adolescent Gynecology, Department of Obstetrics and Gynecology, University of Michigan, Ann Arbor

⁶Divisions of Pediatric Urology and Endourology, Department of Urology, University of Michigan, Ann Arbor

Abstract

Objective: To determine rates and types of peripartum morbidity among delivering women with spina bifida (SB) compared to those without SB. The rates of pregnancy and delivery among women with SB have been significantly increasing. Current knowledge of peripartum outcomes for these women is limited.

Methods: Using 2004-2013 National Inpatient Sample data, we identified all hospitalizations for delivery, distinguishing between women with and without SB. Using a code-based algorithm, we determined whether a complication occurred during the hospitalization. We then fit a series of multivariable logistic models to examine for associations between a complication occurrence during vaginal or cesarean delivery and a woman's SB status.

Results: We identified 38,319,814 weighted admissions for delivery, 9,516 of which were made by women with SB. Women with SB had a significantly higher rate of cesarean delivery than women without this diagnosis (53% v 32%, $p<0.001$). The 46.7% of women with SB who delivered vaginally did not have significantly increased odds of a complication associated with their delivery compared to women without SB [odds ratio (OR) 1.15, 95% confidence interval (CI) 0.99 to 1.34, $p=0.066$]. However, women with SB who underwent a cesarean delivery did have higher odds of morbidity compared to those without (OR 1.49, 95% CI 1.25 to 1.78, $p<0.001$).

Corresponding Author: Courtney L. Shepard, University of Michigan Health System, Divisions of Pediatric Urology and Dow Urologic Health Services Research, Department of Urology, 1500 E. Medical Center Drive, F3055 Taubman Center, Ann Arbor, MI 48109, coshepar@med.umich.edu, 734-615-3038 (Phone), 734-615-3520 (Fax).

Disclosures/Conflicts of Interest: The authors report no conflicts of interest.

Common complications included preterm delivery, urinary tract infection, hematologic event, and blood transfusion.

Conclusions: Compared to women without SB, those with SB deliver more frequently by cesarean section and have higher odds of morbidity associated with cesarean delivery, but not vaginal delivery.

Keywords

myelomeningocele; spina bifida; spina bifida occulta; pregnancy; delivery; complications

Introduction

Over the last several decades, refinements in medical and surgical management have greatly reduced childhood mortality from spina bifida, allowing most to survive well into adulthood.¹ Consequently, adult patients with SB, and the providers who care for them, now face entirely new health challenges related to their disability that remain poorly understood. There is perhaps no single area more under-investigated for this population than the reproductive health of women with SB.

Despite the fact that the number of women with SB giving birth has increased by 56% over the last 10 years,² there are a paucity of studies on peripartum outcomes. Reports are limited to a few small case series, the largest of which describes 69 deliveries in 38 women with SB. These studies suggest a higher rate of complications associated with delivery as would be expected owing to their atypical skeletal anatomy, medial comorbidities, and prior surgical procedures (e.g., urinary tract reconstruction, placement of cerebrospinal fluid shunt). However, without larger scale studies, the risk of various complications remains unknown, hampering preconception counseling and delivery planning.

In this context, we conducted an observational study using national discharge data. During hospitalizations for delivery, we determined the modes of delivery as well as rates and types of peripartum morbidity among women with SB compared to those without this diagnosis. Findings from this study will allow urologists and other health care providers who care for women with SB to facilitate informed discussions about their potential risks of delivery.

Materials and Methods

Data source and study population

For our study, we used data from the National Inpatient Sample (NIS). Developed by the Healthcare Cost and Utilization Project, the NIS is an all-payer database that captures a 20% sample of admissions from participating hospitals. Weighted estimates from the NIS are considered generalizable to all discharges in the United States.³ Using appropriate *International Classification of Diseases, Ninth Revision (ICD-9)*, *Clinical Modification* diagnosis codes and Clinical Classification Software (CCS) groups (Supplementary Table 1), we identified all women admitted to an acute care hospital following an obstetric delivery between 2004 and 2013. Through a previously described algorithm,² we distinguished between admissions for women with and without SB. Women with SB were further

categorized by severity of SB (SB occulta, SB without hydrocephalus, SB with hydrocephalus). We then used CCS groups (Supplementary Table 1) to determine whether these women underwent vaginal or cesarean delivery.

Outcomes

We measured two binary outcomes: 1) the occurrence of an immediate peripartum complication around a vaginal delivery and 2) the occurrence of an immediate peripartum complication around a cesarean delivery. We defined the peripartum period as the time between admission and discharge. Based on author consensus, complications were abstracted using definitions from the Centers for Disease Control and Prevention,⁴ the American Congress of Obstetricians and Gynecologists,⁵ and the American College of Surgeons.⁶ We also identified complications specific to the patients with SB, including malfunction or replacement of a ventriculoperitoneal shunt and common complications of abdominal surgery after bladder augmentation such as cystotomy and enterotomy (Supplementary Table 2).

Statistical analysis

For our initial analytic step, we tested for differences between women with and without SB, who were admitted for delivery. We made comparisons over a variety of sociodemographic characteristics including age on admission, primary payer, and urban-rural residence, level of comorbidity (assessed by the Elixhauser methodology),⁷ and specific comorbidities (Supplementary Table 3). Next, we compared crude rates of peripartum complication between women with and without SB stratified by type of delivery. Additionally, we performed temporal trends analysis to determine if the utilization of cesarean delivery changed over this time period.

We fit two multivariable logistic regression models to determine if a woman's SB status and severity were independently associated with the occurrence of a peripartum complication among women who underwent a vaginal delivery, as well as among women who underwent a cesarean delivery. We adjusted for the patient factors described above.

We performed all analyses using SAS® version 9.4 (Cary, NC). All of our tests were two sided with the probability of type I error set at 0.05. Our institutional review board deemed our study to be exempt from its oversight.

Results

We identified 38,310,298 hospitalizations for delivery among women without SB and 9,516 among women with SB (33.7% SB occulta, 47.5% SB without hydrocephalus, 18.8% SB with hydrocephalus). Among women without SB, 67.7% underwent a vaginal delivery, and 32.3% underwent a cesarean delivery. Among women with SB, 46.7% underwent a vaginal delivery, and 53.3% underwent a cesarean delivery ($p<0.001$ for the difference in distribution between these groups of women). The frequency of cesarean delivery increased with increasing severity of SB (42.6% of all deliveries for women with SB occulta, 56.1% for SB without hydrocephalus, and 65.2% for SB with hydrocephalus). The percentage of women with SB who delivered by cesarean section over the last ten years has not

significantly changed ($p=0.47$ for trend, Supplementary Figure 1). Women with SB were younger and were more likely to have private insurance or Medicare, live outside of a city, and deliver at an urban teaching hospital (Table 1). Women with SB also had higher levels of comorbid illness.

When comparing overall frequencies of specific immediate peripartum outcomes, women with SB were more likely to have several adverse fetal and maternal outcomes compared to women without SB (Table 2). They were more likely to have preterm delivery defined as delivery before 37 weeks gestation (11.3% versus 7.3%, $p<0.001$). Women with SB had a nearly six-fold increased incidence of urinary tract infections (4.6% versus 0.8%, $p<0.001$). In addition, they were more likely to suffer a hematologic event (4.0% versus 2.0%, $p<0.001$) and they were also more likely to undergo blood transfusion (2.3% versus 1.0%, $p<0.001$). Women with SB were also significantly more likely to have neurologic, pulmonary, renal, and cardiac complications, although the frequencies were very low.

The frequency of immediate peripartum adverse outcomes differed by delivery type. For vaginal delivery, women with SB were more likely to have a renal complication or urinary tract infection than women without SB (Table 3). Although they were more likely to sustain trauma to the perineum or vulva (56.3% versus 52.5%, $p=0.021$), they were less likely to require forceps- or vacuum-assisted delivery (0.6% versus 1.5%, $p=0.039$). They were more likely to have a hematologic complication (2.3% versus 1.2%, $p=0.002$), but were no more likely to require a blood transfusion (0.6% versus 0.5%, $p=0.65$). The remainder of their outcomes were similar to women without SB.

With respect to cesarean delivery, women with SB were more likely to suffer from an intra-operative injury. Hematologic complications were more frequent among SB women (5.5% versus 3.7%, $p<0.001$) as were blood transfusions (3.7% versus 1.9%, $p<0.001$). Women with SB were more likely to sustain neurologic, pulmonary, shunt-related, and infectious complications, but the frequency of renal complications among them did not differ from that of women without SB (Table 3).

After adjusting for measured patient and hospital factors, for vaginal delivery, a woman's risk of an immediate peripartum complication was not associated with her SB status [odds ratio (OR), 1.15; 95% confidence interval (CI), 0.99 to 1.34; $p=0.066$] (Figure 1). However, for cesarean delivery, a woman's odds of suffering an immediate peripartum complication were nearly 50% higher if she had SB (OR, 1.49; 95% CI, 1.25 to 1.78; $p<0.001$). When evaluating the odds of complication by SB severity, the increased odds of complication during cesarean delivery was driven by the largest group of women, SB women without hydrocephalus (OR, 1.71; 95% CI, 1.35 to 2.16; $p=0.016$) (Supplementary Figure 2).

Comment

This is the first large-scale study describing obstetric outcomes among mothers with spina bifida, which allows for a broader understanding than the current small case series. We demonstrated that despite the fact that most women live outside a city, most delivered at an urban teaching hospital. This indicates that many women with SB must travel for their care.

There was a significantly higher incidence of cesarean delivery among women with SB compared to the general population, with utilization increasing with increasing severity of SB. The utilization of cesarean delivery has been stable over time. Immediate peripartum complications occur with significantly greater frequency in this population than among women without SB after cesarean section, but not vaginal delivery. These findings serve to inform SB providers when counseling women who are of child-bearing age.

Certain trends have been reported in prior case series which are consistent with our findings. Specifically, the higher incidence of preterm delivery has been noted, although at higher rates than seen in this study.^{8,9} This could be caused, in part, by the inability to sense contractions in patients with higher level lesions or by the increased incidence of urinary tract infection among women with SB during pregnancy, which is associated with preterm labor.⁸ Their increased risk of urinary tract infections is a well-documented phenomenon that may be explained by these patients' underlying predisposition as well as changes during pregnancy causing increased urinary stasis and difficulties with urinary catheterizations.⁹⁻¹² The reported rates of renal complication such as acute renal injury vary widely and likely relates to baseline renal function prior to surgery.¹¹⁻¹⁵ The significantly higher odds of neurologic and pulmonary complications that we observed are consistent with other reports of an increased incidence of seizure episodes, shunt malfunctions, respiratory compromise, and pneumonia after delivery.¹⁶

There are limitations of our study that merit discussion. First, we must acknowledge the risk of selection bias. In our observational study, women with SB who undergo vaginal delivery may differ in unobserved ways from those who had cesarean deliveries, such as differences in level of lesion. The difference in disability based on level of lesion may moderate a woman's odds of complication. The reliability of using ICD9 codes to determine the severity of SB is unclear and providers less familiar with the nuances of SB may have entered these codes. The higher rates of cesarean delivery and complications with delivery among women with SB occulta was unexpected given patients with this diagnosis are usually asymptomatic. These patients may represent a particular subset of women with SB occulta who are particularly symptomatic, or alternatively some of these women may have a neural tube defect but were assigned an incorrect ICD-9 code. Additionally, we are unable to distinguish between planned versus labored cesarean deliveries. The NIS database does not allow us to follow patients before and after discharge, so complications that occurred during pregnancy and in the later post-partum period are unknown. We also cannot determine which patients have previously undergone other surgeries such as bladder reconstruction. Finally, the outcomes of the neonates are unknown, which is an important part of prenatal counseling. This is an area that needs further study.

Despite these limitations, our study has several important clinical implications. It is currently recommended that women with SB undergo vaginal delivery unless there are obstetrical contraindications, or the patient has undergone a reconstructive urethral continence surgery.^{11,12,15} The findings of this study support this recommendation with a cesarean section done for obstetrical or urological reasons only. However, our findings do not imply that a cesarean delivery should not be performed when indicated as this will be the safest type of delivery for some women. Other studies have reported similar rates of cesarean delivery, most

commonly for previous cesarean delivery, pelvic contraction, breech presentation, premature delivery, and preeclampsia.⁸ Greenwell et al reported on 28 deliveries among women who underwent lower urinary tract reconstruction for various congenital anomalies including SB. They found that 43% of deliveries were planned cesarean deliveries and 23% were emergent, most commonly for premature labor and preeclampsia.¹⁴ There is general consensus that a urologist should be present or immediately available for a women with SB who requires a cesarean delivery if she has undergone bladder reconstruction or has other urinary tract anomalies to help decrease the risk of intra-operative injury.^{11,12}

Our role in discussing reproductive function with these women cannot be limited to the perinatal period. It is concerning that 26% of the deliveries in this study occurred by age 22 for women with SB as 59% to 76% of pregnancies in women this age are unintended.¹⁷ While this rate is no different than for women without SB, providers may not be talking about sex with their adolescent and young adult SB patients; indeed, as few as 5% of young women with SB have talked to a physician about this topic.^{18,19} This is especially worrisome as many women with physical disabilities assume sexual education they get in school does not apply to them, yet most adolescent and young adult women with physical disabilities are just as sexually active as their able-bodied counterparts.^{20,21} These women deserve to understand their sexual and reproductive health and have access to family planning options and gynecologic care. As lifelong genitourinary providers who are especially invested in these patients, urologists are best positioned to initiate discussions about sexual and reproductive health with these women and indeed have taken responsibility for starting these conversations.²²⁻²⁴ With intentional collaboration with obstetricians and primary care providers, we can help these women make informed decisions. Additionally, these women should be screened for safety in their relationships as women with physical disabilities are four times more likely to be victims of sexual abuse and are more likely to experience intimate partner violence than able-bodied women.^{25,26}

For women who are pregnant or considering pregnancy, they should be informed while vaginal delivery is safe and possible for women without contraindications, most women with SB undergo cesarean section. The overall risk of complications during delivery is similar to women without SB, but slightly higher during a cesarean section than for women without this diagnosis. With the increased risk of preterm delivery, they should be informed of signs and symptoms that could indicate contractions in women with abnormal abdominal sensation. Due to the high prevalence of urinary tract infections, they should be closely monitored by their urologists and obstetricians to help prevent the infections and ensure the enlarging uterus does not inhibit adequate bladder drainage. There may be a need for prophylactic antibiotics to prevent urinary tract infections and alternative catheterization schedules or approaches during pregnancy.^{8,12,16} Patients should be informed of their increased likelihood of hematologic events and possible need for a blood transfusion. Finally, patients should work with their SB-specific sub-specialists to optimize their medical status before and during pregnancy. Active collaboration between specialists and obstetricians will help to manage and possibly prevent complications associated with delivery.

Conclusions

While delivery among women with SB is generally safe, these women do have a slightly increased risk of peripartum complications, particularly after cesarean section. These are likely related to their underlying comorbidities. A team approach including urologists, obstetricians, and other appropriate subspecialists will be needed to help women with SB make informed decisions and to optimize their obstetrical outcomes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements:

The authors would like to thank Lindsey Admon, MD, MS (National Clinician Scholars Program, University of Michigan Institute for Healthcare Policy and Innovation and the US Department of Veteran Affairs, Ann Arbor) for her contribution to designing this research study and editing the manuscript.

Source of Funding: This research was supported by a T-32 grant from the National Institutes of Health's National Institute of Diabetes and Digestive and Kidney Diseases (grant number 5T32DK007782-17).

References

1. Shin M, Kucik JE, Siffel C, et al. Improved survival among children with spina bifida in the United States. *The Journal of pediatrics*. 12 2012;161(6):1132–1137. [PubMed: 22727874]
2. Shepard CL YP, Hollingsworth JM, Kraft KH. Pregnancy among mothers with spina bifida. *Journal of Pediatric Urology*. 2017.
3. Quality AfHRA. HCUP Databases. Healthcare Cost and Utilization Project (HCUP). Overview of the National (Nationwide) Inpatient Sample (NIS). 2016; <https://www.hcup-us.ahrq.gov/nisoverview.jsp> Accessed 02/13/2017, 2017.
4. Prevention CfDCA. Severe Maternal Morbidity in the United States. 5/22/2017.
5. Kilpatrick SK, Ecker JL. Severe maternal morbidity: screening and review. *American journal of obstetrics and gynecology*. 9 2016;215(3):B17–22. [PubMed: 27560600]
6. Program ACoSNQI. User Guide for the 2014 ACS NSQIP Participant Use Data File. 10 2015.
7. Elixhauser A, Steiner C, Harris DR, Coffey RM. Comorbidity measures for use with administrative data. *Medical care*. 1 1998;36(1):8–27. [PubMed: 9431328]
8. Sterling L, Keunen J, Wigdor E, Sermer M, Maxwell C. Pregnancy outcomes in women with spinal cord lesions. *Journal of obstetrics and gynaecology Canada : JOGC = Journal d'obstetrique et gynecologie du Canada : JOGC*. 1 2013;35(1):39–43.
9. Arata M, Grover S, Dunne K, Bryan D. Pregnancy outcome and complications in women with spina bifida. *The Journal of reproductive medicine*. 9 2000;45(9):743–748. [PubMed: 11027084]
10. Richmond D, Zaharievska I, Bond A. Management of pregnancy in mothers with spina bifida. *European journal of obstetrics, gynecology, and reproductive biology*. 8 1987;25(4):341–345.
11. Vordermark JS, Deshon GE, Agee RE. Management of pregnancy after major urinary tract reconstruction. *Obstetrics and gynecology*. 3 1990;75(3 Pt 2):564–567. [PubMed: 2406671]
12. Thomas JC, Adams MC. Female sexual function and pregnancy after genitourinary reconstruction. *The Journal of urology*. 12 2009;182(6):2578–2584. [PubMed: 19836802]
13. Creagh TA, McInerney PD, Thomas PJ, Mundy AR. Pregnancy after lower urinary tract reconstruction in women. *The Journal of urology*. 10 1995;154(4):1323–1324. [PubMed: 7658529]
14. Greenwell TJ, Venn SN, Creighton S, Leaver RB, Woodhouse CR. Pregnancy after lower urinary tract reconstruction for congenital abnormalities. *BJU Int*. 11 2003;92(7):773–777. [PubMed: 14616465]

15. Rietberg CC, Lindhout D. Adult patients with spina bifida cystica: genetic counselling, pregnancy and delivery. *European journal of obstetrics, gynecology, and reproductive biology*. 11 1993;52(1): 63–70.
16. Jackson AB, Mott PK. Reproductive health care for women with spina bifida. *TheScientificWorldJournal*. 11 26 2007;7:1875–1883.
17. Finer LB, Zolna MR. Declines in Unintended Pregnancy in the United States, 2008–2011. *The New England journal of medicine*. 3 03 2016;374(9):843–852. [PubMed: 26962904]
18. Lee NG, Andrews E, Rosoklija I, et al. The effect of spinal cord level on sexual function in the spina bifida population. *Journal of pediatric urology*. 6 2015;11(3):142.e141–146. [PubMed: 25864616]
19. Gatti C, Del Rossi C, Ferrari A, Casolari E, Casadio G, Scire G. Predictors of successful sexual partnering of adults with spina bifida. *The Journal of urology*. 10 2009;182(4 Suppl):1911–1916. [PubMed: 19695634]
20. Nosek MA HC, Rintala DH, Young ME, Chanpong GF. National study of women with physical disabilities: final report. *Sexuality and disability*. 2001;19(1):5–39.
21. Kahn NF, Halpern CT. Experiences of Vaginal, Oral, and Anal Sex From Adolescence to Early Adulthood in Populations With Physical Disabilities. *The Journal of adolescent health : official publication of the Society for Adolescent Medicine*. 10 19 2017.
22. Woodhouse CR, Lipshultz L, Hwang K, Mouriquand P, Creighton S. Adult care of children from pediatric urology: part 2. *The Journal of urology*. 9 2012;188(3):717–723. [PubMed: 22818132]
23. Joseph DB. Current Urologic Management of Spina Bifida. *AUA Update Series*. 2015 2015;34:247–255.
24. Tanaka S Transition from Pediatric to Adult Health Care in Urology. *AUA Update Series*. 2015;34:257–263.
25. Casteel C, Martin SL, Smith JB, Gurka KK, Kupper LL. National study of physical and sexual assault among women with disabilities. *Injury prevention : journal of the International Society for Child and Adolescent Injury Prevention*. 4 2008;14(2):87–90. [PubMed: 18388227]
26. Barrett KA, O'Day B, Roche A, Carlson BL. Intimate partner violence, health status, and health care access among women with disabilities. *Women's health issues : official publication of the Jacobs Institute of Women's Health*. Mar-Apr 2009;19(2):94–100.

Figure 1.
Odds of complication for women with spina bifida associated with vaginal compared to women without spina bifida.

Table 1.

Personal and hospital characteristics of women hospitalized for a delivery.

Characteristic	No Spina Bifida	Spina Bifida	P-value
Sociodemographic Factors			
Age			0.0043
<18	2042638 (5.3)	440 (4.6)	
18-22	7371181 (19.2)	2021 (21.2)	
22-26	7140991 (18.6)	1891 (19.9)	
26-30	8877900 (23.2)	2280 (24.0)	
30-34	7268459 (19.0)	1737 (18.3)	
>34	5609129 (14.6)	1147 (12.1)	
Primary Payer			<0.001
Self-pay	1128307 (2.9)	138 (1.4)	
Private	19561812 (51.1)	5038 (52.9)	
Medicare	232260 (0.6)	373 (3.9)	
Medicaid	16291052 (42.5)	3635 (38.2)	
No Charge/Other	1096866 (2.9)	334 (3.5)	
Residence			<0.001
Central/Fringe (1 million or greater)	21894088 (57.1)	4971 (52.2)	
50,000-999,999	10786337 (28.2)	2914 (30.6)	
10,000-49,999	3515184 (9.2)	1043 (11.0)	
<10,000	2114689 (5.5)	589 (6.2)	
Hospital Location and Teaching Status			<0.001
Rural	4291728 (11.2)	1014 (10.7)	
Urban non-teaching	16257311 (42.4)	3460 (36.4)	
Urban teaching	17761258 (46.4)	5043 (53.0)	
Patient Factors			
Previous cesarean section	6043535 (15.8)	1732 (18.2)	0.0035
Mean Number of Elixhauser Conditions	0.34	0.67	<0.001
Comorbid Conditions Directly Related to SB			
Chronic Renal Failure	12112 (0.0)	19 (0.2)	<0.001
Chronic Lung Disease	1277928 (3.3)	1039 (10.9)	<0.001
Type of Delivery			
Vaginal Delivery	25945792 (67.7)	4445 (46.7)	<0.001
SB occulta	n/a	1842 (19.3)	
SB no hydrocephalus	n/a	1982 (20.8)	
SB with hydrocephalus	n/a	622 (6.5)	
Caesarean delivery	12364506 (32.3)	5071 (53.3)	<0.001
SB Occulta	n/a	1368 (14.4)	
SB no hydrocephalus	n/a	2536 (26.7)	

Characteristic	No Spina Bifida	Spina Bifida	P-value
SB with hydrocephalus	n/a	1166 (12.2)	

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2.

Frequency of fetal and maternal diagnoses and morbidity overall.

Variable	No Spina Bifida	Spina Bifida	p-value
Fetal Outcomes			
Preterm delivery (<37 wk gestation)	2787426 (7.3)	1080 (11.3)	<0.001
Maternal Antenatal Diagnosis			
Fetopelvic disproportion/obstruction	1861285 (4.9)	502 (5.3)	0.38
Maternal Outcomes			
Outcomes Likely Related to a SB Condition			
Neurologic Event	86371 (0.2)	57 (0.6)	<0.001
Pulmonary Event	60576 (0.2)	89 (0.9)	<0.001
Renal Event	25743 (0.1)	21 (0.2)	0.014
VP shunt malfunction/replacement	1116 (0.0)	*	<0.001
Urinary Tract Infection	291566 (0.8)	439 (4.6)	<0.001
Other Outcomes			
Anesthesia-related complication	143721 (0.4)	44 (0.5)	0.53
Blood Transfusion	368359 (1.0)	217 (2.3)	<0.001
Cardiac Event	88798 (0.2)	62 (0.6)	<0.001
Hematologic Event	763717 (2.0)	383 (4.0)	<0.001
DVT	15889 (0.0)	*	0.17
Delivery-related infections	163148 (0.4)	48 (0.5)	0.61
Hemorrhage	1090237 (2.8)	310 (3.3)	0.28

* Incidence too low to report.

Table 3.

Frequency of fetal and maternal morbidity associated with vaginal and cesarean deliveries.

Variable	No Spina Bifida	Spina Bifida	p-value
VAGINAL DELIVERY OUTCOMES			
Fetal Outcomes			
Fetal distress	1640011 (6.3)	324 (7.3)	0.22
Umbilical cord complication	6382296 (24.6)	1130 (25.4)	0.56
Maternal Outcomes			
Outcomes Likely Related to a SB Condition			
Neurologic event	42264 (0.2)	14 (0.3)	0.24
Pulmonary event	18327 (0.1)	*	0.70
Renal event	9649 (0.0)	*	0.0033
Urinary tract infection	178646 (0.7)	157 (3.5)	<0.001
VP Shunt malfunction/replacement	*	*	n/a
Other Outcomes			
Anesthesia-related Complication	73856 (0.3)	14 (0.3)	0.91
Forceps or vacuum delivery	378255 (1.5)	29 (0.6)	0.039
OB-related trauma to perineum or vulva	13623040 (52.5)	2502 (56.3)	0.021
Blood transfusion	136951 (0.5)	28 (0.6)	0.65
Cardiac event	13928 (0.1)	*	n/a
Hematologic event	310269 (1.2)	103 (2.3)	0.002
DVT	7911 (0.0)	*	0.16
Delivery-related infection	49202 (0.2)	*	0.91
Hemorrhage	821234 (3.2)	179 (4.0)	0.13
CESAREAN DELIVERY OUTCOMES			
Outcomes Likely Related to a SB Condition			
Gastrointestinal tract injury	13806 (0.1)	23 (0.5)	<0.001
Urinary tract injury	33879 (0.3)	71 (1.4)	<0.001
VP Shunt malfunction/replacement	460 (0.0)	*	<0.001
Hysterectomy	28052 (0.2)	14 (0.3)	0.76
Neurologic event	44107 (0.4)	43 (0.8)	0.0072
Pulmonary event	42249 (0.3)	84 (1.7)	<0.001
Renal event	18473 (0.1)	*	0.65
Urinary tract infection	112920 (0.9)	283 (5.6)	<0.001
Other Outcomes			
Anesthesia-related Complication	69865 (0.6)	30 (0.6)	0.88
Blood transfusion	231408 (1.9)	189 (3.7)	<0.001
Cardiac event	74870 (0.6)	62 (1.2)	0.009
Hematologic event	453448 (3.7)	280 (5.5)	<0.001
DVT	7978 (0.1)	*	0.67

Variable	No Spina Bifida	Spina Bifida	p-value
Delivery-related Infection	113946 (0.9)	39 (0.8)	0.59
Hemorrhage	269003 (2.2)	130 (2.6)	0.40
Wound complication	*	*	n/a

* Incidence too low to report.