Case Report

Hodgkin Lymphoma revealed by epidural spinal cord compression

Khalil Ghedira ¹, Nidhal Matar¹, Sofiene Bouali¹, Alia Zehani², Adnen Boubaker¹, Hamedh Jemel¹

¹Neurosurgery Department, Institut National de Neurologie, Faculté de Médecine de Tunis, Tunisia, ²Pathology department, Hôpital La Rabta, Faculté de Médecine de Tunis, Tunisia

Context: Hodgkin Lymphoma is rarely diagnosed as spinal cord compression syndrome. Caused by an epidural mass, this complication is often encountered in a late stage of the disease. We report the case of a 40-year-old man presenting with symptoms of low thoracic spinal cord compression due to an epidural tumor on the MRI. **Findings:** Emergent surgery was undertaken on this patient, consisting in laminectomy and tumor resection. After surgery, pain relief and mild neurological improvement were noticed. The histological study revealed a Hodgkin Lymphoma and the patient was referred to chemotherapy and radiotherapy. **Conclusion:** Though chemotherapy is the gold standard treatment for Hodgkin Lymphoma, surgical spinal decompression may be required in epidural involvement of the disease. Diagnosis may be suspected in the presence of lymphadenopathy and general health decay.

Keywords: Hodgkin, Lymphoma, Epidural, Spine, Surgery

Introduction

Spinal cord compression caused by lymphoma is mostly a late manifestation of the illness. It is also three times as frequent with non-Hodgkin's disease as with Hodgkin's disease. We report the case of a 40-year-old man with epidural spinal cord compression as being the initial presentation of a Hodgkin's Lymphoma, which is a rare condition.

Case description

We report the case of a 40-year-old man, without medical history, who was suffering for two weeks from sphincter disorders consisting of urinary leakage and constipation, associated with numbness of the perineum area and the left foot. Furthermore, he suffered from inflammatory lumbar pain, which had started progressively one year ago, accompanied by asthenia, weight loss, generalized pruritus and night sweats.

At examination, he had a temperature of 37.5°C, a normal blood pressure and heart beat rate, scratch marks on arms and legs and right cervical lymphadenopathy. The neurological examination showed a paraparesis predominant on distal muscles, a brisk right patellar reflex,

a normal plantar reflex, a saddle anesthesia, and a distended bladder. The examination of the spine was normal.

Complete blood count revealed a leukocytosis (34,900/mm³), an inflammatory anemia, a high sedimentation rate (78 mm at the 1st hour) and an elevated CRP (58.6 mg/L). HIV and Wright serology were negative.

The spinal MRI showed an abnormal heterogenous signal of the vertebrae: T1, T2, T10, L1, L3, L5, S1 and ilium and ischium, an epidural circumferential mass around the conus terminalis at the L1-L2 level with extension through the left intervertebral foramen, a second left epidural tumor at the L3-L4 level extending through the intervertebral foramen too and several retroperitoneal and iliac adenopathies (Figure 1).

The patient was operated on and a laminectomy of T12, L1 and L2 was achieved. A white and soft lobulated epiduritis was found leaning on the conus medullaris posteriorly and laterally towards the left. A large resection of the posterior portion of the tumor was done until the conus medullaris seemed free of compression from T12 to L2.

Postoperatively, the patient noticed a pain relief and a mild sensorimotor recovery.

The immunohistopathological study revealed a Hodgkin Lymphoma and the patient was sent for chemotherapy (Figure 2).

Correspondence to: Khalil Ghedira, 643, Résidence Ennawres, rue Tabriz, Cité Ennasr 1, 2037, Ariana, Tunisia. Email: khalil.ghedira@gmail.com

Figure 1 Sagittal T2 (A) and T1 (B) weighted pre-operative MRI showing an epidural soft tissue mass centered at L1 on the left side. The lesion enhances with contrast (B). (C) Axial T1 weighted with gadolinium MRI showing epidural mass with extension into the posterior elements and lateral masses.

Discussion

Frequency

Hodgkin's Lymphoma (HL) is a hematological malignancy that affects lymphatic nodes, accounting for 0.5 to 1% of adult cancers.^{1,2} Extra-nodal involvement of the disease is estimated at 10% of cases.¹⁻⁵ Epidural HL leading to a spinal cord compression occurs in

5% of cases, but usually at an advanced stage of the disease: it reveals the diagnosis in only 0.2% of cases.^{4,6,7}

Pathophysiology

Epidural mass in HL may originate from hematogenous dissemination or more likely develops by direct spread

Figure 2 Microphotographs of the histological (A) and the immunohistochemical (B and C) studies. A: hematoxylin and eosin [H&E], original magnification X 40, shows a tumor proliferation composed by lymphocytes, numerous eosinophilic polynuclear (arrow) and uninucleated Hodgkin cells (arrow head). Antibody staining shows that tumor cells express CD15 (B) and CD30 (C).

from retroperitoneal or thoracic lymph nodes.^{5,6} Otherwise, primary vertebral lesion may occur.⁸ In our case, contiguous spread from retroperitoneal and iliac adenopathy through intervertebral foramen is obviously the origin of the epidural lymphoma. Thoracic spine is the most implicated region, followed by lumbar then cervical spine.^{3,4,8}

Clinical findings

Hodgkin's disease usually presents with painless cervical adenopathy which may be accompanied by weight loss, long-term fever and night sweats.^{2,3} Spinal cord involvement owing to an epidural mass is a rare complication that usually develops in the setting of a diffuse, undiagnosed disease or some time after the initial diagnosis. Then, the disease manifests itself with back pain and neurological deficits depending on the level and location of the lesion.^{3,8}

Imaging

The MRI aspect of HL epiduritis is isointense on T1 and T2 weighted images, with a marked contrast enhancement. A full body CT scan is useful to complete the disease extension assessment.³

Treatment

The gold standard treatment of Hodgkin Lymphoma is chemotherapy, however in case of primary disease presenting with spinal cord compression, surgical debulking is indicated and allows for adequate tissue sampling for a histopathological diagnosis.^{3,6,8} Radiotherapy is useful for further local control of the epidural and paraspinal spread.⁶

Prognosis

If adequately treated, HL involving the spinal epidural space has an overall good prognosis, with 86% of neurological function improvement and 61% of complete clinical response.⁷

Conclusion

The spinal cord compression is a well-known complication with patients with Hodgkin Lymphoma, usually

occurring in the setting of an advanced disease. It may rarely be the initial presentation of the illness and the diagnosis may be misleading. A histological identification of the disease is essential, so a surgical biopsy and decompression should be considered as first-line diagnostic and therapeutic options.

Disclaimer statements

Contributors Only the listed authors contributed to the article.

Funding This study has not profited of any financial support nor sponsorship.

Conflicts of interest None of the authors has any potential conflict of interest.

Ethics approval None.

Disclosures None.

ORCID

Khalil Ghedira <http://orcid.org/0000-0002-5068-3191>

References

- 1 Abu-Bonsrah N, Boah AO, Goodwin CR, Larman T, Crane GM, Sciubba DM. Epidural spinal compression as an initial presentation of Hodgkin lymphoma. *J Clin Neurosci* 2016;26:166–8.
- 2 Uehara M, Takahashi J, Hirabayashi H, Kitahara J, Kamijyo T, Ebara S, *et al.* Hodgkin's disease of the thoracic vertebrae. *Spine J* 2013;13(8):59–63.
- 3 Kasper E, Lam F, Luedi M, Zinn P, Pihan G. Primary epidural lymphocyte-depleted Hodgkin's lymphoma of the thoracic spine - presentation of a rare disease variant. *BMC Neurol* 2012;12:64.
- 4 Gupta V, Srivastava A, Bhatia B. Hodgkin disease with spinal cord compression. *J Pediatr Hematol Oncol* 2009;31(10):771–773.
- 5 Citow JS, Rini B, Wollmann R, Macdonald RL. Isolated, primary extranodal Hodgkin's disease of the spine: case report. *Neurosurgery* 2001;49(2):453–457.
- 6 Liao J, Fu T, Chen W, Jung S. Dumbbell-shaped Hodgkin's disease with cauda equina compression mimicking a herniated intervertebral disc, a case report. *Chang Gung Med J* 2007;30(5):458–63.
- 7 Higgins S, Peschel R. Hodgkin's disease with spinal cord compression. A case report and a review of the literature. *Cancer* 1995;75(1):94–8.
- 8 Rabhi M, Ennibi K, Chaari J, Toloune F. Hodgkin's disease presenting with spinal cord compression. *Rev Neurol* 2007;163(11):1109–12.