Published in final edited form as:

J Immigr Minor Health. 2017 October ; 19(5): 1196–1206. doi:10.1007/s10903-016-0518-8.

Relationships between integration and drug use among deported migrants in Tijuana, Mexico

Danielle Horyniak^{1,2,3}, Miguel Pinedo⁴, Jose Luis Burgos¹, and Victoria D. Ojeda¹

¹Division of Global Public Health, School of Medicine, University of California San Diego, La Jolla, CA, 92093, USA

²Centre for Population Health, Burnet Institute, Melbourne, VIC, 3004, Australia

³School of Public Health and Preventive Medicine, Monash University, Melbourne, VIC, 3004, Australia

⁴Alcohol Research Group, University of California Berkeley, Emeryville, CA, 94608, USA

Abstract

Background: Deported migrants face numerous challenges which may elevate their risk for drug use. We examined relationships between integration and drug use among deported migrants in Tijuana, Mexico.

Methods: A cross-sectional survey conducted at a free health clinic included 255 deported Mexican-born migrants residing in Tijuana 6 months. Multivariable logistic regression examined associations between variables across four integration domains (public participation, social connections, macro-level facilitators and foundations) and recent (past six-month) drug use.

Results: The prevalence of recent drug use was 46%. Having sought work in Tijuana in the past six months, greater household affluence, lifetime history of incarceration in both US and Mexico, and lacking health insurance were independently associated with recent drug use.

Discussion: Policies that support access to employment, adequate housing and healthcare in Mexico, particularly for justice-involved deportees, may facilitate successful integration and reduce potential stressors that may contribute to drug use.

Keywords

Migration; deportation; integration; illicit drug use; Mexico

Corresponding author: Dr. Victoria D. Ojeda, Division of Global Public Health, School of Medicine, University of California San Diego, 9500 Gilman Dr, La Jolla, CA, 92093, USA, Ph. +1 (858) 822 6165, vojeda@uscd.edu.

Author contributions

VO and JLB developed and implemented the overall study that produced the data. VO and DH conceptualised this manuscript. DH performed data analysis and drafted the manuscript. All authors commented and contributed text, and gave approval for the manuscript to be submitted.

Background

Successful settlement of migrants, often termed ‘integration’, incorporates elements of participation in economic, social, cultural and political spheres, and is linked to feelings of identity, belonging, and inclusion [1–5]. Integration is a bi-directional process influenced by both migrants’ willingness and agency to participate in their new community and the host community’s acceptance of migrants [3]. Importantly, not only is migration itself increasingly recognised as a social determinant of health [6], but poor post-migration integration can also increase health vulnerabilities [7–9].

Forced repatriation of migrants from the United States (US) has increased dramatically over the past decade, with over 430,000 non-citizens deported in 2013 [10]. The majority of deported migrants are Mexican nationals, many of whom have lived in the US for decades and have little connection to their birth country [11, 12]. Deported migrants face a range of (re)integration challenges upon their return to Mexico, including social isolation, unemployment, homelessness, economic marginalisation, stigmatisation and exposure to violence [13–16]. Furthermore, emerging research suggests that deported Mexican migrants experience poor physical and psychosocial health, and face significant barriers to health insurance and care [17, 18].

Use of illicit drugs, particularly amphetamines, cocaine, and heroin, cause significant morbidity and mortality globally [19]. Tijuana, Baja California, a primary receiving community for deported Mexican migrants [20, 21], presents a significant risk environment for illicit drug use [22]. The city is located on a prominent drug-trafficking route [23, 24], illicit drugs are readily available, and rates of drug use are above the national average [25]. Although studies have consistently shown that Mexicans with a history of travelling or migrating to the US report higher levels of drug use and dependence compared with those with no such history [26–29], fewer studies have specifically examined drug use among deported migrants. Recent research has estimated prevalence of past-year drug use among deported Mexican migrants at 20%–26% [30, 31] and found that this group are significantly more likely to report illicit drug use compared with voluntary return migrants and non-migrants [30]. Evidence also suggests that deported migrants may increase drug use and try new drugs following deportation, with changes in patterns of drug use linked to social and economic exclusion and post-deportation stressors [32–35]. Given the drug use risk environment in Tijuana and the evidence that deported migrants face extensive (re)integration challenges, we aimed to examine relationships between integration and drug use among deported migrants in this setting.

Conceptual framework

This analysis was guided by Ager and Strang’s conceptual framework for integration [5], which outlines four domains considered central to successful integration: ‘*Markers and Means*’, which encapsulates key areas of participation in the public arena - employment, housing, education and health; ‘*Social Connections*’, which assesses relationships with members of a migrant’s community of origin and with the local host community, and links with government structures and services indicating equality with other residents; ‘*Facilitators*’, which refers to macro-level facilitators of inclusion, specifically language and

cultural knowledge, and safety and stability; and *'Foundations'*, which relates to rights and citizenship.

To adapt this framework to the local context, stigmatised characteristics were incorporated as a specific aspect of the 'Safety and stability' sub-domain of the *'Facilitators'* domain. Stigma, which can be enacted through labelling, negative stereotyping, exclusion and discrimination [36] has been identified as a key issue faced by deported migrants [13, 34, 35, 37, 38]. Host communities are often concerned about migrants bringing crime and disorder to their neighbourhoods [39–41]. Deported migrants may be perceived as gang members and drug users, and are vulnerable to being stigmatised, excluded and discriminated against, potentially impacting their sense of safety and stability. As stigma is known to impact social and labour participation and health inequities [42, 43], we hypothesised that stigmatised characteristics may be key markers of integration associated with drug use in this setting.

Methods

Study methods

This study was conducted at Health Frontiers in Tijuana (HFiT), a free primary care clinic providing care (e.g. HIV and sexually transmitted infection testing and care, wound cleaning, mental health services, screenings for diabetes and high blood pressure) for structurally vulnerable populations, including migrants, homeless and unstably housed populations, female sex workers, and people who use drugs [44].

Interviewer-administered surveys were administered to a convenience sample attending HFiT between January-June 2013. Eligibility criteria were: seeking care at HFiT, age 18 years or older, speaking Spanish or English, and able to provide informed consent. All clinic patients who met the criteria were invited to participate in the study regardless of the clinic services they sought, thus we expect our sample to be fairly representative of the clinic population. The survey elicited information on participants' socio-demographic characteristics, migration and deportation experiences, social, economic and civic participation in Tijuana, sexual and drug-related behaviours, health status, and health service access and utilisation. Participants received refreshments, condoms, and \$US10 for their time. The study was approved by ethics boards at University of California San Diego and the HFiT clinic.

The total sample included 339 Mexican-born participants who reported ever being deported from the US. Our survey measured lifetime and past six-month drug use. To accurately assess drug use post-deportation, analysis was limited to those who had resided in Tijuana for at least six months (n=255, 75% of deported migrants) as among recent arrivals (<6 months) we could not discriminate between pre- and post-deportation drug use.

Measures

The outcome of interest was recent (past six-month) use of any illicit drugs, excluding marijuana (yes vs. no).

From available data, we selected all variables which mapped to the four domains of Ager and Strang's integration framework [5]. Except where indicated, all were measured as binary variables (yes vs. no) and refer to the past six months.

'Markers and means': With respect to employment, included variables were: worked in the formal economy, sought work, experienced barriers to obtaining work, and denied a job. Housing variables were: having ever been denied housing/shelter, mostly lived in unstable housing (defined as anything other than having one's own home/room/apartment, or living in a family or friend's home) and household affluence. We generated a variable to assess household affluence based on participants' responses to a series of yes/no questions describing which goods and facilities they had in their primary residence (e.g. electricity, running water, television); questions were derived from the 2010 Mexican census [45]. From these questions a continuous measure was constructed (range 1–9, higher scores indicate greater affluence). Health variables were: self-rated health (good, very good or excellent vs. fair or poor), and depressive symptoms (NIH PROMIS depression scale score 15 [46]). Information on highest level of education attained was collected, but we could not differentiate between education completed in the US and education completed following deportation, so this variable was considered as a socio-demographic covariate rather than an indicator of integration.

'Social connections': Variables were: currently have any family or friends in Tijuana, separated from family in the US due to deportation, banned from returning to the US, feeling rejected in Tijuana, and visited a doctor. "Being banned from returning to the US" was included as we hypothesised that inability to return to the US may be an important motivator promoting integration.

'Facilitators': Included variables were: low Spanish proficiency, country identify most strongly with (USA vs. Mexico/identify with both countries equally), been threatened or harassed in Tijuana, and been threatened or harassed by police or military in Tijuana. For the stigmatised characteristics sub-domain, variables were: having a visible tattoo and lifetime incarceration history (categorised as never incarcerated, incarcerated in Mexico only, incarcerated in US only, or incarcerated in both Mexico and US). Tattoos have long been considered a marker of social undesirability and engagement in deviant behaviours, such as crime and drug use [47–49]; previous research among deported migrants has found that tattoos are considered a marker of 'Americanisation' and deportation history [37]. Similarly, reports suggest that incarceration history is associated with being perceived by local communities as a drug user, gang member and 'dangerous' outsider [13, 35, 50].

'Foundations': Variables mapped to this domain were: have any form of Mexican identification (birth certificate, driver's licence, passport, consular identification card, Instituto Federal Electoral voting card, or Clave Única de Registro de Población population registry card [equivalent to the US Social Security Number]), and have health insurance.

Additional socio-demographic and migration-related variables included in analysis were: age, sex, birth state, highest level of education completed, duration of residence in Tijuana, total time spent in the US, number of times deported and time since last deportation.

Data Analysis

To characterise the sample, socio-demographic, migration and drug use variables were analysed descriptively and using bivariate logistic regression. To examine the associations between integration and recent drug use we first constructed separate multivariable logistic regression models for each of the four integration domains. We then constructed a final multivariable model to adjust for socio-demographic and migration variables, and relationships between integration variables across domains. We first assessed the data for multicollinearity. We then incorporated integration variables from each domain which were statistically significant ($p < 0.05$) within that domain's multivariable analysis, in order to create the most parsimonious model. Blocks of variables were added sequentially beginning with socio-demographic and migration variables. Integration domains were added in the order presented in Ager and Strang's framework which reflects a social-ecological model of health: Markers & Means (individual-level), Social Connections, Facilitators (social/environmental-level), Foundations (structural-level). Likelihood-ratio tests were used to compare models following addition of each domain. We also conducted a sensitivity analysis including all integration variables regardless of significance within individual models and found that the direction and magnitude of results were largely unchanged (data not shown). Analyses were performed using Stata 13.1 (Statacorp LP, Texas, USA).

Results

Socio-demographic and migration characteristics of the sample

Participants' socio-demographic and migration characteristics are presented in Tables 1. In brief, the sample was predominantly male (88%), with a median age of 43 years (Interquartile Range (IQR): 35–50 years). Twelve per cent of participants were born in Baja California and ten per cent reported that they or a family member spoke an Indigenous language. Fifty-four per cent had completed high school or higher education. Participants had lived in the US for a median of 15 years (IQR: 7–23 years). Sixty per cent had been deported more than once, among whom a median of three deportations (IQR: 2–5) were reported. Fifty-three per cent of the participants experienced their most recent deportation within the past five years. Most participants (72%) reported that they were banned from returning to the US, among whom 25% reported a lifetime ban (data not shown). Thirty-eight per cent of participants reported that they planned to return to the US in the six months following interview.

Post-migration integration characteristics of the sample

Post-migration integration characteristics are presented in Table 2. Within the '*Means and Markers*' domain, although 59% of participants had worked in the formal economy in the past six months, seeking work (57%) and experiencing barriers to obtaining work (62%) were common. Fifty-eight per cent of participants reported living in unstable accommodation in the past six months, with poor housing amenity reported (median score: 5, IQR: 3–6).

Within the '*Social Connections*' domain, despite family separation being common (74%), most participants reported having family and/or friends in Tijuana (65%).

Within the '*Facilitators*' domain, 20% of participants reported being harassed or threatened in Tijuana in the past six months, and 45% had been harassed or threatened by police/military. Stigmatising characteristics were prevalent; 44% of participants reporting having a visible tattoo, and 78% had a history of incarceration.

Within the '*Foundations*' domain, 56% of participants possessed any form of Mexican identification, and 40% had health insurance at the time of interview.

Recent drug use

The prevalence of recent illicit drug use was 46%. Injecting was a common route of administration (88% of recent drug users; Table 3), with crystal methamphetamine and heroin the most commonly used drugs. Among those who had used illicit drugs in the past six months, 61% reported they lived in Mexico when they initiated illicit drug use. Sixty-eight per cent of recent drug users reported being in need of help to control their drug use at the time of interview.

Associations between integration and recent drug use

Table 4 displays the results from multivariable logistic regression analyses examining associations between each individual integration domain and drug use. Within the '*Means and Markers*' domain (Model 1), having sought work in Tijuana and higher household affluence were both negatively associated with recent drug use. Within the '*Social Connections*' domain (Model 2) having family or friends in Tijuana was positively associated with recent drug use and having recently visited a doctor was negatively associated. Within the '*Facilitators*' domain (Model 3) having a visible tattoo and lifetime incarceration in Mexico and in both Mexico and the US were all positively associated with recent drug use. Within the '*Foundations*' domain (Model 4), having any form of health insurance was negatively associated with recent drug use. All individual domain models were statistically significant and fit the data well.

In the final model incorporating integration variables from across all domains (Table 5) and adjusting for socio-demographic and migration variables, two variables from the '*Means and Markers*' domain were negatively associated with recent drug use: having sought work in Tijuana in the past six months (Adjusted Odds Ratio [AOR]: 0.37, 95% Confidence Interval [CI]: 0.18 – 0.77), and household affluence (AOR: 0.82, 95% CI: 0.69–0.96). No '*Social Connections*' variables were associated with recent drug use. Within the '*Facilitators*' domain, history of incarceration in both US and Mexico (AOR: 3.30, 95% CI: 1.09–9.96) was positively associated with recent drug use, but no association was observed with history of incarceration in Mexico alone (AOR: 2.08, 95% CI: 0.71–6.10) or history of incarceration in the U.S. alone (AOR: 2.66, 95% CI: 0.92–7.68). Within the '*Foundations*' domain, having health insurance was negatively associated with recent drug use (AOR: 0.37, 95% CI: 0.17–0.78). Likelihood-ratio tests indicated statistically significant improvement of the model with the addition of each block of variables with the exception of '*Social Connections*' ($p=0.16$).

Discussion

In this study of structurally vulnerable deported migrants in Tijuana, Mexico, illicit drug use was pervasive; 46% of participants reported past six-month drug use, almost double the prevalence of past-year drug use reported among general samples of deported Mexican migrants [30, 31]. Factors which may have contributed to the high prevalence of recent drug use recorded include the location of the clinic where the study was conducted (in Tijuana's red light district, where cheap drugs are readily available and drug use is prevalent), and that some participants may have accessed harm reduction services (e.g. needle exchange) provided by the clinic's collaborator. Importantly, over half our sample reported initiating drug use in Mexico. Injecting drug use and use of heroin and crystal methamphetamine were common among our sample, characteristics associated with high risk of harms including HIV transmission and overdose [51, 52]. Substantial public health effort is already in place to address epidemics of drug use and HIV in Tijuana [53, 54]; our findings suggest that targeted efforts to ensure inclusion of deported migrants in prevention and treatment services may be warranted. Potential strategies include collaboration with government and non-government organisations to incorporate drug use screening, brief intervention and referrals into services provided to deported migrants. Similarly, ensuring deported migrants know about programs to prevent and reduce drug-related harms and can access HIV testing services will benefit the community and help to reduce the HIV epidemic in the region.

Supporting our hypothesis that integration challenges would be associated with drug use, we found that each individual-domain level analysis was significant, and in the final model, the addition of each domain with the exception the *'Social Connections'* domain significantly improved the model. . In the final multivariable model, within the *'Means and Markers'* domain seeking work in Tijuana and greater household affluence were both negatively associated with drug use. This is an interesting finding given that neither obtaining work nor housing was significantly associated with illicit drug use. A possible explanation for this finding is that the process of establishing housing and job prospects (rather than the outcome itself) may facilitate feelings of agency and belonging which may be protective against drug use. The relationship between seeking work and drug use may also reflect that some people may not use drugs while seeking employment if they feel their chances of obtaining employment may be compromised by drug use.

No independent associations were detected between any *'Social Connections'* variables and drug use. This is surprising, given that qualitative research has shown social exclusion to be a key issue faced by deported migrants as well as shaping mental health [34, 35]. Recent research described the familial relationships and sense of belonging cultivated among deported migrants employed in a call centre in Mexico City despite exclusion from the broader community [14], suggesting that the type and quality of social relationships may have variable impacts on drug use risk. The lack of association seen in our study may be due to limitations in the data collected; it would be useful to capture more detailed information about the nature and quality of deported migrants' social connections, for example with family, friends and other deported migrants, in future studies.

Within the ‘*Facilitators*’ domain, recent drug use was positively associated with a history of incarceration in both US and Mexico, but not with a history of incarceration in the US or Mexico alone. This finding is relatively unsurprising given that prisons have been identified as risk environments for drug use [55]. US incarceration may be indicative of a history of drug use, including drug-related behaviours which precipitated deportation [33]. Incarceration in the US has also been associated with changing patterns of drug use post-deportation [32]. Deported migrants with a history of incarceration face multiple markers of stigma [14, 35], potentially increasing vulnerability to drug use. The temporal relationships between incarceration, deportation and drug use remain unclear however, indicating a need for longitudinal research. Importantly, our survey did not capture information about reasons for incarceration, so we cannot differentiate between incarceration in immigration detention prior to deportation and incarceration in prison for other reasons. It would be useful in future research to collect detailed information to enable examination of the possible differences in relationships between drug use and incarceration across these different settings.

Despite the implementation of universal public health insurance (‘*Seguro Popular*’), which has increased coverage in the general Mexican population [56], a significant proportion of our sample was uninsured. This finding is consistent with previous research among deported migrants [18]. Within the ‘*Foundations*’ domain, having health insurance was strongly negatively associated with drug use. A possible explanation for this finding is that uninsured participants may have limited access to appropriate care for mental health conditions, and may turn to substance use as a coping mechanism, as has been reported among other migrant samples [e.g. 57, 58]. Documentation status was not independently associated with drug use, but it is likely that this underpins the health insurance findings. Proof of Mexican citizenship is required for enrolment in ‘*Seguro Popular*’. For those lacking documentation, obtaining it may require travel to a person’s city of origin to obtain original copies of birth documents, a journey some migrants do not have the resources to undertake or may not wish to undertake if they hope to return to the US.

Our findings reflect the complex nature of the integration process, and highlight the need for comprehensive responses to facilitate social, economic and civic participation and reduce health inequalities. Our findings suggest that policies that support access to employment, adequate housing and healthcare are integral to preventing drug use by addressing potential stressors which may contribute to this behaviour. In particular, we highlight the need for policies to enable deported migrants to obtain identification documents and gain access to health insurance. In a promising move, the governments of the states of Guerrero and Nayarit recently signed agreements with the city of Tijuana to help facilitate access to birth certificates for deported migrants from those states [59]. Coordinated binational efforts to provide copies of birth certificates prior to deportation (e.g. via Mexican consulates in the US) may also ease (re)integration challenges.

It is worthwhile reflecting on the assumption that integration is an inherently positive outcome. Acculturation to US culture has been associated with declines in health status and uptake of health-damaging behaviours for Mexican migrants to the US [60–62]. Likewise, it is possible that integration in Tijuana may have unanticipated negative impacts due to the risk environment for drug use; qualitative research has suggested that exposure to the drug

use environment is a facilitator of drug use for deported migrants [33]. Our analysis detected increased odds of drug use with longer duration of residence in Tijuana providing further support for this relationship. Longitudinal research which examines both exposure to the drug use risk environment and changes over time in post-migration integration will be essential in understanding drug use and reducing drug-related harms among deported migrants in Tijuana. Such research could have important implications for policy and practice. For example, if evidence supports a relationship between exposure to the drug use risk environment and subsequent drug use this would highlight an urgent need for early intervention with deported migrants upon arrival in Tijuana. In 2007, Mexico's Federal Institute of Migration established the 'Repatriación Humana' ('Humane Repatriation') program in 11 border cities including Tijuana, which offers basic services (e.g. resettlement information, referrals to shelters, medical consultations) to deportees [63]. Access to this program is, however, limited to newly-arrived deportees. Making such a program more widely available to deportees (e.g. throughout the first six-twelve months post-deportation) may aid them during the transition and resettlement phase. The 'Repatriación Humana' also provides assistance to deported migrants in returning to their cities of origin, including provision of transportation [64]. Providing deported migrants with resources to return home, *if desired*, is important, and may be protective for some migrants. This should however be optional, as each deportee's circumstances are unique, and return to their city of origin may not be desirable or feasible. For example, a recent community report notes that social factors (e.g. unemployment in sending communities, desire to stay close to the border to maintain contact with U.S.-based family and friends) dampen deportees' interest in returning to sending communities [65]. Continued access to support programs which harness the human capital of returning migrants and promotes meaningful participation may reduce the stressors associated with deportation and thus potentially reduce drug use.

Limitations

The study used convenience sampling methods, drawing on a structurally vulnerable sample accessing free healthcare. Deported migrants who were less disadvantaged may have been excluded, resulting in an overestimate of drug use prevalence, and possibly inflating relationships between integration and drug use. Thus, our findings may not be generalizable to the overall population of deported migrants in Tijuana or more broadly.

This secondary analysis leveraged existing data to explore the relationship between integration vis-à-vis drug use, consequently, some components of Ager and Strang's framework were unable to be measured (for example, we lacked variables which mapped to the education sub-domain, and our measures of social connections were only general and did not measure sources of social support in great detail).

Importantly, the integration framework was developed within a refugee resettlement context, thus it may not fully capture the integration experiences of deported migrants or the Tijuana context. Resettled refugee populations are often unfamiliar with their new host country; they may truly initiate the process of integration into a new environment. By contrast, many deportees have ties to their country of origin, including familiarity with the cultural context, language and values. Yet, they may feel, and are often treated, as outsiders [13, 35]. An

important limitation of this integration framework in its application to the deported migrant context is that it is not designed to capture the nuances of re-integration and re-acculturation challenges. Future research would benefit from utilising a combination of integration and risk environment frameworks (e.g. [22, 66]), and considering both individual and contextual-level factors influencing acculturation to assess drug use risk.

As this study was cross-sectional, we cannot ignore the potential for reverse causality, that is, that drug use leads to poor integration outcomes. Irrespective of the direction of this relationship, our findings highlight the importance of program and policy response to address both drug use and social and economic integration among this population. We also cannot draw conclusions about the temporal relationships between deportation and drug use initiation; although over 60% of recent drug users in our study reported initiating drug use in Mexico, it is possible that initiation occurred prior to migration to the US rather than following deportation.

Conclusion

Recent illicit drug use is pervasive among deported migrants in Tijuana, and is associated with a range of negative integration experiences. Policies that support access to employment, housing and healthcare in Mexico, particularly for justice-involved deportees, may facilitate successful integration and reduce potential stressors that may contribute to drug use.

Acknowledgements

The authors would like to thank the study participants for sharing their experiences with us, and our collaborators at PrevenCasa for their continued support. We acknowledge funding from the following sources: Australian Endeavour Awards (#4722_2015 - Horyniak), Australian National Health & Medical Research Council (Early Career Fellowship #1092077 - Horyniak), US National Institute on Alcohol Abuse and Alcoholism (T32AA007240 - Pinedo), US National Institute on Drug Abuse (K01DA025504 - Ojeda), US National Institute of Mental Health (K01MH095680 - Burgos), University of California GloCal Health Fellowship (Ojeda), and Center for US-Mexican Studies Fellowship (Ojeda). The funding bodies played no role in the study design, data analysis or preparation of the manuscript for publication.

References

1. Fozdar F, Hartley L. Civic and Ethno Belonging among Recent Refugees to Australia. *Journal of Refugee Studies* 2014;27(1):126–44.
2. Wille J. Agency and belonging: Southern Sudanese Former Refugees' reflections on life in Australia. *Australasian Review of African Studies* 2011;32(2):80–100.
3. Valtonen K. From the Margin to the Mainstream: Conceptualizing Refugee Settlement Processes. *Journal of Refugee Studies* 2004;17(1):70–96.
4. Ruben R, Van Houte M, Davids T. What Determines the Embeddedness of Forced-Return Migrants? Rethinking the Role of Pre- and Post-Return Assistance. *International Migration Review* 2009;43(4):908–37.
5. Ager A, Strang A. Understanding Integration: A Conceptual Framework. *Journal of Refugee Studies* 2008;21(2):166–91.
6. Castañeda H, Holmes SM, Madrigal DS, Young MED, Beyeler N, Quesada J. Immigration as a Social Determinant of Health. *Annu Rev Public Health* 2015;36(1):375–92. [PubMed: 25494053]
7. Davies AA, Borland RM, Blake C, West HE. The Dynamics of Health and Return Migration. *PLoS Med* 2011;8(6):e1001046. [PubMed: 21738448]

8. Viruell-Fuentes EA, Miranda PY, Abdulrahim S. More than culture: Structural racism, intersectionality theory, and immigrant health. *Soc Sci Med* 2012;75(12):2099–106. [PubMed: 22386617]
9. Gil-González D, Carrasco-Portiño M, Vives-Cases C, Agudelo-Suárez AA, Castejón Bolea R, Ronda-Pérez E. Is health a right for all? An umbrella review of the barriers to health care access faced by migrants. *Ethn Health* 2014;20(5):523–41. [PubMed: 25117877]
10. US Department of Homeland Security. 2013 Yearbook of Immigration Statistics Washington, D.C.: US Department of Homeland Security, 2014.
11. Forced Apart (By the Numbers): Non-Citizens Deported Mostly for Nonviolent Offenses New York: Human Rights Watch, 2009.
12. Hagan J, Eschbach K, Rodriguez N. U.S. Deportation Policy, Family Separation, and Circular Migration. *Int Migr Rev* 2008;42(1):64–88.
13. Wheatley C. Push Back: U.S. Deportation Policy and the Reincorporation of Involuntary Return Migrants in Mexico. *The Latin Americanist* 2011;55(4):35–60.
14. Anderson J. “Tagged as a criminal”: Narratives of deportation and return migration in a Mexico City call center. *Latino Studies* 2015;13(1):8–27.
15. Goldenberg S, Strathdee S, Gallardo M, Patterson TL. “People here are alone, using drugs, selling their body”: Deportation and HIV vulnerability among clients of female sex workers in Tijuana. *Field Actions Science Reports* 2010;Special Issue 2.
16. Boehm DA. US-Mexico Mixed Migration in an Age of Deportation: An Inquiry into the Transnational Circulation of Violence. *Refugee Survey Quarterly* 2011;30(1):1–21.
17. Bojorquez I, Aguilera RM, Ramirez J, Cerecero D, Mejia S. Common Mental Disorders at the Time of Deportation: A Survey at the Mexico-United States Border. *J Immigr Minor Health* 2014;10.1007/s10903-014-0083-y.
18. Fernández-Niño JA, Ramírez-Valdés CJ, Cerecero-García D, Bojorquez-Chapela I. Deported Mexican migrants: health status and access to care. *Rev Saude Publica* 2014;48(3):478–85. [PubMed: 25119943]
19. Degenhardt L, Whiteford HA, Ferrari AJ, et al. Global burden of disease attributable to illicit drug use and dependence: findings from the Global Burden of Disease Study 2010. *The Lancet* 2013;382(9904):1564–74.
20. U.S. Department of Homeland Security. ICE Enforcement and Removal Operations Report: Fiscal Year 2014 U.S. Department of Homeland Security, 2014.
21. Masferrer C, Roberts BR. Going Back Home? Changing Demography and Geography of Mexican Return Migration. *Population Research and Policy Review* 2012;31:465–96.
22. Rhodes T. The ‘risk environment’: a framework for understanding and reducing drug-related harm. *Int J Drug Policy* 2002;13:85–94.
23. Brouwer KC, Case P, Ramos R, et al. Trends in the Production and Trafficking and Consumption of Methamphetamine and Cocaine in Mexico. *Subst Use Misuse* 2006;41:707–27. [PubMed: 16603456]
24. Bucardo J, Brouwer KC, Magis-Rodriguez C, et al. Historical Trends in the Production and Consumption of Illicit Drugs in Mexico: Implications for the Prevention of Blood Borne Infections. *Drug Alcohol Depend* 2005;79:281–93. [PubMed: 16102372]
25. Instituto Nacional de Salud Pública. Encuesta Nacional de Adicciones 2008 Cuernavaca, Mexico: Instituto Nacional de Salud Pública, 2008.
26. Guerrero EG, Villatoro JA, Kong Y, Gamino MB, Vega WA, Mora ME. Mexicans’ use of illicit drugs in an era of drug reform: national comparative analysis by migrant status. *Int J Drug Policy* 2014;25(3):451–7. [PubMed: 24816376]
27. Borges G, Breslau J, Orozco R, et al. A cross-national study on Mexico-US migration, substance use and substance use disorders. *Drug Alcohol Depend* 2011;117(1):16–23. [PubMed: 21296509]
28. Borges G, Medina-Mora ME, Orozco R, Fleiz C, Cherpitel C, Breslau J. The Mexican migration to the United States and substance use in northern Mexico. *Addiction* 2009;104(4):603–11. [PubMed: 19215601]

29. Borges G, Medina-Mora ME, Breslau J, Aguilar-Gaxiola S. The effect of migration to the United States on substance use disorders among returned Mexican migrants and families of migrants. *Am J Public Health* 2007;97(10):1847–51. [PubMed: 17761563]
30. Zhang X, Martinez-Donate AP, Nobles J, Hovell MF, Rangel MG, Rhoads NM. Substance Use Across Different Phases of the Migration Process: A Survey of Mexican Migrants Flows. *J Immigr Minor Health* 2015;10.1007/s10903-014-0109-5.
31. Rangel MG, Martinez-Donate AP, Hovell MF, et al. A two-way road: rates of HIV infection and behavioral risk factors among deported Mexican labor migrants. *AIDS Behav* 2012;16(6):1630–40. [PubMed: 22562390]
32. Robertson AM, Rangel MG, Lozada R, Vera A, Ojeda VD. Male injection drug users try new drugs following U.S. deportation to Tijuana, Mexico. *Drug Alcohol Depend* 2012;120(1–3):142–8. [PubMed: 21835559]
33. Ojeda VD, Robertson AM, Hiller SP, et al. A Qualitative View of Drug Use Behaviors of Mexican Male Injection Drug Users Deported from the United States. *J Urban Health* 2011;88(1):104–17. [PubMed: 21246301]
34. Martin YC. The syndemics of removal: trauma and substance abuse. In: Brotherton DC, Stageman DL, Leyro SP, editors. *Outside Justice: Immigration and the Criminalizing Impact of Changing Policy and Practice* New York: Springer; 2013.
35. Brotherton DC, Barrios L. Displacement and stigma: The social-psychological crisis of the deportee. *Crime, Media, Culture* 2009;5(1):29–55.
36. Link BG, Phelan JC. Conceptualizing Stigma. *Annual Review of Sociology* 2001;27:363–85.
37. Pinedo M, Burgos JL, Ojeda VD. A critical review of social and structural conditions that influence HIV risk among Mexican deportees. *Microbes Infect* 2014;16(5):379–90. [PubMed: 24583278]
38. Schuster L, Majidi N. Deportation Stigma and Re-migration. *Journal of Ethnic and Migration Studies* 2015;41(4):635–52.
39. Fitzgerald J, Curtis KA, Corliss CL. Anxious Publics: Worries About Crime and Immigration. *Comparative Political Studies* 2012;45(4):477–506.
40. Collins J, Reid C. Minority Youth, Crime, Conflict, and Belonging in Australia. *Int Migration & Integration* 2009;10(4):377–91.
41. Simon RJ, Sikich KW. Public Attitudes toward Immigrants and Immigration Policies across Seven Nations. *International Migration Review* 2007;41(4):956–62.
42. Hatzenbuehler ML, Phelan JC, Link BG. Stigma as a fundamental cause of population health inequalities. *Am J Public Health* 2013;103(5):813–21. [PubMed: 23488505]
43. Link BG, Phelan JC. Stigma and its public health implications. *Lancet* 2006;367(9509):528–9. [PubMed: 16473129]
44. Ojeda VD, Eppstein A, Lozada R, et al. Establishing a binational student-run free-clinic in Tijuana, Mexico: a model for US-Mexico border states. *J Immigr Minor Health* 2014;16(3):546–8. [PubMed: 23371839]
45. Instituto Nacional de Estadística y Geografía. Censo de población y vivienda 2010 Aguascalientes, Aguascalientes. Available from: www.inegi.org.mx.
46. Ferraiolo N, Pinedo MM, J., Burgos JL, Vargas-Ojeda AC, Rodriguez M, Ojeda VD. Depressive symptoms among patients at a clinic in the Red Light District of Tijuana, Mexico. *International Journal of Culture and Mental Health* In press;10.1080/17542863.2016.1144776.
47. Adams J. Marked Difference: Tattooing and its Association with Deviance in the United States. *Deviant Behavior* 2009;30(3):266–92.
48. Lane DC. Tat's All Folks: An Analysis of Tattoo Literature. *Sociology Compass* 2014;8(4):398–410.
49. Jennings WG, Fox BH, Farrington DP. Inked into crime? An examination of the causal relationship between tattoos and life-course offending among males from the Cambridge study in delinquent development. *Journal of Criminal Justice* 2014;42(1):77–84.
50. García V, González L. Labor Migration, Drug Trafficking Organizations, and Drug Use: Major Challenges for Transnational Communities in Mexico. *Urban Anthropology and Studies of Cultural Systems and World Economic Development* 2009;38(2–4):303–44. [PubMed: 20927201]

51. Degenhardt L, Mathers B, Guarinieri M, et al. Meth/amphetamine use and associated HIV: Implications for global policy and public health. *Int J Drug Policy* 2010;21(5):347–58. [PubMed: 20117923]
52. Degenhardt L, Hall W. Extent of illicit drug use and dependence, and their contribution to the global burden of disease. *Lancet* 2012;379:55–70. [PubMed: 22225671]
53. Martinez-Donate AP, Hovell MF, Rangel MG, et al. Migrants in transit: the importance of monitoring HIV risk among migrant flows at the Mexico-US border. *Am J Public Health* 2015;105(3):497–509. [PubMed: 25602882]
54. Strathdee SA, Magis-Rodriguez C, Mays VM, Jimenez R, Patterson TL. The emerging HIV epidemic on the Mexico-U.S. border: an international case study characterizing the role of epidemiology in surveillance and response. *Ann Epidemiol* 2012;22(6):426–38. [PubMed: 22626001]
55. Rhodes T, Singer M, Bourgois P, Friedman SR, Strathdee SA. The social structural production of HIV risk among injecting drug users. *Soc Sci Med* 2005;61(5):1026–44. [PubMed: 15955404]
56. Gutierrez JP, Garcia-Saiso S, Dolci GF, Hernandez Avila M. Effective access to health care in Mexico. *BMC Health Serv Res* 2014;14:186. [PubMed: 24758691]
57. Horyniak D, Higgs P, Cogger S, Dietze P, Bofu T. Heavy alcohol consumption among marginalised African refugee young people in Melbourne, Australia: motivations for drinking, experiences of alcohol-related problems and strategies for managing drinking. *Ethn Health* 2015;21(3):284–99. [PubMed: 26169071]
58. Posselt M, Procter N, Galletly C, Crespigny C, de Crespigny C. Aetiology of coexisting mental health and alcohol and other drug disorders: Perspectives of refugee youth and service providers. *Australian Psychologist* 2015;50(2):130–40.
59. Logran importantes acuerdos para migrantes en Congreso Nacional de Migración en Tijuana [Important achievements for migrants at the Migration Congress in Tijuana]. San Diego Red [Internet] 2015 Available from: <http://www.sandiegored.com/noticias/62229/Logran-importantes-acuerdos-para-migrantes-en-Congreso-Nacional-de-Migracion-en-Tijuana/>.
60. Blanco C, Morcillo C, Alegría M, et al. Acculturation and drug use disorders among Hispanics in the U.S. *J Psychiatr Res* 2013;47(2):226–32. [PubMed: 23128062]
61. Carter-Pokras O, Zambrana RE, Yankelovich G, Estrada M, Castillo-Salgado C, Ortega AN. Health status of Mexican-origin persons: do proxy measures of acculturation advance our understanding of health disparities? *J Immigr Minor Health* 2008;10(6):475–88. [PubMed: 18470618]
62. Antecol H, Bedard K. Unhealthy assimilation: Why do immigrants converge to American health status levels? *Demography* 2006;43(2):337–60. [PubMed: 16889132]
63. Government of Mexico. Programa de Repatriación [Repatriation Programs] [28/07/2016] Available from: www.gob.mx/inm/acciones-y-programas/programa-de-repatriacion-12469.
64. Government of Mexico. Procidimiento de Repatriación al Interior de México (PRIM) [Repatriation procedures to the interior of Mexico]. Available from: www.inm.gob.mx/static/repatriacion_h/Repatriacion_H.pdf.
65. Instituto para la Seguridad y la Democracia (Insyde). Diagnóstico del Instituto Nacional de Migración: Hacia un Sistema de Rendición de Cuentas en pro de los Derechos de las Personas Migrantes en México [Assessment of the National Institute of Migration: Towards a System of Accountability for the Rights of Migrants in Mexico] 2013 [27/09/2016] Available from: http://insyde.org.mx/wp-content/uploads/2014/03/Diagnostico_INM_Insyde_2013_Completo
66. Rhodes T. Risk environments and drug harms: a social science for harm reduction approach. *Int J Drug Policy* 2009;20(3):193–201. [PubMed: 19147339]

Socio-demographic and migration characteristics of deported migrants attending a free health clinic in Tijuana, Mexico and bivariate associations with recent drug use¹, 2013

Table 1.

Variable	Total N=255 n (%)	Drug use (past 6m)		Odds Ratio (95% CI)	p-value
		Yes N=117 n (%)	No N=138 n (%)		
Socio-demographics					
Median age (IQR)	43 (35–50)	40 (34–47)	45 (39–52)	0.94 (0.91–0.97)	<0.001
Male sex	225 (88)	105 (90)	120 (87)	1.43 (0.65 – 3.17)	0.376
Born in Baja California	31 (12)	23 (20)	8 (6)	3.97 (1.70 – 9.26)	0.001
Completed high school or higher education	137 (54)	79 (68)	58 (42)	2.87 (1.72–4.79)	<0.001
Migration					
Total time spent in the US					
<1 year	14 (6)	9 (8)	5 (4)	1	
1–5 years	28 (11)	16 (14)	12 (9)	0.74 (0.20–2.79)	0.657
5–10 years	38 (15)	17 (15)	21 (15)	0.45 (0.13–1.60)	0.216
10 years	173 (68)	74 (63)	99 (72)	0.42 (0.13–1.29)	0.129
Not reported	2 (1)	1 (1)	1 (1)	--	
Deported more than once	153 (60)	77 (66)	76 (55)	1.53 (0.92–2.56)	0.103
Time since last deportation					
<1 year	7 (3)	2 (2)	5 (4)	1	
1–5 years	127 (50)	48 (41)	79 (57)	1.52 (0.28–8.14)	0.625
5–10 years	44 (17)	25 (21)	19 (14)	3.29 (0.57–18.83)	0.181
10 years	63 (25)	41 (35)	22 (16)	4.66 (0.83–26.01)	0.079
Not reported	14 (6)	1 (1)	13 (9)	--	
Duration of residence in Tijuana					
6–12 months	33 (13)	6 (5)	27 (20)	1	
1–3 years	77 (30)	24 (21)	53 (38)	2.04 (0.74–5.58)	0.166
3 years	145 (57)	87 (74)	58 (42)	6.75 (2.62–17.37)	<0.001
/ Missing data excluded					

Integration characteristics of deported migrants attending a free health clinic in Tijuana, Mexico, and bivariate associations with recent drug use¹, 2013

Table 2.

Variable	Total N=255 n (%)	Drug use (past 6m)		Odds Ratio (95% CI)	p-value
		Yes N=117 n (%)	No N=138 n (%)		
Means & Markers					
Worked in formal economy last 6m	151 (59)	70 (60)	81 (59)	1.05 (0.63–1.73)	0.854
Sought work in Tijuana last 6m	145 (57)	54 (46)	91 (66)	0.44 (0.27–0.73)	0.002
Experienced barriers to obtaining work last 6m	159 (62)	70 (59)	89 (65)	0.82 (0.49–1.36)	0.444
Denied a job in Tijuana last 6m	105 (41)	47 (40)	58 (42)	0.93 (0.56–1.53)	0.764
Mostly lived in unstable accommodation last 6m	148 (58)	72 (62)	76 (55)	1.31 (0.79–2.15)	0.298
Household affluence score ²	5 (3–6)	4 (0–5)	5 (4–6)	0.78 (0.70–0.87)	<0.001
Denied housing/shelter in Tijuana last 6m	43 (17)	21 (18)	22 (16)	1.15 (0.60–2.22)	0.670
Good/very good/excellent self-rated health	134 (53)	55 (47)	79 (57)	0.67 (0.41–1.11)	0.121
Depressive symptoms (15 on NIH PROMIS scale)	139 (55)	72 (62)	67 (49)	1.70 (1.03–2.80)	0.039
Social Connection					
Has family or friends in Tijuana	165 (65)	83 (71)	82 (59)	1.67 (0.99–2.82)	0.056
Separated from family due to deportation	189 (74)	93 (79)	96 (70)	1.70 (0.95–3.02)	0.073
Banned from returning to USA	183 (72)	91 (78)	92 (67)	1.78 (1.00–3.14)	0.047
Felt rejected in Tijuana last 6m	92 (36)	45 (38)	47 (34)	1.21 (0.72–2.02)	0.466
Visited a doctor in the last 6m	115 (45)	40 (34)	75 (54)	0.44 (0.26–0.73)	0.002
Facilitators					
Low Spanish proficiency	4 (2)	2 (2)	2 (2)	1.18 (0.16–8.53)	0.868
Identify most strongly with USA	68 (27)	27 (23)	41 (30)	0.72 (0.41–1.26)	0.252
Threatened/harassed in Tijuana last 6m	50 (20)	26 (22)	24 (17)	1.36 (0.73–2.52)	0.334
Threatened/harassed by police/military in Tijuana	115 (45)	61 (52)	54 (39)	1.69 (1.03–2.79)	0.038
Have a visible tattoo	111 (44)	66 (56)	45 (33)	2.67 (1.61–4.45)	<0.001
Ever incarcerated, in Mexico only	67 (26)	37 (32)	30 (22)	4.52 (2.03–10.06)	<0.001
Ever incarcerated, in US only	71 (28)	28 (24)	43 (31)	2.39 (1.08–5.29)	0.032
Ever incarcerated, in both Mexico and US	60 (24)	40 (34)	20 (14)	7.33 (3.18–16.88)	<0.001
Foundation					
Has any form of Mexican ID	142 (56)	55 (47)	87 (63)	0.52 (0.32–0.86)	0.011

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Variable	Total N=255 n (%)	Drug use (past 6m)		Odds Ratio (95% CI)	p-value
		Yes N=117 n (%)	No N=138 n (%)		
Has any form of health insurance	101 (40)	31 (27)	70 (51)	0.35 (0.21–0.59)	<0.001

Missing data excluded
Median (Interquartile range)

Table 3.

Drug use characteristics of deported migrants attending a free health clinic in Tijuana, Mexico, who reported recent drug use, 2013

Variable	N=117 n (%)
Country lived in most of the time when first started using drugs	
Mexico	71 (61)
USA	42 (36)
Other	4 (3)
Drug/s used in past six months	
Marijuana	37 (32)
Heroin	47 (42)
Cocaine (powder)	7 (6)
Crack Cocaine	7 (6)
Heroin and cocaine together	14 (12)
Crystal Methamphetamine	66 (60)
Crystal Methamphetamine and heroin together	39 (33)
Crystal Methamphetamine and cocaine together	1 (1)
Other ^I	16 (13)
Number of drug types used in past six months	
One	72 (62)
Two or more	41 (35)
Not reported	4 (3)
Injected drugs in the past six months	
No	9 (11)
Yes	70 (88)
Currently need help controlling drug use	
Don't require help	38 (32)
Require some help	31 (27)
In great need of help	34 (29)
Urgently need help	14 (12)

^I Includes hallucinogens, inhalants, prescription opioids and benzodiazepines

Multivariable logistic regression models examining associations between integration and recent drug use among deported migrants attending a free health clinic in Tijuana, Mexico, 2013

Table 4.

Variable	Model 1 Means & Markers N=250	Model 2 Social Connection N=246	Model 3 Facilitators N=251	Model 4 Foundation N=255
Worked in formal economy last 6m	1.54 (0.87 – 2.73)			
Sought work in Tijuana last 6m	0.38 (0.21 – 0.69)**			
Experienced barriers to obtaining work last 6m	0.81 (0.44 – 1.48)			
Denied a job in Tijuana last 6m	1.19 (0.63 – 2.25)			
Mostly lived in unstable accommodation last 6m	0.87 (0.48 – 1.56)			
Household affluence score ²	0.77 (0.68 – 0.87)**			
Denied housing/shelter in Tijuana last 6m	0.93 (0.42 – 2.03)			
Good/very good/excellent self-rated health	0.82 (0.47 – 1.43)			
Depressive symptoms (15 on NIH PROMIS scale)	1.53 (0.87 – 2.68)			
Has family or friends in Tijuana	1.86 (1.06 – 3.26)*			
Separated from family due to deportation	1.53 (0.83 – 2.83)			
Banned from returning to USA	1.74 (0.95 – 3.16)			
Felt rejected in Tijuana last 6m	1.14 (0.66 – 1.98)			
Visited a doctor in the last 6m	0.41 (0.24 – 0.71)**			
Low Spanish proficiency			1.71 (0.21 – 13.87)	
Identify most strongly with USA			0.68 (0.36 – 1.30)	
Threatened/harassed in Tijuana last 6m			1.04 (0.52 – 2.10)	
Threatened/harassed by police/military in Tijuana			1.31 (0.74 – 2.31)	
Have a visible tattoo			2.24 (1.30 – 3.86)**	
Ever incarcerated, in Mexico only			3.90 (1.71 – 8.92)**	
Ever incarcerated, in US only			2.02 (0.86 – 4.65)	
Ever incarcerated, in both Mexico and US			6.01 (2.49 – 14.52)**	
Has any form of Mexican ID				0.77 (0.44 – 1.35)
Has any form of health insurance				0.39 (0.22 – 0.71)**

Model statistics

Variable	Model 1 Means & Markers N=250	Model 2 Social Connection N=246	Model 3 Facilitators N=251	Model 4 Foundation N=255
Model significance p-value	<0.001	0.001	<0.001	<0.001
Hosmer & Lemeshow goodness-of-fit p-value	0.304	0.619	0.576	0.562
* p<0.05				
** p<0.01				

Table 5.

Multivariate logistic regression model identifying individual integration variables significantly associated with recent drug use among deported migrants attending a free health clinic in Tijuana, Mexico, 2013

Integration domain and variable	Adjusted Odds Ratio ^{1,2,3} (95% CI)	p-value
<i>Means & Markers</i>		
Sought work in Tijuana last 6m	0.37 (0.18 – 0.77)	0.008
Household affluence score ²	0.82 (0.69 – 0.96)	0.025
<i>Social Connection</i>		
Has family or friends in Tijuana	1.04 (0.46 – 2.37)	0.922
Visited a doctor in the last 6m	0.65 (0.26 – 1.37)	0.255
<i>Facilitators</i>		
Threatened/harassed by police/military in Tijuana	1.18 (0.56 – 2.49)	0.667
Have a visible tattoo	1.65 (0.80 – 3.40)	0.171
Ever incarcerated, in Mexico only	2.04 (0.68 – 6.07)	0.201
Ever incarcerated, in US only	2.66 (0.92–7.68)	0.070
Ever incarcerated, in both Mexico and US	3.30 (1.09 –9.96)	0.034
<i>Foundation</i>		
Has any form of health insurance	0.37 (0.17 – 0.78)	0.009

¹ Adjusted for age, sex, birth state, education, total time spent in the US, number of deportations, time since last deportation and duration of residence in Tijuana

² N=229 (missing data excluded)

³ Hosmer-Lemeshow goodness of fit p=0.062