Published in final edited form as:

J Hepatol. 2019 June ; 70(6): 1214–1221. doi:10.1016/j.jhep.2019.01.036.

Clinical Utility of Genomic Analysis in Adults with Idiopathic Liver Disease

Aaron Hakim¹, Xuchen Zhang², Angela DeLisle¹, Elif A. Oral³, Daniel Dykas⁴, Kaela Drzewiecki¹, David N. Assis¹, Marina Silveira¹, Jennifer Batisti¹, Dhanpat Jain², Allen Bale⁴, Pramod K. Mistry^{1,5,#}, and Silvia Vilarinho^{1,2,#,*}

¹Department of Internal Medicine, Section of Digestive Diseases, Yale University School of Medicine, New Haven, CT.

²Department of Pathology, Yale University School of Medicine, New Haven, CT.

³Department of Internal Medicine, Division of Metabolism, Endocrinology and Diabetes, University of Michigan, MI.

⁴Department of Genetics, Yale University School of Medicine, New Haven, CT.

⁵Department of Pediatrics and of Cellular & Molecular Physiology, Yale University School of Medicine, New Haven, CT.

Abstract

Background & Aims: Adult patients suffering from liver disease of unknown cause represent an understudied and underserved population. The use of whole-exome sequencing (WES) for the study of a broader spectrum of non-oncological diseases, among adults, remains poorly studied. We assessed the utility of WES in diagnosis and management of adults with unexplained liver disease despite comprehensive evaluation by a hepatologist and with no history of alcohol overuse.

Methods: We performed WES and deep phenotyping of nineteen unrelated adult patients with idiopathic liver disease recruited at a tertiary academic health care center.

Results: Analysis of the exome in 19 cases identified four monogenic disorders in five unrelated adults. Patient 1 suffered for 18 years from devastating complications of undiagnosed Type 3 Familial Partial Lipodystrophy due to a deleterious heterozygous variant in *PPARG*. Molecular diagnosis enabled initiation of leptin replacement therapy with subsequent normalization of liver

*Correspondence should be address to Silvia Vilarinho, M.D., Ph.D., Departments of Internal Medicine (Digestive Diseases) and of Pathology, Yale University School of Medicine, 333 Cedar Street, LMP1080, New Haven, CT 06510, USA. Telephone: +1-203-737-6063, Fax: +1-203-737-1755, silvia.vilarinho@yale.edu.

Author Contributions: P.K.M. and S.V. developed study concept and design; A.H., X.Z., D.D., K.D., D.J., A.B., S.V. performed research and/or analyzed data; A.D., E.O., D.A., M.S., J.B., D.J., P.K.M., S.V. participated in patient recruitment and/or patient's ascertainment and management; A.H., P.K.M. and S.V. wrote the manuscript and all authors critically revised the manuscript draft.

#Co-senior authors

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflicts of interest: E.O. received grant support and served as a consultant to Aegerion Pharmaceuticals. A.H. reports being previously employed by Great Point Partners, a healthcare investment company. The remaining authors have no conflicts of interest.

transaminases, amelioration of dyslipidemia, and decrease in daily insulin requirements. Patients 2 and 3 were diagnosed with *MDR3* deficiency due to recessive mutations in *ABCB4*. Patient 4 with a prior diagnosis of non-alcoholic steatohepatitis was found to harbor a mitochondrial disorder due to a homozygous pathogenic variant in *NDUFB3*; this finding enabled initiation of disease preventive measures including supplementation with antioxidants. Patient 5 is a lean patient with hepatic steatosis of unknown etiology who was found to have a damaging heterozygous variant in *APOB*.

Conclusions: Genomic analysis yielded an actionable diagnosis in a substantial number (~25%) of selected adult patients with chronic liver disease of unknown etiology. This study supports the use of WES in the evaluation and management of adults with idiopathic liver disease in clinical practice.

LAY SUMMARY

We performed whole exome sequencing in nineteen adult patients with unexplained liver disease after an unrevealing conventional work-up performed by a hepatologist. In five cases, genomic analysis led to a diagnosis and informed treatment and management of the disease.

Keywords

undiagnosed liver disease; whole-exome sequencing; precision medicine; genetic diagnosis; germline mutations

INTRODUCTION

Chronic liver disease (CLD) is a significant health problem affecting more than 4 million people in the United States and leading to over 40,000 deaths annually [1]. CLD is often undiagnosed for many years unless there is awareness of subtle clinical signs, behavioral risk factors and/or investigation of abnormal liver function tests. In many patients, by the time overt manifestations of CLD emerge, liver injury has advanced to result in portal hypertension or hepatic decompensation. The taxonomy of CLD in clinical practice is based broadly on categories of etiology such as exposure to toxins, viral infections, cholestatic, autoimmune, metabolic and select genetic disorders. A significant limitation of this approach is that it precludes consideration of a wider array of underlying genetic disorders masquerading within these broad phenotypes. Additionally, it is estimated that up to 30% of cases of cirrhosis and up to 14% of adults awaiting liver transplantation suffer from liver disease of unknown etiology [2, 3]. These patients often undergo a long and costly odyssey of diagnostic tests, interventions, inappropriate therapies and medical opinions.

Understanding the etiology of CLD may be essential to halt progression of liver dysfunction, as illustrated by the development of a vaccine and anti-viral therapy for hepatitis B, and the highly effective, safe and curative anti-viral therapies for hepatitis C [4]. Advances in human genetics and genomics have created an unprecedented opportunity for gene discovery and diagnosis in the clinic. Specifically, whole-exome sequencing (WES), which consists of sequencing all the ~20,000 human protein-coding genes, currently represents a remarkable balance between cost, time of analysis and information collected, making it attractive and suitable for clinical use and translational research studies. In pediatric cohorts, we [5–7] and

others [8–10] have shown that WES combined with deep clinical phenotyping is an effective and unbiased means to identify rare protein-altering coding variants in individual genes. However, to date, most studies that investigate the use of next generation sequencing technologies in diagnosis and individualization of medical care have been performed in either pediatric or cancer patients. There is paucity of information on the clinical utility of these approaches for a broader spectrum of diseases among adults, such as CLD [11–13]. By using unbiased genomic analysis we also begin to understand parameters of adult clinical presentations that harbor an underlying monogenic cause, and to develop a more comprehensive category of ‘genetic’ liver diseases in adults beyond the traditionally considered disorders such as Wilson’s disease or hemochromatosis. Here, we provide data to support the utility of WES in the diagnosis and management of adults with liver disease of unknown cause with or without involvement of other diseases and/or unusual clinical findings.

PATIENTS and METHODS

Human Subjects

Study protocol was approved by the Yale Human Investigation Committee, and informed consent was obtained in accordance with institutional review board standards. Adults with unexplained liver disease despite a comprehensive evaluation (unrevealing hepatitis viral serologies - including negative HBsAg and anti-HBc, ferritin, iron studies, ceruloplasmin, ANA, alpha-1-antitrypsin phenotype, abdominal imaging, liver biopsy) underwent further investigation using whole-exome sequencing. In some cases we questioned prior diagnosis such as non-alcoholic fatty liver disease (NAFLD) in absence of typical metabolic or body habitus features. Where possible, samples from available family members were also obtained for segregation studies.

DNA isolation, exome capture and sequencing

Genomic DNA was isolated from peripheral blood leucocytes or buccal swabs using standard procedures. DNA fragments containing exonic sequences were captured and sequenced on the Illumina HiSeq platform.

Exome Sequencing Analysis

Exome sequencing data were mapped and aligned to the build 19 of the reference human genome using BWA. Variants were called using GATK [14, 15] and annotated using Annovar [16]. Variants were selected for minor allele frequency <0.01 for homozygous and compound heterozygous variants (recessive inheritance pattern) or $<2 \times 10^{-5}$ for heterozygous variants (dominant inheritance pattern). Allele frequencies were determined using the genome aggregation database (gnomAD) [17], including the Exome Aggregation Consortium database (ExAC), 1000 Genomes, and the National Heart, Lung and Blood Institute’s (NHLBI) Exome Variant Server. Subsequently, protein-altering variants were selected and prioritized based on their predicted deleteriousness. MetaSVM [18] was used to infer the impact of missense variants. Rare protein-altering variants predicted to be deleterious were then selected if they occurred as pathogenic variants described in NCBI

Clin Var, and/or in genes previously associated with liver-related diseases listed in the Online Mendelian Inheritance in Man (OMIM) database (Supplementary Figure 1).

Principal Component Analysis

Principal component analysis (PCA) was performed to determine the ancestry of the patients in our cohort. All tag SNP genotypes were obtained from WES data and used as inputs, along with the same SNPs from subjects in the HapMap project, to perform PCA with EIGENSTRAT software [19].

Sanger Sequencing

Sanger sequencing of the identified *PPARG* variant (p.Gly161Val) in patient 1 was performed by PCR amplification of genomic DNA of the proband and her parents. Sanger sequencing of the identified *ABCB4* variants (p.Arg549Cys and p.Ala934Thr) in patient 2 was performed by PCR amplification of genomic DNA of the proband, her mother and her son. Sanger sequencing of the identified *ABCB4* variant (p.Ter1280Arg) in patient 3 was performed by PCR amplification of genomic DNA of the proband. Sanger sequencing of the identified *NDFUB3* variant (p.Trp22Arg) in patient 4 was performed by PCR amplification of genomic DNA of the proband and his parents. Sanger sequencing of the heterozygous splice-site variant (c.2067+1G>A) in *APOB* in patient 5 was confirmed by PCR amplification of genomic DNA of the proband. Forward and reverse primers for each variant are described in Supplementary Table 1.

WES-based Copy Number Variants (CNVs) detection

A proprietary program using exome capture target region read depth normalized to read depth for each chromosome was used to assess CNVs. The normalized sequence depths for a sample batch of four or greater samples are used to stratify target regions by variance where those regions significantly vary from the batch means. The use of data from a batch of samples allows for the establishment of a baseline reference depth range for each target region. This methodology can detect deletions and duplications of two or more adjacent exons with close to 100% sensitivity and specificity. Single-exon deletions and duplications are flagged, but require confirmation by a second method due to a significant false positive rate. Validation of the method was performed using samples tested by standard, high-resolution array comparative genomic hybridization and exon-array comparative genomic hybridization or PCR.

Orthologues

Full-length orthologous protein sequences from both vertebrate and invertebrates were obtained from GenBank. Protein sequences were aligned using the ClustalW or Clustal Omega algorithm.

RESULTS

Study population characteristics and whole-exome sequencing

Nineteen adults with unexplained liver disease and no history of alcohol overuse were recruited from Yale New Haven Health after an unrevealing conventional work-up performed by a hepatologist. These individuals presented between the ages of 22 and 73 years-old with a variety of liver disorders (Table 1) with or without other comorbidities. We performed individual whole-exome sequencing of germ line DNA isolated from each patient. Targeted bases were sequenced by a mean of 90 reads, with 94% of targeted bases having more than eight independent reads, and 92% having more than fifteen independent reads, conferring high confidence calling of homozygous and heterozygous variants across the exome (Supplementary Table 2). Using unbiased WES, we identified a monogenic disorder in five patients of this adult population cohort, gaining insight into their liver disease pathogenesis and with direct impact on clinical management (Table 2). Fourteen patients remain unexplained after genomic analysis (Supplementary Table 3), including WES-based CNV detection.

Exome sequencing yields a diagnosis and initiation of therapy in a patient who suffered from devastating complications of undiagnosed Familial Partial Lipodystrophy Type 3 for 18 years

Patient 1 is a 33 year-old female with biopsy proven severe (80–90%) hepatic macrovesicular steatosis with periportal and pericentral fibrosis (Figure 1A, B). There was moderate portal inflammation with occasional hepatocyte ballooning, rare poorly- formed Mallory-Denk bodies, ceroid laden macrophages and marked Kupffer cell siderosis. Her past medical history is significant for early onset hyperlipidemia diagnosed in childhood, recurrent episodes of hypertriglyceridemia-induced pancreatitis eventually leading to total pancreatectomy, splenectomy and insulin dependence. She also had history of hypertension and of pre-eclampsia at the age of 29. Her social and family history is non-contributory. She had been seen and evaluated by many expert pediatric and adult physicians at several U.S. tertiary medical centers within the last 18 years and despite a comprehensive work-up, her operational diagnosis was hyperchylomicronemia syndrome although genetic deficiency of lipoprotein lipase or apolipoprotein CM could not be demonstrated.

We performed WES of germ line DNA to investigate a possible underlying genetic defect. Since her biological parents were unaffected, we analyzed her exome data considering both a recessive as well as a dominant pattern of inheritance. Consistent with an unrelated union, no rare homozygous genotypes were observed in the proband. However, she harbored one missense variant (chr3:12434114, G>T, NM_015869, c.482G>T, p.Gly161Val) in *PPARG*, which encodes peroxisome proliferator-activated receptor γ and heterozygous pathogenic variants in this gene have been related to autosomal dominant Familial Partial Lipodystrophy Type 3 (FPLD3). This variant is predicted to be damaging by MetaSVM and it is absent among > 100,000 alleles in the gnomAD database, and therefore is likely to be pathogenic in this patient (Table 3). Sanger sequencing confirmed the heterozygous variant in the proband and further showed that neither parent harbors the variant, revealing that it occurred *de novo* in the patient (Figure 1C). This variant is located in the DNA binding domain of the *PPARG*

protein at a highly conserved position across orthologues (Figure 1D). Moreover, a single case of an older female with clinical features consistent with FPLD3 and harboring the same *PPARG* variant (p.Gly161Val) as patient 1 has been reported [20]. At this point, in light of new genotype knowledge and presumed diagnosis, we reevaluated her clinical and laboratory findings, which were consistent with FPLD3.

PPARG is known to be a key transcription factor in adipocyte differentiation, which explains the lack of adipose tissue in the patient and the striking accumulation of triglycerides in the bloodstream as well as its accumulation in other organs, such as the liver. Given the diagnosis of FPLD3 and decreased overall adiposity, we postulated that patient's leptin levels should be low since leptin is mainly produced by adipocytes. As predicted, her leptin level was significantly low at 1.5 ng/mL (normal range for age and gender = 8.0–38.9 ng/mL) (Figure 1E). This finding not only supported our genetic diagnosis, but led to a new therapeutic intervention. The patient was initiated on leptin replacement therapy in a named compassionate use program. During the ensuing 13 months there was significant amelioration of dyslipidemia: total cholesterol fell from 238 to 130 mg/dL, triglyceride levels fell from 3532 to 267 mg/dL, and HDL cholesterol increased from 8 to 24 mg/dL (Figure 1E). Concomitantly, there was normalization of her liver transaminases: AST and ALT levels decreased from 114 to 19 U/L and from 94 to 17U/L, respectively (Figure 1F). Furthermore, the patient's daily insulin requirements decreased by approximately half compared to doses of insulin required prior to leptin replacement therapy.

MDR3 deficiency diagnosed in adulthood in two unrelated patients

Patient 2 is a 31-year-old female who presented with an acute esophageal variceal hemorrhage in her 22nd week of her second gestation. Of note, the patient had symptomatic cholestasis during her first pregnancy and she was diagnosed with benign intrahepatic cholestasis of pregnancy at that time. On this presentation, she had severe cholestasis (with high γ -glutamyltransferase, GGT) and liver biopsy revealed cirrhosis and cholestasis with ductular proliferation (Figure 2A, B). WES revealed two rare variants predicted to be damaging by MetaSVM (chr7:87069069, C>T, NM_000443, c.1645C>T, p.Arg549Cys and chr7:87041333, G>A, NM_000443, c.2800G>A, p.Ala934Thr) in *ABCB4*, which encodes ATP binding cassette subfamily B member 4, also known as multidrug resistance protein 3 (MDR3) (Table 3). There were no other disease-causing variants in genes known to be associated with liver disease, using recessive or dominant models. Sanger sequencing confirmed both variants in the patient's genomic DNA. Study of available family members revealed that her mother is a heterozygous carrier of p.Arg549Cys whereas her son is heterozygous for the other variant, p.Ala934Thr, supporting that the two variants found in our patient were located *in trans* at the *ABCB4* gene locus (Figure 2C, D), consistent with an autosomal recessive disorder. Both variants are conserved across orthologues (Figure 2E, F). As expected, her total bile acids (cholic acid, deoxycholic acid and chenodeoxycholic acid)

levels have been persistently elevated, ranging from 36 to 105 $\mu\text{mol/L}$ (normal is $<6.8 \mu\text{mol/L}$).

Patient 3 is a 29 year-old male with past medical history significant for liver transplantation at 8 years of age for cirrhosis of unclear etiology who presented with chronic graft rejection in the setting of medical non-adherence. Consistent with known consanguinity, patient 3 harbors twenty-seven rare homozygous variants. Twenty-six of these variants are missense variants in genes unrelated to liver disease by OMIM, and therefore are unlikely to be causal pathogenic variants in this patient. The other homozygous variant encoded a stop lost in *ABCB4* (chr7:87031414, T>A, NM_000443, c.3838T>A, p.Ter1280Arg), which was confirmed by Sanger sequencing (data not shown). This variant has never been reported in general population by gnomAD. However, another substitution in the same nucleotide (NM_000443 c.3838T>C) resulting in identical stop-loss variant in the p.Ter1280 codon has been previously associated with cholestasis and liver failure [21]. Both the c.3838T>A and c.3838T>C nucleotide substitutions result in the extension of the protein by 19 new amino acids. Given new knowledge of the patient's genotype, we reviewed the patient's liver explant. Liver parenchyma showed cirrhosis with patchy macrovesicular steatosis, portal/septal chronic inflammation, marked cholestasis with ductular proliferation and increased hepatocytic copper deposition (Figure 2G, H, I, J), consistent with a genetic diagnosis of MDR3 deficiency.

Recessive variant in NDFUB3 in a patient with elevated transaminases, hepatic steatosis, recurrent avascular necrosis and short stature

Patient 4 is a 32 year-old male with persistent elevation of transaminases (AST and ALT ~3 times upper limit of normal), recurrent avascular necrosis, and short stature (with both parents' being of average height). Liver biopsy showed minimal macrovesicular (small and large droplet fat) steatosis ($<5\%$) (Figure 3A, B). He is the single child of unrelated European-descended parents. WES was remarkable for a very rare homozygous variant in *NDUFB3* (chr2:201943669, T>C, NM_002491, c.64T>C, p.Trp22Arg), which encodes for NADH-dehydrogenase 1 beta complex 3 and consists of the first enzyme in the electron transport chain in mitochondria (Table 3). Both parents were found to be heterozygous for this variant by Sanger sequencing (data not shown). This variant is predicted to be damaging by MetaSVM and it was first reported in a single female infant with lethal complex I mitochondrial deficiency [22] and more recently in 10 children from 8 families, seven of them of Irish ancestry [23]. In this last cohort, all patients have short stature ($<9^{\text{th}}$ percentile) and similar facial dysmorphic features to patient 4, such as a prominent forehead, smooth philtrum and deep-set eyes (Supplementary Figure 2). In contrast to the first patient reported with this homozygous variant, they have a good long-term prognosis, even though some patients presented with an acute metabolic crisis with evidence of an isolated complex I deficiency in muscle [23]. In light of new genotype information, our patient 4 had mitochondrial ETC testing in skeletal muscle biopsy that revealed a deficiency of rotenone sensitive I+III activity despite normal citrate synthase activity, which fulfills a minor criterion of the modified Walker criteria for diagnosis of a respiratory chain disorder ($<30\%$). Liver electron microscopic findings were also suggestive of a mitochondrial abnormality, with hepatocytes showing different sized lipid droplets (Figure 3C). Moreover,

the patient suffers from oculomotor dysfunction with optic nerve anomalies, episodes of lactic acidosis during surgical interventions, and progressive fatigue. He was started on mitochondrial cocktail supplement, including co-enzyme Q10, vitamins B2 and B6.

Lean patient with hepatic steatosis of unknown etiology was found to have a novel damaging heterozygous variant in APOB

Patient 5 is a 23 year-old lean male who presented for evaluation of persistent elevated ALT (2 to 3 times upper limit of normal) for which he underwent a liver biopsy that showed mild macrovesicular steatosis (30%) with minimal lobular inflammation and minimal pericentral sinusoidal fibrosis (Figure 3D). His work-up was also remarkable for a ceruloplasmin of 17mg/dL (normal range =18–36mg/dL). WES analysis revealed no rare variations in *ATP7B* and a pathogenic heterozygous splicing variant in *APOB* (chr2:21250699, c.2067+1G>A), which encodes apolipoprotein B (ApoB). This change was validated by Sanger sequencing (data not shown) and is predicted to abrogate the normal splicing of exon 14 in the *APOB* gene. Heterozygous carriers typically have decreased plasma levels of LDL-cholesterol and ApoB, and may be asymptomatic or have clinical manifestations such as liver steatosis [24, 25]. Consistent with the patient's genotype, his circulating ApoB level was found to be half the lower limit of normal (26 mg/dL compared to normal range of 52 to 109 mg/dL), with low circulating LDL cholesterol (20mg/dL) and triglycerides (19 mg/dL) levels. Thus, this new genotype information explains both clinical and laboratory findings in this patient, and may have implications in his clinical management beyond family counseling. Specifically, fat-soluble vitamin E supplementation has been recommended for patients with homozygous hypobetalipoproteinemia to protect from neurological complications, and modest doses of Vitamin E have been proposed for treatment of heterozygous patients [26], but further studies are required.

DISCUSSION

This study provides evidence that a subset of adult patients who suffer from liver disease of indeterminate etiology with or without other co-morbidities harbor an underlying Mendelian disorder, which may be unrecognized during their entire childhood until genetic testing is performed. These findings have several implications. First, by establishing a diagnosis for a substantial number of undiagnosed cases, we provided new insights into disease pathogenesis. Second, knowledge of genotype led to recognition of unappreciated phenotypic features as well as enablement of new therapeutic and preventive medicine interventions beyond family counseling. For example, genomic analysis in patient 1 led to recognition of phenotypic aspects unappreciated by standard clinical examination by specialists not familiar with the FPLD3 phenotype. It also led to initiation of leptin replacement therapy with striking amelioration of metabolic dysregulation and liver disease. Correct diagnosis additionally allowed appropriate attention on monitoring and prevention of premature coronary artery disease and other cardiovascular risk factors. Third, our data highlight the importance of using WES in the investigation of liver disease of unknown cause so that we may start developing an understanding of what clinical presentations/diseases are genetic and may remain undiagnosed until adulthood. The genetic diagnosis of MDR3 deficiency in patient 2 underscores the silent progression of inherited chronic liver

disease to cirrhosis, portal hypertension and decompensation, remaining unrecognized for decades and with first presentation in adulthood [27]. Fourth, reminiscent of the widely accepted Radiology and Pathology Rounds in clinical practice, this study illustrates the potential clinical value of Genome Rounds in the individual assessment and medical care of adults suffering from liver disease of unknown cause (Figure 4). This approach perfectly exemplifies the mission of the Precision Medicine Initiative [28] launched by U.S. President Barack Obama in January 2015. Physicians recognize that every patient is unique and have always tried to adjust their interventions as best they can to each individual. Now, we start having the technology and knowledge to translate this concept to routine clinical practice in tertiary medical centers across the world. Even in the contemporary taxonomy of liver diseases, there is little understanding of the heterogeneity of disease within each category and distinct subtypes based on their underlying genetic and/or pathobiology. This concept is illustrated by the three cases (patients 1, 4 and 5) who harbor distinct genetic defects affecting different molecular pathways leading to hepatic steatosis and presumed diagnosis of NAFLD, with direct implications in bedside therapeutic and preventive medicine interventions.

In this cohort, most of the diagnosed patients had seen multiple physicians from a diverse array of medical and surgical specialists for several years prior to their diagnosis, such as primary care providers, internists, surgeons, endocrinologists and hepatologists, among others. This suggests that the investigation of unrecognized genetic disorders in adults would have clinical utility among a broad group of adult multispecialty clinical practices. Decades ago, Mendelian genetics mostly relied on family-based studies with very distinct and often severe phenotype(s). However, as illustrated in this study, the absence of family history of similar phenotype should not deter physicians to investigate a genetic cause for the unexplained liver disease since it might arise from a *de novo* variant, which by definition is not inherited from any parent, or result in a recessive inheritance pattern for which both parents are usually healthy carriers and 75% of siblings will be clinically unaffected.

One limitation of this study is a relatively small sample size, and patient recruitment at a tertiary care academic center. Further studies are required to assess the generalizability of these findings in a broader liver disease population. Additionally, the patients in this cohort whose phenotype remains unexplained may have a pathogenic variant not detected by the methodology used, such as variants in the non-coding region of the genome, or in a gene not yet known to be associated with a human disease. In fact, approximately three quarters of human genes have not yet been linked to a human phenotype [29], and for this reason we will continue to re-analyze these patients' WES data regularly. This study's diagnostic yield is comparable to data recently reported in inherited cardiovascular diseases and chronic kidney disease in adults [11, 12].

Collectively, our data support the incorporation of WES in the diagnostic and management algorithms of adults suffering from idiopathic liver disease despite a comprehensive work-up, and underscores its value as a means of developing an understanding of what liver phenotypes of unknown cause in adults with or without involvement of other diseases are genetic. A multidisciplinary Genome Rounds approach (Figure 4) will likely create the basis to develop best practice guidelines for genomic medicine in a variety of non-oncological

medical and surgical specialties, including hepatology. This strategy will shed further light on genetic contributions, and therefore underlying molecular pathogenesis, across different forms of liver disease that are clinically indistinguishable through conventional diagnostic approaches.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

We thank all the patients and their families whose contribution to this study lead to advancing our understanding of liver disease, Dr. Michael Nathanson for his efforts to refer patients to this study and the staff of the Yale Center for Genome Analysis for exome sequencing. Research reported in this publication was supported by the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health under Award Number K08DK113109. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. This work was also supported in part by Yale Liver Center P30DK034989, and AASLD Sheila Sherlock Clinical and Translational Research Award in Liver Disease (to S.V.).

REFERENCES

- [1]. Murphy SL, Xu J, Kochanek KD, Curtin SC, Arias E. Deaths: Final Data for 2015. *Natl Vital Stat Rep* 2017;66:1–75.
- [2]. Czaja AJ. Cryptogenic chronic hepatitis and its changing guise in adults. *Dig Dis Sci* 2011;56:3421–3438. [PubMed: 21647651]
- [3]. Rinaldi L, Nascimbeni F, Giordano M, Masetti C, Guerrero B, Amelia A, et al. Clinical features and natural history of cryptogenic cirrhosis compared to hepatitis C virus-related cirrhosis. *World J Gastroenterol* 2017;23:1458–1468. [PubMed: 28293093]
- [4]. Vilarinho S, Lifton RP. Pioneering a Global Cure for Chronic Hepatitis C Virus Infection. *Cell* 2016;167:12–15. [PubMed: 27634325]
- [5]. Vilarinho S, Choi M, Jain D, Malhotra A, Kulkarni S, Pashankar D, et al. Individual exome analysis in diagnosis and management of paediatric liver failure of indeterminate aetiology. *J Hepatol* 2014;61:1056–1063. [PubMed: 25016221]
- [6]. Vilarinho S, Erson-Omay EZ, Harmanci AS, Morotti R, Carrion-Grant G, Baranoski J, et al. Paediatric hepatocellular carcinoma due to somatic CTNNB1 and NFE2L2 mutations in the setting of inherited bi-allelic ABCB11 mutations. *J Hepatol* 2014;61:1178–1183. [PubMed: 25016225]
- [7]. Vilarinho S, Sari S, Mazzacuvu F, Bilguvar K, Esendagli-Yilmaz G, Jain D, et al. ACOX2 deficiency: A disorder of bile acid synthesis with transaminase elevation, liver fibrosis, ataxia, and cognitive impairment. *P Natl Acad Sci USA* 2016;113:11289–11293.
- [8]. Yang Y, Muzny DM, Reid JG, Bainbridge MN, Willis A, Ward PA, et al. Clinical whole-exome sequencing for the diagnosis of mendelian disorders. *N Engl J Med* 2013;369:1502–1511. [PubMed: 24088041]
- [9]. Lee H, Deignan JL, Dorrani N, Strom SP, Kantarci S, Quintero-Rivera F, et al. Clinical exome sequencing for genetic identification of rare Mendelian disorders. *JAMA* 2014;312:1880–1887. [PubMed: 25326637]
- [10]. Tarailo-Graovac M, Shyr C, Ross CJ, Horvath GA, Salvarinova R, Ye XC, et al. Exome Sequencing and the Management of Neurometabolic Disorders. *N Engl J Med* 2016;374:2246–2255. [PubMed: 27276562]
- [11]. Seidemann SB, Smith E, Subrahmanyam L, Dykas D, Abou Ziki MD, Azari B, et al. Application of Whole Exome Sequencing in the Clinical Diagnosis and Management of Inherited Cardiovascular Diseases in Adults. *Circ Cardiovasc Genet* 2017;10.

- [12]. Lata S, Marasa M, Li Y, Fasel DA, Groopman E, Jobanputra V, et al. Whole-Exome Sequencing in Adults With Chronic Kidney Disease: A Pilot Study. *Ann Intern Med* 2018;168:100–109. [PubMed: 29204651]
- [13]. Splinter K, Adams DR, Bacino CA, Bellen HJ, Bernstein JA, Cheatle-Jarvela AM, et al. Effect of Genetic Diagnosis on Patients with Previously Undiagnosed Disease. *N Engl J Med* 2018.
- [14]. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytzsky A, et al. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res* 2010;20:1297–1303. [PubMed: 20644199]
- [15]. Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine A, et al. From FastQ data to high confidence variant calls: the Genome Analysis Toolkit best practices pipeline. *Curr Protoc Bioinformatics* 2013;43:11 10 11–33. [PubMed: 25431634]
- [16]. Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res* 2010;38:e164. [PubMed: 20601685]
- [17]. Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E, Fennell T, et al. Analysis of protein-coding genetic variation in 60,706 humans. *Nature* 2016;536:285–291. [PubMed: 27535533]
- [18]. Dong C, Wei P, Jian X, Gibbs R, Boerwinkle E, Wang K, et al. Comparison and integration of deleteriousness prediction methods for nonsynonymous SNVs in whole exome sequencing studies. *Human molecular genetics* 2015;24:2125–2137. [PubMed: 25552646]
- [19]. Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D. Principal components analysis corrects for stratification in genome-wide association studies. *Nat Genet* 2006;38:904–909. [PubMed: 16862161]
- [20]. Lau E, Carvalho D, Oliveira J, Fernandes S, Freitas P. Familial partial lipodystrophy type 3: a new mutation on the PPARG gene. *Hormones (Athens)* 2015;14:317–320. [PubMed: 26158656]
- [21]. Chinnaratha MA, Venugopal K, de Boer WB, Macquillan G. Novel mutation in progressive familial intrahepatic cholestasis III. *Liver Int* 2013;33:1293. [PubMed: 23574360]
- [22]. Calvo SE, Compton AG, Hershman SG, Lim SC, Lieber DS, Tucker EJ, et al. Molecular diagnosis of infantile mitochondrial disease with targeted next-generation sequencing. *Sci Transl Med* 2012;4:118ra110.
- [23]. Alston CL, Howard C, Olahova M, Hardy SA, He L, Murray PG, et al. A recurrent mitochondrial p.Trp22Arg NDUFB3 variant causes a distinctive facial appearance, short stature and a mild biochemical and clinical phenotype. *J Med Genet* 2016;53:634–641. [PubMed: 27091925]
- [24]. Rimbert A, Pichelin M, Lecointe S, Marrec M, Le Scouarnec S, Barrak E, et al. Identification of novel APOB mutations by targeted next-generation sequencing for the molecular diagnosis of familial hypobetalipoproteinemia. *Atherosclerosis* 2016;250:52–56. [PubMed: 27179706]
- [25]. Tarugi P, Lonardo A, Gabelli C, Sala F, Ballarini G, Cortella I, et al. Phenotypic expression of familial hypobetalipoproteinemia in three kindreds with mutations of apolipoprotein B gene. *J Lipid Res* 2001;42:1552–1561. [PubMed: 11590210]
- [26]. Sen D, Dagdelen S, Erbas T. Hepatosteatorosis with hypobetalipoproteinemia. *J Natl Med Assoc* 2007;99:284–286. [PubMed: 17393954]
- [27]. Schatz SB, Jungst C, Keitel-Anselmo V, Kubitz R, Becker C, Gerner P, et al. Phenotypic spectrum and diagnostic pitfalls of ABCB4 deficiency depending on age of onset. *Hepatol Commun* 2018;2:504–514. [PubMed: 29761167]
- [28]. Collins FS, Varmus H. A new initiative on precision medicine. *N Engl J Med* 2015;372:793–795. [PubMed: 25635347]
- [29]. Chong JX, Buckingham KJ, Jhangiani SN, Boehm C, Sobreira N, Smith JD, et al. The Genetic Basis of Mendelian Phenotypes: Discoveries, Challenges, and Opportunities. *Am J Hum Genet* 2015;97:199–215. [PubMed: 26166479]

HIGHLIGHTS

Whole exome sequencing led to a diagnosis in 5/19 cases of unexplained liver disease
The 5 cases represented 4 monogenic disorders diagnosed in adulthood Genomic analysis
informed the treatment and management of liver disease

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1.

Liver histology, genetic and laboratory findings in patient 1. (A) Liver parenchyma shows marked steatosis with moderate steatohepatitis (H&E stain, 40x). (B) Trichrome stain of liver biopsy tissue shows portal, periportal and perisinusoidal fibrosis consistent with stage 2 fibrosis (Brunt grading and staging system) (20x). (C) Pedigree depicts proband and unaffected subjects in black and white symbols, respectively, with Sanger sequencing chromatograms of the proband and her unaffected parents. *PPARG* alleles are denoted WT (wild-type) or Mut (p.Gly161Val). The *PPARG* p.Gly161Val variant is heterozygous in the

proband and absent in both parents. (D) Location of Gly-161 in *PPARG* and its conservation across species. The patient's variant is in a highly conserved DNA-binding domain (DBD) of the PPARG protein. Amino acid positions identical to the human reference are highlighted in yellow. N-terminal transactivation domain (AF1), highly conserved DNA-binding domain (DBD), C-terminal ligand-binding domain (LBD). (E) Triglycerides and leptin levels before (no shadow) and after (depicted by a gray shadow) initiation of leptin replacement therapy. (F) ALT/AST levels before (no shadow) and after (depicted by a gray shadow) initiation of leptin replacement therapy. LLN, lower limit of normal; ULN, upper limit of normal.

Figure 2.

Liver histology and genetic findings in patients 2 and 3. (A) Trichrome stain for patient 2 reveals fibrous septa with nodule formation consistent with cirrhosis (10x). (B) Cytokeratin 7 immunohistochemical staining (depicted in brown) for patient 2 shows ductular proliferation (10x). (C) Pedigree depicts patient 2 and unaffected subjects shown in black and white symbols, respectively. (D) Sanger sequencing chromatograms of the proband (patient 2) and her unaffected mother and son. The patient harbors two variants in *ABCB4*, p.Arg549Cys and p.Ala934Thr, whereas her mother and her son solely carry one of these variants each, p.Arg549Cys and p.Ala934Thr, respectively. (E, F) Conservation of Arg-549 and Ala-934 across species, respectively. Amino acid positions identical to the human reference are highlighted in yellow. (G) Liver parenchyma (H&E stain) from patient 3 showing cirrhosis (4x). (H, I) Liver parenchyma (H&E stain) from patient 3 at lower and higher magnification, respectively, revealing portal/septal chronic inflammation, ductular proliferation, and marked cholestasis (10x and 20x). (J) Copper stain of liver parenchyma for patient 3 supporting marked cholestasis (40x).

Figure 3.

Liver biopsies of patients 4 and 5. (A, B) Patient 4 liver parenchyma shows near normal histology with minimal macrovesicular steatosis (small and large droplet fat) (H&E stain) at lower (20x) and higher magnification (40x), respectively. (C) Electron microscopic findings for patient 4 is suggestive of a mitochondrial abnormality, with hepatocytes showing different sized lipid droplets (scale bar, 2 microns). (D) Hepatic parenchyma (H&E stain) from patient 5 shows predominantly macrovesicular steatosis with minimal steatohepatitis (10x).

Figure 4.

Schematic representation of multidisciplinary Genome Rounds in Adult Hepatology. It merges genotype-phenotype information with the goal of recognizing unappreciated phenotypic features by standard clinical examination, providing a diagnosis and new therapeutic options, and establishing adequate family counseling in adults. gDNA, genomic DNA.

Table 1:

Summary of study population characteristics and demographics.

Clinical Category	Patients, <i>n</i>	Mean age (range), <i>yo</i>	Male, <i>n</i>	Female, <i>n</i>	Alive <i>n</i>	Deceased or transplanted <i>n</i>
Cryptogenic cirrhosis	7	39 (29–70)	3	4	5	2
Non-obese NAFLD +/- NASH	6	30 (24–34)	4	2	6	0
Idiopathic cholestasis	4	46 (23–73)	3	1	3	1
HELLP and severe hyperammonemia	1	22	0	1	1	0
Idiopathic noncirrhotic portal hypertension	1	60	1	0	1	0
Total	19	38 (22–73)	11	8	16	3

n, number; HELLP, hemolysis, elevated liver enzymes and low platelet counts; *yo*, years-old

Table 2:

Demographics, clinical features and genetic diagnosis identified in five subjects of our cohort, and its clinical implications.

Patient ID	Age (yo)	Ethnicity	Sex	Presenting Features	Genetic Diagnosis	Clinical Implications
1	33	European	F	Lean NASH, hypertriglyceridemia, recurrent pancreatitis	Familial Partial Lipodystrophy Type 3	Initiation of leptin therapy; preventative cardiovascular measures; family counseling
2	32	African	F	Cryptogenic cirrhosis decompensated by esophageal variceal hemorrhage	MDR3 Deficiency	Family counseling; transplant candidacy
3	29	European	M	Cryptogenic cirrhosis at 8 years-old, now status post liver transplantation	MDR3 Deficiency	Family counseling; retransplant candidacy
4	32	European	M	Non-obese NAFLD, recurrent avascular necrosis, short stature	Mitochondrial complex I deficiency	Supplementation with co-enzyme Q10, vitamins B2 & B5; preventive interventions; family counseling
5	24	Asian	M	Lean NAFLD	Heterozygous Familial Hypobeta-lipoproteinemia	Family counseling; consideration for vitamin E supplementation

Ethnicity was determined by principal component analysis as described in Methods section; yo, years-old; F, female; M, male

Table 3:

Diagnostic genetic variants identified in five subjects in our cohort.

Patient ID	Gene	Inheritance and Effect of Variant	AA or cDNA change	Meta SVM score prediction	gnomAD (overall)	gnomAD (highest frequency)
1	<i>PPARG</i> (NM_015869)	Heterozygous, Missense	p.Gly161Val (c.482G>T)	1.104 (D)	0	0
2	<i>ABCB4</i> (NM_000443)	Compound Heterozygous, Missense	p.Arg549Cys (c.1645C>T)	0.448 (D)	0	0
			p.Ala934Thr (c.2800G>A)	0.676 (D)	1.3e ⁻³	1.4e ⁻² (African)
3	<i>ABCB4</i> (NM_000443)	Homozygous, Stop loss	p.Ter1280Arg (c.3838T>A)	N/A	0	0
4	<i>NDUFB3</i> (NM_002491)	Homozygous, Missense	p.Trp22Arg (c.64T>C)	0.485 (D)	8.2e ⁻⁴	1.3e ⁻³ (European)
5	<i>APOB</i> (NM_000384)	Heterozygous, Splice Site	c.2067+1G>A	N/A	0	0

AA, amino acid; gnomAD, Genome Aggregation Database (includes 123,136 exome and 15,496 whole-genome sequences); MAF, minor allele frequency; MetaSVM scores missense variants on a scale of -2 to 3 , with scores <0 predicted to be tolerated (T) and scores >0 predicted to be damaging (D); yo, years-old; N/A, not applicable.