Published in final edited form as:

J Pediatr. 2019 February ; 205: 35–40.e1. doi:10.1016/j.jpeds.2018.09.038.

Social Risk Factors Impact Hospital Readmission and Outpatient Appointment Adherence for Children with Congenital Heart Disease

Abigail C. Demianczyk, PhD¹, Shashank P. Behere, MD^{2,3}, Deepika Thacker, MD^{2,3}, Maia Noeder, PhD^{1,3}, Emily A. Delaplane, MSW⁴, Christian Pizarro, MD^{2,3}, and Erica Sood, PhD^{1,2,3}

¹Division of Behavioral Health, Nemours/Alfred I. duPont Hospital for Children, Wilmington, DE.

²Nemours Cardiac Center, Nemours/Alfred I. duPont Hospital for Children, Wilmington, DE.

³Department of Pediatrics, Sidney Kimmel Medical College, Thomas Jefferson University, Philadelphia, PA

⁴Department of Patient and Family Services, Nemours/Alfred I. duPont Hospital for Children, Wilmington, DE.

Abstract

Objective: To examine the relations of individual and cumulative social risk factors to hospitalization outcomes and adherence to outpatient cardiology appointments within the first two years of life for congenital heart disease (CHD) survivors.

Study design: Data were extracted for 219 patients who underwent infant cardiac surgery with cardiopulmonary bypass. Cumulative social risk was dichotomized into high social risk (2 risk factors; N=103) versus low social risk (1 risk factor; N=116). Risk of morbidity by procedure was assigned from 1–5 (STS-EACTS Morbidity Scores and Categories). Two-way ANOVAs examined effects of social risk and morbidity risk on length of first surgical hospitalization, number of readmissions and readmission days, subsequent cardiac surgical interventions, and adherence to outpatient cardiology appointments.

Results: An interaction between social risk and morbidity risk was identified for number of readmission days ($F(4,209)=3.07$, $P=.02$, $\eta^2=.06$). Pairwise comparisons demonstrated that among those patients with the lowest risk of morbidity by procedure (morbidity scores of 1 and 2), patients at high social risk had more readmission days than patients at low social risk (Morbidity

---

Corresponding Author: Erica Sood, PhD, Nemours Cardiac Center, Alfred I. duPont Hospital for Children, 1600 Rockland Road, Wilmington, DE 19803. Phone: 302-651-6304. Fax: 302-651-5345. Erica.Sood@nemours.org. No reprints requested.

The authors declare no conflicts of interest.

Portions of this study were presented at the 5th Annual Cardiac Neurodevelopmental Symposium, Philadelphia, PA, June, 2016; the 124th Annual Convention of the American Psychological Association, Denver, CO, August, 2016; and the 67th Annual Scientific Session and Expo of the American College of Cardiology, Orlando, FL, March, 2018.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Score 1: 16.63 ± 34.41 vs 3.02 ± 7.13 days; Morbidity Score 2: 27.68 ± 52.11 vs 2.20 ± 4.43 days). High social risk also predicted significantly worse adherence to cardiology appointments.

Conclusions: Cumulative social risk impacts readmission days for CHD patients with low risk of morbidity by procedure. Social risk assessment can identify families who may benefit from social/behavioral interventions to optimize discharge readiness, CHD home management, and long-term outcomes.

Congenital heart disease (CHD) is the most common birth defect, impacting 9 per 1,000 live births.¹ With decreasing mortality due to advances in surgical and perioperative care,^{2,3} the focus is shifting from *how long* children with CHD are living to *how well* they are living. Although cardiac surgery and hospitalization are medically necessary for many children with CHD,⁴ the burden of prolonged hospital stays and frequent readmissions can impact child and family quality of life,⁵ while also increasing costs.⁶ Prolonged hospital stays and readmissions additionally increase risk for hospital-acquired infection and negatively impact mid- and long-term neurodevelopmental outcomes.^{7,8} In recognition of these issues and in preparation for value-based care, cardiac centers are working to prevent delays in discharge and unplanned readmissions by identifying barriers and risk factors while optimizing discharge readiness and outpatient care.

Social determinants of health are well described in the literature. Studies have shown relationships between individual social risk factors (for example, race/ethnicity or socioeconomic status) and mortality among infants with CHD.^{9–14} Social risk factors also predict morbidity in adult cardiac patients,¹⁵ but few studies have investigated these relationships in a sample of young children with CHD. This study aimed to examine the relationships of individual and cumulative social risk factors to hospitalization data (length of first surgical hospitalization, number of readmissions and readmission days, and subsequent cardiac surgical interventions) and adherence to outpatient cardiology appointments within the first two years of life following cardiac surgery for CHD.

Methods

Retrospective chart reviews for sociodemographic and clinical characteristics were conducted for patients with CHD who underwent cardiac surgery with cardiopulmonary bypass in the first year of life at the Nemours Cardiac Center between January 2009 and September 2013 (N = 335); 116 of these patients were excluded from analyses due to having received cardiac outpatient care at an outside facility precluding evaluation of adherence to appointments and hospital readmissions (n = 69), mortality within the first two years of life (n = 31), heart transplantation (n = 6), discharge from hospital to a long-term care facility (n = 5), or lack of cardiac outpatient care in the first year of life without sufficient information to determine whether the patient had transferred care to another facility (n = 5), resulting in a total sample of 219 participants. Patients who experienced loss of care after the first year of life or long gaps in care were included in analyses. This retrospective study was reviewed and approved by the Nemours Institutional Review Board with waiver of the requirement for obtaining informed consent/parental permission.

Six social risk factors that have previously been shown to impact rates of mortality, morbidity, and/or access to and use of specialty care within CHD or other pediatric patient populations were examined.^{9–14,16–26} These include child race/ethnicity (minority = 1; non-minority = 0), regional poverty level using patient zip code (percentage of families living below the federal poverty level; 40% = 1; <40% = 0), primary medical insurance (Medicaid/CHIP = 1; private/military = 0), maternal age at time of birth (<21 = 1; 21 = 0), primary parental language (non-English = 1; English = 0), and distance to cardiac outpatient clinic (measured distance between hospital/satellite clinic and patient zip code; 25 miles = 1; < 25 miles = 0). A cumulative social risk variable ranging from 0 to 6 was calculated from number of social risk factors present. Participants were then stratified into high and low social risk groups based on the median number of social risk factors for this sample (2 social risk factors = 1, 47% of sample); 1 social risk factors = 0, 53% of sample).

Patients were stratified into groups based on the risk of morbidity associated with their initial and/or definitive cardiac surgery using the empiric Morbidity Scores and Categories proposed by Jacobs et al from the Society of Thoracic Surgeons Congenital Heart Surgery Database.²⁷ Surgical procedures were given a value between 1 and 5 based on the associated rate of major complications and the average postoperative length of stay, with higher scores indicating greater risk of morbidity. For patients who required pulmonary artery banding prior to definitive repair for an otherwise uncomplicated ventricular septal defect, the morbidity score for their definitive repair was recorded. For any other patient with more than one cardiac surgery in the first year of life, the morbidity score for their first surgery was counted.

Hospitalization data examined for this study were length of first surgical hospitalization, number of unplanned and total readmissions, total readmission days, and number of subsequent cardiac surgical interventions after the first surgery from birth to 24 months of age. Unplanned readmissions were defined as admissions to the cardiac floor or cardiac intensive care unit for any reason other than a scheduled procedure or surgery (eg, illness, cardiac specific complications). In-hospital monitoring after cardiac catheterization was not included as a readmission or part of hospital readmission days unless the hospitalization lasted longer than 24 hours. Admissions and surgeries clearly related to other medical conditions (e.g., cleft palate repair) were also not included.

Adherence to outpatient cardiology appointments was measured by calculating the average number of days between suggested and actual return dates for all cardiology appointments from birth to 12 months of age and 12 to 24 months of age. Patients who returned early or on the suggested return date were recorded as having zero days between actual and suggested return appointment. Days between suggested and actual return dates were not calculated or included in analyses if suggested return was not documented in the cardiology note or if patients were readmitted prior to suggested return. Higher values for this measure indicated poorer adherence.

Statistical Analyses

Independent sample *t*-tests were conducted to examine mean differences in hospitalization and outpatient appointment adherence for patients with and without each social risk factor.

To examine the effects of cumulative social risk and surgical morbidity risk, two-way analyses of variance (ANOVAs) were conducted for each hospitalization outcome and outpatient appointment adherence. Follow up analyses of simple main effects for social risk, as well as simple main effects and pairwise comparisons for surgical morbidity risk were conducted for any significant interactions.

Results

The number of patients meeting each social risk factor was as follows: racial/ethnic minority: $N = 89$ (40.6%); high regional poverty level: $N = 26$ (11.9%); primary Medicaid insurance: $N = 122$ (55.7%); maternal age less than 21 years at time of birth: $N = 15$ (6.8%); primarily non-English speaking: $N = 11$ (5.0%); living 25 miles or more from outpatient cardiac care: $N = 57$ (26.0%). High social risk was categorized as having two or more individual social risk factors, criteria met by 103 (47.0%) participants. Risk of morbidity by procedure was as follows: Morbidity score of One: $N = 76$ (34.7%); Two: $N = 43$ (19.6%); Three: $N = 45$ (20.5%); Four: $N = 37$ (16.9%); Five: $N = 18$ (8.2%). There were no significant differences between social risk groups for prevalence of premature birth, confirmed or suspected genetic abnormality, single ventricle anatomy, or age at first surgery (all p s $> .05$) (Table 1).

Means, standard deviations, and statistics for all independent sample t -tests are reported in Table 2 (available at www.jpeds.com). Child racial/ethnic minority status was associated with more unplanned readmissions in the first 2 years of life ($M = 1.58$, $SD = 2.12$ vs. $M = 1.04$, $SD = 1.89$), $t(216) = -1.98$, $p = .05$. Primary Medicaid insurance was associated with a greater number of hospital readmissions ($M = 2.14$, $SD = 2.49$ vs. $M = 1.51$, $SD = 2.15$), $t(216) = -1.99$, $p = 0.05$, and hospital readmission days ($M = 22.91$, $SD = 39.36$ vs. $M = 12.71$, $SD = 22.78$), $t(217) = -2.27$, $p = 0.02$, during the first two years of life. Child racial/ethnic minority status was associated with worse adherence to cardiac outpatient care in the first year of life ($M = 12.87$, $SD = 26.14$ vs. $M = 6.35$, $SD = 14.43$; $t(216) = -2.36$, $p = 0.02$), and primary Medicaid insurance ($M = 40.09$, $SD = 85.81$ vs. $M = 13.41$, $SD = 47.51$; $t(206) = -2.67$, $p = 0.01$) and living in a high poverty area ($M = 57.85$, $SD = 115.31$ vs. $M = 24.94$, $SD = 65.48$; $t(206) = -2.02$, $p = 0.04$) were associated with worse adherence to outpatient care in the second year of life.

A statistically significant interaction was found between social risk and surgical morbidity risk on total days spent in the hospital during readmissions in the first two years of life, $F(4, 209) = 3.07$, $p = 0.02$, partial $\eta^2 = 0.06$. Results are displayed in the Figure. A follow up analysis of the simple main effects of social risk at each level of surgical morbidity risk was performed. For individuals with a morbidity score of one ($F(1, 209) = 3.95$, $p < 0.05$, partial $\eta^2 = 0.02$) and two ($F(1, 209) = 7.17$, $p = 0.01$, partial $\eta^2 = 0.03$), the mean number of hospital readmission days was significantly higher for individuals with high social risk (Morbidity Score 1: 16.63 ± 34.41 days; Morbidity Score 2: 27.68 ± 52.11 days) than low social risk (Morbidity Score 1: 3.02 ± 7.13 days; Morbidity Score 2: 2.20 ± 4.43 days). The interaction between social risk and surgical morbidity risk trended towards significance for number of subsequent cardiac surgical interventions in the first two years of life ($p = 0.07$),

but power was limited due to a small sample size. There were no other significant interactions. (Results and estimated marginal means are reported in Table 3.)

Because interaction effects between social and morbidity risk were not significant for other hospitalization-related outcomes, analyses of main effects were performed. Statistically significant main effects of surgical morbidity risk were found for total length of stay during first surgical hospitalization ($F(4, 209) = 4.23, p = 0.03$, partial $\eta^2 = 0.08$), unplanned readmissions ($F(4, 208) = 2.68, p = 0.03$, partial $\eta^2 = 0.05$), total readmissions ($F(4, 209) = 7.49, p < 0.01$, partial $\eta^2 = 0.13$), and subsequent cardiac surgical interventions ($F(4, 209) = 31.27, p < 0.01$, partial $\eta^2 = 0.37$), such that higher surgical morbidity risk was associated with longer lengths of stay, more readmissions, and more cardiac surgical interventions. All pairwise comparisons were run, and the estimated marginal means and significant differences are reported in Table 4 for surgical morbidity and Table 4 for social risk. There were no significant main effects of social risk for hospitalization outcomes.

There were no significant interactions between social and morbidity risk for adherence to outpatient cardiology appointments, but there was a significant main effect of social risk. Specifically, a statistically significant main effect of social risk was found for average delay in outpatient care during the first year of life, $F(1, 208) = 4.43, p = 0.04$, partial $\eta^2 = 0.02$, such that high social risk was associated with poorer adherence to outpatient follow up compared with the low risk group. There were no other significant main effects of social risk, and no significant main effects of surgical morbidity risk. All results and estimated marginal means are reported in Table 4.

Discussion

Social determinants of health are well described in relation to CHD mortality,^{9–14} but far less explored with regard to morbidity, despite most CHD patients surviving through childhood and adolescence. This study highlights the importance of understanding the impact of both individual and cumulative social risk factors on a variety of CHD outcomes across the spectrum of care. Those patients at low risk of morbidity by procedure but high social risk had an average of 13–25 additional readmission days and thousands of dollars of added medical costs⁶ during the first 2 years of life compared with those with similarly low morbidity risk and low social risk. This finding is consistent with that of Lasa et al who demonstrated a higher prevalence of readmission and cardiac reintervention for minority pediatric patients with low surgical complexity CHD when compared with non-minority patients of similar surgical complexity.¹² Many hospital-based psychosocial supports and intervention programs focus on those patients considered to be at greatest risk for poor outcomes based on surgical morbidity and/or mortality risk (e.g., single ventricle, shunt dependent) and their families. The present study results highlight the need for increased psychosocial supports for patients and families with multiple social risk factors, even when undergoing procedures at low risk for surgical morbidity, as they may actually yield a greater impact on hospital readmissions while also substantially reducing health care costs.

The current results indicate that both individual and cumulative social risk factors are associated with poorer adherence to outpatient appointments during both the first and second

years of life regardless of risk of morbidity by procedure. Efforts to promote adherence to outpatient cardiac care after heart surgery for patients with social risk factors are crucial, as prior research has demonstrated that a history of one or more missed outpatient cardiology appointments is associated with a future loss to cardiac care later in life.³¹

When considered together, these results support the need for enhanced psychosocial supports for families of children with CHD to improve outcomes. These supports could include implementation of universal psychosocial screening, ongoing psychosocial monitoring and support from CHD diagnosis through home management, and referral to community-based services (i.e., Nurse-Family Partnership, Early Head Start) that have demonstrated positive impacts on socioeconomic status, family environment, and child developmental and behavioral outcomes.^{32–35} Although cumulative social risk was a strong predictor of outcomes, current results demonstrate that even a single social risk factor can be associated with poorer outcomes. Therefore, psychosocial screening tools, such as the Psychosocial Assessment Tool 2.0,³⁶ should be used to assess a family's individual risk factors as well as overall level of social risk.³⁷ Increased social work and psychology presence within inpatient and outpatient cardiology settings is needed to match the level of psychosocial support afforded to other critical and chronic illness populations, such as pediatric cancer.³⁸ Psychosocial standards of care based on empirical evidence are also needed to direct services and supports.

Patients who underwent cardiac surgery at Nemours but attended outpatient cardiology appointments with an outside provider due to a variety of factors (e.g., non-par insurance, far distance to hospital or regional satellite clinic) were excluded from this study. It is possible that these patients differed in important ways from those included in the study, thereby impacting generalizability of results. Included patients were followed at both the main hospital as well as regional satellite clinics, but data are not available about ease of scheduling or cardiologist availability. Schedule restrictions could result in slight delays in outpatient appointments at satellite clinics compared with the main hospital depending on the cardiologist's schedule.

Development of hospital-based services and interventions as well as referral to community-based programs that target families at high social risk may help to reduce health disparities for patients with CHD. Future research should include patients from multiple centers within diverse settings where social risk profiles or the impact of social risk may vary (e.g., region of the United States, primarily urban vs. rural setting, single payer vs. private insurance market) and should record when a particular readmission was attributable to one or more social risk factors.

Acknowledgments

Supported by the National Institute of General Medical Sciences of the National Institutes of Health (U54-GM104941 [PI: Binder-Macleod]). The study sponsor was not involved in the study design, the collection, analysis, and interpretation of data, the writing of the report, or the decision to submit the paper for publication.

Abbreviations:

CHD Congenital Heart Disease

References

1. van der Linde D, Konings EE, Slager MA, Witsenburg M, Helbing WA, Takkenberg JJ, et al. Birth prevalence of congenital heart disease worldwide: a systematic review and meta-analysis. *J Am Coll Cardiol* 2011; 58: 2241–7. [PubMed: 22078432]
2. Mahle WT, Spray TL, Wernovsky G, Gaynor JW, and Clark BJ. Survival after reconstructive surgery for hypoplastic left heart syndrome: a 15-year experience from a single institution. *Circulation* 2000; 102: III-136–III-141. [PubMed: 11082376]
3. Jacobs JP, Quintessenza JA, Burke RP, Bleiweis MS, Byrne BJ, Ceithaml EL, et al. Analysis of regional congenital cardiac surgical outcomes in Florida using the Society of Thoracic Surgeons Congenital Heart Surgery Database. *Cardiol Young* 2009; 19: 360–369 [PubMed: 19575843]
4. Hoffman JI, and Kaplan S. The incidence of congenital heart disease. *J Am Coll Cardiol* 2002; 39: 1890–1900. [PubMed: 12084585]
5. Connor JA, Kline NE, Mott S, Harris SK, and Jenkins KJ. The meaning of cost for families of children with congenital heart disease. *J Pediatr Health Care* 2010; 24: 318–325 [PubMed: 20804952]
6. Faraoni D, Nasr VG, and DiNardo JA. Overall hospital cost estimates in children with congenital heart disease: analysis of the 2012 Kid's Inpatient Database. *Pediatr Cardiol* 2016; 37: 37–43. [PubMed: 26184611]
7. Newburger JW, Sleeper LA, Bellinger DC, Goldberg CS, Tabbut S, Lu M, et al. Early developmental outcome in children with hypoplastic left heart syndrome and related anomalies: the Single Ventricle Reconstruction trial. *Circulation* 2012; 125: 2081–2091. [PubMed: 22456475]
8. Fuller S, Nord AS, Gerdes M, Wernovsky G, Jarvik GP, Bernbaum J et al. Predictors of impaired neurodevelopmental outcomes at one year of age after infant cardiac surgery. *Eur J Cardiothorac Surg* 2009; 36: 40–47. [PubMed: 19394849]
9. Boneva RS, Botto LD, Moore CA, Yang Q, Correa A, and Erickson JD. Mortality associated with congenital heart defects in the United States trends and racial disparities, 1979–1997. *Circulation* 2001; 103: 103: 2376–2381. [PubMed: 11352887]
10. Fixler DE, Pastor P, Sigman E, and Eifler CW. Ethnicity and socioeconomic status: impact on the diagnosis of congenital heart disease. *JACC* 1993; 21: 1722–1726. [PubMed: 8496543]
11. Gilboa SM, Salemi JL, Nembhard WN, Fixler DE, and Correa A Mortality resulting from congenital heart disease among children and adults in the United States, 1999 to 2006. *Circulation* 2010; 122: 2254–2263. [PubMed: 21098447]
12. Lasa JJ, Cohen MS, Wernovsky G, and Pinto NM. Is race associated with morbidity and mortality after hospital discharge among neonates undergoing heart surgery? *Pediatr Cardiol* 2013; 34: 415–423. [PubMed: 22930261]
13. Nembhard WN, Salemi JL, Ethen MK, Fixler DE, DiMaggio A, and Canfield MA. Racial/ethnic disparities in risk of early childhood mortality among children with congenital heart defects. *Pediatrics* 2001; 127: e1128–e1138.
14. Anderson BR, Fieldston ES, Newburger JW, Bacha EA, and Glied SA. Disparities in outcomes and resource use after hospitalization for cardiac surgery by neighborhood income. *Pediatrics* 2018; 141: e20172432 [PubMed: 29472494]
15. Mensah GA, Mokdad AH, Ford ES, Greenlund KJ, and Croft JB. State of disparities in cardiovascular health in the United States. *Circulation* 2005; 111: 1233–1241 [PubMed: 15769763]
16. Chang RKR, Chen AY, and Klitzner TS. Factors associated with age at operation for children with congenital heart disease. *Pediatrics* 2000; 105: 1073–1081. [PubMed: 10790465]

17. Ghanayem NS, Allen KR, Tabbutt S, Atz AM, Clabby ML, Cooper DS et al. Interstage mortality after the Norwood procedure: results of the multicenter Single Ventricle Reconstruction trial. *J Thorac Cardiovasc Surg* 2012; 144: 896–906. [PubMed: 22795436]
18. DiBardino DJ, Pasquali SK, Hirsch JC, Benjamin DK, Kleeman KC, Salazar JD et al. Effect of sex and race on outcome in patients undergoing congenital heart surgery: an analysis of the society of thoracic surgeons congenital heart surgery database. *Ann Thorac Surg* 2012; 94: 2054–2060. [PubMed: 22884593]
19. Kucik JE, Nembhard WN, Donohue P, Devine O, Wang Y, Minkovitz CS et al. Community socioeconomic disadvantage and the survival of infants with congenital heart defects. *Am J Public Health* 2014; 104: e150–e157. [PubMed: 25211743]
20. Peiris V, Singh TP, Tworetzky W, Chong EC, Gauvreau K, and Brown DW. Association of socioeconomic position and medical insurance with fetal diagnosis of critical congenital heart disease. *Circ Cardiovasc Qual Outcomes* 2009; 2: 354–360. [PubMed: 20031861]
21. Kucik JE, Cassel CH, Alverson CJ, Donohue P, Tanner JP, Minkovitz CS et al. Role of health insurance on the survival of infants with congenital heart defects. *Am J Public Health* 2014; 104: e62–e70.
22. DeMone JA, Gonzalez PC, Guavreau K, Piercey GE, and Jenkins KJ. Risk of death for Medicaid recipients undergoing congenital heart surgery. *Pediatr Cardiol* 2003; 24: 97–102. [PubMed: 12360394]
23. Mussatto KA, Hollenbeck-Pringle D, Trachtenberg F, Sood E, Sananes R, Pike NA, et al. Utilisation of early intervention services in young children with hypoplastic left heart syndrome. *Cardiol Young* (2017); 28: 1–8. [PubMed: 28803588]
24. Mayer ML, Skinner AC, and Slifkin RT. Unmet need for routine and specialty care: data from the National Survey of Children with Special Health Care Needs. *Pediatrics* 2004; 113: e109–e115. [PubMed: 14754979]
25. Chen E, Matthews KA, and Boyce WT Socioeconomic differences in children's health: how and why do these relationships change with age? *Psychol Bull* 2002; 128: 295–329. [PubMed: 11931521]
26. Flores G, Olson L, and Tomany-Korman SC Racial and ethnic disparities in early childhood health and health care. *Pediatrics* 2005; 115: e183–e193. [PubMed: 15687426]
27. Jacobs ML, O'Brien SM, Jacobs JP, et al. An empirically based tool for analyzing morbidity associated with operations for congenital heart disease. *J Thorac Cardiovasc Surg* 2013; 145: 1046–1057.e1. [PubMed: 22835225]
28. Loccoh EC, Yu S, Donohue J, Lowery R, Butcher J, Pasquali SK et al. Prevalence and risk factors associated with non-attendance in neurodevelopmental follow-up clinic among infants with CHD. *Cardiol Young* 2018; 28: 554–560. [PubMed: 29357956]
29. Samuels RC, Ward VL, Melvin P, Macht-Greenberg M, Wenren LM, Yi J, et al. Missed appointments: factors contributing to high no-show rates in urban pediatrics primary care. *Clin Pediatr* 2015; 54: 976–982.
30. Peres H, Glazer Y, Landau D, Marks K, Abokaf H, Belmaker I, et al. Understanding utilization of outpatient clinics for children with special health care needs in southern Israel. *Matern Child Health J* 2014; 18: 1831–1845. [PubMed: 24414986]
31. Mackie AS, Rempel GR, Rankin KN, Nicholas D, Magill-Evans J. Risk factors for loss to follow-up among children and young adults with congenital heart disease. *Cardiol Young* 2012; 22: 307–315. [PubMed: 22013913]
32. Olds DL, Kitzman HJ, Cole RE, Hanks CA, Arcoleo KJ, Anson EA, et al. Enduring effects of prenatal and infancy home visiting by nurses on maternal life course and government spending: follow-up of a randomized trial among children at age 12 years. *Arch Pediatr Adolesc Med* 2010; 164: 419–424. [PubMed: 20439792]
33. Olds DL, Robinson J, Pettitt L, Luckey DW, Homberg J, Ng RK, et al. Effects of home visits by paraprofessionals and by nurses: age 4 follow-up results of a randomized trial. *Pediatrics* 2004; 114: 1560–1568. [PubMed: 15574615]

34. Olds DL, Kitzman H, Knudtson MD, Anson E, Smith JA, and Cole R. Effect of home visiting by nurses on maternal and child mortality: results of a 2-decade follow-up of a randomized clinical trial. *JAMA Pediatr* 2014; 168: 800–806. [PubMed: 25003802]
35. Goyal NK, Teeters A, and Ammerman RT. Home visiting and outcomes of preterm infants: a systematic review. *Pediatrics* 2013; 132: 502–516. [PubMed: 23940238]
36. Pai ALH, Patino-Fernandez AM, McSherry M, Beele D, Alderfer MA, Reilly AT, et al. The Psychosocial Assessment Tool (PAT2.0): psychometric properties of a screener for psychosocial distress in families of children newly diagnosed with cancer. *J Pediatr Psychol* 2008; 33(1): 50–62. [PubMed: 17609228]
37. Hearps SJ, McCarthy MC, Muscara F, Hearps SJ, Burke K, Jones B, et al. Psychosocial risk in families of infants undergoing surgery for a serious congenital heart disease. *Cardiol Young* 2014; 24(4): 632–639. [PubMed: 23803336]
38. Wiener L, Kazak AE, Noll RB, Patenaude AF, and Kupst MJ. Standards for the psychosocial care of children with cancer and their families: An introduction to the special issue. *Pediatr Blood Cancer* 2015; 62: S419–S424. [PubMed: 26397836]

Figure.
Significant Interactions between Social Risk and Surgical Morbidity Score for Readmission Days

Table 1.

Patient Characteristics

	Low Social Risk N (%)	High Social Risk N (%)	Total Sample N (%)
Clinical Characteristic			
Premature Birth	15 (12.9%)	12 (11.7%)	27 (12.3%)
Confirmed/Suspected Genetic Condition	37 (31.9%)	32 (31.1%)	69 (31.5%)
Single Ventricle Anatomy	19 (16.4%)	13 (12.6%)	32 (14.6%)
Neonatal Surgery	54 (46.6%)	40 (38.8%)	94 (42.9%)
Admissions and Interventions			
1 Unplanned Readmissions	43 (37.1%)	54 (52.4%)	97 (44.3%)
1 Total Readmissions	61 (52.6%)	66 (64.1%)	127 (58.0%)
>1 Cardiac Surgical Interventions	38 (32.8%)	41 (39.8%)	79 (36.1%)

Table 2.

Outcomes for Individual Social Risk Factors (Online Only)

	Racial/Ethnic Minority		Medicaid Insurance		High Poverty Area		Maternal Age <21		Non-English Speaking		>25 Miles from Outpatient	
	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No
	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)
	Statistics	Statistics	Statistics	Statistics	Statistics	Statistics	Statistics	Statistics	Statistics	Statistics	Statistics	Statistics
Surgical Length of Stay	30.34	27.63	30.81	26.11	22.27	29.60	16.67	29.62	14.73	29.47	31.95	27.60
	(45.20)	(35.83)	(41.35)	(37.88)	(21.50)	(41.65)	(13.72)	(40.98)	(12.24)	(40.65)	(37.49)	(40.67)
	$t=-0.49, p=0.62$		$t=-0.87, p=0.39$		$t=0.88, p=0.38$		$t=1.22, p=0.23$		$t=1.20, p=0.23$		$t=-0.71, p=0.48$	
Unplanned Readmissions	1.58	1.04	1.49	0.97	0.88	1.31	1.50	1.24	2.00	1.22	1.16	1.29
	(2.12)	(1.89)	(2.09)	(1.85)	(1.53)	(2.05)	(1.83)	(2.01)	(2.31)	(1.98)	(2.04)	(1.98)
	$t=-1.98, p=0.05$		$t=-1.92, p=0.06$		$t=1.01, p=0.31$		$t=-0.47, p=0.64$		$t=-1.21, p=0.23$		$t=0.42, p=0.68$	
Total Readmissions	2.17	1.65	2.14	1.51	1.62	1.89	2.00	1.85	2.09	1.85	1.75	1.90
	(2.53)	(2.23)	(2.49)	(2.15)	(2.47)	(2.35)	(2.30)	(2.37)	(2.21)	(2.38)	(2.16)	(2.44)
	$t=-1.61, p=0.11$		$t=-1.99, p=0.05$		$t=0.56, p=0.58$		$t=-0.24, p=0.81$		$t=-0.33, p=0.74$		$t=0.39, p=0.70$	
Readmission Days	22.46	15.61	22.91	12.71	15.42	18.79	11.33	18.91	19.64	18.33	21.82	17.19
	(36.99)	(30.52)	(39.36)	(22.78)	(33.66)	(33.42)	(18.80)	(34.18)	(31.16)	(33.57)	(40.83)	(30.39)
	$t=-1.50, p=0.14$		$t=-2.27, p=0.02$		$t=0.48, p=0.63$		$t=0.85, p=0.40$		$t=-0.13, p=0.90$		$t=-0.90, p=0.37$	
Subsequent surgical interventions	0.56	0.56	0.65	0.45	0.54	0.56	0.73	0.55	0.27	0.58	0.60	0.55
	(0.87)	(0.89)	(0.98)	(0.74)	(0.76)	(0.90)	(1.16)	(0.86)	(0.47)	(0.90)	(0.75)	(0.93)
	$t=-0.02, p=0.99$		$t=-1.62, p=0.11$		$t=0.14, p=0.89$		$t=-0.78, p=0.44$		$t=1.11, p=0.27$		$t=-0.35, p=0.73$	
Adherence 0-12m	12.87	6.35	11.04	6.41	11.05	8.70	10.79	8.85	3.76	9.26	11.36	8.14
	(26.14)	(14.43)	(23.44)	(14.94)	(25.68)	(19.40)	(19.25)	(20.30)	(3.87)	(20.68)	(22.82)	(19.19)
	$t=-2.36, p=0.02$		$t=-1.69, p=0.09$		$t=-0.56, p=0.58$		$t=-0.36, p=0.72$		$t=0.88, p=0.38$		$t=-1.03, p=0.30$	
Adherence 12-24m	40.08	20.52	40.09	13.41	57.85	24.94	42.32	27.49	32.23	28.21	30.98	27.48
	(88.45)	(58.80)	(85.81)	(47.51)	(115.31)	(65.48)	(102.28)	(70.55)	(49.79)	(73.85)	(77.71)	(71.00)
	$t=-1.92, p=0.06$		$t=-2.67, p=0.01$		$t=-2.02, p=0.04$		$t=-0.71, p=0.48$		$t=-0.18, p=0.86$		$t=-0.31, p=0.76$	

Note: M = mean; SD = standard deviation; m = months

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3.

Estimated Marginal Means for Social Risk by Surgical Morbidity Score Interactions

	Statistics	Social Risk Level <i>F, p, η^2</i>	Morbidity One <i>M (SE)</i>	Morbidity Two <i>M (SE)</i>	Morbidity Three <i>M (SE)</i>	Morbidity Four <i>M (SE)</i>	Morbidity Five <i>M (SE)</i>
Surgical Length of Stay	$F=1.23, p=0.34$	Low	15.76 (6.03)	18.33 (9.98)	22.83 (6.94)	44.83 (8.06)	60.17 (15.77)
	$\eta^2=0.02$	High	20.40 (6.53)	30.71 (7.30)	44.93 (10.33)	32.57 (10.33)	51.00 (11.15)
Unplanned Readmissions	$F=1.56, p=0.19$	Low	0.54 (0.30)	0.53 (0.50)	1.00 (0.35)	1.52 (0.40)	2.83 (0.79)
	$\eta^2=0.03$	High	1.44 (0.33)	1.86 (0.37)	1.14 (0.52)	0.93 (0.52)	2.58 (0.56)
Total Readmissions	$F=1.64, p=0.16$	Low	0.76 (0.34)	0.80 (0.56)	1.42 (0.39)	2.91 (0.46)	4.00 (0.89)
	$\eta^2=0.03$	High	1.83 (0.39)	2.07 (0.41)	1.71 (0.58)	1.93 (0.58)	4.67 (0.63)
Readmission Days	$F=3.07, p=0.02$	Low	3.02 (4.65)	2.20 (7.68)	10.00 (5.34)	33.26 (6.20)	48.67 (12.14)
	$\eta^2=0.06$	High	16.63 (5.03)	27.68 (5.62)	10.43 (7.95)	14.29 (7.95)	66.75 (8.59)
Subsequent surgical interventions	$F=2.20, p=0.07$	Low	0.15 (0.11)	0.07 (0.18)	0.39 (0.13)	1.39 (0.14)	1.67 (0.28)
	$\eta^2=0.04$	High	0.11 (0.12)	0.39 (0.13)	0.64 (0.18)	0.93 (0.18)	2.08 (0.20)
Adherence 0–12m	$F=0.90, p=0.47$	Low	8.26 (3.12)	1.39 (5.15)	5.11 (3.58)	5.72 (4.16)	0.60 (8.15)
	$\eta^2=0.02$	High	16.11 (3.37)	16.57 (3.77)	7.76 (5.33)	4.65 (5.33)	9.38 (6.02)
Adherence 12–24m	$F=0.41, p=0.80$	Low	12.14 (11.67)	40.21 (18.33)	4.58 (12.96)	20.01 (14.80)	4.43 (28.98)
	$\eta^2=0.01$	High	50.05 (12.36)	62.56 (14.20)	12.28 (18.97)	55.70 (19.69)	10.27 (20.49)

Notes: M = mean; SE = standard error; m = months

Table 4.

Simple Main Effects of Surgical Morbidity and Social Risk with Post-Hoc Probing

	Surgical Morbidity					Social Risk (SR)		
	Stats <i>F</i> , <i>p</i> , η^2	M1 M (SE)	M2 M (SE)	M3 M (SE)	M4 M (SE)	M5 M (SE)	Sig. Diff.	Stats <i>F</i> , <i>p</i> , η^2
Surgical Length of Stay	<i>F</i> =4.23, <i>p</i> =0.03, $\eta^2=0.08$	18.08 (4.45)	24.52 (6.18)	33.88 (6.22)	38.70 (6.22)	55.58 (9.66)	5 vs. 1	<i>F</i> =0.34, <i>p</i> =0.56, $\eta^2<0.01$
Unplanned Readmissions	<i>F</i> =2.68, <i>p</i> =0.03, $\eta^2=0.05$	0.99 (0.23)	1.20 (0.31)	1.07 (0.31)	1.23 (0.33)	2.71 (0.49)	5 vs. 1, 3	<i>F</i> =0.99, <i>p</i> =0.32, $\eta^2=0.01$
Total Readmissions	<i>F</i> =7.49, <i>p</i> <0.01, $\eta^2=0.13$	1.29 (0.25)	1.44 (0.35)	1.57 (0.35)	2.42 (0.37)	4.33 (0.55)	5 vs. 1-4	<i>F</i> =1.81, <i>p</i> =0.18, $\eta^2=0.01$
Readmission Days	<i>F</i> =9.59, <i>p</i> <0.01, $\eta^2=0.16$	9.83 (3.42)	14.94 (4.76)	10.21 (4.79)	23.77 (5.04)	57.71 (7.46)	5 vs. 1-4	<i>F</i> =2.70, <i>p</i> =0.10, $\eta^2=0.01$
Subsequent surgical interventions	<i>F</i> =31.27, <i>p</i> <0.01, $\eta^2=0.37$	0.13 (0.8)	0.23 (0.11)	0.52 (0.11)	1.16 (0.12)	1.88 (0.17)	5 vs. 1-4, 4 vs. 1-3	<i>F</i> =0.87, <i>p</i> =0.35, $\eta^2<0.01$
Adherence 0-12m	<i>F</i> =1.11, <i>p</i> =0.36, $\eta^2=0.02$	12.18 (2.29)	8.98 (3.19)	6.44 (3.21)	5.18 (3.38)	4.99 (5.06)	— —	<i>F</i> =4.43, <i>p</i> =0.04, $\eta^2=0.02$
Adherence 12-24m	<i>F</i> =2.25, <i>p</i> =0.07, $\eta^2=0.04$	31.10 (8.50)	51.39 (11.59)	8.43 (11.49)	37.86 (12.32)	7.35 (17.75)	— —	<i>F</i> =3.72, <i>p</i> =0.06, $\eta^2=0.02$

Notes: M1-M5 = Morbidity Scores 1-5; SR = Social Risk; M = mean; SE = standard error; m = months; Sig. Diff. = Significant Group Differences