Published in final edited form as:

Musculoskelet Sci Pract. 2019 June ; 41: 36–42. doi:10.1016/j.msksp.2019.03.004.

Treatment monitoring as a component of psychologically informed physical therapy: a case series of patients at high risk for persistent low back pain related disability

Jason M. Beneciuk, DPT, PhD, MPH^{a,b}, Lindsay A. Ballengee, PT, DPT^c, and Steven Z. George, PT, PhD, FAPTA^d

^aUniversity of Florida, College of Public Health & Health Professions, Department of Physical Therapy, Box 100154, UFHSC, Gainesville, FL 32610-0154.

^bBrooks Rehabilitation Clinical Research Center, 3901 University Blvd. South, Suite 103, Jacksonville, FL 32216.

^cDepartment of Orthopaedic Surgery, Duke University, 2200 W. Main Street, Suite B-230, Durham, NC 27705.

^dDuke Clinical Research Institute and Department of Orthopaedic Surgery, Duke University, 2400 Pratt Street, Durham NC 27705.

Abstract

Background: Psychologically Informed Physical Therapy (PIPT) aims to identify individuals at high risk for transitioning to chronicity and merge impairment-focused physical therapy with cognitive behavioral therapy principles. Treatment monitoring is an important part of PIPT and involves identifying changes in clinical measures to inform clinical decision making.

Objectives: The purpose of this case series is to describe treatment monitoring using psychological and physical impairment measures for patients identified as ‘high-risk’ for persistent low back pain (LBP) related disability.

Design: Secondary analysis of patients (n = 23) identified as ‘high-risk’ using the STarT Back Tool and enrolled in two-phased, sequential study that evaluated feasibility and generated preliminary PIPT treatment effects for 4-week clinical outcomes.

Method: Physical therapists (n = 5) used psychological [Fear-Avoidance Beliefs Questionnaire (FABQPA, FABQ-W), Tampa Scale for Kinesiophobia (TSK-11), Pain Catastrophizing Scale (PCS) and Fear of Daily Activities Questionnaire (FDAQ)] and the Physical Impairment Index (PII) measures for PIPT treatment monitoring. Clinical outcome measures [Numerical Pain Rating

Corresponding author: Jason Beneciuk at: beneciuk@phhp.ufl.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of Interest: The authors declare no conflicts of interest.

Ethical Approval: All enrolled patients provided informed consent in compliance with the University of Florida Institutional Review Board.

Scale (NPRS), Oswestry Disability Index (ODI)] were administered at intake and 4-weeks later. Linear regression models evaluated independent contribution of intake and 4-week changes in psychological measures and PII scores as predictors of 4-week NPRS and ODI scores.

Results: FABQ-PA and PCS changes provided largest contributions to prediction of 4-week ODI scores. Treatment monitoring measures did not explain additional variability in 4-week NPRS scores after baseline scores were considered.

Conclusions: For patients at high risk for persistent LBP psychological measures consistently performed better as treatment monitoring variables compared to physical impairment measures. Treatment monitoring for PIPT with psychological measures provides opportunities to refine prediction of disability outcomes. Findings from this exploratory case series should be interpreted with caution based on its small sample size and lack of statistical power which prohibits definitive conclusions to be made on any of the treatment monitoring measures.

Keywords

Psychologically Informed Physical Therapy; treatment monitoring; low back pain

Low back pain (LBP) is a major public health concern because it is highly prevalent and the leading cause of disability worldwide.¹ Global prevalence of LBP has increased by 17.3% from 2005 to 2015 and continues to be a leading cause of global years lived with disability since 1990.^{1, 2} Furthermore, approximately 25% of people consulting primary care providers for an episode of acute LBP will develop chronic pain.³ Psychologically informed physical therapy (PIPT) is a secondary prevention approach for LBP that first aims to identify individuals at high risk for transitioning to chronicity and then provides tailored treatment by merging impairment-focused physical therapy with cognitive behavioral therapy principles as needed to reduce that risk.^{4, 5} The primary goal of PIPT is to prevent LBP associated disability by emphasizing: (1) identification of individuals who are at high risk for developing chronic LBP based on the presence of psychological distress and (2) targeted treatment aimed at psychological factors in conjunction with traditional, impairment-based physical therapy.⁴

The Subgroups for Targeted Treatment (STarT) Back Approach involves patient stratification into low, medium or high-risk groups and providing matched treatment based on their prognostic risk profile.⁶ Implementing the STarT Back Approach has resulted in increased generic health benefit and cost savings in primary care settings⁷ and improved clinical outcomes in outpatient physical therapy settings.⁸ Risk group stratification is primarily determined by responses to STarT Back Tool (SBT) psychological items with PIPT treatment approaches indicated for patients identified as 'high-risk'.^{9, 10}

Treatment monitoring is an important component of PIPT that considers the intermediary process between treatment received and clinical outcome.^{5, 11} For a high-risk patient, treatment monitoring should consist of longitudinal tracking to identify changes in risk status or dynamic aspects of psychological distress or impairment based measures (e.g., range of motion, spinal tenderness) to predict poor outcomes for pain and functional status.^{12, 13} Favorable change may indicate which treatment components should continue to be

emphasized; while lack of favorable change may suggest modification in specific treatment components are warranted. Treatment monitoring also has potential for improving clinical outcomes through providing opportunities for enhanced communication with patients.^{14, 15}

Throughout this manuscript, the term ‘*treatment monitoring*’ describes the process of identifying changes in specific psychological or physical impairment measures that occur in conjunction with PIPT treatment provided. Treatment monitoring can be considered a pragmatic, clinically feasible method to inform clinical decision making by identifying potentially relevant treatment mediators (i.e., variables that change as a result of treatment and related to change in outcome of interest) during patient encounters.^{11, 16, 17} An example of treatment monitoring for a high-risk patient would be measurement of reduction of psychological distress during graded exposure¹¹ or increased lumbar flexion range of motion and decreased spinal tenderness during manual therapy and exercise.

Treatment monitoring that is feasible to implement during routine clinical practice is an important element to support clinical decision making since previous studies have identified preliminary key mediators.^{18–20} For example, Smeets et al. reported catastrophizing and perceived control over pain were mediators for outcomes in chronic low back pain.¹⁸ Recent systematic review findings indicate self-efficacy, psychological distress, and pain-related fear to be mediators of pain or disability outcomes for spine related conditions, although it has been acknowledged that these findings are not conclusive and further research is needed in this area.²¹ Recently, Mansell, et al. found changes in pain-related distress and pain intensity to have mediating effects on the relationship between treatment group allocation (i.e., PIPT or current best care) and change in disability outcomes for patients identified as SBT high-risk for persistent LBP related disability.²⁰

The purpose of this exploratory case series is to describe a treatment monitoring process during a 4-week episode of PIPT. This case series included a previously validated physical impairment index and selected psychological measures as treatment monitoring variables. This case series intentionally included patients identified as SBT ‘high-risk’ to distinguish from previous studies in this area that did not account for initial risk status, therefore our limited sample size was not appropriate for formal mediation analyses. The overall goal of this exploratory case series was to provide initial data to inform clinical decision making and provide information for powering future larger studies that would focus on patients identified as SBT high risk.

METHODS

Design Overview

Patients included in this case series were part of a two-phased, sequential study that evaluated feasibility and generated preliminary PIPT treatment effects for 4-week clinical outcomes.⁸ The second phase of that study is most relevant as it consisted of patient outcome collection from participating clinicians. All clinicians and patients were selected from seven outpatient physical therapy clinics.

Risk Stratified Care Training

Clinicians randomized to apply risk stratified care attended three training sessions over the course of 4 weeks. Each session was provided by the study authors (XXX, XXX) and lasted between 2 and 4 hours. Education content focused on PIPT practice principles⁴ and was developed to reflect clinical protocols that have been used in previous studies.^{10, 22, 23} The overall objective of this multifactorial education approach was for clinicians to embrace the biopsychosocial model of pain and disability.²⁴ As a component of stratified care education, physical therapists were trained to implement the STarT Back approach for LBP management.^{6, 7} The STarT Back approach provides an example of stratification based on prognostic risk for persistent LBP related disability that consists of two corresponding components. First, patients are categorized into one of three subgroups (low, medium, or high risk) for persistent LBP related disability using the 9-item STarT Back Screening Tool (SBT).⁹ Second, targeted treatment pathways are matched to each SBT subgroup such that patients categorized as low risk receive minimal care primarily consisting of reassurance and education while patients categorized as medium risk also received traditional physical therapy approaches that restoring function by targeting physical symptoms.²⁵ For patients categorized as high risk, physical therapy is focused on restoring function using PIPT which consisted of patient-centered communication, patient education (e.g., pain neuroscience, activation philosophy), activity based intervention (graded exercise or graded exposure), physical impairment based intervention (guided by recent clinical practice guidelines).^{8, 25} Those interested in further description of the intervention approach used for high risk patients in this study can refer to the primary study methods.⁸ Specific to this case series, physical therapists providing stratified care for high risk patients were also instructed in treatment monitoring using unidimensional psychological measures.⁸

Patient Participants

Consecutive patients referred by a physician to physical therapy for LBP were evaluated by participating physical therapists at outpatient clinic locations. Physical therapists recruited and screened all patients with LBP for study eligibility prior to enrollment. Study inclusion criteria consisted of: 1) adults between the ages of 18 and 65 years seeking physical therapy for LBP (defined as having symptoms at T12 or lower, including radiating pain into the buttocks and lower extremity) and 2) the ability to read and speak the English language. Study exclusion criteria consisted of: 1) the presence of systemic involvement related to metastatic or visceral disease; 2) recent spinal fracture; 3) osteoporosis; or 4) pregnancy. All enrolled patients provided informed consent in compliance with the XXX Institutional Review Board.

Demographic and Historical Information

Study participants completed a standardized questionnaire consisting of items related to age, sex, race, and employment status. Information involving LBP clinical characteristics (i.e., prior surgery, symptom duration, symptom onset, symptom location, work-related LBP) was also obtained.

Determining Risk Status

STarT Back Screening Tool (SBT).—The SBT is a 9-item screening measure used to identify subgroups of patients with LBP based on the presence of modifiable prognostic factors which may be useful in matching patients with targeted interventions.⁹ SBT overall scores (ranging from 0 to 9) and SBT psychosocial subscale scores (ranging from 0 to 5) are determined by summing all positive responses. The SBT then categorizes patients as ‘high-risk’ (psychosocial subscale scores ≥4) in which high levels of psychosocial prognostic factors are present with or without physical factors present, ‘medium-risk’ (overall score >3; psychosocial subscale score <4) in which physical and psychosocial factors are present, but not a high level of psychosocial factors, or ‘low-risk’ (overall score 0–3) in which few prognostic factors are present.⁶ This case series reports on only those patients that met these eligibility criteria, provided informed consent, and were categorized as SBT ‘high-risk’.

Treatment Monitoring Measures

Treatment monitoring measures were collected at baseline and 4 weeks later. These measures are described in more detail below:

Fear-Avoidance Beliefs Questionnaire (FABQ-PA, FABQ-W).—Fear-avoidance beliefs specific to LBP were assessed with the FABQ.²⁶ The FABQ consists of a 4-item FABQ physical activity scale (FABQ-PA, potentially ranging from 0 to 24) and a 7-item FABQ work scale (FABQ-W, potentially ranging from 0 to 42), with higher scores indicating higher levels of fear-avoidance beliefs for both FABQ scales. Both FABQ scales have been found to have acceptable reliability^{26–29} and demonstrated internal consistency.^{26, 29–32} The FABQ-W has demonstrated predictive validity for disability and work loss in patients with LBP.^{29, 33–35}

Pain Catastrophizing Scale (PCS).—The PCS was used to assess the degree of catastrophic cognitions due to LBP.³⁶ Pain catastrophizing has been broadly defined as an exaggerated negative orientation towards actual or anticipated pain experiences.³⁶ The PCS is a 13-item questionnaire with a potential range of 0 to 52, with higher scores indicating higher levels of pain catastrophizing. The PCS has been found to have good test-retest reliability³⁶ and internal consistency.^{37–39} The PCS has also been found to demonstrate several different types of validity.^{36–39}

Tampa Scale of Kinesiophobia (TSK-11).—The TSK-11 was used to assess the degree of fear of movement and injury or re-injury in individuals with LBP.⁴⁰ The TSK-11 is an 11-item questionnaire with a potential range of 11 to 44, with higher scores indicating greater fear of movement and injury or re-injury due to pain. The TSK-11 has been found to have good test-retest reliability and internal consistency.⁴⁰ Predictive and concurrent validity have also been reported for the TSK-11.⁴⁰

Fear of Daily Activities Questionnaire (FDAQ).—The FDAQ was used to quantify fear of ‘specific’ activities commonly reported by patients experiencing LBP.⁴¹ The FDAQ is a 10-item scale with a total range of 0 to 100, with higher scores indicating higher fear of

specific activities. Inclusion of the FDAQ has been shown to improve subgrouping methods for patients with LBP.⁴²

Physical Impairment Index (PII).—The PII was administered by a physical therapist to quantify physical impairment due to LBP.⁴³ The PII consists of 7 different physical examination procedures: 1) lumbar flexion range of motion (ROM); 2) lumbar extension ROM; 3) lumbar lateral flexion ROM; 4) passive straight leg raise ROM; 5) bilateral active straight leg raise; 6) active sit-up; and 7) assessment of spinal tenderness with each procedure scored as a negative (0) or positive (1) for presence of impairment. The PII has a total range of 0 to 7, with higher scores indicating higher levels of physical impairment due to LBP. Good or excellent reliability has been reported for individual PII items and convergent validity has been supported for LBP disability.^{43, 44} Overall PII scores have been reported to be more responsive to change compared to individual test items, with a 1-point change representing minimal detectable change over four-weeks of physical therapy.⁴⁴

Clinical Outcome Measures

Clinical outcome measures were collected at intake and 4-weeks later and are described in more detail below.

Numerical Pain Rating Scale (NPRS)—Pain intensity was rated using a NPRS, ranging from “0” (no pain) to “10” (worst pain imaginable). Participants were asked to rate their current pain intensity, as well as their best and worst level of pain intensity over the past 24 hours. These three pain ratings were averaged and used as the NPRS variable in this study.⁴⁵ The NPRS has been found to have sound psychometric properties^{45–47} with a minimal clinically important difference reported to be 2 points.⁴⁸

Modified Oswestry Disability Questionnaire (ODI)—LBP-related disability was assessed with the ODI, which has 10 items that assesses how LBP affects common daily activities.^{49, 50} The ODI has a range of 0% “no disability due to LBP” to 100% “completely disabled due to LBP”, with higher scores indicating higher disability from LBP. The ODI has been found to have sound psychometric properties^{49, 51, 52} with a minimal clinically important difference reported to be 10 percentage points.⁵³

Data Analysis

All statistical analyses were performed using SPSS version 24.0 (SPSS Inc., Chicago, Illinois). Descriptive statistics were used to provide a summary of data with means and standard deviations calculated for all baseline continuous variables and frequency counts with percentages for categorical variables.

Univariate Associations between Treatment Monitoring and Clinical Outcomes Measures—Bivariate associations using Pearson r correlation coefficients were calculated for 4 week changes in treatment monitoring (FABQ-PA, FABQ-W, PCS, TSK-11, FDAQ, and PII) and clinical outcome (NPRS and ODI) measure scores to assess which observed changes were associated with clinical outcomes. Magnitude of Pearson r correlation

coefficients were interpreted as “little to no relationship” (0.00 to .25), “fair” (.25 to .50), “moderate to good” (.50 to .75), and “good to excellent” (above .75).⁵⁴

Multivariate Treatment Monitoring Associations with Four Week Clinical Outcomes

Separate simple linear regression models were used to evaluate independent associations for intake and 4-week changes in treatment monitoring (FABQ-PA, FABQ-W, PCS, TSK-11, FDAQ, and PII) measure scores with NPRS and ODI scores at 4-weeks. Two separate models were created for each treatment monitoring measure. Model 1 accounted for intake dependent variable scores (i.e., NPRS or ODI depending upon the outcome of interest) and intake psychological measure or PII score (depending upon treatment monitoring variable of interest). Model 2 added 4-week psychological measure or PII change scores to Model 1.

RESULTS

Twenty three participants out of the original 70 (32.9%) that received PIPT were categorized as ‘high risk’ and included in this case series (Table 1). Duration of symptoms for this sample was categorized into three groups (14 days or less [n = 1, 4.3%]; 15 to 90 days [n = 10, 43.5%]; and 91 days or longer [n = 12, 52.2%]). Symptom duration was not associated with NPRS or ODI outcome scores at 4 weeks ($P > 0.05$) and therefore was not included as a covariate in regression models.

Univariate Correlates of Treatment Monitoring and Clinical Outcome Measures

Pearson correlation coefficients are reported in Table 2. Moderate to good associations ($r = .574$ to $.768$, $P < 0.05$) were observed between psychological measure 4 week change scores. Little associations were observed between psychological measures and PII 4 week change scores ($r = .028$ to $.185$; $P > 0.05$). Moderate to good associations ($r = .455$ to $.697$, $P < 0.05$) were observed between 4 week changes in ODI scores and changes in psychological measures while little associations ($r = .010$ to $.260$; $P > 0.05$) between 4 week changes in NPRS scores and changes in treatment monitoring measures were observed.

Treatment Monitoring Multivariate Association with 4 Week Clinical Outcomes

Pain Intensity—Intake NPRS and treatment monitoring measures explained a large amount of variability in 4-week NPRS scores (range: 45.6% to 47.7%, $P < 0.05$) (**Table 3 – Model 1**). The addition of 4-week change in treatment monitoring measures did not explain additional variability in 4-week NPRS scores (**Table 3 – Change from Model 1 to Model 2**).

Disability—Intake ODI and treatment monitoring measures explained a large amount of variability in 4-week ODI scores (range: 30.6% to 38.0%) (**Table 4 – Model 1**). However, the 4-week change in treatment monitoring added to these regression models. Specifically, the addition of 4-week FABQ-PA (37.0%), FABQ-W (20.0%), PCS (27.1%), TSK-11 (16.3%), and FDAQ (15.2%) change scores explained an additional amount of variability ($P < 0.05$) in 4-week ODI scores (**Table 4 – Change from Model 1 to Model 2**).

DISCUSSION

This exploratory case series used psychological and physical impairment measures as treatment monitoring variables for patients with LBP that were categorized as SBT high risk. Our primary finding was that for LBP related disability, psychological measures were consistently better for treatment monitoring in comparison to the PII. Another important finding was that none of the treatment monitoring measures improved prediction of 4-week pain intensity outcomes better than baseline pain intensity scores. Collectively, these results indicate that for patients with LBP categorized as SBT high-risk, treatment monitoring (i.e. 4-week change from baseline) with psychological measures can be used to refine prediction of disability outcomes, but not for pain intensity. These results also suggest that there may be a limited treatment monitoring role for physical impairment measures in this patient population.

The finding that in this small sample of patients psychological measures were better at treatment monitoring in comparison to a PII is not surprising. We intentionally used a previously validated physical impairment index that incorporated different individual measures and has been correlated with recovery of function in patients with LBP.^{43, 44, 49} However, this composite measure did not contribute to regression models in this small cohort. Previous studies have indicated psychological factors to be stronger predictors for LBP outcomes when compared to physical impairment measures.^{55, 56} This exploratory case series adds to the existing literature by considering the PII for treatment monitoring in high risk patients when in previous studies only baseline values were used without consideration of risk status. However, this does not necessarily exclude measuring physical impairment as part of treatment monitoring as this sample was too small to make definitive conclusions.

Furthermore, this case series suggests that general psychological measures may be appropriate for treatment monitoring for high risk patients when compared to measures that elicit more specific beliefs. In this case series, the FDAQ was included as a patient specific measure and it contributed the smallest amount of variance to disability outcomes. Therefore, there may be limited value in capturing and monitoring fear of patient-specific activities over existing general psychological measures. These results align with previous research that has examined specific fear-avoidance activity monitoring to predict disability outcomes.⁵⁷ For example, the Photograph Series of Daily Activities (PHODA) has been found to reliably detect changes over time in pain-related fear of patients with chronic LBP, however not able to identify patients who are likely to improve according to other pain-related fear measures such as the FABQ and TSK.⁵⁸

A primary limitation of this case series was the small sample size of 23 high risk patients with LBP. When focusing on SBT high risk patients, a smaller sample is inevitable as three separate outpatient physical therapy cohorts in the United States have indicated the prevalence of high risk patients ranged from 20–30% of LBP episodes.^{8, 59, 60} There are several limitations related to the sample size including prohibiting use of more sophisticated statistical models for analysis (e.g., latent growth modelling) and the statistical estimates we provided lack precision. Interestingly, there does not appear to be a major issue with statistical power due to sample size. The regression models were generally able to detect

changes of 10% or more, which suggests the treatment monitoring variables identified in this case series may potentially have clinical relevance. However, there are inherent limits with samples this small as it limits the type of statistical models that can be used and estimates will be imprecise. Despite this limitation these data can be used to adequately power larger studies focused on SBT high risk patients.

Additional limitations include not administering psychological or physical impairment measures to low or medium risk patients with LBP. Having these additional measures would have allowed comparison of treatment monitoring potential in different SBT risk categories. Finally, we may have obtained more valuable information if treatment monitoring measures were administered during additional time points during the episode of care. If treatment-monitoring measures were administered weekly, for example, we may have been able to obtain higher granularity data or been able to detect whether linear changes in pain associated psychological distress had occurred.

This case series provides several directions for future research. First, the results suggest there is a role for exploring different psychological measures such as self-efficacy for improved treatment monitoring in predicting pain intensity outcomes. Other studies have found promising results by capturing self-efficacy as a significant predictor of health related quality of life among patients with chronic LBP.⁶¹ Additionally, higher self-efficacy has been associated with a protective effect for severity of pain symptoms in patients with chronic pain.⁶² Therefore, self-efficacy may be a useful treatment-monitoring variable for future pain intensity and disability outcomes, but to the best of our knowledge, it has not been investigated as such in high-risk patients.

Another direction for future research would be to compare whether use of composite psychological measures (e.g., STarT Back Tool, Örebro Musculoskeletal Pain Screening Questionnaire, and OSPRO Yellow Flag Assessment Tool)^{9, 63, 64} can be used to enhance treatment monitoring as previous studies have found changes in these measures to improve outcome prediction for disability outcomes in cohort studies.^{12, 13, 60} Future research could compare these composite measures with unidimensional measures to determine which are most appropriate for treatment monitoring in high-risk patients.

Finally, future research should consider how other non-psychological measures such as socioeconomic status⁶⁵ and race⁶⁶ impact treatment monitoring to provide further insight to the unexplained variance in pain and disability outcomes. In this case series, we included an index consisting of 8 different components including lumbar flexion, extension, pelvic flexion, lateral flexion, straight leg raising, spinal tenderness, bilateral active straight leg raise, and sit-up. However, additional composite impairment or performance based measures such as postural control or the ability to perform a functional task may provide more useful information that is also feasible to assess during routine clinical encounters. For example, improvement in trunk postural control among patients with low back pain has been found to be a significant predictor for LBP related pain and disability outcomes.⁶⁷ Moreover, Timed Up and Go test scores have been found to be associated with pain, disability and lumbar mobility measures.^{68,69}

Conclusion

For high-risk patients with low back pain, psychological measures performed better as treatment monitoring tools for disability outcomes when compared to a physical impairment index. Psychological measures were not better predictors of change in pain intensity over baseline pain intensity scores. Findings from this exploratory case series should be interpreted with caution based on small sample size and lack of statistical power which prohibits definitive conclusions to be made on any of the treatment monitoring measures. Future research is needed to determine whether additional psychological factors (e.g. self-efficacy) improve prediction of pain and disability outcomes in high-risk patients.

Acknowledgements

The authors graciously acknowledge the physical therapists that assisted with data collection at participating Brooks Rehabilitation clinical sites including Trent Harrison, Renata Salvatori, Monika Beneciuk, Raine Osborne, Brian Hagist, Tasha Mouton, Timothy Shreve, Jenny Hagist, Sara Cristello, Sara Bertrand, Nate Moore, and Steve Pompilio. JMB and SZG were supported by a grant from the Patient-Centered Outcomes Research Institute (PCS-1402–10867) while working on this manuscript.

Funding:

This study was funded by the 2012 Brooks Rehabilitation Collaborative Grant. This manuscript was written while JMB received support from the National Institutes of Health – National Center for Medical and Rehabilitation Research and National Institute for Neurological Disorders & Stroke (K12 HD055929). JMB and SZG were supported by a grant from the Patient-Centered Outcomes Research Institute (PCS-1402–10867) while working on this manuscript.

References

1. Global Burden of Disease 2015 Disability-Adjusted Life-Years (DALYs) and Healthy Life Expectancy (HALE) Collaborators. Global, regional, and national disability-adjusted life-years (DALYs) for 315 diseases and injuries and healthy life expectancy (HALE), 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*. 10 8 2016;388(10053): 1603–1658. [PubMed: 27733283]
2. Global Burden of Disease 2015 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*. 10 8 2016;388(10053):1545–1602. [PubMed: 27733282]
3. Chou R, Shekelle P. Will this patient develop persistent disabling low back pain? *Jama*. 4 7 2010;303(13):1295–1302. [PubMed: 20371789]
4. Main CJ, George SZ. Psychologically informed practice for management of low back pain: future directions in practice and research. *Physical therapy*. 5 2011;91(5):820–824. [PubMed: 21451091]
5. Keefe FJ, Main CJ, George SZ. Advancing Psychologically Informed Practice for Patients With Persistent Musculoskeletal Pain: Promise, Pitfalls, and Solutions. *Physical therapy*. 5 1 2018;98(5): 398–407. [PubMed: 29669084]
6. Hay EM, Dunn KM, Hill JC, et al. A randomised clinical trial of subgrouping and targeted treatment for low back pain compared with best current care. The STarT Back Trial Study Protocol. *BMC musculoskeletal disorders*. 2008;9:58. [PubMed: 18430242]
7. Hill JC, Whitehurst DG, Lewis M, et al. Comparison of stratified primary care management for low back pain with current best practice (STarT Back): a randomised controlled trial. *Lancet*. 10 29 2011;378(9802):1560–1571. [PubMed: 21963002]
8. Beneciuk JM, George SZ. Pragmatic Implementation of a Stratified Primary Care Model for Low Back Pain Management in Outpatient Physical Therapy Settings: Two-Phase, Sequential Preliminary Study. *Physical therapy*. 8 2015;95(8):1120–1134. [PubMed: 25858972]

9. Hill JC, Dunn KM, Lewis M, et al. A primary care back pain screening tool: identifying patient subgroups for initial treatment. *Arthritis and rheumatism*. 5 15 2008;59(5):632–641. [PubMed: 18438893]
10. Main CJ, Sowden G, Hill JC, Watson PJ, Hay EM. Integrating physical and psychological approaches to treatment in low back pain: the development and content of the STarT Back trial's 'high-risk' intervention (StarT Back; ISRCTN 37113406). *Physiotherapy*. 6 2012;98(2):110–116. [PubMed: 22507360]
11. Hill JC, Fritz JM. Psychosocial influences on low back pain, disability, and response to treatment. *Physical therapy*. 5 2011;91(5):712–721. [PubMed: 21451093]
12. Beneciuk JM, Fritz JM, George SZ. The STarT Back Screening Tool for prediction of 6-month clinical outcomes: relevance of change patterns in outpatient physical therapy settings. *The Journal of orthopaedic and sports physical therapy*. 9 2014;44(9):656–664. [PubMed: 25098194]
13. George SZ, Beneciuk JM, Lentz TA, et al. Optimal Screening for Prediction of Referral and Outcome (OSPRO) for Musculoskeletal Pain Conditions: Results From the Validation Cohort. *The Journal of orthopaedic and sports physical therapy*. 6 2018;48(6):460–475. [PubMed: 29629615]
14. Shimokawa K, Lambert MJ, Smart DW. Enhancing treatment outcome of patients at risk of treatment failure: meta-analytic and mega-analytic review of a psychotherapy quality assurance system. *Journal of consulting and clinical psychology*. 6 2010;78(3):298–311. [PubMed: 20515206]
15. Simon W, Lambert MJ, Harris MW, Busath G, Vazquez A. Providing patient progress information and clinical support tools to therapists: effects on patients at risk of treatment failure. *Psychotherapy research : journal of the Society for Psychotherapy Research*. 2012;22(6):638–647. [PubMed: 22755547]
16. Vlaeyen JW, Morley S. Cognitive-behavioral treatments for chronic pain: what works for whom? *The Clinical journal of pain*. Jan-Feb 2005;21(1):1–8. [PubMed: 15599126]
17. Kazdin AE. Mediators and mechanisms of change in psychotherapy research. *Annual review of clinical psychology*. 2007;3:1–27.
18. Smeets RJ, Vlaeyen JW, Kester AD, Knottnerus JA. Reduction of pain catastrophizing mediates the outcome of both physical and cognitive-behavioral treatment in chronic low back pain. *The journal of pain : official journal of the American Pain Society*. 4 2006;7(4):261–271. [PubMed: 16618470]
19. Mansell G, Hill JC, Main CJ, Von Korff M, van der Windt D. Mediators of Treatment Effect in the Back In Action Trial: Using Latent Growth Modeling to Take Change Over Time Into Account. *The Clinical journal of pain*. 9 2017;33(9):811–819. [PubMed: 27930393]
20. Mansell G, Hill JC, Main C, Vowles KE, van der Windt D. Exploring What Factors Mediate Treatment Effect: Example of the STarT Back Study High-Risk Intervention. *The journal of pain : official journal of the American Pain Society*. 11 2016;17(11):1237–1245. [PubMed: 27565304]
21. Lee H, Hubscher M, Moseley GL, et al. How does pain lead to disability? A systematic review and meta-analysis of mediation studies in people with back and neck pain. *Pain*. 6 2015;156(6):988–997. [PubMed: 25760473]
22. Moseley GL, Nicholas MK, Hodges PW. A randomized controlled trial of intensive neurophysiology education in chronic low back pain. *The Clinical journal of pain*. Sep-Oct 2004;20(5):324–330. [PubMed: 15322439]
23. Louw A, Puente-dura EL, Mintken P. Use of an abbreviated neuroscience education approach in the treatment of chronic low back pain: a case report. *Physiotherapy theory and practice*. 1 2012;28(1):50–62. [PubMed: 21721995]
24. Pincus T, Kent P, Bronfort G, Loisel P, Pransky G, Hartvigsen J. Twenty-five years with the biopsychosocial model of low back pain-is it time to celebrate? A report from the twelfth international forum for primary care research on low back pain. *Spine*. 11 15 2013;38(24):2118–2123. [PubMed: 23970112]
25. Delitto A, George SZ, Van Dillen LR, et al. Low back pain. *The Journal of orthopaedic and sports physical therapy*. 4 2012;42(4):A1–57.
26. Waddell G, Newton M, Henderson I, Somerville D, Main CJ. A Fear-Avoidance Beliefs Questionnaire (FABQ) and the role of fear-avoidance beliefs in chronic low back pain and disability. *Pain*. 2 1993;52(2):157–168. [PubMed: 8455963]

27. Jacob T, Baras M, Zeev A, Epstein L. Low back pain: reliability of a set of pain measurement tools. *Arch Phys Med Rehabil.* 6 2001;82(6):735–742. [PubMed: 11387576]
28. Pfingsten M, Kroner-Herwig B, Leibing E, Kronshage U, Hildebrandt J. Validation of the German version of the Fear-Avoidance Beliefs Questionnaire (FABQ). *Eur J Pain.* 2000;4(3):259–266. [PubMed: 10985869]
29. Staerkle R, Mannion AF, Elfering A, et al. Longitudinal validation of the fear-avoidance beliefs questionnaire (FABQ) in a Swiss-German sample of low back pain patients. *Eur Spine J.* Jul 2004;13(4):332–340.
30. Kovacs FM, Muriel A, Medina JM, et al. Psychometric characteristics of the Spanish version of the FAB questionnaire. *Spine.* 1 1 2006;31(1):104–110. [PubMed: 16395186]
31. Swinkels-Meewisse IE, Roelofs J, Schouten EG, Verbeek AL, Oostendorp RA, Vlaeyen JW. Fear of movement/(re)injury predicting chronic disabling low back pain: a prospective inception cohort study. *Spine.* 3 15 2006;31(6):658–664. [PubMed: 16540870]
32. Swinkels-Meewisse IE, Roelofs J, Verbeek AL, Oostendorp RA, Vlaeyen JW. Fear-avoidance beliefs, disability, and participation in workers and non-workers with acute low back pain. *The Clinical journal of pain.* 1 2006;22(1):45–54. [PubMed: 16340593]
33. Fritz JM, George SZ. Identifying psychosocial variables in patients with acute work-related low back pain: the importance of fear-avoidance beliefs. *Physical therapy.* 10 2002;82(10):973–983. [PubMed: 12350212]
34. Fritz JM, George SZ, Delitto A. The role of fear-avoidance beliefs in acute low back pain: relationships with current and future disability and work status. *Pain.* 10 2001;94(1):7–15. [PubMed: 11576740]
35. George SZ, Fritz JM, Bialosky JE, Donald DA. The effect of a fear-avoidance-based physical therapy intervention for patients with acute low back pain: results of a randomized clinical trial. *Spine.* 12 1 2003;28(23):2551–2560. [PubMed: 14652471]
36. Sullivan M, Bishop S, Pivik J. The Pain Catastrophizing Scale: development and validation. *Psychological Assessment.* 12 1995;7(4):524–532.
37. Crombez G, Eccleston C, Baeyens F, Eelen P. When somatic information threatens, catastrophic thinking enhances attentional interference. *Pain.* 4 1998;75(2–3):187–198. [PubMed: 9583754]
38. Crombez G, Vlaeyen JW, Heuts PH, Lysens R. Pain-related fear is more disabling than pain itself: evidence on the role of pain-related fear in chronic back pain disability. *Pain.* 3 1999;80(1–2):329–339. [PubMed: 10204746]
39. Osman A, Barrios FX, Gutierrez PM, Kopper BA, Merrifield T, Grittmann L. The Pain Catastrophizing Scale: further psychometric evaluation with adult samples. *J Behav Med.* 8 2000;23(4):351–365. [PubMed: 10984864]
40. Woby SR, Roach NK, Urmston M, Watson PJ. Psychometric properties of the TSK-11: a shortened version of the Tampa Scale for Kinesiophobia. *Pain.* 9 2005;117(1–2):137–144. [PubMed: 16055269]
41. George SZ, Valencia C, Zeppieri G, Jr., Robinson ME. Development of a self-report measure of fearful activities for patients with low back pain: the fear of daily activities questionnaire. *Physical therapy.* 9 2009;89(9):969–979. [PubMed: 19608630]
42. Beneciuk JM, Robinson ME, George SZ. Low back pain subgroups using fear-avoidance model measures: results of a cluster analysis. *The Clinical journal of pain.* 10 2012;28(8):658–666. [PubMed: 22510537]
43. Waddell G, Somerville D, Henderson I, Newton M. Objective clinical evaluation of physical impairment in chronic low back pain. *Spine.* 6 1992;17(6):617–628. [PubMed: 1308095]
44. Fritz JM, Piva SR. Physical impairment index: reliability, validity, and responsiveness in patients with acute low back pain. *Spine.* 6 1 2003;28(11):1189–1194. [PubMed: 12782991]
45. Jensen MP, Turner LR, Turner JA, Romano JM. The use of multiple-item scales for pain intensity measurement in chronic pain patients. *Pain.* 9 1996;67(1):35–40. [PubMed: 8895229]
46. Jensen MP, Turner JA, Romano JM, Fisher LD. Comparative reliability and validity of chronic pain intensity measures. *Pain.* 11 1999;83(2):157–162. [PubMed: 10534586]
47. Bolton JE. Accuracy of recall of usual pain intensity in back pain patients. *Pain.* 12 1999;83(3):533–539. [PubMed: 10568862]

48. Childs JD, Piva SR, Fritz JM. Responsiveness of the numeric pain rating scale in patients with low back pain. *Spine*. 6 1 2005;30(11):1331–1334. [PubMed: 15928561]
49. Fritz JM, Irrgang JJ. A comparison of a modified Oswestry Low Back Pain Disability Questionnaire and the Quebec Back Pain Disability Scale. *Physical therapy*. 2 2001;81(2):776–788. [PubMed: 11175676]
50. Hudson-Cook N, Tomes-Nicholson K, Breen A. A revised Oswestry disability questionnaire In: Roland MO, Jenner JR, eds. *Back Pain: New Approaches to Rehabilitation and Education*. New York, NY: Manchester University Press; 1989:187–204.
51. Fairbank JC, Pynsent PB. The Oswestry Disability Index. *Spine*. 11 15 2000;25(22):2940–2952; discussion 2952. [PubMed: 11074683]
52. Roland M, Fairbank J. The Roland-Morris Disability Questionnaire and the Oswestry Disability Questionnaire. *Spine*. 12 15 2000;25(24):3115–3124. [PubMed: 11124727]
53. Ostelo RW, Deyo RA, Stratford P, et al. Interpreting change scores for pain and functional status in low back pain: towards international consensus regarding minimal important change. *Spine*. 1 1 2008;33(1):90–94. [PubMed: 18165753]
54. Portney LG, Watkins MP. *Foundations of Clinical Research: Applications to Practice*. 3rd Edition ed. Upper Saddle River, NJ: Pearson Education; 2009.
55. Burton AK, Tillotson KM, Main CJ, Hollis S. Psychosocial predictors of outcome in acute and subchronic low back trouble. *Spine*. 3 15 1995;20(6):722–728. [PubMed: 7604349]
56. George SZ, Fritz JM, McNeil DW. Fear-avoidance beliefs as measured by the fear-avoidance beliefs questionnaire: change in fear-avoidance beliefs questionnaire is predictive of change in self-report of disability and pain intensity for patients with acute low back pain. *The Clinical journal of pain*. 2 2006;22(2):197–203. [PubMed: 16428956]
57. Pincus T, Smeets RJ, Simmonds MJ, Sullivan MJ. The fear avoidance model disentangled: improving the clinical utility of the fear avoidance model. *The Clinical journal of pain*. Nov-Dec 2010;26(9):739–746. [PubMed: 20842017]
58. Oliveira CB, Franco MR, Demarchi SJ, et al. Psychometric Properties of the Photograph Series of Daily Activities-Short Electronic Version (PHODA-SeV) in Patients With Chronic Low Back Pain. *The Journal of orthopaedic and sports physical therapy*. 9 2018;48(9):719–727. [PubMed: 29792106]
59. Fritz JM, Beneciuk JM, George SZ. Relationship between categorization with the STarT Back Screening Tool and prognosis for people receiving physical therapy for low back pain. *Physical therapy*. 5 2011;91(5):722–732. [PubMed: 21451094]
60. Beneciuk JM, Bishop MD, Fritz JM, et al. The STarT back screening tool and individual psychological measures: evaluation of prognostic capabilities for low back pain clinical outcomes in outpatient physical therapy settings. *Physical therapy*. 3 2013;93(3):321–333. [PubMed: 23125279]
61. Glattacker M, Heyduck K, Jakob T. Yellow flags as predictors of rehabilitation outcome in chronic low back pain. *Rehabilitation psychology*. 8 2018;63(3):408–417. [PubMed: 30024205]
62. Jackson T, Wang Y, Wang Y, Fan H. Self-efficacy and chronic pain outcomes: a meta-analytic review. *The journal of pain : official journal of the American Pain Society*. 8 2014;15(8):800–814. [PubMed: 24878675]
63. Linton SJ, Nicholas M, MacDonald S. Development of a short form of the Orebro Musculoskeletal Pain Screening Questionnaire. *Spine*. 10 15 2011;36(22):1891–1895. [PubMed: 21192286]
64. Lentz TA, Beneciuk JM, Bialosky JE, et al. Development of a Yellow Flag Assessment Tool for Orthopaedic Physical Therapists: Results From the Optimal Screening for Prediction of Referral and Outcome (OSPRO) Cohort. *The Journal of orthopaedic and sports physical therapy*. 5 2016;46(5):327–343. [PubMed: 26999408]
65. Beneciuk JM, Hill JC, Campbell P, Afolabi E, George SZ, Dunn KM, Foster NE. Identifying Treatment Effect Modifiers in the STarT Back Trial: A Secondary Analysis. *J Pain*. 2017 1;18(1):54–65. [PubMed: 27765643]
66. Katzan IL, Thompson NR, George SZ, Passek S, Frost F, Stilphen M. The use of STarT back screening tool to predict functional disability outcomes in patients receiving physical therapy for

low back pain. *Spine J.* 2018 10 9 pii: S1529-9430(18)31158-6. doi: 10.1016/j.spinee.2018.10.002. [Epub ahead of print].

67. Shahvarpour A, Gagnon D, Preuss R, Henry SM, Lariviere C. Trunk postural balance and low back pain: Reliability and relationship with clinical changes following a lumbar stabilization exercise program. *Gait & posture.* 3 2018;61:375–381. [PubMed: 29448220]
68. Gautschi OP, Smoll NR, Corniola MV, et al. Validity and Reliability of a Measurement of Objective Functional Impairment in Lumbar Degenerative Disc Disease: The Timed Up and Go (TUG) Test. *Neurosurgery.* 8 2016;79(2):270–278. [PubMed: 26702840]
69. Coyle PC, Velasco T, Sions JM, Hicks GE. Lumbar Mobility and Performance-Based Function: An Investigation in Older Adults with and without Chronic Low Back Pain. *Pain medicine (Malden, Mass.).* 1 1 2017;18(1):161–168.

Highlights

- FABQ-PA and PCS changes contributed to prediction of 4-week ODI scores.
- Treatment monitoring measure changes did not predict 4-week NPRS scores.
- Physical Impairment Index changes did not contribute to 4-week clinical outcomes.

Table 1.

Baseline characteristics of study sample (n = 23).

Variable	Value
Age (years)	43.7 (12.5)
Sex, female (n) (%)	15 (65.2%)
Race (n) (%)	
Caucasian	16 (69.6%)
Black or African American	5 (21.7%)
Other	2 (8.7%)
Employment status	
Employed	17 (73.9%)
Unemployed	4 (17.4%)
Retired	2 (8.7%)
Work related LBP (yes)	1 (4.3%)
Symptom duration	
14 days	1 (4.3%)
15 to 90 days	10 (43.5%)
91 days	12 (52.2%)
Psychological Measures	
FABQ-PA (potential range, 0 to 24)	17.6 (5.1)
FABQ-W (potential range, 0 to 42)	23.4 (10.7)
PCS (potential range, 0 to 52)	30.3 (11.1)
TSK-11 (potential range, 11 to 44)	30.3 (7.0)
FDAQ (potential range, 0 to 100)	56.5 (20.6)
Clinical Measures	
NPRS (potential range, 0 to 10)	6.0 (2.0)
ODI (potential range, 0 to 100)	44.6 (15.4)
PIT (potential range, 0 to 7)	4.3 (1.6)

Values are reported as mean point estimates (standard deviation) for continuous variables and frequency counts (percentage) for categorical variables.

LBP = low back pain; FABQ-PA = Fear Avoidance Beliefs Questionnaire (physical activity scale); FABQ-W = Fear Avoidance Beliefs Questionnaire (work scale); PCS = Pain Catastrophizing Scale; TSK-11 = Tampa Scale for Kinesiophobia (11 item version); FDAQ = Fear of Daily Activities Questionnaire; NPRS = Numerical Pain Rating Scale; ODI = Oswestry Disability Index; PIT = Physical Impairment Index.

Table 2.

Relationship between psychological and clinical measure change scores at 4 weeks.

	FABQ-W	PCS	TSK-11	FDAQ	PII	NPRS	ODI
FABQ-PA	.637 **	.694 **	.699 **	.592 **	.168	.010	.697 **
FABQ-W		.616 **	.652 **	.715 **	.028	-.021	.219
PCS			.749 **	.574 **	.169	.184	.651 **
TSK-11				.768 **	.185	.220	.455 *
FDAQ					.065	.260	.204
PII						.258	.250
NPRS							.368

**
P < 0.01;

* P < 0.05. FABQ-PA = Fear Avoidance Beliefs Questionnaire (physical activity scale); FABQ-W = Fear Avoidance Beliefs Questionnaire (work scale); PCS = Pain Catastrophizing Scale; TSK-11 = Tampa Scale for Kinesiophobia (11 item version); FDAQ = Fear of Daily Activities Questionnaire; NPRS = Numerical Pain Rating Scale; ODI = Oswestry Disability Index; PII = Physical Impairment Index.

Table 3.

Prediction of pain intensity (NPRS) at 4 weeks.

	Model 1	Model 2	Change from Model 1 to Model 2
Treatment Monitoring Measure	Intake NPRS + Intake Score	Model 1 + Change Score at 4 Weeks	Additional Variance Explained by Change Score at 4 Weeks
FABQ-PA	%R ² = 45.7 Adjusted %R ² = 39.7	%R ² = 46.3 Adjusted %R ² = 36.8	%R ² = 0.5 P = .686
FABQ-W	%R ² = 46.2 Adjusted %R ² = 40.2	%R ² = 46.6 Adjusted %R ² = 37.1	%R ² = 0.4 P = .723
PCS	%R ² = 45.7 Adjusted %R ² = 39.6	%R ² = 49.9 Adjusted %R ² = 41.1	%R ² = 4.2 P = .248
TSK-11	%R ² = 45.7 Adjusted %R ² = 39.6	%R ² = 52.2 Adjusted %R ² = 43.8	%R ² = 6.6 P = .145
FDAQ	%R ² = 45.6 Adjusted %R ² = 39.6	%R ² = 53.5 Adjusted %R ² = 45.3	%R ² = 7.9 P = .107
PII	%R ² = 47.7 Adjusted %R ² = 41.1	%R ² = 52.8 Adjusted %R ² = 43.4	%R ² = 5.1 P = .221

FABQ-PA = Fear Avoidance Beliefs Questionnaire (physical activity scale); FABQ-W = Fear Avoidance Beliefs Questionnaire (work scale); PCS = Pain Catastrophizing Scale; TSK-11 = Tampa Scale for Kinesiophobia (11 item version); FDAQ = Fear of Daily Activities Questionnaire; NPRS = Numerical Pain Rating Scale; PII = Physical Impairment Index.

Table 4.

Prediction of disability (ODI) at 4 weeks.

	Model 1	Model 2	Change from Model 1 to Model 2
Treatment Monitoring Measure	Intake ODI + Intake Score	Model 1 + Change Score at 4 Weeks	Additional Variance Explained by Change Score at 4 Weeks
FABQ-PA	%R ² = 30.6 Adjusted %R ² = 22.9	%R ² = 67.6 Adjusted %R ² = 61.8	%R ² = 37.0 P < .001
FABQ-W	%R ² = 38.0 Adjusted %R ² = 31.1	%R ² = 58.0 Adjusted %R ² = 50.6	%R ² = 20.0 P = .011
PCS	%R ² = 31.7 Adjusted %R ² = 24.1	%R ² = 58.9 Adjusted %R ² = 51.6	%R ² = 27.1 P = .004
TSK-11	%R ² = 31.8 Adjusted %R ² = 24.3	%R ² = 48.1 Adjusted %R ² = 39.0	%R ² = 16.3 P = .034
FDAQ	%R ² = 31.3 Adjusted %R ² = 23.7	%R ² = 46.5 Adjusted %R ² = 37.1	%R ² = 15.2 P = .042

FABQ-PA = Fear Avoidance Beliefs Questionnaire (physical activity scale); FABQ-W = Fear Avoidance Beliefs Questionnaire (work scale); PCS = Pain Catastrophizing Scale; TSK-11 = Tampa Scale for Kinesiophobia (11 item version); FDAQ = Fear of Daily Activities Questionnaire; ODI = Oswestry Disability Index; PII = Physical Impairment Index.