Brazilian Journal of Physical Therapy

<https://www.journals.elsevier.com/brazilian-journal-of-physical-therapy>

SYSTEMATIC REVIEW

Current evidence does not support whole body vibration in clinical practice in children and adolescents with disabilities: a systematic review of randomized controlled trial

Hércules Ribeiro Leite^{a,*}, Ana Cristina Resende Camargos^b,
Vanessa Amaral Mendonça^a, Ana Cristina Rodrigues Lacerda^a,
Bruno Alvarenga Soares^a, Vinicius Cunha Oliveira^a

^a Discipline of Physical Therapy, Universidade Federal dos Vales do Jequitinhonha e Mucuri (UFVJM), Diamantina, MG, Brazil

^b Discipline of Physical Therapy, Universidade Federal de Minas Gerais (UFMG), Belo Horizonte, MG, Brazil

Received 11 January 2018; received in revised form 4 September 2018; accepted 6 September 2018

Available online 19 September 2018

KEYWORDS

Children;
Adolescent;
Disability;
Whole body vibration;
Exercise

Abstract

Background: Whole body vibration has been used alone or combined with other interventions in rehabilitation of children and adolescents with disabilities; however, there is limited evidence to support this approach.

Objectives: To review the strength, quality, and conclusiveness of evidence supporting the use of whole body vibration in children and adolescents with disabilities.

Methods: Electronic database search included Medline, AMED, Embase, Cochrane, SportDiscus, CINAHL and PEDro from the inception to June 2018. Studies investigating the effects of whole body vibration, alone or combined with other interventions, compared to minimal intervention or other interventions were included. The outcomes measured were: body structure and function (lean body mass, bone mineral density, knee muscle strength and overall stability) and activity and participation (gait speed, walking distance, gross motor function, self-care and mobility).

* Corresponding author at: Departamento de Fisioterapia, Programa de Pós-Graduação em Reabilitação e Desempenho Funcional (PPGREab), Centro Integrado de Pesquisa e Pós-Graduação em Saúde – CIPq-Saúde, Universidade Federal dos Vales do Jequitinhonha e Mucuri, Rodovia MGT 367, Km 583, Alto da Jacuba, n° 5000 – CEP 39100-000, Diamantina, MG, Brazil.

E-mail: hercules.leite@ufvjm.edu.br (H.R. Leite).

Results: Fifteen randomized trials involving 403 participants were included. Methodological quality of eligible trials was moderate (mean of 5.5 points on the 10-point PEDro scale). Overall, whole body vibration was no better than minimal intervention. In all comparisons where additional effect of whole body vibration was better than other interventions, the effect size ranged from low to high in the trials, but ranged from very-low to low quality at short and medium-term follow-up. Sensitivity analysis for health condition and low-quality studies showed impact on trunk bone mineral density of additional effect of whole body vibration at medium-term compared to other interventions.

Conclusion: The low to very-low quality of evidence suggests caution in recommending the use of this approach. New studies could change the findings of this review. PROSPERO registration: CRD42017060704.

© 2018 Associação Brasileira de Pesquisa e Pós-Graduação em Fisioterapia. Published by Elsevier Editora Ltda. All rights reserved.

Introduction

Children and adolescents with disabilities are characterized as having impairments in body structure and function, limitation of activity and social restriction in accordance to International Classification of Functioning, Disability and Health (ICF).^{1–6} Disability is a generic term and may involve neurological, musculoskeletal, metabolic and burned-related conditions in the childhood and adolescence.⁷ Considering different levels of disability, these individuals can present low bone mineral density, altered body composition, decreased levels of muscle strength, instability, reduced walking capacity, reduced gross motor function, and activities of daily living performance.^{8–10} Furthermore, children and adolescents with disabling conditions have reduced levels of habitual activity compared with their typically developing peers, as recommended by guidelines.^{11–14} These individuals are limited to engage in some physical activity programs due many barriers, such as sedentary lifestyles, motor deficits, social and recreation opportunities, or low motivation to be physically active.¹⁵ New approaches have been used to minimize deficiencies and activity limitations of these population,¹⁶ such as the use of whole body vibration (WBV).

WBV has become a very popular treatment for disabilities in children,¹⁷ adolescents¹⁸ and adults¹⁹ over the last decade. WBV is a type of training that uses high-frequency mechanical stimuli, which is generated by a vibrating platform and transmitted through body to load bone and stimulate sensory receptors.¹² Previous studies using WBV showed improvements on muscle strength, lean mass, bone mass, stability, gait capacity and gross motor performance.^{20–23} Although WBV is used in clinical practice by physical therapists worldwide, its effectiveness in children and adolescent with disabilities is still unclear due to limited methodological quality of previous studies.^{24–31}

A previous systematic review investigated effects of WBV on health-related physical fitness in children and adolescents with disabilities.^{32,33} Authors reported that WBV may be effective on disabled children and adolescents' health; however, they included low quality studies (i.e. not randomized controlled trials), did not assess risk of bias of the included trials, and meta-analysis was not conducted.³¹ Moreover, the role of WBV in physical rehabilitation in

children and adolescent with disabilities is uncertain because training can be used alone or in conjunction with other interventions,¹⁴ once there is the possibility that the scientific evidence of one intervention differs from the effect of the other and even from the combination of both.³⁴ Therefore, in an attempt to address these gaps, the research question for this systematic review of randomized controlled trials was: is WBV alone or combined with other interventions more effective than minimal or other interventions in children and adolescent with disabilities?

Methods

This study was a systematic review with meta-analysis. Review followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement,³⁵ and its protocol was prospectively registered at International prospective register of systematic reviews (PROSPERO)³⁶ (CRD42017060704).

Identification and selection of the studies

Search strategies were conducted on Medline, AMED, Embase, Cochrane, SportDiscus, CINAHL and PEDro in September 2016 and updated in June 2018, without language or date restrictions. Descriptors used in our search strategy were related to "randomized controlled trials", "children", "adolescent" and "whole body vibration". We also hand searched identified systematic reviews in the field.

Screening of identified titles and abstracts, and assessment of potential full-texts using our eligibility criteria was conducted by two independent reviewers (ACRC and HRL), with discrepancies resolved by a third reviewer (VCO).

Assessment of characteristic of studies

Quality

The methodological quality of the included trials was assessed by extracting the PEDro score from the Physiotherapy Evidence Database (www.pedro.org.au).³⁷ If the trial was not included in the database the score was assessed by two independent reviewers (ACRC and HRL) using the 0–10 PEDro scale. A third reviewer (VCO) resolved

discrepancies. The PEDro scale assesses 10 criteria and final score ranges from 0 to 10, with greater scores meaning greater methodological quality.³⁷ Reviewers were trained a priori.

Participants

In the current review, disability was defined as a complex phenomenon which covers impairments, activity limitations and participation restrictions according to definition of World Health Organization (WHO).⁶ Studies were eligible if they included randomized controlled trials that investigated the effects of WBV in children and adolescent with disabilities, such as neurological, musculoskeletal, metabolic and burn-related conditions.

Intervention

The experimental intervention was WBV which was defined as any mechanical stimulus characterized by an oscillatory motion generated by vibrating devices using two different systems: (a) reciprocating vertical displacements on the left and right side of a fulcrum; (b) the whole plate oscillating uniformly up and down.^{32,33} Trials were considered for inclusion if they compared WBV to minimal intervention (i.e. no intervention, waiting list, placebo and sham) or additional effect of WBV compared to other interventions.

Outcomes

We were interested on primary (related to body and structure and function: lean body mass, bone mineral density, muscle strength and overall stability) and secondary outcomes (related to activity and participation: gait speed/walking distance, gross motor function, self-care and mobility), according to Classification of Functioning and Disability and Health.⁶

Data analysis

Data were extracted by two independent reviewers (ACRC and HRL), with a third reviewer (VCO) resolving potential discrepancies. Extracted data included characteristics of included studies (i.e. participants, interventions, outcomes, and type of platform), and outcome measures (i.e. means, standard deviations, and sample sizes). Outcome measures were extracted for short-, medium-, and long-term effects. Short-term effect was considered up to twelve weeks after baseline, medium-term was considered over twelve weeks but less than twelve months after baseline, and long-term effect was considered at least twelve months after baseline. When more than one timepoint was available for short-term effect, we considered the one closer to the end of the intervention. When outcome measures were not provided in the study, data were imputed following the Cochrane recommendations.³⁸ For bone mineral density, assessed regions varied across studies, so we arbitrarily decided to extract data, firstly, for leg or trunk. Then, data for other regions were considered in studies assessing other regions (i.e. tibia, femur, lumbar, and total bone mineral density).

Outcome data were pooled in meta-analysis using different health conditions and results were presented as standardized mean difference (SMD) with 95% confidence

intervals (CIs). Between-trial heterogeneity was assessed using I^2 , and random-effects model was used considering clinical heterogeneity and when appropriate (i.e. $I^2 \geq 50\%$).³⁸ To judge the clinical relevance effects of changes provided by WBV, effect sizes were assessed using Cohen's benchmarks: $d < 0.2$ for small; $d > 0.5$ for medium; and > 0.8 for large effects.³⁹ There are systematic reviews that pooled different disabilities⁴⁰ and it is known that this can impact on outcomes.⁴¹ Thus, we conducted sensitivity analysis to investigate impact of methodological quality and health condition on estimated effects.⁴² Software Comprehensive Meta-analysis 2.2.04 (Biostat, Englewood, NJ) was used for all analysis.

The Grading of Recommendations Assessment, Development and Evaluation (GRADE) system was used to summarize the overall quality of the current evidence.⁴³ The four levels of the GRADE system range from high-quality to very low-quality evidence.⁴³ Quality of evidence scoring for each outcome using GRADE started at high-quality evidence, which was downgraded by one point if one of the following criteria was present: (i) poor methodological quality for average PEDro score less than five out of 10; (ii) inconsistency of estimates among trials for $I^2 \geq 50\%$, heterogeneity or absence of pooling; (iii) indirectness when participants were selected by no reliable methods or when their inclusion criteria in any of the analyzed trials was not clear; (iv) imprecision for samples < 300 participants for each outcome⁴⁴; and (v) publication bias or when its analysis was not possible due to small number of trials (i.e. $n < 10$ trials).⁴⁵ Two independent reviewers (HRL and ACRC) assessed current evidence and a third reviewer (VCO) resolved potential discrepancies.

Results

Flow of the studies through the review

Searches identified 2012 references, 73 potential full-texts were assessed for our eligibility criteria, and 15 original studies were included in this review (Fig. 1).

Characteristics of the included studies

Characteristics of included trials are in Table 1. All 15 included randomized controlled trials were published in English between 2004 and 2017.

Quality

Mean methodological quality of included trials was 5.5 out of 10 on the PEDro scale (scores ranged from 4.0 to 8.0). Methodological issues included: absence of concealed allocation ($n = 8$ studies, 53%); absence of report of similar groups at baseline ($n = 2$ studies, 14%); blinding of therapist ($n = 0$ study, 0%), participant ($n = 0$ study, 0%) and assessors ($n = 5$ studies, 33%); more than 15% dropouts ($n = 7$ studies, 53%); and intention-to-treat analysis ($n = 5$ studies, 33%). Detailed assessment of methodological quality of included trials is in Table 2.

Table 1 Descriptive data of the included trials ($n = 15$).

Trial	Source	Participants	Intervention	Outcome measures	Type of platform
WBV compared to minimal intervention					
<i>Neurological conditions</i>					
González-Aguero et al. ⁵⁰ 2013	Adolescents with Down syndrome, boys and girls, recruited from different schools in Aragon (Spain).	<p>$n = 30$</p> <p>Experimental: $n = 16$ Age (yr) mean (SD): 15.3 (2.6)</p> <p>Control: $n = 14$ Age (yr) mean (SD): 15.8 (3.0)</p>	<p>Experimental: WBV 25–30 Hz, 2 mm, squatting position (15–20 min, 3/wk $\times$ 20 wks)</p> <p>Control: no intervention</p>	<p>Lean and fat body mass = DXA</p> <p><i>Medium term effects = 20 wks</i></p>	Power Plate® Pro5 = Vertical vibration platform
Matute-Llorente et al. ⁵⁵ 2015	Adolescents with Down syndrome, boys and girls, recruited from three schools and institutions of Aragón (Spain).	<p>$n = 30$</p> <p>Experimental: $n = 16$ Age (yr) mean (SD): 15.2 (2.5)</p> <p>Control: $n = 14$ Age (yr) mean (SD): 15.5 (3.0)</p>	<p>Experimental: WBV, 25–30 Hz, 2 mm, squatting position, bent knees at 90° (5–10 min, 3/wk $\times$ 20 wks)</p> <p>Control: normal daily routine</p>	<p>BMD and BMC = DXA</p> <p>Bone strength indexes, bone morphometry, vBMD, and trabecular bone microarchitecture of tibia and radius = Stratec XCT-2000 L pQCT scanner</p> <p><i>Medium term effects = 20 wks</i></p>	Power Plate® Pro5 = synchronous vibration platform
<i>Metabolic conditions</i>					
Erceg et al. ⁵¹ 2015	Overweight, Latino boys, recruited from Los Angeles area.	<p>$n = 32$</p> <p>Experimental: $n = 16$ Age (yr) mean (SD): 9 (1)</p> <p>Control: $n = 16$ Age (yr) mean (SD): 9 (1)</p>	<p>Experimental: WBV, 30–40 Hz, low and high amplitude, squat, wide-stance squat, calf raise, lunge and modified push-up (3–20 min, 3/wk $\times$ 10 wk)</p> <p>Control: normal daily routine for the 10 wk trial period</p>	<p>BMD and BMC = DXA</p> <p>Bone biomarkers = osteocalcin and collagen type I C-telopeptide</p> <p><i>Short term effects = 10 wks</i></p>	NEXTgeneration Power Plate® = vibration platforms simultaneously oscillates in the vertical, anterior–posterior and mediolateral planes

Table 1 (Continued)

Trial	Source	Participants	Intervention	Outcome measures	Type of platform
WBV compared to other intervention					
<i>Neurological conditions</i>					
Eid ²⁰ 2015	Children with Down syndrome, boys and girls, recruited from outpatient clinic of Faculty of Physical Therapy, Cairo, Egypt.	<i>n</i> = 30 Experimental: <i>n</i> = 15 Age (yr) <i>mean</i> (<i>SD</i>): 8.9 (0.7) Control: <i>n</i> = 15 Age (yr) <i>mean</i> (<i>SD</i>): 9.3 (0.8)	Experimental: WBV, 25–30 Hz, 2 mm, standing with knees semi-squats (30°) (5–10 min, 3/wk × 24 wks) + Physical Therapy program (1 h, 3/wk × 24 wks) Control: Physical Therapy program (1 h, 3/wk × 24 wks)	Overall stability index, mediolateral stability index and anteroposterior stability index = Biodex Stability System Muscle strength (knee extension and flexion) = Handheld dynamometer <i>Medium term effects</i> = 24 wks	Vibraflex Home Edition II = Side-to-side alternating vertical sinusoidal vibration
El-Shamy and Mohamed ²¹ 2012	Children with spastic diplegic cerebral palsy, boys and girls, recruited from outpatient clinic at the Faculty of Physical Therapy, Cairo, Egypt.	<i>n</i> = 30 Experimental: <i>n</i> = 15 Age (yr) <i>mean</i> (<i>SD</i>): 11.7 (0.8) Control: <i>n</i> = 15 Age (yr) <i>mean</i> (<i>SD</i>): 11.6 (1.1)	Experimental: WBV, static standing position with the knees bent, 25 Hz, 1.7 mm (10 min/day, 5/wk × 24 wks) + Physical Therapy Exercise Program based on neuro-developmental technique Control: Physical Therapy Exercise Program based on neuro-developmental technique	Femur, lumbar spine and total body BMD = DXA <i>Medium term effects</i> = 24 wks	Not specified
El-Shamy ⁴⁶ 2014	Children with spastic diplegic cerebral palsy, boys and girls, recruited from outpatient clinic at the Faculty of Physical Therapy, Cairo, Egypt.	<i>n</i> = 30 Experimental: <i>n</i> = 15 Age (yr) <i>mean</i> (<i>SD</i>): 9.7 (1.2) Control: <i>n</i> = 15 Age (yr) <i>mean</i> (<i>SD</i>): 9.9 (1.1)	Experimental: WBV, 12–18 Hz, 2–4 mm, standing with knees flexed (10–45°) (9 min, 5/wk × 12 wks) + Traditional Physical Therapy Program (1 h, 5/wk × 12 wks) Control: Traditional Physical Therapy Program (1 h, 5/wk × 12 wks)	Muscle strength (knee extension) = Isokinetic dynamometer Overall stability index, mediolateral stability index and anteroposterior stability index = Biodex Stability System <i>Short term effects</i> = 12 wks	Vibraflex Home Edition II = Side-to-side alternating vertical sinusoidal vibration

Table 1 (Continued)

Trial	Source	Participants	Intervention	Outcome measures	Type of platform
Ibrahim et al. ²³ 2014	Children with diplegic spastic cerebral palsy, boys and girls, recruited from the outpatient clinic of the College of Physical Therapy, Cairo University.	<i>n</i> = 30 Experimental: <i>n</i> = 15 Age (yr) <i>mean</i> (<i>SD</i>): 9.6 (1.4) Control: <i>n</i> = 15 Age (yr) <i>mean</i> (<i>SD</i>): 9.6 (1.4)	Experimental: WBV, 12–18 Hz, 4–6 mm, with the knees slightly bent (9 min, 3/wk × 12 wks) + Traditional Physical Therapy Program (1 h, 3/wk × 12 wks) Control: Traditional Physical Therapy Program (1 h, 3/wk × 12 wks)	Knee extensor muscle strength = Handheld dynamometer Spasticity of hip adductors, knee extensors, and ankle plantar flexors = Ashworth scale Walking distance = 6-min walking test Balance in basic mobility maneuvers = TUG Gross motor function = GMFM-88 <i>Short term effects</i> = 12 wks	Power Plate = Side-alternating vibration platform, unsynchronized multidimensional oscillations along the sagittal axis
Ko et al. ⁴⁷ 2016	Children with spastic cerebral palsy, boys and girls, Republic of Korea.	<i>n</i> = 24 Experimental: <i>n</i> = 12 Age (yr) <i>mean</i> (<i>SD</i>): 9.4 (2.7) Control: <i>n</i> = 12 Age (yr) <i>mean</i> (<i>SD</i>): 9.5 (2.2)	Experimental: WBV, 20–24 Hz, 1–2 mm, with their knees flexed at 30° (9 min, 2/wk × 3 weeks) + Traditional Physical Therapy Program (30 min, 2/wk × 3 wks) Control: Traditional Physical Therapy Program (30 min, 2/wk × 3 wks)	Overall stability = Tetrax Interactive Balance System Gait speed = 10 m walk test Joint Position Sense = Tiltmeter (iPhone application) Gait analysis = OptoGait System <i>Short term effects</i> = 3 wks	Galileo System vibration unit = Side-to-side alternating vertical sinusoidal vibration

Table 1 (Continued)

Trial	Source	Participants	Intervention	Outcome measures	Type of platform
Lee and Chon ²² 2013	Children with cerebral palsy with spastic diplegia or quadriplegia, boys and girls, recruited from a local special school and pediatric rehabilitation center.	<i>n</i> = 30 Experimental: <i>n</i> = 15 Age (yr) <i>mean</i> (<i>SD</i>): 10.0 (2.3) Control: <i>n</i> = 15 Age (yr) <i>mean</i> (<i>SD</i>): 9.7 (2.6)	Experimental: WBV, 5–25 Hz, 2–6 mm, squat ranging from 30 to 100 degrees of knee flexion (18 min, 3/wk × 8 wks) + Traditional Physical Therapy Program (30 min, 3/wk × 8 wks) Control: Conventional physical therapy program (30 min, 3/wk × 8 weeks)	Gait speed, stride length, cycle time and sagittal kinematics = Three-dimensional gait analyses (Qualisys) Leg muscle thickness of the leg muscles = ultrasound image <i>Short term effects</i> = 8 wks	Galileo System = Side-to-side alternating vertical sinusoidal vibration
Ruck et al. ⁴⁸ 2010	Children with cerebral palsy, boys and girls, recruited of a primary school for children with special needs (Ecole Victor-Doré, Montreal).	<i>n</i> = 20 Experimental: <i>n</i> = 10 Age (yr) <i>mean</i> : 8.3 Control: 10 Age (yr) <i>mean</i> : 8.1	Experimental: WBV, 12–18 Hz, 2–6 mm, knees and hips between 10° and 45° (9 min, 5/wk × 24 wks) + School physical therapy program Control: School physical therapy program	Gross motor function = GMFM Gait speed = 10 m walk test BMD of the lumbar spine and femur = DXA <i>Medium term effects</i> = 24 wks	Vibraflex Home Edition II® = Side-to-side alternating vertical sinusoidal vibration
Stark et al. ⁴² 2016	Toddlers with cerebral palsy, boys and girls, recruited through the Children's Hospital, University of Cologne, Germany, or via cooperating centers.	<i>n</i> = 24 Experimental: <i>n</i> = 12 Age (months) <i>mean</i> (<i>SD</i>): 18.6 (3.2) Control: <i>n</i> = 12 Age (months) <i>mean</i> (<i>SD</i>): 19.4 (3.2)	Experimental: WBV 12–22 Hz, 2.5 mm, in three exercises: (A) standing still or alternately squatting and standing up, (B) sitting on the platform and (C) "four point position" (9 min, 10/wk × 14 weeks) + Regular physical therapy Control: Regular physical therapy	Gross motor function = GMFM-66 Activities of daily living = PEDI in the self-care and mobility domains Cognitive Development = Bayley II Mental Scale <i>Medium term effects</i> = 14 wks <i>Follow-up</i> = 14 wks	Galileo® system combined with a tilt table = sinusoidal vibration, alternately on the right and left side

Table 1 (Continued)

Trial	Source	Participants	Intervention	Outcome measures	Type of platform
Ward et al. ⁴⁹ 2004	Pre-or post-pubertal neurologic disabled, ambulant children, boys and girls, indicated by consultant community pediatricians.	<i>n</i> = 20 Experimental: <i>n</i> = 10 Age (yr) <i>mean</i> (<i>SD</i>): 6.9 (2.4) Control: <i>n</i> = 10 Age (yr) <i>mean</i> (<i>SD</i>): 11.2 (4.7)	Experimental: WBV 90 Hz, standing (10 min/day, 5/wk × 24 wks) Control: standing on the platform without vibration (10 min/day, 5/wk × 24 wks)	Spine and tibial BMD, diaphyseal cross-sectional bone area, periosteal bone circumference, polar moment of inertia, cortical thickness, and muscle area = 3-D QCT scanner <i>Medium term effects</i> = 24 wks	Vertical ground-based vibration
<i>Musculoskeletal conditions</i>					
Högler et al. ⁵³ 2017	Children with mild to moderate osteogenesis imperfecta, boys and girls, recruited from specialist clinics at tertiary Children's Hospitals in Birmingham, Sheffield and Manchester, as well as through an advertisement placed on the Brittle Bone Society website.	<i>n</i> = 24 Experimental: <i>n</i> = 12 Age (yr) <i>mean</i> : 9.38 Control: <i>n</i> = 12 Age (yr) <i>mean</i> : 6.49	Experimental: WBV, 20–25 Hz, 2–4 mm, knees bent (10–45°, semi-squat and squat position (18 min, 2/day × 20 wks) + Regular physical therapy Control: Regular physical therapy	Lean mass and BMD = DXA Tibial BMD = pQCT scanner Walking distance = 6-min walking test (6MWT) <i>Medium term effects</i> = 20 wks	Galileo M TM = Side-to-side alternating vertical sinusoidal (rotational) vibrations around a fulcrum in the mid-section of the plate.
El-Shamy ⁵² 2017	Boys with hemophilia type A, selected from Al-Noor Hospital, Makkah, Saudi Arabia.	<i>n</i> = 30 Experimental: <i>n</i> = 15 Age (yr) <i>mean</i> (<i>SD</i>): 12.22 (2.33) Control: <i>n</i> = 15 Age (yr) <i>mean</i> (<i>SD</i>): 11.90 (2.74)	Experimental: WBV, 30–40 Hz, 2–4 mm, semi-squats (12–15 min, 3/wk × 12 wks) + Conventional physical therapy program (1 h/d, 3/wk, 12 wks) Control: Conventional physical therapy program (1 h/d, 3/wk, 12 wks)	Muscle strength (knee extension) = Isokinetic dynamometer Femur and lumbar spine BMD = DXA Walking distance = 6-min walking test <i>Short term effects</i> = 12 wks	Power Plate Next Generation Vibration Plat-form = simultaneously oscillates in the vertical, anterior-posterior, and mediolateral planes, predominantly vertical vibration

Table 1 (Continued)

Trial	Source	Participants	Intervention	Outcome measures	Type of platform
<i>Burn-related conditions</i> Edionwe et al. ⁵⁴ 2016	Severely burned children ($\geq 30\%$ total body surface area burns), boys and girls, recruited immediately after discharge from the hospital.	<i>n</i> = 19 Experimental: <i>n</i> = 9 Age (yr) <i>mean (SD)</i> : 11.7 (3.7) Control: <i>n</i> = 10 Age (yr) <i>mean (SD)</i> : 13.1 (4.0)	Experimental: WBV, 30–40 Hz, 2–4 mm, semi-squats (12–15 min, 5/wk $\times$ 6 wk) + Progressive resistance exercise (3/wk) and aerobic conditioning (30 min to 1 h, 5/wk) Control: Progressive resistance exercise (3/wk) and aerobic conditioning (30 min to 1 h, 6/wk)	Lean body mass, BMD and BMC = DXA Muscle strength (knee extension and flexion) = isokinetic dynamometer <i>Short term effects</i> = 6 wks	Power Plate Next Generation = vibration platform simultaneously oscillates in the vertical, anterior–posterior and mediolateral planes

SD, standard deviation; WBV, whole body vibration; wk, week; wks, weeks; DXA, dual-energy X-ray absorptiometry; BMD, bone mineral density; BMC, bone mineral content; vBMD, volumetric bone mineral density; TUG, timed up and go test; GMFM, gross motor function measure; PEDl, pediatric evaluation disability inventory.

Table 2 PEDro scores of included trials ($n = 15$).

Trial	Random allo- cation	Concealed allo- cation	Groups similar at baseline	Participant blind- ing	Therapist blind- ing	Assessor blind- ing	Adequate follow- up	Intention to-treat analysis	Between group dif- ference	Point estimate and vari- ability	Total (0-10)
Edionwe et al. ⁵⁴ 2016	Y	N	Y	N	N	N	Y	N	Y	Y	5
Eid ²⁰ 2015	Y	Y	Y	N	N	Y	Y	Y	Y	Y	8
El-Shamy and Mohamed ²¹ 2012	Y	N	Y	N	N	N	N	N	Y	Y	4
El-Shamy ⁴⁶ 2014	Y	Y	Y	N	N	Y	Y	Y	Y	Y	8
El-Shamy ⁵² 2017	Y	Y	Y	N	N	N	Y	Y	Y	Y	7
Erceg et al. ⁵¹ 2015	Y	N	Y	N	N	N	N	N	Y	Y	4
González-Aguero et al. ⁵⁰ 2013	Y	N	Y	N	N	N	N	N	Y	Y	4
Hogler et al. ⁵³ 2017	Y	Y	Y	N	N	N	N	N	Y	Y	5
Ibrahim et al. ²³ 2014	Y	N	Y	N	N	N	N	N	Y	Y	4
Ko et al. ⁴⁷ 2016	Y	N	N	N	N	N	Y	N	Y	Y	4
Lee and Chon ²² 2013	Y	Y	Y	N	N	Y	Y	N	Y	Y	7
Matute-Llorente et al. ⁵⁵ 2015	Y	N	Y	N	N	N	N	N	Y	Y	4
Ruck et al. ⁴⁸ 2010	Y	Y	Y	N	N	N	N	N	Y	Y	5
Stark et al. ⁴² 2016	Y	Y	Y	N	N	Y	Y	Y	Y	Y	8
Ward et al. ⁴⁹ 2004	Y	N	N	N	N	Y	Y	Y	Y	Y	6
Total	15	7	13	0	0	5	8	5	15	15	5.5/10

N, no; Y, yes.

Figure 1 Flowchart of the trial selection process. AMED, Allied and Complementary Medicine Database; CINAHL, Cumulative Index to Nursing and Allied Health Literature; EMBASE, Excerpta Medica Database; PEDro, Physiotherapy Evidence Database; SportDiscus, Sport Information Resource Center.

Health conditions

Investigated participants included children with neurological (cerebral palsy^{21–23,42,46–49} and Down syndrome^{20,50}), metabolic (overweight),⁵¹ musculoskeletal (hemophilia⁵² and osteogenesis imperfecta⁵³) and burn-related conditions.⁵⁴ Average age ranged from 15.8 months⁵⁰ to 18.6 years old.⁴²

Intervention

Only three trials investigated the effect of WBV compared to minimal intervention^{49–51} and twelve trials investigated additional effects of WBV compared to other intervention.^{20–23,42,46–48,52–55} Trials used different vibrating platforms and dosage parameters. Platform frequencies ranged from 5²² to 90 Hz⁴⁹ and amplitude was less than 6 mm. WBV programs were static (duration per session ranged from 5 to 20 min or dynamic (duration per session ranged from 3 to 20 min, two to five times per week, with total duration ranging from three to twenty four weeks).

Outcomes

The investigated primary outcomes included those related to body structure and function (i.e. lean body mass, bone mineral density, knee strength extension and overall stability) and secondary outcomes those related to activity and participation (i.e. gait speed, walking distance, gross motor function, self-care and mobility). All trials reported short- and medium-term effects only.

Effectiveness of whole body vibration

Effects of WBV compared with minimal intervention and its additional effects to other intervention were analyzed separately by health condition, and results were presented by the ICF's domains: Fig. 2 presents the results for outcomes related to the body structure and function domain (our primary outcomes); and Fig. 3 presents the results for

outcomes related to the activity and participation domains (our secondary outcomes).

Effects of whole body vibration compared to minimal intervention

Body structure and function

Only one trial including 24 children with Down syndrome⁵⁰ showed a large effect of WBV on lean body mass at medium-term follow-up when compared to minimal intervention, SMD was 2.0 (95% CI 1.0–3.0). Other two trials including overweight participants⁵¹ and Down syndrome⁵⁵ showed no effects on total bone mineral density and leg bone mineral density at short- and medium-term follow-up, respectively. SMDs were respectively: -0.2 (95% CI -1.1 to 0.7)⁵¹ and -0.3 (95% CI -1.1 to 0.5).⁵⁵ Current evidence showing a large effect of WBV on lean body mass at medium-term in children with Down syndrome is very-low quality. Furthermore, evidence that additional effect of WBV does not increase total and leg bone mineral density at short- and medium-terms in overweight participants and Down syndrome was very-low as well.

Additional effects of whole body vibration compared to other intervention

Neurological condition

Body structure and function impairment (Fig. 2): pooled estimates showed no additional effects of WBV on leg bone mineral density at medium-term follow up; SMD was -0.1 (95% CI -1.2 to 1.0). However, pooled estimates for trunk bone mineral density showed additional effect at medium-term follow up; SMD was 0.6 (95% CI 0.1 – 1.0). For knee strength extension, the pooled estimates [SMD = 5.1 (95% CI 4.1 – 6.1)] and SMD from one study 0.1 (95% CI 0.2 – 1.7) showed additional effects of WBV on short- and medium-term follow up, respectively. Taken together, the evidence was low quality and indicated that additional effect of WBV provides a medium effect size on trunk bone mineral density, and low and large effect on knee strength extension at

Figure 2 Pooled results of additional effect of whole body vibration (WBV) compared to other interventions on structure and body function impairment. Right side favors WBV added to other interventions.

short and medium-term follow-up, respectively. SMD from one study showed additional effects of WBV on overall stability 1.4 (95% CI 0.6–2.1) at medium-term follow up, but the pooled estimates did not show significant effects at short-term follow up; SMD was 0.4 (95% CI –1.4 to 2.3). The strength of the evidence ranged from low to very-low quality and indicated a large effect size that additional effect of WBV improve overall stability at medium-term follow up but not in short.

Activity and participation limitation (Fig. 3): Pooled estimates showed additional effects of WBV on gait speed at short-term follow up; SMD was 1.2 (95% CI 0.7–1.8). However, one study did not find a positive effect of WBV on gait speed at medium term follow up; SMD was 1.0 (95% CI –0.0 to 2.1). The evidence, which ranged from low to very-low quality, indicated that additional effect of WBV provides a large effect on gait speed at short-term follow up compared to other interventions. SMDs from individuals studies

Figure 3 Pooled results of additional effect of whole body vibration (WBV) compared to other interventions on activity and participation limitation. Right side favors WBV added to other interventions.

investigating cerebral palsy showed a positive effect of WBV on walking distance 3.7 (95% CI 2.4–4.6), standing 1.1 (95% CI 0.4–1.9) and running, walking and jumping gross motor function domains 3.5 (2.4–4.6) at short-term follow up. However, the SMDs from individual studies did not find any effect at medium-term follow up for total gross motor function 0.1 (95% CI –0.7 to 0.9), self-care 0.2 (95% CI –0.5 to 1.0), mobility 0.6 (95% CI –0.2 to 1.4), standing 0.3 (95% CI –0.7 to 1.3) and walking, running and jumping 0.7 (95% CI –0.3 to 1.7) at medium-term follow-up. Taken together, the strength of the evidence was very-low quality that additional effects of WBV provides a large effect on walking distance and gross motor function domains at short-term follow up.

Musculoskeletal condition

Body structure and function impairment (Fig. 2): SMDs from individual studies showed a large effect of additional effect of WBV on lean body mass 1.1 (95% CI 0.2–2.1) in children with osteogenesis imperfecta at medium-term follow-up. For leg (SMD=2.9 [95% CI 1.8–3.9]; –0.1 [95% CI –1.0 to 0.8]) and trunk bone mineral density (SMD=2.9 [95% CI 1.8–3.9]; 0.1 [95% CI –0.7 to 1.0]) the studies showed a large effect of WBV at short-term (hemophilia) and medium-term (osteogenesis imperfecta) follow-up, respectively. One study showed no effect of WBV on knee strength extension in children with hemophilia at short-term follow up; SMD was 6.6 (95% CI 4.8–8.5). The quality of evidence was very low.

Activity and participation limitation (Fig. 3): SMDs from individual studies showed a large effect of

additional effect of WBV on walking distance 1.1 (95% CI 0.3–1.9) at short term follow up in children with hemophilia, but no effect 0.0 (95% CI –0.4 to 1.8) at medium term follow-up in children with osteogenesis imperfecta compared to other interventions. The quality of evidence was very low.

Burn-related condition

Body structure and function impairment: SMD from one study showed a large effect of additional WBV on leg –1.3 (95% CI –2.3 to –0.3) and trunk lean mass –1.6 (–2.6 to –0.6), and trunk bone mineral density 1.7 (95% CI 0.6–2.7), but not for knee strength extension –3.4 (95% CI –4.8 to –2.0) and leg mineral density 0.7 (95% CI –0.2 to 1.7) in burned children at short-term follow-up. The quality of evidence was very low.

Sensitivity analysis

The sensitivity analysis investigated the impact of poor methodological quality (removing trials with PEDro score <5 out of 10) and health condition (removing heterogeneous disabilities). Methodological quality suggested impact on trunk bone mineral density [SMD=0.3 (–0.3 to 1.0)], but not on leg mineral density [SMD=–0.7 (95% CI –1.5 to 0.0)] at medium-term follow up, and knee strength extension [SMD=5.1 (95% CI 3.6–6.6)], overall stability [SMD=–1.4 (95% CI –2.2 to –0.6)] and gait speed [SMD=–1.4 (95% CI –2.2 to –0.6)] at short term-follow up in children with neurological conditions. Health condition sensibility analysis showed impact on trunk bone mineral density [SMD=0.5

(95% CI -0.0 to 1.1]), but not for leg mineral density [SMD = -0.1 (95% CI -2.1 to 1.9) at medium-term follow up in children with neurological condition.

Discussion

This review aimed to summarize the current evidence of the effects of WBV, alone or combined, on deficiencies and limitations in children and adolescents with disabilities at short and medium-term follow-up. The estimates results were obtained from three studies comparing WBV versus minimal intervention and twelve studies comparing additional effect of WBV versus other interventions. In general, WBV compared to minimal intervention either showed no significant benefit in deficiencies of body structure and function, or the quality evidence was very-low quality to be clinically worthwhile. Pooled results indicated that additional effect of WBV provides a low to high effect size on deficiencies and activity limitations compared to other interventions, but these trials showed very-low to low quality evidence.

It has been speculating that WBV stimulates a moving pattern which activates spinal neurons leading to compensatory muscle contraction in leg and trunk.⁵⁶ The muscle performance benefit is probably caused by neurogenic pathways potentiation involving spinal reflexes and muscle activation, called tonic vibration reflex.^{57,58} The mechanism causing strength and power increases from acute vibration is still being debated. However, the current proposal is that vibration provides a mechanical stimulus that causes the muscle fibers to stretch, thereby evoking a natural stretch, which enhances the neuromuscular function through neurogenic excitability and recruitment.^{57,59,60} Thus, this neural activation are related to muscle improvement, similar to those neural changes seen after conventional resistance and power training.^{57,61} Then, WBV training have been associated with increased muscle strength in disabled children with neurological and musculoskeletal conditions.⁵³ Beyond vibration training (vertical or rotation) to be able to improve peak muscle forces (mechanical loading), increase of bone mass are expected probably due to the mechanostat theory.^{56,62} The WBV approach would serve as an anabolic stimulus impeding bone loss in children with neurologic, musculoskeletal and burn-related conditions.^{21,49} In another way, the high-frequency mechanical stimuli of the WBV which are generated by a vibrating platform stimulate sensory receptors⁵⁰ also may be considered appropriate to improve the overall stability in neurological conditions.²⁰

Although secondary outcomes such as gait capacity (gait speed and walking distance) and gross motor function (standing dimension and walking, running and jumping dimension) improved in children with neurologic conditions,^{22,23,48} the extent of its effect and the underlying mechanism are unknown. The increase of primary outcomes like bone mass and muscle strength may be associated.^{56,63}

Due to small number of studies, the selected studies compared WBV with a large range of other interventions (i.e., progressive resistance exercise, aerobic conditioning and conventional physical therapy), disabilities (i.e., cerebral palsy, Down syndrome, osteogenesis imperfecta, hemophilia, overweight and burned children) and protocols settings. Despite the comparison of different outcomes and

protocols, the trials typically showed no significant difference in outcomes between the groups, or a small to high effect in favor of WBV, but the high effect sized found was too wide. Furthermore, the sensitivity analysis of the present study (i.e., by health condition and methodological quality) revealed no impact on the majority of the outcomes, except for trunk bone mineral density in children with neurological conditions. Taking together, the findings limits to include some clinical worthwhile effects.

Previous systematic review³³ showed effects of WBV on health-related physical fitness (body composition, cardiorespiratory fitness, flexibility, muscular strength, and muscular endurance) of children and adolescents with disabilities. Other systematic reviews also suggested positive effects of WBV in children with cerebral palsy,^{27,64,65} osteogenesis imperfecta,⁶⁶ syndrome de down³³ and Duchenne.⁶⁷ Findings for some of the investigated outcomes of interest were not consistent with ours and a potential explanation is the methods. Previous studies did not follow PRISMA statement,³⁵ they included low-quality studies (i.e. not randomized controlled trials),^{32,33} none effect size was estimated for clinical relevance of their findings^{32,33,64-66} they did not investigate the impact of poor methodological quality on the estimated effects,^{27,32,33,65,66} which compromises their clinical implications.

This present study throughout a high sensitive search strategy select eligible trials in all relevant databases, in accordance to Cochrane guidelines.³⁵ Searches were also done by searching of potential trials from hand searching. Although, it is reasonable that some studies might have been published in local databases and, as a consequence, they were not selected in this study.⁶⁸

Due this review included a small number of randomized trials the limited number of included studies and the high heterogeneity of conditions precluded pooling of data for the majority of the outcomes in this review. Futures high quality studies should clarify the impact of WBV (alone or combined) on deficiencies and limitations in children and adolescents stratified by disability type. Furthermore, studies should also evaluate the long-term effects of WBV, as well as investigate the impact of specific protocol settings, such as frequencies, amplitude and cumulative doses of WBV.

The present findings contribute to important information for clinicians, their patients and researchers. Enthusiasm with new expensive interventions suggesting improvements in the structure and body system in children and adolescent with disabilities need to be followed by scientific evidence.

Conclusion

This is the first systematic review with meta-analysis investigating effects of WBV in children and adolescents with disabilities. Current evidence is often very-low quality and recommendation of WBV in clinical practice is weak, mainly compared with minimal intervention. WBV combined with other intervention may improve some outcomes related to the ICF's body structure and function domain, and other outcomes related to the ICF's activity and participation domains short- and medium-terms. Further high-quality studies are warranted as they are likely to change current estimates.

Conflict of interest

The authors declare no conflicts of interest.

Acknowledgments

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

1. Longo E, Galvão ÉRVP, Ferreira HNC, Lindquist ARR, Shikako-Thomas K. Knowledge translation in pediatric rehabilitation: expanding access to scientific knowledge. *Braz J Phys Ther.* 2017;21(6):389–390.
2. Anaby D, Hand C, Bradley L, et al. The effect of the environment on participation of children and youth with disabilities: a scoping review. *Disabil Rehabil.* 2013;35(19):1589–1598.
3. Bjornson KF, Zhou C, Stevenson RD, Christakis D. Relation of stride activity and participation in mobility-based life habits among children with cerebral palsy. *Arch Phys Med Rehabil.* 2014;95(2):360–368.
4. Law M, Finkelman S, Hurley P, et al. Participation of children with physical disabilities: relationships with diagnosis, physical function, and demographic variables. *Scand J Occup Ther.* 2004;11(4):156–162.
5. Beckung E, Hagberg G. Neuroimpairments, activity limitations, and participation restrictions in children with cerebral palsy. *Dev Med Child Neurol.* 2002;44(5):309–316.
6. *International Classification of Functioning, Disability, and Health: Children & Youth Version: ICF-CY.* World Health Organization; 2007.
7. Cambiaso-Daniel J, Malagaris I, Rivas E, et al. Body composition changes in severely burned children during ICU hospitalization. *Pediatr Crit Care Med.* 2017;18(12):e598–e605.
8. MacDonald M, Leichtman J, Esposito P, Cook N, Ulrich DA. The participation patterns of youth with Down syndrome. *Front Public Health.* 2016;4:253.
9. Curtis DJ, Butler P, Saavedra S, et al. The central role of trunk control in the gross motor function of children with cerebral palsy: a retrospective cross-sectional study. *Dev Med Child Neurol.* 2015;57(4):351–357.
10. Dang VM, Colver A, Dickinson HO, et al. Predictors of participation of adolescents with cerebral palsy: a European multi-centre longitudinal study. *Res Dev Disabil.* 2015;36:551–564.
11. Bjornson KF, Belza B, Kartin D, Logsdon R, McLaughlin JF. Ambulatory physical activity performance in youth with cerebral palsy and youth who are developing typically. *Phys Ther.* 2007;87(3):248.
12. Maher CA, Williams MT, Olds T, Lane AE. Physical and sedentary activity in adolescents with cerebral palsy. *Dev Med Child Neurol.* 2007;49(6):450–457.
13. Molin I, Alricsson M. Physical activity and health among adolescents with cerebral palsy in Sweden. *Int J Adolescent Health.* 2009;21(4):623–633.
14. Zwier JN, van Schie PE, Becher JG, Smits D-W, Gorter JW, Dallmeijer AJ. Physical activity in young children with cerebral palsy. *Disabil Rehabil.* 2010;32(18):1501–1508.
15. Menear K. Parents' perceptions of health and physical activity needs of children with Down syndrome. *Downs Syndr Res Pract.* 2007;12(1):60–68.
16. Simon-Martinez C, Mailleux L, Ortibus E, et al. Combining constraint-induced movement therapy and action-observation training in children with unilateral cerebral palsy: a randomized controlled trial. *BMC Pediatr.* 2018;18(1):250.
17. Kilebrant S, Braathen G, Emilsson R, et al. Whole-body vibration therapy in children with severe motor disabilities. *J Rehabil Med.* 2015;47(3):223–228.
18. Villarroya MA, González-Agüero A, Moros T, Gómez-Trullén E, Casajús JA. Effects of whole body vibration training on balance in adolescents with and without Down syndrome. *Res Dev Disabil.* 2013;34(10):3057–3065.
19. van Nes IJ, Latour H, Schils F, Meijer R, van Kuijk A, Geurts AC. Long-term effects of 6-week whole-body vibration on balance recovery and activities of daily living in the post-acute phase of stroke: a randomized, controlled trial. *Stroke.* 2006;37(September (9)):2331–2335.
20. Eid MA. Effect of whole-body vibration training on standing balance and muscle strength in children with Down syndrome. *Am J Phys Med Rehabil.* 2015;4(8):633–643.
21. el-Shamy SM, Mohamed MSE. Effect of whole body vibration training on bone mineral density in cerebral palsy children. *Indian J Physiother Occup Ther.* 2012;6(January–March (1)):139–141.
22. Lee BK, Chon SC. Effect of whole body vibration training on mobility in children with cerebral palsy: a randomized controlled experimenter-blinded study. *Clin Rehabil.* 2013;27(July (7)):599–607.
23. Ibrahim MM, Eid MA, Moawd SA. Effect of whole-body vibration on muscle strength, spasticity, and motor performance in spastic diplegic cerebral palsy children. *Egypt J Med Hum Genet.* 2014;15(April (2)):173–179.
24. Yabumoto T, Shin S, Watanabe T, et al. Whole-body vibration training improves the walking ability of a moderately impaired child with cerebral palsy: a case study. *J Phys Ther Sci.* 2015;27(9):3023–3025.
25. Semler O, Fricke O, Vezyroglou K, Stark C, Schoenau E. Preliminary results on the mobility after whole body vibration in immobilized children and adolescents. *J Musculoskelet Neuronal Interact.* 2007;7(1):77.
26. Reyes ML, Hernandez M, Holmgren LJ, Sanhueza E, Escobar RG. High-frequency, low-intensity vibrations increase bone mass and muscle strength in upper limbs, improving autonomy in disabled children. *J Bone Miner Res.* 2011;26(8):1759–1766.
27. Moreau NG, Bodkin AW, Bjornson K, Hobbs A, Soileau M, Lahasky K. Effectiveness of rehabilitation interventions to improve gait speed in children with cerebral palsy: systematic review and meta-analysis. *Phys Ther.* 2016;96(12):1938–1954.
28. Duquette SA, Guiliano AM, Starmer DJ. Whole body vibration and cerebral palsy: a systematic review. *J Can Chiropr Assoc.* 2015;59(3):245.
29. Saquette M, Carvalho V, Silva C, Conceição C, Gomes-Neto M. The effects of whole body vibration on mobility and balance in children with cerebral palsy: a systematic review with meta-analysis. *J Musculoskelet Neuronal Interact.* 2015;15(2):137–144.
30. Sá-Caputo DC, Costa-Cavalcanti R, Carvalho-Lima RP, et al. Systematic review of whole body vibration exercises in the treatment of cerebral palsy: brief report. *Dev Neurorehabil.* 2016;19(5):327–333.
31. Matute-Llorente Á, González-Agüero A, Gómez-Cabello A, Vicente-Rodríguez G, Mallén JAC. Effect of whole-body vibration therapy on health-related physical fitness in children and adolescents with disabilities: a systematic review. *J Adolesc Health.* 2014;54(4):385–396.
32. Matute-Llorente A, Gonzalez-Aguero A, Gomez-Cabello A, Vicente-Rodríguez G, Casajús Mallén JA. Effect of whole-body vibration therapy on health-related physical fitness in children and adolescents with disabilities: a systematic review. *J Adolesc Health.* 2014;54(4):385–396.

33. Matute-Llorente A, González-Agüero A, Gómez-Cabello A, Vicente-Rodríguez G, Casajús Mallén JA. Effect of whole-body vibration therapy on health-related physical fitness in children and adolescents with disabilities: a systematic review. *J Adolesc Health*. 2014;54(4):385–396.
34. Almeida KM, Fonseca ST, Figueiredo PR, Aquino AA, Mancini MC. Effects of interventions with therapeutic suits (clothing) on impairments and functional limitations of children with cerebral palsy: a systematic review. *Braz J Phys Ther*. 2017.
35. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev*. 2015;4(1):1.
36. Oliveira CB, Elkins MR, Lemes ÍR, et al. A low proportion of systematic reviews in physical therapy are registered: a survey of 150 published systematic reviews. *Braz J Phys Ther*. 2017.
37. Maher CG, Sherrington C, Herbert RD, Moseley AM, Elkins M. Reliability of the PEDro scale for rating quality of randomized controlled trials. *Phys Ther*. 2003;83(8):713.
38. Higgins JP, Green S. *Cochrane Handbook for Systematic Reviews of Interventions*. vol. 4. John Wiley & Sons; 2011.
39. Bosco FA, Aguinis H, Singh K, Field JG, Pierce CA. Correlational effect size benchmarks. *J Appl Psychol*. 2015;100(2):431.
40. Jones L, Bellis MA, Wood S, et al. Prevalence and risk of violence against children with disabilities: a systematic review and meta-analysis of observational studies. *Lancet*. 2012;380(9845):899–907.
41. Harvey LA, Katalinic OM, Herbert RD, Moseley AM, Lannin NA, Schurr K. Stretch for the treatment and prevention of contracture: an abridged republication of a Cochrane Systematic Review. *J Physiother*. 2017;63(2):67–75.
42. Stark C, Herkenrath P, Hollmann H, et al. Early vibration assisted physiotherapy in toddlers with cerebral palsy – a randomized controlled pilot trial. *J Musculoskelet Neuronal Interact*. 2016;16(3):183–192.
43. Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: 1. Introduction—GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol*. 2011;64(4):383–394.
44. Oliveira VC, Ferreira PH, Maher CG, Pinto RZ, Refshauge KM, Ferreira ML. Effectiveness of self-management of low back pain: systematic review with meta-analysis. *Arthritis Care Res*. 2012;64(11):1739–1748.
45. Ioannidis JP, Trikalinos TA. The appropriateness of asymmetry tests for publication bias in meta-analyses: a large survey. *CMAJ*. 2007;176(8):1091–1096.
46. El-Shamy SM. Effect of whole-body vibration on muscle strength and balance in diplegic cerebral palsy: a randomized controlled trial. *Am J Phys Med Rehabil*. 2014;93(2):114–121.
47. Ko MS, Sim YJ, Kim DH, Jeon HS. Effects of three weeks of whole-body vibration training on joint-position sense, balance, and gait in children with cerebral palsy: a randomized controlled study. *Physiother Can*. 2016;68(Spring (2)): 99–105.
48. Ruck J, Chabot G, Rauch F. Vibration treatment in cerebral palsy: a randomized controlled pilot study. *J Musculoskelet Neuronal Interact*. 2010;10(March (1)):77–83.
49. Ward K, Alsop C, Caulton J, Rubin C, Adams J, Mughal Z. Low magnitude mechanical loading is osteogenic in children with disabling conditions. *J Bone Miner Res*. 2004;19(3):360–369.
50. Gonzalez-Aguero A, Matute-Llorente A, Gomez-Cabello A, Casajus JA, Vicente-Rodriguez G. Effects of whole body vibration training on body composition in adolescents with Down syndrome. *Res Dev Disabil*. 2013;34(5):1426–1433.
51. Erceg DN, Anderson LJ, Nickles CM, Lane CJ, Weigensberg MJ, Schroeder ET. Changes in bone biomarkers, BMC, and insulin resistance following a 10-week whole body vibration exercise program in overweight Latino boys. *Int J Med Sci*. 2015;12(6):494–501.
52. El-Shamy S. Effect of whole body vibration training on quadriceps strength, bone mineral density, and functional capacity in children with hemophilia: a randomized clinical trial. *J Musculoskelet Neuronal Interact*. 2017;17(2):19–26.
53. Högler W, Scott J, Bishop N, et al. The effect of whole body vibration training on bone and muscle function in children with osteogenesis imperfecta. *J Clin Endocrinol Metab*. 2017.
54. Edionwe J, Hess C, Fernandez-Rio J, et al. Effects of whole-body vibration exercise on bone mineral content and density in thermally injured children. *Burns*. 2016;42(May (3)): 605–613.
55. Matute-Llorente A, Gonzalez-Aguero A, Gomez-Cabello A, Olmedillas H, Vicente-Rodriguez G, Casajus JA. Effect of whole body vibration training on bone mineral density and bone quality in adolescents with Down syndrome: a randomized controlled trial. *Osteopor Int*. 2015;26(October (10)):2449–2459.
56. Cochrane D. Vibration exercise: the potential benefits. *Int J Sports Med*. 2011;32(02):75–99.
57. Bosco C, Cardinale M, Tarpela O. Influence of vibration on mechanical power and electromyogram activity in human arm flexor muscles. *Eur J Appl Physiol*. 1999;79(4):306–311.
58. Rittweger J, Mutschelknauss M, Felsenberg D. Acute changes in neuromuscular excitability after exhaustive whole body vibration exercise as compared to exhaustion by squatting exercise. *Clin Physiol Funct Imaging*. 2003;23(2):81–86.
59. Cochrane DJ, Hawke EJ. Effects of acute upper-body vibration on strength and power variables in climbers. *J Strength Cond Res*. 2007;21(2):527.
60. Issurin V, Tenenbaum G. Acute and residual effects of vibratory stimulation on explosive strength in elite and amateur athletes. *J Sports Sci*. 1999;17(3):177–182.
61. Bosco C, Colli R, Introini E, et al. Adaptive responses of human skeletal muscle to vibration exposure. *Clin Physiol*. 1999;19(2):183.
62. Frost HM. Bone's mechanostat: a 2003 update. *Anat Rec*. 2003;275(2):1081–1101.
63. Gusso S, Munns CF, Colle P, et al. Effects of whole-body vibration training on physical function, bone and muscle mass in adolescents and young adults with cerebral palsy. *Sci Rep*. 2016;6:22518.
64. Duquette SA, Guiliano AM, Starmer DJ. Whole body vibration and cerebral palsy: a systematic review. *J Can Chiropr Assoc*. 2015;59(3):245–252.
65. Saquetto M, Carvalho V, Silva C, Conceicao C, Gomes-Neto M. The effects of whole body vibration on mobility and balance in children with cerebral palsy: a systematic review with meta-analysis. *J Musculoskelet Neuronal Interact*. 2015;15(2):137–144.
66. Sá-Caputo DC, da F, Dionello C, Frederico ÉHF, et al. Whole-body vibration exercise improves functional parameters in patients with osteogenesis imperfecta: a systematic review with a suitable approach. *Afr J Tradit Complement Altern Med*. 2017;14(3):199.
67. Moreira-Marconi E, Sá-Caputo DC, Dionello CF, et al. Whole-body vibration exercise is well tolerated in patients with Duchenne muscular dystrophy: a systematic review. *Afr J Tradit Complement Altern Med*. 2017;14(4 suppl):2.
68. Parreira PdCS, Costa LdCM, Junior LCH, Lopes AD, Costa LOP. Current evidence does not support the use of Kinesio Taping in clinical practice: a systematic review. *J Physiother*. 2014;60(1):31–39.