Cochrane
Library

Cochrane Database of Systematic Reviews

Oral vaccines for preventing cholera (Review)

Sinclair D, Abba K, Zaman K, Qadri F, Graves PM

Sinclair D, Abba K, Zaman K, Qadri F, Graves PM.

Oral vaccines for preventing cholera.

Cochrane Database of Systematic Reviews 2011, Issue 3. Art. No.: CD008603.

DOI: 10.1002/14651858.CD008603.pub2.

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS FOR THE MAIN COMPARISON	3
BACKGROUND	6
OBJECTIVES	7
METHODS	7
Figure 1.	10
RESULTS	11
Figure 2.	14
DISCUSSION	17
AUTHORS' CONCLUSIONS	18
ACKNOWLEDGEMENTS	18
REFERENCES	19
CHARACTERISTICS OF STUDIES	30
DATA AND ANALYSES	90
Analysis 1.1. Comparison 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes, Outcome 1 Cases of cholera - 1st year of follow up (with meta analysis).	99
Analysis 1.2. Comparison 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes, Outcome 2 Cases of cholera - 2nd year of follow up (with meta analysis).	101
Analysis 1.3. Comparison 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes, Outcome 3 Cases of cholera - 3rd year of follow up (with meta analysis).	102
Analysis 1.4. Comparison 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes, Outcome 4 Cases of cholera - 4th year of follow up (with meta analysis).	103
Analysis 1.5. Comparison 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes, Outcome 5 Cases of cholera by age group - First two years of follow-up.	105
Analysis 1.6. Comparison 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes, Outcome 6 Cases of cholera by age group - First two years of follow-up (sensitivity analysis).	107
Analysis 2.1. Comparison 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis, Outcome 1 Cases of cholera by time of follow-up (3-dose recipients).	108
Analysis 2.2. Comparison 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis, Outcome 2 Cases of cholera by age-group - 1st year of follow-up (3-dose recipients).	109
Analysis 2.3. Comparison 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis, Outcome 3 Cases of cholera by age-group - 2nd year of follow-up (3-dose recipients).	110
Analysis 2.4. Comparison 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis, Outcome 4 Cases of cholera by blood group - First 2 years of follow-up (3-dose recipients).	111
Analysis 2.5. Comparison 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis, Outcome 5 Cases of all cause diarrhoea - 1st year of follow-up (3-dose recipients).	112
Analysis 2.6. Comparison 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis, Outcome 6 Deaths - 1st year of follow-up (3-dose recipients).	112
Analysis 3.1. Comparison 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis, Outcome 1 Cases of cholera by time of follow-up (3-dose recipients).	113
Analysis 3.2. Comparison 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis, Outcome 2 Cases of cholera by age-group - 1st year of follow-up (3-dose recipients).	114
Analysis 3.3. Comparison 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis, Outcome 3 Cases of cholera by age-group - 2nd year of follow-up (3-dose recipients).	115
Analysis 3.4. Comparison 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis, Outcome 4 Cases of cholera by blood group - First 2 years of follow-up (3-dose recipients).	116
Analysis 3.5. Comparison 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis, Outcome 5 Cases of all cause diarrhoea - 1st year of follow-up (3-dose recipients).	117

Analysis 3.6. Comparison 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis, Outcome 6 Deaths - 1st year of follow-up (3-dose recipients).	117
Analysis 4.1. Comparison 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis, Outcome 1 Cases of confirmed cholera by time of follow-up (3-dose recipients).	118
Analysis 4.2. Comparison 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis, Outcome 2 Cases of cholera by age-group - 1st year of follow-up (3-dose recipients).	119
Analysis 4.3. Comparison 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis, Outcome 3 Cases of cholera by age-group - 2nd year of follow-up (3-dose recipients).	120
Analysis 4.4. Comparison 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis, Outcome 4 Cases of cholera by blood group, First 2 years of follow-up (3-dose recipients).	121
Analysis 4.5. Comparison 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis, Outcome 5 Cases of all cause diarrhoea - 1st year of follow-up (3-dose recipients).	122
Analysis 4.6. Comparison 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis, Outcome 6 Deaths - 1st year of follow-up (3-dose recipients).	122
Analysis 5.1. Comparison 5 Whole cell plus recombinant B subunit vaccine (WC-rBS) versus placebo - Subgroup analysis, Outcome 1 Cases of cholera by age group - 1st year of follow-up (2 doses).	123
Analysis 5.2. Comparison 5 Whole cell plus recombinant B subunit vaccine (WC-rBS) versus placebo - Subgroup analysis, Outcome 2 Cases of cholera by age group - 2nd year of follow-up (2 doses plus booster).	124
Analysis 5.3. Comparison 5 Whole cell plus recombinant B subunit vaccine (WC-rBS) versus placebo - Subgroup analysis, Outcome 3 Cases of cholera in military recruits, 4 to 18 weeks follow-up.	125
Analysis 6.1. Comparison 6 Whole cell vaccines (all types) versus placebo - Safety outcomes (first dose), Outcome 1 Adverse events - Whole cell (WC) versus placebo.	126
Analysis 6.2. Comparison 6 Whole cell vaccines (all types) versus placebo - Safety outcomes (first dose), Outcome 2 Adverse events - Whole cell plus B subunit (WC-BS) versus placebo.	128
Analysis 6.3. Comparison 6 Whole cell vaccines (all types) versus placebo - Safety outcomes (first dose), Outcome 3 Adverse events - Whole cell plus recombinant B subunit (WC-rBS) versus placebo.	130
Analysis 6.4. Comparison 6 Whole cell vaccines (all types) versus placebo - Safety outcomes (first dose), Outcome 4 Adverse events - Bivalent whole cell (BivWC) versus placebo.	133
Analysis 7.1. Comparison 7 Whole cell vaccines (all types) versus placebo - Safety outcomes (second dose), Outcome 1 Adverse events - Whole cell plus recombinant B subunit (WC-rBS) versus placebo.	137
Analysis 7.2. Comparison 7 Whole cell vaccines (all types) versus placebo - Safety outcomes (second dose), Outcome 2 Adverse events - Bivalent whole cell (BivWC) versus placebo.	140
Analysis 8.1. Comparison 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes, Outcome 1 Cases of cholera following natural infection - CVD 103HgR versus placebo.	143
Analysis 8.2. Comparison 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes, Outcome 2 Severe cholera following natural infection - CVD 103HgR versus placebo.	144
Analysis 8.3. Comparison 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes, Outcome 3 Death from any cause (except motor accidents).	144
Analysis 8.4. Comparison 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes, Outcome 4 Death from diarrhoea (any organism).	145
Analysis 8.5. Comparison 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes, Outcome 5 Cases of moderate to severe diarrhoea - following artificial challenge.	146
Analysis 8.6. Comparison 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes, Outcome 6 Cases of any diarrhoea -following artificial challenge.	147
Analysis 9.1. Comparison 9 Live attenuated vaccines (all types) versus placebo - Safety outcomes, Outcome 1 Adverse events - CVD 103-HgR versus placebo.	148
Analysis 9.2. Comparison 9 Live attenuated vaccines (all types) versus placebo - Safety outcomes, Outcome 2 Adverse events - Peru 15 versus placebo.	152
Analysis 9.3. Comparison 9 Live attenuated vaccines (all types) versus placebo - Safety outcomes, Outcome 3 Adverse events - VC638 versus placebo.	154
Analysis 10.1. Comparison 10 Live attenuated CVD 103-HgR vaccine versus placebo - Subgroup analysis, Outcome 1 Cases of cholera by age group (age 2-5 years).	157

Analysis 10.2. Comparison 10 Live attenuated CVD 103-HgR vaccine versus placebo - Subgroup analysis, Outcome 2	
Cases of confirmed cholera attending healthcare facilities (age over 5 years).	157
Analysis 10.3. Comparison 10 Live attenuated CVD 103-HgR vaccine versus placebo - Subgroup analysis, Outcome 3	
Cases of cholera within four years and five months, by blood group.	158
Analysis 10.4. Comparison 10 Live attenuated CVD 103-HgR vaccine versus placebo - Subgroup analysis, Outcome 4 Any	
diarrhoea following artificial challenge, by blood group.	159
Analysis 10.5. Comparison 10 Live attenuated CVD 103-HgR vaccine versus placebo - Subgroup analysis, Outcome 5	
Moderate or severe diarrhoea due to V. cholerae after artificial challenge, by blood group.	160
ADDITIONAL TABLES	161
WHAT'S NEW	164
CONTRIBUTIONS OF AUTHORS	164
DECLARATIONS OF INTEREST	164
SOURCES OF SUPPORT	165
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	165
INDEX TERMS	165

Oral vaccines for preventing cholera

David Sinclair¹, Katharine Abba¹, K Zaman², Firdausi Qadri³, Patricia M Graves⁴

¹International Health Group, Liverpool School of Tropical Medicine, Liverpool, UK. ²Child Health Unit, International Centre for Diarrhoeal Disease Research, Bangladesh, Dhaka, Bangladesh. ³International Centre for Diarrhoeal Disease Research, Bangladesh (ICDDR,B), Dhaka, Bangladesh. ⁴EpiVec Consulting, Atlanta, Georgia, USA

Contact address: David Sinclair, International Health Group, Liverpool School of Tropical Medicine, Pembroke Place, Liverpool, L3 5QA, UK. davesinkers@yahoo.com.

Editorial group: Cochrane Infectious Diseases Group.

Publication status and date: Unchanged, published in Issue 5, 2019.

Citation: Sinclair D, Abba K, Zaman K, Qadri F, Graves PM. Oral vaccines for preventing cholera. *Cochrane Database of Systematic Reviews* 2011, Issue 3. Art. No.: CD008603. DOI: 10.1002/14651858.CD008603.pub2.

Copyright © 2019 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Cholera is a cause of acute watery diarrhoea which can cause dehydration and death if not adequately treated. It usually occurs in epidemics, and is associated with poverty and poor sanitation. Effective, cheap, and easy to administer vaccines could help prevent epidemics.

Objectives

To assess the effectiveness and safety of oral cholera vaccines in preventing cases of cholera and deaths from cholera.

Search methods

In October 2010, we searched the Cochrane Infectious Disease Group Specialized Register; Cochrane Central Register of Controlled Trials (CENTRAL); MEDLINE; EMBASE; LILACS; the metaRegister of Controlled Trials (mRCT), and the WHO International Clinical Trials Registry Platform (ICTRP) for relevant published and ongoing trials.

Selection criteria

Randomized or quasi-randomized controlled trials of oral cholera vaccines in healthy adults and children.

Data collection and analysis

Each trial was assessed for eligibility and risk of bias by two authors working independently. Data was extracted by two independent reviewers and analysed using the Review Manager 5 software. Outcomes are reported as vaccine protective efficacy (VE) with 95% confidence intervals (CIs).

Main results

Seven large efficacy trials, four small artificial challenge studies, and 29 safety trials contributed data to this review.

Five variations of a killed whole cell vaccine have been evaluated in large scale efficacy trials (four trials, 249,935 participants). The overall vaccine efficacy during the first year was 52% (95% CI 35% to 65%), and during the second year was 62% (95% CI 51% to 62%). Protective efficacy was lower in children aged less than 5 years; 38% (95% CI 20% to 53%) compared to older children and adults; 66% (95% CI 57% to 73%).

One trial of a killed whole cell vaccine amongst military recruits demonstrated 86% protective efficacy (95% CI 37% to 97%) in a small epidemic occurring within 4 weeks of the 2-dose schedule (one trial, 1426 participants). Efficacy data is not available beyond two years for the currently available vaccine formulations, but based on data from older trials is unlikely to last beyond three years.

The safety data available on killed whole cell vaccines have not demonstrated any clinically significant increase in adverse events compared to placebo.

Only one live attenuated vaccine has reached Phase III clinical evaluation and was not effective (one trial, 67,508 participants). Two new candidate live attenuated vaccines have demonstrated clinical effectiveness in small artificial challenge studies, but are still in development.

Authors' conclusions

The currently available oral killed whole cell vaccines can prevent 50 to 60% of cholera episodes during the first two years after the primary vaccination schedule. The impact and cost-effectiveness of adopting oral cholera vaccines into the routine vaccination schedule of endemic countries will depend on the prevalence of cholera, the frequency of epidemics, and access to basic services providing rapid rehydration therapy.

PLAIN LANGUAGE SUMMARY

Oral vaccines for preventing cholera

Researchers in The Cochrane Collaboration conducted a review of the effect of oral vaccines for preventing cholera. After searching for relevant studies, they identified 48 relevant articles. Their findings are summarized below.

What is cholera and how do vaccines work?

Cholera is a severe form of diarrhoea. People get cholera by drinking water or eating food that has been contaminated with the bacteria (*Vibrio cholera*). Some people only become mildly ill, but some become extremely unwell with watery diarrhoea and vomiting. These people can become dehydrated very quickly and if untreated 25% to 50% can die.

The disease spreads rapidly in poor communities, especially where there is no sanitation or a lack of clean water. In refugee camps or following natural disasters a cholera epidemic can kill many hundreds of people very quickly.

Oral cholera vaccines work by giving people a small dose of the cholera bacteria to swallow. This dose of bacteria has been killed or changed so that it does not cause diarrhoea but is still able to make the person immune to natural cholera. There are three oral cholera vaccines currently available.

What the research says about the effects of using current oral vaccines

Oral cholera vaccines will decrease your risk of getting cholera if you live somewhere where cholera is common, but they won't remove the risk completely.

Oral cholera vaccines probably don't have any major side effects when they are taken, but rare or late complications cannot be excluded.

SUMMARY OF FINDINGS FOR THE MAIN COMPARISON *[Explanation]*

Oral killed whole cell vaccines for preventing cholera						
Patient or population: Adults and children Settings: Endemic areas Intervention: Killed whole cell vaccines administered orally Comparison: Placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Vaccine efficacy (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Not being vaccinated	Being vaccinated				
How many people get cholera during the first 2 years after vaccination?	Children aged less than 5 years		VE 38% (20% to 53%)	29005 (4 studies ⁵)	high ^{1,2,3,4}	Oral cholera vaccine prevents just over one third of cholera illnesses
	90 per 10,000	56 per 10,000 (42 to 72)				
	Older children and adults		VE 66% (57% to 73%)	214066 (4 studies ⁵)	high ^{1,2,3,4}	Oral cholera vaccine prevents two thirds of cholera illnesses
	30 per 100,000	10 per 100,000 (8 to 13)				
How long does the protection last?	3rd year after vaccination; all ages		VE 30% (2% to 50%)	58184 (1 study ⁷)	moderate ⁶	Oral cholera vaccine is probably less effective in the third year
	30 per 10,000	21 per 10,000 (15 to 29)				
	4th year after vaccination; all ages		VE -5% (-84% to 40%)	56613 (1 study ⁷)	moderate ⁶	Oral cholera vaccine is probably ineffective after 4 years

	30 per 100,000	32 per 10,000 (18 to 55)		
Are there any side effects?	All ages		44,924 (14 studies)	moderate⁸ Oral cholera vaccines probably don't have more side effects than a placebo

*The basis for the **assumed risk** (eg the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% CI) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; VE: Vaccine protective efficacy.

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ No study limitations: Clemens 1988 Bangladesh and Taylor 2000 Peru are individually randomized trials with adequate allocation concealment and blinding. Sur 2009 India is a cluster randomized study, and Trach 1997 is a quasi-randomized study without allocation concealment.

² No serious inconsistency: The findings from all three trials (4 comparisons) were remarkably similar and any observed differences between the vaccines is well within the bounds of random error. It should be noted that a protective effect with the most widely available vaccine (WC-rBS/Dukoral®), given in its recommended schedule of two doses, was not shown until after a booster dose at 10 months.

³ No serious indirectness: The trials are from several endemic countries and include both adults and children. This evidence could reasonably be applied to other endemic settings where the background risk of cholera is known and used to calculate an absolute benefit with vaccination.

⁴ No serious imprecision: The finding is of a statistically significant benefit with vaccination. The clinical importance will depend on the incidence of cholera in the population.

⁵ Clemens 1988 Bangladesh: a 3-arm trial of WC (currently unavailable), WC-BS (currently unavailable) vs placebo, Taylor 2000 Peru; WC-rTB (Dukoral®) versus placebo, Trach 1997 Vietnam; vWC (a variant WC vaccine only available in Vietnam) vs placebo, Sur 2009 India; BivWC (Shanchol®) vs placebo.

⁶ Serious indirectness: The exact vaccines used in this trial are no longer available but the current vaccines are very similar in composition. Downgraded by 1.

⁷ Only Clemens 1988 Bangladesh followed participants beyond 2-years.

⁸ Fourteen studies assessed for side effects during the first 2 weeks after vaccination. No individual side effect has been shown to be more common with the oral cholera vaccine than with placebo. This data cannot exclude rare or late complications. Downgraded by 1.

BACKGROUND

Description of the condition

Cholera is an acute intestinal infection, caused by the bacterium *Vibrio cholerae*. Most infected persons do not become ill, although the bacteria are present in the faeces for 7 to 14 days. Over 90% of those who do become ill experience a mild diarrhoeal episode that is indistinguishable from other diarrhoeal illnesses. However, a proportion develop typical cholera symptoms, with sudden onset of profuse watery diarrhoea, usually accompanied by vomiting, which can lead to severe dehydration (WHO 2000a). If untreated, around 25% to 50% of patients with the typical cholera symptoms will die, but if given adequate rehydration treatment the deaths can be reduced to less than 1% (WHO 2000b). In 2005 there were a total of 131,943 reported cases of cholera throughout the world, including 2272 deaths (WHO 2006a). Ninety-five percent of the reported cases were in Africa, but it is likely that many more cases, both in Africa and elsewhere, went unreported.

V. cholerae is transmitted mainly through the ingestion of faecally contaminated water or food, and can spread rapidly especially where there is poverty, poor hygiene and lack of sanitation. It can lead to serious outbreaks; in 2005 the World Health Organization (WHO) confirmed 49 different outbreaks in 36 countries (WHO 2006a), and in vulnerable populations epidemics can be devastating; in July 1994, in the refugee camps of Goma in Zaire, there were 70,000 cases with 12,000 deaths (Sanchez 1997). More recently, large epidemics have occurred in Zimbabwe (WHO 2009), and Haiti (WHO 2010a).

V. cholerae colonise the gut by attaching themselves to receptors in the mucosa of the upper small intestine (Sack 2004). Pathogenicity is mediated by a toxin, composed of two subunits; A and B. The B subunit is involved in binding the bacteria to the epithelial cell surface. It has no toxic effect, but does stimulate the host's immune response. The soluble A subunit is then released into the mucosal cells and causes hypersecretion of fluids and electrolytes, which lead to the typical symptoms of the disease (Girard 2005). Colonisation of the intestine can be inhibited by host antibodies generated in response to previous infection with *V. cholerae*.

There are over two hundred distinct serological groups of *V. cholerae*, classified on the basis of the 'O' antigen present on the cell surface, of which only two are known to cause epidemics: serogroups O1 and O139. *V. cholerae* O1 can be further classified into two biotypes: classical and El Tor. These in turn can each be divided into three serotypes: Ogawa, Inaba and Hikojima (Heymann 2008). The epidemic strains currently in circulation worldwide are the El Tor biotype of *V. cholerae* O1, which was first recognised in Indonesia in 1961 and has now spread to many other countries in Asia, Europe, Africa, and Latin America; and the Bengal strain of *V. cholerae* O139 which began in 1992 in India and Bangladesh, and remains restricted to Asia (WHO 2000b). The classical biotype of *V. cholerae* O1 is also known to cause

epidemics, though these are now uncommon, and non-O1/non-O139 strains occasionally cause sporadic cases of gastroenteritis (Heymann 2008).

There is evidence that persons with blood group O have overall lower risk of cholera, but increased susceptibility to severe cholera (Harris 2005). The mechanism for this effect is not known, but it should be taken into account when assessing vaccine effectiveness.

Description of the intervention

Widespread use of cholera vaccines began in the 1960s. The vaccines then in use were composed of whole *V. cholerae* O1 cells, killed using formalin, phenol or heat, and administered by injection. In the 1970s, these injected whole cell vaccines fell out of favour (Bhadra 1994), as it was perceived that they had a low efficacy (around 50%), provided only short-term immunity (3 to 6 months), and had an unacceptable rate of side effects. A Cochrane review first published in 1998, however, found that the duration and efficacy of the whole cell injected vaccines may have been underestimated: it was 54% at seven months (based on 18 trials) and 46% at one year (based on 14 trials). Protection waned by the second year in children under five, but persisted into the third year for those over the age of five years (Graves 2010). Nevertheless, injected vaccines are no longer in use or available, and attention is now focused on vaccines administered by the oral route.

Two main types of oral vaccines have been investigated in clinical trials: inactivated vaccines (containing killed whole cells of *V. cholerae*), and live attenuated vaccines (containing genetically modified, non-pathogenic strains of *V. cholerae*). In addition, subunit vaccines have been tested which consist only of cell components (antigens). The live attenuated vaccines are usually given as a single dose, whereas killed whole cell vaccines may require two or three doses at one week intervals to produce an adequate immunological response. Three vaccine formulations are currently available (WHO 2010b):

- **WC-rBS (Dukoral®):** A monovalent inactivated vaccine containing killed whole cells of *V. cholerae* O1 plus additional recombinant cholera toxin B subunit. Produced by SBL Vaccine/Crucell, Sweden.
- **BivWC (Shancho®):** A bivalent inactivated vaccine containing killed whole cells of *V. cholerae* O1 and *V. cholerae* O139. Produced by Shantha Biotechnics, India.
- **BivWC (mORCVAX®):** A bivalent inactivated vaccine containing killed whole cells of *V. cholerae* O1 and *V. cholerae* O139. Produced by VABIOTECH, Vietnam and only available in Vietnam.

However, there are many other candidate vaccines at various stages of clinical development (Girard 2005).

How the intervention might work

Vaccines work by stimulating immunity against a pathogen which has been killed, attenuated or otherwise rendered incapable of causing disease, in order to prevent or mitigate the effects of infection with the natural pathogen if it subsequently occurs. The route of administration of a vaccine may influence its immunogenicity and acceptability. Oral vaccines have the potential to stimulate local immunity within the mucosa of the gut, preventing the colonisation and multiplication of *V. cholerae*. Since cholera is transmitted orally, oral vaccines may thus have more direct effect than injected vaccines which stimulate immunity in the blood. Oral vaccines are also potentially easier to administer, more acceptable to patients than injected vaccines, and have a reduced risk of transmitting blood borne infections (Holmgren 2005).

The cholera toxin B subunit contains similar antigens to those found in enterotoxigenic *Escherichia coli* (ETEC); an important cause of diarrhoea in many parts of the world (Huilan 1991), and the most common cause of diarrhoea in people travelling from industrialised to developing countries (Sack 2004). Oral cholera vaccines may therefore provide significant cross-protection against ETEC infection and the vaccine is already licensed in many countries for preventing ETEC diarrhoea in travellers. This aspect of cholera vaccine use will be covered by another Cochrane review on vaccines to prevent ETEC.

Why it is important to do this review

Oral vaccines have been licensed in many countries and are currently used mainly by travellers (Hill 2006). However, there has not been a full review of the relative effectiveness of different types of oral vaccine, the duration of their efficacy, or their adverse effects.

These vaccines may also have an important role in preventing cholera in areas where it is endemic, or in the prevention or control of outbreaks in high risk settings. The killed whole cell vaccine (WC/rBS) has been used in crisis situations in Darfur, Sudan (WHO 2006b), and in Aceh, Indonesia in 2005 after the tsunami (WHO 2006c). It has also been evaluated in an endemic situation in Beira, Mozambique in 2003-2004 (Lucas 2005). The live CVD 103-HgR vaccine was used during a cholera outbreak in Pohnpei, Federated States of Micronesia in 2000 (Calain 2004).

This review is one of a series of three that replaces a previous Cochrane review 'Vaccines for preventing cholera', which was first published in 1998 and updated in 2001. An updated stable review of injected vaccines (Graves 2010) has now replaced the original cholera vaccines review; it will be accompanied by this review of oral vaccines and a further review assessing the effects of vaccines (including cholera vaccine) on infection with ETEC.

OBJECTIVES

To assess the effectiveness and safety of oral cholera vaccines in preventing cases of cholera and deaths from cholera.

METHODS

Criteria for considering studies for this review

Types of studies

Randomized or quasi-randomized controlled trials, including cluster-randomized trials.

Types of participants

Well adults or children (without symptoms of cholera).

Types of interventions

Intervention

Any vaccine that is designed to prevent cholera and is administered by the oral route.

Control

Placebo, control vaccine, no intervention or different dose or schedule of cholera vaccine.

Types of outcome measures

Primary outcomes

- Cases of cholera.
- Deaths from cholera.

Secondary outcomes

- Cases of severe dehydrating diarrhoea.
- Cases of all-cause diarrhoea.
- Deaths from severe dehydrating diarrhoea.
- Deaths from all causes.
- Serious adverse events leading to hospital admission or death.
- Other adverse events.

Search methods for identification of studies

We attempted to identify all relevant studies regardless of language or publication status (published, unpublished, in press, and in progress). The search was conducted in January 2010 and repeated in October 2010.

Electronic searches

Published studies

We searched the following databases using the search terms detailed in [Table 1](#): The Cochrane Infectious Disease Group Specialized Register; the Cochrane Central Register of Controlled Trials (CENTRAL), published in The Cochrane Library; MEDLINE; EMBASE; and LILACS.

Ongoing studies

We also searched the metaRegister of Controlled Trials (mRCT) and the WHO International Clinical Trials Registry Platform (ICTRP) for ongoing trials using “cholera” and “vaccin*” as search terms.

Searching other resources

Researchers, organizations, and pharmaceutical companies

We attempted to contact individual researchers working in the field for unpublished and ongoing trials.

Reference lists

We also checked the reference lists of all studies identified by the above methods for any additional studies relevant to this review.

Data collection and analysis

Selection of studies

Two authors (PG, KA or DS) independently screened all citations and abstracts identified by the search strategy for potentially eligible studies. Full reports of those studies deemed eligible were formally assessed for inclusion in the review using a pre-designed eligibility form based on the inclusion criteria. All reports were scrutinised for evidence of dual publication.

Trials where participants were given an artificial challenge with *V. cholerae* after vaccination (i.e. by ingesting a standardized dose of bacteria), were included but assessed separately from studies assessing efficacy against natural infection. Trials reporting only safety or adverse event data were included and summarized only if primary outcome data (an efficacy trial) for the same vaccine was already available. Trials testing the vaccine for purposes other than safety or prevention of cholera (for example, for prevention

of diarrhoea associated with ETEC, or ‘traveller’s diarrhoea’) were excluded.

Where it was unclear whether a trial should be included we attempted to contact the authors for clarification, and resolved any differences in opinion through discussion. We obtained translated copies of those papers published in languages other than English. The studies which did not meet the criteria for inclusion, and the reasons for their exclusion, are listed in the ‘[Characteristics of excluded studies](#)’ table.

Data extraction and management

For each included trial, two authors (KA, KZ or DS) independently extracted information (using a pre-tested data extraction form) on the characteristics of the trial (study design, study dates and duration, study location, setting, and source of funding); the participants (the inclusion and exclusion criteria); the intervention (the type of vaccine, type of placebo, dose and immunisation schedule); and the outcomes presented in the papers.

For individually randomized trials, two authors independently extracted the number of participants randomized to each group, and the number experiencing the outcome. Data on the number of doses received and the number of participants lost to follow-up has been calculated and recorded for each group.

For cluster-randomized trials, we recorded the number of clusters in the trial, the average (mean) size of clusters, the unit of randomization (e.g. household or institution), and reported estimates of the intracluster correlation coefficient (ICC) for each outcome. If the trial results were adjusted for clustering we extracted the point estimate and the 95% confidence interval (CI), and also the unadjusted data so that we could calculate an adjusted risk ratio to present in a meta-analysis. Where results were not adjusted for clustering, we extracted the same data as for individually randomized trials, and adjusted the results according to known estimates of the ICC.

Adverse event data has been extracted for each individual type of event wherever possible. Where adverse events were reported for more than one dose, the number of people reporting each side effect after each dose has been recorded. Where trials reported the occurrence of adverse events over time following a single dose, if possible we recorded the proportion of people affected during each time period. If the denominator or total number of people affected for each time period is not clear, then events occurring in the first time period (typically 24 hours) after each dose was recorded.

Where data was missing or incomplete we contacted the authors for clarification. In cases of disagreement we double checked the data extraction and resolved the disagreement through discussion.

Assessment of risk of bias in included studies

Two authors (KZ, KA or DS) independently assessed the risk of bias of the individually randomized trials using the ‘The Cochrane

Collaboration's tool for assessing the risk of bias' (Higgins 2008). We followed this guidance to assess whether adequate steps had been taken to reduce the risk of bias across six domains: sequence generation; allocation concealment; blinding (of participants, personnel, and outcome assessors); incomplete outcome data; selective outcome reporting; and other sources of bias. For cluster-randomized trials we also considered the possible effects of particular biases which occur with this study design: recruitment bias, baseline imbalance, loss of clusters, incorrect analysis and comparability to individually randomized trials (Higgins 2008).

For sequence generation and allocation concealment we report the methods used. For blinding we describe who was blinded and the blinding method. For incomplete outcome data we report the percentage and proportion lost to follow up. For selective outcome reporting we state any discrepancies between the methods used

and the results in terms of the outcomes measured or the outcomes reported. For other biases we describe any other trial features that we think could have affected the trials result (e.g. if the trial was stopped early). We also report components of study design or conduct which may have introduced any bias specific to cluster-randomized trials.

We have categorized our judgments as 'yes' (low risk of bias), 'no' (high risk of bias), or 'unclear', and this information has been used to guide the interpretation of the results. Where our judgement for efficacy trials was unclear we attempted to contact the trial authors for clarification and any differences of opinion were resolved through discussion.

The results of this assessment of the risk of bias can be seen in Figure 1.

Figure 1. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding (performance bias and detection bias): Efficacy outcomes	Blinding (performance bias and detection bias): Safety outcomes	Incomplete outcome data (attrition bias): Efficacy outcomes	Incomplete outcome data (attrition bias): Safety outcomes	Selective reporting (reporting bias)	Other bias
Anh 2007	●	●	?	?	●	●	●	●
Begue 1995	●	●	?	?	●	●	●	●
Benítez 1999	?	●	?	●	?	●	●	●
Chen 1996	?	?	?	?	?	●	●	●
Clemens 1987	●	●	?	?	●	●	●	●
Clemens 1988 Bangladesh	●	●	●	●	●	●	●	●
Cohen 2002	●	●	●	●	●	●	●	●
Concha 1995	?	●	?	●	?	●	●	●
Cryz 1990	?	?	?	●	?	●	●	●
García 2005	?	?	?	?	●	●	●	●
Gotuzzo 1993	?	●	?	●	?	●	●	●
Hallander 2002	?	?	?	?	?	●	●	●
Kanungo 2009	●	●	?	●	?	●	●	●
Kotloff 1992	?	?	?	●	?	●	●	●
Lagos 1993	?	●	?	●	?	●	●	●
Lagos 1995	?	●	?	?	?	●	●	●
Lagos 1999	●	●	?	●	?	?	●	●
Mahalanabis 2008	●	●	?	●	?	●	●	●
Migasena 1989a	?	●	?	●	?	●	●	●
Perry 1998	?	?	?	●	?	●	●	●
Qadri 2005	●	?	?	?	?	●	●	●
Qadri 2007	●	●	?	?	?	●	●	●
Richie 2000 Indonesia	●	●	●	●	?	●	●	●
Sack 1997	?	?	?	●	?	●	●	●
Sanchez 1993a	?	?	?	?	?	●	●	●
Sanchez 1994 Peru	●	●	●	?	●	●	●	●
Sanchez 1995 Peru	●	●	●	●	●	●	●	●
Simanjuntak 1993	●	●	?	?	?	●	●	●
Su-Arehawaratana 1992a	?	?	?	?	?	●	●	●
Su-Arehawaratana 1992b	?	?	?	?	?	●	●	●
Suharyono 1992a	●	●	?	●	?	●	●	●
Suharyono 1992b	●	●	?	●	?	●	●	●
Sur 2009 India	●	●	●	●	●	●	●	●
Tacket 1999	●	●	●	●	●	●	●	●
Taylor 1999a	●	●	?	●	?	●	●	●
Taylor 2000 Peru	●	●	●	●	●	●	●	●
Trach 1997 Viet Nam	●	●	●	?	●	?	●	●
Trach 2002	●	●	?	●	?	●	●	●
Valera 2009	?	●	?	?	?	●	●	●

Measures of treatment effect

All the pre-specified outcomes were dichotomous data and are presented as risk ratios with 95% CIs.

For the occurrence of cholera and diarrhoea cases, the overall risk ratio (RR) has been converted to vaccine efficacy (or effectiveness where intention-to-treat analysis was used) using the formula: % Vaccine Efficacy = $(1 - \text{RR}) \times 100\%$.

Unit of analysis issues

Trials including more than two comparison groups have been split and analysed as individual pair-wise comparisons. When conducting meta-analysis we have ensured that participants and cases in the placebo group were not counted more than once, by dividing the placebo cases and participants evenly between the intervention groups.

Cluster-randomized trials have only been included in the meta-analysis after appropriate adjustment for the effect of clustering. The individualized data has been reduced to the 'effective sample size' by dividing the number of events and the number of participants by the 'design effect'. The design effect was calculated as: $1 + (M-1)\text{ICC}$; where M = average cluster size and ICC = intraclass correlation coefficient. We used estimates of the ICC as presented in the relevant papers.

Dealing with missing data

If data from the trial reports were insufficient, unclear, or missing, we attempted to contact the trial authors for additional information. If we judged the missing data to render the result uninterpretable we have excluded the data from the meta-analysis and clearly stated the reason.

The primary analysis is a complete case analysis where the number of evaluable participants at each time point is used as the denominator.

Assessment of heterogeneity

We assessed for heterogeneity between the trials by examining the forest plot to check for overlapping CIs, by using the Chi² test for heterogeneity using a 10% level of significance, and the I² statistic using a value of 50% to represent moderate levels of heterogeneity. A rough guide to interpretation of the I² statistic is given in the Cochrane Handbook section 9.5.2.

Assessment of reporting biases

There were insufficient trials for us to assess the likelihood of small study effects, such as publication bias, by examining the funnel plot for asymmetry.

Data synthesis

We analysed the data using [Review Manager 5](#). Interventions are compared directly using pair-wise comparisons, and meta-analysis has been performed, where appropriate, if there was more than one trial for a particular comparison. For outcomes that are measured at different time points we have stratified the analysis by the time point.

We have combined studies using the Mantel-Haenszel method with the fixed-effect model. When we have combined the results of trials using different vaccines, or where moderate heterogeneity was detected, we have used the random-effects model. For comparisons which included both individually and cluster-randomized studies; we adjusted the data from the cluster-randomized studies to the 'effective sample size' taking into account the design effect, and then combined the data using the Mantel-Haenszel method. If the reported results of cluster-randomized studies had not been adjusted to take into account the effects of clustering, and we were unable to make these adjustments ourselves, the results are simply reported in tables, and not included in the meta-analysis.

Subgroup analysis and investigation of heterogeneity

We conducted the following subgroup analyses where data were available: age (adult and child, or age under 5 years and over 5 years), time period of follow up, blood group (group O versus other blood groups), type of vaccine, vaccine regimen used or doses received, and whether the challenge was artificial or natural.

Sensitivity analysis

We intended to conduct a sensitivity analysis to evaluate the possible effects of incomplete outcome data by carrying out a best-worst case analysis, such that patients who were lost to follow up were assumed to have the event of interest in one sensitivity analysis and then were assumed to not have the event in a second sensitivity analysis. The data to reliably do this were however not available, so the presented data are a complete-case analysis and represent an assessment of vaccine efficacy, rather than effectiveness.

RESULTS

Description of studies

Results of the search

The search identified 204 references, of which 46 were excluded on abstract alone. Full text copies were obtained of 158 and these were formally assessed using the pre-stated inclusion criteria. Overall, 110 were excluded for the reasons displayed in the [Characteristics of excluded studies](#) table.

Included studies

Forty-eight individual papers have contributed to this review describing 39 separate trials. Fourteen of these describe efficacy data from seven large scale field trials, four describe small artificial challenge efficacy studies, and 29 contribute only safety data. For further details see the [Characteristics of included studies](#) table.

Killed whole cell vaccines

Six trials have evaluated the clinical efficacy of five variations of a killed whole cell vaccine ([Clemens 1988 Bangladesh](#); [Sanchez 1994 Peru](#); [Sanchez 1995 Peru](#); [Taylor 2000 Peru](#); [Trach 1997 Viet Nam](#); [Sur 2009 India](#)).

The composition of these vaccines, the dosing schedule, and the population groups included in these trials are shown in [Table 2](#). The individual vaccines represent step-wise developments from the original vaccines used in [Clemens 1988 Bangladesh](#) to the three vaccines commercially available today.

Two of the field trials used a cluster-randomized design ([Trach 1997 Viet Nam](#); [Sur 2009 India](#)). In order to include these trials in a meta analysis, we have converted the data presented in the original papers to risk ratios, and adjusted for the effect of clustering using the ICC presented in [Sur 2009 India](#). The remaining five trials were individually randomized.

Three of these efficacy trials and 11 additional trials contribute to the safety data for these five vaccines.

Live attenuated vaccines

Only one live attenuated vaccine (CVD 103-HgR) has reached the stage of large scale field evaluation ([Richie 2000 Indonesia](#)). The protective efficacy of two other candidate vaccines: Peru 15 and VC638, has been evaluated in small randomized artificial challenge studies ([Cohen 2002](#); [García 2005](#)). The composition, dosing schedule and population groups included in these trials are shown in [Table 3](#).

An additional 18 trials contributed safety data only to the evaluation of these vaccines.

Excluded studies

Eleven of the excluded trials may be eligible for inclusion in later updates of the review, as the only reason for their exclusion was that no trials assessing the clinical efficacy of these vaccines have been

published; we decided to exclude these early-stage trials because data on safety and tolerability alone is of limited use in practice.

Risk of bias in included studies

Allocation

Efficacy studies

One cluster, quasi-randomized study ([Trach 1997 Viet Nam](#)) used alternate open allocation and three out of the six other efficacy trials did not adequately describe the process of sequence generation or allocation concealment ([Richie 2000 Indonesia](#); [Sanchez 1994 Peru](#); [Sanchez 1995 Peru](#)). However, as the effect of unconcealed allocation in vaccine trials is unlikely to be substantial given that all participants are well prior to enrolment, these trials were judged to be at low risk of bias for these criteria.

Safety (and immunogenicity) only studies

Eleven out of the 29 trials only presenting safety data did not adequately describe the process of allocation concealment for us to make a judgement about the risk of bias.

Blinding

Efficacy outcomes

Six of the seven efficacy trials adequately blinded participants and staff involved with the trial ([Clemens 1988 Bangladesh](#); [Richie 2000 Indonesia](#); [Taylor 2000 Peru](#); [Sanchez 1994 Peru](#); [Sanchez 1995 Peru](#); [Sur 2009 India](#)). One trial was unblinded ([Trach 1997 Viet Nam](#)).

Safety outcomes

Most studies used placebos which were of identical appearance to the vaccine, and could be considered at low risk of bias for safety outcomes. In nine studies the use of a placebo was not adequately described to make a judgement and so were classified as 'unclear'.

Incomplete outcome data

Efficacy studies

Three trials adequately addressed incomplete data for cases of cholera (Taylor 2000 Peru; Sanchez 1994 Peru; Sur 2009 India). In one trial it was unclear whether this had been done, but due to the large sample size and active surveillance system used, this was unlikely to have introduced significant bias (Clemens 1988 Bangladesh). In two trials it was unclear how many participants were lost to follow-up (Richie 2000 Indonesia; Trach 1997 Viet Nam).

Safety (and immunogenicity) only studies

Safety only studies were generally of only short duration with minimal losses to follow-up and therefore considered at low risk of bias.

Selective reporting

We found no evidence of selective reporting bias.

Other potential sources of bias

One trial had evidence of possible other bias (Concha 1995). In this trial, 620 individuals who originally consented to participate dropped out because of a political campaign against it.

Effects of interventions

See: [Summary of findings for the main comparison](#) Summary of findings table: Oral killed whole cell vaccines for preventing cholera

Killed whole cell vaccines

Clinical efficacy

Six trials have evaluated the clinical efficacy of five variations of a killed whole cell vaccine (Clemens 1988 Bangladesh; Sanchez 1994 Peru; Sanchez 1995 Peru; Taylor 2000 Peru; Trach 1997 Viet Nam; Sur 2009 India).

These vaccines are similar but not identical in composition (see Table 2). Despite the variation in dosing schedules, the protective efficacy against confirmed cholera of all five vaccines is similar in both the first and second years following vaccination. It should however be noted that protective efficacy with the two-dose schedule of the WC-rBS vaccine (Dukoral) was not demonstrated in Peru until the second year following a booster dose at 10 months (Taylor 2000 Peru).

The per protocol estimates of protective efficacy as reported in the original papers are shown in Table 4. For comparative purposes we have converted all measures of efficacy to cluster adjusted RRs (Sur 2009 India used rate ratio) and presented these in a forest plot (Year 1 of follow-up: four trials, 252,887 participants: VE 52%, 95% CI 35% to 65%, I^2 49%, Analysis 1.1; Year 2 of follow-up: three trials, 130,334 participants: VE 61%, 95% CI 50% to 70%, I^2 0%, Analysis 1.2).

Evidence of protection for time periods of greater than two years after vaccination is only available for the WC and WC-BS vaccine formulations which are not currently available.

The protective efficacy in children aged less than 5 years was lower than that seen in adults when the data was amalgamated over the first two years of follow-up (four trials, participants; Age < 5 years: VE 38%, 95% CI 20% to 53%, Age > 5 years: VE 66%, 95% CI 57% to 73%, Analysis 1.5; Figure 2). This data was calculated by summing the number cases of cholera in the first two years, and using the number of participants completing 2-years of follow-up as the denominator. A sensitivity analysis using the number of participants completing 1-year follow-up as the denominator did not change the result (Analysis 1.6).

Figure 2. Forest plot of comparison: I Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes, outcome: I.5 Cases of cholera by age group - First two years of follow-up.

Whole cell vaccine (WC: not currently available); three doses given 6 weeks apart

One trial conducted in Bangladesh in 1985 (Clemens 1988 Bangladesh) compared the WC vaccine versus placebo in children aged 2 to 15 years and females aged > 15 years.

Protective efficacy against cholera episodes was established within the first 4-months after vaccination (one trial, 41,580 participants: VE 52%, 95% CI -5% to 78%, Analysis 2.1) and maintained until the third year (Year 1: VE 53%, 95% CI 34% to 66%; Year 2: VE 57%, 95% CI 38% to 70%; Year 3: VE 42%, 95% CI 11% to 62%; Year 4: VE -28%, 95% CI -137% to 31%, Analysis 2.1). Protective efficacy was lost in the fourth year of follow-up.

Vaccine efficacy appears to be lower in children age < 5 years (one trial, 41,580 participants; Year 1: Age 2 to 5 years VE 31%, 95% CI -9% to 57%; Age > 5 years VE 67%, 95% CI 44% to 80%, Analysis 2.2; Year 2: Age 2 to 5 years VE 24%, 95% CI -29% to 55%; Age > 5 years VE 73%, 95% CI 55% to 84%, Analysis 2.3). The difference between vaccine and placebo was not shown to be statistically significant at any time point in this group, although

the trend was towards some protection.

There was also a statistically significant difference between vaccine and placebo in cases of severe watery diarrhoea of any cause (Year 1: VE 32%, 95% CI 7% to 51%, Analysis 2.5), any watery diarrhoea (Year 1: VE 33%, 95% CI 18% to 46%, Analysis 2.5), and diarrhoea of any cause (Year 1: VE 22%, 95% CI 8% to 35%, Analysis 2.5). There is a trend towards a protective effect against all-cause death (VE 23%, 95% CI -1% to 42%), and death from non-dysenteric diarrhoea (VE 53%, 95% CI -16% to 81%), but these did not reach statistical significance (Analysis 2.6).

Whole cell plus B subunit vaccine (WC-BS: not currently available); three doses given 6 weeks apart

The same study (Clemens 1988 Bangladesh) also evaluated the WC-BS vaccine.

Protective efficacy against cholera episodes was similarly demonstrated at 4-months after vaccination (one trial, 41,542 participants: VE 79%, 95% CI 38% to 93%, Analysis 3.1) but evidence

of clinical efficacy was lost in the third year after vaccination (Year 1: VE 62%, 95% CI 46% to 74%; Year 2: VE 58%, 95% CI 40% to 71%; Year 3: 18%, 95% CI -21% to 44%; Year 4: 16%, 95% CI -66% to 58%, [Analysis 3.1](#)).

Vaccine efficacy again appears to be lower in children age < 5 years (one trial, 41,542 participants; Year 1: Age 2 to 5 years VE 38%, 95% CI -1% to 62%; Age > 5 years VE 78%, 95% CI 61% to 87%, [Analysis 3.2](#); Year 2: Age 2 to 5 years VE 47%, 95% CI 3% to 71%; Age > 5 years VE 63%, 95% CI 41% to 76%, [Analysis 3.3](#)).

There was also a statistically significant difference between vaccine and placebo in cases of severe watery diarrhoea of any cause (Year 1: VE 51%, 95% CI 31% to 66%, [Analysis 3.5](#)), any watery diarrhoea (Year 1: VE 38%, 95% CI 23% to 50%, [Analysis 3.5](#)), and diarrhoea of any cause (Year 1: VE 26%, 95% CI 12% to 38%, [Analysis 3.5](#)). All-cause death and death from non-dysenteric diarrhoea were also significantly lower in the group given the vaccine (Year 1: All-cause death: VE 26%, 95% CI 3% to 44%; Deaths from non-dysenteric diarrhoea: VE 80%, 95% CI 31% to 94%, [Analysis 3.6](#)).

No statistically significant difference between WC-BS and WC was demonstrated at any time point, although there was a trend towards increased protection with WC-BS during the first 8-months after vaccination ([Analysis 4.1](#)).

Whole cell plus recombinant vaccine (WC-rBS: available as Dukoral®, SBL); two doses given 2 weeks apart +/- a booster dose at 10 months

One large trial in the general population ([Taylor 2000 Peru](#)), and two smaller trials in military recruits ([Sanchez 1994 Peru](#); [Sanchez 1995 Peru](#)) have evaluated the efficacy of the WC-rBS vaccine; [Taylor 2000 Peru](#) did not demonstrate any significant difference between vaccine or placebo during the first year (one trial, 17,799 participants: VE -4%, 95% CI -105% to 48%, [Analysis 5.1](#)). However, following a booster dose at 10 months the vaccine was superior to placebo in the second year of follow-up (1 trial, 14,999 participants: VE 60%, 95% CI 25% to 79%, [Analysis 5.2](#)).

In the second year of follow-up the estimate of vaccine efficacy was highest in those older than 15 years, although there were very few cholera episodes in the youngest age group (one trial, 14,999 participants, Year 2: Age 2-5 years VE 52%, 95% CI -162% to 91%; Age 5 to 15 years VE 47%, 95% CI -44% to 80%; Age 16 to 65 years VE 71%, 95% CI 22% to 89%; [Analysis 5.2](#)).

Both the small trials in military recruits experienced an outbreak of cholera during or shortly after the vaccination schedule. In [Sanchez 1994 Peru](#) the outbreak occurred 2 to 4 weeks after vaccination. A vaccine efficacy of 86% (95% CI 37% to 97%) was demonstrated in those who received the full two dose schedule, but a single dose did not appear protective (one trial, 1563 participants, [Analysis 5.3](#)). In [Sanchez 1995 Peru](#) the outbreak occurred between the first and second vaccine doses, and vaccine efficacy after one dose

approached statistical significance (VE 44%, 95% CI -4% to 70%, [Analysis 5.3](#)).

Variant whole cell vaccine (vWC: available as ORCVAX®, Vabiotech); two doses given 2 weeks apart

One cluster quasi-randomized trial evaluated the efficacy of the vWC vaccine with 1-year follow-up ([Trach 1997 Viet Nam](#)).

Two doses of vaccine were superior to placebo at preventing cholera episodes requiring in-patient care in all age groups (one trial, 119,033 participants, Age 1 to 5 years VE 68%, 95% CI 14% to 88%, Age > 5 years VE 66%, 95% CI 42% to 80%, authors own figures).

The vaccine was protective against severe and non-severe cholera episodes (one trial, 119,033 participants, Severe episodes VE 65%, 95% CI 34% to 81%, Non-severe episodes VE 56%, 95% CI 26% to 74%, authors own figures).

Bivalent whole cell vaccine (BivWC: available as Shanchol®, Shantha Biotechnics); two doses given 2 weeks apart

One cluster-randomized trial evaluated the efficacy of the BivWC vaccine ([Sur 2009 India](#)). Data are presented for two years of follow-up although the trial is ongoing.

The protective efficacy of the BivWC vaccine was statistically significant during the second but not the first year after vaccination (one trial, 66,900 participants in 3478 clusters: Year 1 VE 45%, 95% CI lower bound -5%, Year 2 VE 77%, 95% CI lower bound 55%, authors own figures).

Over two years follow-up the vaccine was protective in all age groups but lowest in the youngest age group (one trial, 66,900 participants: Age 1 to 4.9 years VE 49%, 95% CI lower bound 6%; Age 5 to 14.9 years VE 87%, 95% CI lower bound 54%; Age > 15 years VE 63%, 95% CI lower bound 23%; authors own figures).

Safety

Whole cell vaccine (WC: not currently available); three doses given 6 weeks apart

Safety data were available from 613 participants. No statistically significant differences were shown between vaccine and placebo after the first or second doses (one trial, 613 participants, [Analysis 6.1](#); [Analysis 3.1](#)).

Whole cell plus B subunit vaccine (WC-BS: not currently available); three doses given 6 weeks apart

Safety data were available from 631 participants. No statistically significant differences were shown between vaccine and placebo

after the first or second doses (one trial, 631 participants, [Analysis 6.2](#); [Analysis 3.2](#))

Whole cell plus recombinant vaccine (WC-rBS: available as Dukoral®, SBL); two doses given 2 weeks apart +/- a booster dose at 10 months

Safety data is available on 12,121 participants who received the WC-rBS vaccine in eight placebo-controlled randomized trials ([Begue 1995](#); [Concha 1995](#); [Hallander 2002](#); [Sanchez 1993a](#); [Sanchez 1995 Peru](#); [Taylor 1999a](#); [Taylor 2000 Peru](#); [Trach 2002](#)). The placebo used in seven of these studies was an oral dose of inactivated *E. coli* (K12 strain).

The largest study ([Taylor 2000 Peru](#)) collected reports of adverse events at the time of the second dose. It found very low levels of symptoms (0.2%), and only the figures for diarrhoea were presented (one study, 10,992 participants, [Analysis 6.3](#)). The remaining studies are small. The only statistically significant result was from one study ([Sanchez 1995 Peru](#)) which found a higher rate of stomach gurgling after the second dose of vaccine (seven trials, 23,870 participants, [Analysis 6.3](#); [Analysis 7.1](#)). The symptoms most commonly reported after taking the vaccine were: stomach gurgling (14%), abdominal pain (9%), headache (5%), and these were generally described as mild.

One additional study translated from Chinese ([Chen 1996](#)) evaluated the safety of a locally formulated WC-rBS in 369 schoolchildren and factory workers and reports no significant differences between vaccine and placebo.

Variant whole cell vaccine (vWC: available as ORCVAX®, Vabiotech); two doses given 2 weeks apart

There is no safety data available for this vaccine.

Bivalent whole cell vaccine (BivWC: available as Shanchol®, Shantha Biotechnics); two doses given 2 weeks apart

Safety data is available from 32,190 participants who received the bivalent whole cell vaccine in four randomized controlled trials ([Mahalanabis 2008](#); [Anh 2007](#); [Kanungo 2009](#); [Sur 2009 India](#)). The placebo used in all four trials was an oral dose of inactivated *E. coli* (K12 strain).

The largest study ([Sur 2009 India](#)) only collected data passively, encouraging participants to present for medical care, and found very low levels of symptoms (<0.2%). It did however record 51 serious adverse events but with no differences between the vaccine and placebo groups. The remaining three studies are small. No clinically important differences between the vaccine and placebo have been shown (four trials, 67,414 participants, [Analysis 6.4](#); [Analysis 7.2](#)). Excluding [Sur 2009 India](#), the symptoms most commonly reported were: abdominal pain (7%), headache (7%), fever (4%), and nausea (3%). These were generally described as mild.

Live attenuated vaccines

Efficacy

Only CVD 103-HgR has been evaluated for clinical efficacy against naturally occurring *V. cholera*. The other live attenuated vaccines listed here remain in development.

CVD 103-HgR (not currently available): one dose

CVD 103-HgR has not been shown to give significant clinical protection from natural cholera infection in any age group (one trial, 67,508 participants, [Analysis 8.1](#); [Analysis 8.2](#)), however only one efficacy study has evaluated this vaccine. This study relied on passive surveillance and the number of cholera events was very low ([Richie 2000 Indonesia](#)). There was also no difference in all-cause death, or deaths related to diarrhoea (one study, 67,508 participants, [Analysis 8.3](#); [Analysis 8.4](#)).

A small artificial challenge study in adult volunteers in the USA ([Tackett 1999](#)) did however, demonstrate a protective effect against moderate to severe diarrhoea (one trial, 51 participants: VE 91%, 95% CI 33% to 99%, [Analysis 8.5](#)) and any diarrhoea (VE 80%, 95% CI 56% to 91%, [Analysis 8.6](#)).

Peru 15 (in development): one dose

One artificial challenge study conducted in adult volunteers in the USA ([Cohen 2002](#)) showed a protective effect against moderate to severe diarrhoea (one trial, 36 participants: VE 95%, 95% CI 21% to 100%, [Analysis 8.5](#)) and any diarrhoea (VE 97%, 95% CI 69% to 100%, [Analysis 8.6](#)). Phase III clinical trials are necessary before conclusions on the clinical efficacy of this vaccine can be made.

VC638 (in development): one dose

One small artificial challenge study conducted in adult volunteers in Cuba ([García 2005](#)) demonstrated a protective effect against any diarrhoea (one trial, 21 participants: VE 99%, 95% CI 68% to 100%, [Analysis 8.6](#)), but not severe diarrhoea ([Analysis 8.5](#)). Phase III clinical trials are necessary before conclusions on the clinical efficacy of this vaccine can be made.

Safety

CVD 103-HgR (not currently available): one dose

A total of 1970 participants have received CVD 103-HgR in fifteen included randomized controlled trials ([Cryz 1990](#); [Gotuzzo 1993](#); [Kotloff 1992](#); [Lagos 1993](#); [Lagos 1995](#); [Lagos 1999](#); [Migasena 1989a](#); [Perry 1998](#); [Richie 2000 Indonesia](#);

Simanjuntak 1993; Su-Arehawaratana 1992a; Su-Arehawaratana 1992b; Suharyono 1992a; Suharyono 1992b; Tacket 1999). The placebo used in 14 of these studies was an oral dose of inactivated *E. coli* (K12 strain).

No symptom was shown to be statistically more common in those given the vaccine (15 trials, 1970 participants, [Analysis 9.1](#)). The commonest reported symptoms following vaccination were: malaise (20% but only recorded in two trials), anorexia (12% but only recorded in three trials), headache (13%), abdominal pain (10%), fever (7%), diarrhoea (5%), vomiting (5%). In general these symptoms are reported to be mild. [Su-Arehawaratana 1992a](#) reports one participant developing diarrhoea after vaccination that required them to seek hospital care.

Peru 15 (in development): one dose

A total of 252 participants have received Peru 15 in four randomized controlled trials ([Cohen 2002](#); [Qadri 2005](#); [Qadri 2007](#); [Sack 1997](#)). The placebo used in these trials was the buffer given alone. Headache was the only symptom to be statistically more common with the vaccine (four trials, 419 participants: Headache RR 4.14, 95% CI 1.27 to 13.48, [Analysis 9.2](#)). The commonest reported symptoms during the first few days after vaccination were: nausea (18%), loss of energy (15%), loss of appetite (10%), and headache (10%). Other adverse events were uncommon, and all adverse events were described as mild.

VC638 (in development): one dose

A total of 90 participants have received VC638 in three randomized studies ([García 2005](#); [Benítez 1999](#); [Valera 2009](#)). The placebo used in these trials was the buffer given alone.

No symptom was shown to be statistically more common in those given vaccine (three trials, 137 participants, [Analysis 9.3](#)). The commonest reported symptoms during the first few days were: stomach gurgling (40%), nausea (33%), abdominal pain (32%), headache (19%), and diarrhoea (13%). Other adverse events were uncommon, and all but one adverse event (a moderate headache) were described as mild.

DISCUSSION

Summary of main results

Killed whole cell vaccines

Five variations of a killed whole cell cholera vaccine have been evaluated in large scale clinical trials. The overall vaccine efficacy during the first year was 52% (95% CI 35% to 65%), and during the second year was 61% (95% CI 50% to 70%).

The protective efficacy over 2-years follow-up was lower in children aged less than 5 years (VE 38%, 95% CI 20% to 53%), than that seen in older age-groups (VE 66%, 95% CI 57% to 73%).

Any observed differences in vaccine efficacy between these vaccines is well within the bounds of random error.

Clinical protection against cholera with the older vaccines (WC and WC-BS) was demonstrated within 4-months of the primary schedule and persisted as long as the third year after vaccination. This cannot be reliably extrapolated to the currently available vaccines given the changes in both the immunisation schedule and the composition of the vaccines.

Of the currently available vaccines:

- A two dose regimen of WC-rBS (Dukoral®) was not shown to be clinically effective in adults in Peru until after a third booster dose was given at 10 months. One smaller trial in military recruits in Peru, did demonstrate a high protective efficacy in a small epidemic occurring within 4 weeks of the two dose schedule but extrapolation of this result beyond short term follow-up may be unreliable. Clinical efficacy in children aged less than 5 years has not been demonstrated.

- A two dose regimen of BivWC (Shanchol®) is likely to be effective during the first and second years after vaccination though this only reached statistical significance during the second year, and follow-up in this trial is ongoing. There is a trend towards protection in all age groups but this was not statistically significant in the under 5 year olds.

- The Vietnam variation of BivWC (mORCVAX®) has not been formally evaluated in published clinical trials. It contains the same elements as Shanchol but has a different manufacturing process.

Live attenuated vaccines

The live attenuated vaccines remain in development. The only vaccine to reach Phase III clinical trials and licensure in some countries, CVD 103-HgR, has not been shown to provide a protective effect against clinical cholera episodes; however, it has only been evaluated in one large efficacy trial in which there were few cases in either arm.

Overall completeness and applicability of evidence

The currently available vaccines represent stepwise modifications to the original vaccines developed and studied in Bangladesh in the 1980s. Although changes have occurred in both the composition and the recommended vaccination schedule, they remain similar enough to sensibly combine in a meta-analysis, and this is confirmed by their remarkably similar efficacies. The efficacy data from these older studies and vaccines therefore remains relevant to the assessment of the WC-rBS (Dukoral®) and BivWC (Shanchol®) vaccines available today.

The current recommended schedule for WC-rBS is two doses 2 weeks apart, and three doses 2 weeks apart for children age 2 to 5 years. The two dose schedule (rather than the three doses used in the Bangladesh study) has been adopted based on immunological data, and the observation that two doses of the original WC and WC-BS vaccines were equally effective to three doses in the Bangladesh study (Clemens 1988 Bangladesh). Unfortunately we have been unable to get access to the data to confirm this finding. The lack of protective efficacy with a two-dose schedule seen in the only large scale trial of WC-rBS (Taylor 2000 Peru) has been discussed in the literature with questions raised about the adequacy and accuracy of the cholera surveillance during the first year of follow-up (Clemens 2001; Taylor 2001). Reassuringly two doses were protective in the much smaller military trials, but the number of events was low and the period of follow-up inadequate to make conclusions for the use of the vaccine outside of an acute epidemic situation. Although the two dose schedule of BivWC has been shown to be protective in the first year (though not quite reaching statistical significance), this vaccine is sufficiently different from WC-rBS to restrict the generalisation of this result. The primary analysis used in this review is a complete-case analysis excluding participants who received incomplete vaccine schedules. These findings will therefore tend to overestimate the effectiveness of the vaccine when given outside of trial settings, where vaccine coverage will almost always be considerably less than 100%. This factor should be taken into consideration when planning a cholera vaccination programme.

The best evidence for the use of cholera vaccines in epidemic situations, such as seen in Zimbabwe and Haiti in recent years, comes from the two trials in adult military recruits. Sanchez 1994 Peru demonstrated 86% protective efficacy (95% CI 37% to 97%) in a small epidemic occurring within 4 weeks of the two-dose schedule of WC-rBS. The reactive use of cholera vaccines once an epidemic has begun has been further evaluated through case-control studies (Anh 2011), and modelling exercises (Reyburn 2011), which are outside of the scope of this review (Ryan 2011).

Quality of the evidence

The quality of the evidence was assessed using the GRADE methodology. Overall the quality is moderate to high, meaning that we can have a high degree of confidence in these results, and further research is unlikely to substantially alter the current esti-

mates of protective efficacy. See Summary of findings table 1.

Agreements and disagreements with other studies or reviews

The World Health Organization published a position paper on oral cholera vaccines in 2010 (WHO 2010b). The findings presented here are in broad agreement with this paper.

AUTHORS' CONCLUSIONS

Implications for practice

The currently available oral killed whole cell vaccines can prevent 50 to 60% of cholera episodes during the first 2-years after the primary vaccination schedule. Protective efficacy is unlikely to last more than 3 years and booster doses in line with the manufacturers recommendations will be required.

The impact and cost-effectiveness of adopting oral cholera vaccines into the routine vaccination schedule of endemic countries will depend on the prevalence of cholera among the community, the frequency of epidemics, and the availability or unavailability of adequate facilities to provide rapid rehydration therapy.

Although there is currently little high quality evidence for the effect of vaccines in emergency and epidemic situations, It is likely that cholera vaccines would have an important impact on reducing disease in epidemics, especially where access to clean water and sanitation is difficult to achieve.

Implications for research

The evidence from Peru suggests that countries considering routine vaccination should assess whether the two-dose primary immunization schedule is adequate in their setting.

ACKNOWLEDGEMENTS

The editorial base for the Cochrane Infectious Disease Group is funded by the Department for International Development (DFID), UK, for the benefit of developing countries.

REFERENCES

References to studies included in this review

Anh 2007 *{published data only}*

Anh DD, Canh do G, Lopez AL, Thiem VD, Long PT, Son NH, et al. Safety and immunogenicity of a reformulated Vietnamese bivalent killed, whole-cell, oral cholera vaccine in adults. *Vaccine* 2007; Vol. 25, issue 6:1149–55.

Begue 1995 *{published data only}*

Begue RE, Castellares G, Ruiz R, Hayashi KE, Sanchez JL, Gotuzzo E, Oberst RB, Taylor DN, Svennerholm A-M. Community-based assessment of safety and immunogenicity of the whole cell plus recombinant b subunit (WC/rBS) oral cholera vaccine in Peru. *Vaccine* 1995;13(7):691–4.

Benítez 1999 *{published data only}*

Benítez JA, García L, Silva A, García H, Fando R, Cedré B, et al. Preliminary assessment of the safety and immunogenicity of a new CTXPhi-negative, hemagglutinin/protease-defective El Tor strain as a cholera vaccine candidate. *Infection and Immunity* 1999; Vol. 67, issue 2:539–45.

Chen 1996 *{published data only}*

Chen Q, Yu S, Wang Y. [Community trial for safety and immunogenicity of oral-administered lyophilized rBS-WC cholera vaccine]. *Zhonghua yu fang yi xue za zhi* [Chinese journal of preventive medicine] 1996; Vol. 30, issue 6: 330–3.

Clemens 1987 *{published data only}*

Clemens JD, Stanton BF, Chakraborty J, Sack DA, Khan MR, Huda S, et al. B subunit-whole cell and whole cell-only oral vaccines against cholera: studies on reactogenicity and immunogenicity. *Journal of Infectious Diseases* 1987; 155(1):79–85.

Clemens 1988 Bangladesh *{published data only}*

* Clemens JD, Harris JR, Sack DA, Chakraborty J, Ahmed F, Stanton BF, et al. Field trial of oral cholera vaccine in Bangladesh: results from one year of follow-up. *Journal of Infectious Diseases* 1988;158(1):60–9.
Clemens JD, Harris JR, Sack DA, Chakraborty J, Ahmed F, Stanton BF, et al. Field trial of oral cholera vaccines in Bangladesh. *Southeast Asian Journal of Tropical Medicine and Public Health* 1988;19(3):417–422.
Clemens JD, Sack DA, Chakraborty J, Rao MR, Ahmed F, Harris JR, et al. Field trial of oral cholera vaccines in Bangladesh: evaluation of anti-bacterial and anti-toxic breast-milk immunity in response to ingestion of the vaccines. *Vaccine* 1990;8:469–72.
Clemens JD, Sack DA, Harris JR, Chakraborty J, Khan MR, Huda S, et al. ABO blood groups and cholera: new observations on specificity of risk and modifications of vaccine efficacy. *Journal of Infectious Diseases* 1989;159(4): 770–3.
Clemens JD, Sack DA, Harris JR, Chakraborty J, Khan MR, Stanton BF, et al. Field trial of oral cholera vaccines in Bangladesh. *Lancet* 1986;2(8499):124–7.
Clemens JD, Sack DA, Harris JR, Chakraborty J, Khan MR, Stanton BF, et al. Impact of b subunit killed whole-

cell and killed whole-cell-only oral vaccines against cholera upon treated diarrhoeal illness and mortality in an area endemic for cholera. *Lancet* 1988;1(8599):1375–9.

Clemens JD, Sack DA, Harris JR, van Loon F, Chakraborty J, Ahmed F, et al. Field trial of oral cholera vaccines in Bangladesh: results from three-year follow-up. *Lancet* 1990; 335(8684):270–73.

van Loon FPL, Clemens JD, Chakraborty J, Rao MR, Kay BA, Sack DA, et al. Field trial of inactivated oral cholera vaccines in Bangladesh: results from 5 years of follow-up. *Vaccine* 1996;14(2):162–6.

Cohen 2002 *{published data only}*

Cohen MB, Giannella RA, Bean J, Taylor DN, Parker S, Hoepfer A, et al. Randomized, controlled human challenge study of the safety, immunogenicity, and protective efficacy of a single dose of Peru-15, a live attenuated oral cholera vaccine. *Infection and Immunity* 2002; Vol. 70, issue 4: 1965–70.

Concha 1995 *{published data only}*

Concha A, Giraldo A, Castaneda E, Martinez M, de la Hoz F, Rivas F, et al. Safety and immunogenicity of oral killed whole cell recombinant B subunit cholera vaccine in Barranquilla, Colombia. [Seguridad e inmunogenicidad de la vacuna anticólera de células enteras muertas y de la subunidad B recombinante de la toxina en Barranquilla, Colombia]. *Bol Of Sanit Panam* 1995;119(2):103–12.

* Concha A, Giraldo A, Castañeda E, Martínez M, de la Hoz F, Rivas F, et al. Safety and immunogenicity of oral killed whole cell recombinant B subunit cholera vaccine in Barranquilla, Colombia. *Bulletin of the Pan American Health Organization* 1995; Vol. 29, issue 4:312–21.

Cryz 1990 *{published data only}*

Cryz SJ, Levine MM, Kaper JB, Fürer E, Althaus B. Randomized double-blind placebo controlled trial to evaluate the safety and immunogenicity of the live oral cholera vaccine strain CVD 103-HgR in Swiss adults. *Vaccine* 1990; Vol. 8, issue 6:577–80.

García 2005 *{published data only}*

García L, Jidy MD, García H, Rodríguez BL, Fernández R, Año G, et al. The vaccine candidate *Vibrio cholerae* 638 is protective against cholera in healthy volunteers. *Infection and Immunity* 2005; Vol. 73, issue 5:3018–24.

Gotuzzo 1993 *{published data only}*

Gotuzzo E, Butron B, Seas C, Penny M, Ruiz R, Losonsky G, et al. Safety, immunogenicity, and excretion pattern of single-dose live oral cholera vaccine CVD 103-HgR in Peruvian adults of high and low socioeconomic levels. *Infection and Immunity* 1993; Vol. 61, issue 9:3994–7.

Hallander 2002 *{published data only}*

Hallander HO, Paniagua M, Espinoza F, Askelöf P, Corrales E, Ringman M, et al. Calibrated serological techniques demonstrate significant different serum response rates to an oral killed cholera vaccine between Swedish and Nicaraguan children. *Vaccine* 2002; Vol. 21, issue 1–2:138–45.

Kanungo 2009 {published data only}

Kanungo S, Paisley A, Lopez AL, Bhattacharya M, Manna B, Kim DR, et al. Immune responses following one and two doses of the reformulated, bivalent, killed, whole-cell, oral cholera vaccine among adults and children in Kolkata, India: a randomized, placebo-controlled trial. *Vaccine* 2009; Vol. 27, issue 49:6887–93.

Kotloff 1992 {published data only}

Kotloff KL, Wasserman SS, O'Donnell S, Losonsky GA, Cryz SJ, Levine MM. Safety and immunogenicity in North Americans of a single dose of live oral cholera vaccine CVD 103-HgR: results of a randomized, placebo-controlled, double-blind crossover trial. *Infection and Immunity* 1992; Vol. 60, issue 10:4430–2.

Lagos 1993 {published data only}

Lagos R, Avendaño A, Horwitz I, Prado V, Ferreccio C, Sotomayor V, et al. [Tolerance and immunogenicity of an oral dose of CVD 103-HgR, a live attenuated *Vibrio cholerae* 01 strain: a double-blind study of Chilean adults]. *Revista médica de Chile* 1993; Vol. 121, issue 8:857–63.

Lagos 1995 {published data only}

Lagos R, Avendaño A, Prado V, Horwitz I, Wasserman S, Losonsky G, et al. Attenuated live cholera vaccine strain CVD 103-HgR elicits significantly higher serum vibriocidal antibody titers in persons of blood group O. *Infection and Immunity* 1995; Vol. 63, issue 2:707–9.

Lagos 1999 {published data only}

Lagos R, San Martin O, Wasserman SS, Prado V, Losonsky GA, Bustamante C, et al. Palatability, reactogenicity and immunogenicity of engineered live oral cholera vaccine CVD 103-HgR in Chilean infants and toddlers. *The Pediatric Infectious Disease Journal* 1999; Vol. 18, issue 7: 624–30.

Mahalanabis 2008 {published data only}

Mahalanabis D, Lopez AL, Sur D, Deen J, Manna B, Kanungo S, et al. A randomized, placebo-controlled trial of the bivalent killed, whole-cell, oral cholera vaccine in adults and children in a cholera endemic area in Kolkata, India. *PLoS ONE* 2008; Vol. 3, issue 6:e2323.

Migasena 1989a {published data only}

Migasena S, Pitisuttitham P, Prayurachong B, Suntharasamai P, Supanaranond W, Desakorn V, et al. Preliminary assessment of the safety and immunogenicity of live oral cholera vaccine strain CVD 103-HgR in healthy Thai adults. *Infection and Immunity* 1989; Vol. 57, issue 11: 3261–4.

Perry 1998 {published data only}

Perry RT, Plowe CV, Koumaré B, Bougoudogo F, Kotloff KL, Losonsky GA, et al. A single dose of live oral cholera vaccine CVD 103-HgR is safe and immunogenic in HIV-infected and HIV-noninfected adults in Mali. *Bulletin of the World Health Organization* 1998; Vol. 76, issue 1: 63–71.

Qadri 2005 {published data only}

Qadri F, Chowdhury MI, Faruque SM, Salam MA, Ahmed T, Begum YA, et al. Randomized, controlled study of the

safety and immunogenicity of Peru-15, a live attenuated oral vaccine candidate for cholera, in adult volunteers in Bangladesh. *The Journal of Infectious Diseases* 2005; Vol. 192, issue 4:573–9.

Qadri 2007 {published data only}

Qadri F, Chowdhury MI, Faruque SM, Salam MA, Ahmed T, Begum YA, et al. Peru-15, a live attenuated oral cholera vaccine, is safe and immunogenic in Bangladeshi toddlers and infants. *Vaccine*. 2007; Vol. 25, issue 2:231–8.

Richie 2000 Indonesia {published data only}

Punjabi N, Simanjuntak C, Richie E, Sidharta Y, et al. Large scale, randomized, double blind, placebo controlled field trial to assess the efficacy of a single dose of live oral cholera vaccine CVD 103-HgR. [abstract]. *American Journal of Tropical Medicine and Hygiene* 1998;59(3 Suppl):249–50.
* Richie E, Punjabi NH, Sidharta Y, Peetosutan K, Sukander M, Wasserman SS, et al. Efficacy trial of single dose live oral vaccine CVD 103-HgR in North Jakarta, Indonesia, a cholera endemic area. *Vaccine* 2000;18:2399–410.

Sack 1997 {published data only}

Sack DA, Sack RB, Shimko J, Gomes G, O'Sullivan D, Metcalfe K, et al. Evaluation of Peru-15, a new live oral vaccine for cholera, in volunteers. *The Journal of Infectious Diseases* 1997; Vol. 176, issue 1:201–5.

Sanchez 1993a {published data only}

Sanchez JL, Trofa AF, Taylor DN, Kuschner RA, DeFraites RF, Craig SC, et al. Safety and immunogenicity of the oral, whole cell/recombinant B subunit cholera vaccine in North American volunteers. *The Journal of Infectious Diseases* 1993; Vol. 167, issue 6:1446–9.

Sanchez 1994 Peru {published data only}

* Sanchez JL, Vasquez B, Begue RE, Meza R, Castellares G, Cabezas C, et al. Protective efficacy of oral whole-cell/recombinant-b-subunit cholera vaccine in Peruvian military recruits. *Lancet* 1994;344(8932):1273–76.

Sanchez 1995 Peru {published data only}

* Sanchez JL, Hayashi KE, Kruger HF, Meza R, English CK, Vidal W, et al. Immunological response to vibrio cholerae 01 infection and an oral cholera vaccine among Peruvians. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1995;89(5):542–5.

Simanjuntak 1993 {published data only}

Simanjuntak CH, O'Hanley P, Punjabi NH, Noriega F, Pazzaglia G, Dykstra P, et al. Safety, immunogenicity, and transmissibility of single-dose live oral cholera vaccine strain CVD 103-HgR in 24- to 59-month-old Indonesian children. *The Journal of Infectious Diseases* 1993; Vol. 168, issue 5:1169–76.

Su-Arehawaratana 1992a {published data only}

Su-Arehawaratana P, Singharaj P, Taylor DN, Hoge C, Trofa A, Kuanont K, et al. Safety and immunogenicity of different immunization regimens of CVD 103-HgR live oral cholera vaccine in soldiers and civilians in Thailand. *The Journal of Infectious Diseases* 1992; Vol. 165, issue 6: 1042–8.

Su-Arehawaratana 1992b {published data only}

Su-Arehawaratana P, Singharaj P, Taylor DN, Hoge C, Trofa A, Kuvanont K, et al. Safety and immunogenicity of different immunization regimens of CVD 103-HgR live oral cholera vaccine in soldiers and civilians in Thailand. *The Journal of Infectious Diseases* 1992;**165**(6):1042–8.

Suharyono 1992a {published data only}

Suharyono, Simanjuntak C, Witham N, Punjabi N, Heppner DG, Losonsky G, et al. Safety and immunogenicity of single-dose live oral cholera vaccine CVD 103-HgR in 5-9-year-old Indonesian children. *Lancet* 1992; Vol. 340, issue 8821:689–94.

Suharyono 1992b {published data only}

Suharyono, Simanjuntak C, Witham N, Punjabi N, Heppner DG, Losonsky G, et al. Safety and immunogenicity of single-dose live oral cholera vaccine CVD 103-HgR in 5-9-year-old Indonesian children. *Lancet* 1992;**340**(8821):689–94.

Sur 2009 India {published data only}

* Sur D, Lopez AL, Kanungo S, Paisley A, Manna B, Ali M, et al. Efficacy and safety of a modified killed-whole-cell oral cholera vaccine in India: an interim analysis of a cluster-randomised, double-blind, placebo-controlled trial. *Lancet* 2009; Vol. 374, issue 9702:1694–702.

Tacket 1999 {published data only}

Tacket CO, Cohen MB, Wasserman SS, Losonsky G, Livio S, Kotloff K, et al. Randomized, double-blind, placebo-controlled, multicentered trial of the efficacy of a single dose of live oral cholera vaccine CVD 103-HgR in preventing cholera following challenge with *Vibrio cholerae* O1 El Tor inaba three months after vaccination. *Infection and Immunity* 1999; Vol. 67, issue 12:6341–5.

Taylor 1999a {published data only}

Taylor DN, Cárdenas V, Perez J, Puga R, Svennerholm AM. Safety, immunogenicity, and lot stability of the whole cell/recombinant B subunit (WC/rCTB) cholera vaccine in Peruvian adults and children. *The American Journal of Tropical Medicine and Hygiene* 1999; Vol. 61, issue 6: 869–73.

Taylor 2000 Peru {published data only}

* Taylor DN, Cardenas V, Sanchez JL, Begue RE, Gilman R, Bautista C, et al. Two-year study of the protective efficacy of the oral whole cell plus recombinant B subunit cholera vaccine in Peru. *Journal of Infectious Diseases* 2000; **181**:1667–73.

Trach 1997 Viet Nam {published data only}

Trach DD, Clemens JD, Ke NT, Thuy HT, Son ND, Canh DG, et al. Field trial of a locally produced, killed, oral cholera vaccine in Vietnam. *Lancet* 1997;**349**(9047):231–5.

Trach 2002 {published data only}

Trach DD, Cam PD, Ke NT, Rao MR, Dinh D, Hang PV, et al. Investigations into the safety and immunogenicity of a killed oral cholera vaccine developed in Viet Nam. *Bulletin of the World Health Organization* 2002; Vol. 80, issue 1: 2–8.

Valera 2009 {published data only}

Valera R, Garcia HM, Jidy MD, Mirabal M, Armesto MI, Fando R, et al. Randomized, double-blind, placebo-controlled trial to evaluate the safety and immunogenicity of live oral cholera vaccine 638 in Cuban adults. *Vaccine* 2009; Vol. 27, issue 47:6564–9.

References to studies excluded from this review

Ahmed 2006 {published data only}

Ahmed A, Li J, Shiloach Y, Bobbins JB, Szu SC. Safety and immunogenicity of *Escherichia coli* O157 O-specific polysaccharide conjugate vaccine in 2-5-year-old children. *Journal of Infectious Diseases* 2006; Vol. 193, issue 4: 515–21.

Ahren 1993 {published data only}

Ahren C, Wenneras C, Holmgren J, Svennerholm A-M. Intestinal antibody response after oral immunization with a prototype cholera B subunit-colonization factor antigen enterotoxigenic *Escherichia coli* vaccine. *Vaccine* 1993;**11** (9):929–34.

Albert 2003 {published data only}

Albert MJ, Qadri F, Wahed MA, Ahmed T, Rahman AS, Ahmed F, et al. Supplementation with zinc, but not vitamin A, improves seroconversion to vibriocidal antibody in children given an oral cholera vaccine. *The Journal of Infectious Diseases* 2003; Vol. 187, issue 6:909–13.

Ali 2005 {published data only}

Ali M, Emch M, von Seidlein L, Yunus M, Sack DA, Rao M, et al. Herd immunity conferred by killed oral cholera vaccines in Bangladesh: a reanalysis. *Lancet* 2005;**366** (9479):44–9.

Ali 2008 {published data only}

Ali M, Emch M, Yunus M, Sack D, Lopez AL, Holmgren J, Clemens J. Vaccine protection of Bangladeshi infants and young children against cholera: implications and for vaccine deployment and person-to-person transmission. *Pediatric Infectious Disease Journal* 2008;**27**(1):33–7.

Anonymous 1968 {published data only}

Anonymous. A controlled field trial of the effectiveness of various doses of cholera El Tor vaccine in the Philippines. *Bulletin of the World Health Organization* 1968; Vol. 38, issue 6:917–23.

Anonymous 1973a {published data only}

Anonymous. A controlled field trial of the effectiveness of monovalent classical and El Tor cholera vaccines in the Philippines. *Bulletin of the World Health Organization* 1973; Vol. 49, issue 1:13–9.

Anonymous 1973b {published data only}

Anonymous. A controlled field trial on the effectiveness of the intradermal and subcutaneous administration of cholera vaccine in the Philippines. *Bulletin of the World Health Organization* 1973; Vol. 49, issue 4:389–94.

Azurin 1967 {published data only}

Azurin JC, Cruz A, Pesigan TP, Alvero M, Camena T, Suplido R, et al. A controlled field trial of the effectiveness of cholera and cholera El Tor vaccines in the Philippines.

- Bulletin of the World Health Organization 1967; Vol. 37, issue 5:703–27.
- Azurin 1971** *{published data only}*
Azurin JC, Alvero M. Cholera incidence in a population offered cholera vaccination: comparison of cooperative and uncooperative groups. Bulletin of the World Health Organization 1971; Vol. 44, issue 6:815–9.
- Benenson 1968a** *{published data only}*
Benenson AS, Joseph PR, Oseasohn RO. Cholera vaccine field trials in east Pakistan. 1. Reaction and antigenicity studies. Bulletin of the World Health Organization 1968; Vol. 38, issue 3:347–57.
- Benenson 1968b** *{published data only}*
Benenson AS, Mosley WH, Fahimuddin M, Oseasohn RO. Cholera vaccine field trials in east Pakistan. 2. Effectiveness in the field. Bulletin of the World Health Organization 1968; Vol. 38, issue 3:359–72.
- Bergquist 1997** *{published data only}*
Bergquist C, Johansson EL, Lagergard T, Holmgren J, Rudin A. Intranasal vaccination of humans with recombinant cholera toxin B subunit induces systemic and local antibody responses in the upper respiratory tract and the vagina. Infection and Immunity 1997; Vol. 65, issue 7:2676–84.
- Black 1986** *{published data only}*
* Black RE, Levine MM, Clements ML, Young CR, Svennerholm A-M, Holmgren J, Germanier R. Oral immunization with killed whole vibrio and B subunit or procholeraenoid combination cholera vaccines: immune response and protection from V.cholerae challenge. In: Kuwahara S, Peirce NF editor(s). *Advances in Research on Cholera and Related Diarrheas*. Vol. 3, Tokyo: KTK Scientific Publications, 1986:271–5.
- Black 1987** *{published data only}*
* Black RE, Levine MM, Clements ML, Young CR, Svennerholm A-M, Holmgren J. Protective efficacy in humans of killed whole-vibrio oral cholera vaccine with and without the B subunit of cholera toxin. *Infection and Immunity* 1987;55(5):1116–20.
- Burgasov 1976** *{published data only}*
Burgasov PN, Sumarokov AA, Lelikov VL, Marcuk LM, Fedenev VG, Dzaparidze MN, et al. Comparative study of reactions and serological response to cholera vaccines in a controlled field trial conducted in the USSR. Bulletin of the World Health Organization 1976; Vol. 54, issue 2:163–70.
- Bwanga 1984** *{published data only}*
Bwanga M. [Initial controlled clinical trials of an oral anticholera vaccine during the cholera epidemic in the district of Malemba-Nkulu (Shaba-Zaire)]. Bulletin de la Société de pathologie exotique et de ses filiales 1984; Vol. 77, issue 1:13–6.
- Cash 1974** *{published data only}*
Cash RA, Music SI, Libonati JP, Schwartz AR, Hornick RB. Live oral cholera vaccine: evaluation of the clinical effectiveness of two strains in humans. Infection and Immunity 1974; Vol. 10, issue 4:762–4.
- Cavailler 2006** *{published data only}*
Cavailler P, Lucas M, Perroud V, McChesney M, Ampuero S, Guerin PJ, et al. Feasibility of a mass vaccination campaign using a two-dose oral cholera vaccine in an urban cholera-endemic setting in Mozambique. *Vaccine*. 2006; Vol. 24, issue 22:4890–5.
- Chaicumpa 1998** *{published data only}*
* Chaicumpa W, Chongsanguan M, Kalambaheti T, Wilairatana P, Srimanote P, Makakunkijcharoen Y, et al. Immunogenicity of liposome-associated and refined antigen oral cholera vaccine in Thai volunteers. *Vaccine* 1998;16(7):678–84.
- Chongsanguan 1988** *{published data only}*
Chongsanguan M, Chaicumpa W, Ruangunaporn Y, Looareesuwan S. Immunogenicity of two formulations of oral cholera vaccines in Thai volunteers. *Vaccine* 1991;9(1):53–9.
- Chongsanguan 1991** *{published data only}*
Chongsanguan M, Chaicumpa W, Ruangunaporn Y, Looareesuwan S. Immunogenicity of two formulations of oral cholera vaccines in Thai volunteers. *Vaccine* 1991; Vol. 9, issue 1:53–9.
- Ciznar 1989** *{published data only}*
Ciznar I, Hussain N, Ahsan CR, Kay BA, Clemens JD, Sack DA. Oral cholera vaccines containing B-subunit-killed whole cells and killed whole cells only. I. Cross-reacting antigens of members of family Vibrionaceae and the vaccines. *Vaccine* 1989; Vol. 7, issue 2:111–6.
- Clemens 1986** *{published data only}*
Clemens JD, Jertborn M, Sack D, Stanton B, Holmgren J, Khan MR, Huda S. Effect of neutralization of gastric acid on immune responses to an oral b subunit, killed whole-cell cholera vaccine. *Journal of Infectious Diseases* 1986;154(1):175–8.
- Clemens 1988** *{published data only}*
Clemens JD, Sack DA, Harris JR, Chakraborty J, Neogy PK, Stanton B, et al. Cross-protection by B subunit-whole cell Cholera vaccine against diarrhea associated with heat-labile toxin-producing enterotoxigenic escherichia coli: results of a large-scale field trial. *Journal of Infectious Diseases* 1988;158(2):372–7.
- Clemens 1989a** *{published data only}*
Clemens JD, Harris JR, Kay BA, Chakraborty J, Sack DA, Ansaruzzaman M, et al. Oral cholera vaccines containing B-subunit-killed whole cells and killed whole cells only. II. Field evaluation of cross-protection against other members of the vibronaceae family. *Vaccine* 1989;7(2):117–20.
- Clemens 1989b** *{published data only}*
Clemens JD, Stanton BF, Harris JR, Chakraborty J, Dack DA, Rao MR, et al. Exclusion of clinically atypical or microbiologically mixed diarrhoeal episodes from outcome events in a field trial of oral cholera vaccines. *International Journal of Epidemiology* 1989;18(2):440–5.
- Clemens 1990** *{published data only}*
Clemens JD, Sack DA, Chakraborty J, Rao MR, Ahmed F, Harris JR, et al. Field trial of oral cholera vaccines in

- Bangladesh: evaluation of anti-bacterial and anti-toxic breast-milk immunity in response to ingestion of the vaccines. *Vaccine* 1990;**8**:469–72.
- Clemens 1991** *{published data only}*
 * Clemens JD, van Loon F, Sack DA, Chakraborty J, Rao MR, Ahmed F, et al. Field trial of oral cholera vaccines in Bangladesh: serum vibriocidal and antitoxic antibodies as markers of the risk of cholera. *Journal of Infectious Diseases* 1991;**163**:1235–42.
- Clemens 1992a** *{published data only}*
 Clemens JD, van Loon FF, Rao M, Sack DA, Ahmed F, Chakraborty J, et al. Nonparticipation as a determinant of adverse health outcomes in a field trial of oral cholera vaccines. *American Journal of Epidemiology* 1992; Vol. 135, issue 8:865–74.
- Clemens 1992b** *{published data only}*
 Clemens JD, Sack DA, Rao MR, Chakraborty J, Khan MR, Kay B, et al. Evidence that inactivated oral cholera vaccines both prevent and mitigate vibrio cholerae 01 infections in a cholera-endemic area. *Journal of Infectious Diseases* 1992; **166**:1029–34.
- Clemens 1995** *{published data only}*
 Clemens J, Rao M, Sack D, Ahmed F, Khan MR, Chakraborty J, et al. Impaired immune response to natural infection as a correlate of vaccine failure in a field trial of killed oral cholera vaccines. *American Journal of Epidemiology* 1995;**142**(7):759–64.
- Cohen 1999** *{published data only}*
 Cohen MB, Giannella RA, Losonsky GA, Lang DR, Parker S, Hawkins JA, et al. Validation and characterization of a human volunteer challenge model for cholera by using frozen bacteria of the new *Vibrio cholerae* epidemic serotype, O139. *Infection & Immunity* 1999; Vol. 67, issue 12:6346–9.
- Cohen 2000** *{published data only}*
 Cohen D, Orr N, Haim M, Ashkenazi S, Robin G, Green MA, et al. Safety and immunogenicity of two different lots of the oral, killed enterotoxigenic *Escherichia coli*-cholera toxin B subunit vaccine in young Israeli adults. *Infection and Immunology* 2000;**68**(8):4492–7.
- Cooper 2000** *{published data only}*
 Cooper PJ, Chico ME, Losonsky G, Sandoval C, Espinel I, Sridhara R, et al. Albendazole treatment of children with ascariasis enhances the vibriocidal antibody response to the live attenuated oral cholera vaccine CVD 103-HgR. *The Journal of Infectious Diseases* 2000; Vol. 182, issue 4: 1199–206.
- Cooper 2001** *{published data only}*
 Cooper PJ, Chico M, Sandoval C, Espinel I, Guevara A, Levine MM, et al. Human infection with *Ascaris lumbricoides* is associated with suppression of the interleukin-2 response to recombinant cholera toxin B subunit following vaccination with the live oral cholera vaccine CVD 103-HgR. *Infection and Immunity* 2001; Vol. 69, issue 3:1574–80.
- Coster 1995** *{published data only}*
 Coster TS, Killeen KP, Waldor MK, Beattie DT, Spriggs DR, Kenner JR, et al. Safety, immunogenicity, and efficacy of live attenuated *Vibrio cholerae* O139 vaccine prototype. *Lancet* 1995; Vol. 345, issue 8955:949–52.
- Cryz 1992** *{published data only}*
 Cryz Jr SJ, Levine MM, Losonsky G, Kaper JB, Althaus B. Safety and immunogenicity of a booster dose of *Vibrio cholerae* CVD 103-HgR live oral cholera vaccine in Swiss adults. *Infection and Immunity* 1992; Vol. 60, issue 9: 3916–7.
- Cryz 1995** *{published data only}*
 Cryz SJ, Que JU, Levine MM, Wiedermann G, Kollaritsch H. Safety and immunogenicity of a live oral bivalent typhoid fever (*Salmonella typhi* Ty21a)-cholera (*Vibrio cholerae* CVD 103-HgR) vaccine in healthy adults. *Infection and Immunity* 1995; Vol. 63, issue 4:1336–9.
- Das 1967** *{published data only}*
 Das Gupta A, Sinha R, Shrivastava DL, De SP, Taneja BL, Rao MS, et al. Controlled field trial of the effectiveness of cholera and cholera El Tor vaccines in Calcutta. *Bulletin of the World Health Organization* 1967; Vol. 37, issue 3: 371–85.
- Dearlove 1992** *{published data only}*
 Dearlove CE, Forrest BD, van den Bosch L, La Brooy JT. The antibody response to an oral Ty21a-based typhoid-cholera hybrid is unaffected by prior oral vaccination with Ty21a. *The Journal of Infectious Diseases* 1992; Vol. 165, issue 1:182–3.
- Durham 1998** *{published data only}*
 Durham LK, Longini IM Jr, Halloran ME, Clemens JD, Nizam A, Rao M. Estimation of vaccine efficacy in the presence of waning: application to cholera vaccines. *American Journal of Epidemiology* 1998;**147**(10):948–59.
- Emch 2006** *{published data only}*
 Emch M, Ali M, Park JK, Yunus M, Sack DA, Clemens JD. Relationship between neighbourhood-level killed oral cholera vaccine coverage and protective efficacy: evidence for herd immunity. *International Journal of Epidemiology* 2006;**35**(4):1044–50.
- Emch 2007** *{published data only}*
 Emch M, Ali M, Acosta C, Yunus M, Sack DA, Clemens JD. Efficacy calculation in randomized trials: global or local measures?. *Health Place* 2007;**13**(1):238–48.
- Emch 2009** *{published data only}*
 Emch M, Ali M, Root ED, Yunus M. Spatial and environmental connectivity analysis in a cholera vaccine trial. *Social science & medicine* (1982) 2009;**68**(4):631–7.
- Forrest 1991** *{published data only}*
 Forrest BD, LaBrooy JT. In vivo evidence of immunological masking of the *Vibrio cholerae* O antigen of a hybrid *Salmonella typhi* Ty21a-*Vibrio cholerae* oral vaccine in humans. *Vaccine* 1991; Vol. 9, issue 7:515–20.
- Ganguly 1975** *{published data only}*
 Ganguly R, Clem LW, Bencic Z, Sinha R, Sakazaki R, Waldman RH. Antibody response in the intestinal

- secretions of volunteers immunized with various cholera vaccines. *Bulletin of the World Health Organization* 1975;**52** (3):323–30.
- Gateff 1975** *{published data only}*
Gateff C, Dodin A, Wiart J. [A comparison of the serological effects of classical cholera vaccine and of purified fraction vaccine, with or without simultaneous yellow fever vaccine (author's transl)]. *Annales de microbiologie* 1975; Vol. 126, issue 2:231–46.
- Glass 1989** *{published data only}*
Glass RI, Svennerholm A-M, Stoll BJ, Khan MR, Huda S, Huq MI, Holmgren J. Effects of undernutrition on infection with vibrio cholerae 01 and on response to oral cholera vaccine. *Pediatric Infectious Disease Journal* 1989;**8** (2):105–9.
- Glenn 2007** *{published data only}*
Glenn GM, Villar CP, Flyer DC, Bourgeois AL, McKenzie R, Lavker RM, et al. Safety and immunogenicity of an enterotoxigenic Escherichia coli vaccine patch containing heat-labile toxin: use of skin pretreatment to disrupt the stratum corneum. *Infection and Immunity* 2007; Vol. 75, issue 5:2163–70.
- Graves 2000** *{published data only}*
Graves P, Deeks J, Demicheli V, Pratt M, Jefferson T. Vaccines for preventing cholera. *Cochrane Database of Systematic Reviews* 2001, Issue 1. [Art. No.: CD000974. DOI: 10.1002/14651858.CD000974]
- Gray 1989** *{published data only}*
Gray BN, Walker C, Andrewartha L, Freeman S, Bennett RC. Controlled clinical trial of adjuvant immunotherapy with BCG and neuraminidase-treated autologous tumour cells in large bowel cancer. *Journal of Surgical Oncology* 1989; Vol. 40, issue 1:34–7.
- Gupta 1998** *{published data only}*
Gupta RK, Taylor DN, Bryla DA, Robbins JB, Szu SC. Phase 1 evaluation of Vibrio cholerae O1, serotype Inaba, polysaccharide-cholera toxin conjugates in adult volunteers. *Infection and Immunity* 1998; Vol. 66, issue 7:3095–9.
- Hall 2001** *{published data only}*
Hall ER, Wierzbza TF, Ahren C, Rao MR, Bassily S, Francis W, et al. Induction of systemic antifimbria and antitoxin antibody responses in Egyptian children and adults by an oral, killed enterotoxigenic Escherichia coli plus cholera toxin B subunit vaccine. *Infection and Immunity* 2001; Vol. 69, issue 5:2853–7.
- Holmgren 1989** *{published data only}*
Holmgren J, Clemens J, Sack DA, Sanchez J, Svennerholm A-M. Oral immunization against cholera. *Current Topics in Microbiology and Immunology* 1989;**146**:197–204.
- Holmgren 1992** *{published data only}*
Holmgren J, Svennerholm AM, Jertborn M, Clemens J, Sack DA, Salenstedt R, et al. An oral B subunit: whole cell vaccine against cholera. *Vaccine* 1992;**10**(13):911–4.
- Hotomi 1998** *{published data only}*
Hotomi M, Saito T, Yamanaka N. Specific mucosal immunity and enhanced nasopharyngeal clearance of nontypeable Haemophilus influenzae after intranasal immunization with outer membrane protein P6 and cholera toxin. *Vaccine* 1998; Vol. 16, issue 20:1950–6.
- Islam 2008** *{published data only}*
Islam Z, Maskery B, Nyamete A, Horowitz MS, Yunus M, Whittington D. Private demand for cholera vaccines in rural Matlab, Bangladesh. *Health Policy* 2008; Vol. 85, issue 2:184–95.
- Jertborn 1984** *{published data only}*
Jertborn M, Svennerholm AM, Holmgren J. Gut mucosal, salivary and serum antitoxic and antibacterial antibody responses in Swedes after oral immunization with B subunit-whole cell cholera vaccine. *International Archives of Allergy and Applied Immunology* 1984; Vol. 75, issue 1:38–43.
- Jertborn 1986** *{published data only}*
Jertborn M, Svennerholm A-M, Holmgren J. Saliva, breast milk, and serum antibody responses as indirect measures of intestinal immunity after oral cholera vaccination or natural disease. *Journal of Clinical Microbiology* 1986;**24**(2):203–9.
- Jertborn 1988** *{published data only}*
Jertborn M, Svennerholm A-M, Holmgren J. Five-year immunologic memory in Swedish volunteers after oral cholera vaccination. *Journal of Infectious Diseases* 1988;**157** (2):374–7.
- Jertborn 1992** *{published data only}*
Jertborn M, Svennerholm A-M, Holmgren J. Safety and immunogenicity of an oral recombinant cholera b subunit-whole cell vaccine in Swedish volunteers. *Vaccine* 1992;**10** (2):130–2.
- Jertborn 1993** *{published data only}*
Jertborn M, Svennerholm AM, Holmgren J. Evaluation of different immunization schedules for oral cholera B subunit-whole cell vaccine in Swedish volunteers. *Vaccine* 1993; Vol. 11, issue 10:1007–12.
- Jertborn 1994** *{published data only}*
Jertborn M, Svennerholm AM, Holmgren J. Immunological memory after immunization with oral cholera B subunit-whole-cell vaccine in Swedish volunteers. *Vaccine* 1994; Vol. 12, issue 12:1078–82.
- Jertborn 1996** *{published data only}*
Jertborn M, Svennerholm A-M, Holmgren J. Intestinal and systemic immune responses in human after oral immunization with a bivalent b subunit-01/0139 whole cell cholera vaccine. *Vaccine* 1996;**14**(15):1459–65.
- Jertborn 1998** *{published data only}*
Jertborn M, Ahren C, Holmgren J, Svennerholm AM. Safety and immunogenicity of an oral inactivated enterotoxigenic Escherichia coli vaccine. *Vaccine* 1998; Vol. 16, issue 2–3: 255–60.
- Jertborn 2001** *{published data only}*
Jertborn M, Ahren C, Svennerholm AM. Dose-dependent circulating immunoglobulin A antibody-secreting cell and serum antibody responses in Swedish volunteers to an oral inactivated enterotoxigenic Escherichia coli vaccine. *Clinical and Diagnostic Laboratory Immunology* 2001;**8**(2): 424–8.

Johansson 2001 {published data only}

Johansson EL, Wassen L, Holmgren J, Jertborn M, Rudin A. Nasal and vaginal vaccinations have differential effects on antibody responses in vaginal and cervical secretions in humans. *Infection and Immunity* 2001;69(12):7481–6.

Johansson 2004 {published data only}

Johansson EL, Bergquist C, Edebo A, Johansson C, Svennerholm AM. Comparison of different routes of vaccination for eliciting antibody responses in the human stomach. *Vaccine* 2004; Vol. 22, issue 8:984–90.

Jones 2004 {published data only}

Jones T. Peru-1 5 (AVANT). *Current Opinion in Investigational Drugs* 2004; Vol. 5, issue 8:887–91.

Karlsen 2003 {published data only}

Karlsen TH, Sommerfelt H, Klomstad S, Andersen PK, Strand TA, Ulvik RJ, et al. Intestinal and systemic immune responses to an oral cholera toxoid B subunit whole-cell vaccine administered during zinc supplementation. *Infection and Immunity* 2003; Vol. 71, issue 7:3909–13.

Kenner 1995 {published data only}

Kenner JR, Coster TS, Taylor DN, Trofa AF, Barrera-Oro M, Hyman T, et al. Peru-15, an improved live attenuated oral vaccine candidate for *Vibrio cholerae* O1. *The Journal of Infectious Diseases* 1995; Vol. 172, issue 4:1126–9.

Kilhamn 1998 {published data only}

Kilhamn J, Jertborn M, Svennerholm AM. Kinetics of local and systemic immune responses to an oral cholera vaccine given alone or together with acetylcysteine. *Clinical and Diagnostic Laboratory Immunology* 1998; Vol. 5, issue 2: 247–50.

Kim 2008 {published data only}

Kim D, Canh do G, Poulos C, Thoa le TK, Cook J, Hoa NT, et al. Private demand for cholera vaccines in Hue, Vietnam. *Value Health* 2008; Vol. 11, issue 1:119–28.

Kirk 2005 {published data only}

Kirk MD, Kiedrzyński T, Johnson E, Elymore A, Wainiqolo I. Risk factors for cholera in Pohnpei during an outbreak in 2000: lessons for Pacific countries and territories. *Pacific Health Dialog* 2005; Vol. 12, issue 2:17–22.

Koenig 1998 {published data only}

Koenig MA, Roy NC, McElrath T, Shahidullah M, Wojtyniak B. Duration of protective immunity conferred by maternal tetanus toxoid immunization: Further evidence from Matlab, Bangladesh. *American Journal of Public Health* 1998; Vol. 88, issue 6:903–7.

Kollaritsch 1996 {published data only}

Kollaritsch H, Furer E, Herzog C, Wiedermann G, Que JU, Cryz SJ. Randomized, double-blind placebo-controlled trial to evaluate the safety and immunogenicity of combined *Salmonella typhi* Ty21a and *Vibrio cholerae* CVD 103-HgR live oral vaccines. *Infection and Immunity* 1996; Vol. 64, issue 4:1454–7.

Kollaritsch 1997 {published data only}

Kollaritsch H, Que JU, Kunz C, Wiedermann G, Herzog C, Cryz SJ. Safety and immunogenicity of live oral cholera

and typhoid vaccines administered alone or in combination with antimalarial drugs, oral polio vaccine, or yellow fever vaccine. *The Journal of Infectious Diseases* 1997; Vol. 175, issue 4:871–5.

Kozlowski 1999 {published data only}

Kozlowski PA, Cu-Uvin S, Neutra MR, Flanigan TP. Mucosal vaccination strategies for women. *The Journal of Infectious Diseases* 1999; Vol. 179 Suppl 3:S493–8.

Langevin-Perriat 1988 {published data only}

Langevin-Perriat A, Lafont S, Vincent C, Revillard JP, Mazert MC, Gerfaux G, et al. Intestinal secretory antibody response induced by an oral cholera vaccine in human volunteers. *Vaccine* 1988; Vol. 6, issue 6:509–12.

Lastre 2002 {published data only}

Lastre M, Del Campo J, Cedre B, Valmaseda T, Garcia L, Bracho G, Serrano T, Fando R, Sierra G, Perez O. Mucosal IgA anti-lipopolysaccharide antibodies induced by 638 oral live attenuated candidate vaccine. *Immunologia* 2002;21(1): 3–9.

Lelikov 1974 {published data only}

Lelikov VL, Sumarokov AA, Ivanov NR, Fedenev VG, Poliakov KA. [Study of the immunogenicity and immunological effectiveness of cholera anatoxin (data from a controlled epidemiological experiment). I. The characteristics of cholera anatoxin compared with corpuscular cholera vaccines administered by syringe and by using a jet injector]. *Zhurnal mikrobiologii, epidemiologii, i immunobiologii* 1974, issue 11:25–30.

Levine 1984 {published data only}

Levine MM, Black RE, Clements ML et al. Evaluation in humans of attenuated *Vibrio cholerae* El Tor Ogawa strain Texas Star-SR as a live oral vaccine. *Infection and Immunity* 1984; Vol. 43, issue 2:515–22.

Levine 1988a {published data only}

Levine MM, Kaper JB, Herrington D, Losonsky G, Morris JG, Clements ML, et al. Volunteer studies of deletion mutants of *Vibrio cholerae* O1 prepared by recombinant techniques. *Infection and Immunity* 1988; Vol. 56, issue 1: 161–7.

Levine 1988b {published data only}

Levine MM, Kaper JB, Herrington D, Ketley J, Losonsky G, Tacket CO, et al. Safety, immunogenicity, and efficacy of recombinant live oral cholera vaccines, CVD 103 and CVD 103-HgR. *Lancet* 1988; Vol. 2, issue 8609:467–70.

Lewis 1993 {published data only}

Lewis DJ, Castello-Branco LR, Novotny P, Dougan G, Poulton TA, Griffin GE. Circulating cellular immune response to oral immunization of humans with cholera toxin B-subunit. *Vaccine* 1993; Vol. 11, issue 2:119–21.

Leyten 2005 {published data only}

Leyten EM, Soonawala D, Schultsz C, Herzog C, Ligthelm RJ, Wijnands S, et al. Analysis of efficacy of CVD 103-HgR live oral cholera vaccine against all-cause travellers' diarrhoea in a randomised, double-blind, placebo-controlled study. *Vaccine* 2005;23(43):5120–6.

Lopez 2008 *{published data only}*

Lopez AL, Clemens JD, Deen J, Jodar L. Cholera vaccines for the developing world. *Human Vaccine* 2008; Vol. 4, issue 2:165–9.

Losonsky 1993 *{published data only}*

Losonsky GA, Tacket CO, Wasserman SS, Kaper JB, Levine MM. Secondary *Vibrio cholerae*-specific cellular antibody responses following wild-type homologous challenge in people vaccinated with CVD 103-HgR live oral cholera vaccine: Changes with time and lack of correlation with protection. *Infection and Immunity* 1993; Vol. 61, issue 2: 729–33.

Losonsky 1996 *{published data only}*

Losonsky GA, Yunyongying J, Lim V, Reymann M, Yu Leong Lim, Wasserman SS, et al. Factors influencing secondary vibriocidal immune responses: Relevance for understanding immunity to cholera. *Infection and Immunity* 1996; Vol. 64, issue 1:10–5.

Lucas 2005 *{published data only}*

Lucas ME, Deen JL, von Seidlein L, Wang XY, Ampuero J, Puri M, et al. Effectiveness of mass oral cholera vaccination in Beira, Mozambique. *New England Journal of Medicine* 2005; Vol. 352, issue 8:757–67.

Lucas 2007 *{published data only}*

Lucas ME, Jeuland M, Deen J, Lazaro N, MacMahon M, Nyamete A, et al. Private demand for cholera vaccines in Beira, Mozambique. *Vaccine* 2007; Vol. 25, issue 14: 2599–609.

Mahalanabis 2009 *{published data only}*

Mahalanabis D, Ramamurthy T, Nair GB, Ghosh A, Shaikh S, Sen B, et al. Randomized placebo controlled human volunteer trial of a live oral cholera vaccine VA1.3 for safety and immune response. *Vaccine* 2009; Vol. 27, issue 35: 4850–6.

Martell 2009 *{published data only}*

Martell BA, Orson FM, Poling J, Mitchell E, Rossen RD, Gardner T, et al. Cocaine vaccine for the treatment of cocaine dependence in methadone-maintained patients: a randomized, double-blind, placebo-controlled efficacy trial. *Arch Gen Psychiatry* 2009; Vol. 66, issue 10:1116–23.

María García 2005 *{published data only}*

María García H, Año G, Cedré B, Valmaseda T, Maestre JL, Díaz M, et al. [Selection of attenuated *Vibrio cholerae* strains to obtain oral attenuated candidate vaccines against cholera]. *Revista cubana de medicina tropical* 2005; Vol. 57, issue 2:92–104.

McCormack 1969 *{published data only}*

McCormack WM, Rahman AS, Chowdhury AK, Mosley WH, Phillips RA. Report of the 1966-67 cholera vaccine field trial in rural East Pakistan. 3. The lack of effect of prior vaccination or circulating vibriocidal antibody on the severity of clinical cholera. *Bulletin of the World Health Organization* 1969; Vol. 40, issue 2:199–204.

Migasena 1988 *{published data only}*

* Migasena S, Pitisuttitham P, Supanaranond W, Desakorn V, Prayurahong B, Suntharasamai P. Reactogenicity and

immunogenicity of oral cholera vaccine in Thai volunteers. *Southeast Asian Journal of Tropical Medicine and Public Health* 1988;19(3):423–8.

Migasena 1989b *{published data only}*

Migasena S, Desakorn V, Suntharasamai P, Pitisuttitham P, Prayurahong B, Supanaranond W, et al. Immunogenicity of two formulations of oral cholera vaccine comprised of killed whole vibrios and the B subunit of cholera toxin. *Infection and Immunity* 1989; Vol. 57, issue 1:117–20.

Migasena 1989c *{published data only}*

Migasena S, Pitisuttitham P, Suntharasamai P, Prayurahong B, Supanaranond W, Desakorn V, et al. Comparison of the reactivities and immunogenicities of procholerae and the B subunit of cholera toxin in Thai volunteers. *Infection and Immunity* 1989; Vol. 57, issue 7:1942–5.

Mitra 1990 *{published data only}*

Mitra AK, Rabbani GH. A double-blind, controlled trial of bioflorin (*Streptococcus faecium* SF68) in adults with acute diarrhea due to *Vibrio cholerae* and enterotoxigenic *Escherichia coli*. *Gastroenterology* 1990; Vol. 99, issue 4: 1149–52.

Mosley 1968 *{published data only}*

Mosley WH, Benenson AS, Barui R. A serological survey for cholera antibodies in rural east Pakistan. 2. A comparison of antibody titres in the immunized and control populations of a cholera-vaccine field-trial area and the relation of antibody titre to cholera case rate. *Bulletin of the World Health Organization* 1968; Vol. 38, issue 3:335–46.

Mosley 1969a *{published data only}*

Mosley WH, McCormack WM, Ahmed A, Chowdhury AK, Barui RK. Report of the 1966-67 cholera vaccine field trial in rural East Pakistan. 2. Results of the serological surveys in the study population--the relationship of case rate to antibody titre and an estimate of the inapparent infection rate with *Vibrio cholerae*. *Bulletin of the World Health Organization* 1969; Vol. 40, issue 2:187–97.

Mosley 1969b *{published data only}*

Mosley WH, McCormack WM, Fahimuddin M, Aziz KM, Rahman AS, Chowdhury AK, et al. Report of the 1966-67 cholera vaccine field trial in rural East Pakistan. I. Study design and results of the first year of observation. *Bulletin of the World Health Organization* 1969; Vol. 40, issue 2: 177–85.

Mosley 1970 *{published data only}*

Mosley WH, Woodward WE, Aziz KM, Rahman AS, Chowdhury AK, Ahmed A, et al. The 1968-1969 cholera-vaccine field trial in rural East Pakistan. Effectiveness of monovalent Ogawa and Inaba vaccines and a purified Inaba antigen, with comparative results of serological and animal protection tests. *The Journal of Infectious Diseases* 1970; Vol. 121:Suppl 121:1-9.

Mosley 1972 *{published data only}*

Mosley WH, Aziz KM, Mizanur Rahman AS, Alauddin Chowdhury AK, Ahmed A, Fahimuddin M. Report of the 1966-67 cholera vaccine trial in rural East Pakistan.

- Bulletin of the World Health Organization 1972; Vol. 47, issue 2:229–38.
- Mosley 1973** *{published data only}*
Mosley WH, Aziz KM, Rahman AS, Chowdhury AK, Ahmed A. Field trials of monovalent Ogawa and Inaba cholera vaccines in rural Bangladesh—three years of observation. *Bulletin of the World Health Organization* 1973; Vol. 49, issue 4:381–7.
- Nimbkar 1975** *{published data only}*
Nimbkar YS, Karbhari RS, Cherian S, Chanderkar NG, Bhamaria RP, Ranadive PS, et al. Antibody response to two cholera vaccines in volunteers. *Progress in Drug Research* 1975;19:544–62.
- Olsson 2006** *{published data only}*
Olsson L, Parment PA. Present and future cholera vaccines. *Expert Review of Vaccines* 2006; Vol. 5, issue 6:751–2.
- Oseasohn 1965** *{published data only}*
Oseasohn RO, Benenson AS, Fahimuddin M. Cholera vaccine field trial in rural East Pakistan (first year of observation). *Proceedings of the Cholera Research Symposium* 1965:362–6.
- Paineau 2008** *{published data only}*
Paineau D, Carcano D, Leyer G, Darquy S, Alyanakian MA, Simoneau G, et al. Effects of seven potential probiotic strains on specific immune responses in healthy adults: a double-blind, randomized, controlled trial. *FEMS Immunology and Medical Microbiology* 2008; Vol. 53, issue 1:107–13.
- Pal 1980** *{published data only}*
Pal SC, Deb BC, Sen Gupta PG, De SP, Sircar BK, Sen D, et al. A controlled field trial of an aluminum phosphate-adsorbed cholera vaccine in Calcutta. *Bulletin of the World Health Organization* 1980; Vol. 58, issue 5:741–5.
- Peltola 1977** *{published data only}*
Peltola H, Ruutu P, Palomäki H, Kaukinen K, Aho K. [Intracutaneous vaccination technic with less side effects against typhoid and cholera]. *Duodecim; lääketieteellinen aikakauskirja* 1977; Vol. 93, issue 7:431–8.
- Peltola 1989** *{published data only}*
Peltola H, Siitonen A, Kyrönseppä H, Simula I, Mattila L, Oksanen P, et al. Prevention of travellers' diarrhoea by oral b-subunit/whole-cell cholera vaccine. *Lancet* 1991;338:1285–9.
- Peltola 1991** *{published data only}*
Peltola H, Siitonen A, Kyrönseppä H, Simula I, Mattila L, Oksanen P, et al. Prevention of travellers' diarrhoea by oral B-subunit/whole-cell cholera vaccine. *Lancet* 1991; Vol. 338, issue 8778:1285–9.
- Philippines 1965** *{published data only}*
Philippines Cholera Committee. A controlled field trial of the effectiveness of cholera and cholera El Tor vaccines in the Philippines. Preliminary report. *Bulletin of the World Health Organization* 1965; Vol. 32, issue 5:603–25.
- Pitisuttithum 2001** *{published data only}*
Pitisuttithum P, Cohen MB, Phonrat B, Suthisarnsuntorn U, Bussaratid V, Desakorn V, et al. A human volunteer challenge model using frozen bacteria of the new epidemic serotype, V. cholerae O139 in Thai volunteers. *Vaccine* 2001; Vol. 20, issue 5–6:920–5.
- Qadri 2003** *{published data only}*
Qadri F, Ahmed T, Ahmed F, Bradley Sack R, Sack DA, Svennerholm AM. Safety and immunogenicity of an oral, inactivated enterotoxigenic *Escherichia coli* plus cholera toxin B subunit vaccine in Bangladeshi children 18–36 months of age. *Vaccine* 2003;2(21):19–20.
- Qadri 2004** *{published data only}*
Qadri F, Ahmed T, Wahed MA, Ahmed F, Bhuiyan NA, Rahman AS, et al. Suppressive effect of zinc on antibody response to cholera toxin in children given the killed, B subunit-whole cell, oral cholera vaccine. *Vaccine* 2004; Vol. 22, issue 4:416–21.
- Qadri 2006** *{published data only}*
Qadri F, Ahmed T, Ahmed F, Begum YA, Sack DA, Svennerholm AM. Reduced doses of oral killed enterotoxigenic *Escherichia coli* plus cholera toxin B subunit vaccine is safe and immunogenic in Bangladeshi infants 6–17 months of age: dosing studies in different age groups. *Vaccine* 2006; Vol. 24, issue 10:1726–33.
- Quiding-Jarbrink 2001** *{published data only}*
Quiding-Jarbrink M, Lonroth H, Ahlstedt I, Holmgren J, Svennerholm AM. Human gastric B cell responses can be induced by intestinal immunisation. *Gut* 2001; Vol. 49, issue 4:512–8.
- Rao 2002** *{published data only}*
Rao MR, Blackwelder WC, Troendle JF, Naficy AB, Clemens JD. Sample size determination for phase II studies of new vaccines. *Vaccine* 2002; Vol. 20, issue 27–28:3364–9.
- Rudin 1998** *{published data only}*
Rudin A, Johansson EL, Bergquist C, Holmgren J. Differential kinetics and distribution of antibodies in serum and nasal and vaginal secretions after nasal and oral vaccination of humans. *Infection and Immunity* 1998; Vol. 66, issue 7:3390–6.
- Rudin 1999** *{published data only}*
Rudin A, Riise GC, Holmgren J. Antibody responses in the lower respiratory tract and male urogenital tract in humans after nasal and oral vaccination with cholera toxin B subunit. *Infection and Immunity* 1999; Vol. 67, issue 6:2884–90.
- Sack 1991** *{published data only}*
Sack DA, Clemens JD, Huda S, Harris JR, Khan MR, Chakraborty J, et al. Antibody responses after immunization with killed oral cholera vaccines during the 1985 vaccine field trial in Bangladesh. *Journal of Infectious Diseases* 1991; 164(2):407–11.
- Sack 2007** *{published data only}*
Sack DA, Shimko J, Torres O, Bourgeois AL, Francia DS, Gustafsson B, et al. Randomised, double-blind, safety and efficacy of a killed oral vaccine for enterotoxigenic *E. coli* diarrhoea of travellers to Guatemala and Mexico. *Vaccine* 2007; Vol. 25, issue 22:4392–400.

Sanchez 1993b {published data only}

Sanchez JL, Trofa AF, Taylor DN, Kuschner RA, DeFraites RF, Craig SC, et al. Safety and immunogenicity of the oral, whole cell/recombinant b subunit cholera vaccine in North American volunteers. *Journal of Infectious Diseases* 1993;**167**(6):1446–9.

Sanchez 1994 {published data only}

Sanchez J, Begue R, Gaillour A, Cardenas V, et al. Feasibility of an efficacy trial of the whole cell plus recombinant B subunit (WC/rBS) oral cholera vaccine in Lima, Peru. [abstract]. *American Journal of Tropical Medicine and Hygiene* 1994; Vol. 51:216.

Saroso 1978 {published data only}

Saroso JS, Bahrawi W, Witjaksono H, Budiarso RL, Brotowasisto, Bencic Z, et al. A controlled field trial of plain and aluminium hydroxide-adsorbed cholera vaccines in Surabaya, Indonesia, during 1973–75. *Bulletin of the World Health Organization* 1978; Vol. 56, issue 4:619–27.

Savarino 1998 {published data only}

Savarino SJ, Brown FM, Hall E, Bassily S, Youssef F, Wierzbica T, Peruski L, El-Masry NA, Safwat M, Rao M, Jertborn M, Svennerholm A-M, Lee YJ, Clemens JD. Safety and immunogenicity of an oral, killed enterotoxigenic *E.coli*-cholera toxin B subunit vaccine in Egyptian adults.. *J Infect Dis* 1998;**177**:796–9.

Savarino 1999 {published data only}

Savarino SJ, Hall ER, Bassily S, Brown FM, Youssef F, Wierzbica TF, et al. Oral, inactivated, whole cell enterotoxigenic *Escherichia coli* plus cholera toxin B subunit vaccine: Results of the initial evaluation in children. *Journal of Infectious Diseases* 1999; Vol. 179, issue 1:107–14.

Savarino 2002 {published data only}

Savarino SJ, Hall ER, Bassily S, Wierzbica TF, Youssef FG, Peruski LF, et al. Introductory evaluation of an oral, killed whole cell vaccine in trial to avoid misclassification. *Pediatric Infectious Disease Journal* 2002;**21**(4):322–30.

Sommer 1973 {published data only}

Sommer A, Khan M, Mosley WH. Efficacy of vaccination of family contacts of cholera cases. *Lancet* 1973;**1**(814):1230–2.

SonLa 2007 {published data only}

SonLa Study Group. Using a fingerprint recognition system in a vaccine trial to avoid misclassification. *Bulletin of the World Health Organisation* 2007;**85**(1):64–7.

Stellfeld 2004 {published data only}

Stellfeld M. [Dukoral. Oral vaccine against cholera]. *Ugeskr Laeger* 2004; Vol. 166, issue 47:4251–3.

Sumarokov 1974 {published data only}

Sumarokov AA, Lelikov VL, Dzhaparidze MN, Karaeva LT, Derteva II. Study of the immunogenicity and immunological effectiveness of cholera toxin (data from a controlled epidemiological experiment). II. The characteristics of cholera toxin immunogenicity compared with corpuscular cholera vaccines administered by syringe and by using a jet injector [zuchenie reaktogennosti

i immunologicheskoy éffektivnosti kholerogena–anatoksin (materialy kontroliruemogo épidemiologicheskogo opyta). Soobshchenie II. Kharakteristika immunogennosti kholerogena–anatoksin v sravnenii s korpuskulyarnymi kholernymi vaktsinami pri ikh vvedenii shpritsom i s pomoshchyu. *Zhurnal mikrobiologii, epidemiologii, i immunobiologii* 1974, issue 11:31–7.

Sumarokov 1978 {published data only}

Sumarokov AA, Ivanov NR, Lelikov VL, Dzhaparidze MN, Karaeva LT, Derteva II, et al. Reactogenic properties and immunological efficacy of oral cholera chemical vaccine in a limited controlled revaccination trial in volunteers. *Zhurnal Mikrobiologii, Epidemiologii i Immunobiologii* 1978;**12**:87–92.

Sumarokov 1990 {published data only}

Sumarokov AA, Ivanov NR, Dzhaparidze MN, Reznikov IuB, Rystsova EA, Nikitina GP, et al. [The determination of the optimal inoculation dose of an oral cholera chemical bivalent vaccine in a controlled experiment]. *Zhurnal mikrobiologii, epidemiologii, i immunobiologii* 1990, issue 12:55–62.

Sumarokov 1991 {published data only}

Sumarokov AA, Ivanov NR, Dzhaparidze MN, Rystsova EA, Reznikov IuB, Matusevich LJa, et al. [The characteristics of the reactogenicity and immunological activity of a new cholera bivalent chemical vaccine based on the results of controlled trials]. *Zhurnal mikrobiologii, epidemiologii, i immunobiologii* 1991, issue 7:55–8.

Sumarokov 1993 {published data only}

Sumarokov AA, Dzhaparidze MN, Eliseev IuIu, Nikitina GP, Poliakov KA, Kazakova ES, et al. [The determination of the optimal inoculation dose of an oral cholera bivalent chemical vaccine in a controlled trial of the vaccination of children and adolescents]. *Zhurnal mikrobiologii, epidemiologii, i immunobiologii* 1993, issue 5:55–60.

Suntharasamai 1992 {published data only}

Suntharasamai P, Migasena S, Vongstongsri U, Supanaranond W, Pitituttitham P, Suppeeranan L, et al. Clinical and bacteriological studies of El Tor cholera after ingestion of known inocula in Thai volunteers. *Vaccine* 1992; Vol. 10, issue 8:502–5.

Svennerholm 1981 {published data only}

Svennerholm AM, Hanson LA, Holmgren J, Jalil F, Lindblad BS, Khan SR, et al. Antibody responses to live and killed poliovirus vaccines in the milk of Pakistani and Swedish women. *The Journal of Infectious Diseases* 1981; Vol. 143, issue 5:707–11.

Svennerholm 1984 {published data only}

Svennerholm AM, Gotheffors L, Sack DA, Bardhan PK, Holmgren J. Local and systemic antibody responses and immunological memory in humans after immunization with cholera B subunit by different routes. *Bulletin of the World Health Organization* 1984; Vol. 62, issue 6:909–18.

Tacket 1992 {published data only}

Tacket CO, Losonsky G, Nataro JP, Cryz SJ, Edelman R, Kaper JB, et al. Onset and duration of protective immunity

- in challenged volunteers after vaccination with live oral cholera vaccine CVD 103-HgR. *The Journal of Infectious Diseases* 1992; Vol. 166, issue 4:837–41.
- Tacket 1995a** *{published data only}*
Tacket CO, Losonsky G, Nataro JP, Wasserman SS, Cryz SJ, Edelman R, et al. Extension of the volunteer challenge model to study South American cholera in a population of volunteers predominantly with blood group antigen O. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1995; Vol. 89, issue 1:75–7.
- Tacket 1995b** *{published data only}*
Tacket CO, Losonsky G, Nataro JP, Comstock L, Michalski J, Edelman R, et al. Initial clinical studies of CVD 112 *Vibrio cholerae* O139 live oral vaccine: safety and efficacy against experimental challenge. *The Journal of Infectious Diseases* 1995; Vol. 172, issue 3:883–6.
- Tacket 1998** *{published data only}*
Tacket CO, Taylor RK, Losonsky G, Lim Y, Nataro JP, Kaper JB, et al. Investigation of the roles of toxin-coregulated pili and mannose-sensitive hemagglutinin pili in the pathogenesis of *Vibrio cholerae* O139 infection. *Infection and Immunity* 1998; Vol. 66, issue 2:692–5.
- Taylor 1994** *{published data only}*
Taylor DN, Killeen KP, Hack DC, Kenner JR, Coster TS, Beattie DT, et al. Development of a live, oral, attenuated vaccine against El Tor cholera. *The Journal of Infectious Diseases* 1994; Vol. 170, issue 6:1518–23.
- Taylor 1997** *{published data only}*
Taylor DN, Tacket CO, Losonsky G, Castro O, Gutierrez J, Meza R, et al. Evaluation of a bivalent (CVD 103-HgR/CVD 111) live oral cholera vaccine in adult volunteers from the United States and Peru. *Infection and Immunity* 1997; Vol. 65, issue 9:3852–6.
- Taylor 1999b** *{published data only}*
Taylor DN, Sanchez JL, Castro JM, Lebron C, Parrado CM, Johnson DE, et al. Expanded safety and immunogenicity of a bivalent, oral, attenuated cholera vaccine, CVD 103-HgR plus CVD 111, in United States military personnel stationed in Panama. *Infection and Immunity* 1999; Vol. 67, issue 4:2030–4.
- Thiem 2006** *{published data only}*
Thiem VD, Deen JL, von Seidlein L, Canh do D, Anh DD, Park JK, et al. Long-term effectiveness against cholera of oral killed whole-cell produced in Vietnam. *Vaccine* 2006; 24(20):4297–303.
- Von Seidlein 2007** *{published data only}*
Von Seidlein L, Vu DT, Dang DA, Do GC, Puri M, Gupta V, et al. Using a fingerprint recognition system in a vaccine trial to avoid misclassification. *Bulletin of the World Health Organization* 2007; Vol. 85, issue 1:64–7.
- Wassen 2005** *{published data only}*
Wassen L, Jertborn M. Kinetics of local and systemic immune responses after vaginal immunization with recombinant cholera toxin B subunit in humans. *Clinical and Diagnostic Laboratory Immunology* 2005; Vol. 12, issue 3:447–52.
- Wassen 2006** *{published data only}*
Wassen L, Jertborn M. Influence of exogenous reproductive hormones on specific antibody production in genital secretions after vaginal vaccination with recombinant cholera toxin B subunit in humans. *Clinical and Vaccine Immunology* 2006; Vol. 13, issue 2:202–7.
- Wasserman 1993** *{published data only}*
Wasserman SS, Kotloff KL, Losonsky GA, Levine MM. Immunologic response to oral cholera vaccination in a crossover study: a novel placebo effect. *American Journal of Epidemiology* 1993; Vol. 138, issue 11:988–93.
- Wassén 1996** *{published data only}*
Wassén L, Schön K, Holmgren J, Jertborn M, Lycke N. Local intravaginal vaccination of the female genital tract. *Scandinavian Journal of Immunology* 1996; Vol. 44, issue 4:408–14.
- Wiedermann 2000** *{published data only}*
Wiedermann G, Kollaritsch H, Kundi M, Svennerholm AM, Bjare U. Double-blind, randomized, placebo controlled pilot study evaluating efficacy and reactogenicity of an oral ETEC B-subunit-inactivated whole cell vaccine against travelers' diarrhea (preliminary report). *Journal of Travel Medicine* 2000; Vol. 7, issue 1:27–9.

Additional references

- Anh 2011**
Anh DD, Lopez AL, Thiem VD, Grahek SL, Duong TN, Park JK, et al. Use of Oral Cholera Vaccines in an Outbreak in Vietnam: A Case Control Study. *PLoS Neglected Tropical Diseases* 2011;5(1):e1006.
- Bhadra 1994**
Bhadra RK, Dasgupta U, Das J. Cholera vaccine: developmental strategies and problems. *Indian Journal of Biochemistry and Biophysics* 1994;31:441–48.
- Calain 2004**
Calain P, Chaine J-P, Johnson E, Hawley M-L, O'Leary MJ, Oshitani H, et al. Can oral cholera vaccination play a role in controlling a cholera outbreak?. *Vaccine* 2005;22(19):2444–51.
- Clemens 2001**
Clemens JD, Sack DA. Misleading Negative Findings in a Field Trial of Killed, Oral Cholera Vaccine in Peru (Comment). *Journal of Infectious Diseases* 2001;183:1306–8.
- Girard 2005**
Girard MP, Steele D, Chaignat C-L, Kiény MP. A review of vaccine research and development: human enteric infections. *Vaccine* 2005;24(15):2732–50.
- Graves 2001**
Graves P, Deeks J, Demicheli V, Pratt M, Jefferson T. Vaccines for preventing cholera. *Cochrane Database of Systematic Reviews* 2001, Issue 1. [Art. No.: CD000974. DOI: 10.1002/14651858.CD000974]

Graves 2010

Graves P, Deeks J, Demicheli V, Jefferson T. Vaccines for preventing cholera: killed whole cell or other subunit vaccines (injected). *Cochrane Database of Systematic Reviews* 2010, Issue 1. DOI: 10.1002/14651858.CD000974.

Harris 2005

Harris JB, Khan AI, LaRocque RC, Dorer DJ, Chowdhury F, Faruque ASG, et al. Blood Group, Immunity, and Risk of Infection with *Vibrio cholerae* in an Area of Endemicity. *Infection and Immunity* 2005;**73**(11):7422–7427.

Heymann 2008

Heyman DL. *Control of Communicable Diseases Manual*. 19th Edition. Washington: American Public Health Association, 2008.

Higgins 2008

Higgins JPT, Altman DG (editors). Chapter 8: Assessing risk of bias in included studies.. *Cochrane Handbook of Systematic Reviews of Intervention*. Version 5.0.0. The Cochrane Collaboration, 2008.

Hill 2006

Hill DR, Ford R, Lallo DG. Oral cholera vaccines: use in clinical practice. *Lancet infectious diseases* 2006;**6**:361–573.

Holmgren 2005

Holmgren J, Adamson J, Anjuere F, Clemens J, Czerkinsky C, Flack C-F, et al. Mucosal adjuvants and anti-infection and anti-immunopathology vaccines based on cholera toxin, cholera toxin B subunit and CpG DNA. *Immunology letters* 2005;**97**(2):181–188.

Huilan 1991

Huilan S, Zhen LG, Mathan MM, Mathew MM, Olarte J, Espejo R, et al. Etiology of acute diarrhoea among children in developing countries: a multicentre study in five countries. *Bulletin of the World Health Organization* 1991; **69**:549–55.

Lucas 2005

Lucas ME, Deen JL, von Seidlein L, Wang XY, Ampuero J, Puri M, et al. Effectiveness of mass cholera vaccination in Beira, Mozambique. *New England Journal of Medicine* 2005;**352**(8):757–67.

Review Manager 5 [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.0. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2008.

Reyburn 2011

Reyburn R, Deen JL, Grais RF, Bhattacharya S, Sur D, Lopez AL, et al. The case for reactive mass oral cholera vaccinations. *PLoS Neglected Tropical Diseases* 2011;**5**(1): e952.

Ryan 2011

Ryan ET. The cholera pandemic, still with us after half a century: time to rethink. *PLoS Neglected Tropical Diseases* 2011;**5**(1):e1003.

Sack 2004

Sack DA, Sack RB, Nair GB, Siddique AK. Cholera. *Lancet* 2004;**363**:223–33.

Sanchez 1997

Sanchez J, Taylor DN. Cholera. *Lancet* 1997;**349**:1825–30.

Taylor 2001

Taylor DN, Sanchez J, Cardenas V, Gilman RE, Sadoff J. Misleading negative findings in a field trial of killed, oral cholera vaccine in Peru (Authors reply). *Journal of Infectious Diseases* 2001;**183**:1308–9.

WHO 2000a

World Health Organization. Fact Sheet No. 103: Cholera. <http://www.who.int/mediacentre/factsheets/fs107/en/> Revised March 2000.

WHO 2000b

World Health Organization Department of Communicable Disease Surveillance and Response. WHO report on global surveillance of epidemic-prone infectious diseases. <http://www.who.int/csr/resources/publications/surveillance/en/cholera.pdf> 2000; Vol. Chapter 4:39–43.

WHO 2006a

World Health Organization. Cholera 2005. *Weekly epidemiological record* 2006;**81**:297–308.

WHO 2006b

World Health Organization Global Task Force on Cholera Control. Oral cholera vaccine use in complex emergencies: what next?. Report of a WHO meeting, Cairo, Egypt 14–16 December 2005; Vol. WHO/CDS/NTD/IDM/2006.2.

WHO 2006c

World Health Organization. The use of two-dose oral vaccine in the context of major natural disaster: report of a mass vaccination campaign in Aceh Province, Indonesia 2005. http://www.who.int/topics/cholera/publications/final_tsunami.pdf 2006.

WHO 2009

WHO. Outbreak News: Cholera, Zimbabwe - update. *Weekly Epidemiological Record* 2009;**84**:109–10.

WHO 2010a

WHO. Outbreak news: Cholera, Haiti - update. *Weekly Epidemiological Record* 2010;**85**:489–90.

WHO 2010b

WHO. Cholera vaccines:WHO position paper. *Weekly Epidemiological Record* 2010;**85**:117–28.

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Anh 2007

Methods	Design: A randomized controlled trial (individually randomized) Trial dates and duration: Enrollment from May to June 2005; follow-up for 28 days	
Participants	Sample size: 153 participants enrolled Inclusion criteria: Age 18 to 40 years, healthy male and non-pregnant females, written informed consent Exclusion criteria: History of diarrhoea, anti-diarrhoeal or antibiotic use during the past week, history of diarrhoea and abdominal pain lasting for 2 weeks during the past 6 months	
Interventions	Vaccine: Bivalent killed whole-cell vaccine (BivWC; mORCVAX, VABIOTECH) Placebo: Heat-killed <i>E. coli</i> K12 strain All participants were randomized to receive 2 doses, at an interval of 14 days	
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none">Serious adverse events during 28 days follow-upAdverse events within 3 days of each dose <i>Not included in the review:</i> <ul style="list-style-type: none">Immunological outcomes: Geometric mean-fold rise in serum vibriocidal antibody titres and proportion who develop ≥ 4-fold rises from baseline after one or two dose	
Notes	Location: SonLa Province, Northwest Vietnam Setting: Source of funding: The Bill and Melinda Gates Foundation through the Diseases of Most Impoverished Program administered by the International Vaccine Institute, and the Swedish International Development Cooperation Agency	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: 'A randomization list was prepared by a statistician who otherwise was not involved in the study. Randomization numbers were generated in blocks of four'
Allocation concealment (selection bias)	Low risk	Comment: See other comments, no further description.
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported

Anh 2007 (Continued)

Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'The reformulated vaccine and the placebo were packaged as liquid formulations in identical vials containing five 1.5-ml doses'. 'A physician who was unaware of the study agent received by the subject conducted a structured interview regarding the subjects...symptoms'
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: Nine participants (5.9%) did not receive the second dose of vaccine; 5 were found ineligible and 4 lost to follow-up
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Begue 1995

Methods	Design: Randomized controlled trial (individual randomization) Duration and dates (field work): March 1993
Participants	Sample size: 624 received the first dose of vaccine, 541 received 2 doses Inclusion criteria: Persons aged 2 to 65 years Exclusion criteria: pregnancy
Interventions	Vaccine: Killed whole-cell vaccine plus recombinant cholera toxin subunit (WC-rBS; Dukoral®, SBL, Sweden) Placebo: Inactivated <i>E. coli</i> K12 suspension Vaccine and placebo were administered along with freshly prepared antacid solution. Two doses were given two weeks apart
Outcomes	<i>Included in the review:</i> <ul style="list-style-type: none"> Adverse events after the first dose: participants were observed for one hour and then asked about symptoms at time of the second dose <i>Not included in the review:</i> <ul style="list-style-type: none"> Immunological outcomes: Geometric mean vibriocidal antibody, IgG antitoxin and IgA antitoxin titres pre and post vaccination. Proportion who developed ≥ 2 or ≥ 4 fold increases.
Notes	Location: outskirts of Lima, Peru Setting: Small community of 300 families. Source of funding: US Naval Medical Research and Development Command

Begue 1995 (Continued)

<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "...administer the vaccine or placebo according to a pre-randomized list" Comment: Unclear description but probably low risk of bias
Allocation concealment (selection bias)	Low risk	Comment: Not described but probably low risk of bias
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'Inactivated Escherichia coli K12, identical in appearance to the vaccine, was used as placebo, and was administered orally in the above antacid solution, and in a double blinded manner.'
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported as an outcome
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No loss to follow-up
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Benítez 1999

Methods	Design: Randomized controlled trial (individually randomized) Duration and dates (field work): Not stated
Participants	Sample size: 56 (this paper describes 4 separate small trials with different doses of VC638. A total of 42 received vaccine and 14 placebo) Inclusion criteria: Age 18 to 40 years, male students or workers, good health, informed consent Exclusion criteria: Recent history of diarrhoeal disease or cholera vaccination, taking medication at the time of recruitment
Interventions	Vaccine: VC638 - A live attenuated strain of V. cholerae O1 El Tor Ogawa <ul style="list-style-type: none"> • 2 x 10⁹ CFU • 1 x 10⁹ CFU • 2 x 10⁸ CFU

	<ul style="list-style-type: none">● 4 x 10^{7CFU} Placebo: Buffer alone	
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none">● Adverse events (detected through inpatient observation) <i>Not included in the review:</i> <ul style="list-style-type: none">● Immunological outcomes: Serum vibriocidal geometric mean antibody titres on days 0, 14 and proportion who develop ≥2 or ≥4-fold rises from baseline after one dose	
Notes	Location: La Lisa of Havana, Cuba Setting: Institute of Tropical Medicine Source of funding: None stated	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: Not described as randomised though it seems unlikely that this was not done
Allocation concealment (selection bias)	Low risk	Quote: 'The clinical investigator assigned a letter to each volunteer. The code was kept by the monitor till the end of the experiment and analysis of all samples'
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'The placebo consisted of bicarbonate buffer alone and was indistinguishable from the vaccine preparation. To ensure double-blinding, identical flasks, containing either inoculum or placebo, were coded by an outside monitor'
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No losses recorded during the monitoring of adverse events
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of any other bias

Chen 1996

Methods	Design: Randomized controlled trial (individually randomized) Duration and dates (field work): Jun 1993 to Jan 1994
Participants	Sample size: 369 Inclusion criteria: Students from the primary and secondary school, and factory workers of Jiu-Fu area, Guang-Zhou Exclusion criteria: A history of cholera, or acute diarrhoea in the past 2 weeks
Interventions	Vaccine 1: Killed whole-cell vaccine plus recombinant cholera toxin B subunit (locally formulated) <ul style="list-style-type: none"> • 1x10¹⁰ vibrio cholera whole cells + 5mg rBS Vaccine 1: Killed whole-cell vaccine plus recombinant cholera toxin B subunit (locally formulated) <ul style="list-style-type: none"> • 1x10¹⁰ vibrio cholera whole cells + 1mg rBS Placebo: Buffer alone
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none"> • Adverse events (detected through observation) <i>Not included in the review:</i> <ul style="list-style-type: none"> • Immunological outcomes
Notes	Location: Jiu-Fu Area in Guang-Zhou city Setting: Source of funding: National 638 funds and fund from the Academy of Guang-Dong Province

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Labelled as 'Randomized', no further details.
Allocation concealment (selection bias)	Unclear risk	None described.
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy is not reported
Blinding (performance bias and detection bias) Safety outcomes	Unclear risk	Described as 'double-blind'. No further details
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy is not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	No losses to follow-up reported

Chen 1996 (Continued)

Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Clemens 1987

Methods	Design: Randomized controlled trial (individual randomization) Trial dates and duration: 1984, with short-term follow-up	
Participants	Sample size: 1,257 enrolled and took first dose of vaccine or placebo, 1051 received two doses, and 898 received third doses Inclusion criteria: Children aged 2 to 15 years and women aged over 15 years Exclusion criteria: Pregnancy, people too ill to leave their beds on the day of the vaccination	
Interventions	Vaccine 1: Killed whole cell plus purified cholera B subunit vaccine (WC-BS) Vaccine 2: Killed whole cell vaccine (WC) Placebo 1: Heat-inactivated <i>E. coli</i> K12 strain Placebo 2: Distilled water	
Outcomes	<i>Included in the review:</i> <ul style="list-style-type: none">Adverse events for three consecutive days after each dose <i>Not included in the review:</i> <ul style="list-style-type: none">Immunogenicity	
Notes	<i>Location:</i> Matlab, Bangladesh Setting: Community, within a health and demographic surveillance site Source of funding: United States Agency for International Development (USAID); the government of Japan; the Swedish Agency for Research Cooperation with Developing Countries, and the World Health Organization	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "randomly assigned" Comment: Unclear description but probably low risk of bias
Allocation concealment (selection bias)	Low risk	Quote: "Each of the agents, labelled only as W,X,Y or Z" Comment: Allocation concealed
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported

Clemens 1987 (Continued)

Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: “..study physicians who were kept unaware of the identities of agents received by subjects..”
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No loss to follow-up
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Clemens 1988 Bangladesh

Methods	Design: Randomized controlled trial (individual randomization) Trial dates and duration: Vaccination January to March 1985; follow-up 5 years Surveillance: Passive surveillance system at diarrhoea treatment centres serving the study population
Participants	Number of participants: 89,596 received at least one dose of vaccine or placebo, 62,285 ingested three complete doses Inclusion criteria: children aged 2-15 years and women over the age of 15 Exclusion criteria: Pregnancy, illness requiring bed rest
Interventions	Vaccine 1: Killed whole cell plus purified cholera B subunit vaccine (WC-BS) Vaccine 2: Killed whole cell vaccine (WC) Placebo: <i>Escherichia coli</i> K12 strain placebo (K12) All subjects were randomized to receive three doses, at 6 week intervals. All doses were ingested with antacid
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none"> Cholera infection (faecal excretion of <i>V. Cholerae</i> 01) Symptomatic cholera infection (faecal excretion of <i>V. Cholerae</i> 01 from 48 hours before to 48 hours after a diarrhoea episode) <ul style="list-style-type: none"> Cases of cholera (non-bloody diarrhoea, dehydration and excretion of <i>V. cholerae</i> 01). Cases of cholera, excluding cases that are clinically atypical or associated with mixed infections. Symptomatic and asymptomatic cholera infection detected using active surveillance of among persons residing in the same courtyard as a sentinel cases detected in active surveillance. Participants were surveyed for symptoms and rectal swabs taken and cultured for <i>V. cholerae</i> 01 each day for 7 days. Cases of diarrhoea, classified according to watery and non-watery, and severe and non-severe. Deaths from cholera.

	<ul style="list-style-type: none">• All deaths.• Adverse events within 3 days of first dose and within 3 days of second dose. Cases of diarrhoea and cholera were only included in the analysis if they occurred at least 14 days after the third dose of vaccine or placebo <i>Not included in the review:</i> <ul style="list-style-type: none">• Immunological response in participants with cholera, comparing those receiving placebo and placebo.• Diarrhoeal episodes associated with other <i>Vibrio</i> and <i>Aeromonas</i> species.• Antibacterial and anti-toxic antibody responses in breast milk.• Antibody responses following immunisation.• Cases of cholera by neighbourhood vaccine coverage level (herd immunity).• Diarrhoea associated with ETEC	
Notes	Location: Matlab, Bangladesh Setting: Surveillance study area, served by three diarrhoea treatment centres Source of funding: Bill and Melinda Gates Foundation; U.S. National Institutes of Health; U.S. National Science Foundation; Swedish International Development Cooperation Agency; governments of Korea, Japan and Kuwait, USAID, Word Health Organization	
<i>Risk of bias</i>		
Bias	Authors’ judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote:“After computerisation of the census, we assigned every person in the eligible age-gender categories to letters A, B or C, using simple randomisation”
Allocation concealment (selection bias)	Low risk	Quote: “The agents were identified only by the letters A, B and C” Comment: Allocation concealed
Blinding (performance bias and detection bias) Efficacy outcomes	Low risk	Quote: “During the conduct of the study, the identities of these letter...were unknown to all persons connected with the trial in Bangladesh”
Blinding (performance bias and detection bias) Safety outcomes	Low risk	As above
Incomplete outcome data (attrition bias) Efficacy outcomes	Low risk	Comment: Attrition between the first and third doses was high: 30.5%. The protective effect is reported as being similar in those who only received two doses, so these losses are unlikely to have introduced significant bias

Clemens 1988 Bangladesh (Continued)

Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: There was no missing data for adverse events.
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting.
Other bias	Low risk	No evidence of other bias.

Cohen 2002

Methods	Design: Randomized controlled trial (individual randomization) Trial dates and duration: 2000-2001
Participants	Number of participants: 59 (36 included in the challenge study) Inclusion criteria: Age 18 to 40 years, informed consent, and judged likely to comply with the study requirements Exclusion criteria: Clinically significant abnormalities on urinalysis, full blood count, serum hepatic transaminases, glucose, creatine, urea nitrogen, electrolytes or ECG. Travel to cholera endemic areas in the previous 5 years, history of cholera or ETEC challenge, recent antibiotic use, abnormal stool pattern, regular laxative use, failure to pass psychological screening, allergy to tetracycline or ciprofloxacin, pregnant or breastfeeding, HIV-positive, hep B-positive, hep C-positive, stool culture positive for enteric pathogen.
Interventions	Vaccine: Peru 15 - a live attenuated strain of <i>V. cholerae</i> O1 El Tor Inaba plus 200 ml CeraVacx buffer (Cera Products, Columbia) Placebo: 200ml CeraVacx buffer (Cera Products, Columbia) Challenge: Three months after vaccination, willing participants were given artificial challenge with 10 ⁵ CFU of virulent <i>V. cholerae</i> O1 El Tor Inaba Strain N16961, prepared from a standardised frozen inoculum
Outcomes	<i>Included in the review:</i> <ul style="list-style-type: none"> Adverse events during the first 3 days after the dose (assessed by a self completed diary) Participants who went on to receive artificial challenge were also monitored for diarrhoea, and positive stool culture with the challenge strain; on an inpatient basis <ul style="list-style-type: none"> Any diarrhoea: passage of two or more unformed stools over a 48 hour period that equalled or exceeded 200 g for a single stool, or 300 g or greater in total Moderate or severe cholera: diarrhoea with passage of >3,000 g during the study period plus a positive stool culture for <i>V. cholerae</i> O1 <i>Not included in the review:</i> <ul style="list-style-type: none"> Immunological outcomes: Geometric mean inverse Inaba vibriocidal antibody titres pre and post immunisation and proportion who developed ≥ 4-fold rises from baseline after one dose
Notes	Location: USA Setting: Volunteer study. Outpatient phase for adverse events, inpatients phase for response to artificial challenge Sources of funding: National Institutes of Health, General Clinical Research Centres Program

Cohen 2002 (Continued)

<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: 'sequence generated by SAS PROC PLAN'.
Allocation concealment (selection bias)	Low risk	Comment: The randomization code generated off-site and study blinded until after analysis
Blinding (performance bias and detection bias) Efficacy outcomes	Low risk	Quote: 'Investigators did not know the vaccine status of all volunteers until the data was locked and the code was broken after the challenge was completed'
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'Volunteers were randomly assigned to groups in a double-blind manner. ...A study nurse who was unaware of the group assignment reviewed the (symptom) diary'
Incomplete outcome data (attrition bias) Efficacy outcomes	High risk	Comment: The loss of participants between randomisation and the challenge study (39%) could introduce significant bias
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: The data was complete for the three days of adverse event monitoring
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Concha 1995

Methods	Design: Randomized controlled trial (cluster randomized by household) Duration: Two months, January and February 1992
Participants	Sample size: 1313 received an initial dose of vaccine or placebo, 1165 received two doses Inclusion criteria: People between the ages of 12 months and 64 years who have resided in the study area for at least two months Exclusion criteria: Confirmed or possible pregnancy, illness requiring bed rest, known mental illness or incapacity to give informed consent, diarrhoea at the time either of the two vaccine doses were administered

Interventions	Vaccine: Killed whole-cell vaccine plus recombinant cholera toxin subunit (WC-rBS; Dukoral®, SBL, Sweden) Placebo: killed whole cells of <i>E. coli</i> K12. Both vaccine and placebo were administered with a buffer solution. Two doses were given, two weeks apart	
Outcomes	<i>Included in the review:</i> <ul style="list-style-type: none">Reported symptoms in the three days following ingestion of the vaccine (daily visits using pre-coded forms) <i>Not included in the review:</i> <ul style="list-style-type: none">Immunological outcomes: Geometric mean vibriocidal antibody, IgG antitoxin and IgA antitoxin titres pre and post vaccination.	
Notes	Location: Los Olivios, Barraquilla, Colombia Setting: Households in a poor neighbourhood Source of funding: Pan American Health Organization and World Health Organization. Vaccine donated by the National Bacteriological Laboratory in Stockholm, Sweden	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Households were randomly selected to receive either vaccine or placebo" Comment: The method of sequence generation is unclear
Allocation concealment (selection bias)	Low risk	Quote: "the vaccination team knew the two only as 'vaccine A and 'vaccine B'
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy is not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: "Both agents were administered double-blind; the vaccination team know the two only as "vaccine A" and "vaccine B"....nurses, who .. were unaware of how the agent were distributed..record any. .symptoms"
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No loss to follow-up
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting

Concha 1995 (Continued)

Other bias	High risk	Comment: 620 individuals who originally consented to participate dropped out of the study because of a political campaign against it
------------	-----------	--

Cryz 1990

Methods	Design: Randomized controlled trial (individually randomized) Trial dates and duration: Study dates not given; follow-up 21 days
Participants	Sample size: 50 enrolled Inclusion criteria: Age 21 to 45 years, healthy, informed consent Exclusion criteria: None stated
Interventions	Vaccine: CVD 103-HgR live attenuated vaccine containing: <ul style="list-style-type: none"> • 5 x 10⁸ CFU of lyophilized genetically modified <i>V. cholerae</i> O1 Classical Inaba (569B) Placebo: 5 x 10 ⁸ CFU of heat-killed <i>E. coli</i> K12 strain
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none"> • Adverse events during first 7 days after vaccination (interview on day 7) only diarrhoea and abdominal pain are reported. <i>Not included in the review:</i> <ul style="list-style-type: none"> • Immunological outcome: Geometric mean serum vibriocidal antibody titres on day 0, 10 and 21, proportion who develop ≥ 4 fold rises in serum titres.
Notes	Location: Switzerland Setting: Not stated Source of funding: None stated

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Described as 'randomised', no further details given.
Allocation concealment (selection bias)	Unclear risk	None described
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'A coded sachet containing either vaccine or placebo was mixed with the buffer solution and immediately ingested. 'The appearance of the placebo was iden-

Cryz 1990 (Continued)

		tical to that of the vaccine.'
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No losses are reported during the first week of adverse event surveillance
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

García 2005

Methods	Design: Randomized controlled study with artificial challenge (Individually randomized) Trial dates and duration: Dates not stated; artificial challenge took place 1 month after vaccination	
Participants	Sample size: 45 (21 in challenge study) Inclusion criteria: Age 18 to 40 years, volunteer male workers among the western scientific community, good health, informed consent Exclusion criteria: Recent history of diarrhoeal disease or cholera vaccination, taking any medication at the time of recruitment, any abnormality in clinical laboratory tests (complete blood count, chemistry panel, HIV and Hep C virus antibodies, Hep B virus antigen), stool cultures positive for an enteric pathogen, recent antibiotic use, or psychological incompatibility with accepting quarantine conditions	
Interventions	Vaccine: VC638 - A live attenuated strain of <i>V. Cholerae</i> O1 El Tor Ogawa <ul style="list-style-type: none">1 x 10⁹ CFU plus buffer Placebo: Buffer alone Artificial challenge: 7 x 10 ⁵ CFU of fully virulent El Tor Ogawa strain 3008 (orally).	
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none"><i>V. cholerae</i> diarrhoea following oral challengeAdverse events (inpatient monitoring for 5 days) <i>Not included in the review:</i> <ul style="list-style-type: none">Faecal virus sheddingGeometric mean vibriocidal antibody titres pre and post immunisation, LPS specific IgA, and proportion who developed ≥2-fold rises from baseline after one dose	
Notes	Location: Havana Cuba Setting: Inpatient trials unit, Institute of Tropical Medicine Source of funding: None stated	
Risk of bias		
Bias	Authors' judgement	Support for judgement

García 2005 (Continued)

Random sequence generation (selection bias)	Unclear risk	Comment: Described as 'Randomized', no further details given
Allocation concealment (selection bias)	Unclear risk	Comment: None described.
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Comment: Described as 'double blind', no further details given
Blinding (performance bias and detection bias) Safety outcomes	Unclear risk	Comment: Described as 'double blind', no further details given
Incomplete outcome data (attrition bias) Efficacy outcomes	High risk	Comment: The loss of participants between randomisation and the challenge study (47%) could introduce significant bias
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: The data was complete as participants for the three days of adverse event monitoring
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Gotuzzo 1993

Methods	Design: A randomized controlled trial (individually randomized) Duration: Vaccination from Sept to Dec 1991; follow-up 28 days
Participants	Sample size: 241 enrolled Inclusion criteria: Adults aged 18 to 38 years Exclusion criteria: Pregnancy, antibiotics or diarrhoea within the previous 72 h, previous cholera vaccine
Interventions	Vaccine 1: CVD 103-HgR live attenuated vaccine containing: <ul style="list-style-type: none"> • 5 x 10⁹ lyophilized organisms of a genetically modified <i>V. cholerae</i> O1 Vaccine 2: CVD 103-HgR live attenuated vaccine containing: <ul style="list-style-type: none"> • 5 x 10⁸ lyophilized organisms of a genetically modified <i>V. cholerae</i> O1 Placebo: 5 x 10 ⁸ cells of heat-killed <i>E. coli</i> K-12 strain
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none"> • Adverse events during the first 7 days <i>Not included in the review:</i> <ul style="list-style-type: none"> • Immunological outcomes: Geometric mean rise in vibriocidal antibody titres, and proportion who develop ≥4 fold rises in serum titres from baseline after one or two doses

Gotuzzo 1993 (Continued)

Notes	<p>Location: Peru</p> <p>Setting: 2 groups: high socioeconomic group: medical students and physicians from the Facultad de Medicina, Universidad Peruana Cayetano Heredia, and a low socioeconomic group: selected from Canto Grande, a periurban slum community with poor water and sanitation</p> <p>Source of funding: National Institute of Allergy and Infectious Diseases (NIAID)</p>
-------	---

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: Described as 'randomized', no further details given
Allocation concealment (selection bias)	Low risk	Quote: 'Eligible adults were administered coded preparations sequentially labelled A, B, or C, two of which contained the vaccine and the other a placebo'
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'Double-blind clinical follow-up was maintained for 7 days following vaccination'
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No losses to follow up are reported
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Hallander 2002

Methods	<p>Design: Randomized controlled trial (individually randomized)</p> <p>Duration: Enrolled; follow-up 28 days</p>
Participants	<p>Sample size: 249</p> <p>Inclusion criteria: Age 1 to 12 years, permanent resident in study area, informed consent, good health</p> <p>Exclusion criteria: None stated</p>

Interventions	Vaccine Killed whole-cell vaccine plus recombinant cholera toxin subunit (WC-rBS; Dukoral®, SBL, Sweden) Placebo: heat-killed <i>E. coli</i> K-12 strain (C 600) All participants were randomized to receive 2 doses, 14 days apart
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none"> • Serious adverse events • Adverse events during the first 3 days after each dose (parental interview and diary cards for 3 days) <i>Not included in the review:</i> <ul style="list-style-type: none"> • Immunological outcomes: Geometric mean rise in vibriocidal antibody titres, and proportion who develop ≥ 2 fold rises in serum titres from baseline after one or two doses
Notes	Location: León, Nicaragua Setting: Source of funding: None declared *This paper contained the details of three individual trials: OCV-023, OCV-024 and OCV-028. OCV-023 and OCV-024 were excluded from this review as they used a variation on this vaccine for which no primary efficacy data is available

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: Described as 'randomized', no further details given
Allocation concealment (selection bias)	Unclear risk	Comment: None described.
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Unclear risk	Comment: None described.
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No losses described
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Methods	Design: A randomized controlled trial (individually randomized) Trial dates and duration: Enrollment from June to August 2007; follow-up for 28 days	
Participants	Sample size: 160 patients stratified into adults and children Inclusion criteria: Age 18 to 40 years for adult study, 1 to 18 years for children, healthy male and non-pregnant females, written informed consent Exclusion criteria: Abdominal pain, loss of appetite, nausea, general ill feeling or vomiting within the past 24 h, any diarrhoea within 6 weeks of enrolment, diarrhoea or abdominal pain lasting more than 2 weeks in the past 6 months, antibiotics in the past 2 weeks, anti-diarrhoeal medication or acute disease in the past week, history of serious chronic disease or an immunocompromising condition or therapy	
Interventions	Vaccine: Bivalent killed whole-cell vaccine (BivWC: SHanchol®, Shantha Biotechnics) Placebo: Heat killed <i>E. coli</i> K12 strain All subjects were randomized to receive 2 doses, at an interval of 14 days. All doses were administered via an oral syringe and offered water	
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none">• Serious adverse events during 28 days follow-up• Adverse events within 3 days of each dose <i>Not included in the review:</i> <ul style="list-style-type: none">• Immunological outcomes: Geometric mean-fold rise in serum vibriocidal antibody titres and proportion who develop ≥ 4-fold rises from baseline after one or two doses	
Notes	Location: Kolkata, India Setting: The clinical trial unit of the National Institute of Cholera and Enteric Diseases (NICED) Source of funding: the Bill and Melinda Gates Foundation through the Diseases of the Most Impoverished Program and the Cholera Vaccine Initiative, and the governments of Korea, Kuwait and Sweden through the International Vaccine Institute	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: 'Separate randomization lists for the two age groups were prepared by a statistician in the IVI who was otherwise not involved in the study. Randomization was performed in blocks of four using Visual Fortran 5.0 (Digital USA)'
Allocation concealment (selection bias)	Low risk	Quote: 'Vials were labeled with four-letter codes... the identities of the codes were only known to Shantha staff who labeled the vials and who were otherwise not involved in the study', 'Eligible subjects were assigned to receive either vaccine or placebo

Kanungo 2009 (Continued)

		according to the randomization list. Subjects were assigned sequentially to a number in the randomization list'
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'Study staff and participants were unaware of the identity of the codes during the study period'
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: Five participants (3.1%) were lost to follow-up between the first and second doses. Four withdrew consent and one was found to be ineligible
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Kotloff 1992

Methods	Design: Randomized controlled cross-over trial (individually randomized) Duration: Study dates not given; follow-up 28 days
Participants	Sample size: 94 enrolled Inclusion criteria: Age 18 to 40 years, college students, informed consent Exclusion criteria: Previously lived in a cholera endemic area, antibiotic therapy in previous 2 weeks
Interventions	Vaccine: CVD 103-HgR live attenuated vaccine containing: <ul style="list-style-type: none"> • 5 x 10⁸ CFU of lyophilized genetically modified <i>V. cholerae</i> O1 Classical Inaba (569B) Placebo: 5 x 10 ⁸ CFU of heat-killed <i>E. coli</i> K12 strain All participants were randomized to receive one dose with crossover to receive the alternative arm after 8 days
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none"> • Adverse events during first 7 days after vaccination <i>Not included in the review:</i> <ul style="list-style-type: none"> • Immunological outcome: Geometric mean serum vibriocidal antibody titres, and IgG antitoxin pre and post dose, proportion who develop ≥4 fold rises in serum titres, Excretion of vaccine strain.

Kotloff 1992 (Continued)

Notes	Location: Maryland, USA Setting: College students Source of funding: Swiss Serum and Vaccine Institute	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: 'In a double-blind fashion, subjects were randomly allocated to receive a single dose'
Allocation concealment (selection bias)	Unclear risk	Comment: None described
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'The blind was maintained through analysis of data'.
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No losses to follow up are reported
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Lagos 1993

Methods	Design: A randomized controlled trial (individually randomized) Duration: Vaccination took place Nov to Dec 1991; follow-up 7 days
Participants	Sample size: 81 enrolled Inclusion criteria: Age 18 to 35 years, male conscripts of the Chilean Air Force, employees of the Roberto del Rio Hospital, and medical students of the university of Chile, informed consent Exclusion criteria: Antibiotics or diarrhoea during the previous week, signs or symptoms of any acute disease, any type of chronic ailment
Interventions	Vaccine: CVD 103-HgR live attenuated vaccine containing: <ul style="list-style-type: none"> • 5 x 10⁹ lyophilized organisms of a genetically modified <i>V. cholerae</i> O1 Placebo: 5 x 10 ⁹ heat-killed <i>E. coli</i> K12 strain (C600)

	All participants were randomized to receive one dose	
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none">Adverse events during the first 7 days after the vaccine <i>Not included in the review:</i> <ul style="list-style-type: none">Immunological outcomes: Geometric mean serum vibriocidal antibody titres, and IgG antitoxin pre and post dose, proportion who develop ≥ 4 fold rises in serum titres,Excretion of vaccine strain.	
Notes	Location: Chile Setting: Source of funding: None stated	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: Not described.
Allocation concealment (selection bias)	Low risk	Quote: 'The codes were kept in confidential archives at the Swiss Institute of Sera and Vaccines...until the end of the serological analysis'
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'All observations were made using the double-blind methodology'
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: All participants are included in the adverse event data
Selective reporting (reporting bias)	Low risk	No evidence of selective outcome reporting
Other bias	Low risk	No other bias detected

Lagos 1995

Methods	Design: Randomized controlled trial (individually randomized) Duration: Study dates not given; follow-up 8 days
Participants	Sample size: 349 enrolled Inclusion criteria: Age 5 to 9 years, from public schools in a low-socioeconomic-level community Exclusion criteria: Fever, antibiotic therapy, or chronic disease
Interventions	Vaccine: CVD 103-HgR live attenuated vaccine containing: <ul style="list-style-type: none"> • 5 x 10⁹ lyophilized organisms of a genetically modified <i>V. cholerae</i> O1 Classical Inaba (569B) Placebo: Heat-killed <i>E. coli</i> K12 strain All participants were randomized to receive one dose
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none"> • Adverse events during first 8 days after vaccination <i>Not included in the review:</i> <ul style="list-style-type: none"> • Immunological outcome: Geometric mean serum vibriocidal antibody titres, and IgG antitoxin pre and post dose, proportion who develop ≥ 4 fold rises in serum titres, Excretion of vaccine strain.
Notes	Location: Santiago, Chile Setting: Source of funding: The World Health Organization and NIAID

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: Described as 'randomized', no further details given
Allocation concealment (selection bias)	Low risk	Quote: 'Lyophilized vaccine and placebo were contained in randomized coded aluminum foil sachets. The code remained unbroken until the clinical study, including serology, was completed'
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'Double-blind clinical follow-up to detect adverse reactions was maintained daily for 8 days after the single oral immunization'
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported

Lagos 1995 (Continued)

Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No losses recorded during adverse event monitoring
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Lagos 1999

Methods	Design: Randomized controlled trial (individually randomized) Trial dates and duration: June 1995 to Nov 1997; follow-up 28 days
Participants	Sample size: 312 enrolled Inclusion criteria: Age 3 to 17 months, normal medical history, parental consent Exclusion criteria: Signs and symptoms of acute illness, antibiotic therapy or diarrhoea in previous 2 days
Interventions	Vaccine: CVD 103-HgR live attenuated vaccine containing: <ul style="list-style-type: none"> • 5 x 10⁹ CFU of lyophilized organisms of a genetically modified <i>V. cholerae</i> O1 Placebo: 5 x 10 ⁸ CFU of heat-killed <i>E. coli</i> K12 strain Participants were initially randomized to receive vaccine or placebo. After 14 days all participants in both groups received a dose of vaccine
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none"> • Adverse events during first 7 days after first vaccination (daily home visit and symptom enquiry), <i>Not included in the review:</i> <ul style="list-style-type: none"> • Immunological outcome: Geometric mean serum vibriocidal antibody titres, and IgG antitoxin pre and post dose, proportion who develop ≥4 fold rises in serum titres, Excretion of vaccine strain.
Notes	Location: Santiago, Chile Setting: Well-baby clinics at a semi-rural ambulatory health centre Source of funding: The World Health Organization and NIAID

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: 'For the first dose one-half were randomly allocated in double blind fashion to receive a dose of vaccine'
Allocation concealment (selection bias)	Low risk	Quote: 'The sachets of test product were packed as individual treatments consisting of two sachets labeled with the same number followed by the letter A or B, indicating the appropriate sachet for the first and

Lagos 1999 (Continued)

		second dose of the immunization regimen', 'Each subject received the treatment number matching his/her study identification number'
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Comment: Described as 'double-blind'.
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Unclear risk	Comment: All randomised participants completed the follow-up for adverse events following the first dose
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Mahalanabis 2008

Methods	Design: Randomized controlled trial (individually randomized) Trial dates and duration: Enrollment from Aug to Oct 2005; follow-up for 28 days
Participants	Sample size: 201 participants stratified into adults and children Inclusion criteria: Age 18 to 40 years for adult study, 1 to 17 years for children, healthy male and non-pregnant females, written informed consent Exclusion criteria: Abdominal pain, vomiting, loss of appetite, generalized ill-feeling or nausea during the preceding 24 hours, diarrhoea or history of anti-diarrhoeal or antibiotic use during the past week, history of diarrhoea and abdominal pain lasting for more than 2 weeks during the past 6 months
Interventions	Vaccine: Bivalent killed whole-cell vaccine (BivWC: Shanchol®, Shantha Biotechnics) Placebo: Heat-killed <i>E. coli</i> K12 strain All subjects were randomized to receive 2 doses, at an interval of 14 days. All doses were administered via an oral syringe and offered water
Outcomes	No evidence of other bias
Notes	Location: Kolkata, India Setting: Clinical trial ward of the Infectious Diseases Hospital Source of funding: The Bill and Melinda Gates Foundation through the Diseases of Most Impoverished Program administered by the International Vaccine Institute, and

	the Swedish International Development Cooperation Agency	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: 'Separate randomization lists for adults and children were prepared by a statistician in IVI who was otherwise not involved in the study. Randomization numbers were generated in blocks of 8 using the program Visual Fortran 5.0. (Digital, USA)'
Allocation concealment (selection bias)	Low risk	Quote: 'Study agents were coded using 8 letters (4 for vaccine and 4 for placebo) in the adult trial and 8 different letters in the pediatric trial. Only the code letters on the vials identified the study agents as vaccine or placebo. The codes were revealed to the researchers once recruitment, data collection, and laboratory analyses were complete'
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'All study personnel and participants were blinded to treatment assignment during the duration of the study'
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: Three randomized participants (2.5%) were excluded from the adverse event follow-up, one who declined the first dose and two who received the wrong allocation for the second dose
Selective reporting (reporting bias)	Low risk	No evidence of selective outcome reporting
Other bias	Low risk	No evidence of other bias

Migasena 1989a

Methods	Design: A randomized controlled study (individually randomized) Duration: Dates not given; follow-up 5 days
Participants	Sample size: 24 enrolled Inclusion criteria: Healthy adults age 20 to 30 years, informed consent Exclusion criteria: None stated
Interventions	Vaccine: CVD 103-HgR live attenuated vaccine containing: <ul style="list-style-type: none"> • 5 x 10⁸ lyophilized organisms of a genetically modified <i>V. cholerae</i> O1 Classical Inaba (569B) Placebo: 5 x 10 ⁸ heat-killed <i>E. coli</i> K12 strain (C600) On day 5 all participants began a 5-day course of tetracycline
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none"> • Adverse events within 5 days of the vaccine (Daily interview) <i>Not included in the review:</i> <ul style="list-style-type: none"> • Immunological outcomes: Geometric mean serum vibriocidal antibody titres, and IgG antitoxin pre and post dose
Notes	Location: Thailand Setting: Vaccine Trial Centre in the Faculty of Tropical Medicine Source of funding:

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: Described as 'randomized', no further details given
Allocation concealment (selection bias)	Low risk	Quote: 'The study was carried out in double-blind fashion without the volunteers, the nursing staff, or the clinical investigators knowing the identity of the contents of the packets. A four-letter code for the packets was employed as an extra precaution to maintain double blindness'
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	See above
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported

Migasena 1989a (Continued)

Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No losses are reported
Selective reporting (reporting bias)	Low risk	No evidence of selective outcome reporting
Other bias	Low risk	No other sources of bias identified

Perry 1998

Methods	Design: Randomized controlled cross-over trial (individually randomized) Trial dates and duration: Dates not stated, cross-over after 12 days.	
Participants	Number of participants: 76 Inclusion criteria: Clinically healthy commercial sex workers and students aged 18 to 50 years. Half were HIV-positive and half were age and sex matched HIV-negative. Exclusion criteria: Pregnancy; clinical symptoms of AIDS; previous cholera vaccination; reported having previously had cholera; taken antibiotics within the previous 4 days; current diarrhoea or other acute illness	
Interventions	vaccine: CVD 103-HgR live attenuated vaccine containing: <ul style="list-style-type: none">• 5 x 10⁹ CFU of a genetically modified <i>V. cholerae</i> O1 Placebo: Lactate and aspartame only (these are also constituents of the vaccine) On day 12 those who initially received the placebo now received the vaccine, and vice versa	
Outcomes	<i>Included in the review:</i> <ul style="list-style-type: none">• Adverse events (Active surveillance; daily visits by physicians every day for 6 days and every other day for a further 6 days. Only adverse events prior to crossover are included) <i>Not included in the review:</i> <ul style="list-style-type: none">• Rectal swabs for vaccine virus on days of inoculation, daily for 4 days and on days 6 and 12• Immunological outcomes: Geometric mean vibriocidal antibodies pre and post vaccination. Seroconversion rates (criteria not stated)	
Notes	Location: Mali Setting: Not clear Source of funding: WHO Global Programme on Vaccines, National Institute of Allergy and Infectious Diseases, Centre for Vaccine Development, University of Maryland School of Medicine	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: Described as 'randomized', no further details given

Perry 1998 (Continued)

Allocation concealment (selection bias)	Unclear risk	Comment: Not described
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'identically appearing placebo packets'
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: Four (6%) participants were lost to follow-up during the study
Selective reporting (reporting bias)	Low risk	No evidence of selective outcome reporting
Other bias	Low risk	No other sources of bias identified

Qadri 2005

Methods	Design: Randomized controlled trial (individually randomized) Trial dates and duration: Dates not stated, follow-up 21 days
Participants	Number of participants: 70 Inclusion criteria: Age 18 to 45 years, healthy, willing to participate, informed consent Exclusion criteria: Any chronic disease or recent illness, immunosuppressive conditions during the past 6 months, pregnancy, diarrhoeal illness in the last 6 weeks, febrile illness in the last week or antibiotics in the last 2 weeks, history of any enteric vaccine given in the last month, stool samples positive for any enteric pathogen, food handlers and those cooking for or looking after infants and young children
Interventions	Vaccine: Peru 15 - a live attenuated strain of <i>V. cholerae</i> 01 El Tor Inaba (AVANT Immunotherapeutics Inc, US) containing: <ul style="list-style-type: none"> • 2 x 10⁸ CFU plus buffer Placebo: Buffer only All participants were given doxycycline for four days on day 6 to clear the vaccine strain
Outcomes	<i>Included in the review:</i> <ul style="list-style-type: none"> • Adverse events: reported up to 4 days after vaccination (patients were seen twice daily by a clinician or visited daily at home) <i>Not included in the review:</i> <ul style="list-style-type: none"> • Immunological outcomes: Geometric mean vibriocidal antibody titres, IgA and IgG antitoxin titres and IgA anti-lipopolysaccharide titres; on days 1, 7 and 21, and proportion who developed ≥4-fold rises from baseline after one dose

Qadri 2005 (Continued)

Notes	Location: Dhaka, Bangladesh Setting: Participants recruited from an urban slum, close to inpatient and outpatient facilities of ICCDR,B Source of funding: Diseases of the Most Impoverished Program, Bill and Melinda Gates Foundation, International Vaccine Institute	
<i>Risk of bias</i>		
Bias	Authors’ judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: ‘Randomization was performed by the International Vaccine Institute, South Korea’
Allocation concealment (selection bias)	Unclear risk	Comment: None described.
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Unclear risk	Comment: Described as ‘double-blind’. No further details given
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No losses to follow-up during the 4 days of adverse event reporting
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting.
Other bias	Low risk	No other bias identified

Qadri 2007

Methods	Trial design: Randomized controlled trial (individually randomized) Trial dates and duration: Dates not stated, follow-up 21 days
Participants	Number of participants: 240 Inclusion criteria: age 9 months to 5 years, healthy, parental consent Exclusion criteria: Any chronic disease; any recent illness; any illness or treatment causing immunosuppression in the last 9 months; diarrhoeal illness in the last two weeks; febrile illness in the last week; any enteric vaccine given in the last month; stool samples positive for any enteric pathogen

Interventions	Vaccine 1: Full dose Peru 15 - a live attenuated strain of <i>V. cholerae</i> 01 El Tor Inaba (AVANT Immunotherapeutics Inc, US) <ul style="list-style-type: none">• 2 x 10⁸ CFU plus buffer Vaccine 2: Reduced dose Peru 15 - a live attenuated strain of <i>V. cholerae</i> 01 El Tor Inaba (AVANT Immunotherapeutics Inc, US) <ul style="list-style-type: none">• 2 x 10⁷ CFU plus buffer Placebo: Buffer only All participants were given erythromycin for four days on day 6 to clear the vaccine strain	
Outcomes	<i>Included in the review:</i> <ul style="list-style-type: none">• Adverse events: reported up to 4 days after vaccination (monitored as an inpatient for 12 days then daily up to day 21, data only presented for first 4 days) <i>Not included in the review:</i> <ul style="list-style-type: none">• Immunological outcomes: Geometric mean Inaba and Ogawa vibriocidal antibody titres on days 1 and 7 and proportion who developed ≥4-fold rises from baseline after one dose	
Notes	Location: Dhaka, Bangladesh Setting: Participants recruited from an urban slum, close to inpatient and outpatient facilities of ICCDR,B Source of funding: Bill and Melinda Gates Foundation	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: 'Randomization was carried out by the International Vaccine Institute and sent to the vaccine formulation team.'
Allocation concealment (selection bias)	Low risk	Quote: 'The vaccine formulation team prepared and blinded the vaccine and the placebo according to the randomization list'
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'The results were analyzed sequentially in the order in which they had been completed and unblinded.'
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: Follow-up for adverse events was 100%.

Qadri 2007 (Continued)

Selective reporting (reporting bias)	Low risk	No evidence of selective outcome reporting
Other bias	Low risk	No other bias identified

Richie 2000 Indonesia

Methods	Design: Randomized controlled trial (individually randomized) Trial dates and duration: July 1993 to Dec 1997, 4 years Surveillance: Passive surveillance conducted through four North Jakarta hospitals distributed across the study area
Participants	Sample size: 67,508 Inclusion criteria: Persons aged 2 to 41 years living in the study area Exclusion criteria: Pregnancy, plans to move out of the study area, diagnosis of cancer
Interventions	Vaccine: CVD 103-HgR live attenuated vaccine containing: <ul style="list-style-type: none"> • 5 x 10⁹ lyophilized organisms of a genetically modified <i>V. cholerae</i> O1 Classical Inaba (569B) Placebo: 5 x 10 ⁸ heat-killed <i>E. coli</i> K12 strain (C600)
Outcomes	<i>Included in the review:</i> <ul style="list-style-type: none"> • Cases of cholera • Deaths other than those caused by vehicular accident • Adverse events reported during the three days after ingestion of the vaccine or placebo <i>Not included in the review:</i> <ul style="list-style-type: none"> • Immunological outcomes: Geometric mean vibriocidal antibody titres pre and post dose, and proportion who develop ≥ 4-fold rises after one dose
Notes	Location: 65 communities in North Jakarta, Indonesia Setting: Poor communities with poor sanitation and relative high cholera incidences Source of funding: World Health Organization, National Institute of Allergy and Infectious Diseases, the Swiss Serum and Vaccine Institute, Berne

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "A randomized trial" Comment: Method of randomization unclear
Allocation concealment (selection bias)	Low risk	Quote: "Vaccine..and placebo ..were contained in identical, numbered packets"
Blinding (performance bias and detection bias) Efficacy outcomes	Low risk	Quote: "Surveillance data collected included..vaccine number. ..Patients with diarrhoea who did not state that they were

Richie 2000 Indonesia (Continued)

		participants in the study were also included in the surveillance.”
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: “a double-blind nested study of adverse events was conducted..”
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Comment: No mention of the numbers of people who moved out of the area and were therefore lost of follow up
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No loss to follow up
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Sack 1997

Methods	Design: Randomized controlled trial (individually randomized) Trial dates and duration: Dates not given, 21 days follow-up
Participants	Number of participants: 50 in outpatient study (a smaller inpatient study is also reported but included no outcomes relevant to this review) Inclusion criteria: Age 18 to 50 years, healthy, willing to participate, informed consent Exclusion criteria: chronic illness, immunosuppressive condition, abnormal stool pattern, significant abnormality in screening laboratory hematology and chemistry tests, HIV +ve, hepatitis B surface antigen +ve, pregnancy, travel to a cholera endemic area within 5 years, receipt of cholera vaccine, history of cholera infection or vaccination, previous participation in a cholera or ETEC vaccine trial, use of antibiotics within 7 days of vaccination, food handlers or close contact with children under age 5
Interventions	Vaccine 1: Full dose Peru 15 - a live attenuated strain of <i>V. cholerae</i> 01 El Tor Inaba <ul style="list-style-type: none"> • 2 x 10⁹ CFU plus buffer Vaccine 2: Reduced dose Peru 15 - a live attenuated strain of <i>V. cholerae</i> 01 El Tor Inaba (AVANT Immunotherapeutics Inc, US) <ul style="list-style-type: none"> • 2 x 10⁸ CFU plus buffer Placebo: Buffer only
Outcomes	<i>Included in the review:</i> <ul style="list-style-type: none"> • Adverse events: reported up to 7 days after vaccination (using a self reported symptom diary) <i>Not included in the review:</i> <ul style="list-style-type: none"> • Immunological outcomes: Geometric mean titres of vibriocidal antibody and IgG antitoxin on days 0, 10 and 21, and proportion who developed ≥ 2 and ≥ 4-fold rises from baseline after one dose

Sack 1997 (Continued)

Notes	Location: USA Setting: Students or employee volunteers at John Hopkins University and Hospital Source of funding: National Institute Health, Virus Research Institute	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: Described as 'Randomized'. No further details given
Allocation concealment (selection bias)	Unclear risk	Comment: None described
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'Volunteers were randomized to receive either a 10 ⁹ or 10 ⁸ CFU or a placebo in a double-masked manner', 'To protect the masked code, some volunteers were assigned to 1 or 0.1 mL of placebo'
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No losses are described during the adverse event monitoring
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Sanchez 1993a

Methods	Design: A randomized 2 x 2 factorial controlled trial (individually randomized) Duration: Enrolled during October 1991; follow-up 28 days
Participants	Sample size: 186 enrolled and randomized to four groups* Inclusion criteria: Age 18 to 44 years Exclusion criteria: Pregnancy or planned pregnancy, diarrhoea or fever within the past 5 days, recent use of antimotility or antibacterial agent, foodhandler, chronic gastrointestinal disorder
Interventions	Vaccine: Killed whole-cell vaccine plus recombinant cholera toxin subunit (WC-rBS; Dukoral®, SBL, Sweden) Placebo: Buffer alone

	All participants were randomized to receive 2 doses, at 11 to 16 days apart	
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none">● Adverse events: text summary only <i>Not included in the review:</i> <ul style="list-style-type: none">● Immunological outcomes: Geometric mean-fold rise, and proportion who develop ≥ 2 or ≥ 4-fold rises in serum titres of anti-CT IgA, anti-CT IgG, and vibriocidal antibody from baseline after one or two doses	
Notes	Location:USA Setting: Military personnel Source of funding: None stated * A further single arm study involving 74 participants is reported in this paper but excluded from this review	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: Described as 'randomized', no further details given
Allocation concealment (selection bias)	Unclear risk	Comment: None described.
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Unclear risk	Comment: None described.
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: 6 participants did not receive the second dose as they were unavailable
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Sanchez 1994 Peru

Methods	Design: Randomized controlled trial (individual block randomization in groups of 10) Duration: Enrolled January to March 1994, follow-up to June 1994 Surveillance: Passive surveillance through clinics within the military training centres where the study was conducted	
Participants	Sample size: 1563 enrolled, 1426 received 2 doses of vaccine or placebo Inclusion criteria: 17-65 years volunteers, available for three months Exclusion criteria: Previous cholera vaccination	
Interventions	Vaccine: Killed whole-cell vaccine plus recombinant cholera toxin subunit (WC-rBS; Dukoral®, SBL, Sweden) Placebo: Heat -inactivated <i>E. coli</i> K12 strain Vaccine and placebo were given with freshly prepared antacid solution. Two doses were given, 7 to 14 days apart	
Outcomes	<i>Included in the review:</i> <ul style="list-style-type: none">• Cases of confirmed cholera <i>Not included in the review:</i> <ul style="list-style-type: none">• Cases of severe cholera (cholera with signs of dehydration) as the treatment group is not stated• Cases subgrouped by blood group as the treatment group is not stated	
Notes	Location: Lima, Peru Setting: Military training centres Source of funding: Not stated	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Randomization was done in blocks of 10 to ensure equal study groups" Comment: Unclear description but probably done
Allocation concealment (selection bias)	Low risk	Quote: "Each bottle was identified with a unique code number; vaccine and placebo bottles were pre-coded"
Blinding (performance bias and detection bias) Efficacy outcomes	Low risk	Quote: "The placebo....in a concentration identical in turbidity and appearance to the vaccine preparation" Plus see above
Blinding (performance bias and detection bias) Safety outcomes	Unclear risk	Not applicable as safety data is not reported

Sanchez 1994 Peru (Continued)

Incomplete outcome data (attrition bias) Efficacy outcomes	High risk	Comment: Subjects lost to follow-up after the second dose were assumed to contribute half the period to the denominator analysis. Mentioned that losses to follow-up were similar in both groups Cases of cholera which occurred in participants between study doses were excluded from the primary analysis
Incomplete outcome data (attrition bias) Safety outcomes	Unclear risk	Not applicable as safety data is not reported
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Sanchez 1995 Peru

Methods	Design: Randomized controlled trial (individually randomized) Trial dates and duration: February 1992, 4 weeks Surveillance: Passive surveillance for diarrhoea was performed at the single military medical clinic, where all cases were evaluated
Participants	Sample size: 346 enrolled and received first dose, 307 received two full doses Inclusion criteria: Male Hispanics aged 17-23 years, informed consent Exclusion criteria: Major illness at the time of vaccination, previous cholera vaccine
Interventions	Vaccine: Killed whole-cell vaccine plus recombinant cholera toxin subunit (WC-rBS; Dukoral®, SBL, Sweden) Placebo: suspension of heat-inactivated <i>E. coli</i> K12 strain Vaccines were given with a buffer solution. Two doses were given two weeks apart
Outcomes	<i>Included in the review:</i> <ul style="list-style-type: none"> • Cholera cases • Adverse events within 24 hours of each dose (active surveillance was conducted for 3 days) <i>Not included in the review:</i> <ul style="list-style-type: none"> • Immunological outcomes: Geometric mean serum vibriocidal antibody titre pre and post vaccination, and proportion who develop ≥ 4 fold rises from baseline; anti-cholera toxin IgG titre pre and post vaccination, and proportion who develop ≥ 0.20 rises from baseline
Notes	Location: Ancon, Peru Setting: Military training centre Source of funding: US Army and Navy medical departments(?)
Risk of bias	

Sanchez 1995 Peru (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: 'randomly allocated' Comment: Method of randomization not adequately described but probably done
Allocation concealment (selection bias)	Low risk	Quote: "Each bottle was identified with one of 2 letters; vaccine and placebo preparations were pre-coded"
Blinding (performance bias and detection bias) Efficacy outcomes	Low risk	Quote: 'suspension of heat-inactivated <i>E coli</i> K12 strain, with same appearance as vaccine'.
Blinding (performance bias and detection bias) Safety outcomes	Low risk	See above
Incomplete outcome data (attrition bias) Efficacy outcomes	Low risk	Comment: All participants remained in the study area and were included in the analysis. Identification of cases through passive surveillance, with clinical data collected from all participants with diarrhoea
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No loss to follow-up
Selective reporting (reporting bias)	High risk	Comment: The report states adverse event data were collected for three days after each dose but only symptoms within 24 hours are presented
Other bias	Low risk	No evidence of other bias

Simanjuntak 1993

Methods	Trial design: Randomized controlled trial - initially randomized in pairs one to each treatment arm, later changed to individual randomization Trial dates and duration and dates: 1991 to 1992
Participants	Number of participants: 303 Inclusion criteria: Children aged 24 to 59 months Exclusion criteria: Chronic health problem, receiving antibiotic therapy, acute illness on the scheduled day of vaccination
Interventions	Vaccine: CVD 103-HgR live attenuated vaccine containing: <ul style="list-style-type: none"> • 5 x 10⁹ CFU of a lyophilised genetically modified strain of <i>V. cholerae</i> O1

	Placebo: <ul style="list-style-type: none">• 5 x 10⁸ CFU inactivated <i>E. coli</i> K12 (placebo) Both were given with aspartame sweetener and a buffer.	
Outcomes	<i>Included in the review:</i> <ul style="list-style-type: none">• Adverse events (active surveillance; daily visits by physicians to record complaints and conduct physical examination, up to day nine after vaccination) <i>Not included in the review:</i> <ul style="list-style-type: none">• Stools samples for vaccine virus on day 5• Immunological outcomes: Serum vibriocidal antibody titres on days 0, 9 and 28 and proportion with a ≥ 4-fold rises from baseline.	
Notes	Location: North Jakarta, Indonesia Setting: Villages Source of funding:Consultative group on vaccine development of the national vaccine programme, USA, National Institute of Allergy and Infectious Diseases, US Naval Medical Research and Command	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Comment: Described as randomized. Codes generated by the manufacturer
Allocation concealment (selection bias)	Low risk	Quote: 'Coded preparations looked identical and were only identified by the codes 'N' or 'O'
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'The clinical follow-up as well as the administration of the vaccine was double-blind with neither the clinical staff, the patient or their parents knowing the identity of the preparation'
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No losses during adverse event follow-up are noted.
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Su-Archawaratana 1992a

Methods	Trial design: Randomized controlled trial (individually randomized) Trial dates and duration: February 1988
Participants	Number of participants: 206 (in study 1), Inclusion criteria: Thai soldiers aged 18 to 26, who volunteered for the study Exclusion criteria: Previous parenteral inactivated whole cell vaccine
Interventions	Vaccine: CVD 103-HgR live attenuated vaccine containing: <ul style="list-style-type: none"> • 5 x 10⁸ CFU of a lyophilised genetically modified strain of <i>V. cholerae</i> O1 Placebo: 5 x 10 ⁸ CFU inactivated <i>E. coli</i> K12
Outcomes	<i>Included in the review:</i> <ul style="list-style-type: none"> • Adverse events (examined every day for 7 days) although only diarrhoea is reported <i>Not included in the review:</i> <ul style="list-style-type: none"> • Immunological outcomes: Serum vibriocidal antibodies titres on days 0, 7 and 21, and the proportion who develop a ≥ 4-fold increase.
Notes	Location: Thailand Setting: Field study using volunteers Sources of funding: National Institutes of Health, Swiss Serum and Vaccine Institute, US Agency for International Development

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: Described as 'randomised', no further details
Allocation concealment (selection bias)	Unclear risk	Comment: None described
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Unclear risk	Comment: None described
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No losses are reported
Selective reporting (reporting bias)	Low risk	No evidence of selective outcome reporting
Other bias	Low risk	No other bias identified

Su-Archawaratana 1992b

Methods	Trial design: Randomized controlled crossover trial. Trial dates and duration: June 1991
Participants	Number of participants: 120 Inclusion criteria: Volunteers and Thai soldiers aged 18 to 26 Exclusion criteria: None stated
Interventions	Vaccine: CVD 103-HgR live attenuated vaccine containing: <ul style="list-style-type: none"> • 5 x 10⁸ CFU of a lyophilised genetically modified strain of <i>V. cholerae</i> O1 • 5 x 10⁹ CFU of a lyophilised genetically modified strain of <i>V. cholerae</i> O1 Placebo: 5 x 10 ⁸ CFU inactivated <i>E. coli</i> K12 The trial had 6 arms with each arm crossing over to receive the alternative dose or placebo on day 7 All doses were given with buffer and aspartame sweetener
Outcomes	<i>Included in the review:</i> <ul style="list-style-type: none"> • Adverse events (examined every day for 7 days after each dose) although only diarrhoea is reported <i>Not included in the review:</i> <ul style="list-style-type: none"> • Immunological outcomes: Serum vibriocidal antibodies titres on days 0, 7 and 21, and the proportion who develop a ≥ 4 fold increase.
Notes	Location: Thailand Setting: Field study using volunteers Sources of funding: National Institutes of Health, Swiss Serum and Vaccine Institute, US Agency for International Development

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: Described as 'randomized', no further details
Allocation concealment (selection bias)	Unclear risk	Comment: None described
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Unclear risk	Comment: Described as 'double blind', no further details.
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No losses are reported

Su-Arehawaratana 1992b (Continued)

Selective reporting (reporting bias)	Low risk	No evidence of selective outcome reporting
Other bias	Low risk	No other bias identified

Suharyono 1992a

Methods	Trial design: Randomized controlled trial (4 arms) Trial dates and duration: February to March 1990
Participants	Number of participants: 274 Inclusion criteria: Children aged 5 to 9 years. Written parental consent. Only one child per family was eligible to take part Exclusion criteria: Having a chronic health disorder; receiving antibiotic therapy; acute illness on the scheduled day of vaccination
Interventions	Vaccine: CVD 103-HgR live attenuated vaccine containing: <ul style="list-style-type: none"> • 5×10^6 CFU CVD 103HgR centrifuged • 5×10^7 CFU CVD 103HgR centrifuged • 5×10^8 CFU CVD 103HgR filtered Placebo: 5×10^8 CFU inactivated <i>E. coli</i> K12 strain
Outcomes	<i>Included in the review:</i> <ul style="list-style-type: none"> • Adverse events (Active surveillance; daily visits by study staff for 9 days) <i>Not included in the review:</i> <ul style="list-style-type: none"> • Stools samples for vaccine virus on day 5 • Immunological outcomes: Serum vibriocidal antibodies on days 0, 9 and 28, and the proportion who develop a ≥ 4 fold increase.
Notes	Location: North Jakarta, Indonesia Setting: Small rural village, vaccinated at village health office Source of funding: Consultative group on vaccine development of the national vaccine programme, USA, National Institute of Allergy and Infectious Diseases, Naval Medical Research unit 2, Jakarta, United States Agency for International Development

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: 'Randomly allocated to one of the four letter-coded groups according to a computer-generated randomization sequence'
Allocation concealment (selection bias)	Low risk	Quote: 'Vaccine and placebo packets indistinguishable and identified only by a colour-coded letter'

Suharyono 1992a (Continued)

Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'The clinical follow-up as well as the administration of the vaccine was double-blind with neither the clinical staff, the patient or their parents knowing the identity of the preparation'
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No losses reported.
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Suharyono 1992b

Methods	Trial design: Randomized controlled trial (individually randomized) Trial dates and duration: Sept to Oct 1990	
Participants	Number of participants: 140 Inclusion criteria: As for Suharyono 1992a Exclusion criteria: As for Suharyono 1992a	
Interventions	Vaccine: CVD 103-HgR live attenuated vaccine containing: <ul style="list-style-type: none">• 5 x 10⁹ CFU CVD 103HgR centrifuged• 5 x 10¹⁰ CFU CVD 103HgR centrifuged• 5 x 10⁹ CFU CVD 103HgR filtered• 5 x 10¹⁰ CFU CVD 103HgR filtered• Half of the children in each of these groups were randomized to also receive an extra half dose of buffer Placebo: 5 x 10 ⁸ CFU inactivated <i>E. coli</i> K12 strain	
Outcomes	As for Suharyono 1992a	
Notes	Location: As for Suharyono 1992a Setting: Source of funding: As for Suharyono 1992a	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement

Suharyono 1992b (Continued)

Random sequence generation (selection bias)	Low risk	Quote: 'the child was allocated to receive one of the nine treatment groups, according to a randomised sequence' Comment: Study A in the same paper used a computer to generate the sequence
Allocation concealment (selection bias)	Low risk	Comment: Only clearly described for study A but probably done
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'The clinical follow-up as well as the administration of the vaccine was double-blind with neither the clinical staff, the patient or their parents knowing the identity of the preparation'
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No losses to follow up are reported
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Sur 2009 India

Methods	Design: Randomized controlled trial (cluster randomization) Dates and duration: Vaccination July to September 2006; An interim analysis after 2 years of follow-up Surveillance: Passive surveillance through nine diarrhoea clinics established in the study area, two hospitals serving the study area, and encouragement to private medical practitioners to refer to the treatment centres Method of adjustment for clustering: Robust sandwich variance estimates
Participants	Sample size: 3933 clusters (107,774 individuals) were randomized. 69,328 individuals received at least one dose of vaccine or placebo. The primary analysis includes 66,900 participants who received 2 doses of the vaccine Inclusion criteria: Age > 1 year, written informed consent Exclusion criteria: Pregnancy
Interventions	Vaccine: Bivalent killed whole-cell vaccine (BivWC: Shanchol®, Shantha Biotechnics) Placebo: Heat killed <i>E. coli</i> K12 strain All subjects were randomized to receive 2 doses, at a minimum interval of 14 days. All

	doses were administered via an oral syringe	
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none">● First symptomatic cholera episode detected using a passive surveillance system with confirmation of faecal excretion of <i>V. Cholerae</i> 01 during a non-bloody diarrhoeal episode.● All-cause death● Serious adverse events within 14 days of vaccination● Adverse events within 14 days of each dose. <i>Not included in the review:</i>	
Notes	Location: Kolkata, India Setting: Surveillance study area, served by 9 study clinics, private practitioners and 2 hospitals Source of funding: Bill and Melinda Gates Foundation; Swedish International Development Cooperation Agency; governments of South Korea, Sweden and Kuwait	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: 'An external statistician, who was masked to the identities of the codes, used an SAS version 9.1 computer algorithm to randomly assign dwellings to the four codes in a 1:1:1:1 ratio within each of the strata'
Allocation concealment (selection bias)	Low risk	Quote: 'The vials were labelled with one of four letters, two each for vaccine and placebo. Project staff and study participants were unaware of the identities of the codes'
Blinding (performance bias and detection bias) Efficacy outcomes	Low risk	Quote: 'The vaccine and placebo were identical in appearance'
Blinding (performance bias and detection bias) Safety outcomes	Low risk	See above
Incomplete outcome data (attrition bias) Efficacy outcomes	Low risk	Comment: Attrition between the first and second doses of vaccine were low: 3.6% vaccine group vs 3.4% placebo group
Incomplete outcome data (attrition bias) Safety outcomes	High risk	Comment: Safety data was collected passively with participants requested to present to medical services. Consequently the incidence of adverse event reporting is very low and likely to be an underestimate

Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Tacket 1999

Methods	Design: Randomized controlled trial (individually randomized) with challenge study Trial dates and duration: Dates not stated; challenge study was undertaken 3 months after vaccination
Participants	Sample size: 85 (51 included in challenge study) Inclusion criteria: Age 18 to 40, healthy, informed consent. Exclusion criteria: clinically significant abnormalities on urinalysis, complete blood count, serum hepatic transaminases, glucose, creatinine, blood urea nitrogen, electrolytes, or electrocardiogram, travel to a cholera endemic area in the previous 5 years, abnormal stool pattern or regular use of laxatives, failure to pass a psychological examination, allergy to tetracycline or ciprofloxacin, history of cholera or enterotoxigenic <i>E. coli</i> challenge, history of recent antibiotic use, pregnancy or nursing, positive serology for HIV, hepatitis B antigen, or hepatitis B, stool culture positive for an enteric pathogen
Interventions	Vaccine: CVD 103-HgR live attenuated vaccine containing: <ul style="list-style-type: none"> 2 to 8 x 10⁸ CFU of lyophilized organisms of a genetically modified strain of <i>V. cholerae</i> O1 plus buffer Placebo: heat-inactivated <i>E. coli</i> K12 plus buffer Challenge: 10 ⁵ organisms of <i>V. cholerae</i> O1 El Tor Inaba (N16961)
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none"> Adverse events following vaccine (symptom diary for 3 days) Any diarrhoea following artificial challenge Moderate or severe cholera diarrhoea following artificial challenge <i>Not included in the review:</i> <ul style="list-style-type: none"> Immunological outcomes: Geometric mean serum vibriocidal antibody titre pre and post vaccination, and proportion who develop ≥4 fold rises from baseline; anti-cholera toxin IgG titre pre and post vaccination, and proportion who develop ≥0.20 rises from baseline
Notes	Location: Baltimore and Cincinnati, USA Setting: Hospital Source of funding: National Institute of Allergy and Infectious Diseases, and the Swiss Serum and Vaccine Institute

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: 'For those of blood group O and non-O within each clinical center, subjects were randomized in blocks of four (two to

Tacket 1999 (Continued)

		receive vaccine and two to receive placebo) by using SAS PROC PLAN'
Allocation concealment (selection bias)	Low risk	Quote: 'The code was held by the study sponsor until the database was complete and unalterable'
Blinding (performance bias and detection bias) Efficacy outcomes	Low risk	Quote: 'When suspended in the buffer solution, the placebo was identical in appearance to the vaccine suspension'
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'When suspended in the buffer solution, the placebo was identical in appearance to the vaccine suspension'
Incomplete outcome data (attrition bias) Efficacy outcomes	High risk	Comment: The artificial challenge study included only 60% of those given the vaccine
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No participants were lost to follow-up or excluded during the initial 3 days
Selective reporting (reporting bias)	Low risk	No evidence of selective outcome reporting
Other bias	Low risk	No other bias identified.

Taylor 1999a

Methods	Design: Randomized controlled trial (individually randomized) Duration: Enrollment from Jan to Feb 1995; follow-up for 28 days
Participants	Sample size: 216 enrolled Inclusion criteria: Age 2 to 64 years and residing in the study area, informed consent Exclusion criteria: None stated.
Interventions	Vaccine: Killed whole-cell vaccine plus recombinant cholera toxin subunit (WC-rBS; Dukoral®, SBL, Sweden) Placebo: Heat killed <i>E. coli</i> K12 strain All participants were randomized to receive 2 doses, at a minimum interval of 14 days. All doses were administered via pumps designed to deliver the correct dose
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none"> Adverse events within 3 days of each dose <i>Not included in the review:</i> <ul style="list-style-type: none"> Immunological outcomes: Geometric mean serum vibriocidal antibody titre, proportion who develop ≥ 2 or ≥ 4-fold rises from baseline after one or two doses

Notes	Location: Flores de Villa, southern Lima Setting: Source of funding: The U.S. Army Medical Material and Development Command	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: 'Vaccination teams were assigned to a section of households using pre-randomized forms to enter adults and children in the study'.
Allocation concealment (selection bias)	Low risk	Quote: 'Each bottle was identified with a unique number; vaccine and placebo preparations were pre-coded'
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'The participants and the persons who assessed side effects were blinded to the vaccine code'. 'The placebo consisted of a suspension of heat-inactivated E. coli K12 strain (SBL Vaccin AB) in a concentration that matched the turbidity and appearance of the vaccine preparation'
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: 12 participants were lost to follow-up between doses. Reasons for drop-out were not given but follow-up in the 3 days after each dose was complete
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of any other bias

Taylor 2000 Peru

Methods	Design: Randomized controlled trial (individually randomized) Duration: 1993 to 1995 Surveillance: Active surveillance in the community through twice weekly visits to each household
Participants	Sample size: 21,924 received the first dose, 17,799 received the second dose, and 14,997 received the booster dose Inclusion criteria: Aged 2 to 65 years old and residing in the vaccine trial area Exclusion criteria: None stated.
Interventions	Vaccine: Killed whole-cell vaccine plus recombinant cholera toxin subunit (WC-rBS; Dukoral®, SBL, Sweden) Placebo: Heat-inactivated <i>E. coli</i> K12 strain Vaccine or placebo were given as two doses two weeks apart, followed by a third dose 10 months later
Outcomes	<i>Included in the review:</i> <ul style="list-style-type: none"> Cases of cholera identified through active household surveillance. Rectal swabs were collected from participants with diarrhoea and cultured to test for <i>V. cholerae</i>. Cases of cholera identified through passive surveillance at the health post and hospital serving the area. Level of dehydration in participants with cholera; using WHO definitions of mild, moderate or severe. Adverse events after the first dose: Symptom enquiry at time of second dose <i>Not included in the review:</i> <ul style="list-style-type: none"> Immunological outcomes: Plasma vibriocidal and anti-cholera toxin antibodies at day 14 after the second dose.
Notes	Location: Pampas de San Juan de Miraflores, in the southern outskirts of Lima, Peru Setting: 36 poor marginal neighbourhoods, with a nearby hospital, 4 health posts and 40 neighbourhood rehydration units Source of funding: US Army Medical Materiel and Development Command, Fort Detrick, Maryland

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The trial area was divided into 4 quadrants (A to D), and every eligible person was randomly assigned a vaccine code of 1 or 9 to give a total of 8 possible codes" Comment: Sequence generation unclear but probably low risk
Allocation concealment (selection bias)	Low risk	Quote: "During the study, the vaccine codes were kept locked by.. who was not involved in the study; the codes were not known to any person conducting the trial"

Taylor 2000 Peru (Continued)

Blinding (performance bias and detection bias) Efficacy outcomes	Low risk	Quote: "Suspension of heat-inactivated <i>Escherichia coli</i> K12 strain in a concentration that matched the turbidity and appearance of the vaccine preparation." plus see above
Blinding (performance bias and detection bias) Safety outcomes	Low risk	See above
Incomplete outcome data (attrition bias) Efficacy outcomes	Low risk	Comment: Attrition between the first and second doses was high: 18.8% overall. As well as the per protocol analysis, the authors conducted an intention to treat analysis which did not significantly alter the result
Incomplete outcome data (attrition bias) Safety outcomes	High risk	Comment: Adverse events from the first dose were recorded only for those attending for a second dose. This method is likely to underestimate the true incidence of adverse events if those experiencing a significant event after the first dose are more likely to drop-out
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of other bias

Trach 1997 Viet Nam

Methods	Design: Quasi-randomized controlled trial (alternate allocation, clustered by household) Duration: Vaccination started December 1992, follow-up to December 1993 Surveillance: Passive surveillance through community health centres, polyclinics and hospitals serving the study area Method of adjustment for clustering: Logistic regression models with generalised estimating equations
Participants	Sample size: 134,453 individuals, 22,653 households Inclusion criteria: Residents aged one year or older Exclusion criteria: None
Interventions	Vaccine: Variant killed whole cell vaccine (vWC; National Institute of Hygiene and Epidemiology, Vietnam) Control: No vaccine Two doses were given with a two week interval between them.

Outcomes	<i>Included in the review:</i> <ul style="list-style-type: none">• Cases of cholera requiring inpatient care in hospital or polyclinic (faecal sample yields <i>V. cholerae</i> 01)• Deaths from cholera• Visits to community health centres, polyclinics and hospitals for treatment of diarrhoea	
Notes	Location: Hue, central Vietnam Setting: city community served by 19 health centres, four polyclinics and one regional hospital Sources of funding: Ministry of Health Vietnam, Swedish Agency for Research Cooperation, World Health Organization, USA National Institute of Child Health and Human Development	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: 'Each household.. was given a serial number. Even-numbered households were assigned the vaccine, and odd numbered households were assigned no vaccine' Comment: Alternate allocation is unlikely to significantly bias a vaccine trial
Allocation concealment (selection bias)	Low risk	Comment: Alternate allocation, concealment not possible, but unlikely to introduce significant bias
Blinding (performance bias and detection bias) Efficacy outcomes	High risk	Quote: 'open field trial'
Blinding (performance bias and detection bias) Safety outcomes	Unclear risk	Not applicable - not included as an outcome
Incomplete outcome data (attrition bias) Efficacy outcomes	High risk	Comment: No mention of the participants who may have moved out of the area and therefore been lost to follow up. Cases identified through passive surveillance at the polyclinics and hospitals
Incomplete outcome data (attrition bias) Safety outcomes	Unclear risk	Not applicable as safety data not reported
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting

Other bias	Low risk	No evidence of any other bias
------------	----------	-------------------------------

Trach 2002

Methods	Design: A 3-arm randomized controlled trial (individually randomized) Trial dates and duration: Dates not stated, follow-up 28 days
Participants	Sample size: 71 in adult study, 70 in child study (from included study arms) Inclusion criteria: Adult study: Age 17 to 25 years, Hanoi residents. Child study: Age 1-12 years, attending an elementary school or day care centre in Hanoi Exclusion criteria: Diarrhoea during the preceding week, chronic or recurrent abdominal pain or diarrhoea, pregnancy, steroids or other immunosuppressive medications, antibiotics, or known to have HIV or another immunosuppressive condition
Interventions	Vaccine 1: Killed whole-cell vaccine plus recombinant cholera toxin subunit (WC-rBS; Dukoral®, SBL, Sweden) Vaccine 2: A bivalent vaccine containing: 5×10^{10} formalin-killed <i>V. cholerae</i> O1 Inaba, El Tor biotype cells (strain Phil 6973); 2.5×10^{10} heat-killed <i>V. cholerae</i> O1 Ogawa, classical biotype cells (strain Cairo 50); 2.5×10^{10} formalin-killed <i>V. cholerae</i> O1 Inaba, classical biotype cells (strain 569B); 2.5×10^{10} heat-killed <i>V. cholerae</i> O1 Inaba, classical biotype cells (strain Cairo 48); and 5×10^{10} formalin-killed <i>V. cholerae</i> O139 (strain AI4456): This arm was excluded as the included strains are different from both the vWC vaccine and the BivWC vaccines with efficacy data Placebo: heat-killed <i>E. coli</i> K12 strain
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none"> Adverse events (visited for three consecutive days to ask about AE plus an interview at day 14) Serious adverse events <i>Not included in the review:</i> <ul style="list-style-type: none"> Immunological outcomes: Geometric mean vibriocidal antibody titres pre and post vaccination, and proportion who develop a ≥ 4 fold increase.
Notes	Location: Hanoi, Vietnam Setting: Source of funding: Swedish Agency for Cooperation with Developing Countries; the National Institute of Child Health and Human Development, National Institutes of Health, USA; the World Health Organization; and the Diseases of the Most Impoverished Programme, funded by the Bill and Melinda Gates Foundation

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: 'Consenting eligible subjects in blocks of eight were randomly allocated' Comment: Description is unclear but probably done

Trach 2002 (Continued)

Allocation concealment (selection bias)	Low risk	Quote: 'The vials with the agent for each group were labelled with one of two code letters'. 'The codes were kept secret from all persons involved in the study until freezing of the data set.'
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy outcomes are not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Quote: 'The two vaccines and the placebo were packaged as liquid formulations in identical vials'
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy outcomes are not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: Only three participants were lost to follow-up between doses, although the reasons are not stated
Selective reporting (reporting bias)	High risk	Comment: Nausea was not assessed in the children's study
Other bias	Low risk	No other source of bias identified

Valera 2009

Methods	Design: Randomized controlled trial (individually randomized) Duration and dates (field work):
Participants	Sample size: 36 Inclusion criteria: Age 18 to 40 years, volunteers working in the scientific community, healthy, informed consent Exclusion criteria: Previous history of clinically significant diarrhoea or cholera vaccination, receiving medication at the time of recruitment
Interventions	Vaccine: VC638 - A live attenuated strain of <i>V. cholerae</i> O1 El Tor Ogawa (Final Institute, Havana) <ul style="list-style-type: none"> 1 x 10⁹ CFU plus buffer Placebo: Buffer alone All participants received 300mg of doxycycline on day 5.
Outcomes	<i>Included in review:</i> <ul style="list-style-type: none"> Adverse events (active surveillance for 5 days, then passive up to day 30) <i>Not included in the review:</i> <ul style="list-style-type: none"> Immunological outcomes: Geometric mean serum vibriocidal antibody titres on day 0 and 14, and proportion who develop ≥ 4-fold rises from baseline after one dose

Notes	Location: Havana, Cuba Setting: Unit for Isolation of Biological Risks at Tropical Medicine Institute Source of funding: None stated	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Comment: Described as 'randomized', no further details given
Allocation concealment (selection bias)	Low risk	Quote: 'Vaccine and placebo vials were packaged and coded at random with identical appearance. The code remained unbroken until the end of the study'
Blinding (performance bias and detection bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Blinding (performance bias and detection bias) Safety outcomes	Low risk	Comment: See above
Incomplete outcome data (attrition bias) Efficacy outcomes	Unclear risk	Not applicable as efficacy not reported
Incomplete outcome data (attrition bias) Safety outcomes	Low risk	Comment: No losses during the adverse event monitoring stage
Selective reporting (reporting bias)	Low risk	No evidence of selective reporting
Other bias	Low risk	No evidence of any other bias

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Ahmed 2006	Non-cholera vaccine, no cholera outcomes (abstract)
Ahren 1993	Non-comparative study.
Albert 2003	A non-comparative study, all children received the same vaccine (abstract)

(Continued)

Ali 2005	Relates to Clemens 1988 Bangladesh. No additional data relevant to this review
Ali 2008	Relates to Clemens 1988 Bangladesh. No additional data relevant to this review
Anonymous 1968	Injectable vaccine
Anonymous 1973a	Injectable vaccine
Anonymous 1973b	Injectable vaccine
Azurin 1967	Injectable cholera vaccine
Azurin 1971	Injectable cholera vaccine
Benenson 1968a	Injectable cholera vaccine
Benenson 1968b	Injectable cholera vaccine
Bergquist 1997	Intranasal vaccination (abstract)
Black 1986	Non-randomized study
Black 1987	Non-randomized study
Burgasov 1976	Injectable cholera vaccine
Bwanga 1984	Not randomized
Cash 1974	Non-randomized study
Cavailler 2006	Non-randomized study
Chaicumpa 1998	Non-randomized, immunological outcomes only
Chongsa-nguan 1988	Safety data presented but no trials assess the efficacy of this type of vaccine (lipopolysaccharide)
Chongsa-nguan 1991	Compared two new vaccine candidates for which efficacy data is not currently available
Ciznar 1989	Not a human study (abstract)
Clemens 1986	Immunological outcomes only
Clemens 1988	Relates to Clemens 1988 Bangladesh. No additional data relevant to this review
Clemens 1989a	Relates to Clemens 1988 Bangladesh. No additional data relevant to this review

(Continued)

Clemens 1989b	Relates to Clemens 1988 Bangladesh. No additional data relevant to this review
Clemens 1990	Relates to Clemens 1988 Bangladesh. No additional data relevant to this review
Clemens 1991	Relates to Clemens 1988 Bangladesh. No additional data relevant to this review
Clemens 1992a	Refers to Clemens 1988, no new data relevant to this review
Clemens 1992b	Relates to Clemens 1988 Bangladesh. No additional data relevant to this review
Clemens 1995	Relates to Clemens 1988 Bangladesh. No additional data relevant to this review
Cohen 1999	No vaccine was given (abstract)
Cohen 2000	Non-cholera vaccine, no cholera outcomes
Cooper 2000	Non-comparative study, all participants received the same vaccine
Cooper 2001	Non-comparative study, all participants received the same vaccine
Coster 1995	The paper contains two very small studies. Study 1 is excluded as it has no placebo group. Study 2 is excluded as it was not randomized
Cryz 1992	No control group.
Cryz 1995	No efficacy data for this vaccine (CVD 103-HgR-Ty21a)
Das 1967	Injectable cholera vaccine
Dearlove 1992	Non-cholera vaccine (abstract)
Durham 1998	Relates to Clemens 1988 Bangladesh. No additional data relevant to this review
Emch 2006	Relates to Clemens 1988 Bangladesh. No additional data relevant to this review
Emch 2007	Relates to Clemens 1988 Bangladesh. No additional data relevant to this review
Emch 2009	Relates to Clemens 1988 Bangladesh. No additional data relevant to this review
Forrest 1991	A new vaccine candidate for which efficacy data is not currently available
Ganguly 1975	Immunological outcomes only
Gateff 1975	Injectable cholera vaccine
Glass 1989	Immunological outcomes only

(Continued)

Glenn 2007	Non-cholera vaccine, no cholera outcomes (abstract)
Graves 2000	A Cochrane Review (abstract)
Gray 1989	Not relevant (abstract)
Gupta 1998	Injectable cholera vaccine
Hall 2001	Non-cholera vaccine, no cholera outcomes (abstract)
Holmgren 1989	Relates to Clemens 1988 Bangladesh. No additional data relevant to this review
Holmgren 1992	Relates to Clemens 1988 Bangladesh. No additional data relevant to this review
Hotomi 1998	Intranasal vaccination (abstract)
Islam 2008	Not an RCT; a willingness to pay study
Jertborn 1984	Not randomized
Jertborn 1986	Retrospective, non-comparative study.
Jertborn 1988	Retrospective study.
Jertborn 1992	This small study (41 participants) compared the safety and immunogenicity of the WC-BS and WC-rBS vaccines. As there was no placebo group we could not include the data
Jertborn 1993	Immunogenicity data only
Jertborn 1994	Not randomized
Jertborn 1996	Non-randomised study.
Jertborn 1998	Non-cholera vaccine, no cholera outcomes
Jertborn 2001	Not cholera vaccine
Johansson 2001	Nasal and intravaginal vaccination (abstract)
Johansson 2004	Nasal and intravaginal vaccination (abstract)
Jones 2004	A summary of included studies
Karlsen 2003	All participants received the same vaccine (abstract)
Kenner 1995	Not randomized

(Continued)

Kilhamn 1998	All participants received the same vaccine (abstract)
Kim 2008	A willingness to pay study (abstract)
Kirk 2005	A case-control study (abstract)
Koenig 1998	Non-cholera vaccine, no cholera outcomes (abstract)
Kollaritsch 1996	No efficacy data for this vaccine (CVD 103-HgR-Ty21a)
Kollaritsch 1997	All participants received the same vaccine (abstract)
Kozlowski 1999	Not a relevant comparison. Randomized to oral, intranasal and vaginal vaccination. (abstract)
Langevin-Perriat 1988	Immunological data only
Lastre 2002	Immunological outcomes only
Lelikov 1974	Injectable cholera vaccine
Levine 1984	Not randomized
Levine 1988a	Not randomized
Levine 1988b	Not randomized
Lewis 1993	Not randomized (on abstract)
Leyten 2005	No relevant outcomes
Lopez 2008	Not an RCT. A review (abstract)
Losonsky 1993	Not randomized
Losonsky 1996	Not randomized
Lucas 2005	A case-control study
Lucas 2007	A willingness to pay study (abstract)
Mahalanabis 2009	No efficacy data for this vaccine (VA1.3)
Martell 2009	Non-cholera vaccine, no cholera outcomes
María García 2005	This paper describes multiple small comparative studies (9 volunteers in each treatment arm) with different modifications and dosing of potential vaccine strains including VC638. We were unable to incorporate any of this data

(Continued)

McCormack 1969	Injectable vaccine
Migasena 1988	Non-randomized study
Migasena 1989b	Contains only safety and immunogenicity data. Excluded as no group received placebo
Migasena 1989c	No efficacy study available for these vaccines
Mitra 1990	No cholera vaccine was given. (Abstract)
Mosley 1968	Injectable vaccine
Mosley 1969a	Injectable vaccine
Mosley 1969b	Injectable vaccine
Mosley 1970	Injectable vaccine
Mosley 1972	Injectable vaccine
Mosley 1973	Injectable vaccine
Nimbkar 1975	Immunological outcomes only
Olsson 2006	Not an RCT (abstract)
Oseasohn 1965	Injectable vaccine
Paineau 2008	No vaccine given (abstract)
Pal 1980	Injectable vaccine
Peltola 1977	Intracutaneous vaccine (abstract)
Peltola 1989	Trial assesses oral cholera vaccine for preventing travellers diarrhoea, not cholera
Peltola 1991	No cholera outcomes relevant to this review
Philippines 1965	Injectable cholera vaccine
Pitisuttithum 2001	No cholera vaccine was given. A study to validate an artificial cholera challenge model
Qadri 2003	Not cholera vaccine
Qadri 2004	All participants received the same vaccine
Qadri 2006	Non-cholera vaccine, no cholera outcomes (abstract)

(Continued)

Quiding-Jarbrink 2001	All participants received the same vaccine
Rao 2002	Not an RCT (Abstract)
Rudin 1998	Not an appropriate comparison group; oral versus nasal vaccination (abstract)
Rudin 1999	Not an appropriate comparison group; oral versus nasal vaccination (abstract)
Sack 1991	Relates to Clemens 1988 Bangladesh. No additional data relevant to this review
Sack 2007	Refers to Clemens 1988, no new data relevant to this review
Sanchez 1993b	Data on adverse effects not presented in usable form.
Sanchez 1994	A preliminary report from Taylor 2000 Peru. Contains no additional data
Saroso 1978	Injectable cholera vaccine
Savarino 1998	Non-cholera vaccine, no cholera outcomes
Savarino 1999	Non-cholera vaccine, no cholera outcomes
Savarino 2002	Non-cholera vaccine, no cholera outcomes
Sommer 1973	Randomized controlled study, but vaccine given after exposure to cholera in family members
SonLa 2007	Non-comparative study
Stellfeld 2004	A review article. Not an RCT
Sumarokov 1974	Injectable vaccine (abstract)
Sumarokov 1978	No clinical efficacy data is available for this vaccine. Oral tablet containing cholera toxin, Ogawa and Inaba O-antigens and <i>Vibrio cholerae</i> exoenzymes.
Sumarokov 1990	No clinical efficacy data is available for this vaccine. Oral tablet containing cholera toxin, Ogawa and Inaba O-antigens and <i>Vibrio cholerae</i> exoenzymes.
Sumarokov 1991	No clinical efficacy data is available for this vaccine. Oral tablet containing cholera toxin, Ogawa and Inaba O-antigens and <i>Vibrio cholerae</i> exoenzymes.
Sumarokov 1993	No clinical efficacy data is available for this vaccine. Oral tablet containing cholera toxin, Ogawa and Inaba O-antigens and <i>Vibrio cholerae</i> exoenzymes.
Suntharasamai 1992	No vaccine given
Svennerholm 1981	No cholera vaccine given (abstract)

(Continued)

Svennerholm 1984	No efficacy data for this vaccine (B subunit alone)
Tacket 1992	Not randomized
Tacket 1995a	No vaccine given
Tacket 1995b	No efficacy data for this vaccine (CVD 112)
Tacket 1998	No efficacy data for this vaccine (CVD 112)
Taylor 1994	No efficacy data for these vaccines (Peru 14, Peru 5, Peru 3)
Taylor 1997	No efficacy data for this vaccine (CVD 103-HgR/CVD112)
Taylor 1999b	No efficacy data for this vaccine (CVD 103-HgR/CVD112)
Thiem 2006	Case control study
Von Seidlein 2007	A study of a fingerprint recognition system used during a cholera vaccine trial. Contains no relevant outcomes
Wassen 2005	Vaginal vaccination (abstract)
Wassen 2006	Vaginal vaccination (abstract)
Wasserman 1993	Immunogenicity data only
Wassén 1996	Intravaginal vaccination (abstract)
Wiedermann 2000	Non cholera vaccine, no cholera outcomes

DATA AND ANALYSES

Comparison 1. Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Cases of cholera - 1st year of follow up (with meta analysis)	4	249935	Risk Ratio (M-H, Random, 95% CI)	0.48 [0.35, 0.65]
1.1 Whole cell vaccine (WC); 3 doses, 6 weeks apart	1	31162	Risk Ratio (M-H, Random, 95% CI)	0.47 [0.33, 0.69]
1.2 Whole cell plus B subunit vaccine (WC-BS); 3 doses, 6 weeks apart	1	31124	Risk Ratio (M-H, Random, 95% CI)	0.38 [0.25, 0.56]
1.3 Whole cell plus recombinant B subunit vaccine (WC-rBS); 2 doses, 2 weeks apart	1	17799	Risk Ratio (M-H, Random, 95% CI)	1.04 [0.52, 2.05]
1.4 Variant whole cell vaccine (vWC); 2 doses, 2 weeks apart	1	111928	Risk Ratio (M-H, Random, 95% CI)	0.36 [0.23, 0.56]
1.5 Bivalent whole cell vaccine (BivWC); 2 doses, 2 weeks apart	1	57922	Risk Ratio (M-H, Random, 95% CI)	0.55 [0.26, 1.17]
2 Cases of cholera - 2nd year of follow up (with meta analysis)	3	130334	Risk Ratio (M-H, Random, 95% CI)	0.39 [0.30, 0.50]
2.1 Whole cell vaccine (WC); 3 doses, 6 weeks apart	1	30011	Risk Ratio (M-H, Random, 95% CI)	0.43 [0.28, 0.65]
2.2 Whole cell plus B subunit vaccine (WC-BS); 3 doses, 6 weeks apart	1	30008	Risk Ratio (M-H, Random, 95% CI)	0.42 [0.28, 0.63]
2.3 Whole cell plus recombinant B subunit vaccine (WC-rBS); 2 doses, 2 weeks apart plus booster at 10 months	1	14997	Risk Ratio (M-H, Random, 95% CI)	0.40 [0.21, 0.75]
2.4 Variant whole cell vaccine (vWC); 2 doses, 2 weeks apart	0	0	Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
2.5 Bivalent whole cell vaccine (BivWC); 2 doses, 2 weeks apart	1	55318	Risk Ratio (M-H, Random, 95% CI)	0.22 [0.11, 0.48]
3 Cases of cholera - 3rd year of follow up (with meta analysis)	1	58174	Risk Ratio (M-H, Random, 95% CI)	0.70 [0.50, 0.98]
3.1 Whole cell vaccine (WC); 3 doses, 6 weeks apart	1	29114	Risk Ratio (M-H, Random, 95% CI)	0.59 [0.36, 0.97]
3.2 Whole cell plus B subunit vaccine (WC-BS); 3 doses, 6 weeks apart	1	29060	Risk Ratio (M-H, Random, 95% CI)	0.81 [0.51, 1.29]

3.3 Whole cell plus recombinant B subunit vaccine (WC-rBS); 2 doses, 2 weeks apart plus booster at 10 months	0	0	Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
3.4 Variant whole cell vaccine (vWC); 2 doses, 2 weeks apart	0	0	Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
3.5 Bivalent whole cell vaccine (BivWC); 2 doses, 2 weeks apart	0	0	Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
4 Cases of cholera - 4th year of follow up (with meta analysis)	1	56613	Risk Ratio (M-H, Random, 95% CI)	1.05 [0.60, 1.84]
4.1 Whole cell vaccine (WC); 3 doses, 6 weeks apart	1	28357	Risk Ratio (M-H, Random, 95% CI)	1.28 [0.59, 2.76]
4.2 Whole cell plus B subunit vaccine (WC-BS); 3 doses, 6 weeks apart	1	28256	Risk Ratio (M-H, Random, 95% CI)	0.84 [0.37, 1.91]
4.3 Whole cell plus recombinant B subunit vaccine (WC-rBS); 2 doses, 2 weeks apart plus booster at 10 months	0	0	Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
4.4 Variant whole cell vaccine (vWC); 2 doses, 2 weeks apart	0	0	Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
4.5 Bivalent whole cell vaccine (BivWC); 2 doses, 2 weeks apart	0	0	Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
5 Cases of cholera by age group - First two years of follow-up	4	243071	Risk Ratio (M-H, Random, 95% CI)	0.43 [0.33, 0.56]
5.1 Age < 5 years	4	29005	Risk Ratio (M-H, Random, 95% CI)	0.62 [0.47, 0.80]
5.2 Age > 5 years	4	214066	Risk Ratio (M-H, Random, 95% CI)	0.34 [0.27, 0.43]
6 Cases of cholera by age group - First two years of follow-up (sensitivity analysis)	4	248140	Risk Ratio (M-H, Random, 95% CI)	0.43 [0.33, 0.56]
6.1 Age < 5 years	4	29773	Risk Ratio (M-H, Random, 95% CI)	0.62 [0.47, 0.80]
6.2 Age > 5 years	4	218367	Risk Ratio (M-H, Random, 95% CI)	0.34 [0.27, 0.43]

Comparison 2. Whole cell vaccine (WC) versus placebo - Subgroup analysis

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Cases of cholera by time of follow-up (3-dose recipients)	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
1.1 First four months after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.2 First year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.3 Second year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.4 Third year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

1.5 Fourth year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2 Cases of cholera by age-group - 1st year of follow-up (3-dose recipients)	1	41580	Risk Ratio (M-H, Random, 95% CI)	0.48 [0.23, 0.98]
2.1 Age 2 to 5 years	1	7730	Risk Ratio (M-H, Random, 95% CI)	0.69 [0.43, 1.09]
2.2 Age > 5 years	1	33850	Risk Ratio (M-H, Random, 95% CI)	0.33 [0.20, 0.54]
3 Cases of cholera by age-group - 2nd year of follow-up (3-dose recipients)	1	40017	Risk Ratio (M-H, Random, 95% CI)	0.45 [0.16, 1.25]
3.1 Age 2 to 5 years	1	7419	Risk Ratio (M-H, Random, 95% CI)	0.76 [0.45, 1.29]
3.2 Age > 5 years	1	32598	Risk Ratio (M-H, Random, 95% CI)	0.27 [0.16, 0.45]
4 Cases of cholera by blood group - First 2 years of follow-up (3-dose recipients)	1	41580	Risk Ratio (M-H, Fixed, 95% CI)	0.45 [0.36, 0.58]
4.1 Blood Group O	1	13465	Risk Ratio (M-H, Fixed, 95% CI)	0.53 [0.37, 0.76]
4.2 All other blood groups	1	28115	Risk Ratio (M-H, Fixed, 95% CI)	0.41 [0.29, 0.56]
5 Cases of all cause diarrhoea - 1st year of follow-up (3-dose recipients)	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
5.1 Severe watery diarrhoea	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
5.2 Any watery diarrhoea	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
5.3 Any diarrhoea	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
6 Deaths - 1st year of follow-up (3-dose recipients)	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
6.1 All cause deaths	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
6.2 Deaths from non-dysenteric diarrhoea (adult females only)	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Comparison 3. Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Cases of cholera by time of follow-up (3-dose recipients)	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
1.1 First four months after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.2 First year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.3 Second year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.4 Third year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.5 Fourth year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2 Cases of cholera by age-group - 1st year of follow-up (3-dose recipients)	1	41542	Risk Ratio (M-H, Random, 95% CI)	0.38 [0.14, 1.03]

2.1 Age 2 to 5 years	1	7558	Risk Ratio (M-H, Random, 95% CI)	0.62 [0.38, 1.01]
2.2 Age > 5 years	1	33984	Risk Ratio (M-H, Random, 95% CI)	0.22 [0.13, 0.39]
3 Cases of cholera by age-group - 2nd year of follow-up (3-dose recipients)	1	40014	Risk Ratio (M-H, Fixed, 95% CI)	0.42 [0.29, 0.60]
3.1 Age 2 to 5 years	1	7273	Risk Ratio (M-H, Fixed, 95% CI)	0.53 [0.29, 0.96]
3.2 Age > 5 years	1	32741	Risk Ratio (M-H, Fixed, 95% CI)	0.37 [0.23, 0.59]
4 Cases of cholera by blood group - First 2 years of follow-up (3-dose recipients)	1	41542	Risk Ratio (M-H, Fixed, 95% CI)	0.40 [0.31, 0.51]
4.1 Blood Group O	1	13453	Risk Ratio (M-H, Fixed, 95% CI)	0.48 [0.33, 0.70]
4.2 All other blood groups	1	28089	Risk Ratio (M-H, Fixed, 95% CI)	0.34 [0.24, 0.48]
5 Cases of all cause diarrhoea - 1st year of follow-up (3-dose recipients)	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
5.1 Severe watery diarrhoea	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
5.2 Any watery diarrhoea	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
5.3 Any diarrhoea	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
6 Deaths - 1st year of follow-up (3-dose recipients)	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
6.1 All cause deaths	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
6.2 Deaths from non-dysenteric diarrhoea (adult females only)	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Comparison 4. Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Cases of confirmed cholera by time of follow-up (3-dose recipients)	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
1.1 First four months after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.2 Four to eight months after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.3 First year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.4 Second year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.5 Third year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.6 Fourth year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2 Cases of cholera by age-group - 1st year of follow-up (3-dose recipients)	1	82085	Risk Ratio (M-H, Fixed, 95% CI)	1.12 [0.80, 1.55]
2.1 Age 2 to 5 years	1	7628	Risk Ratio (M-H, Fixed, 95% CI)	1.10 [0.65, 1.86]
2.2 Age > 5 years	1	33820	Risk Ratio (M-H, Fixed, 95% CI)	1.48 [0.77, 2.85]

2.3 Eight to 12 months after vaccination	1	40637	Risk Ratio (M-H, Fixed, 95% CI)	0.92 [0.53, 1.60]
3 Cases of cholera by age-group - 2nd year of follow-up (3-dose recipients)	1	40007	Risk Ratio (M-H, Fixed, 95% CI)	1.01 [0.66, 1.55]
3.1 Age 2 to 5 years	1	7344	Risk Ratio (M-H, Fixed, 95% CI)	1.44 [0.77, 2.71]
3.2 Age > 5 years	1	32663	Risk Ratio (M-H, Fixed, 95% CI)	0.73 [0.40, 1.33]
4 Cases of cholera by blood group, First 2 years of follow-up (3-dose recipients)	1	41448	Risk Ratio (M-H, Fixed, 95% CI)	1.14 [0.85, 1.54]
4.1 Blood group O	1	13422	Risk Ratio (M-H, Fixed, 95% CI)	1.10 [0.72, 1.68]
4.2 Any other blood group	1	28026	Risk Ratio (M-H, Fixed, 95% CI)	1.19 [0.79, 1.79]
5 Cases of all cause diarrhoea - 1st year of follow-up (3-dose recipients)	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
5.1 Severe watery diarrhoea	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
5.2 Any watery diarrhoea	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
5.3 Any diarrhoea	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
6 Deaths - 1st year of follow-up (3-dose recipients)	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
6.1 All cause deaths	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
6.2 Deaths from non-dysenteric diarrhoea (adult females only)	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Comparison 5. Whole cell plus recombinant B subunit vaccine (WC-rBS) versus placebo - Subgroup analysis

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Cases of cholera by age group - 1st year of follow-up (2 doses)	1	17799	Risk Ratio (M-H, Fixed, 95% CI)	1.04 [0.52, 2.05]
1.1 Age 2 to 5 years	1	2368	Risk Ratio (M-H, Fixed, 95% CI)	3.91 [0.44, 34.90]
1.2 Age 6 to 15 years	1	6782	Risk Ratio (M-H, Fixed, 95% CI)	0.70 [0.22, 2.19]
1.3 Age 16 to 65 years	1	8649	Risk Ratio (M-H, Fixed, 95% CI)	0.98 [0.37, 2.60]
2 Cases of cholera by age group - 2nd year of follow-up (2 doses plus booster)	1	14997	Risk Ratio (M-H, Fixed, 95% CI)	0.40 [0.21, 0.75]
2.1 Age 2 to 5 years	1	2040	Risk Ratio (M-H, Fixed, 95% CI)	0.48 [0.09, 2.62]
2.2 Age 6 to 15 years	1	6049	Risk Ratio (M-H, Fixed, 95% CI)	0.53 [0.20, 1.44]
2.3 Age 16 to 65 years	1	6908	Risk Ratio (M-H, Fixed, 95% CI)	0.29 [0.11, 0.78]
3 Cases of cholera in military recruits, 4 to 18 weeks follow-up	2		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
3.1 Cases of cholera - Occurring after the second dose	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3.2 Cases of cholera - Occurring between the first and second dose	2		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Comparison 6. Whole cell vaccines (all types) versus placebo - Safety outcomes (first dose)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Adverse events - Whole cell (WC) versus placebo	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Abdominal pain	1	613	Risk Ratio (M-H, Fixed, 95% CI)	1.17 [0.80, 1.70]
1.2 Severe abdominal pain	1	613	Risk Ratio (M-H, Fixed, 95% CI)	0.51 [0.09, 2.77]
1.3 Diarrhoea	1	613	Risk Ratio (M-H, Fixed, 95% CI)	1.50 [0.95, 2.36]
1.4 Watery diarrhoea	1	613	Risk Ratio (M-H, Fixed, 95% CI)	1.46 [0.75, 2.84]
1.5 Subjective fever	1	613	Risk Ratio (M-H, Fixed, 95% CI)	1.77 [0.86, 3.65]
1.6 Nausea	1	613	Risk Ratio (M-H, Fixed, 95% CI)	1.92 [0.83, 4.46]
1.7 Vomiting	1	613	Risk Ratio (M-H, Fixed, 95% CI)	2.39 [0.62, 9.15]
1.8 Other symptoms requiring bedrest	1	613	Risk Ratio (M-H, Fixed, 95% CI)	1.02 [0.06, 16.28]
2 Adverse events - Whole cell plus B subunit (WC-BS) versus placebo	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
2.1 Abdominal pain or stomach cramps	1	631	Risk Ratio (M-H, Fixed, 95% CI)	1.17 [0.80, 1.70]
2.2 Severe abdominal pain	1	631	Risk Ratio (M-H, Fixed, 95% CI)	0.48 [0.09, 2.62]
2.3 Diarrhoea	1	631	Risk Ratio (M-H, Fixed, 95% CI)	1.35 [0.85, 2.13]
2.4 Watery diarrhoea	1	631	Risk Ratio (M-H, Fixed, 95% CI)	1.31 [0.67, 2.57]
2.5 Subjective fever	1	631	Risk Ratio (M-H, Fixed, 95% CI)	1.14 [0.52, 2.51]
2.6 Nausea	1	631	Risk Ratio (M-H, Fixed, 95% CI)	1.45 [0.60, 3.50]
2.7 Vomiting	1	624	Risk Ratio (M-H, Fixed, 95% CI)	1.21 [0.46, 3.22]
2.8 Other symptoms requiring bedrest	1	631	Risk Ratio (M-H, Fixed, 95% CI)	0.97 [0.06, 15.37]
3 Adverse events - Whole cell plus recombinant B subunit (WC-rBS) versus placebo	7		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
3.1 Abdominal pain or stomach cramps	6	2878	Risk Ratio (M-H, Random, 95% CI)	1.11 [0.70, 1.74]
3.2 Stomach gurgling	3	1219	Risk Ratio (M-H, Random, 95% CI)	1.16 [0.90, 1.49]
3.3 Diarrhoea	7	23870	Risk Ratio (M-H, Random, 95% CI)	1.04 [0.73, 1.49]
3.4 Fever	4	941	Risk Ratio (M-H, Random, 95% CI)	0.31 [0.08, 1.26]
3.5 Nausea	4	2213	Risk Ratio (M-H, Random, 95% CI)	1.41 [0.32, 6.13]
3.6 Vomiting	4	2049	Risk Ratio (M-H, Random, 95% CI)	1.46 [0.40, 5.33]
3.7 Headache	4	2488	Risk Ratio (M-H, Random, 95% CI)	0.72 [0.37, 1.40]
3.8 Loss of appetite	2	390	Risk Ratio (M-H, Random, 95% CI)	0.74 [0.17, 3.18]
3.9 Dizziness	1	1313	Risk Ratio (M-H, Random, 95% CI)	1.04 [0.41, 2.69]
3.10 Any adverse event	2	21616	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.71, 1.28]
3.11 Any serious adverse event	2	21133	Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
3.12 Other	1	624	Risk Ratio (M-H, Random, 95% CI)	0.89 [0.35, 2.29]
4 Adverse events - Bivalent whole cell (BivWC) versus placebo	4		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
4.1 Diarrhoea	4	67414	Risk Ratio (M-H, Fixed, 95% CI)	0.80 [0.42, 1.55]
4.2 Abdo pain	4	67414	Risk Ratio (M-H, Fixed, 95% CI)	1.09 [0.63, 1.88]
4.3 Gas	1	160	Risk Ratio (M-H, Fixed, 95% CI)	0.65 [0.19, 2.22]
4.4 Loss of appetite	3	514	Risk Ratio (M-H, Fixed, 95% CI)	1.20 [0.35, 4.13]

4.5 Nausea	4	67414	Risk Ratio (M-H, Fixed, 95% CI)	0.80 [0.38, 1.67]
4.6 Vomiting	4	67414	Risk Ratio (M-H, Fixed, 95% CI)	1.12 [0.57, 2.21]
4.7 Fever	4	67414	Risk Ratio (M-H, Fixed, 95% CI)	1.62 [0.84, 3.10]
4.8 Headache	3	514	Risk Ratio (M-H, Fixed, 95% CI)	0.98 [0.55, 1.75]
4.9 General ill feeling	3	514	Risk Ratio (M-H, Fixed, 95% CI)	1.70 [0.61, 4.77]
4.10 Rash	1	66900	Risk Ratio (M-H, Fixed, 95% CI)	1.64 [0.27, 9.83]
4.11 Weakness	1	66900	Risk Ratio (M-H, Fixed, 95% CI)	0.27 [0.03, 2.45]
4.12 Itch	1	66900	Risk Ratio (M-H, Fixed, 95% CI)	3.29 [0.34, 31.58]
4.13 Cough	1	66900	Risk Ratio (M-H, Fixed, 95% CI)	1.10 [0.15, 7.77]
4.14 Dizziness	1	66900	Risk Ratio (M-H, Fixed, 95% CI)	1.10 [0.07, 17.51]

Comparison 7. Whole cell vaccines (all types) versus placebo - Safety outcomes (second dose)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Adverse events - Whole cell plus recombinant B subunit (WC-rBS) versus placebo	5		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Abdominal pain or stomach cramps	5	2061	Risk Ratio (M-H, Fixed, 95% CI)	0.95 [0.69, 1.32]
1.2 Stomach gurgling	2	556	Risk Ratio (M-H, Fixed, 95% CI)	7.82 [4.36, 14.03]
1.3 Diarrhoea	5	2061	Risk Ratio (M-H, Fixed, 95% CI)	1.20 [0.70, 2.05]
1.4 Fever	4	896	Risk Ratio (M-H, Fixed, 95% CI)	1.19 [0.34, 4.20]
1.5 Nausea	3	1438	Risk Ratio (M-H, Fixed, 95% CI)	1.04 [0.26, 4.06]
1.6 Vomiting	4	1859	Risk Ratio (M-H, Fixed, 95% CI)	2.13 [0.54, 8.44]
1.7 Headache	3	1674	Risk Ratio (M-H, Fixed, 95% CI)	1.26 [0.79, 1.99]
1.8 Loss of appetite	2	387	Risk Ratio (M-H, Fixed, 95% CI)	6.80 [0.36, 129.27]
1.9 Dizziness	1	1165	Risk Ratio (M-H, Fixed, 95% CI)	4.79 [0.54, 42.75]
1.10 Any adverse event	0	0	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.11 Any serious adverse event	0	0	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.12 Other	0	0	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2 Adverse events - Bivalent whole cell (BivWC) versus placebo	4		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
2.1 Diarrhoea	4	67397	Risk Ratio (M-H, Fixed, 95% CI)	1.17 [0.53, 2.57]
2.2 Abdo pain	4	67397	Risk Ratio (M-H, Fixed, 95% CI)	1.41 [0.64, 3.12]
2.3 Gas	1	155	Risk Ratio (M-H, Fixed, 95% CI)	0.68 [0.12, 3.93]
2.4 Loss of appetite	3	497	Risk Ratio (M-H, Fixed, 95% CI)	9.12 [0.50, 166.49]
2.5 Nausea	4	67397	Risk Ratio (M-H, Fixed, 95% CI)	5.08 [1.12, 22.92]
2.6 Vomiting	4	67397	Risk Ratio (M-H, Fixed, 95% CI)	0.91 [0.41, 2.01]
2.7 Fever	4	67397	Risk Ratio (M-H, Fixed, 95% CI)	0.97 [0.46, 2.04]
2.8 Headache	3	497	Risk Ratio (M-H, Fixed, 95% CI)	1.24 [0.59, 2.62]
2.9 General ill feeling	3	497	Risk Ratio (M-H, Fixed, 95% CI)	1.22 [0.36, 4.15]
2.10 Rash	1	66900	Risk Ratio (M-H, Fixed, 95% CI)	0.37 [0.01, 8.96]
2.11 Weakness	1	66900	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.12 Itch	1	66900	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.13 Cough	1	66900	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.14 Dizziness	1	66900	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Comparison 8. Live attenuated vaccines (all types) versus placebo - Efficacy outcomes

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Cases of cholera following natural infection - CVD 103HgR versus placebo	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
1.1 First year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.2 Second year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.3 Third year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2 Severe cholera following natural infection - CVD 103HgR versus placebo	1	67508	Risk Ratio (M-H, Fixed, 95% CI)	1.58 [0.61, 4.07]
3 Death from any cause (except motor accidents)	1	67508	Risk Ratio (M-H, Fixed, 95% CI)	1.03 [0.83, 1.28]
4 Death from diarrhoea (any organism)	1	67508	Risk Ratio (M-H, Fixed, 95% CI)	0.75 [0.26, 2.17]
5 Cases of moderate to severe diarrhoea - following artificial challenge	3	108	Risk Ratio (M-H, Fixed, 95% CI)	0.08 [0.02, 0.34]
5.1 CVD 103HgR	1	51	Risk Ratio (M-H, Fixed, 95% CI)	0.09 [0.01, 0.67]
5.2 Peru 15	1	36	Risk Ratio (M-H, Fixed, 95% CI)	0.05 [0.00, 0.79]
5.3 VC638	1	21	Risk Ratio (M-H, Fixed, 95% CI)	0.15 [0.01, 2.86]
6 Cases of any diarrhoea -following artificial challenge	3	108	Risk Ratio (M-H, Fixed, 95% CI)	0.14 [0.07, 0.28]
6.1 CVD 103HgR	1	51	Risk Ratio (M-H, Fixed, 95% CI)	0.20 [0.09, 0.44]
6.2 Peru 15	1	36	Risk Ratio (M-H, Fixed, 95% CI)	0.07 [0.01, 0.52]
6.3 VC638	1	21	Risk Ratio (M-H, Fixed, 95% CI)	0.05 [0.00, 0.80]

Comparison 9. Live attenuated vaccines (all types) versus placebo - Safety outcomes

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Adverse events - CVD 103-HgR versus placebo	12		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Diarrhoea	12	3320	Risk Ratio (M-H, Random, 95% CI)	1.09 [0.81, 1.47]
1.2 Fever	8	2516	Risk Ratio (M-H, Random, 95% CI)	0.84 [0.57, 1.23]
1.3 Vomiting	9	2866	Risk Ratio (M-H, Random, 95% CI)	1.22 [0.84, 1.79]
1.4 Nausea	3	1474	Risk Ratio (M-H, Random, 95% CI)	1.10 [0.67, 1.80]
1.5 Seizure	1	1077	Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
1.6 Itching	1	1077	Risk Ratio (M-H, Random, 95% CI)	2.00 [0.61, 6.61]
1.7 Rash	3	1489	Risk Ratio (M-H, Random, 95% CI)	0.94 [0.26, 3.49]
1.8 Abdominal pain	7	2155	Risk Ratio (M-H, Random, 95% CI)	1.12 [0.86, 1.46]
1.9 Headache	3	1243	Risk Ratio (M-H, Random, 95% CI)	1.20 [0.91, 1.58]

1.10 Anorexia	3	478	Risk Ratio (M-H, Random, 95% CI)	1.06 [0.48, 2.36]
1.11 Malaise	2	434	Risk Ratio (M-H, Random, 95% CI)	0.88 [0.61, 1.26]
1.12 Borborygmus	1	81	Risk Ratio (M-H, Random, 95% CI)	0.89 [0.59, 1.35]
1.13 Liquid stools	1	81	Risk Ratio (M-H, Random, 95% CI)	1.37 [0.33, 5.72]
2 Adverse events - Peru 15 versus placebo	4		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
2.1 Loss of appetite	1	59	Risk Ratio (M-H, Fixed, 95% CI)	0.63 [0.16, 2.55]
2.2 Loss of energy	1	59	Risk Ratio (M-H, Fixed, 95% CI)	1.43 [0.32, 6.41]
2.3 Abdominal cramps	3	369	Risk Ratio (M-H, Fixed, 95% CI)	2.92 [0.62, 13.82]
2.4 Headache	3	349	Risk Ratio (M-H, Fixed, 95% CI)	4.14 [1.27, 13.48]
2.5 Vomiting	2	299	Risk Ratio (M-H, Fixed, 95% CI)	5.01 [0.26, 96.01]
2.6 Nausea	1	59	Risk Ratio (M-H, Fixed, 95% CI)	1.66 [0.38, 7.26]
2.7 Diarrhoea	3	369	Risk Ratio (M-H, Fixed, 95% CI)	2.44 [0.12, 48.45]
2.8 Gas	1	70	Risk Ratio (M-H, Fixed, 95% CI)	2.27 [0.10, 53.81]
2.9 Fever	2	310	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.10 Respiratory symptoms	1	50	Risk Ratio (M-H, Fixed, 95% CI)	0.94 [0.25, 3.47]
2.11 Gastrointestinal symptoms	1	50	Risk Ratio (M-H, Fixed, 95% CI)	1.5 [0.72, 3.14]
3 Adverse events - VC638 versus placebo	3		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
3.1 Abdominal pain	3	137	Risk Ratio (M-H, Fixed, 95% CI)	1.94 [0.94, 4.02]
3.2 Nausea	1	36	Risk Ratio (M-H, Fixed, 95% CI)	4.0 [0.56, 28.40]
3.3 Diarrhoea	3	137	Risk Ratio (M-H, Fixed, 95% CI)	2.05 [0.65, 6.48]
3.4 Headache	3	137	Risk Ratio (M-H, Fixed, 95% CI)	2.30 [0.83, 6.36]
3.5 General discomfort	1	36	Risk Ratio (M-H, Fixed, 95% CI)	2.6 [0.13, 50.25]
3.6 Borborygmus	3	137	Risk Ratio (M-H, Fixed, 95% CI)	1.23 [0.77, 1.95]
3.7 Vomiting	2	101	Risk Ratio (M-H, Fixed, 95% CI)	1.05 [0.05, 24.33]
3.8 Fever	2	101	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3.9 Heartburn	1	56	Risk Ratio (M-H, Fixed, 95% CI)	1.0 [0.23, 4.40]
3.10 Malaise	1	56	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

Comparison 10. Live attenuated CVD 103-HgR vaccine versus placebo - Subgroup analysis

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Cases of cholera by age group (age 2-5 years)	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
1.1 First year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.2 Second year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
1.3 Third year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2 Cases of confirmed cholera attending healthcare facilities (age over 5 years)	1		Risk Ratio (M-H, Fixed, 95% CI)	Totals not selected
2.1 First year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2.2 Second year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]

2.3 Third year after vaccination	1		Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
3 Cases of cholera within four years and five months, by blood group	1	67508	Risk Ratio (M-H, Fixed, 95% CI)	0.82 [0.54, 1.24]
3.1 Blood group O	1	24303	Risk Ratio (M-H, Fixed, 95% CI)	0.60 [0.34, 1.08]
3.2 All other blood groups	1	43205	Risk Ratio (M-H, Fixed, 95% CI)	1.15 [0.63, 2.10]
4 Any diarrhoea following artificial challenge, by blood group	1	51	Risk Ratio (M-H, Fixed, 95% CI)	0.17 [0.07, 0.43]
4.1 Blood group O	1	23	Risk Ratio (M-H, Fixed, 95% CI)	0.30 [0.13, 0.73]
4.2 Blood group non-O	1	28	Risk Ratio (M-H, Fixed, 95% CI)	0.08 [0.01, 0.54]
5 Moderate or severe diarrhoea due to <i>V. cholerae</i> after artificial challenge, by blood group	1	51	Risk Ratio (M-H, Fixed, 95% CI)	0.12 [0.02, 0.64]
5.1 Blood group O	1	23	Risk Ratio (M-H, Fixed, 95% CI)	0.13 [0.02, 1.00]
5.2 Blood group non-O	1	28	Risk Ratio (M-H, Fixed, 95% CI)	0.10 [0.01, 1.72]
6 Additional adverse event data			Other data	No numeric data

Analysis 1.1. Comparison 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes, Outcome 1 Cases of cholera - 1st year of follow up (with meta analysis).

Review: Oral vaccines for preventing cholera

Comparison: 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes

Outcome: 1 Cases of cholera - 1st year of follow up (with meta analysis)

(1) For the purpose of this meta analysis the events and participants in the Placebo arm of Clemens 1985 Bangladesh have been divided equally between the two intervention arms

(2) Trach 1997- This result has been adjusted from the crude figures given in the original paper to give the effective sample size; using an estimate of the ICC of -0.063 and a mean cluster size of 5.3

(3) Sur 2009- This result has been adjusted from the crude figures given in the original paper to give the effective sample size; using an ICC of -0.0085 and a mean cluster size of 19.2

Analysis 1.2. Comparison 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes, Outcome 2 Cases of cholera - 2nd year of follow up (with meta analysis).

Review: Oral vaccines for preventing cholera

Comparison: 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes

Outcome: 2 Cases of cholera - 2nd year of follow up (with meta analysis)

(1) For the purpose of this meta analysis the events and participants in the Placebo arm of Clemens 1985 Bangladesh have been divided equally between the two intervention arms

(2) Sur 2009- This result has been adjusted from the crude figures given in the original paper to give the effective sample size; using an ICC of -0.0085 and a mean cluster size of 19.2

Analysis 1.3. Comparison 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes, Outcome 3 Cases of cholera - 3rd year of follow up (with meta analysis).

Review: Oral vaccines for preventing cholera

Comparison: 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes

Outcome: 3 Cases of cholera - 3rd year of follow up (with meta analysis)

(Continued ...)

Analysis 1.4. Comparison 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes, Outcome 4 Cases of cholera - 4th year of follow up (with meta analysis).

Review: Oral vaccines for preventing cholera

Comparison: 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes

Outcome: 4 Cases of cholera - 4th year of follow up (with meta analysis)

(Continued . . .)

Analysis 1.5. Comparison 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes, Outcome 5 Cases of cholera by age group - First two years of follow-up.

Review: Oral vaccines for preventing cholera

Comparison: 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes

Outcome: 5 Cases of cholera by age group - First two years of follow-up

- (1) WC vs Placebo: For the purpose of this meta analysis the events and participants in the Placebo arm of Clemens 1985 Bangladesh have been divided equally between the two intervention arms
- (2) WC-BS vs Placebo: For the purpose of this meta analysis the events and participants in the Placebo arm of Clemens 1985 Bangladesh have been divided equally between the two intervention arms
- (3) Trach 1997- This result has been adjusted from the crude figures given in the original paper to give the effective sample size; using an estimate of the ICC of -0.063 and a mean cluster size of 5.3
- (4) Sur 2009- This result has been adjusted from the crude figures given in the original paper to give the effective sample size; using an ICC of -0.0085 and a mean cluster size of 19.2
- (5) WC vs Placebo: For the purpose of this meta analysis the events and participants in the Placebo arm of Clemens 1985 Bangladesh have been divided equally between the two intervention arms
- (6) WC-BS vs Placebo: For the purpose of this meta analysis the events and participants in the Placebo arm of Clemens 1985 Bangladesh have been divided equally between the two intervention arms
- (7) Trach 1997- This result has been adjusted from the crude figures given in the original paper to give the effective sample size; using an estimate of the ICC of -0.063 and a mean cluster size of 5.3
- (8) Sur 2009- This result has been adjusted from the crude figures given in the original paper to give the effective sample size; using an ICC of -0.0085 and a mean cluster size of 19.2

Analysis 1.6. Comparison 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes, Outcome 6 Cases of cholera by age group - First two years of follow-up (sensitivity analysis).

Review: Oral vaccines for preventing cholera

Comparison: 1 Whole cell vaccines (all types) versus placebo - Primary efficacy outcomes

Outcome: 6 Cases of cholera by age group - First two years of follow-up (sensitivity analysis)

(1) For the purpose of this meta analysis the events and participants in the Placebo arm of Clemens 1985 Bangladesh have been divided equally between the two intervention arms

(2) Trach 1997- This result has been adjusted from the crude figures given in the original paper to give the effective sample size; using an estimate of the ICC of -0.063 and a mean cluster size of 5.3

(3) Sur 2009- This result has been adjusted from the crude figures given in the original paper to give the effective sample size; using an ICC of -0.0085 and a mean cluster size of 19.2

(4) For the purpose of this meta analysis the events and participants in the Placebo arm of Clemens 1985 Bangladesh have been divided equally between the two intervention arms

(5) Trach 1997- This result has been adjusted from the crude figures given in the original paper to give the effective sample size; using an estimate of the ICC of -0.063 and a mean cluster size of 5.3

(6) Sur 2009- This result has been adjusted from the crude figures given in the original paper to give the effective sample size; using an ICC of -0.0085 and a mean cluster size of 19.2

Analysis 2.1. Comparison 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis, Outcome 1 Cases of cholera by time of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis

Outcome: 1 Cases of cholera by time of follow-up (3-dose recipients)

Analysis 2.2. Comparison 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis, Outcome 2 Cases of cholera by age-group - 1st year of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis

Outcome: 2 Cases of cholera by age-group - 1st year of follow-up (3-dose recipients)

Analysis 2.3. Comparison 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis, Outcome 3 Cases of cholera by age-group - 2nd year of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis

Outcome: 3 Cases of cholera by age-group - 2nd year of follow-up (3-dose recipients)

(1) For the purpose of this meta analysis the events and participants in the Placebo arm of Clemens 1985 Bangladesh have been divided equally between the two intervention arms

(2) For the purpose of this meta analysis the events and participants in the Placebo arm of Clemens 1985 Bangladesh have been divided equally between the two intervention arms

Analysis 2.4. Comparison 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis, Outcome 4 Cases of cholera by blood group - First 2 years of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis

Outcome: 4 Cases of cholera by blood group - First 2 years of follow-up (3-dose recipients)

Analysis 2.5. Comparison 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis, Outcome 5 Cases of all cause diarrhoea - 1st year of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis

Outcome: 5 Cases of all cause diarrhoea - 1st year of follow-up (3-dose recipients)

Analysis 2.6. Comparison 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis, Outcome 6 Deaths - 1st year of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 2 Whole cell vaccine (WC) versus placebo - Subgroup analysis

Outcome: 6 Deaths - 1st year of follow-up (3-dose recipients)

Analysis 3.1. Comparison 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis, Outcome 1 Cases of cholera by time of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis

Outcome: 1 Cases of cholera by time of follow-up (3-dose recipients)

Analysis 3.2. Comparison 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis, Outcome 2 Cases of cholera by age-group - 1st year of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis

Outcome: 2 Cases of cholera by age-group - 1st year of follow-up (3-dose recipients)

Analysis 3.3. Comparison 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis, Outcome 3 Cases of cholera by age-group - 2nd year of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis

Outcome: 3 Cases of cholera by age-group - 2nd year of follow-up (3-dose recipients)

(1) For the purpose of this meta analysis the events and participants in the Placebo arm of Clemens 1985 Bangladesh have been divided equally between the two intervention arms

(2) For the purpose of this meta analysis the events and participants in the Placebo arm of Clemens 1985 Bangladesh have been divided equally between the two intervention arms

Analysis 3.4. Comparison 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis, Outcome 4 Cases of cholera by blood group - First 2 years of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis

Outcome: 4 Cases of cholera by blood group - First 2 years of follow-up (3-dose recipients)

Analysis 3.5. Comparison 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis, Outcome 5 Cases of all cause diarrhoea - 1st year of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis

Outcome: 5 Cases of all cause diarrhoea - 1st year of follow-up (3-dose recipients)

Analysis 3.6. Comparison 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis, Outcome 6 Deaths - 1st year of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 3 Whole cell vaccine plus B subunit (WC-BS) versus placebo - Subgroup analysis

Outcome: 6 Deaths - 1st year of follow-up (3-dose recipients)

Analysis 4.1. Comparison 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis, Outcome 1 Cases of confirmed cholera by time of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis

Outcome: 1 Cases of confirmed cholera by time of follow-up (3-dose recipients)

Analysis 4.2. Comparison 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis, Outcome 2 Cases of cholera by age-group - 1st year of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis

Outcome: 2 Cases of cholera by age-group - 1st year of follow-up (3-dose recipients)

(1) For the purpose of this meta analysis the events and participants in the Placebo arm of Clemens 1985 Bangladesh have been divided equally between the two intervention arms

(2) For the purpose of this meta analysis the events and participants in the Placebo arm of Clemens 1985 Bangladesh have been divided equally between the two intervention arms

Analysis 4.3. Comparison 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis, Outcome 3 Cases of cholera by age-group - 2nd year of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis

Outcome: 3 Cases of cholera by age-group - 2nd year of follow-up (3-dose recipients)

(1) For the purpose of this meta analysis the events and participants in the Placebo arm of Clemens 1985 Bangladesh have been divided equally between the two intervention arms

(2) For the purpose of this meta analysis the events and participants in the Placebo arm of Clemens 1985 Bangladesh have been divided equally between the two intervention arms

Analysis 4.4. Comparison 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis, Outcome 4 Cases of cholera by blood group, First 2 years of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis

Outcome: 4 Cases of cholera by blood group, First 2 years of follow-up (3-dose recipients)

Analysis 4.5. Comparison 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis, Outcome 5 Cases of all cause diarrhoea - 1st year of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis

Outcome: 5 Cases of all cause diarrhoea - 1st year of follow-up (3-dose recipients)

Analysis 4.6. Comparison 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis, Outcome 6 Deaths - 1st year of follow-up (3-dose recipients).

Review: Oral vaccines for preventing cholera

Comparison: 4 Whole cell vaccine (WC) versus Whole cell vaccine plus B subunit (WC-BS) - Subgroup analysis

Outcome: 6 Deaths - 1st year of follow-up (3-dose recipients)

Analysis 5.1. Comparison 5 Whole cell plus recombinant B subunit vaccine (WC-rBS) versus placebo - Subgroup analysis, Outcome 1 Cases of cholera by age group - 1st year of follow-up (2 doses).

Review: Oral vaccines for preventing cholera

Comparison: 5 Whole cell plus recombinant B subunit vaccine (WC-rBS) versus placebo - Subgroup analysis

Outcome: 1 Cases of cholera by age group - 1st year of follow-up (2 doses)

Analysis 5.2. Comparison 5 Whole cell plus recombinant B subunit vaccine (WC-rBS) versus placebo - Subgroup analysis, Outcome 2 Cases of cholera by age group - 2nd year of follow-up (2 doses plus booster).

Review: Oral vaccines for preventing cholera

Comparison: 5 Whole cell plus recombinant B subunit vaccine (WC-rBS) versus placebo - Subgroup analysis

Outcome: 2 Cases of cholera by age group - 2nd year of follow-up (2 doses plus booster)

Analysis 5.3. Comparison 5 Whole cell plus recombinant B subunit vaccine (WC-rBS) versus placebo - Subgroup analysis, Outcome 3 Cases of cholera in military recruits, 4 to 18 weeks follow-up.

Review: Oral vaccines for preventing cholera

Comparison: 5 Whole cell plus recombinant B subunit vaccine (WC-rBS) versus placebo - Subgroup analysis

Outcome: 3 Cases of cholera in military recruits, 4 to 18 weeks follow-up

(1) 18 weeks follow-up

Analysis 6.1. Comparison 6 Whole cell vaccines (all types) versus placebo - Safety outcomes (first dose), Outcome 1 Adverse events - Whole cell (WC) versus placebo.

Review: Oral vaccines for preventing cholera

Comparison: 6 Whole cell vaccines (all types) versus placebo - Safety outcomes (first dose)

Outcome: 1 Adverse events - Whole cell (WC) versus placebo

(Continued ...)

Analysis 6.2. Comparison 6 Whole cell vaccines (all types) versus placebo - Safety outcomes (first dose), Outcome 2 Adverse events - Whole cell plus B subunit (WC-BS) versus placebo.

Review: Oral vaccines for preventing cholera

Comparison: 6 Whole cell vaccines (all types) versus placebo - Safety outcomes (first dose)

Outcome: 2 Adverse events - Whole cell plus B subunit (WC-BS) versus placebo

(Continued ...)

Analysis 6.3. Comparison 6 Whole cell vaccines (all types) versus placebo - Safety outcomes (first dose), Outcome 3 Adverse events - Whole cell plus recombinant B subunit (WC-rBS) versus placebo.

Review: Oral vaccines for preventing cholera

Comparison: 6 Whole cell vaccines (all types) versus placebo - Safety outcomes (first dose)

Outcome: 3 Adverse events - Whole cell plus recombinant B subunit (WC-rBS) versus placebo

(Continued ...)

(Continued . . .)

(... Continued)

- (1) Begue 1995: Ages 2 to 65 years, 2 days AE monitoring
- (2) Concha 1995: Ages 1 to 65 years, 3 days AE monitoring
- (3) Hallander 2002: Ages 1 to 12 years, 3 days AE monitoring
- (4) Sanchez 1995: Ages 17 to 23, 24 hours AE monitoring
- (5) Taylor 1999b: Age 2 to 65 years, 3 days AE monitoring after each dose.
- (6) Trach 2002: Ages 1 to 12 years and 17 to 25 years, 3 days AE monitoring
- (7) Taylor 2000 Peru: Ages 2 to 65 years, reporting symptoms during the period between doses and children 14 days (collected at time of second dose)

Analysis 6.4. Comparison 6 Whole cell vaccines (all types) versus placebo - Safety outcomes (first dose), Outcome 4 Adverse events - Bivalent whole cell (BivWC) versus placebo.

Review: Oral vaccines for preventing cholera

Comparison: 6 Whole cell vaccines (all types) versus placebo - Safety outcomes (first dose)

Outcome: 4 Adverse events - Bivalent whole cell (BivWC) versus placebo

(Continued ...)

(Continued . . .)

(Continued . . .)

- (1) Anh 2007: Ages 18 to 40, 3 days of AE monitoring after each dose
- (2) Kanungo 2009: Ages 1 to 40 yrs, 3 days of AE monitoring after each dose
- (3) Mahalanabis 2008: Ages 1 to 40, 3 days of AE monitoring after each dose
- (4) Sur 2009: Age >1 yr; passively reporting symptoms within 14 days of the 1st dose

Analysis 7.1. Comparison 7 Whole cell vaccines (all types) versus placebo - Safety outcomes (second dose), Outcome 1 Adverse events - Whole cell plus recombinant B subunit (WC-rBS) versus placebo.

Review: Oral vaccines for preventing cholera

Comparison: 7 Whole cell vaccines (all types) versus placebo - Safety outcomes (second dose)

Outcome: 1 Adverse events - Whole cell plus recombinant B subunit (WC-rBS) versus placebo

(Continued ...)

(Continued . . .)

Analysis 7.2. Comparison 7 Whole cell vaccines (all types) versus placebo - Safety outcomes (second dose), Outcome 2 Adverse events - Bivalent whole cell (BivWC) versus placebo.

Review: Oral vaccines for preventing cholera

Comparison: 7 Whole cell vaccines (all types) versus placebo - Safety outcomes (second dose)

Outcome: 2 Adverse events - Bivalent whole cell (BivWC) versus placebo

(Continued ...)

(... Continued)

(1) Sur 2009 includes age >1 yr reporting symptoms within 14 days of the 1st dose

Analysis 8.1. Comparison 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes, Outcome 1 Cases of cholera following natural infection - CVD 103HgR versus placebo.

Review: Oral vaccines for preventing cholera

Comparison: 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes

Outcome: 1 Cases of cholera following natural infection - CVD 103HgR versus placebo

Analysis 8.2. Comparison 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes, Outcome 2 Severe cholera following natural infection - CVD 103HgR versus placebo.

Review: Oral vaccines for preventing cholera

Comparison: 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes

Outcome: 2 Severe cholera following natural infection - CVD 103HgR versus placebo

Analysis 8.3. Comparison 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes, Outcome 3 Death from any cause (except motor accidents).

Review: Oral vaccines for preventing cholera

Comparison: 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes

Outcome: 3 Death from any cause (except motor accidents)

Analysis 8.4. Comparison 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes, Outcome 4 Death from diarrhoea (any organism).

Review: Oral vaccines for preventing cholera

Comparison: 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes

Outcome: 4 Death from diarrhoea (any organism)

Analysis 8.5. Comparison 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes, Outcome 5 Cases of moderate to severe diarrhoea - following artificial challenge.

Review: Oral vaccines for preventing cholera

Comparison: 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes

Outcome: 5 Cases of moderate to severe diarrhoea - following artificial challenge

Analysis 8.6. Comparison 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes, Outcome 6 Cases of any diarrhoea -following artificial challenge.

Review: Oral vaccines for preventing cholera

Comparison: 8 Live attenuated vaccines (all types) versus placebo - Efficacy outcomes

Outcome: 6 Cases of any diarrhoea -following artificial challenge

Analysis 9.1. Comparison 9 Live attenuated vaccines (all types) versus placebo - Safety outcomes, Outcome 1 Adverse events - CVD 103-HgR versus placebo.

Review: Oral vaccines for preventing cholera

Comparison: 9 Live attenuated vaccines (all types) versus placebo - Safety outcomes

Outcome: 1 Adverse events - CVD 103-HgR versus placebo

(Continued ...)

- (1) Cryz 1990: Age 21 to 45yrs, 7 days AE monitoring, vaccine dose: 5×10^8
- (2) Gotuzzo 1993: age 18 to 38 years, 7 days AE monitoring, vaccine doses: 5×10^8 , and 5×10^9 were used
- (3) Lagos 1993: Age 18 to 35 years, 7 days AE monitoring, vaccine dose: 5×10^9
- (4) Lagos 1995: Children age 5 to 9 years, 9 days AE monitoring, vaccine dose: 5×10^9
- (5) Lagos 1999: Children age 3 to 17 months, 7 days AE monitoring, vaccine dose: 5×10^9
- (6) Richie 2000: Age 2 to 41 years, 3 days AE monitoring, vaccine dose: 5×10^9
- (7) Simanjuntak 1993: Children age 2 to 5 years, 9 days AE monitoring, vaccine dose: 5×10^9
- (8) Su-Arehawaratana 1992a: Age 18 to 26 years, 7 days AE monitoring, vaccine dose: 5×10^8
- (9) Su-Arehawaratana 1992b: Age 18 to 26 years, 7 days AE monitoring, vaccine doses: 5×10^8 , and 5×10^9
- (10) Suharyono 1992a: Children aged 5 to 9 years, 9 days AE monitoring, vaccine doses: 5×10^6 , 5×10^7 , and 5×10^8
- (11) Suharyono 1992b: Children aged 5 to 9 years, 9 days AE monitoring, vaccine doses: 5×10^9 , and 1×10^{10}
- (12) Tacket 1999: Ages 18 to 40 years, 3 days AE monitoring, vaccine dose: 2.8×10^8

Analysis 9.2. Comparison 9 Live attenuated vaccines (all types) versus placebo - Safety outcomes, Outcome 2 Adverse events - Peru 15 versus placebo.

Review: Oral vaccines for preventing cholera

Comparison: 9 Live attenuated vaccines (all types) versus placebo - Safety outcomes

Outcome: 2 Adverse events - Peru 15 versus placebo

(Continued ...)

(1) Cohen 2002: Age 18 to 40 years, only records adverse events occurring on the day of vaccination although a diary was completed for 3 days

(2) Qadri 2005: Age 18 to 45 years, 4 days AE monitoring, all AEs are described as mild

(3) Qadri 2007: Age 9 months to 5 years, 4 days AE monitoring, all AE are described as mild.

(4) Sack 2007: Age 18 to 50 years, 3 days AE monitoring, all AE described as mild

Analysis 9.3. Comparison 9 Live attenuated vaccines (all types) versus placebo - Safety outcomes, Outcome 3 Adverse events - VC638 versus placebo.

Review: Oral vaccines for preventing cholera

Comparison: 9 Live attenuated vaccines (all types) versus placebo - Safety outcomes

Outcome: 3 Adverse events - VC638 versus placebo

(Continued . . .)

(Continued . . .)

(1) Benítez 1999: Age 18 to 40 years, AE monitoring for 120 hours, all adverse events are described as mild

(2) García 2005: Age 18 to 40 years, 5 days AE monitoring, all were mild except one headache described as moderate.

(3) Valera 2009: Age 18 to 40 years, 3 days AE monitoring, all adverse events are described as mild

Analysis 10.1. Comparison 10 Live attenuated CVD 103-HgR vaccine versus placebo - Subgroup analysis, Outcome 1 Cases of cholera by age group (age 2-5 years).

Review: Oral vaccines for preventing cholera

Comparison: 10 Live attenuated CVD 103-HgR vaccine versus placebo - Subgroup analysis

Outcome: 1 Cases of cholera by age group (age 2-5 years)

Analysis 10.2. Comparison 10 Live attenuated CVD 103-HgR vaccine versus placebo - Subgroup analysis, Outcome 2 Cases of confirmed cholera attending healthcare facilities (age over 5 years).

Review: Oral vaccines for preventing cholera

Comparison: 10 Live attenuated CVD 103-HgR vaccine versus placebo - Subgroup analysis

Outcome: 2 Cases of confirmed cholera attending healthcare facilities (age over 5 years)

Analysis 10.3. Comparison 10 Live attenuated CVD 103-HgR vaccine versus placebo - Subgroup analysis, Outcome 3 Cases of cholera within four years and five months, by blood group.

Review: Oral vaccines for preventing cholera

Comparison: 10 Live attenuated CVD 103-HgR vaccine versus placebo - Subgroup analysis

Outcome: 3 Cases of cholera within four years and five months, by blood group

Analysis 10.4. Comparison 10 Live attenuated CVD 103-HgR vaccine versus placebo - Subgroup analysis, Outcome 4 Any diarrhoea following artificial challenge, by blood group.

Review: Oral vaccines for preventing cholera

Comparison: 10 Live attenuated CVD 103-HgR vaccine versus placebo - Subgroup analysis

Outcome: 4 Any diarrhoea following artificial challenge, by blood group

Analysis 10.5. Comparison 10 Live attenuated CVD 103-HgR vaccine versus placebo - Subgroup analysis, Outcome 5 Moderate or severe diarrhoea due to V. cholerae after artificial challenge, by blood group.

Review: Oral vaccines for preventing cholera

Comparison: 10 Live attenuated CVD 103-HgR vaccine versus placebo - Subgroup analysis

Outcome: 5 Moderate or severe diarrhoea due to V. cholerae after artificial challenge, by blood group

Analysis 10.6. Comparison 10 Live attenuated CVD 103-HgR vaccine versus placebo - Subgroup analysis, Outcome 6 Additional adverse event data.

Additional adverse event data

Study	Adverse event monitoring	Adverse events reporting	Results
Kotloff 1992	Monitored daily for 7 days	Data presented in an unusable form	'Among volunteers who experienced symptoms, the complaints were mild' 'All episodes of fever were low grade. No subject exceeded the minimum definition of diarrhea (four stools within 24 h) or vomiting (one episode of emesis) or met these criteria for more than 1

Additional adverse event data (Continued)

			day'
Migasena 1989a	Seen daily for 5 days	Text summary only	'No significant adverse reactions, including fever, diarrhea, vomiting, anorexia, or abdominal cramps were observed in any participant during the 7-day period of observation'
Perry 1998	Seen daily for 6 days after each inoculation of vaccine or placebo	Data presented is from a crossover trial where all participants took vaccine and placebo 12 days apart	'No significant difference was seen in reported diarrhoea, fever or vomiting following vaccine or placebo'
Su-Arehawaratana 1992a	Monitored daily for 7 days	Numerical data is only provided for diarrhoea	'No increased rate of diarrhoeal episodes or other gastrointestinal adverse reactions was observed among vaccine than among placebo recipients'
Su-Arehawaratana 1992b	Monitored daily for seven days after each dose	Numerical data is only provided for diarrhoea	'No increased rate of diarrhoeal episodes or other gastrointestinal adverse reactions was observed among vaccine than among placebo recipients'

ADDITIONAL TABLES

Table 1. Detailed Search Strategy

Search set	CIDG SR^	CENTRAL	MEDLINE^^	EMBASE^^	LILACS^^
1	cholera	cholera	cholera	cholera	cholera
2	Vaccin*	Vaccin*	Vaccin*	Vaccin\$	Vaccin\$
3	1 or 2	1 or 2	1 or 2	1 or 2	1 or 2
4		CHOLERA VACCINES	CHOLERA VACCINES	CHOLERA- VACCINE	
5		3 or 4	3 or 4	3 or 4	
6			Limit 5 to human	Limit 5 to humans	

Table 2. The vaccine composition, dosing and participants in field trials of killed whole cell vaccines

Vaccine code (Trade name)	Item	Vaccine composition, dosing and population used in the field trials
WC (not currently available)	Composition	Three strains of <i>V. cholerae</i> O1: <ul style="list-style-type: none"> • 2.5×10^{10} heat-killed <i>V. cholerae</i> classical Inaba whole cells (strain Cairo 48) • 2.5×10^{10} heat-killed <i>V. cholerae</i> classical Ogawa whole cells (strain Cairo 50) • 2.5×10^{10} formalin-killed <i>V. cholerae</i> El Tor Inaba whole cells (strain Phil 6973) • 2.5×10^{10} formalin-killed <i>V. cholerae</i> classical Ogawa whole cells (strain Cairo 50)
	Dosing schedule	Three doses, at 6 week intervals
	Field trial	Clemens 1988 Bangladesh : 41580 participants in primary analysis
	Population	Children aged 2-15 years and women over the age of 15
WC-BS (not currently available)	Composition	As for WC with additional: <ul style="list-style-type: none"> • 1 mg purified cholera B subunit
	Dosing schedule	Three doses, at 6 week intervals
	Field trial	Clemens 1988 Bangladesh : 41,542 participants in primary analysis
	Population	Children aged 2-15 years and women over the age of 15
WC/rBS (Dukoral®)	Composition	As for WC-BS except 1 mg purified cholera B subunit is replaced with: <ul style="list-style-type: none"> • 1 mg recombinant cholera B subunit
	Dosing schedule	Two doses, 2 weeks apart Taylor 2000 Peru also gave a booster dose at 10 months
	Field trials	Sanchez 1994 Peru : (1426 participants in primary analysis), Sanchez 1995 Peru : (307 participants), Taylor 2000 Peru : (17,799 participants)
	Population	Sanchez 1994 Peru and Sanchez 1995 Peru : Military recruits Taylor 2000 Peru : Adults and children aged 2 to 65 years
vWC (ORCVAX®)	Composition	Four strains of <i>V. cholerae</i> O1. As for WC except the 2.5×10^{10} formalin-killed <i>V. cholerae</i> classical Ogawa whole cells (strain Cairo 50) are replaced with: <ul style="list-style-type: none"> • 2.5×10^{10} formalin-killed <i>V. cholerae</i> O1 Inaba, classical biotype cells (strain 569B)
	Dosing schedule	Two doses, 2 weeks apart

Table 2. The vaccine composition, dosing and participants in field trials of killed whole cell vaccines (Continued)

BivWC (Shanchol®)	Field trial	Trach 1997 Viet Nam : 114879 participants in primary analysis
	Population	Adults and children aged > 1 year
	Composition	Three strains of <i>V. cholerae</i> O1 plus one strain of <i>V. cholerae</i> O139: <ul style="list-style-type: none"> • 600 ELISA units of LPS of formalin-killed <i>V. cholerae</i> O1 El Tor Inaba (strain Phil 6973), • 300 ELISA units of LPS of heat-killed <i>V. cholerae</i> O1 Classical Ogawa (strain Cairo 50), • 300 ELISA units of LPS of formalin-killed <i>V. cholerae</i> O1 Classical Ogawa (strain Cairo 50), • 300 ELISA units of LPS of heat-killed <i>V. cholerae</i> O1 Classical Ogawa (strain Cairo 48), and • 600 ELISA units of LPS of formalin-killed <i>V. cholerae</i> O139 (strain 4260B).
	Dosing schedule	Two doses, 2 weeks apart
	Field trial	Sur 2009 India : 66,900 participants in primary analysis
	Population	Adults and children aged > 1 year, living in Kolkata, India

WC = killed whole cell, BS = cholera toxin B subunit, rBS = recombinant cholera toxin B subunit, LPS = Lipopolysaccharide, ELISA = Enzyme-linked immunosorbent assay

Table 3. The vaccine composition, dosing and participants in efficacy trials of live attenuated vaccines

Vaccine code (Trade name)	Item	Vaccine composition, dosing and population used in the field trials
CVD103-HGR (not currently available)	Composition	Richie 2000 Indonesia : 5 x 10 ⁹ lyophilized organisms of a genetically modified <i>V. cholerae</i> O1 Classical Inaba (569B) Tacket 1999 : 2 to 8 x 10 ⁸ CFU of lyophilized organisms of a genetically modified strain of <i>V. cholerae</i> O1 plus buffer
	Dosing schedule	A single dose
	Field trial Artificial challenge study	Richie 2000 Indonesia : 67,508 participants Tacket 1999 : 51 participants
	Population	Richie 2000 Indonesia : Age 2 to 41 years in Jakarta, Indonesia Tacket 1999 : Adults aged 18 to 40 in USA

Table 3. The vaccine composition, dosing and participants in efficacy trials of live attenuated vaccines (Continued)

Peru15 (not currently available)	Composition	5 x 10 ⁸ CFU of a live attenuated strain of <i>V. cholerae</i> O1 El Tor Inaba plus 200ml CeraVax buffer (Cera Products, Columbia)
	Dosing schedule	A single dose
	Artificial challenge study	Cohen 2002 : 36 participants
	Population	Volunteers aged 18 to 40 in USA
VC638 (not currently available)	Composition	1 x 10 ⁹ CFU of a live attenuated strain of <i>V. cholerae</i> O1 El Tor Ogawa plus buffer
	Dosing schedule	A single dose
	Artificial challenge study	García 2005 : 21 participants
	Population	Volunteer males aged 8 to 40 in Cuba

CFU = Colony forming units

WHAT'S NEW

Date	Event	Description
3 August 2011	Amended	Plain language summary amended.

CONTRIBUTIONS OF AUTHORS

This review is an update of a previous review ([Graves 2001](#)). The protocol was revised with input from KA, KZ, FQ, PG and DS. The search results were screened by KA, PG and DS. KA, KZ and DS extracted data. KA wrote the first draft which was revised by DS with additional input from all authors.

DECLARATIONS OF INTEREST

We certify that we have no affiliations with or involvement in any organisation or entity with a direct financial interest in the subject matter of the review (e.g. employment, consultancy, stock ownership, honoraria, expert testimony).

SOURCES OF SUPPORT

Internal sources

- No sources of support supplied

External sources

- Department for International Development, UK.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

Since the publication of the original review, several changes have occurred in standard Cochrane methodology which were not in the original review. Notably; the method of assessing risk of bias has changed, and summary of findings tables incorporating the GRADE methodology for assessing the quality of evidence have been added. The current methodology for these additions is described in the methods section.

INDEX TERMS

Medical Subject Headings (MeSH)

Administration, Oral; Cholera [*prevention & control]; Cholera Vaccines [*administration & dosage; adverse effects]; Randomized Controlled Trials as Topic; Vaccines, Attenuated [administration & dosage; adverse effects]

MeSH check words

Adult; Child; Humans