Accuracy of Real-Time Delivery Room Resuscitation Documentation

Claire E. Fishman¹, Danielle D. Weinberg², Ashley Murray³, and Elizabeth E. Foglia^{1,2}

¹Department of Pediatrics, University of Pennsylvania School of Medicine, Philadelphia, PA USA

²Division of Neonatology, The Children's Hospital of Philadelphia, Philadelphia, PA USA

³Department of Nursing, Hospital of the University of Pennsylvania, Philadelphia, PA USA

Abstract

Objective: To assess the accuracy of real-time delivery room resuscitation documentation

Design: Retrospective observational study

Setting: Level 3 academic Neonatal Intensive Care Unit

Participants: Fifty infants with video recording of neonatal resuscitation

Main Outcome Measures: Vital sign assessments and interventions performed during resuscitation. The accuracy of written documentation was compared with video gold standard.

Results: Timing of initial heart rate assessment agreed with video in 44/50 (88%) records; the documented heart rate was correct in 34/44 (77%) of these. Heart rate and oxygen saturation were documented at 5 minutes of life in 90% of resuscitations. Of these, 100% of heart rate and 93% of oxygen saturation values were correctly recorded. Written records accurately reflected the mode(s) of respiratory support for 89–100%, procedures for 91–100%, and medications for 100% of events.

Conclusion: Real-time documentation correctly reflects interventions performed during delivery room resuscitation but is less accurate for early vital sign assessments.

Introduction

Accurate recording of delivery room resuscitation is essential for complete medical documentation and supports research and quality assurance interventions. The Neonatal Resuscitation Program recommends that resuscitation events are documented as they occur and supplemented with a retrospective narrative summary (1). In many settings, documentation occurs after resuscitation, but retrospective recording is often inaccurate (2). Real-time documentation, in which a scribe records assessments and interventions as they occur, may provide a more accurate method of recording neonatal resuscitation. The

Corresponding author: Elizabeth E. Foglia MD MSCE, 3400 Spruce Street, 8th floor, Ravdin Building, Philadelphia, PA 19104, USA, 215-662-3228, foglia@email.chop.edu.

Conflicts of Interest

The authors disclose no conflicts of interest.

objective of this study was to assess the accuracy of real-time documentation of clinical assessments and interventions during newborn resuscitation after birth.

Methods

Study design and population

This was a retrospective observational study conducted at the Hospital of the University of Pennsylvania, a level 3 academic Neonatal Intensive Care Unit, from June 20, 2017 through December 10, 2017. High-risk resuscitations are conducted in a specialized infant resuscitation room and are routinely video recorded for quality assurance. We planned to enroll 50 infants who underwent resuscitation with complete video recordings, consistent with previous studies (2). Parental consent for video recording is obtained as part of the general hospital admission form. The University of Pennsylvania Institutional Review Board reviewed and exempted this study as a quality assurance initiative.

Equipment

Video recordings: A B-Line LiveCapture ultraportable unit (B-Line Medical, Washington DC) synchronizes video and audio feed from 2 mounted cameras in the infant resuscitation room and 1 video stream from the vital sign monitor. Digital files are securely stored on the hospital server for 4 weeks before they are automatically deleted.

Physiologic Monitoring: High-risk resuscitations are monitored using electrocardiogram and pulse oximetry with a Phillips Intellivue monitor (Phillips, Amsterdam, Netherlands).

Resuscitation Documentation: A neonatal nurse documents infant demographics, vital signs, respiratory interventions and settings, medications, and procedures in real time on a paper form, which becomes part of the medical record (Supplemental Figure). Recording nurses were aware of video recording but were unaware that the accuracy of written documentation was being assessed in the current study.

Measures

Time and date of birth were obtained from each infant's electronic medical record; the remaining variables were abstracted from each video recording. Vital sign assessments included the first heart rate (HR) ascertained during resuscitation- from the vital sign monitor or as announced by a provider's clinical assessment. Pulse oximetry oxygen saturation (SpO₂) was not abstracted for the initial assessment, as this value is often not available in the first minute of resuscitation. At 5 minutes of life (MOL), HR and SpO₂ were abstracted from the vital sign monitor.

We collected the following resuscitation variables: mode(s) of respiratory support, maximum pressure setting(s) for each mode of support, and maximum fraction of inspired oxygen (FiO₂) during resuscitation. In our hospital, team members verbally announce all changes to respiratory support (including pressure and FiO₂ settings) to the recording nurse, who often stands at a slight distance from the team and cannot easily visualize the equipment settings. We used both the visual and audio components of the video to abstract these data for

respiratory interventions. In addition, we collected data for all procedures performed and medications administered.

We collected the same variables from the paper medical record for the duration of the resuscitation captured by video recording. To account for minor differences between clocks (1), we recorded the range of vital signs documented for the duration of time inclusive of the minute before and after 5 MOL.

Data Analysis

For each measure, medical record documentation was compared with video recording gold standard, and the completeness and percent agreement for each variable was reported. Because a range of HR and SpO₂ values were recorded for the 5 MOL assessment, we considered values with any overlap between the video and medical record to agree.

Results

There were 57 resuscitations with video recordings during the study period. We excluded 7, due to incomplete video recordings. Of 50 resuscitations analyzed, 82% occurred on a weekday and 60% occurred during the daytime (07:00–19:00).

Timing of the first ascertained HR agreed with video in 44/50 (88%) records; the documented HR value was correct in 34/44 (77%) cases. When inaccurate, the documented HR was lower than video gold standard in 4/10 and higher in 6/10 cases. The 5 MOL HR and SpO₂ values were documented in 90% of records and were accurate in 100% and 93% of these, respectively (Table). Documented SpO₂ was lower than video in 2 records (–3% and –24%) and higher in 1 (+4%).

The mode of respiratory support was documented more accurately than the maximum pressure settings used (Table). Medication and fluid administration and invasive procedures were accurately recorded in 100% of cases when they occurred. There was only one instance when an intervention (positive pressure ventilation) was documented in the record but did not occur.

Discussion

We sought to assess the accuracy of real-time documentation of delivery room resuscitation. Vital sign documentation was often inaccurate in the first minute of resuscitation and improved by 5 MOL. Real-time documentation correctly captured most resuscitative interventions but reflected the specific details of respiratory support less accurately.

Although accurate recording of delivery room resuscitation is important for complete medical documentation, research, and quality assurance initiatives, resuscitation documentation is often incomplete and/or inaccurate (2–4). Schilleman et al. reported that retrospective documentation accurately reflected the mode of respiratory support in 83%, FiO₂ use in 57%, and initial HR assessment in 37% delivery room resuscitations (2). In contrast, documentation of each of these measures was more frequently accurate in our

study. These results suggest that real-time documentation represents an improvement over retrospective documentation for neonatal resuscitation.

Su et al. compared the accuracy of real-time documentation during pediatric cardiac arrest resuscitation with video and monitor recordings. While the interventions in place were recorded correctly in 100% of resuscitations, paper records accurately reflected the patient's initial rhythm/condition in only 7/15 (47%) of events (3). Similarly, we found that the initial HR was the assessment that was least accurately documented in the written record. It is possible that the initial minutes of resuscitation may be more chaotic and cause discrepancies in the quality of data capture for initial physiological assessments, even with real-time documentation.

Video recording captures more objective and consistent information than both retrospective and in-person documentation (2–4). However, technical and implementation barriers limit the reliability of video recording (5), as evident by the 7 resuscitations excluded from our study due to incomplete recordings. A combination of both video recording and real-time documentation may provide the most complete and accurate method to document neonatal resuscitation.

One study limitation is that we relied on verbal announcement of respiratory settings, rather than downloading objective recorded data. However, this is the method used to communicate this information to the documenting nurse in our hospital. While it is possible these verbalizations do not accurately reflect the actual respiratory settings, we would expect data recorded on the resuscitation form to correspond to the data abstracted from the video, as they were based on the same verbalizations.

Conclusion

Real-time documentation correctly records interventions performed during delivery room resuscitation. Vital sign documentation is often inaccurate at the start of resuscitation and improves by 5 MOL. Documentation in real-time may represent an improvement over retrospective documentation.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

Dr. Foglia is supported by a NICHD Career Development Award, K23HD084727.

List of Abbreviations

FiO₂	fraction of inspired oxygen
HR	heart rate
MOL	minute of life

SpO₂ pulse oximetry oxygen saturation

References

1. Weiner GM, Zaichkin J (Eds.), Textbook of Neonatal Resuscitation, 7th edition. Elk Grove Village, IL: American Academy of Pediatrics 2016: 12.
2. Schilleman K, Witlox RS, van Vonderen JJ, et al. Auditing documentation on delivery room management using video and physiological recordings. *Arch Dis Child Fetal Neonatal Ed.* 2014;99(6):F485–90. [PubMed: 25125582]
3. Su L, Waller M, Kaplan S, Watson A, Jones M, Wessel DL. Cardiac resuscitation events: one eyewitness is not enough. *Pediatric Critical Care Medicine.* 2015;16(4):335–42. [PubMed: 25651046]
4. Grundgeiger T, Albert M, Reinhart D, et al. Real-time tablet-based resuscitation documentation by the team leader: evaluating documentation quality and clinical performance. *Scandinavian Journal of Trauma, Resuscitation, and Emergency Medicine.* 2016; 24:51.
5. Shivananda S, Twiss J, el-Gouhary E, et al. Video recording of neonatal resuscitation: a feasibility study to inform widespread adoption. *World J Clin Pediatr.* 2017;6(1): 69–80.

What is already known on this topic:

1. Accurate recording of delivery room resuscitation is essential for complete medical documentation and supports research and quality assurance interventions
2. Retrospective documentation of assessments and interventions during neonatal resuscitation is often incomplete and/or inaccurate.

What this study adds:

1. Real-time documentation accurately captures interventions performed during delivery room resuscitation.
2. Real-time documentation of vital signs is often inaccurate at the start of resuscitation and improves by 5 minutes of life.
3. Real-time documentation may represent an improvement over retrospective documentation.

Table:

Accuracy of real-time documentation of delivery room assessments and interventions

Resuscitation Characteristics	Documentation Complete	Documentation Accurate if Completed
Demographic Information		
Birth date	98% (49/50)	100% (49/49)
Birth time	98% (49/50)	98% (48/49)
Vital Signs		
First heart rate assessment *	88% (44/50)	77% (34/44)
Heart rate at MOL 5	90% (45/50)	100% (45/45)
SpO ₂ value at MOL 5	90% (45/50)	93% (42/45)
Interventions		
CPAP	89% (34/38)	74% (25/34)
PPV-mask	100% (28/28)	93% (26/28)
PPV-ETT	92% (12/13)	67% (8/12)
Maximal FiO ₂ during resuscitation	100% (50/50)	82% (41/50)
Delayed cord clamping	91% (20/22)	N/A
Intubation	100% (13/13)	N/A
Chest tube placement	100% (1/1)	N/A
Thoracentesis	100% (1/1)	N/A
Surfactant	100% (5/5)	N/A
Normal saline fluid bolus	100% (1/1)	N/A

* Complete documentation of first heart rate assessment refers to concordant timing (± 1 minute) of first ascertained heart rate between written record and video recording.

Abbreviations: CPAP: continuous positive airway pressure; ETT: endotracheal tube; FiO₂: fraction of inspired oxygen; MOL: minute of life; PPV: positive pressure ventilation