Published in final edited form as:

Nat Cell Biol. 2016 July 1; 18(7): 777–789. doi:10.1038/ncb3378.

Protracted p53-independent stimulation of p21^{WAF1/Cip1} fuels genomic instability by deregulating the replication licensing machinery

Panagiotis Galanos^{#1}, Konstantinos Vougas^{#1,2}, David Walter³, Alexander Polyzos², Apolinar Maya-Mendoza⁴, Emma J. Haagensen⁵, Antonis Kokkalis², Fani-Marlen Roumelioti², Sarantis Gagos², Maria Tzetzis⁶, Begoña Canovas⁷, Ana Igea⁷, Emanuel Kanavakis⁶, Dimitris Kletsas⁸, Igor Roninson⁹, Spiros D. Garbis¹⁰, Angel Nebreda⁷, Dimitris Thanos², Paul Townsend^{11,12}, J. Julian Blow⁵, Claus Storgaard Sørensen³, Jiri Bartek^{4,13,**}, and Vassilis G. Gorgoulis^{1,2,11,12,**}

¹Molecular Carcinogenesis Group, Department of Histology and Embryology, School of Medicine, University of Athens, Athens, Greece

²Biomedical Research Foundation of the Academy of Athens, Athens, Greece

³Biotech Research and Innovation Centre (BRIC), University of Copenhagen, Copenhagen, Denmark

⁴Genome Integrity Unit, Danish Cancer Society Research Centre, Copenhagen, Denmark

⁵Centre for Gene Regulation & Expression, College of Life Sciences, University of Dundee, Dundee, UK

⁶Department of Medical Genetics, Medical School, University of Athens, Athens, Greece

⁷Institute for Research in Biomedicine (IRB Barcelona), Barcelona, Spain

⁸Laboratory of Cell Proliferation and Ageing, Institute of Biology, National Centre for Scientific Research 'Demokritos', Athens, Greece

⁹Cancer Center, Ordway Research Institute, Albany, NY, USA

¹⁰Cancer and Clinical Experimental Science Units, Faculty of Medicine, Institute for Life Sciences, Center for Proteome Research, University of Southampton, Southampton, UK

¹¹Faculty Institute of Cancer Sciences, University of Manchester, Manchester Academic Health Science Centre, Manchester, UK

¹²Manchester Centre for Cellular Metabolism, University of Manchester, Manchester Academic Health Science Centre, Manchester, UK

^{**}To whom correspondence should be addressed: Dr Vassilis G Gorgoulis, Dept Histology-Embryology, Medical School, NKUA, 75 Mikras Asias Str, Goudi, Athens, GR-11146, Greece, Tel.: 0030-2107462352, vgorg@med.uoa.gr, vgorgoulis@gmail.com or: Dr. Jiri Bartek, Danish Cancer Society Research Center, Strandboulevarden 49, DK-2100 Copenhagen, Denmark, Tel.: +45 – 35 25 73 57, jb@cancer.dk.

In the memory of Ioannis Terrovitis

Preliminary data of this work have been presented as a keynote presentation at the 2013 Cold Spring Harbor meeting "Eukaryotic DNA Replication & Genome Maintenance"

¹³Institute of Molecular and Translational Medicine, Faculty of Medicine and Dentistry, Palacky University, Olomouc, Czech Republic

These authors contributed equally to this work.

Abstract

The cyclin-dependent kinase inhibitor p21^{WAF1/Cip1} is the prototype downstream effector of the tumor suppressor protein p53. Yet, evidence from human cancer and mice models, imply that p21^{WAF1/Cip1}, under certain conditions, can exercise oncogenic activity. The mechanism behind this behavior is still obscure. Within this context we unexpectedly noticed, predominantly in p53 mutant human cancers, that a subset of highly atypical cancerous cells expressing strongly p21^{WAF1/Cip1} demonstrated also signs of proliferation. This finding suggests either tolerance to high p21^{WAF1/Cip1} levels or that p21^{WAF1/Cip1} *per se* guided a selective process that led to more aggressive off-springs. To address the latter scenario we employed p21^{WAF1/Cip1}-inducible p53-null cellular models and monitored them over a prolonged time period, using high-throughput screening means. After an initial phase characterized by stalled growth, mainly due to senescence, a subpopulation of p21^{WAF1/Cip1} cells emerged, demonstrating increased genomic instability, aggressiveness and chemo-resistance. At the mechanistic level unremitting p21^{WAF1/Cip1} production “saturates” the CRL4^{CDT2} and SCF^{Skp2} ubiquitin ligase complexes reducing the turnover of the replication licensing machinery. Dereglulation of replication licensing triggered replication stress fuelling genomic instability. Conceptually, the above notion should be considered when anti-tumor strategies are designed, since p21^{WAF1/Cip1} responds also to p53-independent signals, including various chemotherapeutic compounds.

Introduction

Accruing evidence point out that a number of molecules involved in key cellular processes display bimodality in cancer i.e. they can act either as tumor suppressors or as oncoproteins (**Supplemental Table 1**). This odd phenomenon is attributed to the so called “cellular or environmental context” that configures their behavior. The mechanistic basis underlying this context-dependent duality is vague in most cases and its explication is essential for rationally designed therapeutic strategies.

The cyclin-dependent kinase inhibitor (CDKI) p21^{WAF1/Cip1} is a pivotal downstream effector of the master tumor-suppressor protein p53, mediating mainly G1 growth arrest in response to various stimuli. This function is primarily dependent on the ability of p21^{WAF1/Cip1} to inhibit cyclin-dependent kinase-2 (Cdk2) (Abbas and Dutta 2009). In spite of its profound p53-dependent role in halting cellular proliferation, several reports, in human cancer and mice models, suggest that p21^{WAF1/Cip1} can manifest oncogenic activities (**Supplemental Table 1**). In some of these studies the oncogenic function was credited to the non-conventional cytoplasmic localization of p21^{WAF1/Cip1} which binds and inhibits the activity of proteins directly involved in apoptosis (Roninson 2002; Pateras et al. 2009). However, in most cases the underlying mechanism remains speculative. It is also interesting that while p53 is frequently mutated in cancer (Rivlin et al. 2011), p21^{WAF1/Cip1} is rarely affected genetically (Abbas and Dutta 2009; Warfel and El-Deiry 2013). The latter would be logical

if p21^{WAF1/Cip1} operated exclusively within the p53 signaling cascade. Nevertheless, p21^{WAF1/Cip1} is activated by a wide range of p53-independent signals and stimuli, including growth factors, nuclear receptors, chemicals and drugs (Abbas and Dutta 2009) (Supplemental Fig. S1).

We report that constitutive expression of p21^{WAF1/Cip1}, in a p53 loss of function environment, causes replication stress and triggers genomic instability by deregulating the replication licensing machinery. Replication licensing is a fundamental biological process that assures that replication takes place once per cell cycle (Abbas et al. 2013). The replication licensing factors (RLFs) ORC, Cdt1 and Cdc6 accumulate during late M and G1 phases forming together with the MCM2-7 helicase the pre-replication complex, licensing the genome for replication. Upon entry into S-phase Cdk activity increases, the replication origins are fired initiating the replication process, while the RLFs are targeted for degradation (unlicensed state) (Takeda and Dutta 2005). Deregulation of the replication licensing process is linked with genomic instability and promotion of malignant behavior, mainly via a process termed re-replication (Blow and Gillespie 2008; Negrini et al. 2010; Halazonetis et al. 2008). Aberrant expression of RLFs is reported in various common malignancies such as head and neck, lung and colon cancer (Karakaidos et al. 2004; Liontos et al. 2007).

Results

1 In advanced stage cancer, a subset of atypical cells expressing p21^{WAF1/Cip1} demonstrated signs of proliferation

In an array of tumors, that have not undergone previous chemo- or radio-therapy, we observed an unusual relationship between p21^{WAF1/Cip1} and the proliferation marker Ki67 that drove our attention. While the anticipated mutual exclusive p21^{WAF1/Cip1}/Ki67 expression pattern was evident, in line with p21^{WAF1/Cip1}'s growth inhibitory properties, we noticed that a number of large cancerous cells with giant nuclei co-expressed p21^{WAF1/Cip1} and Ki67 (Fig. 1). In pathology, identification of cancerous cells with such morphological features is considered a sign of adverse prognosis (Rosai and Ackerman 2011). Notably, most of the carcinomas examined displayed p53 alteration (Karakaidos et al. 2004; Liontos et al. 2007; Velimezi et al. 2013). The irregular p21^{WAF1/Cip1}/Ki67 co-localization led us to hypothesize either tolerance to high p21^{WAF1/Cip1} levels or that somehow constitutive p21^{WAF1/Cip1} expression guided a selective process resulting in the emergence of a subpopulation of p21^{WAF1/Cip1} expressing cancerous cells that regained their proliferative capacity and possibly during this process became more aggressive.

2 p53-independent stimulation of p21^{WAF1/Cip1} up-regulates key replication licensing factors

To address the later scenario we employed a doxycycline-inducible p21^{WAF1/Cip1} expression system (Tet-ON) introduced in the p53 null cells Saos2 (Tet-ON p21^{WAF1/Cip1} Saos2) (Bates et al. 1998) and monitored, subsequent to p21^{WAF1/Cip1} induction, their behavior over a prolonged period (Supplemental Fig. S2). To avoid heterogeneous p21^{WAF1/Cip1} expression from bulk cultures we isolated p21^{WAF1/Cip1}-inducible clones and experimented with those

demonstrating the strongest induction. Periodically (every 5 days) p21^{WAF1/Cip1} expression was evaluated by *in situ* analysis. In parallel, the transcriptome and proteome landscapes were examined at distinct time-points (12hs, 48hs and 96hs) in an attempt to unravel pathways/networks that could exercise over time an “oncogenic” effect (**Supplemental Tables 2,3**). Representative results obtained by the high-throughput assays were confirmed independently by quantitative real-time PCR and immune-blotting (Fig. 2, Supplemental Fig. S3).

As expected and previously reported (Bates et al. 1998), the cells reduced their growth rate and progressively most of them acquired a senescent phenotype that peaked around day 10 (Supplemental Fig. S4, Supplemental Video 1). In accordance with the phenotypical changes and growth kinetics, two-way Gene-Ontology biological-process enrichment analyses, followed by false discovery rate correction (Supplemental Fig. S2), revealed at the transcriptomic and proteomic level, suppression of central factors involved in “mitosis” and “mitosis surveillance” (Fig. 2a). Surprisingly, however, proteome analysis disclosed also a prominent up-regulation of the RLFs, Cdt1, Cdc6 and ORC ($p=1.5 \times 10^{-6}$); with Cdt1 protein raise being the earliest biochemical alteration among all measured transcriptome and proteome changes (**Supplemental Tables 2,3**). The protein abundance raise of the RLFs, was not accompanied by a significant mRNA increase, implying regulation at the post-transcriptional level (Fig. 2). Since Cdt1 and p21^{WAF1/Cip1} are targeted for degradation by the same ubiquitin ligase complexes, CRL4^{CDT2} and SCF^{Skp2} (Abbas et al. 2008; Kim et al. 2008; Havens and Walter 2011), we reasoned that unremitted p21^{WAF1/Cip1} production could sequester their enzymatic activity leading eventually to Cdt1 accumulation (Supplemental Fig. S5a). In support to this notion, SET8, an H4 methyltransferase required for chromosome compaction in mitosis, and E2F1, targets of CRL4^{CDT2} (Jorgensen et al. 2011) and Skp2 (Marti et al. 1999), respectively, were up-regulated after p21^{WAF1/Cip1} induction (Supplemental Fig. S5a,S6). Moreover, transfection with a mutant p21^{WAF1/Cip1} that harbors three substitutions in its PIP degron motif abrogating its interaction with PCNA, which is essential for CRL4^{CDT2} to target p21^{WAF1/Cip1}, did not augment Cdt1 expression, strongly favoring the “saturation/competition” mechanism (Fig. 2b). In addition, the cell cycle profile imposed by constitutive p21^{WAF1/Cip1} expression, characterized by a G1-phase increase and an S-phase decline, could contribute to the reduced RLF protein turn-over observed (Supplemental Fig. S5b). Along the same line, blocking protein synthesis with cyclohexamide, demonstrated a dramatic enhancement of Cdc6 protein stability, in p21^{WAF1/Cip1} expressing cells (Supplemental Fig. S5c). Reduced Cdc6 protein turn-over could be attributed to low activity of the E3-ligase APC^{Cdh1} that targets Cdc6 for degradation (Duursma and Agami 2005). Indeed, the levels of the E3-ligase APC substrate recognition and activating modules Cdh1/Fzr1 were decreased, supporting the above assumption (Supplemental Fig. S5c). Increased Cdc6 stability was also associated with Cdk2-mediated phosphorylation at Serine 54 (Cdc6-pS54) that protects it, as previously reported, from rapid destruction (Supplemental Fig. S5c) (Duursma and Agami 2005). Despite the fact that p21^{WAF1/Cip1} expression reduced Cdk2 activity (Supplemental Fig. S7) low Cdh1/Fzr1 levels appears to tilt the balance in favor of Cdc6-pS54 accumulation (Supplemental Fig. S5c). Notably, in the presence of cycloheximide and the proteasome inhibitor MG132, the protein levels of Cdh1/Fzr1 were not restored implying regulation at

the transcriptional level. To this end we examined the mRNA levels of Cdh1/Fzr1 and found them suppressed under conditions of p21^{WAF1/Cip1} expression (Supplemental Fig. S5d).

Since the p53 checkpoint was shown to protect from re-replication (Vaziri et al. 2003), we reasoned that the status of p53 would define the ability of p21^{WAF1/Cip1} to regulate the levels of Cdt1 and Cdc6. Indeed, challenging cellular systems bearing inactive p53, with p21^{WAF1/Cip1}, resulted in the up-regulation of these replication licensing factors (Fig. 2c). On the other hand, p21^{WAF1/Cip1} expression in a cellular environment with wild-type p53 suppressed their expression (Fig. 2c,d). It was suggested, but not shown, that p53 shields the organism from cells undergoing re-replication by triggering apoptosis (Vaziri et al. 2003). In line with this notion, we noticed that the p21^{WAF1/Cip1}-IPTG-ON HT1080 cells, harboring wtp53, experience massive apoptosis, after p21^{WAF1/Cip1} induction, which was accompanied by a dramatic decrease of Cdt1 and Cdc6 expression, whereas silencing of p53 suppressed apoptosis, up-regulated Cdt1, and restored Cdc6 levels (Vaziri et al. 2003).

3 Activation of p21^{WAF1/Cip1}, in p53 null cells, triggers replication stress in a Cdt1/Cdc6-dependent manner

Re-replication, is a form of replication stress and is considered, as mentioned, the basis of genomic instability driven by inappropriate expression of the RLFs, Cdc6 and Cdt1 (Blow and Gillespie 2008; Abbas et al. 2013). Although various mechanistic aspects of how re-replication challenges the stability of the genome are still obscure, it is well ascertained that it leads to DNA damage and DNA damage response (DDR) activation (Blow and Gillespie 2008; Abbas et al. 2013). Following p21^{WAF1/Cip1} induction, MCM2-7 chromatin loading increased robustly indicating that Cdt1 and Cdc6 up-regulation is functional (Supplemental Fig. S8). Flow cytometric analysis of propidium iodide stained cells (DNA content) and double co-stained cells for DNA content and DNA synthesis (Ethynyl-deoxyUridine: EdU) revealed a cellular sub-population with DNA content greater than 4n, indicative of re-replication (Fig. 2e; Supplemental Video 2). Subsequently, DNA damage, as assessed by single cell gel electrophoresis (comet assay) under alkaline conditions (total number of DNA lesions) and pulse field gel electrophoresis (PFGE), was evident, tracked by signs of DDR, as depicted by increase recruitment of 53BP1 in foci and H2AX phosphorylation (γ -H2AX) (Fig. 3a,b,c, Supplemental Fig. S6). Strikingly, re-replication, DNA damage, as well as the DDR, were Cdc6- and Cdt1-dependent further suggesting that re-replication was the source of replication stress (Fig. 3b,c). Finally, re-replication and DNA damage was significantly lesser when the p21^{PCNA} mutant was employed keeping in line with the proposed “saturation/competition” mechanism (Supplemental Fig. S5,S9).

It seems paradoxical that p21^{WAF1/Cip1} could trigger replication stress, given its role as a potent inhibitor of cell cycle progression. Nevertheless, fork progression analysis by DNA fiber spreading (DNA combing), using 20 min consecutive labeling pulses of CldU (red) and IdU (green), showed that replication fork progression did not cease, but its speed was significantly reduced (Fig. 3d). In addition, replication fork asymmetry was observed possibly related with the presence of DNA lesions impeding symmetrical bi-directional movement (Fig. 3d). Consistent with this notion, flow cytometry analysis with double co-staining for γ -H2AX, DNA content (Propidium Iodide: PI) and DNA synthesis (Ethynyl-

deoxyUridine: EdU) showed that, following p21^{WAF1/Cip1} induction, DNA damage accumulated mainly in cells incorporating EdU, whereas depletion of Cdc6 and Cdt1 profoundly suppressed the accumulation of DNA damage in S-phase (Fig. 3e). Markedly, the cells expressing p21^{WAF1/Cip1} demonstrated a focal PCNA pattern characterized by small dots that represents a trait of early S-phase (Essers et al. 2005)(Supplemental Fig. S10). This finding suggests that DNA damage occurs at a time sensitive slot, as current data demonstrate the presence of early replication fragile sites that are prone to breakage under conditions of replication stress (Barlow et al. 2013).

4 p21^{WAF1/Cip1}-mediated replication intermediate lesions are processed by MUS81-EME1 and repaired by a Rad52-dependent mechanism

To further characterize p21^{WAF1/Cip1}-mediated replication stress we examined for single-stranded DNA (ssDNA) formation, a common intermediate at replication-associated lesions. To this end, p21^{WAF1/Cip1} expressing cells were incubated with BrdU for 24hs, under non-denaturing conditions, allowing anti-BrdU staining to disclose ssDNA (Beck et al. 2010). In line with our hypothesis, *in situ* analysis showed a strong co-localization between BrdU staining and p21^{WAF1/Cip1} expression (Fig. 4a), that was also associated with an increased number of foci formed by the ssDNA-binding protein RPA (Fig. 4b). The ssDNA could be generated by replicative helicase – polymerase uncoupling (stalled fork), template-strand resection after double strand break (DSB) formation, unequal branch migration of stalled fork that yield reversed forks structures (chicken foot), or ssDNA gaps on template DNA (Petermann and Helleday 2010; Neelsen et al. 2013a; Couch et al. 2013)(Supplemental Fig. S11). To further discriminate whether ssDNA is from the template or the newly synthesized (nascent) strands short BrdU pulses (20 min) were used. The analysis did not show BrdU staining, suggesting that the source of the ssDNA is the template strand (Supplemental Fig. S12). Regardless of the way ssDNA is formed, it represents a structure of DNA replication intermediate lesions that need to be resolved for replication to restart. It has been reported that after long periods of replication inhibition double strand breaks (DSBs), generated by the structure-specific endonuclease complex of MUS81-EME1*, are required for restart (Hanada et al. 2007; Petermann and Helleday 2010). Based on this fact we hypothesized that sustained p21^{WAF1/Cip1} expression phenocopies the latter state involving MUS81-EME1. Indeed, MUS81-EME1 depletion caused a significant decrease of the DNA damage inflicted by p21^{WAF1/Cip1} expression, as well as reduction of the EdU positive cells harboring signs of DNA damage (Fig. 4c,d; Supplemental Fig. S13). Recent reports present MUS81-EME1 as a central player in oncogene-induced genotoxicity (Neelsen et al. 2013b; Murfuni et al. 2013), promoting homologous recombination (HR)-mediated repair of inactivated (collapsed) forks (Petermann and Helleday 2010). Surprisingly, we noticed that silencing of the homologous repair recombinase Rad51 resulted in decreased γ -H2AX levels (Fig. 4e). This finding implies a negative control over an alternative, Rad51-independent, repair process. Rad51 was reported to exert such an effect preventing Rad52-dependent DNA annealing and repair (Wu et al. 2008). In fact, suppression of Rad52 was followed by increased γ -H2AX expression and cell death (Fig. 4f,g), suggesting that Rad52 guided the repair process. Rad52 is possibly involved in microhomology-mediated repair pathways, which are considered mutagenic (Ottaviani et al. 2014) and according to latest data collapsed forks are recovered by error-prone replicative-based mechanisms challenging genome stability (Iraqi et al. 2012). An

interesting observation was that Rad51 levels were reduced upon p21^{WAF1/Cip1} induction. Rad51 is in short supply and under certain stressogenic conditions such as hypoxia it is repressed by E2F4/p130 complexes. Given that p53-dependent gene repression via p21^{WAF1/Cip1} is mediated by recruitment of such complexes (Benson et al. 2013), we hypothesized that protracted p21^{WAF1/Cip1} expression mimics such circumstances and examined whether the promoter of Rad51 is occupied by E2F4. Indeed it did (Fig. 4h), providing a mechanistical explanation why Rad52 is the preferable repair choice in this setting.

5 Deregulated up-regulation of Cdt1 and Cdc6 links p53-independent activation of p21^{WAF1/Cip1} with senescence

The results so far showed that sustained p21^{WAF1/Cip1} expression triggered senescence, a well established anti-tumor response (Bartkova et al. 2006; Halazonetis et al. 2008) that nonetheless, in a p53 null environment, harbored the seeds of genomic instability, in the form of deregulated licensing factors. p21^{WAF1/Cip1}-imposed senescence was to a large extent mediated by Cdt1 and Cdc6, apparently as a result of the DNA damage checkpoints they activated (Fig. 5a,b) (Bartkova et al. 2006; Lontos et al. 2007). In a previous study we noticed that p73, the p53 homologue, responds to the DDR pathway (Lontos et al. 2009). As inappropriate expression of Cdt1 and Cdc6 caused replication stress and DNA damage we asked whether p73 could operate as the downstream effector of the emerging p21^{WAF1/Cip1}-RLFs-DDR-signaling route, stimulating senescence. We found that p21^{WAF1/Cip1} induction elicited p73 up-regulation, in a Cdt1/Cdc6-dependent manner, that was essential for executing the senescent program (Fig. 5a,b). According to the oncogene-induced DNA damage concept for cancer development continuous formation of DNA breaks exerts a selective pressure on the DDR-induced anti-tumor barriers, eventually breaching them, leading to genomic instability and tumor progression (Halazonetis et al. 2008). If this holds true, then at some point the p21^{WAF1/Cip1} expressing cells will bypass the senescence barrier generating more aggressive offsprings. In addition, apart from fueling genomic instability (Sideridou et al. 2011), Cdc6 overexpression could confer to tumorigenesis by repressing the *INK4/ARF* locus (Gonzalez et al. 2006). Indeed, p21^{WAF1/Cip1} activation led to down-regulation of both products of the *INK4/ARF* locus, p16^{INK4A} and p14^{ARF}, apparently as a result of Cdc6 up-regulation (Fig. 5c). Suppression of p16^{INK4A} removes an indispensable factor for maintaining the senescent phenotype (Beauséjour et al. 1999); thus rendering p21^{WAF1/Cip1}-mediated senescence possibly reversible.

6 Protracted expression of p21^{WAF1/Cip1}, in a p53 loss of function environment, bypasses senescence, fuels genomic instability, promoting aggressive behavior

Monitoring the behavior of the cells we interestingly observed that after the 10th day of switching on p21^{WAF1/Cip1} the senescent phenotype gradually declined. Concurrently, a sub-population of non-senescent p21^{WAF1/Cip1} positive cells emerged (Fig. 6a) demonstrating, later than the 20th day of induction, BdrU and EdU incorporation values similar to those seen in the control counterparts (Fig. 6b,c; Supplemental Video 3). Likewise, around the same time period, the mutual exclusive relationship between cyclin A, an established late S/G2 marker (Woo and Poon 2003; Pines and Hunter 1991), and p21^{WAF1/Cip1} was reduced, and accompanied by a double p21^{WAF1/Cip1}/cyclin A positive sub-population (Fig. 6d).

These results strongly suggest that a fraction of p21^{WAF1/Cip1} expressing cells evaded from p21^{WAF1/Cip1}-induced senescence and arrest, re-entering the cell cycle (from now on these cells are termed “*escaped*”). To support this observation, cdk2 activity as well as phosphorylation of the stimulatory site T160 of Cdk2 that were suppressed upon p21^{WAF1/Cip1} induction, were restored (Fig. 6e, Supplemental Fig. S14). Strikingly, the “*escaped*” cells showed a dramatic reduction of p73 expression that concurs with the decline of the senescent phenotype (Fig. 6e). Another interesting finding was that most of their nuclei were significantly larger than those of the un-induced cycling control cells (Fig. 6f, Supplemental Fig. S15, Supplemental Video 4), a feature that was also observed *in vivo* (Fig. 1). DNA damage formation was also reduced in the “*escaped*” cells implying that an extensive repair process took place (Fig. 6g). The involvement of the MUS81-EME1 - Rad52 repair route (Fig. 4c,d) and the increased presence of micronuclei (Fig. 7b) that are considered surrogate markers of chromosomal instability, defective DDR and repair (Terradas et al. 2009), point out, as mentioned already above, that the latter was error-prone. To gain a more precise, genome wide, view of this likely outcome we performed array comparative genomic hybridization (aCGH) analysis, deep sequencing and M/FISH/SKY combined with inverted-DAPI banding between the “*escaped*” and the un-induced cells (Fig. 7, Supplemental Fig. S16-S18; **Supplemental Tables 4-5**). In total six independent biological replicates were performed and the duration of each experiment was 30 days to produce an adequate amount of “*escaped*” cells for the analysis. To avoid cell culture propagation effects that could mask the effect of p21^{WAF1/Cip1} activation the OFF cells were cultured for the same period (Supplemental Fig. S16a). In aCGH the un-induced cells were used as reference; thus eliminating the noise showing only the differences between the two states (OFF vs ON). Cumulatively the results from all three experimental procedures clearly disclosed that the genome context of the “*escaped*” cells acquired chromosomal aberrations, in the form of gains and losses that ranged in size from approximately 1.75Kb to 92Mb (Fig. 7a, Supplemental Figs S16b,c, S17, S18; **Supplemental Tables 4-5**), as well as novel translocations (Fig. 7c, Supplemental Fig. S16c). Interestingly among the genetic lesions found were alterations concentrated in the same chromosome bands (3p14.3, 3p14.2, 10p15.3) consisting of alternating regions of gains or losses with retention regions in between (Fig. 7a, **Supplemental Table 4**), most probably representing events of chromoanasythesis or chromothripsis, respectively (Holland and Cleveland 2012). One of the most noteworthy observations of this multifaceted experimental approach was the strong concordance between the findings of the cytogenetic, aCGH and deep sequencing analyses (Fig. 7d, Supplemental Fig. S16-S18). Given that each experimental procedure took place at a different time period we can conclude that p21^{WAF1/Cip1} steers a “deterministic” set of genetic events that may play a role in the behavior of the “*escaped*” cells. In line with this notion the transcriptome of the “*escaped*” cells (Supplemental Fig S19,S20), using Monte-Carlo simulation, demonstrated a specific non-random correlation with the genomic alterations found in these cells ($p < 2.2 \times 10^{-16}$). This finding suggests that a causality inference can be safely made since deregulation of the replication licensing machinery and particularly Cdt1 was the first biochemical event to occur after p21^{WAF1/Cip1} stimulation; deducing that genomic instability led to the observed changes in the transcriptome of the “*escaped*” cells. Although p21^{WAF1/Cip1} is not a transcription factor it has been reported to enhance transcription, in certain cases by modulating the activity of transcription cofactors

(Perkins et al. 1997). Nevertheless, the fact that only 42 (7.6%) of the 553 genes found differentially expressed in the “escaped” cells were detected in earlier time-points (12, 48 and 96 hrs) provides further evidence that p21^{WAF1/Cip1}-mediated transcription is most unlikely to be the driving force behind the emerged transcriptome of the “escaped” cells (Supplemental Fig. S21).

The most important finding of the present study was that the descendent/“escaped” clones demonstrated significantly higher anchorage-independent growth and were more invasive when evaluated in comparison with the un-induced cells (Fig. 6h,i). Even more they tolerated treatment with the genotoxic drugs, doxorubicin and cisplatin much more efficiently whereas, no significant difference in the response to taxol, a microtubule polymer stabilizer and mitotic inhibitor, was noticed (Fig. 6j). The enhanced resistance to doxorubicin and cisplatin remained even when p21^{WAF1/Cip1} activation was switched off in the “escaped” cells for 10 days, clearly denoting that this feature was independent from a potential p21^{WAF1/Cip1}-mediated indirect transcriptional effect. As a final point, a number of the transcriptionally altered genes are reported to be functionally connected with aggressive behavior such as, invasion, metastasis and cancer “stemness” providing the ground to interpret the acquired aggressive phenotypic features (Supplemental Fig. S19) (Supplemental Table 6).

Discussion

The present report advances our understanding of how p21^{WAF1/Cip1} can exhibit oncogenic activities. The key factor in tipping the balance from the tumor suppressor function to that of an oncogene was p53 inactivation. Our data suggest a novel function for p53; that of protecting the cells from the adverse effects of p21^{WAF1/Cip1}.

Dissecting p21^{WAF1/Cip1} from the control of p53 demonstrated the ability of the former, when stimulated by p53-independent signals, to deregulate the replication licensing machinery. We provide evidence that continuous production of p21^{WAF1/Cip1} saturates its degradation modules, CRL4^{CDT2} and SCF^{Skp2}, leaving their other targets, including Cdt1 and E2F1/2 unabated to perform their function. Since Cdt1 and Cdc6 represent E2F targets (Karakaidos et al. 2004; Schlisio et al. 2002) a putative consequence of this mechanism could be the formation of positive feedback loops boosting Cdt1 and Cdc6 expression (Fig. 7f). By up-regulating the pivotal replication licensing factors Cdt1, Cdc6 and Orc, the cells expressing p21^{WAF1/Cip1} acquire the capacity to re-replicate. Our work sheds new light on an earlier mechanistic observation, noticing that p21^{WAF1/Cip1} can promote “endo-reduplication” (Niculescu et al, 1998). It also goes one step further demonstrating experimentally that p21^{WAF1/Cip1}-mediated re-replication guides a selective process giving rise to descendant cells with more aggressive features compared to their progenitors (Fig. 7f). Re-replication is a form of replication stress in which the temporal separation of origin licensing and firing is lost. This feature leads to replication fork collapse, DNA damage and eventually genomic instability, if not repaired in an error-free manner (Blow and Gillespie 2008). In comparison to endo-reduplication (multiple consecutive S-phases with no passage through mitosis), which naturally takes place in certain types (Porter 2008), re-replication does not appear to occur as part of any physiological developmental program. Within this

context, the p53 checkpoint was shown to hamper re-replication (Vaziri et al. 2003). It was also suggested that cells undergoing re-replication could be eliminated by apoptosis (Vaziri et al. 2003). This claim was clearly demonstrated in the p21^{WAF1/Cip1}-IPTG-ON HT1080 cellular model (Fig. 2c), that harbors wt p53, further supporting our concept. The fact that the turn-over of p21^{WAF1/Cip1} and Cdt1 is controlled by the same E3-ubiquitin ligases underscores the significance of p53 since its inactivation confiscates a protective mechanism. Having in mind that p21^{WAF1/Cip1} mutations are extremely rare events in cancer (Abbas and Dutta 2009), it is apparent that human cancers, with loss of p53 function, are at risk of suffering additional deleterious genetic alterations by protracted operation of p21^{WAF1/Cip1}, activated through p53-independent signaling routes. Among the p53-independent stimuli shown to induce p21^{WAF1/Cip1} are growth factors such as EGF, FGF2 and TGF- β frequently reported to be overexpressed in various types of human cancer (Abbas and Dutta 2009).

The ensuing involvement of the HJ resolvase MUS81-EME-1 and the recombinase Rad52 points towards a replication-based error prone repair process. MUS81-EME-1 is a key player in oncogene-induced replication stress (Neelsen et al. 2013b; Murfun et al. 2013), whereas reduction of Rad51 generated the conditions for a switch from a high-fidelity homologous recombination to a lower fidelity process mediated by Rad52. Rad52-dependent recombination requires much less homology (micro-homology) than Rad51 facilitating illegitimate events during repair (Ottaviani et al. 2014; Hastings et al. 2009; Wu et al. 2008) (Fig. 7f). This notion was confirmed by aGCH and molecular cytogenetics unraveling an altered genome landscape in the “escaped” cells. Characteristically among the lesions observed some strongly resembled the recently described complex chromosomal rearrangement of chromoanagenesis that appears to result from replicative template-switching events (Fig. 7)(Zhang et al. 2013). Chromothripsis was another complex chromosomal rearrangement viewed (Fig. 7), considered to be the outcome of NHEJ (Stephens et al. 2011), implying that other repair pathways, possibly non-replicative ones, may be involved in p21^{WAF1/Cip1}-driven genomic instability. In such a case, lesions that are not fixed during S-phase or the subsequent G2 by micro-homology-mediated procedures may be repaired in the next G1-phase by its preferred method, NHEJ (Ottaviani et al. 2014). In parallel to re-replication, p21^{WAF1/Cip1} can further promote genomic instability by suppressing, as reported, mismatch and nucleotide excision repair (Abbas et al. 2009).

A question that always emerges when genomic alterations are acquired is whether they represented a passenger or a driver event. The fact that the transcriptomic changes showed a highly significant correlation ($p < 2.2 \times 10^{-16}$) with the genomic ones strongly supports the latter notion. Among the transcripts found deranged were that of IGF1, FGFR2, MMP13 and CD44 well established for their role in cancer progression, possibly accounting for the aggressive behavior of the “escaped” cells (Supplemental Fig. S19,S20). Another up-regulated factor was ID2 shown to antagonize the suppressive effects of p16^{INK4A} and p21^{WAF1/Cip1} (Sikder et al. 2003). That p21^{WAF1/Cip1}-driven genome changes constitute part of a selection trajectory to promote survival and long term evolution was further illustrated by the finding that the descendent/“escaped” clones tolerated DNA damage much more efficiently when exposed to the genotoxic drugs, doxorubicin and cisplatin. Even when exogenous stimulation of p21^{WAF1/Cip1} was switched off the resistance to the DNA damaging agents remained impervious. This observation indisputably depicts that the

“escaped” cells adapted to DNA damage inflicted by p21^{WAF1/Cip1}-dependent re-replication and that this trait is independent from any potential p21^{WAF1/Cip1}-mediated indirect transcriptional effect.

Conclusively our results highlight a “dark side” of p21^{WAF1/Cip1} that should be taken into consideration when designing therapeutic strategies, particularly in p53-null tumors, as a number of agents used in clinical oncology, such as dexamethazone, can activate p21^{WAF1/Cip1} in a p53-independent manner (Cha et al. 1998) with potential detrimental effects to patients.

Material and Methods

Tumor specimens

Formalin-fixed, paraffin-embedded sections from 20 surgically removed head-neck carcinomas, 30 urothelial bladder carcinomas and 30 lung carcinomas were analyzed, after local ethical committee approval. Patients had not undergone chemo- or radiotherapy before surgical resection. The majority of clinical samples have been previously described (Liontos et al. 2007; Evangelou et al. 2013).

Cell lines and culture treatments

Saos2 Tet-ON p21^{WAF1/Cip1} (human osteosarcoma), H1299 p21^{WAF1/Cip1}-Ponesterone-ON (human lung carcinoma) and HT1080 p21^{WAF1/Cip1}-9 (human fibrosarcoma) carrying Tet-ON, ponesterone and isopropyl-b-D-thiogalactoside (IPTG)-inducible p21^{WAF1/Cip1} respectively (Bates et al. 1998; Chang et al. 1999; Wang et al. 1999), MCF7 (human breast cancer) and MDA-MB-231 (human breast cancer) cell lines were maintained in DMEM (Invitrogen) with 10% FCS (Invitrogen), 2mM l-glutamine (Invitrogen), and 100 µg/ml penicillin and streptomycin (Invitrogen) at 37°C and 5% CO₂. Growth and maintenance of Li-Fraumeni Syndrome (LFS) fibroblasts was as previously described (Bischoff et al. 1990).

For p21^{WAF1/Cip1} expression induction, Saos2 p21^{WAF1/Cip1} cells were treated with 2µg/ml Doxycycline (Applichem), while HT1080 p21^{WAF1/Cip1}-9 cells were treated with 100µM IPTG (Ambion). MCF7 and MDA-MB-231 cells were treated with 5 ng/ml TGF-β.

Microphotographs were obtained on an inverted microscope (Axiovert S100; Carl Zeiss) with CP-Achromat objectives and a charge-coupled device IRIS high-resolution color video camera (SSC-C370P; Sony), whereas the Image Pro Plus v3.0 (Media Cybernetics) was used for image acquisition.

siRNA transfections and retrovirus infections

Cdc6, Cdt1, Rad52 (Invitrogen), Rad51 (Thermo Scientific) and Mus81 (Santa Cruz) siRNA gene silencing was performed as previously described (Liontos et al. 2009). Briefly, 3×10⁵ cells plated in 60mm dishes were transfected using Lipofectamine 2000 (Invitrogen) with the appropriate RNAi pool (set of three siRNAs from the manufacturer) or the corresponding RNAi negative control (Invitrogen) as per manufacturer’s guidelines. Cells were harvested 48h after transfection for further analysis.

LFS cells were transiently infected with the pBabe-p21^{WAF1/Cip1} or the corresponding control vector using the Phoenix helper-free retrovirus producer cell line as described before (Sideridou et al. 2011). H1299 cells were transiently infected with the pMSCV, pMSCV-p21^{WAF1/Cip1} or pMSCV-p21^{PCNA} (a mutant p21^{WAF1/Cip1} harboring Q144, M147, F150 substitutions to A in its PIP degron motif) vectors using the Phoenix helper-free retrovirus producer cell line (Abbas et al. 2008).

Immunohistochemistry (IHC)

The following antibodies were used: p21^{WAF1/Cip1} (1:200, Santa Cruz) and Ki67 (1:100, Dako). IHC was performed using the UltraVision LP Detection System (Thermo Scientific) as per manufacturer's instructions. Paraffin sections (4 μ m) were deparaffinised in xylene and rehydrated in a graded series of ethanol-aqueous solutions. Antigen retrieval was carried out in 10mM citrate buffer (pH 6.0) by heating the slides for 25min in a microwave oven. Endogenous peroxidase activity was blocked with 3% hydrogen peroxide in TBS for 10min. Primary antibodies were incubated overnight. Evaluation and controls were as previously described (Evangelou et al. 2013).

Indirect Immunofluorescence (IF)

Indirect IF analysis was performed as previously published (Liontos et al. 2007). Briefly, cells grown on coverslips were fixed in 4% formaldehyde and permeabilized by 0.1% Triton X-100 in two 15 min consecutive steps, at RT. After washing with PBS, cells were blocked for 30min in 10% FCS. FFPE tissues sections were used after dewaxing and rehydration as described in the IHC paragraph. Primary antibodies used were: p21^{WAF1/Cip1} (1:200, Santa Cruz), Ki-67 (1:300, Dako), histone H2A.X phosphorylated at serine 139 (1:300, Millipore), Cyclin A (1:200, Santa Cruz), 53BP1 (1:200, Abcam), PCNA (1:100, Santa Cruz), Rad51 (1:200, Santa Cruz,) and hRPA32 (1:1000, Genetex). Cells were then incubated with primary antibodies at 4°C overnight. Secondary antibodies Alexa Fluor® 488 goat anti-rabbit (1:500, Invitrogen) and Alexa Fluor® 568 goat anti-mouse (1:500, Invitrogen) were applied for 60min at RT, followed by final wash in PBS. Counterstaining was performed with 100 ng/ml of 4,6-diamidino-2-phenylindole (DAPI)(Sigma-Aldrich). Image acquisition of multiple random fields was automated on a ScanR screening station (Olympus) and analyzed by using ScanR (Olympus) analysis software, or a Zeiss Axiolab fluorescence microscope equipped with a Zeiss Axiocam MRm camera and Achroplan objectives while image acquisition was performed with AxioVision software 4.7.1.

Total protein extraction and immunoblotting

Total protein extracts—Total protein extraction was performed as described before (Liontos et al. 2007). Briefly, cells were homogenized in 50 mmol/L HEPES pH 7.5, 150 mmol/L NaCl, 15 mmol/L β -mercaptoethanol, 0.5 mmol/L phenylmethanesulfonyl fluoride (PMSF), 0.1% Nonidet P-40 (Sigma) and centrifuged at 1000xg at 4°C for 10min. The supernatant was collected and adjusted to 1 μ g/ml protease and phosphatase inhibitors (Pierce). The pellet was resuspended in 10mM HEPES pH 7.9, 1.5mM MgCl₂, 10mM KCl, 0.5mM DTT, and 1.5mM PMSF, incubated on ice for 30min and centrifuged at 1000xg at

4°C for 10min. The supernatant fraction containing histones was collected. Total protein and histone concentration was evaluated using the Bio-Rad Protein Assay (Bio-Rad).

Biochemical cell fractionation—Cells (2.5×10^6) were collected and re-suspended in 200µl buffer A (10mM Hepes pH7.9, 10mM KCl, 1.5mM MgCl₂, 0.34M Sucrose, 10% glycerol, 1mM DTT, protease and phosphatase inhibitors)(Supplemental Fig. S8a). Following, 0.1% Triton X-100 was added and then incubated for 6min on ice. Next lysates were centrifuged at 1300g/4°C for 5min. The supernatant (S1) fraction was clarified by centrifugation for 15min at 20000g/4°C to remove cell debris and insoluble aggregates. The resultant supernatant, soluble fraction was used as cytosolic (S2) one. The pellet from the initial lysate was first washed with buffer A and then subjected to further lysis in buffer B (3mM EDTA, 0.2mM EGTA, 1mM DTT, protease and phosphatase inhibitors) for 30min on ice. Subsequently, a centrifugation of 5min at 1700g/4°C was applied. The supernatant was collected as the soluble nuclear (S3) fraction. The pellet was washed in buffer A and then centrifuged for 5 min at 1700g/4°C. Finally, the pellet was re-suspended in buffer A to obtain the chromatin fraction (P3)(Supplemental Fig. S8a).

Primary antibodies employed were: p21^{WAF1/Cip1} (1:400, Santa Cruz), Cdc6 (1:1000, Millipore), Cdt1 (1:400, Santa Cruz), H2AX (1:1000, Abcam), γH2AX (1:1000, Millipore), p73 (1:1000, NeoMarkers), Mus81 (1:400, Santa Cruz), Cdk2 (1:1000, Santa Cruz), Cdk2-pT160 (1:1000, Cell Signalling), Cdk2-pY15 (1:1000, Santa Cruz), PLK1 (1:1000, Santa Cruz), TOP2A (1:1000, Santa Cruz), β-actin (1:1000, Cell Signalling), p14^{ARF} (1:100, Abcam), Cyclin A (1:1000, Santa Cruz), histone H3-pS10 (1:1000, Abcam), Cdk1 (1:400, Santa Cruz), Rad51 (1:500, Santa Cruz), Rad52 (1:500, Santa Cruz), MCM7 (1:1000, Abcam) and p53 (1:1000, Santa Cruz).

Thirty µg of protein from total extracts or 1µg of histones from each sample were adjusted with Laemmli Buffer (Sigma) and loaded on PAGE gels. Gel electrophoresis and transfer to PVDF membrane (Millipore) were performed according to standard protocols. Blots were blocked for 1h in 5% non-fat milk in TBS-0.1% (v/v) Tween-20 solution at RT.

Subsequently, membranes were incubated overnight with primary antibody solution in 0.5% non-fat milk in TBS with 0.1% (v/v) Tween-20 at RT, followed by a 45min incubation with alkaline phosphatase conjugated anti-mouse or anti-rabbit secondary antibody (1:5000 dilution)(Promega) at RT. Signal development was performed with nitro blue tetrazolium/5-bromo-4-chloro-3-indolylphosphate (NBT/BCIP) solution (Molecular Probes) as previously described (Sideridou et al. 2011).

CDK2 kinase activity

For each sample, 50mg of total-cell protein extract was precleared for 2h at 4°C with 5mg of rabbit immunoglobulin G (anti-Cdk2) or 5 mg of mouse immunoglobulin G (anti-cyclin B1) prebound to protein A-Sepharose (Millipore). Precleared lysates were transferred to new microcentrifuge tubes and incubated with anti-Cdk2, or anti-cyclin B1 with mixing for 2h at 4°C. Protein A-Sepharose was added and the samples were mixed for 2h at 4°C. The immunoprecipitates were washed twice with TBS and twice with kinase buffer (100mM Tris pH 7.4, 20mM MgCl₂, 2mM dithiothreitol) before incubation with 5mg of histone H1

(Boehringer Mannheim) and 15nM ATP for 10min at RT. Samples were incubated with Malachite Green Reagent for 30min (Cdk2 and Cyclin B1). The kinase assay mixtures were quantified with optical density at 620nm by subtracting negative control.

Flow cytometric analysis (FACS)

For cell cycle analysis, cells were collected by trypsinization and fixed in 70% ethanol at 4°C for 15min. After fixation, cells were washed with PBS, and the DNA was stained with propidium iodide (50µg/ml), in the presence of 5mM MgCl₂ and 10µg/ml RNase A in 10mM Tris HCl pH 7.5. DNA content was assessed on a FACS Calibur (Becton-Dickinson).

For BrdU pulse-chase proliferation assays, cells were pulse labeled with 10µM BrdU (Roche) for 1h, fixed in 70% ethanol, and incubated in 2M HCl for 30min. Cells were incubated with mouse antibody against BrdU (1:100) for 1h. For EdU analysis, cells were either pulsed for 10min with 10µM EdU or 24h. Cells were fixed in 70% ethanol and incubated with mouse anti-g-H2AX (1:100, Millipore) or mouse anti-p21^{WAF1} (1:100, SantaCruz) for 30min, followed by further 15min incubation with Alexa Fluor 488 anti-mouse IgG (1:100, Invitrogen) or anti-mouse E-Phycoerythrin (1:100, Invitrogen). EdU was detected with a Click-iT EdU Cell Proliferation Assay kit (Invitrogen). For MCM2 staining, cells were washed in wash buffer (1% (w/v) BSA in PBS) and unbound MCM2-7 was extracted in CSK buffer (10 mM HEPES, 100mM NaCl, 3mM MgCl₂, 1mM EGTA, 300mM sucrose, 1% BSA, 0.2% Triton-X100, 1mM DTT, 1mM PMSF, 10ng/ml Pepstatin, 10 ng/ml Leupeptin and 10 ng/ml Aprotinin) on ice for 10 min. Extracted cells were fixed in 2% (v/v) paraformaldehyde in PBS for 10 min at 37°C, washed twice and stored in wash buffer at 4°C until staining. CSK extracted fixed cells were permeabilised in 70% ethanol, washed and incubated for 1h at RT with mouse anti-human MCM2 (1:500, BD Biosciences). Next cells were washed and incubated for 30 min at RT in dark with Alexa Fluor 488 goat anti-mouse IgG (1:500, Life Technologies), before being washed twice and resuspended in 50 µg/ml 7-AAD (7-aminoactinomycin D, Life Technologies) diluted in wash buffer. Samples were analysed using FACS Canto (Becton Dickinson) and BD FACS DIVA software (BD Biosciences). Data analysis was performed using Flowjo (v7.6.5, Tree Star Inc.).

Comet Assay

Comet assay was performed as previously described (Sideridou et al. 2011). Briefly, 10⁵ Saos2 p21^{WAF1/Cip1} cells induced or non-induced with doxycyclin were seeded in 60mm dishes, respectively. Two days later cells were trypsinized, centrifuged, resuspended in 2ml PBS and kept on ice for 10min. Viable cells were counted using Trypan blue and PBS was added to adjust the number of cells to 10⁵ in 500µl of PBS. 50µl of PBS containing cells were then mixed with equal volume of low-melting agarose 1,7% (w/v) and were embedded in plugs. Plugs were incubated in 50ml of lysis solution (100mM TrisHCl pH 10.0, 100mM EDTA, 2.5M NaCl, 1% (v/v) Triton X100, 10% (v/v) DMSO) for 1h on ice and in dark. After lysis, plugs were washed twice in 1xTBE for 1h on ice in dark. Finally, plugs were washed and incubated in ice-cold alkaline denaturation buffer (300mM NaOH, 1mM EDTA, pH 13) for 45 min on ice in the dark. Plugs were mounted onto 1% agarose coated slides that were placed into a 30-cm horizontal constant-field gel electrophoresis chamber in ice-cold

alkaline denaturation buffer for 20min at 0.7 V/cm and at 4°C. After electrophoresis, slides were dehydrated in ice-cold ethanol (100%) for 10min and allowed to dry in the dark. After 24h slides were rehydrated in 5ml of deionized water for 10min and 40µl of 1x SYBR Green (Invitrogen) were applied on each plug. Cells were observed under a fluorescence microscope equipped with a monochrome CCD camera. Analysis was conducted using the Cometscore software (Tritek).

Pulse-field gel electrophoresis (PFGE)

PFGE was performed as previously described (Beck et al, 2010). Briefly cells were embedded in a 0.8% agarose plugs (2.5×10^5 cells/plug), digested in lysis buffer [100mM EDTA, 1% (w/v) sodium lauryl sarcosine, 0.2% (w/v) sodium deoxycholate and 1mg/ml proteinase K] at 37°C for 48h and washed in 10mM Tris-HCl pH8.0 and 100mM EDTA. Electrophoresis was performed at 15°C in 0.8% (w/v) PFGE certified agarose (Bio-Rad Laboratories) with Tris-borate/EDTA buffer employing a CHEF DR III apparatus (9h, 120°, 5.5V/cm, 30-18s switch time)(Bio-Rad). The gel was stained with ethidium bromide.

DNA fiber spreadings

DNA fiber assay was performed as previously described with slight modifications (Bartkova et al. 2006). Briefly, Saos2 p21^{WAF1/Cip1} cells were grown in the presence or absence of doxycyclin for 4 days and then pulsed-labeled with 25µM CldU for 20min, and then labelled with 250µM IdU for 20min. The cells were then harvested and lysed on glass slides in spreading buffer. The DNA was denatured and stained with rat anti-BrdU/CldU (1:1000, Immunologicals Direct) and mouse anti-IdU/BrdU (1:500, Becton Dickinson) primary antibodies.

Senescence staining

Saos2 p21^{WAF1} cells were Doxycyclin induced at the time points indicated (Fig. 5), as described above. Subsequently, cells grown on coverslips were transfected with control siRNA, or anti-Cdc6 and/or anti-Cdt1 siRNAs. Control (OFF) and induced cells were fixed in 1% paraformaldehyde and then processed for Sa-β-gal or Sudan Black B staining and counterstained with nuclear fast red, as described elsewhere (Georgakopoulou et al, 2013). Only cytoplasmic staining was scored as positive signal.

MTT Assay

Cytotoxicity was estimated by the MTT assay (Eliades et al. 2009). Briefly, cells were plated in 96-well, flat-bottomed microplates at a density of approximately 15,000 cells/cm², in DMEM containing 10% FBS. 24h after plating, the medium was replaced containing the chemotherapeutic agents at the concentrations indicated. After 72h of incubation, the medium was replaced with MTT dissolved at a final concentration of 1 mg/ml in serum-free, phenol-red-free DMEM, for further 4h incubation. Then, the MTT formazan was solubilized in isopropanol, and the optical density was measured at a wavelength of 550nm and a reference wavelength of 690nm.

Isolation of nucleic acids

DNA extraction: cells were lysed in 50mM Tris-HCl (pH 8.0), 150mM NaCl, 5mM EDTA, 1% SDS in the presence of proteinase K (0.1mg/ml) until completely dissolved. DNA was extracted with phenol/chloroform and RNase (Sigma) digestion, followed by ethanol precipitation (Liontos et al. 2007).

RNA was extracted with the RNeasy Mini Kit (Qiagen). cDNA was generated with Superscript® II Reverse Transcriptase (Invitrogen) and oligo-dT (Invitrogen).

cDNA preparation and real time RT-PCR

Real-time reverse transcription PCR (RT-PCR) analysis employed the Platinum SYBR Green qPCR SuperMix-UDG (Invitrogen) and was run on an ABI Prism 7300 cycler (Applied Biosystems)(Evangelou et al. 2013). Primer sequences and annealing temperatures are provided in **Supplemental Table 7**. Results are presented as n-fold changes for the various time points after p21^{WAF1/Cip1} induction versus the values of the non-induced sample. Mean value was calculated from three independent measurements.

Soft agar assay

Dishes of 60mm were layered with 3ml 0.7% (w/v) SeaPlaque Low Melting Point Agarose (Lonza) dissolved in serum-containing medium (Sideridou et al. 2011). Saos2 Tet-ON p21^{WAF1/Cip1} cells (25×10^3) were then mixed with 1.6 ml of 0.35% (w/v) warm agar (42°C) in serum-containing medium and plated on the solidified agarose layer. Agar was added weekly and foci were enumerated on day 45. Experiments were performed in three independent replicates.

Invasion assay

Saos2 Tet-ON p21^{WAF1/Cip1} cells were seeded on top of matrigel-coated transwell (Beckton Dickinson) at 5,000 cells per 24-well with serum-free medium (Sideridou et al. 2011). Transwells were embedded into complete (full-serum) culture medium and cells were allowed to invade for 24h. Finally, cells at the top side of the well were scraped, and cells (invading) on the bottom side of the well were stained with Giemsa, photographed and counted. Three independent measurements were averaged and the corresponding s.e. is also reported.

Chromatin immunoprecipitation (ChIP) assay

ChIP assay was performed as previously described (Sideridou et al. 2011). A 100 bp fragment in the *Rad51* promoter and a 110bp amplicon, located approx 1000bp from the transcription start site (Fig. 4h), were amplified. Primers and annealing temperatures are provided in **Supplemental Table 7**. As inputs we used products that corresponded to PCR reactions containing 1% of the total chromatin extract used in the immunoprecipitation reactions.

Proteomics

Protein Digestion and iTRAQ Labeling—Cell pellets were dissolved in 200 μ L dissolution buffer [0.5M triethylammonium bicarbonate (TEAB), 0.05% SDS] with vortex mixing, 5min heating at 90°C and pulsed probe sonication for 20sec. Non dissolved cellular debris was separated from the protein solution with centrifugation at 13,000 rpm for 10min. For each sample a total protein amount of 100 μ g was measured with Bradford assay (Bio-Rad Protein Assay) and diluted with the addition of dissolution buffer up to a final 20 μ L volume. The reduction, cysteine blocking and iTRAQ labeling were conducted in accordance to manufacturer protocols (AB Sciex) as reported before (Delehouze et al. 2013; Papachristou et al. 2013). The pooled whole sample mixture was split in two equal parts, lyophilized and stored at -20°C.

High-pH Reverse Phase (RP) Peptide Fractionation—To one-half of the iTRAQ labeled sample described above, peptide fractionation was performed with reverse phase C₁₈-liquid chromatography (XBridge, 150x4.6 mm, 3.5 μ m; Waters) under alkaline conditions using the Dionex P680 pump equipped with a PDA-100 photodiode array detector as described before (Papachristou et al. 2013). Fractions were collected in 1min intervals and lyophilized to dryness and stored at -20°C until LC-MS analysis.

Hydrophilic Interaction Chromatography (HILIC) peptide fractionation—The second half of the iTRAQ labeled peptides was fractionated with Hydrophilic Interaction Chromatography (Core-Shell Kinetex, 150x2.1 mm, 2.6 μ m, 100 Å column) (Phenomenex) under acidic conditions using the Dionex P680 pump equipped with a PDA-100 photodiode array detector using gradient conditions as described by the authors (Garbis. et al. 2011; Delehouze et al. 2013). Fractions were collected in a peak dependent manner based on the absorbance signal response at 280nm, lyophilized to dryness and stored at -20°C until LC-MS analysis.

LC-MS Analysis—All LC-MS experiments were performed with nano-Ultra-high performance liquid chromatography (Ultimate 3000, Thermo Fisher Scientific) hyphenated with a nanospray ionization / ultra high-resolution LTQ FT-Orbitrap-Elite mass spectrometer (Thermo Fisher Scientific) system. Individual high-pH RP and HILIC peptide fractions were reconstituted in 30 μ L loading solution (aqueous 0.1% formic acid, 2% acetonitrile). Sample volumes of 2 μ L volume were injected and loaded for 8min on the Acclaim PepMap 100, 100 μ m x 2 cm C18, 5 μ m, 100 Å trapping column with the loading pump at a flow rate of 5 μ L/min. The trapped peptides were then gradient separated with reverse phase C₁₈-nano-capillary liquid chromatography (Acclaim PepMap RSLC, 75 μ m x 25 cm, 2 μ m, 100 Å) (Thermo Fisher Scientific) under acidic conditions, and nano-spray ionized with the PicoTip emitter (FS360-20-10-D-20-C7, New Objective), as previously described (Papachristou et al. 2013). Gaseous phase transition of the separated peptides was achieved with positive ion electrospray ionization applying a voltage of 2.5 kV. For every MS survey scan, the top 10 most abundant multiply charged precursor ions between m/z ratio 350 and 1900 and intensity threshold 500 counts were selected with FT mass resolution of 240,000 and subjected to HCD fragmentation with an isolation window of 1.2 Da. Tandem mass spectra were acquired with FT resolution of 15,000 within m/z range of 100-1900. Normalized

collision energy was set to 35 and already targeted precursors were dynamically excluded for further isolation and activation for 30sec with 5ppm mass tolerance.

Database search—The HCD tandem mass spectra collected from RP and HILIC fractions were submitted to Sequest search engine implemented on the Proteome Discoverer software v1.3.0.339 for peptide and protein identifications. All spectra were searched against a UniProt Fasta file containing 20,200 human reviewed entries. The Sequest node included the following parameters: Precursor Mass Tolerance 10 ppm, Fragment Mass Tolerance 20 mmu, Dynamic Modifications were Oxidation of M (+15.995 Da), Deamidation of N, Q (+0.984 Da), Phosphorylation of S (+79.966 Da) and Static Modifications were iTRAQ8plex at any N-Terminus, K, Y (+304.205 Da) and Methylthio at C (+45.988 Da). The Percolator node was used for the determination of the confidence level for peptide identifications with decoy database searching. Strict FDR was set to 0.01, relaxed FDR was set to 0.05 and validation was based on q-Value. The Reporter Ion Quantifier node included a custom iTRAQ 8plex (Thermo Scientific Instruments) Quantification Method, integration window tolerance 20 ppm and integration method Most Confident Centroid. Protein ratios were normalized to protein median and peptides with missing iTRAQ channels were excluded from relative protein quantification. Phosphorylation localization probability was estimated with the phosphoRS node.

High-throughput whole genome analyses

aCGH analysis—Genomic DNA from 30days induced and non-induced Saos2 p21^{WAF1/Cip1} cells was extracted using the BioRobot® M48 System (Qiagen) and the MagAttract® DNA Blood Midi M48 Kit (Qiagen). Quality and quantity of the DNA samples was determined on a NanoDrop ND-1000 spectrophotometer.

Agilent Human Genome CGH 4x180K and 1x1MK microarrays were used (Agilent Technologies). The 4x180K platform is composed of more than 170,000 60-mer oligonucleotide probes with average spatial resolution of 13-25 Kb (NCBI build 37, hg19). The 1x1MK platform is composed of more than 963,000 60-mer oligonucleotide probes for the mapped genes or unique DNA sequences with an average spatial resolution of 2.1 Kb (NCBI build 37, hg19). Labelling and hybridization was carried out according to manufacturer's guidelines. Data were processed using Feature Extraction 10.7.3.1 and analysed using Cytogenomics 2.7.22.0 software (Agilent) with the following settings: Algorithm: ADM-1, Threshold: 6.7, with a minimum of 4 probes for a region to be included. Centralization and fuzzy zero corrections were applied to remove putative variant intervals with small average log₂ ratios.

Deep (next generation) sequencing (NGS)—Genomic DNA from 30 days induced and non-induced Saos2 p21^{WAF1/Cip1} cells was used for whole genome sequencing. The library preparation and the whole genome sequencing were carried out in EMBL Genecore facility according to the Illumina platform.

Whole genome sequencing was performed in “OFF” and “ON” cells achieving a 30x coverage of the human genome. Paired-end 2x100 bp was performed with the use of Illumina Hi-seq 2000. Quality control of the reads was performed with the use of fastqc

software (**Ramirez et al. 2013**) and alignment to the human genome (GRCH37/hg19 version) was performed with bowtie 2 algorithm (**Langmead and Salzberg 2012**) with the use of the following code (-D 5 -R 1 -N 1 -L 22 -i S0,2,..50 – trim3 25 -I 0 -X 500). Samtools (**Li et al. 2009**) was used to convert sam files to bam and for sorting bam files. Breakdancer software (**Chen et al. 2009**) was utilized in order to identify SV (intra- & inter-chromosomal translocations, deletions, insertions and inversions). Taking into consideration the spacing between the two paired ends in the sequencing process we filtered out SVs with less than 500 bp length (deletion, insertion, inversion).

Use of breakdancer with default parameters led to the identification of 103 new inter-chromosomal translocations in “ON” cells versus “OFF” cells.

In order to identify microhomology regions in the 103 inter-chromosomal translocations we used the coordinates from the breakdancer output and extended 30bp on both sides of the breakpoint-junction. Clustal W was used for aligning the regions around the 103 breakpoint junctions. Microhomology regions identified on the breakpoint spanned from 1-9 bp.

Expression microarray analysis—Total RNA was isolated from non-induced and 12h, 48h and 96h induced Saos2-p21^{WAF1/Cip1} Tet ON cells, using RNeasy Total RNA kit (Qiagen). Microarray analysis was performed by the microarray unit of CBM Core Facility Italy (<http://www.cbm.fvg.it>) using Illumina’s Whole-Genome Expression Beadchip. This platform targets more than 47,000 probes derived from the National Center for Biotechnology Information Reference Sequence (NCBI) RefSeq Release 38 and other sources. Integrity of total RNA was evaluated using capillary electrophoresis (Bioanalyzer 2100, Agilent) and quantified using a Nanodrop 1000. Aliquots of RNA (250 ng) samples were amplified according to the specifications of the Illumina® TotalPrep™ RNA Amplification Kit (Ambion) to produce a pool of biotin-labeled RNA corresponding to the polyadenylated (mRNA) fraction. The cRNA samples were applied to whole-genome HumanHT-12 v4.0 arrays (Illumina) and hybridized according to manufacturer’s specification. Each array on the BeadChip targets over 29,000 transcripts using 3-micron beads bearing covalently attached 50-base oligonucleotide probes. Each probe interrogates a single gene, and each bead type is represented with an average 30-fold redundancy on every array. The BeadChips were scanned with the Illumina’s Beadarray system scanner (Illumina). The hybridization images signal intensity was extracted and background subtracted using Illumina Inc. BeadStudio software v3.3.7. Data were checked for the Illumina internal quality control.

To compare the non-induced and “escaped” Saos2-p21^{WAF1/Cip1} Tet-ON cells gene expression profile, total RNA was extracted using Trizol (Life Technologies) and chloroform/isoamyl alcohol (49:1). All RNA samples had a 28S/18S rRNA ratio close to 2 on 1.5% agarose gels, and absorbance ratios 260/280 nm between 1.9 and 2.1. Targets were prepared using the GeneChip Whole Transcriptome (WT) Plus reagent kit and hybridized to Affymetrix GeneChip Human Transcriptome array 2.0 (HTA 2.0) that contains over 6 million probes representative of 245,000 coding transcripts and 40,000 non-coding transcripts, as well as 339,000 probe sets covering exon-exon junctions.

Molecular Cytogenetics

Saos2 Tet-ON p21^{WAF1/Cip1} cell cultures were exposed to colcemid (0.1 µg/ml) (Gibco) for 12h, at 37°C in 5% CO₂. Cells were harvested by trypsinization (Gibco), suspended in medium, and spun down (10min, 1000 rpm). Supernatant was removed and 5ml of 0.075M KCl at RT was added drop by drop. Cells were incubated for 20min at RT, and then 1ml of fixative [3×methanol (Applichem)/1×CH₃COOH (Merck)] was added. Cells were spun down (10min at 1000 rpm), supernatant was removed, fixative was added and the cells were re-centrifuged for 10min at 1000 rpm. Finally, cells were dropped onto wet microscope slides and left to air-dry. For analysis, we combined inverted 4',6'-diamidino-2-phenylindole (DAPI) Banding, and molecular karyotyping by M-FISH (MetaSystems). Multicolor FISH was performed according to manufacturer's protocols (MetaSystems). For inverted DAPI banding, slides were counterstained and mounted with 0.1 µg/ml DAPI in Vectashield antifade medium (Vector Laboratories). Analyses were performed using a 63× magnification lens on a fluorescent Axio-Imager Z1, Zeiss microscope, equipped with a MetaSystems charge-coupled device (CCD) camera and the MetaSystems Isis software.

Bioinformatic analysis

Transcriptome and Proteome analyses from non-induced, 12h, 48h and 96h induced Saos2-p21^{WAF1/Cip1} Tet-ON cells—Protein and gene ratios (time-point/time-0) were log(2) transformed and centered. The 95% confidence intervals of the averaged log₂-ratios were calculated for each time-point to determine the statistically significant thresholds of over-expression and suppression. Time-series log₂-ratio sets for each protein and gene were analyzed by ANOVA to detect a statistically significant change anywhere within the three time-points taking into account that each time point had 2 replicates for proteomics and 3 replicates for micro-arrays. Proteins and genes whose log₂-ratio was above or below the aforementioned over-expression and suppression thresholds respectively and their ANOVA p-value was <0.05 were considered regulated. All calculations were performed with R. The “Gene-Set Enrichment Analysis” algorithm with False-Discovery-Rate (FDR) correction was applied on the Gene-Ontology Biological-Process sets through the Ariadne Genomics Pathway Studio v9.0 to detect which functional groups were non-randomly impacted (FDR adjusted p-values <0.05 were considered significant). Pathway visualization for all data-sets was performed with Ariadne Genomics Pathway Studio v9.0.

Transcriptome analysis from non-induced and “escaped” Saos2-p21^{WAF1/Cip1} Tet-ON cells—The Affymetrix .CEL files generated were analyzed using Partek Genomics Suite. Following normalization with the Robust Multi-array Average (RMA) algorithm, a full set of quality control analysis (which led to the exclusion of one sample) was performed followed by Power Analysis, Principal Component Analysis, and one-way ANOVA analyses for a range of different fold (2-5) and false discovery rate (0.001-0.05) thresholds. For the Hierarchical Clustering of significantly changed transcripts the level of each transcript was standardized to mean 0 and standard deviation of 1.

The significantly changed transcripts identified for thresholds of 2fold and 0.01FDR were analyzed using the Ingenuity Pathway Analysis software. Specifically, all direct and indirect relationships were considered, all data sources, and only experimentally observed

information. Five levels of analysis were performed: Functional Categorization, Canonical Pathway analysis, Network analysis, Regulator Effects analysis, and Upstream Regulator analysis.

Deep (next generation) sequencing (NGS) and aCGH data comparison—To compare the NGS data with the aCGH one regarding DNA copy number aberrations in “escaped” (ON) cells versus control (OFF) cells, the NGS data were processed as follows: a) Genomic regions presenting less than 10-times coverage were filtered out to ensure data high-quality, b) The log₂ of the ratio of the normalized reads in the “escaped” cells over the normalized number of reads in the control-cells was calculated, c) For each chromosome the aforementioned log₂-ratios underwent DNA copy-number segmentation analysis utilizing the circular binary segmentation algorithm through the Bioconductor package “DNAcopy” (Venkatraman and Olshen 2007). All statistical analysis was carried out with R.

Pathway Analysis—Proteins and genes from proteomics and transcriptomics respectively were imported into Ariadne Pathway Studio v9.0 and analyzed for biological context against Ariadne's canonical cell-process pathways utilizing the in-built Gene Set Enrichment Analysis procedure which uses the Mann-Whitney U-Test to evaluate the pathways enrichment p-values that were adjusted according to the FDR correction for multiple testing. The confidence-level was 95%; thus p-values <0.05 were considered significant.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We would like to thank Drs A. Kotsinas, K Evangelou, M Sideridou and A Georgakilas for their valuable support to this work. We would like to thank Prof A. Dutta for kindly providing the vectors with wt and PIP mutated domain of p21^{WAF1/Cip1}, Dr G. Blandino for the H1299 p21^{WAF1/Cip1}-Ponesterone-ON cells and Dr Z. Lygerou for the secondary antibodies employed in the IF analyses. This work received funding from the European Union's Seventh Framework Programme for research, technological development and demonstration under grant agreement No 284460 (INsPiRE) and the Greek GSRT programme of Excellence II (Aristeia II).

Abbreviations

RLFs	Replication licensing factors
MUS81-EME1	methymethanesulphonate-MMS and ultraviolet-sensitive81-MUS81 and essential meiotic endonuclease1-EME1
EGF	epidermal growth factor
FGF2	fibroblast growth factor-2
TGF-β	transforming growth factor-β
HJ	Holliday Junction
NHEJ	non-homologous end joining

PIP

PCNA-Interacting-Protein motif

References

- Abbas T, Dutta A. p21 in cancer: intricate networks and multiple activities. *Nat Rev Cancer*. 2009; 9:400–14. [PubMed: 19440234]
- Abbas T, Keaton MA, Dutta A. Genomic instability in cancer. *Cold Spring Harb Perspect Biol*. 2013; 5:a012914. [PubMed: 23335075]
- Abbas T, Sivaprasad U, Terai K, Amador V, Pagano M, Dutta A. PCNA-dependent regulation of p21 ubiquitylation and degradation via the CRL4Cdt2 ubiquitin ligase complex. *Genes Dev*. 2008; 22:2496–506. [PubMed: 18794347]
- Barlow JH, Faryabi RB, Callén E, Wong N, Malhowski A, Chen HT, Gutierrez-Cruz G, Sun HW, McKinnon P, Wright G, Casellas R, et al. Identification of early replicating fragile sites that contribute to genome instability. *Cell*. 2013; 152:620–32. [PubMed: 23352430]
- Bartkova J, Rezaei N, Lontos M, Karakaidos P, Kletsas D, Issaeva N, Vassiliou LV, Kolettas E, Niforou K, Zoumpourlis VC, Takaoka M, et al. Oncogene-induced senescence is part of the tumorigenesis barrier imposed by DNA damage checkpoints. *Nature*. 2006; 444:633–7. [PubMed: 17136093]
- Bates S, Ryan KM, Phillips AC, Vousden KH. Cell cycle arrest and DNA endoreduplication following p21Waf1/Cip1 expression. *Oncogene*. 1998; 17:1691–703. [PubMed: 9796698]
- Beauséjour CM, Krtolica A, Galimi F, Narita M, Lowe SW, Yaswen P, Campisi J. Reversal of human cellular senescence: roles of the p53 and p16 pathways. *EMBO J*. 2003; 22:4212–22. [PubMed: 12912919]
- Beck H, Nahse V, Larsen MS, Groth P, Clancy T, Lees M, Jorgensen M, Helleday T, Syljuasen RG, Sorensen CS. Regulators of cyclin-dependent kinases are crucial for maintaining genome integrity in S phase. *J Cell Biol*. 2010; 188:629–638. [PubMed: 20194642]
- Benson EK, Mungamuri SK, Attie O, Kracikova M, Sachidanandam R, Manfredi JJ, Aaronson SA. p53-dependent gene repression through p21 is mediated by recruitment of E2F4 repression complexes. *Oncogene*. 2013 Oct 7.
- Bischoff FZ, Yim SO, Pathak S, Grant G, Siciliano MJ, Giovanella BC, Strong LC, Tainsky MA. Spontaneous abnormalities in normal fibroblasts from patients with Li-Fraumeni cancer syndrome: aneuploidy and immortalization. *Cancer Res*. 1990; 50:7979–84. [PubMed: 2253239]
- Blow JJ, Gillespie PJ. Replication licensing and cancer--a fatal entanglement? *Nat Rev Cancer*. 2008; 8:799–806. [PubMed: 18756287]
- Cha HH, Cram EJ, Wang EC, Huang AJ, Kasler HG, Firestone GL. Glucocorticoids stimulate p21 gene expression by targeting multiple transcriptional elements within a steroid responsive region of the p21waf1/cip1 promoter in rat hepatoma cells. *J Biol Chem*. 1998; 273:1998–2007. [PubMed: 9442036]
- Chang BD, Xuan Y, Broude EV, Zhu H, Schott B, Fang J, Roninson IB. Role of p53 and p21waf1/cip1 in senescence-like terminal proliferation arrest induced in human tumor cells by chemotherapeutic drugs. *Oncogene*. 1999; 18:4808–18. [PubMed: 10490814]
- Couch FB, Bansbach CE, Driscoll R, Luzwick JW, Glick GG, Bétous R, Carroll CM, Jung SY, Qin J, Cimprich KA, Cortez D. ATR phosphorylates SMARCA1 to prevent replication fork collapse. *Genes Dev*. 2013; 27:1610–23. [PubMed: 23873943]
- Delehouzé C, Godl K, Loaëc N, Bruyère C, Desban N, Oumata N, Galons H, Roumeliotis TI, Giannopoulou EG, Grenet J, Twitchell D, et al. *Oncogene*. 2013
- Dimri GP, Lee X, Basile G, Acosta M, Scott G, Roskelley C, Medrano EE, Linskens M, Rubelj I, Pereira-Smith O, et al. A biomarker that identifies senescent human cells in culture and in aging skin in vivo. *Proc Natl Acad Sci U S A*. 1995; 92:9363–7. [PubMed: 7568133]
- Duursma A, Agami R. p53-Dependent regulation of Cdc6 protein stability controls cellular proliferation. *Mol Cell Biol*. 2005; 25:6937–47. [PubMed: 16055707]
- Eliades T, Pratsinis H, Athanasiou AE, Eliades G, Kletsas D. Cytotoxicity and estrogenicity of Invisalign appliances. *Am J Orthod Dentofacial Orthop*. 2009; 136:100–3. [PubMed: 19577155]

- Essers J, Theil AF, Baldeyron C, van Cappellen WA, Houtsmuller AB, Kanaar R, Vermeulen W. Nuclear dynamics of PCNA in DNA replication and repair. *Mol Cell Biol.* 2005; 25:9350–9. [PubMed: 16227586]
- Evangelou K, Bartkova J, Kotsinas A, Pateras IS, Lontos M, Velimezi G, Kosar M, Liloglou T, Trougakos IP, Dyrskjot L, Andersen CL, et al. The DNA damage checkpoint precedes activation of ARF in response to escalating oncogenic stress during tumorigenesis. *Cell Death Differ.* 2013; 20:1485–97. [PubMed: 23852374]
- Garbis SD, Roumeliotis TI, Tyritzis SI, Zorpas KM, Pavlakis K, Constantinides CA. A novel multidimensional protein identification technology approach combining protein size exclusion prefractionation, peptide zwitterion-ion hydrophilic interaction chromatography, and nano-ultraperformance RP chromatography/nESI-MS2 for the in-depth analysis of the serum proteome and phosphoproteome: application to clinical sera derived from humans with benign prostate hyperplasia. *Anal Chem.* 2011; 83:708–18. [PubMed: 21174401]
- Georgakopoulou EA, Tsimaratou K, Evangelou K, Fernandez Marcos PJ, Zoumpourlis V, Trougakos IP, Kleitsas D, Bartek J, Serrano M, Gorgoulis VG. Specific lipofuscin staining as a novel biomarker to detect replicative and stress-induced senescence. A method applicable in cryo-preserved and archival tissues. *Aging (Albany NY).* 2013; 5:37–50. [PubMed: 23449538]
- Gonzalez S, Klatt P, Delgado S, Conde E, Lopez-Rios F, Sanchez-Cespedes M, Mendez J, Antequera F, Serrano M. Oncogenic activity of Cdc6 through repression of the INK4/ARF locus. *Nature.* 2006; 440:702–6. [PubMed: 16572177]
- Gorgoulis VG, Zacharatos P, Kotsinas A, Mariatos G, Liloglou T, Vogiatzi T, Foukas P, Rassidakis G, Garinis G, Ioannides T, Zoumpourlis V, et al. Altered expression of the cell cycle regulatory molecules pRb, p53 and MDM2 exert a synergetic effect on tumor growth and chromosomal instability in non-small cell lung carcinomas (NSCLCs). *Mol Med.* 2000; 6:208–37. [PubMed: 10965496]
- Halazonetis TD, Gorgoulis VG, Bartek J. An oncogene-induced DNA damage model for cancer development. *Science.* 2008; 319:1352–5. [PubMed: 18323444]
- Hanada K, Budzowska M, Davies SL, van Drunen E, Onizawa H, Beverloo HB, Maas A, Essers J, Hickson ID, Kanaar R. The structure-specific endonuclease Mus81 contributes to replication restart by generating double-strand DNA breaks. *Nat Struct Mol Biol.* 2007; 14:1096–104. [PubMed: 17934473]
- Hastings PJ, Ira G, Lupski JR. A microhomology-mediated break-induced replication model for the origin of human copy number variation. *PLoS Genet.* 2009; 5:e1000327. [PubMed: 19180184]
- Havens CG, Walter JC. Mechanism of CRL4(Cdt2), a PCNA-dependent E3 ubiquitin ligase. *Genes Dev.* 2011; 25:1568–82. [PubMed: 21828267]
- Holland AJ, Cleveland DW. Chromoanagenesis and cancer: mechanisms and consequences of localized, complex chromosomal rearrangements. *Nat Med.* 2012; 18:1630–8. [PubMed: 23135524]
- Iraqui I, Chekkal Y, Jmari N, Pietrobon V, Fréon K, Costes A, Lambert SA. Recovery of arrested replication forks by homologous recombination is error-prone. *PLoS Genet.* 2012; 8:e1002976. [PubMed: 23093942]
- Jørgensen S, Eskildsen M, Fugger K, Hansen L, Larsen MS, Kousholt AN, Syljuåsen RG, Trelle MB, Jensen ON, Helin K, Sørensen CS. SET8 is degraded via PCNA-coupled CRL4(CDT2) ubiquitylation in S phase and after UV irradiation. *J Cell Biol.* 2011; 192:43–54. [PubMed: 21220508]
- Karakaidos P, Taraviras S, Vassiliou LV, Zacharatos P, Kastrinakis NG, Kougiou D, Kouloukoussa M, Nishitani H, Papavassiliou AG, Lygerou Z, Gorgoulis VG. Overexpression of the replication licensing regulators hCdt1 and hCdc6 characterizes a subset of non-small-cell lung carcinomas: synergistic effect with mutant p53 on tumor growth and chromosomal instability-evidence of E2F-1 transcriptional control over hCdt1. *Am J Pathol.* 2004; 165:1351–65. [PubMed: 15466399]
- Kim Y, Starostina NG, Kipreos ET. The CRL4Cdt2 ubiquitin ligase targets the degradation of p21Cip1 to control replication licensing. *Genes Dev.* 2008; 22:2507–19. [PubMed: 18794348]
- Lontos M, Koutsami M, Sideridou M, Evangelou K, Kleitsas D, Levy B, Kotsinas A, Nahum O, Zoumpourlis V, Kouloukoussa M, Lygerou Z, et al. Deregulated overexpression of hCdt1 and hCdc6 promotes malignant behavior. *Cancer Res.* 2007; 67:10899–909. [PubMed: 18006835]

- Liontos M, Niforou K, Velimezi G, Vougas K, Evangelou K, Apostolopoulou K, Vrtel R, Damalas A, Kontovazenitis P, Kotsinas A, Zoumpourlis V, et al. Modulation of the E2F1-driven cancer cell fate by the DNA damage response machinery and potential novel E2F1 targets in osteosarcomas. *Am J Pathol.* 2009; 175:376–91. [PubMed: 19541929]
- Mailand N, Diffley JF. CDKs promote DNA replication origin licensing in human cells by protecting Cdc6 from APC/C-dependent proteolysis. *Cell.* 2005; 122:915–26. [PubMed: 16153703]
- Marti A, Wirbelauer C, Scheffner M, Krek W. Interaction between ubiquitin-protein ligase SCF^{SKP2} and E2F-1 underlies the regulation of E2F-1 degradation. *Nat Cell Biol.* 1999; 1:14–9. [PubMed: 10559858]
- Murfuni I, Nicolai S, Baldari S, Crescenzi M, Bignami M, Franchitto A, Pichierri P. The WRN and MUS81 proteins limit cell death and genome instability following oncogene activation. *Oncogene.* 2013; 32:610–20. [PubMed: 22410776]
- Neelsen KJ, Zanini IM, Mijic S, Herrador R, Zellweger R, Ray Chaudhuri A, Creavin KD, Blow JJ, Lopes M. Deregulated origin licensing leads to chromosomal breaks by rereplication of a gapped DNA template. *Genes Dev.* 2013a; 27:2537–42. [PubMed: 24298053]
- Neelsen KJ, Zanini IM, Herrador R, Lopes M. Oncogenes induce genotoxic stress by mitotic processing of unusual replication intermediates. *J Cell Biol.* 2013b; 200:699–708. [PubMed: 23479741]
- Negrini S, Gorgoulis VG, Halazonetis TD. Genomic instability-an evolving hallmark of cancer. *Nat Rev Mol Cell Biol.* 2010; 11:220–8. [PubMed: 20177397]
- Ottaviani D, LeCain M, Sheer D. The role of microhomology in genomic structural variation. *Trends Genet.* 2014; 30:85–94. [PubMed: 24503142]
- Papachristou EK, Roumeliotis TI, Chrysagi A, Trigoni C, Charvalos E, Townsend PA, Pavlakis K, Garbis SD. The shotgun proteomic study of the human ThinPrep cervical smear using iTRAQ mass-tagging and 2D LC-FT-Orbitrap-MS: the detection of the human papillomavirus at the protein level. *J Proteome Res.* 2013; 12:2078–89. [PubMed: 23510160]
- Pateras IS, Apostolopoulou K, Niforou K, Kotsinas A, Gorgoulis VG. p57KIP2: "Kip"ing the cell under control. *Mol Cancer Res.* 2009; 7:1902–19. [PubMed: 19934273]
- Perkins ND, Felzien LK, Betts JC, Leung K, Beach DH, Nabel GJ. Regulation of NF-kappaB by cyclin-dependent kinases associated with the p300 coactivator. *Science.* 1997; 275:523–7. [PubMed: 8999795]
- Petermann E, Helleday T. Pathways of mammalian replication fork restart. *Nat Rev Mol Cell Biol.* 2010; 11:683–7. [PubMed: 20842177]
- Pines J, Hunter T. Human cyclins A and B1 are differentially located in the cell and undergo cell cycle-dependent nuclear transport. *J Cell Biol.* 1991; 115:1–17. [PubMed: 1717476]
- Porter ACG. Preventing DNA over-replication: a Cdk perspective. *Cell Div.* 2008; 3:3. [PubMed: 18211690]
- Rivlin N, Brosh R, Oren M, Rotter V. Mutations in the p53 Tumor Suppressor Gene: Important Milestones at the Various Steps of Tumorigenesis. *Genes Cancer.* 2011; 2:466–74. [PubMed: 21779514]
- Roninson IB. Oncogenic functions of tumour suppressor p21(Waf1/Cip1/Sdi1): association with cell senescence and tumour-promoting activities of stromal fibroblasts. *Cancer Lett.* 2002; 179:1–14. [PubMed: 11880176]
- Rosai J. Rosai and Ackerman's Surgical Pathology. Elsevier; London: 2011.
- Sakellariou D, Chiourea M, Raftopoulou C, Gagos S. Alternative lengthening of telomeres: recurrent cytogenetic aberrations and chromosome stability under extreme telomere dysfunction. *Neoplasia.* 2013; 15:1301–13. [PubMed: 24339742]
- Schlisio S, Halperin T, Vidal M, Nevins JR. Interaction of YY1 with E2Fs, mediated by RYBP, provides a mechanism for specificity of E2F function. *EMBO J.* 2002; 21:5775–86. [PubMed: 12411495]
- Sideridou M, Zakopoulou R, Evangelou K, Liontos M, Kotsinas A, Rampakakis E, Gagos S, Kahata K, Grabusic K, Gkouskou K, Trougakos IP, et al. Cdc6 expression represses E-cadherin transcription and activates adjacent replication origins. *J Cell Biol.* 2011; 195:1123–40. [PubMed: 22201124]

- Sikder HA, Devlin MK, Dunlap S, Ryu B, Alani RM. Id proteins in cell growth and tumorigenesis. *Cancer Cell*. 2003; 3:525–30. [PubMed: 12842081]
- Stephens PJ, Greenman CD, Fu B, Yang F, Bignell GR, Mudie LJ, Pleasance ED, Lau KW, Beare D, Stebbings LA, McLaren S, et al. Massive genomic rearrangement acquired in a single catastrophic event during cancer development. *Cell*. 2011; 144:27–40. [PubMed: 21215367]
- Takeda DY, Dutta A. DNA replication and progression through S phase. *Oncogene*. 2005; 24:2827–43. [PubMed: 15838518]
- Terradas M, Martín M, Tusell L, Genescà A. DNA lesions sequestered in micronuclei induce a local defective-damage response. *DNA Repair (Amst)*. 2009; 8:1225–34. [PubMed: 19683478]
- Vaziri C1, Saxena S, Jeon Y, Lee C, Murata K, Machida Y, Wagle N, Hwang DS, Dutta A. A p53-dependent checkpoint pathway prevents rereplication. *Mol Cell*. 2003; 11:997–1008. [PubMed: 12718885]
- Velimezi G, Lontos M, Vougas K, Roumeliotis T, Bartkova J, Sideridou M, Dereli-Oz A, Kocylowski M, Pateras IS, Evangelou K, Kotsinas A, et al. Functional interplay between the DNA-damage-response kinase ATM and ARF tumour suppressor protein in human cancer. *Nat Cell Biol*. 2013; 15:967–77. [PubMed: 23851489]
- Venkatraman ES, Olshen AB. A faster circular binary segmentation algorithm for the analysis of array CGH data. *Bioinformatics*. 2007; 23:657–63. [PubMed: 17234643]
- Wang Y, Blandino G, Givol D. Induced p21waf expression in H1299 cell line promotes cell senescence and protects against cytotoxic effect of radiation and doxorubicin. *Oncogene*. 1999; 18:2643–9. [PubMed: 10353608]
- Warfel NA, El-Deiry WS. p21WAF1 and tumorigenesis: 20 years after. *Curr Opin Oncol*. 2013; 25:52–8. [PubMed: 23159848]
- Woo RA, Poon RY. Cyclin-dependent kinases and S phase control in mammalian cells. *Cell Cycle*. 2003; 2:316–24. [PubMed: 12851482]
- Wu Y1, Kantake N, Sugiyama T, Kowalczykowski SC. Rad51 protein controls Rad52-mediated DNA annealing. *J Biol Chem*. 2008; 283:14883–92. [PubMed: 18337252]
- Zhang CZ, Leibowitz ML, Pellman D. Chromothripsis and beyond: rapid genome evolution from complex chromosomal rearrangements. *Genes Dev*. 2013; 27:2513–30. [PubMed: 24298051]

Figure 1. The p21^{WAF1/Cip1} and Ki67 are co-expressed in a subset of atypical cells of high grade/poorly differentiated, advanced human carcinomas.

Serial-section immunohistochemical (IHC) analysis and double immunofluorescent (IF) analysis showed co-expression of p21^{WAF1/Cip1} and the mitotic marker Ki67 in a subset of large cancer cells with giant nuclei in **a)** head and neck squamous cell carcinomas (HNSCC) (N=20), **b)** lung squamous cell carcinomas (N=30) (inset depicts a large atypical cell with p21^{WAF1/Cip1}/Ki67 co-expression) and **c)** urothelial carcinomas (N=30). IHC and IF: Black and white thin arrows denote p21^{WAF1/Cip1} and Ki67 co-expressing cells, respectively; IF: white and yellow thick arrows depict cells with mutual exclusive p21^{WAF1/Cip1} and Ki67 expression, respectively. Bars in IHC panels: 50 μ m, IF panels: 50 μ m.

Figure 2. Prolonged stimulation of p21^{WAF1/Cip1} in p53 null environment up-regulates the Replication Licensing Factors (RLFs) at the protein level.

a. Heat-maps from the transcriptomic and proteomic analyses, at days 2 and 4, after p21^{WAF1/Cip1} induction in Saos2 p21^{WAF1/Cip1} Tet-ON cells. Set of significant genes that are overexpressed and underexpressed are shown, respectively. Schematic representation depicting significant genes that are up-regulated and down-regulated, along with their biochemical function, at day 4 upon p21^{WAF1/Cip1} induction in Saos2 p21^{WAF1/Cip1} TetON cells. **b.** A specific p21^{WAF1/Cip1} mutant (p21^{PCNA}: harboring Q144, M147, F150

substitutions to A in its PIP degron motif) abrogating its interaction with PCNA shifts CRL4^{CDT2}-mediated degradation towards Cdt1. Lysates from Saos2 cells, after transductions with corresponding retroviral-vectors, were separated by SDS-PAGE and immunoblotted to detect the indicated proteins. (Mock: pMSCV, p21^{WAF1}; pMSCV-p21^{WAF1/Cip1}, p21^{PCNA}: pMSCV-p21^{PCNA}) **c.** The status of p53 defines the ability of p21^{WAF1/Cip1} to regulate the levels of Cdt1 and Cdc6. Immunoblots (IBs) for Cdt1 and Cdc6 in Li-Fraumeni fibroblasts, H1299 p21^{WAF1/Cip1}-Ponesterone-ON and HT1080 p21^{WAF1/Cip1}-IPTG-ON cells challenged with p21^{WAF1/Cip1}. Actin serves as a loading control. **d.** Flow cytometry analysis (FACS) of HT1080 p21^{WAF1/Cip1}-IPTG-ON cells showed that induction of apoptosis as well as Cdt1 and Cdc6 expression are p53-dependent upon p21^{WAF1/Cip1} induction. Corresponding IBs for p53, Cdt1 and Cdc6 in the manipulated HT1080 p21^{WAF1/Cip1}-IPTG-ON cells. Actin serves as a loading control. **e.** FACS analysis of Saos2 p21^{WAF1/Cip1} Tet-ON cells showed an accumulation of cells with >4N DNA content (re-replication) after p21^{WAF1/Cip1} induction that is Cdt1 and Cdc6 dependent.

Figure 3. Sustained p21^{WAF1/Cip1} expression triggers replication stress and accumulation of DNA damage in a Cdt1/Cdc6-dependent manner in S-phase.

a. Prolonged p21^{WAF1/Cip1} expression causes DNA damage. Lysates from Saos2 p21^{WAF1/Cip1} Tet-ON cell, after treatment with 1μg/ml Doxycycline for 96h, were separated by SDS-PAGE and immunoblotted to detect the indicated proteins and phosphorylation levels. To detect DNA double strand breaks, cells were collected and subjected to pulsed field gel electrophoresis (PFGEs). **b.** p21^{WAF1/Cip1} – dependent Cdc6 and Cdt1 overexpression produces DNA damage. Comet assays showed DNA breaks in cells induced for the indicated time points and after siRNA treatments against Cdc6, Cdt1 or both. Red lines in magnifications of insets label comet (moment) tails for length comparison. Bars: 50μm. **c.** p21^{WAF1/Cip1} expression, in cells with non-functional p53, activated the DDR pathway in a Cdc6 and Cdt1-dependent manner. Immunofluorescence and immunoblot analyses of the status of DNA damage response markers (53BP1, γH2AX) in p21^{WAF1/Cip1} induced cells for the corresponding time points and treated with siRNAs against Cdc6, Cdt1 or both. Bars: 5μm, except in magnifications (circled or elipsoid shapes): 7.5 μm. **d.** Sustained p21^{WAF1/Cip1} expression reduced replication fork speed and affected replication fork symmetry. Saos2 p21^{WAF1/Cip1} Tet-ON non-induced cells versus induced ones for 96h and after 20min consecutive labeling pulses of CldU (red) and IdU (green) were subjected to DNA fiber spreading analysis. **e.** Protracted p21^{WAF1/Cip1} expression inflicts DNA damage in S-phase. Flow cytometry analysis (FACS) of p21^{WAF1/Cip1} induced cells for 96h and co-

stained for γ H2X/Propidium Iodide (PI), with or without anti-Cdc6/Cdt1 siRNA targeting, and p21^{WAF1/Cip1} induced cells for the indicated time points and co-stained for EdU/ γ H2X.

Figure 4. Extended p21^{WAF1/Cip1} over-expression mediates replication intermediate lesions accumulation that are processed by MUS81-EME1 and repaired by a Rad52-dependent mechanism.

a-b. Single-strand DNA (ssDNA) production in p21^{WAF1/Cip1} over-expressing cells. p21^{WAF1/Cip1} expression was induced for 96h with 1μg/ml Doxycycline. To detect ssDNA, cells were cultured in medium containing 10 μM BrdU for 24h, fixed and stained with antibodies against BrdU without DNA denaturation (a). Induced cells at 96h were also incubated with anti-RPA antibody (b). **c-d.** p21^{WAF1/Cip1} mediated DNA damage is processed by MUS81 resolvase. Comet assays showed DNA breaks in p21^{WAF1/Cip1} cells induced for 96h and after siRNA treatments against MUS81 (c). Red lines in magnifications of insets label comet (moment) tails for length comparison. Bars: 50μm. IB analysis depicts the efficiency of MUS81 siRNA treatment (c). Flow cytometry analysis (FACS) of p21^{WAF1} induced cells for 96h and co-stained for Edu/γH2X, with or without anti-MUS81 siRNA

targeting (d). **e.** Silencing of the homologous repair recombinase Rad51 resulted in decreased γ -H2AX levels. Lysates from Saos2 p21^{WAF1/Cip1} Tet-ON cell, after treatment with 1 μ g/ml Doxycycline for 96h, were separated by SDS-PAGE and immunoblotted to detect the indicated proteins and phosphorylation levels. **f-g.** Suppression of Rad52 was followed by increased γ -H2AX expression (f,g) and cell death (g). Lysates from Saos2 p21^{WAF1/Cip1} Tet-ON cell, after treatment with 1 μ g/ml Doxycycline for 96h, were separated by SDS-PAGE and immunoblotted to detect the indicated proteins and phosphorylation levels. FACS depicted increased γ -H2AX expression and cell death. **h.** Rad51 promoter is occupied by the E2F4 repressing transcriptional factor upon p21^{WAF1/Cip1} induction. Chromatin immunoprecipitation (ChIP) for E2F4 binding to the Rad51 promoter in non-induced and induced p21^{WAF1/Cip1} cells.

a. Sustained p21^{WAF1/Cip1} expression triggers senescence. p21^{WAF1/Cip1} expression was induced for 96h with 1μg/ml doxycycline. Cells grown on coverslips were stained to assess the senescent phenotype applying the Sudan Black-B protocol (Georgakopoulou et al. 2013) and SA-b-gal (Dimri et al. 1995). Cells were treated with siRNAs against Cdc6, Cdt1, Cdc6/Cdt1 and p73. **b.** IBs depict p73 status upon siRNAs targeting Cdc6 and Cdt1, as well as the efficiency of anti-p73 treatment. **c.** Sustained p21^{WAF1/Cip1} expression reduces p14^{ARF} and p16^{INK4A} protein levels. Bars: 20 μm

Figure 6. Prolonged p21^{WAF1/Cip1} expression, in cells with p53 loss of function, overrides the senescence barrier leading to aggressive behavior.

a. Morphological features observed by reverse-phase contrast microscopy of escaped cells (20 days of p21^{WAF1/Cip1} expression). Bars: 15μm. **b.** BrdU incorporation is restored to almost similar levels to non-induced cells after bypass of senescence. Doxycyclin (1μg/ml) treated cells, up to 20 days, were cultured in medium complemented with 10 μM BrdU at 24h before collection, fixed and IF stained with antibodies against BrdU and p21^{WAF1/Cip1}. **c.** EdU incorporation increases in p21^{WAF1/Cip1} expressing cells after 20 days of continuous

induction. FACS analysis of p21^{WAF1/Cip1} induced cells for 96h and 20 days that were co-stained for EdU and p21^{WAF1/Cip1}. **d.** Appearance of a significant sub-population of Cyclin A and p21^{WAF1/Cip1} positive cells at 20 days of induction. Double IF analysis of induced cells for Cyclin A and p21^{WAF1/Cip1} at indicated time points. Bars: 50µm. **e.** Restoration of Cdk2 activity and reduction of p73 levels in cells “escaping” senescence (see also Supplemental Fig. S14). Cell lysates were separated by SDS-PAGE and immunoblotted to detect the indicated proteins and phosphorylation levels at 4 and 20 days of induction. **f.** Escaped cells depict larger nuclei than non-induced ones. Nuclei stained with DAPI. Histogram depicts average values in the two groups: OFF versus 20 days ON. Bars: 7.5µm. **g.** DNA damage was significantly reduced in escaped cells. Comet assays showed DNA breaks in cells induced for the indicated time points. Red lines in magnifications of insets label comet (moment) tails (TM) for length comparison. Bars: 50µm. **h-i.** Escaped Saos2 p21^{WAF1/Cip1} TetON cells (45 days of p21^{WAF1/Cip1} expression) form more and larger colonies than the cells with non-induced (OFF) p21^{WAF1/Cip1} in soft agar assay ($P < 0.001$) (h). They also exhibit increased invasion capability (i). **j.** Escaped cells (20d) exhibit increased genotoxic drug tolerance. Histogram depicting increased IC₅₀ values by escaped cells upon treatment with Doxorubicin and Cis-platinum.

Figure 7. $p21^{WAF1/Cip1}$ expressing cells that have overridden (escaped) the senescence barrier demonstrate genomic instability.

a. High resolution array-Comparative Genomic Hybridization (array-CGH) analysis (Agilent G3 CGH 1M arrays) between “escaped” $p21^{WAF1/Cip1}$ expressing cells and non-induced ones at the time point of 30 days. Gains and losses in the genome are depicted as blue and red colored regions along the chromosome ideograms, respectively (upper panel). Enlarged inset on left depicts two narrow subchromosomal areas exhibiting alternating regions of gains or losses with retention regions in between (highlighted in blue and red

shadowed rectangles, respectively), that possibly indicate events of chromoanasythesis and chromothrypsis, respectively. Along these areas there are large blocks of chromosomal losses located in the fragile site *FRA3B*. Inset on right denotes a representative result for a large scale subchromosomal loss. **b.** Prolonged p21^{WAF1/Cip1} expression leads to increase frequency of micronuclei in “escaped” cells. Arrows depict micronuclei. Histogram shows quantification of micronuclei in p21^{WAF1/Cip1} induced cells. **c.** Circos diagram depicting novel (N: 103) chromosomal rearrangements in “escaped” p21^{WAF1/Cip1} expressing cells revealed by deep sequencing (human chromosomes are located at the perimeter). **d.** Representative results showing the high correlation between the aCGH, deep sequencing (Next Generation Sequencing: NGS) and cytogenetic analyses results in “escaped” p21^{WAF1/Cip1} cells. **e.** Proposed model depicting prolonged p53-independent p21^{WAF1/Cip1} oncogenic action (see text for details).