Published in final edited form as:

Mar Pollut Bull. 2016 June 15; 107(1): 224–232. doi:10.1016/j.marpolbul.2016.03.068.

Proteomic analysis of bleached and unbleached *Acropora palmata*, a threatened coral species of the Caribbean

Martha Ricaurte¹, Nikolaos V. Schizas^{*#1}, Pawel Ciborowski², and Nawal M. Boukli^{*#3}

¹Department of Marine Sciences, University of Puerto Rico, Mayagüez, Call Box 9000, Mayagüez, PR 00681, USA

²Mass Spectrometry and Proteomics Core Facility, Durham Research Center, Room 3048, 985800. University of Nebraska Medical Center (UNMC), Omaha, NE 68198-5800, USA

³Biomedical Proteomics Facility, Microbiology and Immunology Department, Universidad Central del Caribe, Bayamón, PR 00960, USA

These authors contributed equally to this work.

Abstract

There has been an increase in the scale and frequency of coral bleaching around the world due mainly to changes in sea temperature. This may occur at large scales, often resulting in significant decline in coral coverage. In order to understand the molecular and cellular basis of the ever-increasing incidence of coral bleaching, we have undertaken a comparative proteomic approach with the endangered Caribbean coral *Acropora palmata*. Using two-dimensional gel electrophoresis (2D-GE) and tandem mass spectrometry (MS/MS), we identified 285 and 321 expressed proteins in bleached and unbleached *A. palmata* colonies, respectively, in southwestern Puerto Rico. Overall the expression level of 38 proteins was significantly different between bleached and unbleached corals. A wide range of proteins was detected and categorized, including transcription factors involved mainly in heat stress/UV responses, immunity, apoptosis, biomineralization, the cytoskeleton, and endo-exophagocytosis. The results suggest that for bleached *A. palmata*, there was an induced differential protein expression response compared with those colonies that did not bleach under the same environmental conditions.

Graphical abstract

Proposed model highlighting the bleaching response pathways highlighting key protein signatures with differences in abundance in bleached *Acropora palmata* compared with unbleached *A. palmata*.

Corresponding Author nawal.boukli@uccaribe.edu (NMB).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Introduction

Bleaching, the partial or complete loss of photosynthetic zooxanthellae from coral tissues, occurs mainly when corals are stressed after exposure to elevated sea water temperatures (Aronson and Precht, 2006; Hoegh-Guldberg, 1999; Hoegh-Guldberg, et al., 2007; Hughes, 2003). Increments of one or two degrees Celsius are sufficient to initiate expulsion of zooxanthellae by the host (DeSalvo et al., 2008; Lesser, 2007; Wilkinson, 2008). Depending on the degree of temperature elevation and the time that it remains above normal, coral bleaching can be a reversible phenomenon without major consequences for many coral communities (Baker et al., 2008; Graham et al., 2015), or can result in mass mortality with substantial ecological consequences (Goreau et al., 2000; Pratchett et al., 2008).

Based on sporadic observations, geographically restricted coral bleaching events have occurred since the 1950s. An increase in the severity, frequency, and geographical extent of these events, began approximately in 1982 (Lesser, 2007). All tropical regions are affected by bleaching but one of the most affected is the Caribbean that has suffered several recent and well-documented mass-bleaching events in 1987, 1998, 2003, 2005, and 2010 (Alemu and Clement, 2014; Aronson and Precht, 2006; Ananko et al., 2005; Eakin et al., 2010; Hoegh-Guldberg, 1999; Hughes, 2003; Weil, 2004; Weil et al., 2006). During 2015–2016, a third global bleaching event threatened many reef areas in the tropical Pacific, Indian, and Atlantic Oceans (Witze, 2015). Environmental models predict that bleaching events will increase in frequency and intensity (Hoegh-Guldberg, 1999) due to rising tropical sea surface temperatures. As a consequence of consecutive bleaching events in combination with disease outbreaks (Bruno, 2007; Eakin et al., 2010; Weil, 2004) the dominant reef-building corals of the Caribbean (*Acropora palmata*, *A. cervicornis*, *Orbicella annularis*, and *O. faveolata*), have been disappearing rapidly. The rapid decline of *A. palmata*, perhaps the most iconic scleractinian coral of the Caribbean, has been so alarming that the species was listed as “threatened” under the US Endangered Species Act in 2006 (Miller et al., 2009; Precht et al., 2004). In December 2012, the National Marine Fisheries Service (NMFS) proposed reclassifying *Acropora palmata* as an endangered species (77 FR 73219) but decided against the reclassification in August, 2014.

Proteomics is one of the modern tools to evaluate the effects of environmental stress at the posttranslational levels of proteins in marine organisms (Tomanek, 2011). Bleaching is a response to thermal stress (Brown et al., 1996; Douglas, 2003; Gates et al., 1992; Hoegh-Guldberg, 1999; Hughes, 2003; Pratchett et al., 2013) and has been studied by monitoring changes in gene expression (DeSalvo et al., 2010, 2008; Pinzon et al., 2015; Smith-Keune and Dove, 2008; Voolstra et al., 2009; Yuyama et al., 2012) and western blotting of specific proteins.

To gain further insight into the molecular and cellular basis of coral bleaching, we performed a comparative proteomic analysis between unbleached and bleached colonies in the endangered Caribbean coral *A. palmata* during the mass-bleaching event of August/September 2010. We validated the protein expression of *A. palmata* tissue collected from southwestern Puerto Rico using western blots for a selected set of differentially expressed proteins.

Materials and Methods

Sampling Location

After the massive bleaching in 2005, in which over 80% of Caribbean corals became bleached and over 40% died at many locations (Eakin et al., 2012), another severe event followed during 2010. The latter event persisted for more than 3 months, with the highest water temperatures (Fig. 1) registered with the moored autonomous pCO₂ system (MAPCO₂) of NOAA Pacific Marine Environmental Laboratory (<http://pmel.noaa.gov/co2/story/La+Parguera>). During 2010, tissues from *Acropora palmata* colonies were collected for protein extraction from the fore reef of Collado Cay (17°57'9.06"N, 67° 4'43.35"W), La Parguera, located on the southwest coast of Puerto Rico. The ecology of coral reefs in La Parguera, a marine reserve, has been the subject of numerous studies (Ballantine et al., 2008; Morelock et al., 2001). All coral tissues were collected under permit No. O-VS-PVS15-SJ-00668–22042014 issued to the first author by the Puerto Rico Department of Natural and Environmental Resources.

Protein collection and extraction

Small coral fragments were removed from three unbleached and three bleached *A. palmata* colonies with a hammer and chisel. The tissues were collected at a depth of 1 to 2 m and at least 3 m apart to reduce the possibility of sampling genetic clones. The samples were immediately placed in plastic bags underwater and were transported back to the laboratory in a cooler with ice. Immediately upon return, the corals were broken into 2-cm² pieces and placed in a mortar for grinding in 1 mL⁻¹ of Rehydration Buffer (9.5M Urea, 2% CHAPS, 1% DTT). The homogenized tissue was placed in a 1.5 mL⁻¹ tube and lysed by sonication on ice, three times every 30 seconds, followed by slow shaking in a Bio-Rad laboratory shaker, at room temperature for 1 hour. Samples were centrifuged in a refrigerated centrifuge (Eppendorf 5810R) at 12,000 rpm for 30 minutes at 4°C, and the supernatant was placed in a new sterile 1.5 mL⁻¹ Eppendorf tube. The protein concentration was estimated by the Bradford method (Bradford, 1976).

First-dimension step, isoelectric focusing (IEF)

Two hundred micrograms of protein from unbleached and bleached *A. palmata* were mixed in rehydration buffer (Bio-Rad) overnight on immobilized pH gradient (IPG) strips (pH 310), which were then subjected to first-dimension Isoelectric Focusing (IEF) in a Bio-Rad protein IEF cell (50 μ A/strip) for a total of 20,000 Vh, as previously described in (Boukli et al., 2011, 2012).

High-resolution two-dimensional gel electrophoresis (2D-GE)

After completion of the run, strips were reduced and alkylated for 2×15 min each in equilibration buffer I (6 mol L⁻¹ urea, 2% SDS, 0.05 mol L⁻¹ Tris-HCl, pH 8.8, 50% glycerol, and 2% [w/v] dithiothreitol [DTT]) followed by equilibration buffer II (6 mol L⁻¹ urea, 2% SDS 0.05 mol L⁻¹, Tris-HCl, pH 8.8, 50% glycerol and 2.5% [w/v] Iodoacetamide). The second-dimension separation was performed with a 4–20% gradient SDS-polyacrylamide gel, to increase the range of molecular weights and sharpening bands (15 mA/gel for 15 min, then 30 mA/gel). Gels with unbleached and bleached *A. palmata* were stained with BioSafe Coomassie blue G-250 dye (Bio-Rad).

Image analyses

The images of the stained gels were taken, in a Bio-Rad Versa Doc Model 1000 imaging system at 200 dpi resolution. All images were analyzed for spot detection and matching with Bio-Rad PD-Quest™ software, version 7.4.0. The software calculated individual spot “volumes” by density/area integration; then, to eliminate gel-to-gel variation, individual spot volumes for each gel were normalized relative to the total spot volume of that gel. Normalized spot volume data from each experimental set were exported to Microsoft Excel, where differences in expression of spots between the unbleached and bleached coral groups were analyzed using analysis of variance (ANOVA), with $P < 0.05$ as the criterion for statistical significance. Protein spots determined to be significantly different in abundance 2 fold between the two cell lysates bleached and unbleached *acropora palmata* were calculated in corresponding fold number by PD Quest software. These significantly differentially expressed 2D spots were selected and marked for excision. All experiments were performed in biological and technical triplicates.

Protein in-gel digestion and data analysis

Spots of interest were manually excised from Coomassie blue-stained gels using a sterile scalpel and kept in sterile microcentrifuge tubes for destaining with 40% methanol/10% acetic acid, reduced with 10 mmol L⁻¹ DTT in 50 mmol L⁻¹ ammonium bicarbonate (AB), then alkylated with 55 mmol L⁻¹ iodoacetamide in 50 mmol L⁻¹ AB. The resulting gel fragments were rinsed with 50 mmol L⁻¹ AB and 50% acetonitrile (ACN) and dried under a stream of nitrogen. Samples were digested with 5% trypsin proteomics grade (w/w) (Sigma-Aldrich) overnight at 37°C and combined with an equal volume of saturated cyano-4-hydroxycinnamic acid in 50% ACN/0.1% trifluoroacetic acid. Half the mixture from each spot was applied to a MALDI target plate. MALDI TOF-MS/MS was performed at the University of Texas, Medical Branch, Proteomics Facility (Galveston, Texas). Following MALDI MS analysis, Applied Biosystems GPS Explorer™ software (version 3.6) was used

in conjunction with MASCOT software to search the NCBIInr databases to identify proteins differing in abundance between unbleached and bleached *A. palmata*. Protein match probabilities were determined by using expectation values and/or the MASCOT search engine (Matrix Science). The proteins were identified by searching the raw tandem MS data in the International Protein Index (IPI) database with a Sequest search algorithm and by using the Proteome Discoverer 1.1 (Thermo Electron, San Jose, CA). The database search parameters were the following: database, UniProt KB; mass-type, monoisotopic precursor and fragment; enzyme, trypsin; threshold, 100; peptide tolerance, 1.5 Da; and fragment ion tolerance, 1.0. The modifications considered in these analyses were the following: Moxidation, C-carbamidomethylation, and N/Q-deamidation. The search results were filtered with cross-correlation (XCorr) via charge states (+1, 1.5; +2, 2.0; +3, 2.5; +4, 3.0), delta correlation ($C_n > 0.1$), molecular weight (m/z 500–2000 Da), and peptide mass accuracy (± 1.5 Da) parameters. The criteria involved in the selection of proteins in each gel spot were the comparison of experimental isoelectric point (pI), molecular weight, and sequence coverage of proteins with that of the theoretical pI and molecular weight of the proteins detected on 2D-GE gels. However, proteins identified with a single peptide were also selected in view of their high XCorr numbers and high-quality MS/MS spectra with b and y ion series. All the tandem MS/MS data were further verified manually to identify proteins with high confidence.

2D-PAGE western blot analyses

Protein lysates (200 μ g) from unbleached and bleached *A. palmata* were resolved by 2DPAGE and subsequently transferred to nitrocellulose membranes for western blot analysis by electrophoresis at 100 V for 1 h in a Criterion™ blotter (Bio-Rad) according to the manufacturer's instructions. Each gel was electroblotted for 3.5 hours to supported nitrocellulose. Membranes were blocked and incubated overnight in 5% non-fat dried milk in Tris-buffered saline with each of the five antibodies (1:1000 dilution, Sigma-Aldrich). The antibodies selected correspond to key proteins identified by peptide mass fingerprinting (myosin, actin, CD63 antigen, calmodulin, and collagen, Fig. 4), which were selected according to their functions. After thorough washing, the membranes were incubated for another 60 minutes with their corresponding secondary antibodies (1:10,000 dilution, Sigma-Aldrich) in 5% non-fat dried milk and washed three additional times with TBS-T. Immunoreactive 2D spots were visualized by chemiluminescence using a SuperSignal® West Femto Substrate (Pierce, Thermo-Scientific) and the blot images were captured on Gel Doc™ XR+ and ChemiDoc™ XRS+ gel documentation systems using Image Lab™ software version 4.2 (Bio-Rad). Following western blot analyses, immunogenic proteins were matched to the corresponding proteins in Coomassie blue-stained gels.

Results

Bleaching-induced changes in *A. palmata* proteome

Proteomic analysis revealed that the majority of differences in protein expression were located at molecular weights in the range of 10–55 kDa and pH 3–10 (Fig. 2). Because it was not feasible to sequence all the resolved proteins, the selection criteria for sequencing was based on gel image quantitative analysis of the 2D-GE (Fig. 2), with the goal of

obtaining a broad sampling of the proteins downregulated and upregulated during bleaching in *A. palmata*. Differentially expressed protein spots were analyzed using PD-Quest™ software, selected for subsequent mass spectrometry analysis, and excised from the gel for protein identification by tandem MS (MS/MS). Thirty-eight proteins were significantly differentially expressed, based on the selection criterion of a 2 fold difference at $p < 0.05$ and were selected from two categories: (i) bleaching-induced up-regulation (13 spots) and (ii) bleaching-induced down-regulation (25 spots; Fig. 2, Table 1). To confirm protein identifications, BLASTP analysis was performed and the amino acid sequences were matched to the corresponding available *A. palmata* sequences. Proteins analyzed by peptide mass fingerprinting were grouped according to 12 categories and their functions were linked in the model proposed in Fig. 3. Most of the downregulated proteins (66%) in unbleached colonies were involved in biomineralization, immunity, transcription and endo-exophagocytosis. About 5% of the proteins were attributed to apoptosis after upregulation with bleaching. Other downregulated proteins were related to immune and metabolic dysfunctions, such as ATP synthase subunit 8 ($\downarrow$ 2.29 fold), toll-like receptor 2 protein ($\downarrow$ 2.25 fold), integrin beta ($\downarrow$ 2.34 fold) and galaxin proteins ($\downarrow$ 2.18 fold). On the other hand, upregulated proteins (34.21%) were related to stress, UV response, apoptosis, the cytoskeleton, and cell cycle functions (Table 1). Actin increased 2.6 fold and calmodulin was uniquely expressed in bleached *A. palmata*. Both proteins play an important role in coordinating cytoskeletal and cell cycle functions, as highlighted in the model in Fig. 3. Other proteins were related to transcription factors, oxidative phosphorylation and cytoskeletal functions, with downregulation of proteins such as PaxC, ectonucleoside triphosphate diphosphohydrolase 1, gelsolin, and myosin (Table 1). All exhibited higher expression levels (4.35-, 2.33-, 2.3-, and 2.26-fold greater, respectively) in unbleached colonies than bleached colonies. It should be emphasized that all *A. palmata* colonies studied in this project affected by the 2010 bleaching event recovered and appeared to remain healthy during the next three months (October- December of 2010). Interestingly, photosystem I P700 chlorophyll-a apoprotein A1 was found to have a dual function (stress response and endo-exophagocytosis), indicating that this protein is a good candidate marker to signal coral stress when the temperature changes.

Statistical analysis

The standard deviation of a 2D spot was estimated by the PD-Quest™ software at a 2.0-fold change. Each protein was analyzed in technical and biological triplicates. Data were tested for normality with a Shapiro-Wilk test at a significance level of $p > 0.05$, and the criteria for normality were met in all the proteins expressed for unbleached and bleached *A. palmata*. The homogeneity of the variance among samples was evaluated with Levene's test. The resulting p-value for Levene's test was < 0.05 , indicating that there was a difference between the variances in unbleached and bleached *A. palmata*. The normality and equal variances within time points and the significance of differences between time points was assessed using two-way ANOVA and pairwise post-hoc testing via the Shapiro-Wilks test, with a significance level of $p > 0.05$. This analysis revealed that the differences in protein expression level of the 38 proteins differentially expressed between bleached and unbleached corals were significant.

2D-GE western blot of five selected differentially expressed proteins

The 2D-GE western blot analysis confirmed the differential expression of five key proteins: myosin, collagen, CD63, calmodulin, and actin (Fig. 4). These protein functions were related to the cytoskeleton, the cell cycle, and endo-exophagocytosis cellular responses and contained proteins that were uniquely expressed in bleached *A. palmata*, such as calmodulin and actin (Fig. 4), suggesting that the physiological functions of the bleached corals were affected during stress (Fig. 3).

Discussion

Proteomic studies are increasingly used to study the effects of marine environmental stressors (e.g. Anthony et al., 2009; DeSalvo et al., 2008; Tomanek, 2010). Previous studies employing gene expression analysis were undertaken for *Porites astreoides* in different thermally stressed environments (Hayward et al., 2011) and for *Orbicella faveolata*, in which the transcriptome before and after bleaching was compared (Pinzon et al., 2015). Although, there are no published studies on the whole proteome of the endangered species *A. palmata* that reveal the protein expression profile before, during, and after bleaching in situ. We are bridging this knowledge gap by identifying the proteins differentially expressed in *A. palmata* in response to a temperature stress in the fore reef of Collado Cay, La Parguera, Puerto Rico. This heat stress is the result of an increase in temperatures from 27.5 °C to 30.8 °C in 2010, with a sustained temperature of 29.5 °C, during the months of July, August, September, and October of 2010 (Fig. 1). These high temperatures lasted long enough to cause bleaching (Glynn, 1993). A large scale mass-bleaching event occurred in 2005. The second and most recent mass-bleaching event of the first decade of the new millennium occurred in 2010 throughout the Caribbean (DeSalvo et al., 2008; Doney et al., 2009; Eakin et al., 2010; Fitt et al., 2009; Miller et al., 2009; Palmer et al., 2011; Prada et al., 2010).

The effect of heat stress on *A. palmata* induced a higher number of differentially expressed proteins (25 spots) in unbleached colonies than in bleached colonies (13 spots) as revealed by the proteomic results (Table 1). More than 66% of the differentially expressed proteins were downregulated, such as the cytoskeletal proteins gelsolin and myosin under the skeletal surface organic-matrix, suggesting that the cytoskeleton underwent rearrangement of its adhesion properties during bleaching, as observed by others (DeSalvo et al., 2008; Douglas, 2003; Jones, 2008). We believe that downregulation of the expressed proteins (56.78%) in bleached *A. palmata* compromised the innate coral immune system under heat stress. Proteins such as apextrin (↓ 2.54 fold), ATP synthase subunit 8 (↓ 2.29 fold), ectonucleoside triphosphate diphosphohydrolase 1 (↓ 2.33 fold), and toll-like receptor (↓ 2.25 fold) are key proteins in the immunity cascade that occur in the ancestral cnidarian (Miller et al, 2007). The heat stress induced upregulation of protein functions affecting UV responses, apoptosis, the cytoskeleton, the cell cycle, and stress responses. The upregulated proteins correspond to heat shock transcription factor proteins (↑ 2.4 fold), green fluorescent protein (↑ 2.32 fold), NADH ubiquinone oxidoreductase chain 5 (↑ 2.43 fold), histone H2B (↑ 2.4 fold), activin 1 (↑ 2.2 fold), and photosystem I P700 chlorophyll-a apoprotein A1 (↑ 2.47 fold). This suggests, that these proteins are upregulated with a high metabolic cost for *A. palmata* during bleaching. Based on these findings, the comparative proteomic analysis between

unbleached and bleached *A. palmata* suggests that heat stress is activated via a metabolic route in response to high temperature and UV radiation (Fig. 3). Due to heat stress-induced bleaching, a defense mechanism is activated, leading to differential protein expression. Among key upregulated proteins, there are several reactive oxygen species (ROS) markers, such as NADH-ubiquinone oxidoreductase, activin 1 and cytochrome oxidase. We suspect that the ROS response overwhelmed the antioxidant defenses, leading to various manifestations of oxidative stress highlighted by upregulation of heat shock proteins, activin 1, cytochrome oxidase, cytochrome b, cytochrome P450, and thioredoxin reductase, with respective fold changes of 2.40, 2.20, 2.43, 2.25, and 2.67. Entry of the coral into oxidative stress causes metabolic dysfunctions (Jones et al., 1998), such as lipid peroxidation, protein oxidation, damage to the coral cell cycle, inhibition of enzymes, damage to nucleic acids, and activation of the apoptosis pathway (DeSalvo et al., 2008; Kültz, 2005; Voolstra et al., 2009), ultimately leading to coral death. Some of the upregulated proteins revealed by peptide mass fingerprinting (Table 1) are proapoptotic markers (Danial et al., 2010; DeSalvo et al., 2008), such as apextrin and ectonucleoside triphosphate diphosphohydrolase 1 (with 2.54- and 2.33- fold changes, respectively). Another key proapoptotic protein identified in this study is caspase 8. This protein released by the mitochondria appeared to be uniquely expressed in *A. palmata* during bleaching (see Fig. 2 and Table 1), and it is known to be one of the key upregulated proapoptotic markers, indicating that corals are under stress (DeSalvo et al., 2008; Li et al., 1998; Richier et al., 2008). Green fluorescent protein (GFP) and heat shock proteins (HSPs), known to respond during photoacclimation (Chalfie et al., 2007) and during short thermal stress (Fitt et al., 2009), respectively, were two differentially overexpressed proteins (Table 1) that are potential candidate markers for assessing the physiological state of the coral during bleaching due to thermal stress. Their respective upregulation is believed to trigger an immune response through the photoinhibition of *A. palmata* (Fig. 3). Moreover, HSP proteins bind to denatured proteins stabilizing new proteins to ensure that cells prevent/limit photodamage and therefore might prevent photoinhibition-dependent *A. palmata* bleaching.

In addition, cytoskeletal proteins important for the cell cycle, such as gelsolin, actin, and myosin, were differentially expressed in this study. These proteins are known to be activated during thermal stress at the intracellular and extracellular level by activating the immune response (Palmer et al., 2011). Among other proteins that were uniquely expressed are cyclin and calmodulin. Both proteins are known to play a key role in endo-exophagocytosis processes (Tomanek, 2010). Also, the significant overexpression of calmodulin in *A. palmata* is believed to be instrumental to biomineralization and other calcium-dependent processes through its interaction with several proteins during the development of corals (DeSalvo et al., 2008; ReyesBermudez et al., 2012).

In the proposed metabolic model (Fig. 3), we speculate that during elevated temperature, the intracellular calcium concentration activates caspase 8 in the proteolytic processes, affecting mitochondria internally. This leads to an overload of Ca^{+2} ions, increasing oxidative stress. Other crucial proteins that are released from the mitochondria are the cytochrome oxidase subunits, cytochrome b, and cytochrome P450, all components of the mitochondrial electron transport chain and known to be pro-apoptotic markers. The three forms of cytochromes

were found to be differentially expressed, as they trigger caspases, which in turn induce bleaching and finally lead to apoptosis in *A. palmata* tissue.

Our findings predict that the extracellular matrix of bleached *A. palmata* is changing physiologically when calcium carbonate is forming, reducing the biomineralization capacity (DeSalvo et al., 2008), eventually leading to either apoptosis (Fig. 3) or to recovery if the stressor temperature is removed. During the process of biomineralization, important proteins involved in calcification (Wirshing and Baker, 2014), such as galaxin and minicollagen were highly and uniquely expressed in bleached corals.

Conclusions

The comparative proteomic analysis undertaken in this study confirmed that a differential protein expression profile occurs in bleached compared with unbleached *A. palmata*. Proteins related to thermal heat stress were highly overexpressed in the bleached colonies as well as protein signatures related to stress, UV response, apoptosis and cell cycle functions. Based on our criteria of a two-fold or greater change in expression, the results indicated that temperature significantly regulated key components of the stress induced apoptotic response. We speculate that major metabolic functions are altered during the processes of ROS production, amino acid synthesis, and endo-exophagocytosis (Fig. 3). The latter two processes are involved in transport function during skeletal coral growth, indicating that bleaching seems to affect *A. palmata* growth. We feel it is noteworthy that we found eight uniquely expressed proteins including minicollagen, and proteins related to growth, detoxification, endo-exophagocytosis, and catabolic processes under bleaching conditions.

This study allowed the identification of key changes in protein expression due to bleaching of *A. palmata*. Among the potential marker candidates for bleaching stress are green fluorescent proteins (GFP), heat shock proteins, and caspase 8. This latter protein, which is uniquely present and expressed only in bleached *A. palmata*, is a pro-apoptotic marker that may signal the beginning of coral death. Other uniquely expressed proteins during coral colony bleaching include cyclin and calmodulin. These two proteins, which play a key role in coral defense, represent potentially useful markers to disentangle the molecular mechanisms underlying bleaching.

We propose a network model of protein-protein interactions to identify and prioritize candidate biomarkers when studying bleaching as a defense mechanism triggered in *A. palmata* during changes in sea temperatures. It is worth emphasizing that this is the first proteomic study conducted on the endangered coral species *A. palmata* during a bleaching event in the Caribbean.

Finally, the comparative proteomic analysis used in this present project may open new avenues in the study of coral gene expression and the interactions with the environment. Since climate change can potentially affect all reef organisms, comparative proteomic approaches can also be applied to other marine invertebrates under stress.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

The authors wish to thank Nery Rivera, Gloriene González, Emmanuel García, and Kathy Trujillo for their technical assistance and Cristina López for her assistance in statistical procedures used in the manuscript. We thank David Ballantine for improving the language of the manuscript. The proteomics studies were supported by the National Institutes of Health Research Centers in Minority Institutions (RCMI) Program and Biomedical Proteomics Facility Grant G12MD007583 awarded to NMB at Universidad Central del Caribe in Bayamón and the Puerto Rico and Seed Money awarded by the UPRM College of Arts & Sciences, and SeaGrant Puerto Rico Grant FRS-535265 awarded to NVS. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute on Minority Health and Health Disparities or the National Institutes of Health.

References

- Alemu JB, Clement Y, 2014 Mass Coral Bleaching in 2010 in the Southern Caribbean. PLoS ONE 9, e83829. doi:10.1371/journal.pone.0083829. [PubMed: 24400078]
- Ananko EA, Podkolodny NL, Stepanenko IL, Podkolodnaya OA, Rasskazov DA, Miginsky DS, Likhoshvai VA, Ratushny AV, Podkolodnaya NN, Kolchanov NA 2005 GeneNet in 2005. Nucleic Acids Res 33, D425–D427. doi:10.1093/nar/gki077. [PubMed: 15608230]
- Anthony KRN, Hoogenboom MO, Maynard JA, Grottoli AG, Middlebrook R 2009 Energetics approach to predicting mortality risk from environmental stress: a case study of coral bleaching. Funct Ecol 23, 539–550. doi:10.1111/j.1365-2435.2008.01531.x.
- Aronson RB, Precht WF 2006 Conservation, precaution, and Caribbean reefs. Coral Reefs 25, 441–450. doi:10.1007/s00338-006-0122-9.
- Ballantine D, Appeldoorn R, Yoshioka P, Weil E, Armstrong R, Garcia J, Otero E, Pagan F, Sherman C, Hernandez-Delgado E, Bruckner A, Lilyestrom C 2008 Biology and Ecology of Puerto Rican Coral Reefs, in: Coral Reefs of the USA Coral Reefs of the World pp. 375–406. doi: 10.1007/978-1-4020-6847-8_9.
- Baker AC, Glynn PW, Riegl B 2008 Climate change and coral reef bleaching: An ecological assessment of long-term impacts, recovery trends and future outlook. Estuar Coast Shelf Sci 80, 435–471.
- Boukli N, Shetty V, Cubano L, Ricaurte M, Coelho-Dos-Reis J, Nickens Z, Shah P, Talal AH, Philip R, Jain P 2012 Unique and differential protein signatures within the mononuclear cells of HIV-1 and HCV mono-infected and co-infected patients. Clin Proteomics 9, 11. doi:10.1186/1559-0275-9-11. [PubMed: 22958358]
- Boukli NM, Delgado B, Ricaurte M, Espino AM 2011 *Fasciola hepatica* and *Schistosoma mansoni*: Identification of Common Proteins by Comparative Proteomic Analysis. J Parasitol 97, 852–861. doi:10.1645/GE-2495.1. [PubMed: 21506812]
- Bradford MM 1976 A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Anal Biochem 72, 248–254. doi: 10.1016/0003-2697(76)90527-3. [PubMed: 942051]
- Brandt ME, McManus JW 2009 Disease incidence is related to bleaching extent in reefbuilding corals. Ecology 90, 2859–2867. [PubMed: 19886494]
- Brown BE, Dunne RP, Chansang H 1996 Coral bleaching relative to elevated seawater temperature in the Andaman Sea (Indian Ocean) over the last 50 years. Coral Reefs 15, 151–152. doi:10.1007/s003380050036.
- Bruno JF, Selig ER, Casey KS, Page CA, Willis BL, Harvell CD, Sweatman H, Melendy AM 2007 Thermal stress and coral cover as drivers of coral disease outbreaks. PLoS Biology 5 (6), 1220–1227.
- Chalfie M, Tu Y, Euskirchen G, Ward W, Prasher D 1994 Green fluorescent protein as a marker for gene expression. Science 263, 802–805. doi:10.1126/science.830329. [PubMed: 8303295]

- Danial NN, Gimenez-Cassina A, Tondera D 2010 Homeostatic Functions of BCL-2 Proteins beyond Apoptosis, in: *Advances in Experimental Medicine and Biology* pp. 1–32. doi: 10.1007/978-1-4419-6706-0_1.
- DeSalvo MK, Voolstra CR, Sunagawa S, Schwarz JA, Stillman JH, Coffroth MA, Szmant AM, Medina M 2008 Differential gene expression during thermal stress and bleaching in the Caribbean coral *Montastraea faveolata*. *Mol Ecol* 17, 3952–71. doi:10.1111/j.1365-294X.2008.03879.x. [PubMed: 18662230]
- DeSalvo MK, Sunagawa S, Voolstra CR, Medina M 2010 Transcriptomic responses to heat stress and bleaching in the elkhorn coral *Acropora palmata*. *Mar Ecol Progr Ser* 402, 97–113.
- Doneyk SC, Fabry VJ, Feely RA, Kleypas JA 2009 Ocean acidification: the other CO₂ problem. *Ann Rev Mar Sci* 1, 169–92. doi:10.1146/annurev.marine.010908.163834.
- Douglas AE 2003 Coral bleaching—how and why? *Mar Pollut Bull* 46, 385–92. doi:10.1016/S0025-326X(03)00037-7. [PubMed: 12705909]
- Eakin CM, Morgan JA, Heron SF, Smith TB, Liu G, Alvarez-Filip L, Baca B., Bartels E., Bastidas C, Bouchon C, Brandt M, Bruckner AW, Bunkley-Williams L, Cameron A, Causey BD, Chiappone M, Christensen TR, Crabbe MJ, Day O, de la Guardia E, Díaz-Pulido G, DiResta D, Gil-Agudelo DL, Gilliam DS, Ginsburg RN, Gore S, Guzmán HM, Hendee JC, Hernández-Delgado EA, Husain E, Jeffrey CF, Jones RJ, Jordán-Dahlgren E, Kaufman LS, Kline DI, Kramer PA, Lang JC, Lirman D, Mallela J, Manfrino C, Maréchal JP, Marks K, Mihaly J, Miller WJ, Mueller EM, Muller EM, Orozco Toro CA, Oxenford HA, Ponce-Taylor D, Quinn N, Ritchie KB, Rodríguez S, Ramírez AR, Romano S, Samhouri JF, Sánchez JA, Schmahl GP, Shank BV, Skirving WJ, Steiner SC, Villamizar E, Walsh SM, Walter C, Weil E, Williams EH, Roberson KW, Yusuf Y 2010 Caribbean Corals in Crisis: Record Thermal Stress, Bleaching, and Mortality in 2005. *PLoS ONE* 5(11), e13969. doi:10.1371/journal.pone.0013969. [PubMed: 21125021]
- Fitt WK, Gates RD, Hoegh-Guldberg O, Bythell JC, Jatkar A, Grottolie AG, Gomez M, Fisherc P, Lajuenessef TC, Pantosc O, Iglesias-Prieto R, Franklindh DJ, Rodriguesi LJ, Torregianik JM, van Woesikj R, Lesser MP 2009 Response of two species of Indo-Pacific corals, *Porites cylindrica* and *Stylophora pistillata*, to short-term thermal stress: The host does matter in determining the tolerance of corals to bleaching. *J Exp Mar Bio Ecol* 373, 102–110. doi:10.1016/j.jembe.2009.03.011.
- Gates RD, Baghdasarian G, Muscatine L 1992 Temperature Stress Causes Host Cell Detachment in Symbiotic Cnidarians: Implications for Coral Bleaching. *Biol Bull* 182, 324332. doi: 10.2307/1542252.
- Glynn PW 1991 Coral reef bleaching in the 1980s and possible connections with global warming. *Trends in Ecology and Evolution* 6, 175–179. [PubMed: 21232450]
- Glynn PW 1993 Coral reef bleaching: ecological perspectives. *Coral Reefs* 12, 1–17. doi:10.1007/BF00303779
- Goreau T, McClanahan T, Hayes R, Strong A 2000 Conservation of coral reefs after the 1998 global bleaching event. *Conserv Biol* 14: 5–15.
- Graham NAJ, Jennings S, MacNeil MA, Mouillot D, Wilson SK 2015 Predicting climate-driven regime shifts versus rebound potential in coral reefs. *Nature* 518, 94–97. [PubMed: 25607371]
- Hayward DC, Hetherington S, Behm CA, Grasso LC, Forêt S, Miller DJ, Ball EE 2011 Differential gene expression at coral settlement and metamorphosis—a subtractive hybridization study. *PLoS One* 6, e26411. doi:10.1371/journal.pone.0026411. [PubMed: 22065994]
- Hoegh-Guldberg O 1999 Climate change, coral bleaching and the future of the world's coral reefs. *Mar Freshw Res* 50, 839. doi:10.1071/MF99078.
- Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell CD, Sale PF, Edwards AJ, Caldeira K, Knowlton N, Eakin CM, IglesiasPrieto R, Muthiga N, Bradbury RH, Dubi A, Hatzios ME, 2007 Coral reefs under rapid climate change and ocean acidification. *Science* 318, 1737–1742. [PubMed: 18079392]
- Hughes TP, Baird AH, Bellwood DR, Card M, Connolly SR, Folke C, Grosberg R, Hoegh-Guldberg O, Jackson JBC, Kleypas J, Lough JM, Marshall P, Nyström M, Palumbi SR, Pandolfi JM, Rosen B, Roughgarden J 2003 Climate change, human impacts, and the resilience of coral reefs. *Science* 301, 929–933. doi:10.1126/science.1085046. [PubMed: 12920289]

- Jones RJ, Hoegh-Guldberg O, Larkum AWD, Schreiber U 1998 Temperature-induced bleaching of corals begins with impairment of the CO₂ fixation mechanism in zooxanthellae. *Plant, Cell Environ* 21, 1219–1230. doi:10.1046/j.1365-3040.1998.00345.x.
- Kültz D 2005 Molecular and evolutionary basis of the cellular stress response. *Annu Rev Physiol* 67, 225–57. doi:10.1146/annurev.physiol.67.040403.103635. [PubMed: 15709958]
- Lesser MP, 2007 Coral reef bleaching and global climate change: can corals survive the next century? *Proc Natl Acad Sci U S A* 104, 5259–60. doi:10.1073/pnas.0700910104. [PubMed: 17374718]
- Li H, Zhu H, Xu C, Yuan J 1998 Cleavage of BID by Caspase 8 Mediates the Mitochondrial Damage in the Fas Pathway of Apoptosis. *Cell* 94, 491–501. doi:10.1016/S0092-8674(00)81590-1. [PubMed: 9727492]
- Miller DJ, Hemmrich G, Ball EE, Hayward DC, Khalturin K, Funayama N, Agata K, Bosch TCG 2007 The innate immune repertoire in Cnidaria--ancestral complexity and stochastic gene loss. *Genome Biol* 8, R59. doi:10.1186/gb-2007-8-4-r59. [PubMed: 17437634]
- Miller J, Muller E, Rogers C, Waara R, Atkinson A, Whelan KRT, Patterson M, Witcher B 2009 Coral disease following massive bleaching in 2005 causes 60% decline in coral cover on reefs in the US Virgin Islands. *Coral Reefs* 28, 925–937. doi:10.1007/s00338009-0531-7.
- Morelock J, Ramírez WR, Bruckner AW, Carlo M 2001 Status of coral reefs southwest Puerto Rico. *Caribb J Sci Spec Publ*. No. 4.
- Palmer CV, McGinty ES, Cummings DJ, Smith SM, Bartels E, Mydlarz LD 2011 Patterns of coral ecological immunology: variation in the responses of Caribbean corals to elevated temperature and a pathogen elicitor. *J Exp Biol* 214, 4240–9. doi:10.1242/jeb.061267. [PubMed: 22116768]
- Pinzon JH, Kamel B, Burge CA, Harvell CD, Medina M, Weil E, Mydlarz LD 2015 Whole transcriptome analysis reveals changes in expression of immune-related genes during and after bleaching in a reef-building coral. *R Soc Open Sci* 2, 140214–140214. doi:10.1098/rsos.140214. [PubMed: 26064625]
- Prada C, Weil E, Yoshioka PM 2010 Octocoral bleaching during unusual thermal stress. *Coral Reefs* 29, 41–45. doi:10.1007/s00338-009-0547-z.
- Pratchett M, Munday P, Wilson S, Graham N, Cinner J, Bellwood D, Jones G, Polunin N, McClanahan T 2008 Effects Of Climate-Induced Coral Bleaching On Coral-Reef Fishes -Ecological And Economic Consequences. *Oceanography and Marine Biology: An annual review* 46, 251–296.
- Pratchett MS, McCowan D, Maynard JA, Heron SF 2013 Changes in bleaching susceptibility among corals subject to ocean warming and recurrent bleaching in Moorea, French Polynesia. *PLoS One* 8, e70443. doi:10.1371/journal.pone.0070443. [PubMed: 23922992]
- Precht WF, Robbart ML, Aronson RB 2004 The potential listing of *Acropora* species under the US Endangered Species Act. *Mar Pollut Bull* 49, 534–6. doi:10.1016/j.marpolbul.2004.08.017. [PubMed: 15476831]
- Reyes-Bermudez A, Miller DJ, Sprungala S 2012 The Neuronal Calcium Sensor Protein Acrocalcin: A Potential Target of Calmodulin Regulation during Development in the Coral *Acropora millepora*. *PLoS ONE*, 7(12), e51689 10.1371/journal.pone.0051689. [PubMed: 23284743]
- Richier S, Rodriguez-Lanetty M, Schnitzler CE, Weis VM 2008 Response of the symbiotic cnidarian *Anthopleura elegantissima* transcriptome to temperature and UV increase. *Comp Biochem Physiol Part D. Genomics Proteomics* 3, 283–9. doi:10.1016/j.cbd.2008.08.001. [PubMed: 20494848]
- Roth MS, Latz MI, Goericke R, Deheyn DD 2010 Green fluorescent protein regulation in the coral *Acropora yongei* during photoacclimation. *J Exp Biol* 213, 3644–55. doi:10.1242/jeb.040881. [PubMed: 20952612]
- Schwarz JA, Brokstein PB, Voolstra CR, Terry AY, Miller DJ, Szmant AM, Coffroth MA, Medina M 2008 Coral Life History and Symbiosis: functional genomic resources for two reef building Caribbean corals, *Acropora palmata* and *Montastraea faveolata*. *BMC Genomics* 9, 97. doi: 10.1186/1471-2164-9-97. [PubMed: 18298846]
- Smith-Keune C, Dove S, 2008 Gene expression of a green fluorescent protein homolog as a host-specific biomarker of heat stress within a reef-building coral. *Mar Biotechnol* 10: 166–80. doi: 10.1007/s10126-007-9049-6. [PubMed: 17994304]

- Tomanek L 2010 Variation in the heat shock response and its implication for predicting the effect of global climate change on species' biogeographical distribution ranges and metabolic costs. *J Exp Biol* 213, 971–979. doi:10.1242/jeb.038034. [PubMed: 20190122]
- Tomanek L 2011 Environmental proteomics: changes in the proteome of marine organisms in response to environmental stress, pollutants, infection, symbiosis, and development. *Ann Rev Mar Sci* 3, 373–99. doi:10.1146/annurev-marine-120709-142729.
- Voolstra CR, Schnetzer J, Peshkin L, Randall CJ, Szmant AM, Medina M 2009 Effects of temperature on gene expression in embryos of the coral *Montastraea faveolata*. *BMC Genomics* 10, 627. doi: 10.1186/1471-2164-10-627. [PubMed: 20030803]
- Weil E 2004 Coral Reef Diseases in the Wider Caribbean, in: Rosenberg E and Loya Y (eds). *Coral Health and Disease* Springer-Verlag, Berlin, 35–68 p.
- Weil E, Smith G, Gil-Agudelo DL 2006 Status and progress in coral reef disease research. *Dis Aquat Organ* 69,1–7. doi:10.3354/dao069001. [PubMed: 16703761]
- Wilkinson C 1998 Status of coral reefs of the world: 1998 Australian Institute of Marine Science.
- Wirshing HH, Baker AC 2014 Molecular evolution of calcification genes in morphologically similar but phylogenetically unrelated scleractinian corals. *Mol Phylogenet Evol* 77, 281–295. doi: 10.1016/j.ympev.2014.04.015. [PubMed: 24780747]
- Witze A 2015 Corals worldwide hit by bleaching. *Nature* doi:10.1038/nature.2015.18527.
- Yuyama I, Harii S, Hidaka M 2012 Algal symbiont type affects gene expression in juveniles of the coral *Acropora tenuis* exposed to thermal stress. *Mar Environ Res* 76, 41–7. doi:10.1016/j.marenvres.2011.09.004. [PubMed: 22001189]

Highlights:

- A comparative proteomic approach was undertaken with the endangered Caribbean coral *Acropora palmata*
- 38 significantly differentially expressed proteins were identified by peptide mass fingerprinting in bleached and unbleached *A. palmata* colonies.
- Bleached *A. palmata* key protein signatures were involved in heat stress/UV responses, immunity, apoptosis, biomineralization, cytoskeletal components, and endoexophagocytosis as compared to those colonies of *A. palmata* that did not bleach under the same environmental conditions.
- This is a significant contribution to the fields of marine biology and climate change since this is the first proteomics study in the iconic, and endangered Caribbean coral species, *Acropora palmata*.

Figure 1. Mean sea surface temperatures at depths 0 m and 3 m depth registered during the massbleaching event of 2010 in La Parguera, Puerto Rico. Light blue bar represents the maximum temperature registered on August and September.

Figure 2.

Two-dimensional gels showing differentially expressed proteins in *Acropora palmata*. (A) Unbleached *A. palmata*, (B) Bleached *A. palmata*. The red circles on the 2D gels display the differentially expressed protein spots (fold change ≥ 2 , $P \leq 0.05$). Downregulated and upregulated proteins are indicated with downward and upward arrows, respectively. Examples of common proteins compared between bleached and unbleached *A. palmata* are indicated with yellow squares and capital letters.

Figure 3.

Proposed model highlighting the bleaching response pathways highlighting key protein signatures with differences in abundance in bleached *Acropora palmata* compared with unbleached *A. palmata*. Downregulated proteins are indicated with downward arrows and upregulated proteins are indicated with upward arrows.

Figure 4.
2D-GE western blot of five key differentially expressed proteins (myosin, collagen, CD63, calmodulin, and actin) in *Acropora palmata*.

Acropora palmata proteins identified by MS/MS/Mascot search engine (Matrix Science) and BLAST analysis. The reported proteins have a 100% homology match with the peptides identified by mass spectrometry.

Table 1.

Spot No.	Protein Name	Accession No.	Molecular activity	pI ^a	M _w ^b	P _c ^d	M _s ^d	E _e	SD NB ^f	SD B ^g	H ⁱ
Stress Response											
1	Heat shock transcription factor	ACH53605.1	Proteins respond to oxidative damage, metal toxicity.	6.38	8.96	1	85	↑	0.15	0.212.40
2	Green fluorescent protein FP512	ACH89428	Light-harvesting complex.	8.61	25.92	13	85	↑	0.31	0.412.43
3	NADH-ubiquinone Oxidoreductase chain 5	Q8SJB8	ATP binding.	9.06	67.57	2	68	↑	0.32	0.462.43
4	RecName: Full=Histone H2B	P35067.1	Choline dehydrogenase activity.	10.43	13.8	8	71	↑	0.15	0.212.40

Spot No.	Protein Name	Accession No.	Molecular activity	pI	M _w	Pc	M _s	E _d	SD NB _f	SD Fold change _h	H ⁱ
5	Actinin 1 protein [Acropora digitifera]	BAQ19091.1	Transferase activity, oxidative stress.	8.71	48.39	1	7	0	0.49	0.592.20
6	Photosystem I P700 chlorophyll A apoprotein A1	AAN39416.1	Photosynthesis.	6.38	51.37	6	6	9	0.23	0.342.47
7	Heat Shock protein Hsp-16.2	KJD18266.1	Refolding of denatured proteins	5.06	17.93	7	6	7	0.12	0.162.36
UV response											
8	TPA: fluorescent protein 5, partial [A. digitifera]	FAA00742.1	TPA: fluorescent protein 5, partial [A. digitifera]	7.75	21.18	5	8	3	0.13	0.292.90

Spot No.	Protein Name	Accession No.	Molecular activity	pI	M _w	P _c	M _s	E _d	SD NB _f	Fold change _h	H ⁱ
9	ATPase subunit 6 (ATP synthase subunit 6)	Q9TBW1	ATPase subunit 6 (ATP Synthase subunit 6).	7.92	25.37	11	87	↑	0.41	0.622.08
Amino acid synthesis											
10	Adi-cysteine synthase	F2Z9Y4	Amino acid synthesis related genes.	8.04	7.36	92	92	↓	0.47	0.382.17
Transcription factors											
11	PaxC homeodomain protein [A. palmata]	Q9NBX3	Sequence-specific DNA binding.	4.56	4.79	30	90	↓	0.87	0.202.50
12	PaxC [A. palmata]	B6DC09	DNA binding domains.	5.01	2.99	10	52	↓	0.79	0.562.40

Spot No.	Protein Name	Accession No.	Molecular activity	pI	M _w	Pc	MS	E _d	SD NB _f	Fold change _h	H ⁱ
13	Mannose-binding lectin, partial [A. millepora]	B3V952	Carbohydrate binding.	4.64	22.4	2	1	1	0.32	0.192.25
Immunity-related											
14	Apextrin	A0T3F5	Innate immune response.	5.08	96.0	1	4	5	0.34	0.222.09
15	Ectonucleoside Triphosphate Diphosphohydrolase 1	ABD97868.1	G-protein coupled receptor signaling pathway.	6.85	65.0	1	3	1	0.40	0.302.33
16	ATP synthase subunit 8	NP_612827.1	Hydrogen ion transmembrane transporter activity.	9.51	88.52	6	3	9	0.70	0.542.29

Spot No.	Protein Name	Accession No.	Molecular activity	pI	M _w	Pc	M _s	E _e	SD NB _f	Fold change _h	H ⁱ
17	Toll-like receptor 2	KJD20730.1	Transmembrane signaling receptor activity.	8.69	2592	6	4	↑	0.44	0.352.25
Apoptosis											
18	Caspase 8 [A. palmata]	ADG23096.1	Cysteine-type endopeptidase activity and apoptotic processes	5.94	5079	10	72	↑	0.0	0.19UE ⁺
19	Galaxin	ADI50285.1	Soluble protein of the organic matrix, pro-apoptotic.	5.45	3512	66	66	↓	1.00	0.552.07
Biomining											
20	Galaxin-like 2	ADI50285.1	Biomining related genes. Rich in amino acids containing genes.	4.82	6432	11	53	↓	0.84	0.712.18

Spot No.	Protein Name	Accession No.	Molecular activity	pI	M _w	P _c	M _s	E _e	SD NB _f	Fold change _h	H ⁱ
21	Integrin beta	AAB66910.1	Cell-matrix adhesion.	6.41	87.64	7	254	↓	0.39	0.292.34
Cytoskeletal											
22	Gelsolin	XP_012559373.1	Phagocytosis, engulfment	5.83	40.5	1	91	↓	0.69	0.532.30
23	Myosin	XP_001637809.1	Calcium-regulated, actinmodulating protein.	6.82	3.27	2	75	↓	0.43	0.342.26
24	Actin	ABY40470.1	Calcium-regulated, actinsevering protein.	5.26	17.33	1	76	↑	0.05	0.132.60
Cell cycle											

Spot No.	Protein Name	Accession No.	Molecular activity	pI	M _w	Pc	Ms	E _d	SD NB _f	SD Fold change _h	H ⁱ
25	Cyclin dependent kinase like 2	BAG71255.1	Cell cycle related genes.	5.54	21.89	9	439	↑	0.0	0.73 UE*
26	Calmodulin-like protein [A. millepora]	ACY07618.1	Genes for signal transduction.	4.74	19.66	54	344	↑	0.0	0.66 UE*
Oxidative phosphorylation											
27	Cytochrome oxidase subunit I [A. palmata]	BAG71255.1	Component of the respiratory chain that catalyzes the reduction of oxygen to water.	5.54	21.89	2	18	↓	0.78	0.32 2.43
28	Cytochrome b [A. palmata]	AAD20279.1	Respiratory electron transport chain.	8.41	36.55	2	15	↓	0.60	0.48 2.25

Spot No.	Protein Name	Accession No.	Molecular activity	pI	M _w	P _c	M _s	E _e	SD NB _f	Fold change ^h	H ⁱ
29	Cytochrome P450 74A [A. palmata]	B4YFB4	Electron carrier activity.	8.99	49.80	2	1	1	0.57	0.342.67
Anti-oxidant proteins											
30	Thioredoxin reductase [A. millepora]	AFI99106.2	Oxidoreductase.	6.37	74.30	5	7	7	0.21	0.192.09
Endo-exo phagocytosis-related											
31	CD 63 antigen	O96961	Growth Regulation.	4.34	26.71	7	5	4	0.26	0.0 UE*
32	Light-harvesting protein	CB183412.1	Hydrolase activity.	8.01	28.97	6	4	9	0.12	0.0 UE*

Author Manuscript

Spot No.	Protein Name	Accession No.	Molecular activity	pI ^a	M _w ^b	P _c ^d	M _S ^d	E _e	SD NB _f	Fold change ^h	H ⁱ
38	Mini-collagen	Q8MUZ4	Collagen.	6.18	5.61	9	241	↓		0.0 UE*

^a Isoelectric point

^b Molecular mass

^c Peptide quantity

^d MS score

^e Protein expression: Upregulated (↑) and downregulated (↓)

^f Standard deviation of unbleached *A. palmata*

^g Standard deviation of bleached *A. palmata* h Fold change

ⁱ Histograms: Black bars correspond to unbleached *A. palmata* and striped bars correspond to bleached *A. palmata*.

* Uniquely expressed protein