CORRESPONDENCE

Hematological Malignancies in Adults with a Family Predisposition

by PD Dr. med. Tilmann Bochtler, Dr. med. Georg-Martin Haag, Prof. Dr. med. Sarah Schott, PD Dr. med. Matthias Kloor, Prof. Dr. med. Alwin Krämer, and Prof. Dr. med. Carsten Müller-Tidow in issue 50/2018

Supplemental Human Genetics Point of View

In their review, Bochtler et al. (1) describe the importance of genetic predispositions for hematological neoplasias. From a human genetics perspective, we would like to add the following clinically relevant aspects.

We note that, in contrast to the authors' statement, the German Gene Diagnostics Act (*Gendiagnostikgesetz*, *GenDG*) does not require genetic counseling for diagnostic genetic examinations. In order to avoid delays in the diagnosis, which is in part essential for therapy, diagnostic genetic examinations can be carried out with the patient's written informed consent following appropriate information by a physician. Upon notification of the results, genetic counseling should be offered.

After identification of DNA variants in hematological neoplasms, DNA from fingernails is not suitable for demonstrating the constitutional origin of identified variants. In patients who have received hematopoietic stem cell transplantation, donor DNA has been detected in the genome of the recipient's fingernails (2).

Only sequencing the coding genes and flanking intronic regions is not sufficient when screening for predisposing gene changes. Rather, sequencing must be supplemented by adequate investigations for larger copy number changes in order to detect causative constitutional deletions or duplications, as recommended for *RUNX1*, for example (3).

In addition, genetic studies should also take into account relevant, deeply intronic regions (for example, in *GATA2*) or the possibility of revertant somatic mosaicism (for example, in *SAMD9* / *SAMD9L*).

In contrast to the authors' statement, the risk of developing breast cancer for female carriers of Fanconi anemia, for instance women who have heterozygous *BRCA2* or *PALB2* variants, can be increased significantly higher than threefold (4).

In summary, the challenges that were excellently described by Bochtler et al. show the importance of interdisciplinary cooperation. This cooperation would support early identification of a genetic predisposition for hematological neoplasms in patients, as well as future improvements in the diagnosis and care of those affected.

DOI: 10.3238/arztebl.2019.0251a

References

1. Bochtler T, Haag GM, Schott S, Kloor M, Krämer A, Müller-Tidow C: Hematological malignancies in adults with a family predisposition. *Dtsch Arztebl Int* 2018; 115: 848–54.
2. Imanishi D, Miyazaki Y, Yamasaki R, et al.: Donor-derived DNA in fingernails among recipients of allogeneic hematopoietic stem-cell transplants. *Blood* 2007; 110: 2231–4.
3. Ripperger T, Tawana K, Kratz C, Schlegelberger B, Fitzgibbon J, Steinemann D: Clinical utility gene card for: Familial platelet disorder with associated myeloid malignancies. *Eur J Hum Genet*. 2016; 24: doi: 10.1038/ejhg.2015.278.
4. Easton DF, Pharoah PD, Antoniou AC, et al.: Gene-panel sequencing and the prediction of breast-cancer risk. *N Engl J Med* 2015; 372: 2243–57.

Dr. med. Tim Ripperger, PhD

Institut für Humangenetik, Medizinische Hochschule Hannover

Prof. Dr. rer. nat. Thomas Illig

Institut für Humangenetik, Medizinische Hochschule Hannover
Hannover Unified Biobank, Medizinische Hochschule Hannover

Prof. Dr. med. Brigitte Schlegelberger

Institut für Humangenetik, Medizinische Hochschule Hannover
Kordinatorin, Subnet Familial Leukemia, European Reference Network PaedCan
Schlegelberger.Brigitte@mh-hannover.de

In Reply:

We thank Dr. med. Ripperger, Professor Illig, and Professor Schlegelberger for additions and corrections to our review article from a human genetics perspective (1).

We did not intend to question in any way that the treating clinician can, is allowed to and should initiate cytogenetic and molecular genetic tests in hematological neoplasms, which are already an integral part of clinical routine diagnostics. However, for purposeful germline diagnostics we consider prior genetic counseling (for definition, see German Gene Diagnostics Act) to be necessary in the interest of the patient; for predictive or presymptomatic diagnostics, it is required by law.

DNA from fingernails is listed as possible germline material in the highly detailed expert recommendations from the University of Chicago, but we share the concerns of our Hanoverian colleagues.

We fully agree with the assessment that the sequencing of coding gene regions must be supplemented with additional analyses.

The range we gave for relative risk for breast cancer in female carriers of Fanconi anemia is based on the literature that we cited. However, we acknowledge that other publications have actually shown higher relative risks, such that the relative risk in our review should be revised upwards. DOI: 10.3238/arztebl.2019.0251b

References

1. Bochtler T, Haag GM, Schott S, Kloor M, Krämer A, Müller-Tidow C: Hematological malignancies in adults with a family predisposition. *Dtsch Arztebl Int* 2018; 115: 848–54.

On behalf of the authors

PD Dr. med. Tilmann Bochtler

Medizinische Klinik V, Hämatologie, Onkologie und Rheumatologie,
Universitätsklinikum Heidelberg,
Klinische Kooperationseinheit Molekulare Hämatologie/Onkologie,
Deutsches Krebsforschungszentrum (DKFZ)
Heidelberg
Tilmann.Bochtler@med.uni-heidelberg.de

Conflict of interest statement

The authors of each contribution declare that no conflict of interest exists.