Published in final edited form as:

Occup Environ Med. 2019 June ; 76(6): 357–359. doi:10.1136/oemed-2018-105427.

Let's take the heat out of the CKDu debate: more evidence is needed

Neil Pearce¹ and Ben Caplin²

¹Department of Medical Statistics, London School of Hygiene and Tropical Medicine, United Kingdom

²Centre for Nephrology, Division of Medicine, University College London, London, UK

There is considerable controversy about the causes of the epidemics of Chronic Kidney Disease of undetermined cause (CKDu) which are affecting agricultural communities in Central America, South Asia, and possibly other parts of the world. In this editorial, we argue that currently none of the suggested hypotheses have enough evidence, either for or against them, to draw firm conclusions.

What is CKDu and who gets it?

The common clinical features of CKDu are impaired kidney function in the absence of diabetes, primary glomerular nephritis, or structural abnormality. CKDu exists in Central America[1] and South Asia (India and Sri Lanka)[2, 3], and mainly affects male rural agricultural workers. It may also be occurring in other tropical/subtropical parts of the world, but standardized data are not available for comparison[4]. We will therefore focus on the evidence from Central America and South Asia.

CKDu was first described in Central America in 2002[5], and in the rural population of the North Central Province of Sri Lanka in 2000[6]. However, the estimates in Sri Lanka are not comparable with those from Central America, as in the Sri Lankan prevalence studies to date, serum creatinine has only been performed in those with proteinuria, i.e. an albumin-creatinine ratio (ACR) ≥ 30 mg/g, whereas CKDu typically presents with either no or low-grade proteinuria in Central America[7]. Therefore, valid prevalence estimates can currently only be obtained by identifying renal impairment in random population surveys[4]. Nonetheless, other potential problems of international comparisons arise because the estimated Glomerular Filtration Rate (eGFR) calculated from the serum creatinine is a function of ethnicity[8], exercise, muscle mass and meat consumption.

Causes of CKDu

Perhaps the most clearly established risk factor in both Central America and South Asia, is that CKDu is more common in agricultural communities[9]. In Central America, CKDu

Address for correspondence: Professor Neil Pearce, Department of Medical Statistics, Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, United Kingdom, neil.pearce@lshtm.ac.uk, Website: <http://www.lshtm.ac.uk>.

occurs frequently in sugar cane workers, but also in other occupational groups, including other agricultural workers, fishermen, miners, and construction workers[10]; it also occurs at a lower frequency in women, most of whom have not worked in agriculture. Furthermore, in a recent study in Nicaragua we found that marked decline in kidney function was occurring in a subgroup of participants, whereas the rest of the participants were maintaining normal kidney function[11]; the key exposures in this high-risk subgroup are yet to be identified, but it involved a wide range of agricultural jobs (not just sugarcane), and also involved women who did not work in agriculture.

A large number of specific causes, related to agriculture, have been suggested for CKDu. Heat/dehydration, infection/inflammation and pesticides are the main hypotheses that have been proposed for Central America, whereas in South Asia the emphasis has been on the possible roles of water contamination/metals and/or pesticides.[12] Other possible causes include non-steroidal anti-inflammatory drugs (NSAIDs), population genetics and alcohol [1]. However, a recent systematic review in Central America found that the only risk factors with strong evidence were male sex, family history of CKD, high water intake, and low altitude; data on the other hypotheses which have been proposed, including self-reported heat stress, pesticide exposure, alcohol, and NSAIDs, have little or no support from epidemiological studies[1, 13].

Each of these proposed causes is discussed in detail below.

Heat stress and dehydration

Heat stress and dehydration has become the favoured explanation for the CKDu epidemic in Central America. The report of the 2015 Regional Workshop[14] concluded that “there is growing evidence for a causal role of strenuous work, heat and insufficient rehydration as risk factors for MeN”, although it was also noted that “it is also quite possible that other factors play a role in the disease, perhaps in combination with heat stress and dehydration” and that “exposures to specific agro-chemicals or other yet-unknown toxins need further evaluation”.

The case for heat stress nephropathy has been summarised in a recent review[15], which argues that “recurrent heat exposure with physical exertion and inadequate rehydration can lead to CKD”, and that the excesses in male sugarcane workers arise because they work in extreme conditions of heat exposure and work intensity, as do agricultural workers in many other countries[16]. The authors argue that similar patterns have been observed in South Asia, although heat stress and dehydration are not the currently favoured causal hypotheses in this region [17]. In response, Herath et al [9] give the case against heat-stress nephropathy. They argue that there is sparse evidence of acute kidney injury (AKI) among high risk manual workers, and little evidence that repeated moderate elevations in serum creatinine will lead to CKD, that the animal studies involve extreme forms of AKI which are not observed in workplace-based epidemiological studies, that CKDu is not seen in workers exposed to heat in most tropical regions, that the disease is seen in people not exposed to heat stress in the affected regions, and that in general there is inadequate evidence for it being the initiating and/or main cause of a major global epidemic of CKDu. Similarly, Ordunez et al[18] have concluded that the overall time trends and mortality patterns for

CKD in Central America suggest that the heat/dehydration hypothesis cannot fully explain the epidemic. VanDervort et al[19] have reached similar conclusions regarding the spatial distribution of CKDu in El Salvador.

Heat stress/dehydration is the only hypothesis for which there is also evidence from an intervention study. This was conducted under extremely difficult conditions in El Salvador[20, 21], resulting in great difficulties in following the non-intervention comparison group. The study clearly shows a reduction in heat stress and cross-shift changes in serum creatinine in the intervention group, but the differences between the intervention and comparison groups in cross-harvest measures are small and not statistically significant; furthermore, both groups showed a reduction in eGFR over time which is consistent with a seasonal effect, which has also been observed in the neighbouring country of Nicaragua[11].

Pesticides

Pesticides have been proposed as the cause(s) of CKDu in both Central America and South Asia, although they have perhaps received more attention in the latter context [22]. Given the link with agricultural communities, it is not surprising that pesticides have been suggested as possible causes, particularly since many pesticides in common use are known human nephrotoxins[23]. Nevertheless, the 1st International Mesoamerican Nephropathy (MeN) Workshop in 2012 concluded that pesticides were an unlikely cause of CKDu, and that heat stress and dehydration were the more likely key etiologic factors[24]. Our own systematic review also found little evidence that pesticides were the main cause of CKDu in Central America[1]. More generally, Valcke et al[25] identified 21 relevant analytic studies, but concluded that most had inadequate information on pesticide exposure. The four studies with stronger designs, from Sri Lanka, India and the USA, all showed clear associations, but for different pesticides in each study, and three of these were in areas without CKDu epidemics. They conclude that pesticides can cause AKI, but that there is no clear evidence that they are causing the CKDu epidemics. Jayasumana et al[26] have hypothesized a joint effect of glyphosate and metals in water in Sri Lanka, but no associations have been seen in other regions; glyphosate is widely used in Central America, but one study involving six-month follow-up did not show kidney effects in pesticide applicators[27].

Metals and other contaminants in water

Metals and other contaminants in water have received particular attention as possible causes of CKDu in South Asia[28], particularly with regards to the possible roles of arsenic, cadmium, fluoride and aluminium. However, studies in both Central America[29] and South Asia[23] have consistently found low levels of these contaminants in water and/or urine in CKDu-affected populations. The potential role of silica has been of particular concern in India[30] although these findings have yet to be replicated. Jayasumana et al[31] hypothesized that arsenic was the likely cause of CKDu in Sri Lanka, but studies have generally not found high levels of arsenic in drinking water and/or urine[23]. Similarly, fluoride in water has been raised as a possible cause of CKDu in Sri Lanka[23], but this hypothesis remains highly speculative.

Infection/inflammation

Evidence for an infective/inflammatory cause rests on seroprevalence studies in those affected [10] and case series from both Sri Lanka and Nicaragua describing an acute febrile syndrome in patients presenting to health services with renal dysfunction[32–34]. The acute inflammation observed on renal biopsy in these studies is certainly consistent with a response to an infectious aetiology or alternative inflammatory precipitant. However, whether these cases represent “Clinical Evidence of Acute Mesoamerican Nephropathy” [33], as described in the Nicaraguan studies, or rather AKI in response to an unspecified infectious illness occurring in those at high risk of renal decline due to an existing CKDu of a non-infectious cause cannot be distinguished with this type of study design. More recently, there has been evidence linking hantavirus infection to CKDu cases in Sri Lanka[35] and it has been suggested that the excess mortality among children and adolescents indicates a role of early and long-term exposure to infectious and/or other toxic agents[18, 36].

Other possible causes

Evidence regarding other possible causes is largely anecdotal. These include NSAIDs, alcohol, and intake of fructose-rich soft drinks. Currently, there is little good evidence from population-based studies which would allow these hypotheses to be either confirmed or rejected[23].

Multifactorial hypotheses

Finally, several reviews[9, 15, 25, 36, 37] have concluded that the cause(s) of CKDu are likely to be multifactorial, basically because no single cause has been discovered, and therefore it is possible that two or more of the various possible causes may work in combination. In fact, most diseases are multifactorial[38], but the causes of epidemics usually are not. For example, the epidemics of lung cancer in the 20th century were caused by increased exposure to just one of the causes (i.e. smoking), whereas other contributing factors (e.g. genetics) had not changed, and therefore cannot account for the 20th century epidemics. The likelihood of a single factor playing a key role in the CKDu epidemics, is perhaps enhanced by our recent findings from Nicaragua, in which rapid decline in eGFR was confined to a subgroup of the population[11]. It is also noteworthy that there are well-known examples where the cause(s) of a particular epidemic, or of regional differences in disease risk, took some years or decades to be discovered, and involved several competing hypotheses.

One of the most interesting examples is the epidemic of Balkan Endemic Nephropathy (BEN)[39], which is essentially another type of CKDu. Over a period of 50 years, the hypotheses on the causes of BEN included mycotoxins, metals, viruses, and trace-element insufficiencies[39]. It now seems clear that the main cause was chronic dietary exposure to aristolochic acid (AA) which is the principal component of *Aristolochia clematitis* which grows as a weed in wheat fields in the Balkans[39]. Thus, our lack of success in discovering the cause(s) of CKDu to date does not mean that the disease is multifactorial; it is still possible that one of the hypothesized risk factors, or something we have not yet thought of, is the key driving factor behind the epidemics.

Where to from here?

So even if we don't know the cause(s), why don't we just intervene anyway? The main possible interventions would involve better working conditions, lower pesticide exposure, prevention of infection and cleaner water supplies. All of these should be supported because they will improve working conditions and prevent disease in general, even if their likely impacts on CKDu are currently uncertain. An equally high priority is better research taking an agnostic approach. Clearly, a number of different research methodologies will ultimately contribute to solving this mystery, but in the table we suggest a few key epidemiological questions that address the missing pieces of the jigsaw. What is needed is not further studies or reviews which try and sell the case that a particular risk factor, or combination of risk factors, is the culprit. What is needed instead is smart epidemiology, in which we try and distinguish between the various hypotheses, and develop new ones.

Acknowledgements

This work was supported by grants from the UK Medical Research Council (MR/P02386X/1) and the Colt Foundation. The Centre for Global NCDs is supported by the Wellcome Trust Institutional Strategic Support Fund, 097834/Z/11/B. We wish to thank Nalika Gunawardena, Anthony Newman-Taylor, Agnes Soares, George Davey Smith and Jan Vandenbroucke for their comments on the draft manuscript.

References

1. Gonzalez-Quiroz, ea. What do epidemiological studies tell us about the chronic kidney disease of undetermined cause in Maso-America? A systematic review and meta-analysis. *Chronic Kidney Journal*. 2018; 11:496–506.
2. Anon. Prevalance and risk factors for CKDu in the district of Anuradhapura. Ministry of Health; Colombo, Sri Lanka: 2017.
3. O'Callaghan-Gordo C, Shivashankar R, Anand S, Ghosh S, Glaser J, Gupta R, Jakobsson K, Kondal D, Krishnan A, Mohan S, Mohan V, et al. Prevalence of and risk factors for chronic kidney disease of unknown aetiology in India. *BMJ Open*. 2019
4. Caplin B, Jakobsson K, Glaser J, Nitsch D, Jha V, Singh AK, Correa-Rotter R, Pearce N. International Collaboration for the Epidemiology of eGFR in Low and Middle Income Populations – Rationale and core protocol for the Disadvantaged populations eGFR epidemiology study (DEGREE). *BMC Nephrology*. 2016; 18:1.doi: 10.1186/s12882-016-0417-1
5. Trabanino RG, Aguilar R, Silva CR, Mercado MO, Merino RL. [End-stage renal disease among patients in a referral hospital in El Salvador]. *Rev Panam Salud Publica*. 2002; 12(3):202–6. [PubMed: 12396639]
6. Lanerolle RD, Nanayakkara S, Sherifdeen AH, Sheriff R. Demographic characteristics of end stage renal disease in Sri Lanka. *Journal of the Ceylon College of Physicians*. 2000; 33:3.
7. Torres C, Aragon A, Gonzalez M, Lopez I, Jakobsson K, Elinder CG, Lundberg I, Wesseling C. Decreased kidney function of unknown cause in Nicaragua: a community-based survey. *Am J Kidney Dis*. 2010; 55(3):485–96. [PubMed: 20116154]
8. Stevens LA, Claybon MA, Schmid CH, Chen J, Horio M, Imai E, Nelson RG, Van Deventer M, Wang HY, Zuo L, Zhang YL, et al. Evaluation of the Chronic Kidney Disease Epidemiology Collaboration equation for estimating the glomerular filtration rate in multiple ethnicities. *Kidney Int*. 2011; 79(5):555–62. [PubMed: 21107446]
9. Herath C, Jayasumana C, De Silva P, De Silva PHC, Siribaddana S, De Broe ME. Kidney Diseases in Agricultural Communities: A Case Against Heat-Stress Nephropathy. *Kidney International Reports*. 2018; 3(2):271–280. [PubMed: 29725631]
10. Riefkohl A, Ramirez-Rubio O, Laws RL, McClean MD, Weiner DE, Kaufman JS, Galloway RL, Shadomy SV, Guerra M, Amador JJ, Sanchez JM, et al. *Leptospira* seropositivity as a risk factor

- for Mesoamerican Nephropathy. *International Journal of Occupational and Environmental Health*. 2017; 23(1):1–10. [PubMed: 28209095]
11. Gonzalez-Quiroz M, Smpokou E-T, Silverwood RJ, Camacaho A, Faber D, Garcia BLR, Oomatia A, Hill M, Glaser J, Le Blod J, Wesseling C, et al. Marked decline in kidney function amongst apparently healthy young adults at risk of Mesoamerican nephropathy. *Journal of the American Society of Nephrology*. 2018; 29:2200–2212. [PubMed: 29907699]
 12. Correa-Rotter R, Wesseling C, Johnson RJ. CKD of unknown origin in Central America: the case for a Mesoamerican nephropathy. *Am J Kidney Dis*. 2014; 63(3):506–20. [PubMed: 24412050]
 13. Lunyera J, Mohottige D, Von Isenburg M, Jeuland M, Patel UD, Stanifer JW. CKD of Uncertain Etiology: A Systematic Review. *Clin J Am Soc Nephrol*. 2016; 11(3):379–85. [PubMed: 26712810]
 14. Wegman, D; Crowe, J; Hogstedt, C; Jakobsson, K; Wesseling, C, editors. *Mesoamerican Nephropathy: Report from the Second International Research Workshop on MeN; Heredia, Costa Rica: SALTRA; 2015.*
 15. Glaser J, Lemery J, Rajagopalan B, Diaz HF, Garcia-Trabanino R, Taduri G, Madero M, Amarasinghe M, Abraham G, Anutrakulchai S, Jha V, et al. Climate Change and the Emergent Epidemic of CKD from Heat Stress in Rural Communities: The Case for Heat Stress Nephropathy. *Clinical Journal of the American Society of Nephrology*. 2016; 11(8):1472–1483. [PubMed: 27151892]
 16. Moyce S, Mitchell D, Armitage T, Tancredi D, Joseph J, Schenker M. Heat strain, volume depletion and kidney function in California agricultural workers. *Occupational and Environmental Medicine*. 2017; 74(6):402–409. [PubMed: 28093502]
 17. Anon. International Expert Consultation on CDKu- 2016. Paper 7: The potential role of heat stress, heavy labour and dehydration in the causation and progression of chronic kidney disease (including CKDu). Colombo, Sri Lanka: 2016.
 18. Ordunez P, Nieto FJ, Martinez R, Soliz P, Giraldo GP, Mott SA, Hoy WE. Chronic kidney disease mortality trends in selected Central America countries, 1997–2013: clues to an epidemic of chronic interstitial nephritis of agricultural communities. *Journal of Epidemiology and Community Health*. 2018; 72(4):280–286. [PubMed: 29437864]
 19. VanDervort DR, Lopez DL, Orantes CM, Rodriguez DS. Spatial Distribution of Unspecified Chronic Kidney Disease in El Salvador by Crop Area Cultivated and Ambient Temperature. *Medic Review*. 2014; 16(2):31–38.
 20. Bodin T, Garcia-Trabanino R, Weiss I, Jarquin E, Glaser J, Jakobsson K, Lucas RA, Wesseling C, Hogstedt C, Wegman DH, Group WEPW. Intervention to reduce heat stress and improve efficiency among sugarcane workers in El Salvador: Phase 1. *Occup Environ Med*. 2016; 73(6):409–16. [PubMed: 27073211]
 21. Wegman D. Intervention to diminish dehydration and kidney damage among sugarcane workers. *Scandinavian Journal of Work Environment and Health*. 2018
 22. Anon. International Expert Consultation on CDKu- 2016. Paper 5: Interventions and research priorities on agrochemicals in relation to CKDu in Sri Lanka. Colombo, Sri Lanka: 2016.
 23. Anon. Report of the International Expert Consultation on Chronic Kidney Disease of unknown etiology (CKDu) in Sri Lanka. World Health Organisation, Country Office for Sri Lanka; Colombo, Sri Lanka: 2016.
 24. Wesseling C, Crowe J, Hogstedt C, Jakobsson K, Lucas R, Wegman DH, First International Research Workshop on the Mesoamerican N. Resolving the enigma of the mesoamerican nephropathy: a research workshop summary. *Am J Kidney Dis*. 2014; 63(3):396–404. [PubMed: 24140367]
 25. Valcke M, Levasseur ME, da Silva AS, Wesseling C. Pesticide exposures and chronic kidney disease of unknown etiology: an epidemiologic review. *Environmental Health*. 2017; 16:49. [PubMed: 28535811]
 26. Jayasumana C, Gunatilake S, Siribaddana S. Simultaneous exposure to multiple heavy metals and glyphosate may contribute to Sri Lankan agricultural nephropathy. *Bmc Nephrology*. 2015; 16
 27. Laws RL, Brooks DR, Amador JJ, Weiner DE, Kaufman JS, Ramirez-Rubio O, Riefkohl A, Scammell MK, Lopez-Pilarte D, Sanchez JM, Parikh CR, et al. Biomarkers of Kidney Injury

- Among Nicaraguan Sugarcane Workers. *American Journal of Kidney Diseases*. 2016; 67(2):209–217. [PubMed: 26454687]
28. Anon. International Expert Consultation on CDKu- 2016. Paper 6: Interventions and research priorities on water and heavy metals in relation to CKDu in Sri Lanka. Colombo, Sri Lanka: 2016.
 29. McLean, M, Amador, JJ, Laws, RL, Kaufman, JS, Weiner, DE, Rodriguez, JMS, Rubio, OR, Brooks, DR. Biological sampling report: investigating biomarkers of kidney injury and chronic kidney disease among workers in Western Nicaragua. Boston University School of Public Health; Boston: 2012.
 30. Khandare AL, Reddy YS, Balakrishna N, Rao GS, Gangadhar T, Arlappa N. Role of drinking water with high Silica and Strontium in Chronic Kidney Disease: An Exploratory Community-based Study in an Indian Village. *Indian Journal of Community Health*. 2015; 27(1):95–102.
 31. Jayasumana C, Paranagama PA, Amarasinghe MD, Wijewardane KMRC, Dahanayake KS, Fonseka SI. Possible link of chronic arsenic toxicity with chronic kidney disease of unknown etiology in Sri Lanka. *Journal of Natural Sciences Research*. 2013; 3:64–73.
 32. Badurdeen Z, Nanayakkara N, Ratnatunga NVI, Wazil AWM, Abeysekera TDJ, Rajakrishna PN, Thinnaratchchi JP, Kumarasiri R, Welagedera DD, Rajapaksha N, Alwis APD. Chronic kidney disease of uncertain etiology in Sri Lanka is a possible sequel of interstitial nephritis! Clinicopathological profile of symptomatic, newly-diagnosed CKDu patients. *Clinical Nephrology*. 2016; 86:106–109.
 33. Fischer RSB, Mandayam S, Chavarria D, Vangala V, Nolan MS, Garcia LL, Palma L, Garcia F, Garcia-Trabanino R, Murray KO. Clinical evidence of acute Mesoamerican Nephropathy. *American Journal of Tropical Medicine and Hygiene*. 2017; 97:1247–1256. [PubMed: 28722618]
 34. Fischer RSB. Early detection of acute interstitial nephritis in the genesis of Mesoamerican nephropathy. *Kidney International*. 2017
 35. Gamage CD, Yoshimatsu K, Sarathkumara YD, Kulendiran T, Nanayakkara N, Arikawa J. Serological evidence of hantavirus infection in Girandurukotte, an area endemic for chronic kidney disease of unknown aetiology (CKDu) in Sri Lanka. *International Journal of Infectious Diseases*. 2017; 57:77–78. [PubMed: 28212862]
 36. Hoy, W, Ordunez, P, editors. Epidemic of Chronic Kidney Disease in agricultural communities in Central America. Pan-American Health Organisation (PAHO); Washington, DC: 2017.
 37. Jayasumana C, Gunatilake S, Senanayake P. Glyphosate, hard water and nephrotoxic metals: are they the culprits behind the epidemic of chronic kidney disease of unknown etiology in Sri Lanka? *Int J Environ Res Public Health*. 2014; 11(2):2125–47. [PubMed: 24562182]
 38. Pearce N. Epidemiology in a changing world: variation, causation and ubiquitous risk factors. *International Journal of Epidemiology*. 2011; 40(2):503–512. [PubMed: 21247886]
 39. Stiborova M, Arlt VM, Schmeiser HH. Balkan endemic nephropathy: an update on its aetiology. *Archives of Toxicology*. 2016; 90(11):2595–2615. [PubMed: 27538407]

Table
key epidemiological questions about CKDu

-
- Who gets this disease? Does it occur in all tropical/subtropical regions of the world, or only in some?
 - Does this look like the same disease in different parts of the world? i.e. are we experiencing a single epidemic in different parts of the world, or are different epidemics occurring in parallel?
 - What is the prevalence in various parts of the world which have exposures to some of the risk factors, but not others, e.g. in areas which are hot and involve strenuous work, but not pesticide exposure, and vice versa?
 - Within at risk regions, which population subgroups are most at risk? Is it always rural male agricultural workers (and sugarcane workers in particular)?
 - Within the at risk populations, what determines who suffers from kidney function decline, and can we identify them (e.g. with appropriate biomarkers) prior to the development of established CKD (at a time point when exposure assessment or kidney biopsy might provide more insight)?
-