Research

Cite this article: Pusuluk O, Farrow T, Deliduman C, Vedral V. 2019 Emergence of correlated proton tunnelling in water ice. *Proc. R. Soc. A* **475**: 20180867. <http://dx.doi.org/10.1098/rspa.2018.0867>

Received: 8 December 2018

Accepted: 12 February 2019

Subject Areas:

quantum physics, low temperature physics, chemical physics

Keywords:

quantum information, measures of quantum coherence, measures of quantum correlations, open quantum systems, quantum master equation, low-temperature proton tunnelling in water ice

Author for correspondence:

Onur Pusuluk

e-mail: pusuluk@itu.edu.tr

Electronic supplementary material is available online at <https://dx.doi.org/10.6084/m9.figshare.c.4487114>.

Emergence of correlated proton tunnelling in water ice

Onur Pusuluk^{1,2}, Tristan Farrow^{3,4},

Cemsinan Deliduman⁵ and Vlatko Vedral^{3,4}

¹Department of Physics, Koç University, Sarıyer, İstanbul 34450, Turkey

²Department of Physics, İstanbul Technical University, Maslak, İstanbul 34469, Turkey

³Department of Physics, University of Oxford, Parks Road, Oxford OX1 3PU, UK

⁴Centre for Quantum Technologies, National University of Singapore, 3 Science Drive 2, Singapore 117543, Singapore

⁵Department of Physics, Mimar Sinan Fine Arts University, Bomonti, İstanbul 34380, Turkey

OP, 0000-0002-9167-7273; TF, 0000-0003-3342-3133

Several experimental and theoretical studies report instances of concerted or correlated multiple proton tunnelling in solid phases of water. Here, we construct a pseudo-spin model for the quantum motion of protons in a hexameric H₂O ring and extend it to open system dynamics that takes environmental effects into account in the form of O–H stretch vibrations. We approach the problem of correlations in tunnelling using quantum information theory in a departure from previous studies. Our formalism enables us to quantify the coherent proton mobility around the hexagonal ring by one of the principal measures of coherence, the l_1 norm of coherence. The nature of the pairwise pseudo-spin correlations underlying the overall mobility is further investigated within this formalism. We show that the classical correlations of the individual quantum tunnelling events in long-time limit is sufficient to capture the behaviour of coherent proton mobility observed in low-temperature experiments. We conclude that long-range intra-ring interactions do not appear to be a necessary condition for correlated proton tunnelling in water ice.

1. Introduction

Hydrogen bonding (or H-bonding) is the subject of extensive literature due to its central importance in many natural phenomena in physical, chemical and biological systems. The first attempts that reveal the underlying physics behind it go back to the 1950s [1,2], and since then, quantum aspects of the nature of this weak interaction are still being hotly debated. In the meantime, most of the demystification attempts have focused on the H-bonds using water as an explanatory model [3–6].

Let us designate an H-bonded system $X_1 - H \cdots X_2$, where the single covalent bond $X_1 - H$ is a proton-donating bond, X_1 is the proton-donor and X_2 is the proton-acceptor. One controversial issue about the role of non-trivial quantum effects in such a system is the extent of the covalency of the $H \cdots X_2$ interaction [3], i.e. charge transfer from the lone pair orbital of the proton-acceptor ($\sigma_{LP(X_2)}$) to the unoccupied antibonding orbital of the proton-donating bond ($\sigma_{X_1-H}^*$). Although the covalent contribution to the attractive energy of H-bonds in water is comparable to the electrostatic contribution, the amount of charge transfer itself is of the order of just a few millielectrons [4].

Apart from this intermolecular charge transfer, non-trivial quantum effects also enter into the physics of H-bonding in the form of proton tunnelling back and forth between donor and acceptor. According to diabatic state models [5,6], nuclei of H atoms are likely to tunnel through H-bonds between water monomers. Several *ab initio* studies examined this probability in water ice as well. First and foremost, proton tunnelling was found to be responsible for the pressure-driven phase transitions from proton-ordered ice VIII to proton-disordered ice VII around 100 K [7], and is believed to drive the transition from proton-disordered ice I_h to proton-ordered ice XI in a microscopic model [8].

However, spontaneous single proton tunnellings violate the so-called Bernal-Fowler *ice rules* [9,10] which state that (i) each water molecule is linked to four other ones through H-bonds in such a way that (ii) it behaves as a proton-donor in half of these four bonds and a proton-acceptor in the remaining ones. These local constraints are expected to lead to correlations between individual proton tunnellings. Consistent with this expectation, the likelihood of correlated proton tunnelling in water ice was recently reported by successive low-temperature experiments such as incoherent quasi-elastic neutron scattering measurements on ice I_h and I_c [11], scanning tunnelling microscopy of cyclic water tetramer [12], and high precision measurements of the complex dielectric constant of ice XI [13]. Additionally, the trace of correlations of the individual quantum tunnelling events in water ice has been theoretically explored using several models. One-particle density matrix analysis confirmed the presence of proton correlations in ice VII, but not in ice VIII and I_h [14]. On the contrary, concerted tunnelling of six protons in ice I_h appeared to occur at low temperatures in path integral simulations [15] and in lattice-based calculations [16].

Here, we introduce a pseudo-spin model for the quantum motion of protons in a hexameric H_2O ring. Unlike considerations of pseudo-spins by previous studies [8,16], we do not attempt to impose the collective six proton tunnelling by effectively incorporating a single matrix element into the Hamiltonian, or to map the problem onto a lattice gauge theory. Instead, we develop an extension of the model to open system dynamics and approach the correlation problem from the standpoint of quantum information theory. Temperature dependence of proton correlations in (athermal) equilibrium are monitored by well-known measures of quantumness such as l_1 norm of coherence [17], relative entropy of coherence [17], entanglement of formation [18], concurrence [19], quantum discord [20,21] and geometric measure of discord [22].

2. Model and methods

Although their unit cells belong to different space groups, the basic structures of both ice I_h and XI can be visualized as a hexameric box whose planes are either chair-form or boat-form 3-d hexamers. To reduce the complexity, we restrict our model to a 2-d hexagonal ring with a hydroxyl ion (OH^-) resides in each vertex, as shown in figure 1a. Rigid rotations of the vertices are not taken

Figure 1. (a) The hexagonal lattice. It is made of OH⁻ ions in fixed positions relative to each other. Edges can be interpreted as H-bonds. Protons are allowed to live on the enumerated locations. Ice rules are satisfied only by 2 of the six-proton configurations given in (b,c). Transitions between them require simultaneous relocations of the six protons within the ring in the directions depicted by two-sided arrows. Any other configuration accessible from (b) or (c) by successive proton relocations in H-bonds is called an ionic defect. Bjerrum defects occur in all the other configurations including the remaining six-proton configurations. (Online version in colour.)

into account because of the high energy cost assumed in microscopic models [8] and predicted in experiments [13]. Each edge linking two vertices represents an H-bond and includes two equally likely locations for H⁺ ions. These locations (enumerated in figure 1a) can be regarded as a crystal lattice in which H⁺ ions, or protons, move according to the Hamiltonian

$$H_{\text{Hex}} = \sum_{j=1}^{12} W_j n_j - \sum_{j=1}^{12} J_{j,j+1} (a_j^\dagger a_{j+1} + a_j a_{j+1}^\dagger) + \sum_{j=1}^{12} V_{j,j+1} n_j n_{j+1} + \lambda, \quad (2.1)$$

where subscripts are in mod 12, $n_j = a_j^\dagger a_j$ is the proton number operator at lattice site j , $a_j^\dagger$ and a_j are, respectively, proton creation and annihilation operators that obey the following anti-commutation relations

$$\{a_j, a_k\} = \{a_j^\dagger, a_k^\dagger\} = 0, \{a_j, a_k^\dagger\} = \delta_{jk}. \quad (2.2)$$

On-site energy W_j can be taken as the total potential felt by a proton at j th site, i.e. the sum of a Morse potential describing the single (covalent) bond with the adjacent OH⁻ ion and a Coulomb potential representing the electrostatic attraction to the opposite OH⁻ ion. $J_{j,j+1}$ stands for the orbital interactions which causes proton tunnelling. Its expected dependence on the geometry implies that $J_{j,j+1}^{(\text{edge})} \gg J_{j,j+1}^{(\text{vertex})}$ where the former is the intermolecular proton tunnelling coefficient for the neighbouring sites occupying the same edge, while the latter is the intramolecular proton tunnelling coefficient for the successive sites close to the same vertex. As $J_{j,j+1}^{(\text{edge})}$ should already be quite small compared to other coefficients, we can neglect $J_{j,j+1}^{(\text{vertex})}$. This assumption guarantees the absence of any quantum correlation between the quantum tunnelling events through individual hydrogen bonds in the isolated hexamer. $V_{j,j+1}$ is introduced to penalize two-proton cases associated with the violation of ice rules. Edge sharing sites and vertex sharing sites have different penalty coefficients as well as different tunnelling coefficients. The presence of two protons on the same edge is called as a Bjerrum defect, whereas occupation of both sites near the same vertex is called as an ionic defect. As Bjerrum defects require more energy, $V_{j,j+1}^{(\text{edge})} \gg V_{j,j+1}^{(\text{vertex})}$. Finally, λ is a constant responsible for the total intermolecular interactions between vertices, such as Pauli repulsion, van der Waals interaction and London dispersion.

Symmetry of the lattice provides that $W_j = W$, $J_{j,j+1}^{(\text{edge})} = J$, $V_{j,j+1}^{(\text{edge})} = V_{\text{inter}}$ and $V_{j,j+1}^{(\text{vertex})} = V_{\text{intra}}$.

So far, we have focused only on nearest-neighbour interactions and neglected the further interactions with other neighbours, including the concerted tunnelling of the six protons arising from the collective overlap of the orbitals. The effect of long-range interactions on the proton

dynamics in water ice was in fact proposed to be negligible at low temperatures [8]. However, unlike long-range intra-ring interactions, the long-range inter-ring interactions are expected to be non-negligible for the single-ring dynamics. The electrostatic and topological interactions with adjacent rings should at least have significant effects on the parameters W , J , V_{inter} and V_{intra} . We assume that the effects of other rings on each of these parameters can be approximated, respectively, by a single averaged effect. In what follows W , J , V_{inter} and V_{intra} are recounted as effective parameters that include mean-field averages.

To obtain a pseudo-spin Hamiltonian by preserving the anti-commutation relations (given in equation (2.2)), we apply the Jordan–Wigner transformation for a_j , $a_j^\dagger$ and n_j in equation (2.1) in such a way below

$$\left. \begin{aligned} a_j &= \exp \left(-i\pi \sum_{k=1}^{j-1} \sigma_-^{(k)} \sigma_+^{(k)} \right) \sigma_+^{(j)}, \\ a_j^\dagger &= \exp \left(+i\pi \sum_{k=1}^{j-1} \sigma_-^{(k)} \sigma_+^{(k)} \right) \sigma_-^{(j)}, \\ n_j &= \sigma_-^{(j)} \sigma_+^{(j)}, \end{aligned} \right\} \quad (2.3)$$

where $\sigma_-^{(j)} = |1_j\rangle\langle 0_j|$ and $\sigma_+^{(j)} = |0_j\rangle\langle 1_j|$ with the convention for Pauli z operator that $\sigma_z^{(j)} = |0_j\rangle\langle 0_j| - |1_j\rangle\langle 1_j|$. Note that in contrast to the standard application of the Jordan–Wigner transformation on electron transport phenomena, the creation of a proton at the j th site is an energy lowering process here.

After writing (2.3) in terms of Pauli operators, i.e. $n_j = (1/2)(I^{(j)} - \sigma_z^{(j)})$ and $\sigma_\pm^{(j)} = (1/2)(\sigma_x^{(j)} \pm i\sigma_y^{(j)})$, we substitute it into (2.1) and arrive at the following pseudo-spin Hamiltonian

$$\begin{aligned} H_{\text{Hex}} &= + \sum_{j=1}^6 J_x \left(\sigma_x^{(2j-1)} \otimes \sigma_x^{(2j)} + \sigma_y^{(2j-1)} \otimes \sigma_y^{(2j)} \right) + \sum_{j=1}^6 J_z^{(\text{inter})} \left(\sigma_z^{(2j-1)} \otimes \sigma_z^{(2j)} \right) \\ &\quad + \sum_{j=1}^6 J_z^{(\text{intra})} \left(\sigma_z^{(2j)} \otimes \sigma_z^{(2j+1)} \right) + \sum_{j=1}^{12} B \sigma_z^{(j)} + \tilde{\lambda}, \end{aligned} \quad (2.4)$$

where the superscripts of the Pauli matrices are in mod 12, $J_x = -J/2$, $J_z^{(\text{inter})} = V_{\text{inter}}/4$, $J_z^{(\text{intra})} = V_{\text{intra}}/4$, $B = -(2W + V_{\text{inter}} + V_{\text{intra}})/4$ and $\tilde{\lambda} = \lambda + 6W + 3(V_{\text{inter}} + V_{\text{intra}})/2$. Note that these parameters have some contributions from the mean-field averages of the effects of the surrounding hexamers and the construction of the Hamiltonian guarantees the absence of any quantum correlation between the quantum tunnelling events through individual hydrogen bonds in the isolated hexamer. Also note that the same pseudo-spin formalism described above has been recently used in [23] to investigate the role of proton tunnelling in biological catalysis.

The most general quantum state of pseudo-spins can be described using density matrix formalism such that each computational basis state represents a different configuration of protons. For example, configurations in figure 1b,c are, respectively, represented by basis states $|0101010101\rangle$ and $|1010101010\rangle$, so that the protons can exist in any coherent (or incoherent) superposition of these states during the dynamical evolution of the closed (or open) system. Suppose that the proton residing at the site $j = 2$ in the former configuration moves to the site $j = 1$ by either classical hopping or quantum tunnelling between the initial and final times t_i and t_f . In the case of thermally activated classical hopping, it leaves the site $j = 2$ at $t = t_i$, then enters into the exterior space between the sites, and finally reaches the site $j = 1$ at $t = t_f$. Since we do not take into account its presence in the exterior space between the sites, it disappears at the site $j = 2$ at $t = t_i$ and reappears at the site $j = 1$ at $t = t_f$ in our model, i.e. $\rho(t) = |0\rangle\langle 0|$, where $|0\rangle \equiv |0001010101\rangle$ and $t_i < t < t_f$. Conversely, it will never enter into the exterior space between the sites in the course of its motion during tunnelling, but it will be delocalized between both sites, i.e. $\rho(t) = |\psi\rangle\langle\psi|$, where $|\psi\rangle = \alpha|1\rangle + \beta|2\rangle$, $|1\rangle \equiv |1001010101\rangle$, $|2\rangle \equiv |0101010101\rangle$, $|\alpha|^2 + |\beta|^2 = 1$ and $t_i < t < t_f$.

In this case, it is found at the first site with a probability of $|\alpha|^2$ if its location is measured at any time between t_i and t_f . The decoherence taking place at the end of quantum tunnelling, at $t = t_f$, converts the coherent superposition $|\psi\rangle\langle\psi|$ into incoherent superposition $|\alpha|^2|1\rangle\langle 1| + |\beta|^2|2\rangle\langle 2|$ removing the off-diagonal elements from the density matrix.

The unidirectional tunnelling process described above can be captured by a snapshot of the state within our pseudo-spin formalism, that is to say, we can detect it by evaluating the off-diagonal elements of the pseudo-spin density matrix at a single time. The advantages of our approach go beyond this. Consider the static structures of pseudo-spins, e.g. the ground state during closed-system dynamics or the thermal state during open-system dynamics. When such a stationary state equals to $|\psi\rangle\langle\psi|$, it means that the proton under consideration delocalizes between the first and second sites over a long time period. This can be regarded as a quantum tunnelling of the proton back and forth between these two sites since the state never collapses onto the basis states, which represent the proton localizations at their respective sites (see the same usage of the term in [13,14]). By contrast, we cannot claim a back and forth tunnelling event when the steady state is found to be $|\alpha|^2|1\rangle\langle 1| + |\beta|^2|2\rangle\langle 2|$. Hence, in addition to unidirectional proton tunnelling events in the short time limit, the bidirectional proton tunnelling events in the long-time limit are also described by well-defined density matrices.

Mixed states like the incoherent superpositions above cannot be generated from pure initial states during the closed-system dynamics governed by the Hamiltonian (2.4). But they can be generated from coherent superposition states as a result of environmental decoherence that will be described in what follows.

(a) Open system dynamics

It is hard to draw a generic model of the environment for the motion of protons through H-bonds. Such a model should include at least three kinds of vibrations as each individual H-bond is defined by three geometric parameters, e.g. length of the proton-donating bond, donor–acceptor separation and bond angle. However, a minimalistic model consisting of just the periodic oscillations associated with the lengths of proton-donating bonds seems to be sufficient to describe the low-temperature dynamics of protons in a hexameric H₂O ring as in the following. These oscillations can be incorporated into our model as independent thermal baths around lattice sites with individual self-Hamiltonians

$$H_B^{(j)} = \sum_k \hbar \omega_{j,k} b_{j,k}^\dagger b_{j,k}, \quad (2.5)$$

where $b_{j,k}^\dagger$ and $b_{j,k}$ are phonon creation and annihilation operators associated with the k th oscillator mode at the j th site. We assume that the equilibrium positions of the protons are linearly coupled to the positions of the phonons through

$$H_I = \sum_j n_j \sum_k \left(g_{j,k} b_{j,k}^\dagger + g_{j,k}^* b_{j,k} \right) \propto \sum_j \sigma_z^{(j)} \sum_k \left(g_{j,k} b_{j,k}^\dagger + g_{j,k}^* b_{j,k} \right). \quad (2.6)$$

It is important to realize that the local interaction described above induces the entanglement of each pseudo-spin with the positions of associated phonons. In the absence of spin-spin coupling ($J_x = 0$), the dynamics of the pseudo-spins are fully separated from each other, and each pseudo-spin undergoes a pure dephasing process.

Before extending this discussion to the case of non-vanishing inter-spin coupling, let us first examine the role of memory effects in open system dynamics. The Born–Markov approximation can be justified only if the state of pseudo-spins varies over a time scale much longer than the lifetime of the environmental excitations. The vibration of the O–H bond in O–H $\cdots$ O systems has a period of $\simeq 10$ fs, which corresponds to a stretch harmonic frequency of $\simeq 3500$ cm^{−1}. Unlike the short-lived (≈ 1 ps) H-bonds in liquid water [24], H-bonds survive sufficiently long in ice I_h and the jump time of protons in these bonds is larger than tens of fs, e.g. is equal to 3.7 ps at 5 K

[11]. So, we can describe the picosecond evolution of the pseudo-spins' state ρ on the basis of a Markovian master equation in the following Lindblad form [25]

$$\frac{d\rho}{dt} = -\frac{i}{\hbar}[H_{\text{Hex}} + \hbar H_{\text{LS}}, \rho] + \mathcal{D}(\rho), \quad (2.7)$$

where the Lamb shift Hamiltonian provides a unitary contribution to the open dynamics and reads

$$H_{\text{LS}} = \sum_{\omega} \sum_{j,j'} S_{jj'}(\omega) A_j^{\dagger}(\omega) A_{j'}(\omega), \quad (2.8)$$

whereas the dissipator is defined by

$$\mathcal{D}(\rho) = \sum_{\omega} \sum_{j,j'} \gamma_{jj'}(\omega) \left(A_{j'}(\omega) \rho A_j^{\dagger}(\omega) - \frac{1}{2} \{A_j^{\dagger}(\omega) A_{j'}(\omega), \rho\} \right), \quad (2.9)$$

with $\omega = \epsilon_m - \epsilon_{m'}$. Here, ϵ_m 's are the eigenvalues of pseudo-spin Hamiltonian given in (2.4) and noise operators $A_j(\omega)$ are the eigenoperators of this self-Hamiltonian

$$A_j(\omega) = \sum_{\epsilon_m - \epsilon_{m'} = \omega} |\epsilon_{m'}\rangle \langle \epsilon_{m'}| A_{\alpha} |\epsilon_m\rangle \langle \epsilon_m|, \quad (2.10)$$

where A_j are the Hermitian operators coupled to the environment, i.e. Pauli z operators as introduced in equation (2.6). Coefficients $S_{jj'}(\omega)$ and $\gamma_{jj'}(\omega)$ are, respectively, the imaginary part and half of the real part of the one-sided Fourier transformation of the thermal bath correlation function given by

$$\Gamma_{jj'}(\omega) = \frac{1}{\hbar^2} \int_0^{\infty} ds e^{i\omega s} \left\langle B_j^{\dagger}(s) B_{j'}(0) \right\rangle_{\text{th}}, \quad (2.11)$$

where $B_j(s)$ are the interaction picture representations of the bath operators (included in (2.6)) and $J(\omega)$ is the spectral density function encapsulating all the effects of the environment. We assume that each pseudo-spin is associated with an independent environment, $\Gamma_{jj'} = \Gamma_j \delta_{jj'}$. Furthermore, we focus on a symmetric lattice at a constant temperature T that makes these individual baths identical, so $\Gamma_{jj} = \Gamma$.

The normal modes of lattice vibrations are more complicated in real water ice. The lattice sites, especially the pair of sites sharing the same vertex, are so close to each other that it is expected to find correlations between them. The presence of the correlations between the individual baths may result in the emergence of quantum correlations between the quantum tunnelling events in the course of open system dynamics. However, we would like to restrict our analysis to the importance of classical correlations between the quantum tunnelling events on the overall proton mobility. Hence, we ignore the correlations between the individual baths as well as the Hamiltonian parameter $J_{j,j+1}^{(\text{vertex})}$.

(b) Measures of quantum correlations

Quantum coherence is the degree of quantum superposition found in a generic state ρ with respect to a given orthogonal basis $\{|m\rangle\}$. One of the most widely used measures that satisfy all the requirements for a proper measure of quantum coherence is the l_1 norm of coherence [17], defined as

$$C_{l_1}[\rho] = \sum_{m \neq m'} |\langle m | \rho | m' \rangle|. \quad (2.12)$$

Off-diagonal elements of the density matrix ρ are related to the transitions between computational basis states, and each basis state represents a different configuration of the protons in our model. Hence, the l_1 norm of the pseudo-spins' state quantifies the quantum coherent proton mobility when the proton number is fixed. As an example, $|m\rangle$ equals to $|b\rangle \equiv |01\rangle \otimes^6$ and $|c\rangle \equiv |10\rangle \otimes^6$ for the configurations respectively depicted in figure 1b,c. A transition from one of these states to the other requires simultaneous relocations of the six protons to their

adjacent empty sites in H-bonds. Any quantum coherent superposition $|\psi\rangle = \alpha|b\rangle + \beta|c\rangle$ with $|\alpha|^2 + |\beta|^2 = 1$ represents such a motion if it takes place in the form of concerted tunnelling of six protons between the corresponding configurations. The extent of the quantum character of this motion of the protons is reflected in the density matrix by the off-diagonal elements $\langle b|\rho|c\rangle = \alpha\beta^*$ and $\langle c|\rho|b\rangle = \alpha^*\beta$. The sum of the absolute values of these elements is maximum when $\alpha = \beta = 1/\sqrt{2}$, and this corresponds to a quantum state in which the protons can be found in each of the two configurations with a probability of 1/2 if their locations are measured.

Another proper measure of coherence is the relative entropy of coherence [17]:

$$C_R^{\text{IC}}[\rho] = \min_{\varsigma \in \text{IC}} (S[\rho \parallel \varsigma]) = S[\rho_d] - S[\rho], \quad (2.13)$$

where the minimum is taken over the set of incoherent states (IC) that are diagonal in the basis $\{|m\rangle\}$, $S[\rho \parallel \varsigma]$ is the quantum relative entropy that equals to $-\text{tr}[\rho(\log_2 \rho - \log_2 \varsigma)]$, $S[\rho]$ is the von Neumann entropy that equals to $-\text{tr}[\rho \log_2 \rho]$ and ρ_d is the diagonal part of the density matrix ρ . That is to say, $C_R^{\text{IC}}[\rho]$ measures the distinguishability of a density matrix with a modified copy in which the off-diagonal elements are removed by a full dephasing process. Whereas $C_l[\rho]$ takes into account distinct tunnelling pathways independently of each other, $C_R^{\text{IC}}[\rho]$ does not discriminate between these pathways that rearrange the proton configuration and highlights the overall non-classical mobility, hence showing a holistic picture.

Quantum correlations also arises from the superposition principle. Non-local correlations found in non-separable quantum superposition states are known as quantum entanglement. Entanglement of formation [18] is a good measure of entanglement for a generic bipartite state ρ and is defined as

$$E_F[\rho] = \min \left(\sum_i Q^{(i)} E_E[|\psi_i\rangle\langle\psi_i|] \right), \quad (2.14)$$

where the minimum is taken over all the possible pure state decompositions that realize $\rho = \sum_i Q^{(i)} |\psi_i\rangle\langle\psi_i|$, and E_E is the entropy of entanglement, the unique measure of entanglement for pure bipartite states that equals to the von Neumann entropy of the reduced state of one of the two subsystems, i.e. $E_E[\rho] = S[\rho_{1(2)}] = (S \circ \text{tr}_{2(1)})[\rho]$. Although it is hard to compute $E_F[\rho]$ for a general state, even numerically, an explicit formula in the form of a binary entropy can be derived in the case of two-qubit systems:

$$E_F[\rho] = -f \log_2(f) - (1-f) \log_2(1-f), \quad (2.15)$$

where $f = f(C) = (1 + \sqrt{1 - C^2})/2$ and C is an entanglement monotone, called concurrence [19]. This entanglement monotone is defined as

$$C[\rho] = \max \left(0, \sqrt{\lambda_1} - \sqrt{\lambda_2} - \sqrt{\lambda_3} - \sqrt{\lambda_4} \right), \quad (2.16)$$

where λ_i are the eigenvalues of the operator $\rho(\sigma_y \otimes \sigma_y) \rho^* (\sigma_y \otimes \sigma_y)$ in decreasing order. In addition to providing an explicit formula for the entanglement of formation, the concurrence can also be used to reveal one of the most fundamental properties of entanglement known as the monogamy of entanglement [19], which imposes a trade-off between the amount of entanglement between different subsystems in a composite system.

As each edge in figure 1a represents an H-bond, the entanglement of formation of the reduced state of pseudo-spins lying on the same edge quantifies the entanglement generated by proton tunnelling through the corresponding H-bond. Besides this, non-zero entanglement of formation between a pseudo-spin pair lying on different edges indicates the inter-bond entanglement between the protons that belong to corresponding H-bonds. The trade-off between intra-bond and inter-bond entanglements can then be investigated using the concurrence. However, quantum correlations are not limited to quantum entanglement, e.g. separable mixed states can possess non-classical correlations known as quantum discord [20,21] when the orthogonality condition on local bases breaks down at least in one subsystem and provides local indistinguishability. The

original definition of quantum discord [21] is grounded in the difference between two different quantum generalizations of mutual information:

$$\begin{aligned}\delta_{1:2} &= I[\rho] - J[\rho] \equiv (-S[\rho] + S[\rho_1] + S[\rho_2]) - \max_{\{M_i^{(2)}\}} (S[\rho_1] - S[\rho_1|\{M_i^{(2)}\}]) \\ &= \min_{\{M_i^{(2)}\}} \left(S[\rho_2] + \sum_i P_i S[\rho_{1|M_i^{(2)}}] - S[\rho] \right),\end{aligned}\quad (2.17)$$

where $S[\rho_1|\{M_i^{(2)}\}]$ is the quantum conditional entropy of the first subsystem, given the complete measurement $\{M_i^{(2)}\}$ on the second subsystem and $\rho_{1|M_i^{(2)}} = \text{tr}_2[M_i \rho M_i^\dagger]/P_i$ are post-measurement states of the first subsystem with corresponding probabilities $P_i = \text{tr}_2[M_i^\dagger M_i \rho]$.

Quantum discord is the measure of non-classical correlations which includes entanglement as a subset since the first quantum generalization of mutual information measures the total correlations between subsystems, whereas the second generalization $J[\rho]$ quantifies only classical correlations [20]. In other words, quantum information theory can describe all correlations contained in a complex system in a rigorous way. Classical correlations generated by classical proton hopping through an H-bond can be quantified by the mutual information J between the pseudo-spins lying on the edge corresponding to this bond. Moreover, classical correlations between a pair of protons that belong to different H-bonds in the H₂O hexamer can be captured by $J[\rho_{jj'}]$, where $\rho_{jj'}$ is the reduced state of pseudo-spins j and j' that lie on the edges corresponding to these H-bonds. Note that the classical motion of protons does not appear during the closed system dynamics in our model, but arises from the interaction with the environment.

Since any analytic expression is unknown for the mutual information-based measure $\delta_{1:2}$ in a generic system, it is hard to evaluate it. However, an explicit formula of a distance-based measure known as the geometric measure of discord [22] is available for a generic two-qubit system which can be written in the Bloch representation:

$$\rho = \frac{1}{4} \left(\mathbb{I}_1 \otimes \mathbb{I}_2 + \sum_{i=1}^3 x_i \mathbb{I}_1 \otimes \sigma_i^{(2)} + \sum_{i=1}^3 y_i \sigma_i^{(1)} \otimes \mathbb{I}_2 + \sum_{i,i'=1}^3 T_{ii'} \sigma_i^{(1)} \otimes \sigma_{i'}^{(2)} \right), \quad (2.18)$$

where σ_i stands for the Pauli sigma matrices. Using this representation, the geometric measure of discord can be calculated as:

$$D_{G,1:2}^{\text{ZD}} = \frac{1}{4} \left(\|x\|^2 + \|T\|^2 - k_{\max} \right), \quad (2.19)$$

where x equals to (x_1, x_2, x_3) with $x_i = \text{tr}[\rho(\mathbb{I}_1 \otimes \sigma_i^{(2)})]$, $\|T\|^2 = \text{tr}[T^T T]$ with $T = T_{ii'}|i\rangle\langle i'|$, $k_{\max}$ is the largest eigenvalue of the matrix $K = \mathbf{x}\mathbf{x}^T + T^T T$. This measure actually minimizes the distance of a given state ρ to the set of states with zero discord (ZD) using the metric squared Hilbert–Schmidt norm, $\|\rho - \rho'\|^2 = \text{tr}[\rho - \rho']^2$.

3. Implementation of the model

Almost all of the parameters of the pseudo-spin Hamiltonian (2.4) except J_x can be determined using the tools of quantum chemistry by taking into account all the details of the electronic structure of water ice, e.g. performing a number of different *ab initio* density functional calculations for some of the possible 2^{12} proton configurations. It is also possible to construct a realistic spectral density function based on the molecular dynamics simulations or the density functional theory calculations. However, this would not only increase the demand for computational cost of our model but also reduce its explanatory power since the model parameters are assumed to have contributions from the mean-field averages of the effects of the surrounding hexamers. On the contrary, the present work aims to construct a simple and physically insightful model with the minimum number of parameters that can be estimated from comparisons of the predictions of the model with experimental results. In what follows, we show

that the steady state of equation (2.7) depends on only two parameters in contrast to previous multi-parameter models and the quantum information theoretic analysis of this state is enough to give a quantitative description of the experimental data.

(a) Extension to the physical system

Before elaborating on the final steady state of equation (2.7), we first find a reasonable map between the actual 3-d structures of water ice and the present 2-d model of a single hexamer. If we extended the model by connecting multiple hexamers in 3-d as in the water ice, we would first replace OH^- ions with O atoms in the vertices and increase the number of equally likely locations for protons close to each vertex from two to four, i.e. the local constraints on a single hexamer would change. Keeping this difference in mind, we infer the ordering dynamics in multi-hexamer real structures from the underlying proton dynamics in single hexamers and construct the map between the model and physical system based on the single-hexamer proton relocation events occurring in them.

Both of the ice I_h and XI obey the ice rules. However hexagonal rings of ice XI possess a global proton order which is absent in ice I_h , i.e. H_2O hexamers sharing the same 3-d form and the same orientation have also the same proton configuration. This proton order cannot be preserved in the presence of proton relocation unless each of these hexamers simultaneously switches into another proton configuration through a collective motion of the six protons. Thus, not only each of the 3-d hexamers, but also the whole ice XI crystal being constituted by them is allowed to be found only in two different configurations. The switch between these two configurations can be mapped to the transition between $|01\rangle^{\otimes 6}$ and $|10\rangle^{\otimes 6}$ pseudo-spin states (respectively, depicted in figure 1b,c) as both of them require concerted six-proton relocation.

$|01\rangle^{\otimes 6}$ and $|10\rangle^{\otimes 6}$ pseudo-spin states span the whole subspace in which the single hexamer satisfies the ice rules. In a sense, we assume that this ice rule preserving subspace in the model corresponds to the proton-ordered phase in hexagonal water ice.

Ice I_h is composed of 3-d hexamers that also fulfil the ice rules but do not show global correlation. Proton configuration of these hexamers can be achieved from the ones in ice XI by the proton relocation events occurring through H-bonds and keeping the proton number in each hexamer fixed at six. In our model, similar proton relocation events bring the states living inside the ice rule preserving subspace into another subspace spanned by 62 pseudo-spins representing the ionic defects. Thus, this defective pseudo-spin subspace can be assumed to coincide with the proton-disordered phase in hexagonal water ice.

(b) Asymptotic limit of the model

Here and in the following, we consider the steady-state solution of equation (2.7). The chosen interaction with the environment does not bring the system into a thermal equilibrium in general. On the contrary, it divides 2^{12} -dimensional Hilbert space $\mathcal{H}$ into subspaces $\mathcal{H}_{\mathcal{J}}$ each of which is independently invariant under $\{A_j\}_{j=1}^{12}$ operators. In the asymptotic limit, it provides a detailed balance only inside these subspaces as below:

$$\rho_{\infty} = \sum_{\mathcal{J}} \frac{P(\mathcal{J})}{Z(\mathcal{J})} \sum_{|\epsilon_m\rangle \in \mathcal{H}_{\mathcal{J}}} e^{-\beta \epsilon_m} |\epsilon_m\rangle \langle \epsilon_m|, \quad (3.1)$$

where

$$P(\mathcal{J}) = \sum_{|\epsilon_m\rangle \in \mathcal{H}_{\mathcal{J}}} \langle \epsilon_m | \rho(t=0) | \epsilon_m \rangle \quad (3.2)$$

and

$$Z(\mathcal{J}) = \sum_{|\epsilon_m\rangle \in \mathcal{H}_{\mathcal{J}}} e^{-\beta \epsilon_m}, \quad (3.3)$$

with $\beta = 1/k_B T$.

One of the subspaces $\mathcal{H}_{\mathcal{J}}$ consists of two special kinds of pseudo-spin states mentioned in the previous subsection, i.e. 2 pseudo-spin states obeying the ice rules and 62 pseudo-spin states corresponding to ionic defects. As these state sets are assumed to map to the proton-ordered and disordered phases of the hexagonal water ice, respectively, we label their union by $\mathcal{H}_{\text{ice}}$. If the initial state $\rho(t=0)$ lives only in this 64-dimensional subspace, the state of the pseudo-spins that relax to equilibrium still stays inside the same subspace as follows:

$$\rho_{\infty}^{\text{ice}} = \sum_{|\epsilon_m\rangle \in \mathcal{H}_{\text{ice}}} e^{-\beta \epsilon_m} |\epsilon_m\rangle \langle \epsilon_m| \quad / \quad \sum_{|\epsilon_m\rangle \in \mathcal{H}_{\text{ice}}} e^{-\beta \epsilon_m}. \quad (3.4)$$

It is straightforward to show that the athermal attractor above essentially depends on two free parameters, J_x and $J_z^{(\text{intra})}$ as all the energy eigenvalues of subspace $\mathcal{H}_{\text{ice}}$ have a common functional dependence on the remaining coefficients included in (2.4).

(c) Characterization of proton ordering/disordering

A temperature-dependent transition between two different phases can be characterized by the presence of two different steady states above and below the phase transition temperature. Although our master equation has a unique steady-state solution denoted by $\rho_{\infty}^{\text{ice}}(T)$, it shows different features above and below the phase transition temperature range which are reflected by (i) $P_{\text{BF}}(T)$, the probability of pseudo-spins to be found inside the ice-rule preserving subspace and (ii) $S[\rho_{\infty}^{\text{ice}}(T)]$, the von Neumann entropy of pseudo-spins.

In our model, each basis state represents a different configuration of the protons. As explained in §§3a, the 2-d ice rule preserving subspace spanned by $\{|01\rangle \otimes^6, |10\rangle \otimes^6\}$ can be mapped to the XI phase of water ice based on the proton relocation dynamics. Thus, the change in probability $P_{\text{BF}}(T)$ can be used as an indicator of the proton-disordering phase transition, e.g. it should be close to unity in XI phase and show a decrease during the transition to I_h phase.

The presence of unit probability below the phase transition temperature range means that only four particular elements of $\rho_{\infty}^{\text{ice}}(T)$ can be non-zero. No matter how small a deviation from unity in $P_{\text{BF}}(T)$ is, $64 \times 64 - 4 = 4092$ more elements of $\rho_{\infty}^{\text{ice}}(T)$ can take a non-zero value. This corresponds to an enlargement in the dimension of the effective Hilbert space from 2 to 64, which allows the violation of ice rules that was mapped to the proton-disordered phase in §§3a. Hence, there is no need to a significant decline in $P_{\text{BF}}(T)$ within the phase transition temperature range to indicate the proton-ordered/disordered transition, but it is sufficient for it to gradually deviate from unity which means a turning point behaviour. When it shows a sharp decline above the phase transition temperature range, we have effectively two different steady states above and below the range, each of which has different numbers of non-zero elements and lives in a different Hilbert space.

The von Neumann entropy measures the amount of disorder, uncertainty, or unpredictability of a generic quantum state. Furthermore, each basis state of pseudo-spins represents a different proton configuration in our model. Thus, $S[\rho_{\infty}^{\text{ice}}(T)]$ directly quantifies the proton-disorder.

4. Results

(a) Model parameters

The values of the parameters used here were estimated from comparisons of the predictions of the model with the previous experiments carried out on the ice I_h /XI transition. Actually, it is not easy to observe this transition as protons are expected to become classically immobile around 100–110 K where a glass transformation occurs [26–28]. First, calorimetric measurements [29,30] overcame this problem by using alkali hydroxides as catalyzer and caught the transition at 72 K. According to recent complex dielectric constant measurements of pure water ice reported in [28], hexagonal ice undergoes a phase transition from proton-disordered ice I_h to proton-ordered ice XI at 58.9 K, whereas a reverse transformation occurs at 73.4 K. Since a charge movement should increase the imaginary part of the dielectric constant ϵ'' , these phase transitions were determined

Figure 2. Estimation of the parameters based on the predictions of the model with experiments. $\text{XI} \rightarrow \text{I}_\text{h}$ and $\text{I}_\text{h} \rightarrow \text{XI}$ phase transitions of hexagonal ice, respectively, occur at 58.9 K and 73.4 K [28]. Vertical solid lines (coloured blue and red) pinpoint these temperatures. The dashed curve (in dark blue) is $P_\text{BF}(T)$, the probability of pseudo-spins to be found inside the ice rule preserving subspace spanned by $\{|01\rangle \otimes^6, |10\rangle \otimes^6\}$ when they are prepared in the steady state ρ_∞^ice . We set J_x and $J_z^{(\text{intra})}$, respectively, to -0.5 meV and $+10$ meV to observe the violation of ice rules around the phase transition temperatures. The dotted curve (in dark red) shows the temperature dependence of the von Neumann entropy of ρ_∞^ice . (Online version in colour.)

by detecting the anomalies in the cooling and warming curves of $\varepsilon''(T)$. Whereas a clear peak was observed in the warming curve of $d\varepsilon''/dT$ at 73.4 K, a discontinuous change occurred in the slope of cooling curve of $d\varepsilon''/dT$ at 58.9 K. The former was reflected in the warming curve of $\varepsilon''(T)$ as a continuous increase followed by a change in the slope, while the latter corresponded to a smooth change in the slope of the cooling curve of $\varepsilon''(T)$. See figure 3 in [28] for more details.

As opposed to experimental data on $\varepsilon''(T)$, no hysteresis is expected between the warming and cooling curves of $P_\text{BF}(T)$ since we focus on the steady-state solution of the master equation and do not allow the system under consideration to be driven out of equilibrium due to quantum fluctuations. Moreover, as our model is restricted to a single hexamer, a sharp discontinuity in $P_\text{BF}(T)$ is unlikely to occur during phase transition. Hence, instead of a first-order phase transition, we anticipate observing a smooth change in the slope of $P_\text{BF}(T)$ similar to that of the cooling curve of $\varepsilon''(T)$ shown in figure 3-a in [28] and in figure 2 in [13]. However, even this smooth change in our finite size system can be treated as evidence of a proton-disordering phase transition since it reflects the true microscopic mechanism driving the proton-disordering process (see §§3a and 3c for details).

In this respect, we fixed our free parameters J_x and $J_z^{(\text{intra})}$, respectively, to -0.5 meV and $+10$ meV to reproduce the expected trend of $P_\text{BF}(T)$ using the steady state ρ_∞^ice as shown in figure 2. The state of pseudo-spins stays inside the ice rule preserving subspace with a unit probability while $T < 58.9$ K. The states outside this subspace violate the ice rules and gradually become available between the vertical lines (in blue and red). Above the temperature of ≈ 73.4 K, $P_\text{BF}(T)$ shows a sharp decline that corresponds to an ever-increasing population of ice rule violating states. Details of the procedure used in this parameter estimation are given in Section II-A in the electronic supplementary material.¹

The behaviour of the von Neumann entropy of $\rho_\infty^\text{ice}(T)$ supports the arguments above for the fixed values of J_x and $J_z^{(\text{intra})}$. According to figure 2, it remains at unity until 58.9 K. Note that the pseudo-spins live inside the subspace spanned by $\{|01\rangle \otimes^6, |10\rangle \otimes^6\}$ in the same temperature range. Then this unit disorder is possible only if the pseudo-spins can exist in only two orthogonal states living inside this subspace with an equal probability, i.e. the configurations given in figure 1b,c or two of their coherent superpositions orthogonal to each other are equally likely

¹See the electronic supplementary material for details of parameter estimation procedure, parameter sensitivity test and master equation solutions

Figure 3. First derivative of the free energy with respect to temperature. Since we do not consider any change in the locations of OH^- ions with temperature, we do not include a pressure–volume term in the free energy, i.e. $F = U - k_B TS$, where S is the von Neumann entropy. (Online version in colour.)

Figure 4. Sensitivity of the behaviour of probability $P_{\text{BF}}(T)$ to the fixed values of free parameters J_x (a) and $J_z^{(\text{intra})}$ (b). (Online version in colour.)

for the protons. $S[\rho_\infty^{\text{ice}}(T)]$ rises slowly with further increases of temperature until 73.4 K. Hence, a smooth change occurs in the proton-disorder around 58.9–73.4 K. After that, the slope of $S[\rho_\infty^{\text{ice}}(T)]$ is approximately constant, which shows a rapid increase in the proton-disorder. Note that the first derivative of free energy of $\rho_\infty^{\text{ice}}(T)$ has the same temperature dependence with $P_{\text{BF}}(T)$ and $S[\rho_\infty^{\text{ice}}(T)]$ as shown in figure 3.

Estimation of the free parameters may still look arbitrary at first glance. However, if we decrease the value of J_x , $P_{\text{BF}}(T)$ cannot get close to unity at any temperature (figure 4a), which means that the ice rules are always violated and the system can never end up in XI phase. On the contrary, a change in the value of $J_z^{(\text{intra})}$ sets the temperature at which the ice rules begin to be violated, apart from the experimentally determined phase transition temperatures (figure 4b). Hence, the expected temperature dependence of P_{BF} exhibits a sensitivity to our free parameters, i.e. deviations from the fixed values of either J_x or $J_z^{(\text{intra})}$ that are much smaller than the energy of an H-bond rule out any prediction, preventing the appearance of a slow decline in $P_{\text{BF}}(T)$ from unity around 58.9–73.4 K. Also, this behaviour cannot reappear when the second parameter is also allowed to deviate from its fixed value at the same time. See electronic supplementary material, section II-B (see footnote 1) for the details.

Note that the values of $J_x = -0.5$ meV and $J_z^{(\text{intra})} = +10$ meV are fixed in this way, and do not only stand for the bare coefficients of a single hexamer but also have contributions from the mean-field averages of the effects of the surrounding hexamers.

Figure 5. Coherent quantum effects on proton mobility in a H_2O hexamer. The first two vertical solid lines (coloured blue and red) indicate the phase transition temperatures [28], whereas the last vertical line (in black) is anchored to the temperature of the glass transition at which proton mobility is expected to diminish [26–28]. (Online version in colour.)

(b) Quantum aspects of the proton mobility

We summarized several theoretical and experimental findings suggesting the likelihood of proton tunnelling in hexagonal water ice in §1. Here, we address two of them using the tools of quantum information theory described in §§2b.

The first claim concerns the role of proton tunnelling in I_h/XI phase transition. In [8], the motion of the protons was first mapped into a pseudo-spin model, as we do, but then converted to a gauge theory problem. This description allowed the authors to characterize the ordered and disordered phases, respectively, by confined and deconfined behaviours of ionic defects in the ground state of the system. It was then found that the phase transition under consideration is possible only if the protons tunnel through H-bonds with a rate greater than a critical value (see figure 6-b in [8]). Since the protons are expected to become classically immobile around 100–110 K where a glass transformation occurs [26–28], this is a reasonable claim. Our predictions shown in figure 4a suggest that it may be also unlikely for the hexagonal water ice to end up in XI phase unless the tunnelling rate is less than another critical value. Thus, further investigation of the quantum aspects of the proton mobility in I_h/XI transition using quantum information theory may yield new knowledge about this topic.

The second claim that will be addressed here is the presence of concerted six-proton tunnelling at low temperatures in XI phase. Dielectric constant measurements that determined the phase transition temperatures in pure water ice [28] were extended down to 5 K in [13], and an anomaly was observed in the cooling and warming curves of $\varepsilon''(T)$ in the form of a minimum around 20 K. The monotonic behaviour of the real part of the dielectric constant observed in the same data and disappearance of the anomaly in the repeat measurements on heavy ice were explained by the back and forth tunnelling of protons in groups of six. As mentioned before, the unit disorder of $\rho_{\infty}^{\text{ice}}$ at low temperatures (figure 2) may indicate the presence of two equally likely superpositions of the configurations given in figure 1b,c. Also, each such superposition represents the correlated tunnelling of six protons, and quantum information theory is able to study the nature and extent of this correlation (see §§ 2b).

To provide a first insight into the quantum aspects of proton mobility in a hexameric H_2O loop, we apply the l_1 norm and relative entropy of coherence on $\rho_{\infty}^{\text{ice}}$ as shown in figure 5. $C_R^{\text{IC}}[\rho_{\infty}^{\text{ice}}]$, which quantifies the distinguishability of $\rho_{\infty}^{\text{ice}}$ from its completely decohered version, remains constant throughout the XI phase and steadily decreases with increasing temperature. Hence, there is a rise in the loss of collective quantumness in the global proton mobility starting with the transition from XI phase to I_h phase which lasts thereafter. Conversely, $C_{l_1}[\rho_{\infty}^{\text{ice}}]$, which is the

Figure 6. Pairwise classical correlations. The dashed curve (in black) corresponds to the correlations in a single H-bond. Correlations between two protons, each of which belongs to a different H-bond, is shown by the dotted curve (in dark red). (Online version in colour.)

sum of quantum coherences in individual transitions between proton configuration pairs, shows a different behaviour with respect to temperature. It increases with the $\text{XI} \rightarrow \text{I}_h$ phase transition and reaches a peak around glass transition. Actually, this behaviour seems to be consistent with the experimental data related to real proton mobility, which indicates a local maximum between 60–110 K (see figure 3-a in [28] and figure 2 in [13]) at where the protons are expected to be classically immobile.

The deviation of the temperature dependence of $C_{l_1}[\rho_{\infty}^{\text{ice}}]$ from the experimental data below 60 K is related to the finite size of our model. Although the increase in $\varepsilon''(T)$ during the cooling from 20 K to 5 K arises from the increasing number of protons involved in the correlated six-proton tunnelling events [13], our results are limited to a single hexamer including only six protons. The loss of similarity between the curves of $C_{l_1}[\rho_{\infty}^{\text{ice}}(T)]$ and $\varepsilon''(T)$ above 110 K also originates from the restrictions on our model. The rise in $\varepsilon''(T)$ after the glass transition [13,28] is likely to stem from thermally activated proton hopping, which is expected to suppress quantum coherent proton mobility at these temperatures but outside our scope. Note that the classical motion of the protons enters into our model in the form of incoherent superpositions of pseudo-spins that are generated from coherent superpositions as a result of decoherence.

On the other hand, the uptick in $C_{l_1}[\rho_{\infty}^{\text{ice}}]$ between the blue and red solid lines in figure 5 offers fresh insights about the importance of coherent proton mobility on the proton ordering dynamics [8], i.e. although the tunnelling coefficient is fixed initially, there is an increase in the amount of coherence generated by tunnelling events during the phase transition from proton-disordered ice I_h to proton-ordered ice XI. Also, pinning of $C_{l_1}[\rho_{\infty}^{\text{ice}}(T)]$ at a non-zero value below 60 K eliminates the possibility that the pseudo-spins exist in a maximal mixture of the basis states $|01\rangle^{\otimes 6}$ and $|10\rangle^{\otimes 6}$. Hence, the presence of unit disorder below 60 K in figure 4a should come from a maximal mixture of two orthogonal superpositions of these basis states, i.e. $\rho_{\infty}^{\text{ice}} = 1/2|\psi_+\rangle\langle\psi_+| + 1/2|\psi_-\rangle\langle\psi_-|$, where $|\psi_{\pm}\rangle$ are two coherent superpositions such as $(|01\rangle^{\otimes 6} \pm |10\rangle^{\otimes 6})/\sqrt{2}$. Note that each of the superposition states involved in this mixture can be interpreted as a concerted tunnelling of six protons back and forth between the configurations represented by the states $|01\rangle^{\otimes 6}$ and $|10\rangle^{\otimes 6}$. This observation is in accordance with the dielectric anomaly measured in the form of a minimum around 20 K [13] where we suspect a concerted quantum tunnelling of six protons could be occurring in each hexamer.

A deeper understanding of the overall behaviour of $C_{l_1}[\rho_{\infty}^{\text{ice}}(T)]$ requires investigation of the classical and quantum correlations between pseudo-spin pairs, respectively, shown in figures 6 and 7. Only classical correlations appear between the pseudo-spins lying on different edges according to these figures. This is actually what we expect to see as we prevent the formation

Figure 7. Pairwise quantum correlations in a single H-bond. There is no quantum correlation found between proton pairs belonging to different H-bonds. The upper dot-dashed curve (in dark red) is the concurrence between pseudo-spins lying on the same edge, while the lower dashed curve (in dark red) is the entanglement of formation between them. Their quantum correlations beyond entanglement are measured by quantum discord and its geometric measure, which are displayed, respectively, as the lower dot-dashed and the upper dashed curves (in dark blue). (Online version in colour.)

of quantum correlations between these pseudo-spin pairs by setting the Hamiltonian parameter $J_{j,j+1}^{(\text{vertex})}$ to zero and taking the O–H stretch vibrations as independent from each other. What is unexpected about these results is that the probability change observed in figure 2 resembles the temperature-dependent behaviour of the classical correlations in figure 6, where the mutual information I of the reduced state of the corresponding pseudo-spins is constant around unity throughout the XI phase and starts falling down during the XI/I_h transition. This means that there is an approximately maximal amount of classical correlations between the motions of two different protons that belong to different H-bonds below 60 K.

Beside this, classical correlations between edge sharing pseudo-spins seem to be fixed at unity independent of the temperature. Hence, classical correlations generated by the proton motion in an individual H-bond are invariant under any change in temperature. On the other hand, regardless of the measure that is used to quantify quantum correlations between edge sharing pseudo-spin pairs, quantum correlations generated by proton tunnelling through individual H-bonds are found to be quite low. However, each curve in figure 7 shows a similarity with coherent proton dynamics described by $C_l[\rho_\infty^{\text{ice}}(T)]$ in figure 5. Thus, although the quantum correlations in H-bonds seem to be insignificant when compared to their classical counterparts, temperature dependence of quantum coherent proton mobility still originates from them.

Based on these observations, we surmise that all the individual proton tunnellings throughout six H-bonds found in a single hexamer become classically correlated at low temperatures. These correlations start to weaken during the phase transition from proton-ordered phase ice XI to proton-disordered phase ice I_h. At the same time, quantum correlations between H-bonded atoms become stronger, reaching a maximum during the glass transition around 110–120 K.

5. Future directions

The effect of O–H stretch vibrations is incorporated into the model in the form of Holstein-type local phonon-proton couplings. Such local interactions can originate from the second quantization of the small site displacements after expanding on-site energies W_j around some reference set of coordinates. However, transfer integrals $J_{j,j+1}^{(\text{edge})}$ are also likely to be perturbed by both O–H and O–O vibrations. Moreover, these perturbations should have a different kind of nature, which can be described by non-local phonon-proton couplings well known as Peierls-type interaction. Unlike the Holstein-type interaction, this interaction is not necessarily destructive

and can facilitate proton tunnelling between sites. In fact, the fluctuations of the O–H and O–O bond lengths are expected to have a significant effect on the proton dynamics in real water ice structures. A natural direction to pursue future work is a new quantum master equation approach to open system dynamics of the protons in the presence of both local and non-local phonon couplings. Such a mixed Holstein–Peierls model will reveal the competition between the local and non-local phonon couplings that may be key to understanding this system.

Our first attempts to develop a mixed Holstein–Peierls model using a two-qubit model system given in the electronic supplementary material (see footnote 1) show that although non-local couplings change the dynamics of the system, the static structure of the final steady state remains same. Hence, the bond fluctuations appear not to affect the predictions of the present model unless we do not move on to probe the true proton dynamics around the hexagonal ring in real-time.

The simple pseudo-spin model approach can be readily employed together with the quantum chemical techniques treating electrons and protons quantum mechanically. This additional technique will allow us to give a realistic estimate of both the parameters of the self-Hamiltonian and the form of the spectral density function. It will then be possible to probe the true proton dynamics around the hexagonal ring in real-time. This offers a new opportunity that previous studies did not yield. However, our assumptions about the Markovianity of the open system dynamics may not be justified in this case as we will not work in the long-time regime. Thus, this direction also includes description of the proton dynamics using a non-Markovian evolution.

Besides this, our one-qubit pseudo-spin representation of proton locations is suitable to extend the present model to include nuclear spin degrees of freedom, which are usually ignored by the current modelling approaches in the water literature. Actually, the experimentally determined ratio of ortho/para spin states of isolated single water molecules exhibits a temperature dependence which shares similarities with the proton mobility under consideration in this study. In this respect, we are planning to extend the present model to investigate the possible effects of nuclear spins on proton-ordering dynamics in water ice.

6. Conclusion

We constructed a simple pseudo-spin model to investigate both the possibility and the nature of concerted six-proton tunnelling in a hexameric H_2O employing the tools of quantum information theory. We demonstrated that the static structure of the final steady state of the chosen master equation depends on only two parameters and the quantum information theoretic analysis of this state is sufficient to give a quantitative description of experimental data.

The role of the external environment on the concerted six-proton tunnelling was clearly unveiled as this tunnelling process was not imposed by the self-Hamiltonian but emerged naturally in the long-time limit of the low-temperature dynamics of the open system. Thus, phonon-assistance was found to be central in driving the concerted proton tunnelling up to the temperature of the phase transition from ice XI to ice I_h . Moreover, it was found to be associated with the emergence of ice rules governing the arrangement of atoms in water ice.

Remaining within the framework of the pseudo-spin model enabled us to approach the correlation problem using the tools of quantum information theory. In turn, we inferred that the l_1 norm of coherence [17] is sufficient to capture the behaviour of coherent proton mobility observed in experiments [13,28]. We also discriminated between quantum and classical correlations in concerted proton tunnelling. It was found that the correlations between six proton tunnelling events are not inherently quantum in character. Instead, individual tunnelling events were allowed to be classically correlated only. Low rates and strong correlations were observed for quantum tunnelling events below a critical temperature corresponding to phase transition. Beyond this critical temperature, simulations showed a weakening in the correlations, but an increase in rates. Overall this induces a total increment in the coherent proton mobility until achieving a full proton-disorder.

The finding sheds light on the nature of correlations between the individual tunnelling events that cannot be addressed by previous studies. This also paves the way to investigating the proton dynamics in real-time to advance our understanding of many-proton tunnelling in water ice.

Data accessibility. This work does not have any experimental data.

Authors' contributions. All the authors conceived the idea, derived the technical results, discussed all stages of the project, and prepared the manuscript and figures.

Competing interests. The authors declare no conflict of interest.

Funding. O.P. thanks TUBITAK 2214-Program for financial support. T.F. and V.V. thank the Oxford Martin Programme on Bio-Inspired Quantum Technologies, the EPSRC and the Singapore Ministry of Education and National Research Foundation for financial support.

Acknowledgements. O.P. thanks Alptekin Yıldız for many insightful discussions about the dielectric constant measurements.

References

- Hadži D, Thompson HW (eds). 1959 *Hydrogen bonding*. Oxford, UK: Pergamon Press.
- Pauling L. 1960 *The nature of the chemical bond*, 3rd ed. Ithaca, NY: Cornell University Press.
- Grabowski SJ. 2011 What is the covalency of hydrogen bonding? *Chem. Rev.* **111**, 2597–2625. (doi:10.1021/cr800346f)
- Elgabarty H, Khaliullin RZ, Kühne TD. 2015 Covalency of hydrogen bonds in liquid water can be probed by proton nuclear magnetic resonance experiments. *Nat. Commun.* **6**, 8318. (doi:10.1038/ncomms9318)
- McKenzie RH. 2012 A diabatic state model for donor-hydrogen vibrational frequency shifts in hydrogen bonded complexes. *Chem. Phys. Lett.* **535**, 196–200. (doi:10.1016/j.cplett.2012.03.064)
- McKenzie RH, Bekker C, Athokpam B, Ramesh SG. 2014 Effect of quantum nuclear motion on hydrogen bonding. *J. Chem. Phys.* **140**, 174508. (doi:10.1063/1.4873352)
- Benoit M, Marx D, Parrinello M. 1998 Tunnelling and zero-point motion in high-pressure ice. *Nature* **392**, 258–261. (doi:10.1038/32609)
- Castro-Neto AH, Pujol P, Fradkin E. 2006 Ice: a strongly correlated proton system. *Phys. Rev. B* **74**, 024302. (doi:10.1103/PhysRevB.74.024302)
- Bernal JD, Fowler RH. 1933 A theory of water and ionic solution with particular reference to hydrogen and hydroxyl ions. *J. Chem. Phys.* **1**, 515–548. (doi:10.1063/1.1749327)
- Pauling L. 1935 The structure and entropy of ice and of other crystals with some randomness of atomic arrangement. *J. Am. Chem. Soc.* **57**, 2680–2684. (doi:10.1021/ja01315a102)
- Bove LE, Klotz S, Paciaroni A, Sacchetti F. 2009 Anomalous proton dynamics in ice at low temperatures. *Phys. Rev. Lett.* **103**, 165901. (doi:10.1103/PhysRevLett.103.165901)
- Meng X, Guo J, Peng J, Chen J, Wang Z, Shi J-R, Li X-Z, Wang E-G, Jiang Y. 2015 Direct visualization of concerted proton tunnelling in a water nanocluster. *Nat. Phys.* **11**, 235–239. (doi:10.1038/nphys3225)
- Yen F, Gao T. 2015 Dielectric anomaly in ice near 20 K: evidence of macroscopic quantum phenomena. *J. Phys. Chem. Lett.* **6**, 2822–2825. (doi:10.1021/acs.jpcllett.5b00797)
- Lin L, Morrone JA, Car R. 2011 Correlated tunneling in hydrogen bonds. *J. Stat. Phys.* **145**, 365–384. (doi:10.1007/s10955-011-0320-x)
- Drechsel-Grau C, Marx D. 2014 Quantum simulation of collective proton tunneling in hexagonal ice crystals. *Phys. Rev. Lett.* **112**, 148302. (doi:10.1103/PhysRevLett.112.148302)
- Benton O, Sikora O, Shannon N. 2016 Classical and quantum theories of proton disorder in hexagonal water ice. *Phys. Rev. B* **93**, 125143. (doi:10.1103/PhysRevB.93.125143)
- Baumgratz T, Cramer M, Plenio MB. 2014 Quantifying coherence. *Phys. Rev. Lett.* **113**, 140401. (doi:10.1103/PhysRevLett.113.140401)
- Bennett CH, DiVincenzo DP, Smolin J, Wootters WK. 1996 Mixed-state entanglement and quantum error correction. *Phys. Rev. A* **54**, 3824. (doi:10.1103/PhysRevA.54.3824)
- Coffman V, Kundu J, Wootters WK. 2000 Distributed entanglement. *Phys. Rev. A* **61**, 052306. (doi:10.1103/PhysRevA.61.052306)
- Henderson L, Vedral V. 2001 Classical, quantum and total correlations. *J. Phys. A: Math. Gen.* **34**, 6899–6905. (doi:10.1088/0305-4470/34/35/315)
- Ollivier H, Zurek WH. 2001 Quantum discord: a measure of the quantumness of correlations. *Phys. Rev. Lett.* **88**, 017901. (doi:10.1103/PhysRevLett.88.017901)

22. Dakić B, Vedral V, Brukner Č. 2010 Necessary and sufficient condition for nonzero quantum discord. *Phys. Rev. Lett.* **105**, 190502. (doi:10.1103/PhysRevLett.105.190502)
23. Pusuluk O, Farrow T, Deliduman C, Burnett K, Vedral V. 2018 Proton tunneling in hydrogen bonds and its implications in an induced-fit model of enzyme catalysis. *Proc. R. Soc. A* **474**, 20180037. (doi:10.1098/rspa.2018.0037)
24. Keutsch FN, Saykally RJ. 2001 Water clusters: untangling the mysteries of the liquid, one molecule at a time. *Proc. Natl Acad. Sci. USA* **98**, 10 533–10 540. (doi:10.1073/pnas.191266498)
25. Breuer HP, Petruccione F. 2002 *The theory of open quantum systems*. New York, NY: Oxford University Press. 130–137.
26. Wooldridge PJ, Devlin JP. 1988 Proton trapping and defect energetics in ice from FT-IR monitoring of photoinduced isotopic exchange of isolated D₂O. *J. Chem. Phys.* **88**, 3086–3091. (doi:10.1063/1.453952)
27. Suga H. 1997 A facet of ice sciences. *Thermochim. Acta* **300**, 117–126. (doi:10.1016/S0040-6031(96)03121-8)
28. Yen F, Chi ZH. 2015 Proton ordering dynamics of H₂O ice. *Phys. Chem. Chem. Phys.* **17**, 12 458–12 461. (doi:10.1039/C5CP01529D)
29. Tajima Y, Matsuo T, Suga H. 1982 Phase transition in KOH-doped hexagonal ice. *Nature* **299**, 810–812. (doi:10.1038/299810a0)
30. Tajima Y, Matsuo T, Suga H. 1984 Calorimetric study of phase transition in hexagonal ice doped with alkali hydroxides. *J. Phys. Chem. Solids* **45**, 1135–1144. (doi:10.1016/0022-3697(84)90008-8)