Published in final edited form as:

J Vasc Nurs. 2019 June ; 37(2): 78–85. doi:10.1016/j.jvn.2018.11.004.

Reduced length of stay and 30-day readmission rate on an inpatient vascular surgery service

Brittany O. Aicher, MD^a, Erin Hanlon, PA-C, MSPAS^b, Sarah Rosenberger, MSN, CRNP, CCNS^b, Shahab Toursavadkahi, MD^a, and Robert S. Crawford, MD^c

^aDivision of Vascular Surgery, Department of Surgery, University of Maryland School of Medicine, Baltimore, MD

^bDivision of Advanced Practice Providers, Department of Nursing, University of Maryland Medical Center, Baltimore, MD

^cCenter for Aortic Disease, University of Maryland Medical Center, Baltimore, MD

Abstract

Background: As the cost of care for patients with specific diagnoses becomes fixed, hospitals must develop systems to reduce length of stay and optimize the utilization of hospital resources while maintaining a high quality of care. The goal of this study is to evaluate the implementation and efficacy of a system designed to reduce average length of stay on a vascular surgery service.

Study Design: To effectively reduce our average length of stay we restructured patient rounds, implemented multi-disciplinary rounds, introduced clinical pathways to post-operative care, and expanded outpatient management of post-operative patients. 1697 adult vascular surgery patients discharged while under the medical direction of a vascular surgeon between 7/1/2013 and 6/30/2016 were included in the study.

Results: Improving communication with critical staff and utilizing procedural space outside of the main operating rooms led to a 2.8-day reduction in length of stay (10.8 vs 8.0, $P < 0.001$). There was a trend toward a reduction in 30-day readmission rate (12% vs. 10%, respectively; $P = 0.01$) and no significant difference in case-mix index as a measure of illness severity (2.5 vs. 2.4, respectively; $P = 0.15$). Length of stay reductions were heterogeneous among the types of vascular disease studied, with greater improvements seen in patients undergoing lower extremity amputation, lower extremity angiogram, and endovascular aneurysm repair for non-ruptured abdominal aortic aneurysm. Less pronounced differences were observed in patients undergoing carotid artery endarterectomy/stenting and lower extremity bypasses.

Conclusions: Restructuring team rounds and instituting a multidisciplinary approach to discharge planning produced significant reductions in length of stay without a deleterious effect on patient care that may impact hospital profitability.

Address Correspondence to: Brittany O. Aicher, MD, Department of Surgery, University of Maryland Medical Center, 22 S. Greene Street, S8B02, Baltimore, MD 21201, baicher@som.umaryland.edu, Phone: 715-212-7082, Fax: 410-328-0401.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

Length of stay; cost of hospitalization; multi-disciplinary rounds

Introduction

Hospitals across the United States have suffered financial difficulties, with over 50% reporting financial losses during the recession of 2008–2009, placing the ballooning cost of healthcare under increasing scrutiny [1]. Profits are further complicated by a national shift from pay per admission to pay for performance. In 1971, Maryland implemented a system for hospital reimbursement. Established by state law, the Maryland Health Services Cost Review Commission (MHSCRC) is an independent agency that sets reimbursement rates for all patients. Under this model, third-party payers are charged the same rate for the same care. The rates at individual hospitals are adjusted to account for each hospital's wages, charity care, severity of patient illness, volume, proportion of Maryland Residents seeking care at the institution, and known rate of uncompensated care. In 2012 changes were made to incentivize quality and efficiency and a financial penalty for all-cause readmissions within 30-days of discharge was imposed [2]. The MHSCRC is recognized as an alternative method to implement performance based health care reimbursement and Maryland continues to function with this financial model on a waiver status from the Affordable Care Act. As the cost of caring for a patient with a specific diagnosis becomes fixed, hospitals are developing systems to reduce the average length of stay (ALOS). Tertiary and quaternary centers that provide complex and emergent surgical care are anticipated to have the greatest impact from these systems, making this an important area of improvement at our center [3,4].

On surgical services, attempts to reduce ALOS commonly focus on maximizing preoperative, outpatient evaluation, introducing post-operative pathways, and educating healthcare providers on the importance of reducing ALOS [5]. Our patient population is comprised of a large volume of patients who present with acute illness from other hospitals and require emergent surgery, obviating pre-admission evaluation. Given this clinical setting, we believed the ALOS on our vascular surgery service could be reduced through other changes in practice, which are described here.

Methods

A hospital billing database including all admissions to the University of Maryland Medical Center (UMMC) was utilized to select adult vascular surgery patients discharged between July 1, 2013 and June 30, 2016 while under the medical direction of a vascular surgeon. The implementation of Epic electronic medical record occurred in November 2015. Therefore, to ensure accurate reporting, post-Epic data begins as of December 1, 2015 discharges. We identified 1723 patients who presented either electively or emergently with diagnoses of thoracic and abdominal aortic aneurysms, carotid artery disease, critical limb ischemia, visceral artery aneurysm, and mesenteric ischemia, among others. 26 patients with length of stay of 0 days were excluded. This study was approved by the Institutional Review Board at UMMC.

Intervention

UMMC is a 757-bed tertiary care center in Baltimore, Maryland. The Division of Vascular Surgery is comprised of twelve faculty attending surgeons, six rotating fellows, and two advanced practice providers (APP). General Surgery residents rotate on the service for 1–2 month blocks. Patients are primarily located on a medical-surgical telemetry unit with trained vascular nurses and a multidisciplinary team of a case manager, social worker, and allied health professionals. The interventions targeted improving communication between team members, introducing clinical pathways to post-operative care, optimizing the utilization of existing hospital resources, and expanding post-operative outpatient management.

To improve communication, the surgical and multidisciplinary patient rounds were restructured. At 6 AM daily, a vascular surgery fellow led surgical team rounds comprised of general surgery residents, a vascular surgery attending, and APP. Patients' daily plans were developed and tasks were assigned for team members to execute. Multidisciplinary rounds (MDR) with the APP, charge nurse, bedside nurses, case managers, social workers, and allied health professionals were restructured to occur at 8:30 AM. The APP acted as the liaison to communicate surgical team assessment from fellow led rounds, augment plans based on surgical and allied health driven data, and to delineate clear expectations for patients' care including diagnostic imaging, laboratory tests, surgical planning, and discharge preparation. On weekends, morning rounds included the charge nurse and discharge needs were discussed with the on-call case manager as needed. Discharge planning was discussed with patients and their families frequently to manage their expectations and prepare for outpatient recovery.

Expanding the management capabilities of the dedicated vascular surgery patient care unit further streamlined post-operative care. The medical-surgery unit at UMMC is comprised of 12 beds. Additional training was provided to the nurses to expand the patient population that could be cared for on the patient care unit. This included training in monitoring arterial lines, insulin infusions, and performing frequent neurovascular checks (up to every 2 hours). This streamlined patients' post-operative courses by directing patients to a unit with nursing staff trained in vascular medicine and surgery and invested in early initiation of discharge planning.

Clinical pathways are standardized, evidence-based, interdisciplinary care management plans that outline an appropriate sequence of clinical interventions, milestones, and expected outcomes for a comparable patient population. Clinical pathways aim to increase quality of care by promoting safety, elucidating expected post-operative patient progression, and reducing ALOS. Pathways were developed with a focus on multidisciplinary care and implemented to direct the inpatient hospital course.

Redistribution of existing resources enabled the hospital to meet the needs of the growing vascular division. UMMC has a dedicated endovascular catheterization lab that had been primarily devoted to cardiac procedures. The catheterization lab nursing and ancillary staff were trained and supplied with the equipment necessary to accommodate other endovascular procedures including diagnostic and therapeutic angiogram, thrombolysis, and inferior vena

cava filter placement. These procedures, once limited to the main operating rooms, were redirected to the catheterization lab, decreasing the load on the main operating rooms and increasing the availability of operating room time for complex endovascular and open surgical procedures.

In planning for reducing ALOS, we recognized there would be an abbreviated time for patients and their families to become proficient in self-care and for the surgical team to monitor for post-operative complications such as wound infection. Therefore, a multidisciplinary team of outpatient nurses and APP developed a multifaceted approach to address common issues that arise shortly after discharge. All patients discharged to home received a standardized telephone questionnaire within 72-hours. This survey was modified from previously published tools [6–8] to identify problems that may lead to readmission including questions regarding wound healing, signs or symptoms of infection, and general well-being (Figure 1). If a patient provided a response that raised concern for a post-hospital complication or readmission, a provider would determine if further intervention was warranted in an APP led clinic. An underutilized clinical space within the hospital was repurposed to accommodate these visits without overloading the surgeons' outpatient clinical space. Patients could be scheduled for a same-day visit to obtain labs and diagnostic imaging, evaluate wound care, and coordinate home health services.

Procedural Analysis

After excluding patients with length of stay of zero days and greater than 21 days, we grouped patients based on the International Classification of Diseases, Ninth Revisions, Clinical Modification (ICD-9-CM) and the ICD-10-PCS codes. The codes were grouped into 5 categories (carotid artery disease, aortic surgery, amputations, lower extremity open surgery, and angiogram, with aortic aneurysms differentiated as ruptured or non-ruptured; Table 1). Patients undergoing both lower extremity amputation and bypass were only included in the open lower extremity group. Patients undergoing aortic surgery in addition to another operation were only included in the aortic surgery group. Patients in the angiography group did not have other operations performed during the operation of record. 1,064 patients were included.

Economic Analysis

We obtained financial data for each patient's inpatient hospital course (direct variable expense, direct fixed expense, total cost of hospitalization). Indirect expenses, which include overhead costs incurred in maintaining infrastructure (*e.g.*, administrative costs) and environment (*e.g.*, electricity, water, maintenance, rent) were excluded as they are distributed across all departments and are not necessarily billed in proportion to a patient's use of services. Furthermore, these costs are fixed and are not likely to vary by the volume of vascular cases performed or the complexity of any single case making them less valuable contributors to the overall assessment of the profitability of a particular admission. Direct variable expenses are costs that directly pertain to the provision of patient care and vary with patient volume. These include salaries for nurses, technicians, allied health professionals, and other staff, prosthetic implants, pharmaceuticals, operating room costs, laboratory tests, and radiology, making them dependent upon ALOS and operative intervention provided.

Direct fixed expenses pertain to the provision of patient care and do not vary in relation to patient volume. Finally, costs accrued in the outpatient setting or in subsequent hospitalizations were not included in this analysis.

Statistical Analysis

Means and standard error of the mean were calculated and compared. Continuous variables were compared using two-tailed Student's t-test for normally distributed variables. Chi-squared test was used to compare readmission rates. Data are reported as mean $\pm$ standard error of the mean. Statistical significance was determined at a *P* value < 0.05 . Statistical analyses and figure generation were performed using GraphPad Prism® 5 (GraphPad Software, San Diego, California, USA).

Results

1723 patients were identified in our search, of which 1697 were included in our analysis (631 prior to intervention, 7/1/13–6/30/14; 1066 post-intervention, 7/1/14–6/30/16). There was no difference in patient volume between the two groups, as measured by the number of discharges per month (54 ± 8 vs 47 ± 7 , *P* = 0.19). Case-mix index (CMI) is a reflection of the acuity of the patient population reported during a specific time-frame. There was no difference in CMI between the study groups (2.5 ± 0.07 vs 2.37 ± 0.05 , *P* = 0.151; Figure 2). The ALOS for the studied population decreased significantly with implementation of the aforementioned interventions (10.8 ± 0.5 vs 8.0 ± 0.2 days, *P* < 0.001 ; Figure 2). This reduction remained significant when adjusted for CMI (4.7 ± 0.2 vs 3.8 ± 0.1 days, *P* < 0.001 ; Table 2). During the study period 187 patients were readmitted in less than or equal to 30-days with a trend toward reduction in this measurement after the intervention (81 pre-intervention vs. 106 postintervention, *P* = 0.010; Table 2).

The reduction in length of stay was not equally distributed between the surgical procedures reviewed. Patients undergoing carotid artery stent, carotid endarterectomy, lower extremity amputations, angiography, EVAR for non-ruptured AAA, or aorta-iliac-femoral bypass all experienced significant reductions in length of stay. Those undergoing open lower extremity operations such as bypass or endarterectomy, TEVAR, or aortic surgery for a ruptured aneurysm did not experience a reduction in length of stay (Table 3).

The mean cost of inpatient care prior to intervention was $\$32,896 \pm 1,413$ per patient, which was not significantly different after the intervention (*P* = 0.70; Table 2). The mean direct variable expense prior to intervention was $\$23,801 \pm 1,059$ per patient, which did not differ after our intervention (*P* = 0.81), and the mean direct fixed expense was $\$9,095 \pm 373$ per patient, which did not change (*P* = 0.43).

Discussion

This study found that implementing MDR, APP management of inpatients, post-surgical pathways, TSC, and utilizing procedural space outside of the main operating rooms significantly decreased our ALOS, but was not associated with a financial benefit. Causation with any intervention independent of the others cannot be determined, however. As we have

previously proposed, in Medical Centers such as ours, with a high preponderance of non-elective vascular surgery, specific management, training, and resource capabilities may further reduce ALOS and increase hospital profitability [9]. To our knowledge, no similar group of interventions has been documented in a similar vascular surgery population.

MDR is emerging as a tool to improve team communication, patient outcomes, and ALOS. This is well documented on trauma services where social and patient placement needs frequently delay discharge [10,11] and more recently in vascular surgery [12]. Others have assessed the impact of twice daily attending rounds and 24-hour chief resident availability to nursing, showing an improved sense of safety and team approachability without a difference in inpatient mortality, cardiac arrest, reoperation, or re-admission [13]. APP led rounds have emerged as another means to decrease ALOS and improve nurse and physician satisfaction [14]. Our experience is that APP are essential to the safe, efficient care of surgical inpatients and provide a continuity of care that is difficult to achieve with residents frequently changing services and having responsibilities in locations other than the patient care unit.

Clinical pathways are widely used in medicine to improve outcomes, decrease costs, and improve ALOS and have been reported in the vascular literature as far back as 1997 [15,16]. Post-operative pathways were implemented at our institution to improve team efficiency. Key to the success of these tools is the support of APP and nurses to promote adherence and recognize when deviation occurs. More recently, an urban tertiary care hospital published a comparison of acute aortic disease patient outcomes before and after implementation of clinical pathways, noting improved outcomes, decreased mortality, and decreased time to intervention [17]. Our experience was that some diagnosis groups had more predictable recovery and patients were less likely to deviate from the clinical pathway; these included patients undergoing the following procedures: carotid artery stent/endarterectomy, lower extremity amputations, angiography, EVAR for non-ruptured AAA, or aorta-iliac-femoral bypass. Lower extremity bypass and endarterectomy, TEVAR, and aortic surgery for ruptured aneurysm were more likely to have patient specific circumstances that influenced ALOS and less likely to follow a prescribed pathway of recovery.

Importantly, attempts to reduce ALOS must be balanced by interventions to ensure patients are not discharged only to be readmitted at a later date. As part of the MHSCRC, all cause readmissions to any Maryland hospital are centrally tracked and reported. Therefore, readmission penalties include not only those patients who are admitted back to the original facility but additionally those admitted to other Maryland hospitals. A tertiary center is challenged with a broad geography of patients served. Discharge plans must adapt to accommodate the available resources in the county in which the patient resides and address that patients may seek medical attention and guidance from other facilities if recovery expectations are not clear and early home interventions are not convenient. We believe readmissions were minimized thorough discharge planning coordinated by multiple team members early in each patient's hospitalization and through close follow-up in the form of telephone nurse assessment questionnaires to identify problems within 72-hours of discharge. This system addressed potential problems before a serious complication arose, identified complications early, and ensured patients had appropriate follow-up, lab work, and home health services. Vascular surgery patients are commonly discharged on new

medications that are associated with increased rates of readmission (*e.g.*, anticoagulants, antiplatelet agents, antibiotics) and have high rates of wound complications, that are commonly managed by negative pressure therapy or complex and frequent dressing changes [18]. An intervention that ensures patients are comfortable with their discharge care, do not have lingering questions, and can be seen if needed in a prompt fashion facilitated early discharge from the hospital.

One limitation of this study is that we did not attempt to quantify the effect of these interventions on the cost of outpatient care. Increased home health utilization and wound supplies likely impact the ambulatory cost both to insurance providers as well as out of pocket costs to patients. Another intervention that was not included but could have further aided our efforts would be to ensure regularly scheduled follow up appointments were made prior to patient discharge from the hospital. Unfortunately, we were unable to assess readmissions to other hospitals and were limited to readmissions to our hospital.

Our study period coincides with the introduction of the Critical Care Resuscitation Unit, a six-bed intensive care unit in the R Adams Cowley Shock Trauma Center with the goal of increasing adult critical care transfers to UMMC and improving outcomes. In the year after this unit opened, overall transfers increased 65%, and those of critically ill surgical patients increased by 94% [19]. Within our Division, specifically, there was nearly a four-fold increase in the number of patients transferred to the vascular surgery service. The number of aortic operations performed at our hospital more than doubled (92 vs. 219), and was distributed between EVAR, TEVAR, and open aortic aneurysm repair. The number of patients with the diagnosis of ruptured aortic aneurysm increased from 10 to 26. Increased critical care transfers and aortic operations infer a rise in the severity of illness treated. Despite this influx of patients with critical vascular issues, it is notable that there was no significant change in CMI during the study period. This has a significant impact on cost as the MHSCRC uses CMI to rate adjust reimbursement for specific diagnoses or reduce financial penalties associated with complications.

CMI was designed by the Centers for Medicare and Medicaid to calculate hospital payments as they relate to resource consumption. It is often used, as in this study, as a surrogate for illness severity. Therefore, CMI is higher for patients that utilize more resources, not necessarily patients with more severe disease. Furthermore, CMI is highly dependent upon provider documentation and coding accuracy and can be affected by the skillset of the coders or training providers and coders [20–23]. Of note, no project to improve the accuracy of coding was undertaken in the Division of Vascular Surgery during this time. Future projects in our Division may target accurately capturing comorbidities to improve coding.

Finally, despite a decrease in ALOS, there was no significant impact on cost of inpatient stay, which others have similarly reported [24]. In analyses of cost of stay per day, for most patients, the costs that are directly attributable to the last day of a hospital stay are an economically insignificant component of the total cost of stay, suggesting there is a limit to savings that can be achieved with ALOS reductions. Patients need to spend a reasonable amount of time hospitalized and with reductions in ALOS, costs and procedures that had been incurred at the end of a patient's stay are likely transferred to an earlier day in the

hospitalization. It is only possible to reduce hospital expenditures when a reduced volume of inpatient days allows cost-saving measures to be performed such as decreasing staffing levels by closing inpatient nursing units. These measures reduce overall fixed costs, resulting in measurable savings.

Associated with this, endovascular technologies broadened the options for treatment of many vascular lesions and have reduced ALOS as compared to traditional interventions [25–29]. In addition to shortened ALOS, endovascular intervention is credited with decreased perioperative morbidity and hospital acquired infections [25,26]. Despite these advantages, the necessary imaging, procedural, and supply costs of endovascular therapies far exceeds the cost of traditional open surgery [27,28]. Walker, *et al.* [29] proposed an alternative reimbursement model that stratifies vascular lesions using the TransAtlantic Intersociety Consensus II (TASC II) scoring system, imploring that the cost of treating lower extremity vascular disease be driven by the complexity of the lesion. This is not accounted for in a Diagnosis-Related Group reimbursement model and may further complicate pay-for-performance reimbursement.

A strength of this study is that it is an analysis of a unique state-regulated all-payer hospital reimbursement system. This reduces variance inherent when individual contracts with hospitals and third party payers determine payments levels, which is the case in many states. In those situations, the reimbursements for Medicare are typically lower than that for commercial payers relative to the costs of treatment, forcing hospitals to implement “cost shifting,” or increasing prices to private insurers. Furthermore, outside of this system, uninsured patients are usually charged high prices that no insurer would actually pay, making a hospital unable to collect payment. In a fixed reimbursement model such as ours, hospitals must charge the same rate regardless of insurance type or lack of insurance.

A limitation of our study is the use of direct costs for clinical services. This does not consider indirect costs. Our analysis is further limited to the acute inpatient period, excluding follow-up care and outpatient rehabilitation. Re-hospitalization outside of our center is also unknown. The data examined was obtained from an administrative database, which hinders our ability to comment on patient entry point to the hospital and adverse events such as death or unplanned surgical intervention. Finally, we did not examine overall revenue during the study period. Thus, the total cost to the health care system is understated, however our focus was on assessing ALOS and the associated financial implications during the acute period of care.

As a specialty, vascular surgery has evolved significantly in the past two decades. The advent of endovascular therapies changed the landscape of general vascular practice. These changes are similar to those experienced in general surgery with the arrival of laparoscopy. Both specialties have seen a segregation of specialists into those with primarily minimally invasive/outpatient practices and those with largely inpatient based patient populations, the latter generally based at large, tertiary care centers. In general surgery, this resulted in the birth of acute care surgery as a unique specialty. This work, and others by our group [9,30], highlight that a team dedicated to caring for patients with acute vascular diseases that demand intense resource utilization can optimize outcomes, achieve efficiency, and contain

costs. It is not unreasonable to think this model will continue to sprout at high volume/high acuity centers and evolve into its own academic practice and identity, mirroring what has occurred in general surgery.

Conclusions

Our findings support MDR, early discharge planning, and early identification of patients who will undergo procedures associated with significant lifestyle changes that complicate their discharge. This led to a 2.8-day reduction in ALOS and reduced 30-day readmission in an urban, tertiary teaching hospital. Reductions in ALOS were not equal across surgical procedures, suggesting the need for different strategies to improve the delivery of care in different patient populations. Key to the success of these interventions is educating the individuals involved in patient care to expected care pathway, recognizing when patients are not advancing appropriately, initiating discharge planning early, and maintaining close follow-up after discharge. We seek to provide an outline that both large, tertiary centers and community-based hospitals can use to provide high quality acute vascular surgical care with reduced variation in care, reduced length of stay, optimized outcomes, and minimized 30-day readmissions.

Acknowledgements

The authors thank Christina Varricchio, Senior Manager of Finance, Decision Support, University of Maryland Medical Center, for providing expertise in statistical analysis and hospital finances. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Abbreviations

ALOS	average length of stay
AOID	aortoiliac occlusive disease
APP	advanced practice provider
CMI	case-mix index
EVAR	endovascular repair of abdominal aortic aneurysm
LE	lower extremity
MDR	multidisciplinary rounds
MHSCRC	Maryland Health Services Cost Review Commission
TEVAR	thoracic endovascular aortic repair
TSC	transitional surgery clinic
UMMC	University of Maryland Medical Center

References

- [1]. Fox M. US hospital profits fall to zero: Thomson Reuters. Reuters 2009.

- [2]. Haber S, Beil H, Adamache W, et al. Evaluation of the Maryland All-Payer Model. Centers Medicare Medicaid Serv 2016.
- [3]. Cots F, Mercadé L, Castells X, Salvador X. Relationship between hospital structural level and length of stay outliers: Implications for hospital payment systems. *Health Policy (New York)* 2004;68:159–68.
- [4]. Mehra T, Muller CTB, Volbracht J, et al. Predictors of high profit and high deficit outliers under SwissDRG of a tertiary care center. *PLoS One* 2015;10:1–18.
- [5]. Muluk SC, Painter L, Sile S, et al. Utility of clinical pathway and prospective case management to achieve cost and hospital stay reduction for aortic aneurysm surgery at a tertiary care hospital. *J Vasc Surg* 1997;25:84–93. [PubMed: 9013911]
- [6]. Inman DM, Maxson PM, Johnson KM, et al. Impact of follow-up educational telephone calls on patients after radical prostatectomy: Finding value in low-margin activities. *Urol Nurs* 2011;31:83–91.
- [7]. Donaho EK, Hall AC, Gass JA, et al. Protocol-driven allied health post-discharge transition clinic to reduce hospital readmissions in heart failure. *J Am Heart Assoc* 2015;4:1–11.
- [8]. Brooke BS, De Martino RR, Girotti M, et al. Developing strategies for predicting and preventing readmissions in vascular surgery. *J Vasc Surg* 2012;56:556–562. [PubMed: 22743022]
- [9]. Harris DG, Herrera AV, Drucker CB, et al. Defining the burden, scope, and future of vascular acute care surgery. *J Vasc Surg* 2017;66:1511–17. [PubMed: 28662926]
- [10]. Sen A, Xiao Y, Lee SA, et al. Daily multidisciplinary discharge rounds in a trauma center: A little time, well spent. *J Trauma - Inj Infect Crit Care* 2009;66:880–887.
- [11]. Dutton RP, Cooper C, Jones A, et al. Daily multidisciplinary rounds shorten length of stay for trauma patients. *J Trauma* 2003;55:913–919. [PubMed: 14608165]
- [12]. Houghton JSM, Urriza Rodriguez D, Weale AR, et al. Delayed discharges at a major arterial centre: A 4-month cross-sectional study at a single specialist vascular surgery ward. *BMJ Open* 2016;6:e011193.
- [13]. Johnston MJ, Arora S, King D, Darzi A. Improving the quality of ward-based surgical care with a human factors intervention bundle. *Ann Surg* 2016;267:73–80.
- [14]. Mower-Wade D, Pirrung JM. Advanced practice nurses making a difference: Implementation of a formal rounding process. *J Trauma Nurs* 2010;17:69–71; quiz 72–3. [PubMed: 20559052]
- [15]. Every NR, Hochman J, Becker R, et al. Critical pathways : a review. Committee on Acute Cardiac Care, Council on Clinical Cardiology, American Heart Association. *Circulation* 2000;101:461–5. [PubMed: 10653841]
- [16]. Patterson RB, Whitley D, Porter K. Critical pathways and cost-effective practice. *Semin Vasc Surg* 1997;10:113–8. [PubMed: 9203263]
- [17]. Shin KC, Lee HS, Park JM, et al. Outcomes before and after the Implementation of a Critical Pathway for Patients with Acute Aortic Disease. *Yonsei Med J* 2016;57:626. [PubMed: 26996561]
- [18]. Zdyb E, Courtney D, Malik S, et al. Impact of discharge anticoagulation education by Emergency Department pharmacists at a Tertiary Academic Medical Center. *J Emerg Med* 2017;53:896–903. [PubMed: 28941557]
- [19]. Scalea TM, Rubinson L, Tran Q, et al. Critical Care Resuscitation Unit: An Innovative Solution to Expedite Transfer of Patients with Time-Sensitive Critical Illness. *J Am Coll Surg* 2015;222:614–21.
- [20]. Grogan EL, Speroff T, Deppen SA, et al. Improving documentation of patient acuity level using a progress note template. *J Am Coll Surg* 2004;199:468–75. [PubMed: 15325618]
- [21]. Rosenstein AH, O'Daniel M, White S, Taylor K. Medicare's Value-Based Payment Initiatives: Impact on and Implications for Improving Physician Documentation and Coding. *Am J Med Qual* 2009;24:250–258. [PubMed: 19461068]
- [22]. Courtright E, Diener I, Russo R. Clinical documentation improvement (CDI). Each word carries its own meaning for health care providers. *Case Manager* 2004;15:46–9.
- [23]. Zalatio O, Ranasinghe M, Harbaugh RE, Iantosca M. Impact of improved documentation on an academic neurosurgical practice. *J Neurosurg* 2014;120:756–63. [PubMed: 24359011]

- [24]. Taheri P a, Butz D a, Greenfield LJ. Length of stay has minimal impact on the cost of hospital admission. *J Am Coll Surg* 2000;191:123–130. [PubMed: 10945354]
- [25]. Orr NT, Davenport DL, Minion DJ, Xenos ES. Comparison of perioperative outcomes in endovascular versus open repair for juxtarenal and pararenal aortic aneurysms: A propensitymatched analysis. *Vascular* 2017;25:339–45. [PubMed: 27903931]
- [26]. Hajibandeh SS, Hajibandeh SS, Antoniou SA, et al. Percutaneous access for endovascular aortic aneurysm repair: A systematic review and meta-analysis. *Vascular* 2016;24:638–648. [PubMed: 27000385]
- [27]. Forbes TL, DeRose G, Kribs S, Harris KA. A cost-effectiveness analysis of standard versus endovascular abdominal aortic aneurysm repair. *Can J Surg* 2002;45:420–4. [PubMed: 12500916]
- [28]. IMPROVE Trial Investigators, Powell JT, Sweeting MJ, et al. Endovascular or open repair strategy for ruptured abdominal aortic aneurysm: 30 day outcomes from IMPROVE randomised trial. *BMJ* 2014;348:f7661. [PubMed: 24418950]
- [29]. Walker KL, Nolan BW, Columbo JA, et al. Lesion complexity drives the cost of superficial femoral artery endovascular interventions. *J Vasc Surg* 2015;62:998–1002. [PubMed: 26206581]
- [30]. Harris D, Olson S, Kalsi R, et al. Early treatment at a referral center improves outcomes for patients with acute vascular disease. *Ann Vasc Surg* 2017;In press:e27–8.

Highlights:

- Reductions in average length of stay were made by restructuring patient rounds, implementing multi-disciplinary rounds, introducing clinical pathways to post-operative care, and expanding outpatient management of post-operative patients.
- This multidisciplinary approach to discharge planning resulted in a significant decrease in length of stay, with important economic implications to the hospital without a deleterious effect on patient care

Figure 1:
Transitional Surgery Clinic (TSC) workflow. APP- advanced practice provider.

Figure 2:

Vascular Surgery average length of stay (ALOS), measured in days, and case-mix index (CMI) are based on: A) inpatients discharged within the time frame as laid out, and B) inpatients with a vascular surgeon as their Physician of Record at time of discharge. CMI is a reflection of the acuity of the patient population reported during a specific time-frame.

Table 1:

Patient procedure categories.

Diagnosis	Patients (n)	ICD-9-CM Codes	ICD-10-PCS Codes
Carotid artery disease	113	38.12, 39.7, 0.63	03[7,C,H],[H,J,K,L,M,N]-
Aortic surgery	311	39.71, 39.73, 38.44, 39.52, 38.45, 39.25	04[1,7,B,C,H,L,P,Q,R,S,U,V,W],[0,1,2,3,4,5,9,A,C,D]-, 0470[0,3,4]-, 04[L,R,U,V]0-, 02[U,V]W-
Ruptured aortic aneurysm	36	441.1, 441.3, 441.5, 441.6	I71.1, I71.3, I71.5, I71.8
Lower extremity amputation	112	84.11, 84.12, 84.13, 84.14, 84.15, 84.16, 84.17, 84.3	0Y67-, 0Y68-, 0Y6C-, 0Y6D-, 0Y6F-, 0Y6G-, 0Y6H-, 0Y6J-, 0Y6M-, 0Y6N-, 0Y6P-, 0Y6Q-, 0Y6R-, 0Y6S-, 0Y6T-, 0Y6U-, 0Y6V-, 0Y6W-, 0Y6X-, 0Y6Y-
Lower extremity open surgery	151	33.14, 38.18, 38.08, 38.48, 38.68	The following procedures were separated by open vs. percutaneous or percutaneous endoscopic:
Angiogram	341	39.5, 39.79, 39.9, 0.4, 0.41, 0.43, 0.45, 0.46, 0.47, 0.48, 0.49, 0.55, 0.6, 38.7, 88.4/41/42/43/44/45/47/48/49/51, 99.1	04,[1,7,B,C,H,L,P,Q,R,S,U,V,W], [E,F,H,J,K,L,M,N,P,Q,R,S,T,U,Y]-, 031[5,6]0[9,A,J,K,Z] [6,7,8,9,B,C],02C[Q,R,-] B31[0,1,2,3,4,5,6,7,8,6,B,C,D,F,G,H,J,K,L,N,P,Q,S,T,U]-, B41 [0,2,3,4,5,6,7,8,9,B,C,D,F,G,J]-, 3E0[4,6][0,3]17

Table 2:

Length of stay analysis.

	Control (n=631)	Intervention (n=1066)	<i>P value</i>
Patient Volume, no. of discharges/month	54 ± 8	47 ± 7	0.19
Case-mix Index, mean ± SEM	2.5 ± 0.07	2.37 ± 0.05	0.15
Length of Stay, mean ± SEM, days	10.8 ± 0.5	8.0 ± 0.2	<.001
Case-mix Index Adjusted Length of Stay, mean ± SEM, days	4.7 ± 0.2	3.8 ± 0.1	<.001
30-day Readmission, no., %	81, 12	106, 10	<i>0.01</i>
Cost of Inpatient Care (direct variable + direct fixed costs), USD	32,896 ± 1,413	31,974 ± 1,020	0.59
Direct Fixed Expense, USD	9,095 ± 373	8,674 ± 260	0.34
Direct Variable Expense, USD	23,801 ± 1,059	23,300 ± 784	0.7

SEM- standard error of the mean, USD- US dollars (\$).

Table 3:

Average length of stay for commonly performed procedures. All data reported as: Mean (Standard Error of the Mean). * $P < 0.1$; ** $P < 0.05$.

	Carotid Disease		Lower Extremity Open Procedure		Lower Extremity Amputations		Angiogram	
	Pre (n=40)	Post (n=86)	Pre (n=39)	Post (n=112)	Pre (n=44)	Post (n=68)	Pre (n=128)	Post (n=185)
Age (years)	67.0 (1.5)	66.7 (1.1)	66.7 (2.3)	61.3 (1.4)*	57.6 (1.9)	62.7 (1.7)*	59.9 (1.3)	59.0 (1.0)
CMI	1.58 (0.11)	1.35 (0.05)	3.02 (0.20)	2.68 (0.09)*	2.57 (0.16)	2.19 (0.14)*	2.10 (0.10)	2.11 (0.08)
ALOS (days)	3.9 (0.6)	2.6 (0.3)**	8.5 (0.8)	7.8 (0.4)	11.2 (0.7)	8.1 (0.5)**	8.1 (0.4)	5.9 (0.3)**
CMI adjusted ALOS (days)	2.5 (0.3)	1.9 (0.2)*	3.0 (0.3)	3.0 (0.2)	4.62 (0.25)	4.12 (0.24)	4.6 (0.3)	3.4 (0.2)**
Direct Variable Cost (USD)	8,587 (675)	9,953 (550)	20,928 (1,957)	23,807 (1,430)	16,164 (1,415)	14,153 (981)	19,220 (982)	17,409 (796)
Total Cost (USD)	21,243 (1,503)	22,772 (1,236)	42,378 (3,701)	47,353 (2,496)	34,365 (2,806)	30,081 (1,971)	36,451 (1,811)	32,653 (1,334)*

	All Aortic Surgery		EVAR, non-ruptured		TEVAR, non-ruptured		Aortoiliac Occlusive Disease	
	Pre (n=92)	Post (n=219)	Pre (n=20)	Post (n=59)	Pre (n=17)	Post (n=57)	Pre (n=11)	Post (n=26)
Age (years)	66.8 (1.4)	67.6 (0.9)	73.5 (1.6)	72.5 (1.5)	65.6 (2.6)	67.7 (1.8)	52.3 (4.2)	60.7 (2.2)*
CMI	3.36 (0.17)	3.11 (0.09)	3.02 (0.39)	2.39 (0.09)**	3.39 (0.31)	3.06 (0.12)	3.15 (0.37)	3.47 (0.28)
ALOS (days)	9.4 (0.5)	7.7 (0.3)**	7.1 (1.0)	4.8 (0.6)**	9.1 (4.4)	8.2 (0.5)	12.3 (1.1)	9.1 (0.8)**
CMI adjusted ALOS (days)	3.14 (0.22)	2.61 (0.11)**	2.69 (0.38)	1.96 (0.21)**	2.79 (0.24)	2.76 (0.18)	4.23 (0.45)	2.87 (0.28)**
Direct Variable Cost (USD)	36,806 (2,120)	36,072 (1,490)	38,954 (3,267)	34,670 (2,605)	47,176 (6,395)	50,561 (3,150)	25,651 (3,723)	23,950 (1,787)
Total Cost (USD)	63,389 (3,169)	62,580 (2,293)	62,335 (5,576)	51,222 (3,683) *	76,804 (9,156)	82,882 (4,850)	53,585 (6,488)	49,105 (15,724)

CMI- case-mix index, ALOS- average length of stay, USD- US dollars (\$), EVAR- endovascular repair of abdominal aortic aneurysm, TEVAR- thoracic endovascular aortic repair