Published in final edited form as:

Psychiatry Res Neuroimaging. 2019 June 30; 288: 51–59. doi:10.1016/j.psychres.2019.04.011.

Anterior cingulate volume reductions in abstinent adolescent and young adult cannabis users: Association with affective processing deficits

Kristin E. Maple, Alicia M. Thomas, Megan M. Kangiser, and Krista M. Lisdahl*

Department of Psychology, University of Wisconsin—Milwaukee, 2441 E. Hartford Ave., Milwaukee, WI, USA.

1. INTRODUCTION

Cannabis use is increasing amongst young adults, with 20% using within the past month (Johnston et al., 2016). Adolescents and emerging adults may be especially vulnerable to the negative effects of cannabis use due to the substantial neurodevelopment occurring during this period (Jacobus and Tapert, 2014; Lisdahl et al., 2013, 2014). Cannabis use has a high rate of comorbidity with affective disorders that often emerge during adolescence and young adulthood (Agrawal et al., 2011; Dorard et al., 2008; Wittchen et al., 2007). A possible mechanism underlying psychiatric comorbidities and mood symptoms in young cannabis users is abnormalities in the affective neural network, resulting in deficits in affective processing. One commonality of many disorders (including anxiety, depression, bipolar disorder, and psychosis) comorbid with cannabis use is a deficit in facial emotion processing (Bourke et al., 2010; Mogg et al., 2000; Morris et al., 2009; Samane, 2013), which is a critical aspect of functioning in human social networks (Phillips et al., 2003).

Cannabis use has been associated with abnormal facial emotion processing. Cannabis users are less accurate in recognizing and discriminating between emotions (Bayrakci et al., 2015; Hindocha et al., 2014; Huijbregts et al., 2014). An additional consideration is that cannabis users' accuracy in identifying emotions depends upon the intensity at which the emotion is displayed. Yet, studies probing the importance of emotion intensity have yielded mixed findings. One study concluded that cannabis users had more difficulty identifying more subtle emotions (Platt et al., 2010), while another reported that cannabis users were less accurate than controls only in recognizing more overt, unambiguous emotions (Hindocha et

*Corresponding Author: krista.medina@gmail.com; Phone: (414) 229-7159.

CONTRIBUTORS

KEM was involved in data collection, developing the study concept, data processing, performing data analyses, interpreting the results, and writing the manuscript under KML's supervision. KML designed the study's methods and worked closely with KEM to develop the study concept, perform data analyses, interpret the results, and draft the manuscript. AMT was involved in data collection and data processing. MMK assisted in data collection, data processing, and preparation of the final manuscript.

CONFLICTS OF INTEREST

None

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

al., 2014). Another study found emotion recognition deficits only in more frequent and recent cannabis users (Huijbregts et al., 2014). Functional magnetic resonance imaging (MRI) studies have extended this behavioral work. During facial emotion tasks, cannabis use has been associated with abnormal anterior cingulate cortex (ACC) and amygdala activation (Cornelius et al., 2010; Gruber et al., 2009; Spechler et al., 2015).

Facial processing is complex, recruiting a distributed network of visual, limbic, temporoparietal, and prefrontal areas, as well as the cerebellum (Fusar-Poli et al., 2009b; Haxby et al., 2000; Ishai et al., 2004, 2005; Kesler-West et al., 2001; Said et al., 2011). Haxby and colleagues' (2000) model proposes a core system of face perception, consisting of the fusiform gyrus and superior temporal sulcus. In addition to this core system of facial processing, an extended system consists of a distributed network of regions that are recruited to derive meaning from facial expressions, including facial emotion (Haxby et al., 2000). This extended system includes areas such as the medial prefrontal cortex (PFC) (Fusar-Poli et al., 2009b; Kesler-West et al., 2001), orbital frontal cortex (OFC) (Ishai et al., 2005), inferior frontal gyrus (Ishai et al., 2005; Kesler-West et al., 2001), ACC (Wu et al., 2016), amygdala (Fusar-Poli et al., 2009b; Gur et al., 2002; Morris et al., 1996), hippocampus (Gur et al., 2002; Ishai et al., 2005), and cerebellum (Fusar-Poli et al., 2009b). During processing of emotional versus neutral unfamiliar faces, the entire face perception network is recruited more intensely (Ishai et al., 2004, 2005; Pessoa et al., 2002). Thus, all regions in the network are considered important in facial emotion recognition.

Cannabis use has been associated with structural differences in regions associated with emotional face processing. Cannabis users have exhibited structural abnormalities in volume and thickness within cortical regions including the frontal cortex (medial OFC (mOFC), lateral OFC (lOFC), ACC), temporal lobe, and fusiform gyrus (Battistella et al., 2014; Cheetham et al., 2012; Churchwell et al., 2010; Chye et al., 2017; Epstein and Kumra, 2015; Jacobus et al., 2014; Jarvis et al., 2008; Koenders et al., 2016; Lisdahl et al., 2016; Lopez-Larson et al., 2011; Mashhoon et al., 2015; Mata et al., 2010; Medina et al., 2009; Price et al., 2015; Thames et al., 2017). Cannabis use has also been linked with volume differences in subcortical regions such as the amygdala, hippocampus, and cerebellum (Ashtari et al., 2011; Battistella et al., 2014; Cousijn et al., 2012; Demirakca et al., 2011; Gilman et al., 2014; Jarvis et al., 2008; Koenders et al., 2016; Matochik et al., 2005; McQueeney et al., 2011; Medina et al., 2010; Medina et al., 2007; Nurmedov et al., 2015; Pagliaccio et al., 2015; Schacht et al., 2012; Solowij et al., 2013; Yucel et al., 2008). Typically, cannabis users, compared to non-users, have smaller volumes in these brain regions, though some studies in younger adolescent samples have found larger volumes in cannabis users (e.g., McQueeney et al., 2011; Medina et al., 2009, 2010). Despite the aforementioned literature, some studies have reported null findings for each of these brain regions (e.g., Tzilos et al., 2005; Weiland et al., 2015), suggesting a greater need to understand individual differences, such as gender, moderating the relationship between cannabis use and brain structure.

Males and females recruit different networks while processing emotional faces (Filkowski et al., 2017; Kempton et al., 2009; Whittle et al., 2011). Further, female affective circuitry may be more susceptible to the negative effects of cannabis use (for review, see Crane et al., 2013). Within adolescent female compared to male rats, increased cannabinoid receptor-1

(CB1) downregulation is observed in frontolimbic regions (Rubino et al., 2008). Additionally, relative to males, female rats exposed to tetrahydrocannabinol (THC), the main psychoactive component in cannabis, are more vulnerable to its anxiety and depression producing properties (Rubino et al., 2008). In humans, female cannabis users have exhibited functional disadvantages associated with differences in PFC, IOFC, and amygdala volumes compared to their male counterparts (Chye et al., 2017; McQueeney et al., 2011; Medina et al., 2009). Thus, cannabis-using females may process facial emotions differently from male users, and this may be related to underlying brain structure in frontolimbic areas.

Yet, to date, no study has investigated the relationship between brain structure, gender, and facial affective processing in abstinent cannabis users. The current study is novel in its focus on cannabis use and brain structure in facial emotion processing regions. More specifically, it is unclear in the existing literature if abnormal brain structure in cannabis users has functional consequences related to facial emotion processing. To address this gap in the literature, the primary aim of the current study was to examine the relationship between cannabis use and brain volumes in regions underlying facial emotion processing. We also investigated whether gender moderated this relationship. We hypothesized that cannabis use would be associated with reduced volumes, and that females would exhibit the strongest relationship between cannabis use and reduced volumes. For our secondary aim, we assessed whether differences in brain volumes associated with cannabis use and cannabis x gender interaction were related to deficits in facial emotion processing. We expected that brain volume would be related to poorer facial emotion processing within the cannabis-using group.

2. METHODS

2.1 Participants

Participants included 20 cannabis using and 35 non-using individuals (28 males and 27 females), ages 16–26 years, from a larger neuroimaging study (PI: Lisdahl, NIH R01 DA030354). Individuals were recruited through flyers posted in the community. In the current analysis, cannabis users were required to have at least 40 uses in the past year and at least 50 lifetime uses. Non-users were required to have fewer than or equal to 5 lifetime uses. Individuals were excluded for left-handedness, MRI contraindications, birth complications, birth <33 weeks gestation, neurological disorders, head trauma with >2 minute loss of consciousness, learning or intellectual disability, vision or hearing impairments, major health problems, independent DSM-IV-TR psychological disorder, current use of psychotropic medication, use of more than 10 cigarettes per day, heavy other drug use (>25 lifetime uses of non-cannabis drugs), pregnancy, and failure to demonstrate abstinence from all substances on the day of MRI scanning (as indicated by blood alcohol concentration of >.000, positive urine toxicology, and/or continuous sweat patch testing).

2.2 Design and procedure

The Institutional Review Boards at the University of Wisconsin-Milwaukee and the Medical College of Wisconsin (MCW) approved all aspects of this study. Interested individuals from the community underwent phone screening to determine eligibility, during which

comprehensive substance use history (lifetime) was obtained with the Customary Drinking and Drug Use Record (CDDR) (Brown et al., 1998; Stewart and Brown, 1995). Information regarding youth psychiatric history was gathered via a Diagnostic and Statistical Manual of Mental Disorders—Fourth Edition (DSM-IV) semi-structured interview (Sheehan et al., 1998) administered to both participants and parents.

Informed consent was obtained from eligible participants prior to the study. All participants (cannabis users and non-users) underwent three weeks of monitored abstinence prior to completing the affective processing tasks (Session #4) and MRI scan (Session #5), which occurred 24–48 hours apart. At weekly sessions (Sessions #1, 2, 3) during the three weeks prior to the affective processing tasks and MRI, participant abstinence was evaluated using urine toxicology (One Step Drug Screen Test Dip Card Panel; Innovacon, Inc., San Diego, CA) and continuous sweat toxicology (PharmChek Drugs of Abuse Patch; PharmChem Inc., Fort Worth, TX), which measured levels of THC and other substances. In addition, at each session, participants underwent breathalyzer testing (Alco-Sensor IV; Intoximeters, Inc., St. Louis, MO) for recent alcohol use.

2.3 Measures

2.3.1 Demographic information—Participants completed a background questionnaire outlining demographic variables including age, gender, ethnicity, and education.

2.3.2 Substance use—During each session, participants reported the last instance of any substance use. In addition, participants completed the Timeline Follow Back (TLFB; Sobell and Sobell, 1992). This is a highly reliable and valid measure of past year substance use, using holidays and other memory cues. As part of completing the TLFB, participants were asked to report the quantity of specific substances used on each day during the past year. Substance use was measured in standard units (e.g., joints for cannabis). Lifetime and past 3-month substance use was measured using the CDDR.

2.3.3 Affective processing—During Session #4, participants completed an affective battery (PennCNP: <https://pennncnp.med.upenn.edu/about.html>) containing three tasks of interest to the current analysis: Emotion Recognition, Emotion Discrimination, and Emotional Acuity tasks. In Emotion Recognition, participants were presented with a series of faces and instructed to identify which emotion each face was expressing. Possible answer choices included happy, sad, anger, fear, and no emotion. The facial stimuli are described by Gur et al. (2002) and Pinkham et al. (2008). In the Emotion Discrimination task, participants were presented with one pair of faces for each trial. Each pair of faces contained two pictures of the same individual, with or without a subtle (computer-generated) change in facial expression, which possibly indicated a difference in intensity between the two facial emotions. During each trial, the participant decided which picture expressed the specified emotion (happy or sad) more intensely, or whether they were equal. In the Emotional Acuity task, individual faces were randomly presented, and participants were instructed to rate the facial expression's emotional valence on a 7-point scale: very sad, moderately sad, somewhat sad, neutral, somewhat happy, moderately happy, or very happy. Facial stimuli for

the Emotion Discrimination and Emotional Acuity tasks are described by Erwin et al. (1992).

2.4 MRI data acquisition and pre-processing

2.4.1 MRI acquisition—Participants were scanned on a 3T scanner (GE Healthcare, Waukesha, WI) at MCW. Structural image acquisition took approximately 15 minutes. A T1-weighted, 3-D anatomical brain scan was obtained with spoiled gradient-recalled at steady-state (SPGR) pulse sequence (TR=8.2 ms, TE= 3.4 s., TI= 450 and flip angle of 12°). The in-plane resolution of the anatomical images was 256×256 with a square field of view (FOV) of 240 mm. Slice thickness was 1 mm., with 150 slices acquired.

2.4.2 MRI pre-processing—Structural images were processed using FreeSurfer 5.3's semi-automated processing stream (<http://surfer.nmr.mgh.harvard.edu>). All T1-weighted 3D anatomical datasets underwent motion correction, non-parametric non-uniform intensity normalization, Montreal Neurologic Institute transformation, removal of non-brain materials, and skull-stripping. This was followed by whole-brain segmentation of white and gray matter and calculation of cortical and subcortical volumes (Fischl et al., 2002, 2004), using the Desikan-Killiany atlas (Desikan et al., 2006) in FreeSurfer. All automated FreeSurfer steps were inspected for processing errors, and manual edits were made as needed. For each case, automatic segmentation and parcellation masks were manually edited for accuracy, using multiple views for visual inspection. While FreeSurfer's atlases include several regions, only volumes of select cortical and subcortical regions known to be associated with facial emotion processing were included in the current analysis.

2.5 Data analysis

All analyses were conducted using Statistical Package for the Social Sciences (SPSS). Demographic and other drug use was examined using Analysis of Variance (ANOVA) and Chi-Square analyses to detect differences between groups. Variables that either differed between groups or have known relationships with affective processing or brain structure were statistically controlled for in the regressions. Covariates included gender, past year alcohol use, past year nicotine use, and intracranial volume (ICV).

2.5.1 Principal components analysis (PCA) for emotional processing tasks—

Due to the mixed findings regarding cannabis use and facial emotion processing, many variables were of potential interest to the current study. In order to minimize the number of individual variables investigated, and therefore the number of tests run, we conducted principal components analyses (PCA) to identify the most salient constructs for investigation. We conducted three PCAs to extract components from each of the Emotion Recognition, Emotion Acuity, and Emotion Discrimination tasks. Accuracy and response time variables were selected for each task and entered into PCA (Tables 1, 2, 3).

2.5.2 Primary analysis—Across all 55 cannabis users and non-users, standard entry regression with two blocks (including appropriate covariates) was used to determine the relationship between past year cannabis use, cannabis x gender interaction, and brain structure. Dependent variables included bilateral mOFC, IOFC, Pars triangularis (subregion

of inferior frontal gyrus), superior temporal, fusiform, rostral ACC (rACC), caudal ACC (cACC), amygdala, hippocampus, and cerebellum volumes. Main effects and covariates were entered into the first block of the regressions; interaction variables were also included in the second block. In order to assess if any participants disproportionately influenced significant regression models, DFBETA analyses were conducted for past year cannabis use and cannabis $\times$ gender interactions. DFBETA analyses determined that no outliers were present. False Discovery Rate (FDR) corrections (Benjamini and Hochberg, 1995 method) examining the relationships of cannabis and cannabis $\times$ gender interactions with brain volume variables were run separately for subcortical and cortical variables within each brain hemisphere (R/L). Additionally, f^2 was used to assess effect sizes for multiple regression analyses (small= 0.02–0.14, medium= 0.15–0.34, and large= >0.35) (Cohen et al., 1988).

2.5.3 Secondary analysis: brain-behavior relationships—We conducted brain-behavior analyses on brain regions that were significantly associated with past year cannabis use or cannabis $\times$ gender interactions. For significant regions, we ran Pearson correlations to determine whether brain abnormalities were related to emotional processing accuracy, within the cannabis-using group. For all analyses, significance was determined if $p < 0.05$. DFBETA analyses determined that no participants disproportionately influenced significant correlation models. FDR corrections were run separately for each of the three affective processing tasks, for each significant brain region.

3. RESULTS

3.1 Demographic & drug use information (Table 4)

3.1.1 Cannabis—ANOVAs and chi-square tests revealed no significant difference between cannabis users and non-users in age ($F(1,53) = 2.25, p = 0.14$), race ($\chi^2(5) = 5.80, p = 0.33$), ethnicity ($\chi^2(2) = 1.26, p = 0.53$), gender ($\chi^2(1) = 1.04, p = 0.31$), education ($F(1,53) = 0.20, p = 0.65$), and reading ability ($F(1,53) = 0.67, p = 0.42$). Cannabis users and non-users significantly differed on measures of lifetime cannabis use ($F(1,53) = 26.97, p = 0.000003$), number of days per month of cannabis use in the past three months ($F(1,53) = 120.71, p < 0.000001$), past year cannabis use ($F(1,53) = 41.05, p < 0.000001$), past year alcohol use ($F(1,53) = 26.29, p = 0.000004$), and past year nicotine use ($F(1,53) = 8.46, p = 0.005$).

3.1.2 Gender—ANOVAs and chi-square tests revealed no significant difference between males and females in age ($F(1,53) = 0.06, p = 0.80$), race ($\chi^2(5) = 8.74, p = 0.12$), ethnicity ($\chi^2(2) = 3.77, p = 0.15$), education ($F(1,53) = 0.03, p = 0.86$), reading ability ($F(1,53) = 2.0, p = 0.16$), age of onset of weekly cannabis use ($F(1,18) = 0.40, p = 0.85$), length of cannabis abstinence ($F(1,20) = 0.69, p = 0.42$), lifetime cannabis use ($F(1,53) = 1.81, p = 0.19$), number of days per month of cannabis use in the past three months ($F(1,53) = 1.21, p = 0.28$), past year cannabis use ($F(1,53) = 0.97, p = 0.33$), past year alcohol use, ($F(1,53) = 2.46, p = 0.12$), and past year nicotine use ($F(1,53) = 2.60, p = 0.11$).

3.2 Principal components analysis

Three PCAs with varimax rotation were conducted to reduce variables from each of the tasks (Emotion Recognition, Acuity, and Discrimination) in the PennCNP affective battery. Variables with loadings ≥ 0.6 were considered to define a component.

3.2.1 Emotion recognition—For the Emotion Recognition task, nine variables yielded four components meeting Kaiser's criterion (eigenvalues >1) (Table 1): Recognition Time (33.42% of variance), Sad Correct (15.48% of variance), Happy Correct (13.03% of variance), and Anger Correct (12.10% of variance).

3.2.2 Emotional acuity—For the Emotional Acuity task, twelve variables yielded three components meeting Kaiser's criterion. Components included Acuity Time (37.04% of variance), Acuity Neutral (26.61% of variance), and Acuity Intensity (11.27% of variance) (Table 2).

3.2.3 Emotion discrimination—For the Emotion Discrimination task, eight variables yielded two components meeting Kaiser's criterion (Table 3): Discrimination Time (48.28% of variance) and Discrimination Correct (33.54% of variance).

3.3 Primary results

Greater past year cannabis use was significantly related to smaller left rACC volumes [$t(48) = -3.09$, $\beta = -0.33$, $p = 0.003$, $r^2 = 0.14$; FDR corrected: $p = 0.02$] (Figures 1, 2). Gender did not moderate this relationship. Within the left rACC regression model, the only covariate related to left rACC volumes was intracranial volume, which was positively associated with left rACC volume [$t(48) = 5.16$, $\beta = 0.63$, $p = 0.000005$].

Following FDR correction, past year cannabis use and cannabis $\times$ gender interaction were not significantly associated with volumes in any other regions.

3.4 Secondary results: brain-behavior relationships

Within cannabis users, smaller left rACC volumes were significantly associated with lower Discrimination Correct scores [$r = 0.56$, $p = 0.01$; FDR-corrected: $p = 0.02$] (Figure 3). No other affective measures were significantly associated with left rACC volumes within the cannabis-using group.

4. DISCUSSION

To our knowledge, this is the first study to assess relationships between cannabis use, brain structure, and affective processing. We found that increased past year cannabis use was dose-dependently associated with smaller left rACC volumes in adolescents and young adults; gender did not significantly moderate this relationship. Notably, within cannabis-users, smaller left rACC volumes were significantly correlated with lower emotion discrimination accuracy.

Our finding that cannabis use was related to reduced left rACC volumes is consistent with other studies finding relationships between cannabis use and reduced volume and thickness

in the ACC (Hill et al., 2016; Jacobus et al., 2014; Lisdahl et al., 2016; Rais et al., 2010; Rapp et al., 2013; Szeszko et al., 2007). Notably, many of the aforementioned studies (Rais et al., 2010; Rapp et al., 2013; Szeszko et al., 2007) included samples with psychiatric comorbidities (e.g., ADHD, psychosis). However, one study (Hill et al., 2016) that was conducted in participants without other psychopathology found that cannabinoid-1 (*CNR1*) receptor genotype moderates the relationship between level of cannabis exposure and reduced right ACC volumes. The current study builds upon the *right* ACC findings of Hill and colleagues (2016) by demonstrating a dose-dependent relationship between cannabis use and *left rostral* ACC. Therefore, though consistent with previous results of reduced ACC gray matter, the current study is novel in finding reduced rACC (but not cACC) volumes, in relation to cannabis use. This is consistent with literature supporting the rACC's involvement in emotional processing and the cACC's mediation of attention (Bush et al., 2000). Further, within the present study, smaller rACC volumes were only in the left hemisphere. Left unilateral reductions in the ACC are not a consistent finding in the literature (Hill et al., 2016; Lisdahl et al., 2016; Rapp et al., 2013; Szeszko et al., 2007) and thus, perhaps specific to the rostral portion of the ACC. The ACC itself is rich in CB1 receptors (Herkenham et al., 1990) and chronic cannabis use leads to downregulation of CB1 receptors in the cingulate cortex (Hirvonen et al., 2012). CB1 downregulation may lead to reduced dendritic branching or neuronal atrophy, resulting in gray matter reductions (Lisdahl et al., 2014). The current study's findings suggest that rACC gray matter reductions may persist even after 3 weeks of abstinence. The present rACC structural findings also build upon studies showing relationships between both acute and chronic cannabis exposure and abnormal ACC activation and functional connectivity during emotion processing tasks (e.g., Gruber et al., 2009; Fusar-Poli et al., 2009a, 2010; Gorka et al., 2015; Wesley et al., 2016).

The ACC undergoes substantial structural and functional development during adolescence and young adulthood (Lichenstein et al., 2016; Pfeifer and Peake, 2012; Segalowitz and Dywan, 2009; Tamnes et al., 2013). Abnormal adolescent development of rACC structural and functional connectivity is associated with depression and anxiety (Clauss et al., 2014; Ho et al., 2017; Lichenstein et al., 2016; Swartz et al., 2014). Therefore, given the current study's findings, it is possible that adolescent cannabis use interrupts typical rACC development, increasing risk for affective disorders such as depression and anxiety. However, a longitudinal study would be necessary to confirm causal relationships between cannabis use, rACC structure, and development of affective disorders.

It is worth considering why Discrimination Correct was the only affective measure significantly related to the left rACC. Perhaps the rACC serves a crucial function in an aspect of the Emotion Discrimination task that is not present in the other tasks. For example, it is possible that the rACC is critical in comparing multiple emotional stimuli, such as faces (Munro et al., 2007). Additionally, research suggests a role for the rACC in shifting attention toward or away from emotional faces (Klumpp et al., 2012), which may be important if attending to multiple faces simultaneously. Another possibility is that the Emotion Discrimination task was more difficult (increased executive functioning load) compared to the Emotion Recognition and Emotional Acuity tasks, and thus more sensitive to rACC abnormalities. In functional neuroimaging studies of cannabis users, aberrant activation patterns often occur without differences in task performance (Lisdahl et al., 2014),

suggesting that cannabis users may be able to compensate by adjusting which regions are recruited for a particular task. Thus, perhaps due to simplicity of the Emotion Recognition and Emotional Acuity tasks, cannabis users in our sample were able to rely more heavily on other brain areas besides the rACC while completing those tasks, but not while completing the more demanding Emotion Discrimination task.

There are some additional important factors to consider when interpreting the current findings. Length of cannabis abstinence was much longer in our sample compared with many other studies (mean length of abstinence 25 days). In most studies, users are abstinent for an average of 2–3 days, and research suggests there may be significant cognitive recovery with abstinence (Hanson et al., 2010; Jacobus et al., 2012). Huijbregts and colleagues (2014) found that current cannabis users performed poorly on emotion recognition tasks (recognition and discrimination) relative to abstinent cannabis users and non-users. On the other hand, some structural and functional deficits persist in cannabis users after weeks of abstinence (Ganzer et al., 2016); one study that found emotion processing difficulties after three months of abstinence (Bayrakci et al., 2015). Therefore, the results of the current study suggest a mechanism (rACC structural differences) by which emotion processing deficits can persist in cannabis users. More research is needed to investigate specific emotion processing abilities (e.g., recognition vs. discrimination) in chronic cannabis use at varying lengths of abstinence. Another potential limitation is that age of cannabis use onset was not analyzed in the current study. Early age of onset in cannabis users has been related to poorer cognitive function as well as differences in brain activation and structure (Gruber et al., 2012, 2014; Pope et al., 2003; Sagar et al., 2015). However, it has been suggested that frequency of use over time, or dose-dependent effects such as those in the current analysis, likely have a greater impact on neurocognitive outcomes (Ganzer et al., 2016). Additionally, only one relationship (between cannabis and left rACC volumes) survived correction for multiple comparisons. As such, the other significant associations found may be spurious and must be interpreted with caution. Importantly, the current study only achieved 67% power at the effect size of the left rACC finding ($f^2 = 0.14$), and thus we cannot rule out Type I error (or Type II error for the other brain regions). Greater statistical power in future studies would increase the likelihood of detecting interactions between gender and cannabis use. Finally, given the cross-sectional nature of this study, more work is needed to determine causal relationships.

We found an association between greater past year cannabis use and smaller left rACC volumes, even after 3 weeks of abstinence. Further, within abstinent cannabis users, smaller left rACC volumes were associated with poorer accuracy on an emotion discrimination task. Thus, chronic cannabis use during adolescence and young adulthood, a time of continued neurodevelopment (Casey et al., 2008), may have a negative, lasting impact on affective circuitry. This could result in difficulty discriminating subtle differences in emotions, which may negatively influence mood and emotional well-being (Phillips et al., 2003). Alternatively, those with smaller left rACC volumes and poorer emotion discrimination abilities may be more likely to use cannabis. This could be one mechanism underlying the comorbidity of cannabis use and psychiatric disorders (Gater et al., 1998; Rosen et al., 2006; Wittchen et al., 2007). Longitudinal work is needed to understand the

causal relationships between substance use, comorbid disorders, gender, brain structure, and affective processing.

Acknowledgments

FUNDING

This research was funded by the National Institute on Drug Abuse (R01DA030354; PI: Lisdahl, K.M.). Manuscript preparation was supported by the National Institute of Health (U01DA041025; PI: Lisdahl, K.M.).

REFERENCES

- Agrawal A, Nurnberger JI, Lynskey MT, Bipolar Genome Study, 2011 Cannabis involvement in individuals with bipolar disorder. *Psychiatry Res.* 185, 459–461. [PubMed: 20674039]
- Ashtari M, Avants B, Cyckowski L, Cervellione KL, Roofeh D, Cook P et al., 2011 Medial temporal structures and memory functions in adolescents with heavy cannabis use. *J. Psychiatr. Res.* 45, 1055–1066. [PubMed: 21296361]
- Battistella G, Fornari E, Annoni JM, Chtioui H, Dao K, Fabritius M et al., 2014 Long-term effects of cannabis on brain structure. *Neuropsychopharmacology* 39, 2041–2048. [PubMed: 24633558]
- Bayrakci A, Sert E, Zorlu N, Erol A, Saricicek A, Mete L, 2015 Facial emotion recognition deficits in abstinent cannabis dependent patients. *Compr. Psychiatry* 58, 160–164. [PubMed: 25550274]
- Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J R Stat Soc* 57(1):289–300.
- Bourke C, Douglas K, Porter R, 2010 Processing of facial emotion expression in major depression: a review. *Aust. N. Z. J. Psychiatry* 44, 681–696. [PubMed: 20636189]
- Bush G, Luu P, Posner MI, 2000 Cognitive and emotional influences in anterior cingulate cortex. *Trends Cogn. Sci* 4, 215–222. [PubMed: 10827444]
- Casey BJ, Jones RM, Hare TA, 2008 The adolescent brain. *Ann. N. Y. Acad. Sci* 1124, 111–126. [PubMed: 18400927]
- Cheetham A, Allen NB, Whittle S, Simmons JG, Yucel M, Lubman DI, 2012 Orbitofrontal volumes in early adolescence predict initiation of cannabis use: a 4-year longitudinal and prospective study. *Biol. Psychiatry* 71, 684–692. [PubMed: 22129756]
- Churchwell JC, Lopez-Larson M, Yurgelun-Todd DA, 2010 Altered frontal cortical volume and decision making in adolescent cannabis users. *Front. Psychol* 1, 225. [PubMed: 21833280]
- Chye Y, Solowij N, Suo C, Batalla A, Cousijn J, Goudriaan AE et al., 2017 Orbitofrontal and caudate volumes in cannabis users: a multi-site mega-analysis comparing dependent versus non-dependent users. *Psychopharmacology (Berl)* 234, 1985–1995. [PubMed: 28364340]
- Clauss JA, Avery SN, VanDerKlok RM, Rogers BP, Cowan RL, Benningfield MM et al., 2014 Neurocircuitry underlying risk and resilience to social anxiety disorder. *Depress. Anxiety* 31, 822–833. [PubMed: 24753211]
- Cohen JE (1988) *Statistical power analysis for the behavioral sciences*. Lawrence Erlbaum Associates, Inc., Hillsdale, N.J.
- Cornelius JR, Aizenstein HJ, Hariri AR, 2010 Amygdala reactivity is inversely related to level of cannabis use in individuals with comorbid cannabis dependence and major depression. *Addict. Behav* 35, 644–646. [PubMed: 20189314]
- Cousijn J, Wiers RW, Ridderinkhof KR, van den Brink W, Veltman DJ, Goudriaan AE, 2012 Grey matter alterations associated with cannabis use: results of a VBM study in heavy cannabis users and healthy controls. *Neuroimage* 59, 3845–3851. [PubMed: 21982932]
- Crane NA, Schuster RM, Fusar-Poli P, Gonzalez R, 2013 Effects of cannabis on neurocognitive functioning: recent advances, neurodevelopmental influences, and sex differences. *Neuropsychol. Rev* 23, 117–137. [PubMed: 23129391]
- Demirakca T, Sartorius A, Ende G, Meyer N, Welzel H, Skopp G et al., 2011 Diminished gray matter in the hippocampus of cannabis users: possible protective effects of cannabidiol. *Drug Alcohol Depend.* 114, 242–245. [PubMed: 21050680]

- Desikan RS, Segonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D et al., 2006 An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage* 31, 968–980. [PubMed: 16530430]
- Dorard G, Berthoz S, Phan O, Corcos M, Bungener C, 2008 Affect dysregulation in cannabis abusers: a study in adolescents and young adults. *Eur. Child Adolesc. Psychiatry* 17, 274–282. [PubMed: 18301941]
- Epstein KA, Kumra S, 2015 Altered cortical maturation in adolescent cannabis users with and without schizophrenia. *Schizophr. Res* 162, 143–152. [PubMed: 25600549]
- Erwin RJ, Gur RC, Gur RE, Skolnick B, Mawhinney-Hee M, Smailis J, 1992 Facial emotion discrimination: I. Task construction and behavioral findings in normal subjects. *Psychiatry Res.* 42, 231–240. [PubMed: 1496055]
- Filkowski MM, Olsen RM, Duda B, Wanger TJ, Sabatinelli D, 2017 Sex differences in emotional perception: Meta analysis of divergent activation. *Neuroimage* 147, 925–933. [PubMed: 27988321]
- Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C et al., 2002 Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. *Neuron* 33, 341–355. [PubMed: 11832223]
- Fischl B, van der Kouwe A, Destrieux C, Halgren E, Segonne F, Salat DH et al., 2004 Automatically parcellating the human cerebral cortex. *Cereb. Cortex* 14, 11–22. [PubMed: 14654453]
- Fusar-Poli P, Allen P, Bhattacharyya S, Crippa JA, Mechelli A, Borgwardt S et al., 2010 Modulation of effective connectivity during emotional processing by Delta 9-tetrahydrocannabinol and cannabidiol. *Int. J. Neuropsychopharmacol* 13, 421–432. [PubMed: 19775500]
- Fusar-Poli P, Crippa JA, Bhattacharyya S, Borgwardt SJ, Allen P, Martin-Santos R et al., 2009 Distinct effects of {delta}9-tetrahydrocannabinol and cannabidiol on neural activation during emotional processing. *Arch. Gen. Psychiatry* 66, 95–105. [PubMed: 19124693]
- Fusar-Poli P, Placentino A, Carletti F, Landi P, Allen P, Surguladze S et al., 2009 Functional atlas of emotional faces processing: a voxel-based meta-analysis of 105 functional magnetic resonance imaging studies. *J. Psychiatry Neurosci* 34, 418–432. [PubMed: 19949718]
- Ganzer F, Broning S, Kraft S, Sack PM, Thomasius R, 2016 Weighing the Evidence: A Systematic Review on Long-Term Neurocognitive Effects of Cannabis Use in Abstinent Adolescents and Adults. *Neuropsychol. Rev* 26, 186–222. [PubMed: 27125202]
- Gater R, Tansella M, Korten A, Tiemens BG, Mavreas VG, Olatawura MO, 1998 Sex differences in the prevalence and detection of depressive and anxiety disorders in general health care settings: report from the World Health Organization Collaborative Study on Psychological Problems in General Health Care. *Arch. Gen. Psychiatry* 55, 405–413. [PubMed: 9596043]
- Gilman JM, Kuster JK, Lee S, Lee MJ, Kim BW, Makris N et al., 2014 Cannabis use is quantitatively associated with nucleus accumbens and amygdala abnormalities in young adult recreational users. *J. Neurosci* 34, 5529–5538. [PubMed: 24741043]
- Gorka SM, Phan KL, Lyons M, Mori S, Angstadt M, Rabinak CA, 2016 Cannabinoid modulation of frontolimbic activation and connectivity during volitional regulation of negative affect. *Neuropsychopharmacology* 41, 1888–1896. [PubMed: 26647971]
- Gruber SA, Dahlgren MK, Sagar KA, Gonenc A, Lukas SE, 2014 Worth the wait: effects of age of onset of marijuana use on white matter and impulsivity. *Psychopharmacology (Berl)* 231, 1455–1465. [PubMed: 24190588]
- Gruber SA, Rogowska J, Yurgelun-Todd DA, 2009 Altered affective response in marijuana smokers: an FMRI study. *Drug Alcohol Depend.* 105, 139–153. [PubMed: 19656642]
- Gruber SA, Sagar KA, Dahlgren MK, Racine M, Lukas SE, 2012 Age of onset of marijuana use and executive function. *Psychol. Addict. Behav* 26, 496–506. [PubMed: 22103843]
- Gur RC, Sara R, Hagendoorn M, Marom O, Hughett P, Macy L et al., 2002 A method for obtaining 3-dimensional facial expressions and its standardization for use in neurocognitive studies. *J. Neurosci. Methods* 115, 137–143. [PubMed: 11992665]
- Gur RC, Schroeder L, Turner T, McGrath C, Chan RM, Turetsky BI et al., 2002 Brain activation during facial emotion processing. *Neuroimage* 16, 651–662. [PubMed: 12169250]

- Hanson KL, Winward JL, Schweinsburg AD, Medina KL, Brown SA, Tapert SF, 2010 Longitudinal study of cognition among adolescent marijuana users over three weeks of abstinence. *Addict. Behav* 35, 970–976. [PubMed: 20621421]
- Haxby JV, Hoffman EA, Gobbini MI, 2000 The distributed human neural system for face perception. *Trends Cogn. Sci* 4, 223–233. [PubMed: 10827445]
- Herkenham M, Lynn AB, Little MD, Johnson MR, Melvin LS, de Costa BR et al., 1990 Cannabinoid receptor localization in brain. *Proc. Natl. Acad. Sci. U. S. A.* 87, 1932–1936. [PubMed: 2308954]
- Hill SY, Sharma V, Jones BL, 2016 Lifetime use of cannabis from longitudinal assessments, cannabinoid receptor (CNR1) variation, and reduced volume of the right anterior cingulate. *Psychiatry Res.* 255, 24–34.
- Hindocha C, Wollenberg O, Carter Leno V, Alvarez BO, Curran HV, Freeman TP, 2014 Emotional processing deficits in chronic cannabis use: a replication and extension. *J. Psychopharmacol* 28, 466–471. [PubMed: 24646810]
- Hirvonen J, Goodwin RS, Li CT, Terry GE, Zoghbi SS, Morse C et al., 2012 Reversible and regionally selective downregulation of brain cannabinoid CB1 receptors in chronic daily cannabis smokers. *Mol. Psychiatry* 17, 642–649. [PubMed: 21747398]
- Ho TC, Sacchet MD, Connolly CG, Margulies DS, Tymofiyeva O, Paulus MP et al., 2017 Inflexible functional connectivity of the dorsal anterior cingulate cortex in adolescent major depressive disorder. *Neuropsychopharmacology* 42, 2434–2445. [PubMed: 28553837]
- Huijbregts SC, Griffith-Lendering MF, Vollebergh WA, Swaab H, 2014 Neurocognitive moderation of associations between cannabis use and psychoneuroticism. *J. Clin. Exp. Neuropsychol* 36, 794–805. [PubMed: 25116129]
- Ishai A, Pessoa L, Bickle PC, Ungerleider LG, 2004 Repetition suppression of faces is modulated by emotion. *Proc. Natl. Acad. Sci. U. S. A.* 101, 9827–9832. [PubMed: 15210952]
- Ishai A, Schmidt CF, Boesiger P, 2005 Face perception is mediated by a distributed cortical network. *Brain Res. Bull* 67, 87–93. [PubMed: 16140166]
- Jacobus J, Goldenberg D, Wierenga CE, Tolentino NJ, Liu TT, Tapert SF, 2012 Altered cerebral blood flow and neurocognitive correlates in adolescent cannabis users. *Psychopharmacology (Berl)* 222, 675–684. [PubMed: 22395430]
- Jacobus J, Squeglia LM, Sorg SF, Nguyen-Louie TT, Tapert SF, 2014 Cortical thickness and neurocognition in adolescent marijuana and alcohol users following 28 days of monitored abstinence. *J. Stud. Alcohol. Drugs* 75, 729–743. [PubMed: 25208190]
- Jacobus J, Tapert SF, 2014 Effects of cannabis on the adolescent brain. *Curr. Pharm. Des* 20, 2186–2193. [PubMed: 23829363]
- Jarvis K, DelBello MP, Mills N, Elman I, Strakowski SM, Adler CM, 2008 Neuroanatomic comparison of bipolar adolescents with and without cannabis use disorders. *J. Child Adolesc. Psychopharmacol* 18, 557–563. [PubMed: 19108660]
- Johnston LD, O'Malley PM, Bachman JG, Schulenberg JE, Miech RA, 2016 Monitoring the Future national survey results on drug use, 1975–2015: Volume II, college students and adults ages 19–55. Institute for Social Research, The University of Michigan, Ann Arbor, MI.
- Kempton MJ, Haldane M, Jogia J, Christodoulou T, Powell J, Collier D et al., 2009 The effects of gender and COMT Val158Met polymorphism on fearful facial affect recognition: a fMRI study. *Int. J. Neuropsychopharmacol* 12, 371–381. [PubMed: 18796186]
- Kesler-West ML, Andersen AH, Smith CD, Avison MJ, Davis CE, Kryscio RJ et al., 2001 Neural substrates of facial emotion processing using fMRI. *Brain Res. Cogn. Brain Res* 11, 213–226. [PubMed: 11275483]
- Klump H, Angstadt M, Phan KL, 2012 Shifting the focus of attention modulates amygdala and anterior cingulate cortex reactivity to emotional faces. *Neurosci. Lett* 514, 210–213. [PubMed: 22425719]
- Koch K, Stegmaier S, Schwarz L, Erb M, Reinl M, Scheffler K et al., 2018 Neural correlates of processing emotional prosody in unipolar depression. *Hum. Brain Mapp*
- Koenders L, Cousijn J, Vingerhoets WA, van den Brink W, Wiers RW, Meijer CJ et al., 2016 Grey matter changes associated with heavy cannabis use: a longitudinal sMRI study. *PLoS One* 11, e0152482. [PubMed: 27224247]

- Lichenstein SD, Verstynen T, Forbes EE, 2016 Adolescent brain development and depression: A case for the importance of connectivity of the anterior cingulate cortex. *Neurosci. Biobehav. Rev* 70, 271–287. [PubMed: 27461914]
- Lisdahl KM, Gilbert ER, Wright NE, Shollenbarger S, 2013 Dare to delay? The impacts of adolescent alcohol and marijuana use onset on cognition, brain structure, and function. *Front. Psychiatry* 4, 53. [PubMed: 23847550]
- Lisdahl KM, Tamm L, Epstein JN, Jernigan T, Molina BS, Hinshaw SP et al., 2016 The impact of ADHD persistence, recent cannabis use, and age of regular cannabis use onset on subcortical volume and cortical thickness in young adults. *Drug Alcohol Depend.* 161, 135–146. [PubMed: 26897585]
- Lisdahl KM, Wright NE, Kirchner-Medina C, Maple KE, Shollenbarger S, 2014 Considering cannabis: the effects of regular cannabis use on neurocognition in adolescents and young adults. *Curr. Addict. Rep* 1, 144–156. [PubMed: 25013751]
- Lopez-Larson MP, Bogorodzki P, Rogowska J, McGlade E, King JB, Terry J et al., 2011 Altered prefrontal and insular cortical thickness in adolescent marijuana users. *Behav. Brain Res* 220, 164–172. [PubMed: 21310189]
- Mashhoon Y, Sava S, Sneider JT, Nickerson LD, Silveri MM, 2015 Cortical thinness and volume differences associated with marijuana abuse in emerging adults. *Drug Alcohol Depend.* 155, 275–283. [PubMed: 26249265]
- Mata I, Perez-Iglesias R, Roiz-Santianez R, Tordesillas-Gutierrez D, Pazos A, Gutierrez A et al., 2010 Gyrification brain abnormalities associated with adolescence and early-adulthood cannabis use. *Brain Res.* 1317, 297–304. [PubMed: 20045399]
- Matochik JA, Eldreth DA, Cadet JL, Bolla KI, 2005 Altered brain tissue composition in heavy marijuana users. *Drug Alcohol Depend.* 77, 23–30. [PubMed: 15607838]
- McQueeney T, Padula CB, Price J, Medina KL, Logan P, Tapert SF, 2011 Gender effects on amygdala morphometry in adolescent marijuana users. *Behav. Brain Res.* 224, 128–134. [PubMed: 21664935]
- Medina KL, McQueeney T, Nagel BJ, Hanson KL, Yang TT, Tapert SF, 2009 Prefrontal cortex morphometry in abstinent adolescent marijuana users: subtle gender effects. *Addict. Biol* 14, 457–468. [PubMed: 19650817]
- Medina KL, Nagel BJ, Tapert SF, 2010 Abnormal cerebellar morphometry in abstinent adolescent marijuana users. *Psychiatry Res.* 182, 152–159. [PubMed: 20413277]
- Medina KL, Schweinsburg AD, Cohen-Zion M, Nagel BJ, Tapert SF, 2007 Effects of alcohol and combined marijuana and alcohol use during adolescence on hippocampal volume and asymmetry. *Neurotoxicol. Teratol* 29, 141–152. [PubMed: 17169528]
- Mogg K, Millar N, Bradley BP, 2000 Biases in eye movements to threatening facial expressions in generalized anxiety disorder and depressive disorder. *J. Abnorm. Psychol* 109, 695–704. [PubMed: 11195993]
- Morris JS, Frith CD, Perrett DI, Rowland D, Young AW, Calder AJ et al., 1996 A differential neural response in the human amygdala to fearful and happy facial expressions. *Nature* 383, 812–815. [PubMed: 8893004]
- Morris RW, Weickert CS, Loughland CM, 2009 Emotional face processing in schizophrenia. *Curr. Opin. Psychiatry* 22, 140–146. [PubMed: 19553867]
- Munro GE, Dywan J, Harris GT, McKee S, Unsal A, Segalowitz SJ, 2007 ERN varies with degree of psychopathy in an emotion discrimination task. *Biol. Psychol* 76, 31–42. [PubMed: 17604898]
- Nurmedov S, Metin B, Ekmen S, Noyan O, Yilmaz O, Darcin A et al., 2015 Thalamic and cerebellar gray matter volume reduction in synthetic cannabinoids users. *Eur. Addict. Res* 21, 315–320. [PubMed: 26021304]
- Pagliaccio D, Barch DM, Bogdan R, Wood PK, Lynskey MT, Heath AC et al., 2015 Shared predisposition in the association between cannabis use and subcortical brain structure. *JAMA Psychiatry.* 72, 994–1001. [PubMed: 26308883]
- Pessoa L, McKenna M, Gutierrez E, Ungerleider LG, 2002 Neural processing of emotional faces requires attention. *Proc. Natl. Acad. Sci. U. S. A* 99, 11458–11463. [PubMed: 12177449]

- Pfeifer JH, Peake SJ, 2012 Self-development: integrating cognitive, socioemotional, and neuroimaging perspectives. *Dev. Cogn. Neurosci* 2, 55–69. [PubMed: 22682728]
- Phillips ML, Drevets WC, Rauch SL, Lane R, 2003 Neurobiology of emotion perception I: The neural basis of normal emotion perception. *Biol. Psychiatry* 54, 504–514. [PubMed: 12946879]
- Pinkham AE, Sasson NJ, Calkins ME, Richard J, Hughett P, Gur RE et al., 2008 The other-race effect in face processing among African American and Caucasian individuals with schizophrenia. *Am. J. Psychiatry* 165, 639–645. [PubMed: 18347000]
- Platt B, Kamboj S, Morgan CJ, Curran HV, 2010 Processing dynamic facial affect in frequent cannabis-users: evidence of deficits in the speed of identifying emotional expressions. *Drug Alcohol Depend.* 112, 27–32. [PubMed: 21036306]
- Pope HG, Gruber AJ, Hudson JI, Cohane G, Huestis MA, Yurgelun-Todd D, 2003 Early-onset cannabis use and cognitive deficits: what is the nature of the association? *Drug Alcohol Depend.* 69, 303–310. [PubMed: 12633916]
- Price JS, McQueeney T, Shollenbarger S, Browning EL, Wieser J, Lisdahl KM, 2015 Effects of marijuana use on prefrontal and parietal volumes and cognition in emerging adults. *Psychopharmacology (Berl)* 232, 2939–2950. [PubMed: 25921032]
- Rais M, van Haren NE, Cahn W, Schnack HG, Lepage C, Collins L et al., 2010 Cannabis use and progressive cortical thickness loss in areas rich in CB1 receptors during the first five years of schizophrenia. *Eur. Neuropsychopharmacol* 20, 855–865. [PubMed: 20863671]
- Rapp C, Walter A, Studerus E, Bugra H, Tamagni C, Rothlisberger M et al., 2013 Cannabis use and brain structural alterations of the cingulate cortex in early psychosis. *Psychiatry Res.* 214, 102–108. [PubMed: 24054726]
- Rosen JL, Miller TJ, D'Andrea JT, McGlashan TH, Woods SW, 2006 Comorbid diagnoses in patients meeting criteria for the schizophrenia prodrome. *Schizophr. Res* 85, 124–131. [PubMed: 16650735]
- Rubino T, Viganò D, Realini N, Guidali C, Braida D, Capurro V et al., 2008 Chronic delta 9-tetrahydrocannabinol during adolescence provokes sex-dependent changes in the emotional profile in adult rats: behavioral and biochemical correlates. *Neuropsychopharmacology* 33, 2760–2771. [PubMed: 18172430]
- Sagar KA, Dahlgren MK, Gonenc A, Racine MT, Dremar MW, Gruber SA, 2015 The impact of initiation: Early onset marijuana smokers demonstrate altered Stroop performance and brain activation. *Dev. Cogn. Neurosci* 16, 84–92. [PubMed: 25936584]
- Said CP, Haxby JV, Todorov A, 2011 Brain systems for assessing the affective value of faces. *Philos. Trans. R. Soc. Lond. B. Biol. Sci* 366, 1660–1670. [PubMed: 21536552]
- Samame C, 2013 Social cognition throughout the three phases of bipolar disorder: a state-of-the-art overview. *Psychiatry Res.* 210, 1275–1286. [PubMed: 24075306]
- Schacht JP, Hutchison KE, Filbey FM, 2012 Associations between cannabinoid receptor-1 (CNR1) variation and hippocampus and amygdala volumes in heavy cannabis users. *Neuropsychopharmacology* 37, 2368–2376. [PubMed: 22669173]
- Segalowitz SJ, Dywan J, 2009 Individual differences and developmental change in the ERN response: implications for models of ACC function. *Psychol. Res* 73, 857–870. [PubMed: 19023593]
- Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E et al., 1998 The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J. Clin. Psychiatry* 59 Suppl 20, 57.
- Solowij N, Walterfang M, Lubman DI, Whittle S, Lorenzetti V, Styner M et al., 2013 Alteration to hippocampal shape in cannabis users with and without schizophrenia. *Schizophr. Res* 143, 179–184. [PubMed: 23201308]
- Song Z, Zhang M, Huang P, 2016 Aberrant emotion networks in early major depressive disorder patients: an eigenvector centrality mapping study. *Transl. Psychiatry* 6, e819. [PubMed: 27219345]
- Spechler PA, Orr CA, Chaarani B, Kan KJ, Mackey S, Morton A et al., 2015 Cannabis use in early adolescence: Evidence of amygdala hypersensitivity to signals of threat. *Dev. Cogn. Neurosci* 16, 63–70. [PubMed: 26347227]
- Stewart DG, Brown SA, 1995 Withdrawal and dependency symptoms among adolescent alcohol and drug abusers. *Addiction.* 90:627–635. [PubMed: 7795499]

- Stuhrmann A, Suslow T, Dannlowski U, 2011 Facial emotion processing in major depression: a systematic review of neuroimaging findings. *Biol. Mood Anxiety Disord* 1, 10. [PubMed: 22738433]
- Swartz JR, Phan KL, Angstadt M, Klumpp H, Fitzgerald KD, Monk CS, 2014 Altered activation of the rostral anterior cingulate cortex in the context of emotional face distractors in children and adolescents with anxiety disorders. *Depress. Anxiety* 31, 870–879. [PubMed: 24995682]
- Szeszko PR, Robinson DG, Sevy S, Kumra S, Rupp CI, Betensky JD et al., 2007 Anterior cingulate grey-matter deficits and cannabis use in first-episode schizophrenia. *Br. J. Psychiatry* 190, 230–236. [PubMed: 17329743]
- Tamnes CK, Walhovd KB, Torstveit M, Sells VT, Fjell AM, 2013 Performance monitoring in children and adolescents: a review of developmental changes in the error-related negativity and brain maturation. *Dev. Cogn. Neurosci* 6, 1–13. [PubMed: 23777674]
- Thames AD, Kuhn TP, Williamson TJ, Jones JD, Mahmood Z, Hammond A, 2017 Marijuana effects on changes in brain structure and cognitive function among HIV+ and HIV- adults. *Drug Alcohol Depend.* 170, 120–127. [PubMed: 27889592]
- Troisi A, Pasini A, Saracco M, Spalletta G, 1998 Psychiatric symptoms in male cannabis users not using other illicit drugs. *Addiction* 93, 487–492. [PubMed: 9684387]
- Tzilos GK, Cintron CB, Wood JB, Simpson NS, Young AD, Pope HG et al., 2005 Lack of hippocampal volume change in long-term heavy cannabis users. *Am. J. Addict* 14, 64–72. [PubMed: 15804878]
- Weiland BJ, Thayer RE, Depue BE, Sabbineni A, Bryan AD, Hutchison KE, 2015 Daily marijuana use is not associated with brain morphometric measures in adolescents or adults. *J. Neurosci* 35, 1505–1512. [PubMed: 25632127]
- Wesley MJ, Lile JA, Hanlon CA, Porrino LJ, 2016 Abnormal medial prefrontal cortex activity in heavy cannabis users during conscious emotional evaluation. *Psychopharmacology (Berl)* 233, 1035–1044. [PubMed: 26690589]
- Whittle S, Yucel M, Yap MB, Allen NB, 2011 Sex differences in the neural correlates of emotion: evidence from neuroimaging. *Biol. Psychol* 87, 319–333. [PubMed: 21600956]
- Wittchen HU, Frohlich C, Behrendt S, Gunther A, Rehm J, Zimmermann P et al., 2007 Cannabis use and cannabis use disorders and their relationship to mental disorders: a 10-year prospective-longitudinal community study in adolescents. *Drug Alcohol Depend.* 88 Suppl 1, 60.
- Wu M, Kujawa A, Lu LH, Fitzgerald DA, Klumpp H, Fitzgerald KD et al., 2016 Age-related changes in amygdala-frontal connectivity during emotional face processing from childhood into young adulthood. *Hum. Brain Mapp.* 37, 1684–1695. [PubMed: 26931629]
- Yucel M, Solowij N, Respondek C, Whittle S, Fornito A, Pantelis C et al., 2008 Regional brain abnormalities associated with long-term heavy cannabis use. *Arch. Gen. Psychiatry* 65, 694–701. [PubMed: 18519827]

Highlights

- Cannabis was associated with smaller left rostral anterior cingulate cortex volumes.
- Gender did not moderate the relationship between cannabis use and volumes.
- Rostral anterior cingulate volumes were related to affective processing.

Figure 1. Increased past year cannabis use significantly predicted smaller left rACC volumes [$t(48) = -3.09$, $\beta = -0.33$, $p = 0.003$, $r^2 = 0.14$; FDR corrected: $p = 0.02$].

Figure 2.

Increased cannabis use significantly predicted smaller left rACC volumes [$t(48) = -3.09$, $\beta = -0.33$, $p = 0.003$, $r^2 = 0.14$; FDR corrected: $p = 0.02$].

Figure 3.

Within the cannabis-using group, smaller left rACC volumes significantly predicted lower Discrimination Correct scores [$r = 0.56$, $p = 0.01$; FDR-corrected: $p = 0.02$].

Table 1.

Emotion Recognition Principal Components Analysis Factor Loadings

Affective Variable	Component Labels			
	Recognition Time	Sad Correct	Happy Correct	Anger Correct
Anger Correct	−0.021	−0.001	−0.027	0.939
Fear Correct	−0.433	0.554	0.298	0.289
Sad Correct	0.096	0.913	−0.063	−0.078
Happy Correct	0.030	0.001	0.964	−0.025
Anger Correct Median Response Time	0.659	−0.310	0.106	0.280
Fear Correct Median Response Time	0.822	−0.058	−0.303	−0.049
Happy Correct Median Response Time	0.649	−0.116	−0.078	−0.122
Neutral Correct Median Response Time	0.775	0.366	0.120	0.110
Sad Correct Median Response Time	0.823	0.077	0.164	−0.097

Note. Bolded values denote variable loadings of 0.6. All response times in milliseconds.

Table 2.

Emotional Acuity Principal Components Analysis Factor Loadings

Affective Variable	Component Labels		
	Acuity Time	Acuity Neutral	Acuity Intensity
Total Correct	−0.131	0.894	0.416
Very Happy Correct	−0.125	−0.107	0.807
Happy Neutral Correct	−0.122	0.854	−0.124
Neutral Correct	−0.214	0.923	−0.082
Sad Neutral Correct	−0.190	0.755	0.050
Very Sad Correct	0.209	0.163	0.650
Correct Trials Median Response Time	0.962	−0.189	0.085
Correct Very Happy Trials Median Response Time	0.804	0.071	−0.180
Correct Happy Neutral Trials Median Response Time	0.836	−0.190	−0.017
Correct Neutral Trials Median Response Time	0.914	−0.262	0.083
Correct Sad Neutral Trials Median Response Time	0.865	−0.143	0.183
Correct Very Sad Trials Median Response Time	0.646	−0.199	0.016

Note. Bolded values denote variable loadings of ≥ 0.6 . All response times in milliseconds. The —Acuity Neutral component has high loadings from variables indicating total correct identifications as well as correct identification of more neutral emotions. The —Acuity Intensity component has high loadings from variables indicating correct identification of more extreme or intense emotions.

Table 3.

Emotion Discrimination Principal Components Analysis Factor Loadings

Affective Variable	Component Labels	
	Discrimination Time	Discrimination Correct
Total Correct	0.091	0.991
Happy Trials Correct	0.147	0.886
Sad Trials Correct	−0.008	0.856
Correct Happy Trials Median Response Time	0.910	−0.001
Incorrect Happy Trials Median Response Time	0.732	0.387
Correct Sad Trials Median Response Time	0.882	−0.014
Incorrect Sad Trials Median Response Time	0.859	0.182
Total Correct Trials Median Response Time	0.975	−0.007

Note. Bolded values denote variable loadings of .6. All response times in milliseconds.

Table 4.**Participant Demographics**

	CAN Females (n=8)	CNT Females (n=19)	CAN Males (n=12)	CNT Males (n=16)
	% or $M \pm SD$ (range)	% or $M \pm SD$ (range)	% or $M \pm SD$ (range)	% or $M \pm SD$ (range)
Race (% Caucasian)	37.5%	73.7%	83.3%	68.8%
Ethnicity (% Non-Hispanic)	62.5%	78.9%	91.7%	93.7%
Age (years)	21.4 $\pm$ 2.0 (19–24)	20.8 $\pm$ 2.5 (16–25)	22.0 $\pm$ 2.4 (18–26)	20.5 $\pm$ 3.2 (16–25)
Education (years)	14.1 $\pm$ 1.4 (12–16)	14.1 $\pm$ 1.9 (11–18)	14.4 $\pm$ 1.8 (11–18)	14.0 $\pm$ 3.0 (9–19)
WRAT-4 Word Reading (raw score)	58.8 $\pm$ 7.9 (41–67)	61.2 $\pm$ 4.2 (53–69)	62.0 $\pm$ 4.0 (55–69)	62.5 $\pm$ 3.8 (55–68)
Age of weekly cannabis use onset (years)	17.5 $\pm$ 0.9 (16–19)	-	17.6 $\pm$ 1.7 (15–20)	-
Lifetime cannabis use (uses)	797.8 $\pm$ 723.7 (101–2314) *	0.7 $\pm$ 1.2 (0–3) *	1339.8 $\pm$ 1550.0 (250–6000) *	0.6 $\pm$ 1.4 (0–5) *
Past year cannabis use (joints)	318.9 $\pm$ 301.3 (44.7–879.3) *	0.0 $\pm$ 0.0 (0–0) *	382.3 $\pm$ 361.4 (59–1394) *	.4 $\pm$ 1.3 (0–4.8) *
Number of cannabis uses/month in past 3 months	18.4 $\pm$ 12.1 (0–30) *	0.0 $\pm$ 0.0 (0–0) *	20.5 $\pm$ 10.0 (3–30) *	0.1 $\pm$ 0.3 (0–1) *
Length of cannabis abstinence (days)	26.0 $\pm$ 6.8 (22–40)	-	25.0 $\pm$ 9.3 (19–54)	-
Past year alcohol use (standard drinks)	339.6 $\pm$ 349.9 (0–883) *	59.4 $\pm$ 102.6 (0–450) *	433.5 $\pm$ 275.7 (70.5–897) *	115.5 $\pm$ 187.6 (0–698.5) *
Past year nicotine use (cigarettes)	46.4 $\pm$ 84.9 (0–232.3) *	.8 $\pm$ 2.8 (0–12) *	264.3 $\pm$ 448.2 (0–1088.5) *	0.1 $\pm$ 0.3 (0–1) *
Left Rostral Anterior Cingulate Cortex (rACC) volumes (mm ³)	2449.0 $\pm$ 490.3 (1584–3180)	2777.6 $\pm$ 527.9 (1917–3658)	3125.9 $\pm$ 421.4 (2324–3741)	3257.3 $\pm$ 472.7 (2574–4322)

Note. WRAT-4= Wide Range Achievement Test –4th edition Word Reading subtest. CAN= Cannabis-using. CNT= Non-using control.

* Differences in CAN vs. CNT within gender= $p < 0.05$.