Published in final edited form as:

Trends Pharmacol Sci. 2019 May 6; 40(6): 362–364. doi:10.1016/j.tips.2019.04.010.

Designer drugs for designer receptors: unlocking the translational potential of chemogenetics

Ofer Yizhar¹ and J. Simon Wiegert²

¹Department of Neurobiology, Weizmann Institute of Science, Rehovot, Israel

²Research Group Synaptic Wiring and Information Processing, Center for Molecular Neurobiology Hamburg, Falkenried 94, 20251 Hamburg, Germany

Abstract

Chemogenetic techniques allow selective manipulation of neurons by activating engineered actuator proteins with otherwise-inert effector molecules. A recent study describes the co-evolution of highly potent actuator-effector pairs based on a clinically approved anti-smoking drug. These tools allow selective excitation or inhibition of neurons in the living brain with high specificity and no detectable side-effects.

Naturally-occurring receptor proteins, such as ligand-gated ion channels, have evolved over billions of years to be activated efficiently and exclusively by highly selective small-molecule ligands. The discovery and synthesis of exogenous ligands has massively influenced biomedical research and strongly boosted development of efficient medical treatments. This has spurred the race for the discovery of novel, highly selective compounds. However, even the most selective drug will always affect all cells and tissues expressing its target receptor, often leading to undesired side-effects. This is a particularly difficult problem in treating disorders of the central nervous system (CNS), with its incredible diversity of cell types and circuits and the semi-permeable blood-brain barrier (BBB). Many drugs do not efficiently pass the BBB, and instead exert off-target effects elsewhere in the body.

Chemogenetic tools are two-component actuator systems designed to overcome these limitations, allowing pharmacological modulation of specific cell populations through a combination of genetic expression of an engineered receptor and delivery of a specific ligand designed to only activate the exogenous receptor [1]. Chemogenetic receptors typically constitute ligand-gated ion channels or G-protein coupled receptors that have been designed to lose affinity to their natural ligand and instead become activated by a highly specific, but otherwise inert small-molecule agonist that can be delivered either locally or systemically.

Although chemogenetic tools have only been used in basic research to date, their translational appeal is clear: if genetic expression of the engineered receptor is feasible in the targeted cell population, systemic delivery of its ligand can circumvent any side effects

associated with conventional pharmacotherapy. To fulfill this promise, a clinically-relevant chemogenetic actuator system needs to satisfy several requirements: it should (1) allow high-affinity activation of the engineered receptor, (2) trigger minimal activation of off-target receptors, (3) easily cross the BBB and (4) ideally be based on an Food and Drug Administration (FDA)-approved drug, thus reducing the “energy barrier” for drug approval and translation to the clinic. Putting aside the relatively limited experience with long-term expression of heterologous transgenes in the human CNS, currently available ligands fall short of meeting these requirements, often suffering from low potency [2] or limited selectivity [3]. Moreover, some chemogenetic ligands have been shown to trigger side effects due to metabolic conversion to other neuroactive compounds [4, 5].

To address this challenge, Magnus et al. went back to the scratchboard (or chemical bench) and re-designed the so-called Pharmacologically Selective Actuator Molecules (PSAMs) together with highly specific and selective Pharmacologically Selective Effector Molecules (PSEMs)[2], yielding a highly specific and extremely potent chemogenetic system for selective control of neuronal excitability in the mammalian central nervous system [6]. The study, which includes proof-of-concept experiments in mice and a non-human primate, constitutes a major leap in the race for clinically applicable strategies to control the activity of particular populations of neurons with high selectivity and precision.

PSAMs are chimeric receptors consisting of a ligand binding domain (LBD) of the $\alpha 7$ nicotinic acetylcholine receptor ($\alpha 7$ -nACh) fused to either the chloride-conducting pore of a glycine receptor ($\alpha 7$ -GlyR) for neuronal inhibition or a cation-conducting pore of the serotonin (5HT) receptor 3 ($\alpha 7$ -5HT3) for neuronal excitation (Fig. 1A). Using structure-guided mutagenesis of the LBD, the authors reduced the potency of its natural agonist ACh 13-fold while increasing it 390-fold for Varenicline, an FDA-approved cholinergic agonist used to treat nicotine addiction (Fig. 1B). This yielded potent PSAMs, termed PSAM⁴-5HT3 and PSAM⁴-GlyR that are fully activated by low-nanomolar concentrations of Varenicline, but not by physiological concentrations of acetylcholine (Fig. 1A, B).

Using PSAM⁴-GlyR, Magnus et al. were able to continuously inhibit spiking of cultured neurons for more than 2 weeks (Fig. 1C). Moreover, Varenicline could easily be washed-out to restore normal spiking without any obvious long-lasting change of neuronal excitability. This is remarkable, given that chronic manipulations of neuronal activity typically lead to homeostatic adaptations working against the manipulation [7]. Conversely, a depolarized membrane potential was maintained with PSAM⁴-5HT3 for more than 2 weeks (Fig. 1C). Due to its highly efficient brain penetrance and long lifetime, systemically injected Varenicline displayed robust neuronal silencing in PSAM⁴-GlyR-expressing mice or a primate at doses as low as 0.1 mg per kg bodyweight.

Although the two PSAM⁴ variants were potently activated by Varenicline at low doses, the authors set out to generate PSEMs with improved PSAM⁴-selectivity over Varenicline’s “natural” targets. For this, they used again the molecular structure of the LBD as blueprint to evolve molecular analogs of Varenicline with improved binding to the LBD of PSAM⁴ and reduced binding to natural receptors (Fig. 1D). These ultrapotent PSEMs (uPSEMs) were active in the sub-nanomolar range and showed no detectable agonism for typical Varenicline

targets such as $\alpha 2\beta 4$ -, 5HT3- or $\alpha 7$ -nACh-receptors. These uPSEMs retained their water solubility and brain penetrance upon intraperitoneal injections, making systemic brain delivery *in vivo* straightforward. This is confirmed by a lowest effective dose of 0.03 mg/kg for the variant uPSEM⁷⁹³. Since uPSEMs are based on the clinically approved anti-smoking drug Varenicline, they constitute a promising step towards clinical chemogenetic applications.

As proof-of-concept of the robustness of their approach, the authors demonstrate that their new PSAM⁴-uPSEM-system is suitable to reliably suppress neuronal activity of so-called place cells in the mouse hippocampus. These neurons typically fire strong bursts of action potentials when an animal traverses a certain location. Fixing mice on a treadmill studded with spatial marks, these cells can be directly visualized under a microscope. When uPSEM⁷⁹² was injected systemically, spiking of such place cells was completely abrogated, demonstrating directly the strong potency of this chemogenetic toolset.

The high potency and selectivity of the PSAM⁴-uPSEM-system will make it an indispensable research tool for many neuroscientific questions. It appears especially well suited for chronic circuit manipulations, where other systems fail due to side effects or low potency. Nonetheless, there may be cases where the use of PSAMs is limited. Since PSAM⁴-GlyR is a chloride channel, neuronal silencing depends on the electrochemical chloride gradient, which in most cases is indeed inhibitory at the somatodendritic compartment of mature neurons. However, activating PSAM⁴-GlyR at axonal terminals might lead to a depolarization and subsequent neurotransmitter release due to a depolarizing axonal chloride gradient [8]. Furthermore, the cellular homeostasis of chloride ions could be altered in certain neurological disorders [9]. In these cases, G-protein coupled effectors that belong to the family of “Designer receptors exclusively activated by designer drugs” (DREADDs), such as hM4D [10] could still prove to be a better choice. Ultimately, development of a K⁺-selective PSAM might overcome all remaining limitations, allowing efficient and compartment-independent neuronal silencing.

In summary, the new addition to the chemogenetic toolbox presented by Magnus et al. will open up new avenues for cell-type specific manipulations of neuronal activity – particularly in intact animals. Building on the clinically-approved drug Varenicline, which is well-tolerated, water-soluble, has a long half-life and penetrates the brain efficiently, the new ultrapotent PSEMs tailored to the highly selective LBD of the next-generation PSAMs, might be the first of the chemogenetic actuators to reach clinical testing, once transgene delivery to the CNS becomes feasible in a clinical setting.

References

1. Sternson SM, Roth BL. Chemogenetic tools to interrogate brain functions. *Annu Rev Neurosci.* 2014; 37:387–407. [PubMed: 25002280]
2. Magnus CJ, et al. Chemical and genetic engineering of selective ion channel-ligand interactions. *Science.* 2011; 333(6047):1292–6. [PubMed: 21885782]
3. Chen X, et al. The first structure-activity relationship studies for designer receptors exclusively activated by designer drugs. *ACS Chem Neurosci.* 2015; 6(3):476–84. [PubMed: 25587888]

4. Gomez JL, et al. Chemogenetics revealed: DREADD occupancy and activation via converted clozapine. *Science*. 2017; 357(6350):503–507. [PubMed: 28774929]
5. Jann MW, et al. Rapid formation of clozapine in guinea-pigs and man following clozapine-N-oxide administration. *Arch Int Pharmacodyn Ther*. 1994; 328(2):243–50. [PubMed: 7710309]
6. Magnus CJ, et al. Ultrapotent chemogenetics for research and potential clinical applications. *Science*. 2019
7. Keck T, et al. Integrating Hebbian and homeostatic plasticity: the current state of the field and future research directions. *Philos Trans R Soc Lond B Biol Sci*. 2017; 372(1715)
8. Mahn M, et al. High-efficiency optogenetic silencing with soma-targeted anion-conducting channelrhodopsins. *Nat Commun*. 2018; 9(1):4125. [PubMed: 30297821]
9. Doyon N, et al. Chloride Regulation: A Dynamic Equilibrium Crucial for Synaptic Inhibition. *Neuron*. 2016; 89(6):1157–72. [PubMed: 26985723]
10. Armbruster BN, et al. Evolving the lock to fit the key to create a family of G protein-coupled receptors potently activated by an inert ligand. *Proc Natl Acad Sci U S A*. 2007; 104(12):5163–8. [PubMed: 17360345]

Figure 1. A novel, highly selective and ultra-sensitive chemogenetic toolkit for manipulation of neuronal activity.

(A) Pharmacologically Selective Actuator Molecules (PSAMs) are chimera consisting of a ligand binding domain (LBD) from the $\alpha 7$ -nicotinic acetylcholine receptor and an ion pore domain (IPD) of the glycine receptor (GlyR, anion-conducting) or the serotonin receptor (5HT, cation-conducting). (B) The acetylcholine binding pocket of the $\alpha 7$ -nAChR-LBD (symbolizing the lock) is mutated to become selective for the Pharmacologically Selective Effector Molecule (PSAM) Varenicline, a clinically approved anti-smoking drug. (C) Expression of PSAM⁴-GlyR and PSAM⁴-5HT in defined neuronal populations allows selective down- or upregulation of neuronal activity with nanomolar concentrations of Varenicline and ultrapotent PSEMs (uPSEMs). (D) uPSEMs (symbolizing the keys to fit the binding pocket of the LBD) are derived from Varenicline and optimized to fit the PSAM⁴ binding pocket with higher specificity than Varenicline.