Published in final edited form as:

J Bone Miner Metab. 2019 July ; 37(4): 648–657. doi:10.1007/s00774-018-0969-1.

Calcium restriction during lactation has minimal effects on post-weaning mineral metabolism and bone recovery

Ryan D. Ross^{1,2}, Matthew J. Meagher¹, and D. Rick Sumner^{1,2}

¹Department of Cell & Molecular Medicine, Rush University Medical Center, Chicago, IL

²Department of Orthopedic Surgery, Rush University Medical Center, Chicago, IL

Abstract

Dietary calcium (Ca) restriction during lactation in the rat, which induces intra-cortical and endocortical remodeling, has been proposed as a model to study bone matrix maturation in the adult skeleton. The purpose of this study was to assess the effects of dietary Ca restriction during lactation on post-weaning mineral metabolism and bone formation. Mated female Sprague-Dawley rats were randomized into groups receiving either 0.6% Ca (lactation/normal Ca) or 0.01% Ca (lactation/low Ca) diets during lactation. Virgin animals fed normal Ca were used as controls (virgin/normal Ca). At the time of weaning, animals on the low Ca diet were returned to normal Ca and cohorts of all three groups were sacrificed at days 0, 1, 2, 7, and 14 post-weaning. Lactation caused bone loss, particularly at the endocortical surface, but the amount was not affected by dietary Ca. Rats in the lactation/low Ca group had increased cortical porosity compared to the other groups, particularly within the size range of secondary osteons. Dietary Ca restriction during lactation did not affect post-weaning bone formation kinetics or serum Ca and phosphate levels. In both lactation groups, there was a transient increase in phosphate and fibroblast growth factor 23 (FGF23) post-weaning, which trended toward virgin/normal Ca levels over time. Thus, the additional challenge of low dietary Ca during lactation to induce intra-cortical remodeling in the rat has minimal effects on bone formation kinetics and mineral metabolism during the post-weaning period, providing further justification for this model to study matrix maturation in the adult skeleton.

Introduction

Lactation represents a challenge to the mother to provide the necessary supply of calcium (Ca) to the developing neonate [1]. The primary mechanism by which Ca is mobilized to the breast milk during lactation is through an upregulation of skeletal remodeling, which is uncoupled to favor resorption [2]. Increased skeletal resorption can be substantial, with losses of 30–40% in femoral ash weight in lactating rats [3,4]. After weaning, bone remodeling shifts in favor of new bone formation [5,6], leading to large regions of rapidly forming bone.

Statement of Animal Rights

All applicable institutional and/or national guidelines for the care and use of animals were followed. The articles does not contain any studies with human participants performed by any of the authors.

By taking advantage of the highly anabolic state following weaning, our laboratory has recently demonstrated the usefulness of the low Ca during lactation model to study matrix maturation throughout the cortical bone, including in Haversian-like remodeling units [7]. Despite the extensive bone loss due to the combination of lactation and Ca deficiency, the post-weaning bone matrix maturation process was largely unaffected by this dietary deprivation and the circulating Ca and phosphate levels were not different compared to virgin control animals fed an adequate Ca diet [7].

Adaptations in mineral metabolism associated with lactation have been described in some detail in previous reviews [1,8,9]. The post-weaning recovery phase, however, is not as well-characterized and to our knowledge there have been no investigations comparing the post-lactation recovery process in animals fed adequate Ca to those on a restricted Ca diet during lactation. Therefore, the current study aimed to investigate bone formation kinetics and mineral metabolism following weaning in rats fed normal (0.6% Ca) and low Ca (0.01% Ca) diets during lactation. Due to the anticipated large extent of bone loss associated with the combination of lactation and a low Ca diet, we hypothesized that animals fed a low Ca diet during lactation would have reduced bone formation and delayed normalization of mineral metabolism in the post-weaning recovery period.

Methods and Materials

Study Design.

A total of 75, 12 week old female Sprague-Dawley rats were purchased from Envigo Labs. 50 of the 75 animals were bred at 10 weeks of age and shipped at E15, 6–7 days before parturition, while the remaining 25 animals were age-matched virgin controls. The animals were then individually housed in the Rush University Comparative Research Center under an institutionally approved protocol.

Upon arrival, all animals were immediately placed on a normal Ca diet (0.6% Ca, 0.4% phosphate, Teklad diets, TD.97191). Similar to our previous study [7], the experimental design included two phases: induction (lactation) and recovery (post-weaning). Induction was initiated when the animals in the timed pregnancy groups gave birth, which occurred at 13 weeks of age (± 2 days). At parturition, the 50 dams were randomly assigned to (a) either the normal Ca (0.6% Ca, 0.4% phosphate) or a low Ca (0.01% Ca, 0.4% phosphate, Tekland diets, TD.9507) diet for the duration of the lactation period (i.e., the induction phase) and (b) one of 5 recovery times (days 0, 1, 2, 7 or 14 post-weaning). Virgin animals were randomly assigned to time-matched control groups. At parturition, litters were normalized to 12 pups per dam and all fostered pups were accepted without incident. On day 17 of lactation, the pups were weaned. The animals in the lactation/low Ca group were returned to the normal Ca diet. One animal in the lactation/low Ca group died during parturition and another rat in the lactation/normal Ca died one day after weaning, meaning that most time points had 5 animals but two had 4 animals (Table 1).

During the course of the recovery period, three animals in the lactation/normal Ca and one animal in the lactation/low Ca group developed mammary gland infections (mastitis) attributed to wounds that developed during the course of lactation. On day 2 of recovery, all

animals received a prophylactic three-day course of subcutaneous antibiotics, 100 mg/kg cefazolin. Endpoints derived from animals with mastitis were compared to non-infected animals of the same group to determine whether the localized infection impacted the study results. No outliers in any of the endpoints were noted in these animals and they were, therefore, included in the study.

Calcein (25 mg/kg, Sigma Aldrich) was injected subcutaneously to label forming bone with variable injection strategies depending on the sacrifice time. Animals sacrificed on the first day of recovery were given a single calcein injection at weaning (day 0). Animals sacrificed on day 2 were given calcein injections at days 0 and 1 of recovery. Animals sacrificed on day 7 were given calcein injections on days 3 and 6, and animals sacrificed on day 14 were given calcein injections on days 7 and 12.

At sacrifice, the left femurs were fixed in 70% ethanol. Blood was collected via cardiac puncture using a 25 G needle, allowed to clot at room temperature for ~30 mins and centrifuged at 1,000g for 15 mins at 4°C (Eppendorf 5415R) to separate serum.

Circulating Mineral Metabolism Biomarkers.

Ionic calcium (BioVision) and phosphate (BioVision) were assessed by colorimetric assays. Circulating mineral metabolism biomarkers were assessed using commercially available ELISA assays, including parathyroid hormone (PTH, UCSNK), parathyroid hormone related hormone (PTHrP, UCSNK), calcitonin (UCSNK), and full length fibroblast growth factor 23 (FGF23, Immuntopics). All markers were assessed at days 0, 1, 2, 7, and 14 of recovery.

Micro-Computed Tomography (μ CT).

μ CT scanning was performed using a tube voltage of 70 kVp, a current of 57 μ A, an integration time of 1500 ms, and a voxel size of 2 μ m (μ CT50, Scanco Medical). Cortical geometry and porosity were assessed in a 500 μ m thick transverse cross-sectional region of interest in the distal femoral diaphysis, located at 70% of the total femoral length. This anatomical location was previously identified as a site with significant endocortical and intra-cortical remodeling in lactation/low Ca animals [7]. Reconstructed regions were analyzed using a lower and upper threshold value of 270 and 1000, which corresponds to 617 and 3,000 mg hydroxyapatite per cm^3 (mgHA/ cm^3). Specific μ CT variables measured included total area, marrow area, cortical area, and cortical porosity.

Cortical porosity was classified into three distinct variables; total cortical porosity, osteonal porosity, and lacunar porosity. Total cortical porosity is reported using conventional analysis and nomenclature [10]. The total pore volume within the size range of secondary osteons, was determined by obtaining the total volume of porosity measured that had pore diameters consistent with secondary osteons, 30–60 μ m, and normalizing by the total cortical bone volume analyzed. The chosen diameter range of 30 to 60 μ m is consistent with histological characterization of the size of secondary osteons [11,12] in rats and is a marker of intra-cortical remodeling. By definition, secondary bone is formed in place of previously resorbed bone [13]. Therefore, pores on the size of osteons were referred to as secondary osteons because in our previous publication we demonstrated that the osteons were formed in places

where cortical bone had been resorbed and further, they presented with cement lines and concentric lamellae [7].

The average lacunar volume was determined by quantifying the total pore volume for pores with a diameter between 8 and 20 μm and normalizing by the total cortical bone volume analyzed. The average diameter of osteonal and lacunar porosity was determined from porosity size distributions of pore diameter vs the number of voxels of porosity within that diameter range. The average diameter was calculated as the weighted mean of the distribution gated to the diameter range of lacunae (8–20 μm) and osteons (30–60 μm).

Section Preparation.

After μCT scanning, femurs were dehydrated in a graded series of alcohol solutions and embedded in an epoxy resin (EpoThin, Buehler). 1 mm thick slabs at the femoral ROI were cut from the embedded specimens (Isomet 5000, Buehler) and fixed to plastic slides. The slabs were ground to a thickness of roughly 500 μm and polished to a mirror finish (Phoenix 4000, Buehler), using a series of colloidal diamond suspensions (9, 3, 1 μm average particle size, Metadi, Buehler) with ethylene glycol as a lubricant [14].

Dynamic Histomorphometry.

Dynamic histomorphometry was performed at the endocortical surface of the distal femoral diaphysis using fluorescent microscopy (Nikon Eclipse 80i) and commercial software (OsteoMeasure, OsteoMetrics). The primary outcome measures were the mineralizing surface per bone surface (MS/BS), mineral apposition rate (MAR), and bone formation rate per bone surface (BFR/BS). The MAR and BRR/BS parameters are dependent on double labels, and therefore it was not possible to assess these parameters in the animals sacrificed at days 0 and 1 of recovery, as these animals only received one calcein injection. MS/BS was determined using both single and double labels ($\text{MS/BS} = \text{double label} + \frac{1}{2} \text{single label}$) [15]. In cases where double labels were not clearly present, an imputed MAR value of 0.3 was assumed so that BFR/BS could be calculated.

Statistics.

The study was designed with the intention of running two-way analysis of variance (ANOVA) tests to assess the effects of group (virgin/normal Ca, lactation/normal Ca, and lactation/low Ca) and time (length of time post-weaning or recovery) and to explore interactions between group and time. To ensure that the ANOVA assumptions were met, we first tested the normality of the residuals from the two-way ANOVA output using the Shapiro-Wilk test. All of the μCT variables (cortical area, medullary area, total area, and cortical porosity) were normally distributed, justifying the use of the ANOVAs, followed by post-hoc comparisons using Tukey's HSD. Bone formation (MS/BS, MAR, and BFR/BS) and mineral metabolism measurements (ionic Ca, PTH, PTHrP, calcitonin, phosphate, and FGF23 concentrations) were found to be non-normally distributed. Therefore non-parametric test were used. Group and time effects were tested separately using the Kruskal-Wallis test followed by post-hoc between-group comparisons using Dunn's test. Finally, non-parametric Spearman's correlation analysis was performed to test the associations between mineral metabolism marker levels and indices of bone formation.

Results

Cortical area was reduced in both lactation groups compared to the virgin/normal Ca controls at each time point during recovery (Fig 1a). From day 0 to day 14, both lactation groups had increases in cortical area, leading to a significant overall time effect. However, there was no change in cortical area in the virgin/normal Ca controls during the recovery process, leading to a significant statistical interaction term between group and time, as the three groups demonstrated different trends over time. There were consistent between-group differences for medullary area, with significantly greater values in both lactation groups compared to the virgin/normal Ca controls at each time point (Fig. 1b). There were no significant time or interaction effects for the medullary area. Dietary Ca had less effect on total area (Fig. 1c), although the lactation/low Ca group had lower values than the virgin/normal Ca group at days 0, 1 and 2 and the lactation/normal Ca group at day 1. Similarly, there were no significant time or interaction effects for the total area.

Cortical porosity was elevated in the lactation/low Ca group compared to the other two groups at days 2 and 14, but did not change over time (Fig. 2a). The lactation/low Ca group had elevated osteonal-sized porosity compared to the lactation/normal Ca group at days 1, 7 and 14 (Fig. 2b). Osteonal porosity decreased in the lactation/low Ca group over time, but remained relatively constant in the other two groups. There was a significant time effect in the osteonal diameter with both lactation groups showing decreasing diameter over the recovery period (Fig. 2c). When the porosity data were gated to reflect lacunar-sized pores, both lactation groups had lower lacunar porosity compared to the virgin/normal controls at days 1 and 2 (Fig. 2d). By day 14, the lactation/low Ca lacunar porosity was significantly greater than the lactation/normal Ca or virgin/normal Ca groups. The average lacunar diameter was consistently greater in the lactation/low Ca group (Fig. 2e).

Bone formation indices were assessed at the endocortical surface of the distal femoral diaphysis and demonstrated a high degree of variability, necessitating the use of non-parametric tests. MAR was significantly elevated in the lactation/low Ca group compared to the virgin/normal Ca controls at day 14 (Fig. 3a). MS/BS was elevated in the lactation/low Ca group, reaching significance at day 7 (Fig 3b). Endocortical BFR/BS was elevated in the lactation/low Ca group compared to the lactation/normal Ca at day 7 (Fig. 3c).

Metabolic biomarkers of Ca metabolism were largely unaffected by lactation or the amount of Ca in the diet (Fig. 4). Circulating ionic Ca levels varied over time in the virgin/normal Ca and the lactation/low Ca groups, with a rapid decline on day 1, followed by a stabilization of the calcium levels. The lactation/normal Ca group did not show a significant time effect. At day 0, there were no between-group differences in the level of circulating ionic calcium, however at day 1 Ca levels were slightly elevated in the lactation/low Ca compared to the virgin/normal Ca group. There were no between-group differences at any of the other time points measured.

Circulating PTH levels did not vary over time for any of the groups, nor were there any between-group differences (Fig. 4b). The PTHrP levels in all three groups increased over time (Fig. 4c), however due to high within-group variance, the only significant time effect

was in the lactation/normal Ca group ($p=0.006$). Calcitonin levels did not change over time, nor were there any between-group differences (Fig. 4d).

Circulating phosphate metabolism markers were impacted by lactation, but not by dietary Ca (Fig. 5). Phosphate varied over time in all three groups (Fig. 5a). In both lactation groups the phosphate levels were reduced compared to the virgin/normal Ca controls at day 0, but were elevated at days 1 and 2, with no between-group differences in phosphate at days 7 and 14.

Circulating full length FGF23 levels demonstrated a high degree of within-group variability. Despite this variability, there was a significant time effect in the lactation/normal Ca group, with the other two groups failing to reach significance (Fig. 5b). The lactation/low Ca group had significantly reduced FGF23 at day 0 compared to both the virgin/normal Ca controls and the lactation/normal Ca group. However, by FGF23 levels in both lactation groups were significantly elevated compared to the virgin/normal Ca controls at days 1, 2 and 14, although the levels by day 14 were much more similar to the virgin/normal Ca control levels than at the preceding time points.

Correlations between biomarker levels and the bone formation measurements were significant for PTHrP level and both the mineralizing surface per bone surface (Spearman's $\rho=0.273$, $p\text{-value}=0.042$) and the bone formation rate ($\rho=0.298$, $p=0.026$, Supplemental Table 1). Phosphate levels were negatively correlated with the mineralizing surface ($\rho=-0.329$, $p=0.014$) and nearly significantly associated with the bone formation rate ($\rho=-0.258$, $p=0.057$). None of the other biomarkers had significant correlations with the bone formation variables.

Discussion

To accommodate the high calcium demands of the suckling pups, lactating dams upregulate skeletal resorption leading to loss of bone mass [6,16–27]. After weaning, the dams experience a highly anabolic bone formation period wherein the lost bone mass is largely recovered [5,22,28,29]. In the current study we examined this post-weaning anabolic period by focusing on the cortical bone area and porosity, endocortical bone formation kinetics and mineral metabolism markers as a function of dietary Ca during lactation. Contrary to our hypothesis, the post-weaning bone formation kinetics and markers of mineral metabolism were largely unaffected by the amount of dietary Ca during lactation.

Similar to our previous study, at the end of lactation we found reduced cortical area in the lactation/low Ca animals, which is primarily driven by an increase in the medullary area [7]. The increased medullary area is due to elevated endocortical remodeling, which is known to occur during lactation in rats [17,22,30]. Surprisingly, the extent of cortical bone loss was not affected by dietary calcium during lactation, which is seemingly contrary to previously published results [31]. de Winter et al. [31] reported significantly reduced cortical bone area in lactating dams fed a low calcium diet over 7, 14, and 21 day lactation periods compared to those fed adequate calcium. It is unclear why these two studies had different findings, although it is worth noting that de Winter et al. measured cortical geometry at the midshaft, while in the current study we evaluated the distal femoral diaphysis due to our previous

finding that this location is better-suited for studying both endocortical and intra-cortical remodeling [7].

Although cortical area was not significantly different between the two lactation groups, animals in the lactation/low Ca group had significantly increased cortical porosity compared to the lactation/normal Ca animals. The increase in cortical porosity can be attributed to the development of more secondary osteons due to intra-cortical remodeling based on our gating of pore size, which is consistent with our previous experiment using the lactation/low Ca model [7]. Interestingly, the porosity levels in the lactation/normal Ca group were not different from the virgin/normal Ca controls. Hagaman et al. similarly found increased cortical porosity in rats fed a low calcium diet during lactation, but no change in those fed adequate Ca during lactation [30]. Combined, these studies indicate that lactation alone primarily induces endocortical bone resorption, while the additional challenge of low dietary Ca during lactation triggers intra-cortical remodeling.

Osteocytic osteolysis is thought to play an important role in the maintenance of Ca levels during lactation [32,33]. In the current study, the average lacunar diameter was slightly elevated in the lactation/low Ca group, which is consistent with the presence of osteocytic osteolysis. Osteocytic osteolysis is a process that has been described in mouse models of lactation [32,34,35], but to our knowledge there have been no published reports in the rat model.

The total fraction of porosity characteristic of lacunae was surprisingly lower in the lactation groups compared to the virgin controls. Although the current mechanism is unknown, it is possible that resorption in the lactation model targets regions of particularly high osteocyte density, leading to overall lower lacunar porosity.

Maternal mineral metabolism during pregnancy, lactation, and weaning remains a complex field of study with some conflicting findings in animal models [1,8]. One of the mineral metabolism adaptations thought to be required to upregulate skeletal remodeling during lactation is increased production of PTH [18,19,25,36–39]. Yet, our study found that PTH levels were not different between groups at the point of weaning. Despite the rapid bone formation following weaning, PTH levels have been reported to rapidly return to control levels following weaning [40,41], which is consistent with our findings.

PTHrP levels are elevated during lactation and are likely involved in the increased skeletal remodeling [42]. The response of PTHrP in rodents following weaning is not well-studied. Lactating women have a significant reduction in PTHrP levels 4 months after cessation of breast feeding [43]. Therefore, it is not clear how rapidly PTHrP levels respond to weaning. The current study suggests that there is an increase in PTHrP levels during the first 14 days after weaning. It is possible that PTHrP is involved in post-weaning bone accrual, as PTHrP signaling in osteoblasts has been shown to increase during bone formation [44,45]. Indeed, circulating PTHrP levels were found to be significantly correlated with the bone formation rate and mineralizing surface per bone surface.

Circulating phosphate has previously been shown to increase following abrupt weaning [46,47], which is consistent with the current study. Despite the transient increase, phosphate

metabolism was normalized by 14 days following weaning in the present study. The pattern of FGF23 levels in our study following weaning is consistent with the role of FGF23 in the regulation of phosphate [48,49]. Specifically, as the phosphate levels rose during the post-weaning recovery phase, FGF23 levels increased to promote phosphate excretion in the kidney, thereby normalizing the circulating phosphate levels.

We only examined the first 14 days of the post-weaning recovery period so we could focus on the early changes in mineral metabolism and bone formation. From the μ CT data, it is apparent that the rats had not fully recovered bone mass by day 14, and therefore it is possible that several of the biomarkers, such as PTHrP and FGF23, would show additional dynamic behavior in longer follow-up periods.

Although there were no obvious effects on mineral metabolism in animals that developed mastitis, we cannot fully rule out effects of this localized inflammation. Mastitis is a relatively common occurrence in lactating animals, particularly in dairy cows [50], and in the current study was likely caused by a relatively large litter size (12 pups per dam). As mastitis was only evident upon the second day post-weaning, it is also possible that the abrupt weaning strategy contributed to its development. Abrupt weaning was chosen as a method to capture the rapid transition from resorption to formation and has been used in previous studies [46,51], but it is possible that lengthening the weaning period would have prevented mastitis development.

In the current study we investigated the effects of dietary Ca restriction during lactation on post-weaning bone formation and mineral metabolism. Surprisingly, dietary Ca restriction during lactation did not affect the extent of cortical bone loss, although the combination of lactation and a low Ca diet induced more intra-cortical remodeling than lactation alone. Similarly, dietary calcium during lactation had no effect on post-weaning bone formation, Ca metabolism, or phosphate metabolism. Regardless of dietary calcium, lactation caused transiently elevated phosphate and FGF23 levels post-weaning, which trended toward virgin/normal Ca levels over the 14 day recovery period. We conclude that the recovery process following lactation is largely insensitive to the concentration of dietary Ca provided during lactation and provides further justification for the use of the lactation/low Ca model as a means to investigate matrix maturation during intra-cortical and endocortical remodeling in the adult skeleton.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

The authors would like to thank Meghan Moran, Brittany Wilson, and Maleeha Mashiatulla for their help with the animal husbandry and tissue collections. Micro-Computed Tomography was performed at the Rush University microCT/Histology Core. Research reported in this publication was supported by the National Institute of Arthritis and Musculoskeletal and Skin Diseases of the National Institutes of Health under Award Number R21AR065604. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

1. Kovacs CS (2016) Maternal Mineral and Bone Metabolism During Pregnancy, Lactation, and Post-Weaning Recovery. *Physiol Rev* 96:449–547 doi:10.1152/physrev.00027.2015 [PubMed: 26887676]
2. Kovacs CS (2005) Calcium and bone metabolism during pregnancy and lactation. *J Mammary Gland Biol Neoplasia* 10:105–18
3. Brommage R, DeLuca HF (1985) Regulation of bone mineral loss during lactation. *Am J Physiol* 248:E182–E87
4. Ellinger GM, Duckworth J, Dalgarno AC, Quenouille MH (1952) Skeletal changes during pregnancy and lactation in the rat: effect of different levels of dietary calcium. *British Journal of Nutrition* 6:235–53 [PubMed: 12978211]
5. Bowman BM, Siska CC, Miller SC (2002) Greatly increased cancellous bone formation with rapid improvements in bone structure in the rat maternal skeleton after lactation. *J Bone Miner Res* 17:1954–60
6. Miller SC, Bowman BM (2004) Rapid improvements in cortical bone dynamics and structure after lactation in established breeder rats. *Anat Rec A Discov Mol Cell Evol Biol* 276:143–49
7. Ross RD, Sumner DR (2017) Bone Matrix Maturation in a Rat Model of Intra-Cortical Bone Remodeling. *Calcif Tissue Int* 101:193–203 doi:10.1007/s00223-017-0270-7 [PubMed: 28374176]
8. Kovacs CS (2011) Calcium and bone metabolism disorders during pregnancy and lactation. *Endocrinol Metab Clin North Am* 40:795–826
9. Kovacs CS, Kronenberg HM (1997) Maternal-fetal calcium and bone metabolism during pregnancy, puerperium, and lactation. *Endocr Rev* 18:832–72
10. Bouxsein ML, Boyd SK, Christiansen BA, Guldberg RE, Jepsen KJ, Muller R (2010) Guidelines for assessment of bone microstructure in rodents using micro-computed tomography. *Journal of Bone and Mineral Research* 25:1468–86 [PubMed: 20533309]
11. Duranova H, Martiniakova M, Omelka R, Grosskopf B, Bobonova I, Toman R (2014) Changes in compact bone microstructure of rats subchronically exposed to cadmium. *Acta Vet Scand* 56:64 doi:10.1186/s13028-014-0064-0 [PubMed: 25279860]
12. Martiniakova M, Chovancova H, Omelka R, Grosskopf B, Toman R (2011) Effects of a single intraperitoneal administration of cadmium on femoral bone structure in male rats. *Acta Vet Scand* 53:49 doi:10.1186/1751-0147-53-49 [PubMed: 21884588]
13. Burr DB, Akkus O (2014) Chapter 1 - Bone Morphology and Organization. In: Burr DB, Allen MR (eds) *Basic and Applied Bone Biology* Academic Press, San Diego, pp 3–25 doi:10.1016/B978-0-12-416015-6.00001-0
14. Donnelly E, Baker SP, Boskey AL, van der Meulen MC (2006) Effects of surface roughness and maximum load on the mechanical properties of cancellous bone measured by nanoindentation. *J Biomed Mater Res A* 77:426–35
15. Parfitt AM, Drezner MK, Glorieux FH, Kanis JA, Malluche H, Meunier PJ, Ott SM, Recker RR (1987) Bone histomorphometry: standardization of nomenclature, symbols, and units. *Journal of Bone and Mineral Research* 2:595–610 [PubMed: 3455637]
16. Ellinger GM, Duckworth J, Dalgarno AC, Quenouille MH (1952) Skeletal changes during pregnancy and lactation in the rat: effect of different levels of dietary calcium. *British Journal of Nutrition* 6:235–53 [PubMed: 12978211]
17. Miller SC, Shupe JG, Redd EH, Miller MA, Omura TH (1986) Changes in bone mineral and bone formation rates during pregnancy and lactation in rats. *Bone* 7:283–87 [PubMed: 3768208]
18. Garner SC, Peng TC, Hirsch PF, Boass A, Toverud SU (1987) Increase in serum parathyroid hormone concentration in the lactating rat: effects of dietary calcium and lactational intensity. *J Bone Miner Res* 2:347–52 doi:10.1002/jbmr.5650020412 [PubMed: 3455618]
19. Gonen E, Sahin I, Ozbek M, Kovalak E, Yologlu S, Ates Y (2005) Effects of pregnancy and lactation on bone mineral density, and their relation to the serum calcium, phosphorus, calcitonin and parathyroid hormone levels in rats. *J Endocrinol Invest* 28:322–26
20. Hirsch PF, Hagaman JR (1986) Reduced bone mass in calcitonin-deficient rats whether lactating or not. *J Bone Miner Res* 1:199–206 doi:10.1002/jbmr.5650010206 [PubMed: 3503538]

21. Honda A, Kurabayashi T, Yahata T, Tomita M, Matsushita H, Takakuwa K, Tanaka K (2000) Effects of pregnancy and lactation on trabecular bone and marrow adipocytes in rats. *Calcif Tissue Int* 67:367–72 [PubMed: 11136534]
22. Miller SC, Halloran BP, DeLuca HF, Jee WS (1982) Role of vitamin D in maternal skeletal changes during pregnancy and lactation: a histomorphometric study. *Calcif Tissue Int* 34:245–52 [PubMed: 6809285]
23. Peng TC, Garner SC, Kusy RP, Hirsch PF (1988) Effect of number of suckling pups and dietary calcium on bone mineral content and mechanical properties of femurs of lactating rats. *Bone & Mineral* 3:293–304 [PubMed: 3228614]
24. Rasmussen P (1977) Calcium deficiency, pregnancy, and lactation in rats. Microscopic and microradiographic observations on bones. *CalcifTissue Res* 23:95–102
25. Tojo Y, Kurabayashi T, Honda A, Yamamoto Y, Yahata T, Takakuwa K, Tanaka K (1998) Bone structural and metabolic changes at the end of pregnancy and lactation in rats. *Am J ObstetGynecol* 178:180–85
26. Warnock GM, Duckworth J (1944) Changes in the skeleton during gestation and lactation in the rat. *Biochem J* 38:220–4 [PubMed: 16747783]
27. Wong KM, Singer L, Ophaug RH (1980) Metabolic aspects of bone resorption in calcium-deficient lactating rats. *CalcifTissue Int* 32:213–19
28. Vajda EG, Bowman BM, Miller SC (2001) Cancellous and cortical bone mechanical properties and tissue dynamics during pregnancy, lactation, and postlactation in the rat. *BiolReprod* 65:689–95
29. Bowman BM, Miller SC (1999) Skeletal mass, chemistry, and growth during and after multiple reproductive cycles in the rat. *Bone* 25:553–59 [PubMed: 10574575]
30. Chen H, Hayakawa D, Emura S, Ozawa Y, Okumura T, Shoumura S (2002) Effect of low or high dietary calcium on the morphology of the rat femur. *Histology & Histopathology* 17:1129–35 [PubMed: 12371141]
31. de Winter FR, Steendijk R (1975) The effect of a low-calcium diet in lactating rats; observations on the rapid development and repair of osteoporosis. *CalcifTissue Res* 17:303–16
32. Qing H, Ardeshirpour L, Pajevic PD, Dusevich V, Jahn K, Kato S, Wysolmerski J, Bonewald LF (2012) Demonstration of osteocytic perilacunar/canalicular remodeling in mice during lactation. *JBone MinerRes* 27:1018–29
33. Wysolmerski JJ (2013) Osteocytes remove and replace perilacunar mineral during reproductive cycles. *Bone* 54:230–6 doi:10.1016/j.bone.2013.01.025 [PubMed: 23352996]
34. Nango N, Kubota S, Hasegawa T, Yashiro W, Momose A, Matsuo K (2016) Osteocyte-directed bone demineralization along canaliculi. *Bone* 84:279–88 doi:10.1016/j.bone.2015.12.006 [PubMed: 26709236]
35. Kaya S, Basta-Pljakic J, Seref-Ferlengez Z, Majeska RJ, Cardoso L, Bromage TG, Zhang Q, Flach CR, Mendelsohn R, Yakar S, Fritton SP, Schaffler MB (2017) Lactation-Induced Changes in the Volume of Osteocyte Lacunar-Canalicular Space Alter Mechanical Properties in Cortical Bone Tissue. *J Bone Miner Res* 32:688–97 doi:10.1002/jbmr.3044 [PubMed: 27859586]
36. Anderson JJ, Garner SC, Mar MH, Boass A, Toverud SU, Parikh I (1990) The ovariectomized, lactating rat as an experimental model for osteopenia: calcium metabolism and bone changes. *Bone & Mineral* 11:43–53 [PubMed: 2268738]
37. Boass A, Garner SC, Schultz VL, Toverud SU (1997) Regulation of serum calcitriol by serum ionized calcium in rats during pregnancy and lactation. *J Bone MinerRes* 12:909–14
38. Garner SC, Boass A, Toverud SU (1989) Hypercalcemia fails to suppress elevated serum parathyroid hormone concentrations during lactation in rats. *J Bone MinerRes* 4:577–83
39. Miller SC, Bowman BM (2007) Rapid inactivation and apoptosis of osteoclasts in the maternal skeleton during the bone remodeling reversal at the end of lactation. *AnatRec(Hoboken)* 290:65–73
40. Pike JW, Parker JB, Haussler MR, Boass A, Toverud SV (1979) Dynamic changes in circulating 1,25-dihydroxyvitamin D during reproduction in rats. *Science* 204:1427–9 [PubMed: 451573]
41. Garner SC, Peng TC, Toverud SU (1988) Modulation of serum parathyroid hormone and ionized calcium concentrations during reproduction in rats fed a low calcium diet. *J Bone MinerRes* 3:319–23

42. Sowers MF, Hollis BW, Shapiro B, Randolph J, Janney CA, Zhang D, Schork A, Crutchfield M, Stanczyk F, Russell-Aulet M (1996) Elevated parathyroid hormone-related peptide associated with lactation and bone density loss. *Jama* 276:549–54 [PubMed: 8709404]
43. Lippuner K, Zehnder HJ, Casez JP, Takkinen R, Jaeger P (1996) PTH-related protein is released into the mother's bloodstream during lactation: evidence for beneficial effects on maternal calcium-phosphate metabolism. *J Bone Miner Res* 11:1394–9 doi:10.1002/jbmr.5650111004 [PubMed: 8889837]
44. Datta NS, Abou-Samra AB (2009) PTH and PTHrP signaling in osteoblasts. *Cell Signal* 21:1245–54 doi:10.1016/j.cellsig.2009.02.012 [PubMed: 19249350]
45. Kovacs CS (2014) The Role of PTHrP in Regulating Mineral Metabolism During Pregnancy, Lactation, and Fetal/Neonatal Development. *Clinical Reviews in Bone and Mineral Metabolism* 12:142–64
46. Blahosova A, Neradilova M, Velicky J, Titlbach M, Marsikova L, Reisenauer R (1974) Dynamics of changes of calcium and phosphorus metabolism in relation to the morphology of parafollicular thyroid cells in rats during lactation and forced weaning. *Endokrinologie* 63:122–36 [PubMed: 4474077]
47. Garner SC, Boass A, Toverud SU (1990) Parathyroid hormone is not required for normal milk composition or secretion or lactation-associated bone loss in normocalcemic rats. *J Bone Miner Res* 5:69–75
48. Martin A, David V, Quarles LD (2012) Regulation and function of the FGF23/klotho endocrine pathways. *Physiol Rev* 92:131–55 doi:10.1152/physrev.00002.2011 [PubMed: 22298654]
49. Razzaque MS (2009) The FGF23-Klotho axis: endocrine regulation of phosphate homeostasis. *Nat Rev Endocrinol* 5:611–9 doi:10.1038/nrendo.2009.196 [PubMed: 19844248]
50. Kumar N, Manimaran A, Kumaresan A, Jeyakumar S, Sreela L, Mooventhan P, Sivaram M (2017) Mastitis effects on reproductive performance in dairy cattle: a review. *Trop Anim Health Prod* 49:663–73 doi:10.1007/s11250-017-1253-4 [PubMed: 28283873]
51. Hodnett DW, DeLuca HF, Jorgensen NA (1992) Intestine, bone, and mammary gland contributions to maternal plasma calcium increase after abrupt weaning. *Proc Soc Exp Biol Med* 199:332–6 [PubMed: 1539036]

Fig. 1.

μ CT-derived parameters of cortical geometry measured at the distal femoral diaphysis during the post-weaning recovery period. (a) Cortical area, (b) medullary area, and (c) total area are presented for the virgin/normal Ca (open circles), lactation/normal Ca (gray square), and lactation/low Ca (black triangle) as individual animals. Results from the two-way ANOVA are presented in the legends, with between-group post-hoc differences ($p < 0.05$) indicated by horizontal bars.

Fig. 2.

μCT-derived parameters of cortical porosity measured at the distal femoral diaphysis. (a) Cortical porosity, (b) osteonal pore volume (between 30–60 μm diameter), and (c) lacunar pore volume (between 8–20 μm diameter) are presented for the virgin/normal Ca (open circles), lactation/normal Ca (gray square), and lactation/low Ca (black triangle) as individual animals. Results from the two-way ANOVA are presented in the legends, with between group post-hoc differences ($p < 0.05$) indicated by horizontal bars.

Fig. 3.

Dynamic histomorphometric parameters of bone formation at the endocortical surface. The (a) bone formation rate per bone surface (BFR/BS), (b) mineral apposition rate (MAR), and (c) mineralizing surface per bone surface (MS/BS) are presented for the virgin/normal Ca (open circles), lactation/normal Ca (gray square), and lactation/low Ca (black triangle) as individual animals. Time effects for each group (Krusal-Wallis) are presented in the legends of each figure. Between group differences at each time-point were assessed using a non-parametric Mann-Whitney U test and significance ($p<0.05$) are indicated with a bracket above the corresponding time-point.

Fig. 4.

Calcium metabolism measured after weaning in the recovery phase. The circulating levels of (a) ionic Ca, (b) calcitonin, (c) PTH, and (d) PTHrP are presented in the virgin/normal Ca (open circles), lactation/normal Ca (gray square), and lactation/low Ca (black triangle) as individual animals. Time effects for each group (Krusal-Wallis) are presented in the legends of each figure. Between group differences at each time-point were assessed using a non-parametric Mann-Whitney U test and significance ($p<0.05$) are indicated with a bracket above the corresponding time-point.

Fig. 5.

Phosphate metabolism measured after weaning in the recovery phase. The circulating levels of (a) phosphate, and (b) intact FGF23 are presented in the virgin/normal Ca (open circles), lactation/normal Ca (gray square), and lactation/low Ca (black triangle) as individual animals. Time effects for each group (Krusal-Wallis) are presented in the legends of each figure. Between group differences at each time-point were assessed using a non-parametric Mann-Whitney U test and significance ($p<0.05$) are indicated with a bracket above the corresponding time-point.

Table 1:

Study Design

Group	Induction (Lactation)	Recovery (days)				
		0	1	2	7	14
Virgin/Normal Ca	25	5	5	5	5	5
Lactation/Normal Ca	25	5	5	4	5	5
Lactation/Low Ca	25	5	5	5	4	5

The table values in the shaded cells represent the final group sizes after animal losses described in the Animal Husbandry section.