Research

Cite this article: Schubert BW *et al.* 2019 Yucatán carnivorans shed light on the Great American Biotic Interchange. *Biol. Lett.* **15**: 20190148.
<http://dx.doi.org/10.1098/rsbl.2019.0148>

Received: 26 February 2019
Accepted: 8 April 2019

Subject Areas:

palaeontology, biogeography, evolution, taxonomy and systematics

Keywords:

Arctotherium, *Procyon*, Yucatán, GABI, late Pleistocene

Author for correspondence:

Blaine W. Schubert
e-mail: schubert@etsu.edu

Electronic supplementary material is available online at <https://dx.doi.org/10.6084/m9.figshare.c.4472486>.

Palaeontology

Yucatán carnivorans shed light on the Great American Biotic Interchange

Blaine W. Schubert¹, James C. Chatters², Joaquin Arroyo-Cabres³, Joshua X. Samuels¹, Leopoldo H. Soibelzon⁵, Francisco J. Prevosti^{6,7}, Christopher Widga¹, Alberto Nava⁸, Dominique Rissolo⁹ and Pilar Luna Erreguerena⁴

¹Center of Excellence in Paleontology and Department of Geosciences, East Tennessee State University (ETSU), Johnson City, TN 37614, USA

²Applied Paleoscience and DirectAMS, 10322 NE 190th Street, Bothell, WA 98011, USA

³Laboratorio de Arqueozoología, Subdirección de Laboratorios y Apoyo Académico, and ⁴Subdirección de Arqueología Subacuática, INAH, Moneda 16, Col. Centro, CdMx 06060, Mexico

⁵División Paleontología de Vertebrados, Museo de La Plata, 1900 La Plata, Argentina

⁶Centro Regional de Investigaciones Científicas y Transferencia Tecnológica de La Rioja (CRILAR), Provincia de La Rioja, UNLaR, SEGEMAR, UNCa, CONICET, Entre Ríos y Mendoza s/n, 5301 – Anillaco, La Rioja, Argentina

⁷Departamento de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de La Rioja (UNLaR), Av. Luis M. de la Fuente S/N, 5300 La Rioja, Argentina

⁸Bay Area Underwater Explorers, Berkeley, CA, USA

⁹Qualcomm Institute, University of California, San Diego, CA, USA

BWS, 0000-0002-6061-1677

The Great American Biotic Interchange is considered to be a punctuated process, primarily occurring during four major pulses that began approximately 2.5 Ma. Central America and southeastern Mexico have a poor fossil record of this dynamic faunal history due to tropical climates. Exploration of submerged caves in the Yucatán, particularly the natural trap Hoyo Negro, is exposing a rich and remarkably well-preserved late Pleistocene fauna. Radiometric dates on megafauna range from approximately 38 400–12 850 cal BP, and extinct species include the ursid *Arctotherium wingei* and canid *Procyon troglodytes*. Both genera were previously thought to be indigenous to and confined to South America and appear to represent an instance of large placental mammals, descended from North American progenitors, migrating back north across the Panama Isthmus. This discovery expands the distribution of these carnivorans greater than 2000 km outside South America. Their presence along with a diverse sloth assemblage suggests a more complex history of these organisms in Middle America. We suggest that landscape and ecological changes caused by latest Pleistocene glaciation supported an interchange pulse that included *A. wingei*, *P. troglodytes* and *Homo sapiens*.

1. Introduction

North and South America were geographically separated for most of the Cenozoic, resulting in distinct vertebrate communities. Faunal interchange between the continents began in the late Cenozoic, and most dispersals occurred after the Panama Isthmus developed. This biogeographic phenomenon, known as the Great American Biotic Interchange (GABI), went in both directions [1–6] and enabled the expansion of placental mammals to South America, including the focus of this report, ursids and canids. Recently, researchers hypothesized the isthmus developed approximately 10 Myr earlier than previously thought, 13–15 Ma [7], well before the first large-scale faunal interchange around 2.5 Ma [5], suggesting a lag between continental connection and extensive migration [8].

Figure 1. Specimen recovery in Hoyo Negro and map showing locations. (a) DP 5864 *Arctotherium wingei* cranium being recovered. (b) DP 5865 *A. wingei* cranium after overlying plant material was brushed away. *Tremarctos ornatus* distribution (in red) [22]. Photos by Roberto Chávez-Arce.

The concept of an earlier isthmus and interchange lag has been challenged, citing insufficient evidence for a well-developed isthmus prior to 3 Ma [9–11].

Fossil evidence of GABI is well represented in the United States and parts of Mexico but sparse in southeastern Mexico and Central America, hereafter referred to as Middle America, where tropical conditions are unfavourable for preservation [4,12–15]. Thus, a significant geographical and temporal gap exists in understanding the fauna in the interchange region itself. New discoveries from submerged caves in the Yucatán are changing this, bringing to light an underworld of exquisitely preserved fossils from the late Pleistocene, when sea level and the water table were significantly lower during glaciations. Publications on these underwater discoveries have focused on early human skeletons [16,17], new mammalian genera (two ground sloths and a peccary) [18–20] and a new species interpreted to be a jaguar-like cat [21].

Hoyo Negro (HN), a submerged pit inside the Sac Actun cave system, is the focus of this paper. The site is located on the eastern Yucatán Peninsula, Quintana Roo, Mexico (figure 1). Hoyo Negro represents a collapse karst feature accessible from three submerged horizontal passages with floors that are approximately 12 m below sea level (mbsl). Flooding history of the cave has been reconstructed, indicating the base of

HN was flooded by 9850 cal BP, and upper passages leading to HN were submerged by 8100 cal BP [23]. Hoyo Negro drops from approximately 12 mbsl at the rim to at least 55 mbsl at the floor. The HN pit is bell shaped, approximately 62 m wide at the base, and served as a natural trap for animals moving through the cave in the late Pleistocene. Mammals discovered on the surface of the HN floor include multiple ground sloth species (including the newly named *Nohochichak xibalbahkah*) [18], tapirs, sabertooth cats, cougars, gomphotheres, bears, canids and a relatively complete human skeleton that dates to terminal Pleistocene [16]. In addition, bones and trackways of extinct fauna are known from the upper passages.

Here we report two extinct carnivorans from HN, identified earlier as the bear *Tremarctos* and coyote *Canis latrans* [16]. Representative specimens have now been collected and we revise identifications to the ursid *Arctotherium wingei* and canid *Protocyon troglodytes*. Both genera and species were previously unknown outside South America.

2. Material and methods

Highly trained technical divers performed on-site photography, filming and collecting based on direction from researchers.

Collecting focused on diagnostic specimens, particularly well-preserved skulls. Our emphasis here is systematic identifications; more-detailed skeletal descriptions are in progress. Identifications were made based on current literature. The abbreviation 'DP' is for Departamento de Prehistoria, the older name for the collection at the Instituto Nacional de Antropología e Historia (INAH), México. See supplementary material for details on photography, collecting, preparation and radiometric dating, as well as another figure of the crania (electronic supplementary material, figure S1).

3. Results and discussion

(a) Age and chronology

An accelerator mass spectrometer (AMS) radiocarbon analysis on extracted protein from *Arctotherium* dentine (from DP 5865) resulted in an age of $11\,298 \pm 138$ rcy BP (D-AMS 1176911). When calibrated using Calib 7.1 [24] with intCal13 calibration dataset [25], the two-sigma age range is 12 850–13 430 cal BP. This date represents the only protein-based radiocarbon date from the site and first absolute date for *Arctotherium wingei*. While no dates on *Protcyon* have been successful, systematic dating attempts on this taxon and others from HN continue. Currently, the inclusive age range of HN large mammals is approximately 38 400–12 850 cal BP, with the oldest date on gomphothere [16].

(b) *Arctotherium*

With multiple individuals identified, the bear is the most abundant large mammal from HN. Individuals range in age from subadult to adult and represent the South American short-faced bear, *Arctotherium* (figure 1 and figure 2). The following combined characters diagnose the genus: notch absent on dorsal margin of the foramen magnum, the distance between C1 and P4 less than mesio-distal length of P4, P1–P3 compressed and arranged in a triangle, M1 metastyle reduced or absent [26]. Portions of two individuals are reported here, DP-5864 and DP-5865.

Arctotherium is part of Tremarctinae, an endemic American subfamily. Besides *Arctotherium*, this group includes *Plionarctos*, *Tremarctos* and *Arctodus*. *Plionarctos* is recorded from the late Miocene to early Pliocene and is only known from the United States [27]. *Tremarctos* appeared in the Pliocene of North America and now occurs only in South America [22,28]. *Arctodus* appeared in the late Pliocene, and by the late Pleistocene it occurred across the United States, much of Canada and portions of Mexico [29,30]. *Arctodus* and *Tremarctos* went extinct in North America near the end of the Pleistocene [31].

The oldest record of Ursidae in South America is early Pleistocene, with the appearance of *Arctotherium* at 0.98–1.76 Ma [32]. Ursids are considered to be part of GABI 2, a dispersal pulse that occurred around 1.8 Ma [5]. Five species of *Arctotherium* are recognized, *A. angustidens* (early–middle Pleistocene), *A. vetustum* (middle Pleistocene) and three middle–late Pleistocene species, *A. bonariense*, *A. tarijense* and *A. wingei* [26,32]. *Arctotherium angustidens* was particularly large [33], but over time this genus is represented by smaller forms, culminating in *A. wingei*, the smallest [32,34].

Disagreements have occurred over the evolutionary relationship of *Arctotherium*, *Tremarctos* and *Arctodus*, with earlier morphological interpretations placing *Arctotherium* and *Arctodus* as sister taxa [26]. Recent genetic analyses indicate

Arctotherium is more closely related to *Tremarctos*, and therefore *Arctotherium* and *Arctodus* share morphologies that may relate to their large size and/or dietary behaviour [35]. Based on our understanding of the group, *Arctotherium* arose from *Tremarctos*, or a common ancestor that excluded *Arctodus*.

The HN bear is identified as *A. wingei*, the only *Arctotherium* species known from northern South America, with occurrences in Bolivia, Brazil and Venezuela (figure 1) [32,36,37]. While partial *A. wingei* skulls and isolated teeth are reported from South America, the material is poorly represented. The remains from HN represent the best-preserved and most-extensive record of *Arctotherium* from any locality and they exhibit a high degree of intraspecific morphological variation. These specimens are assigned to *A. wingei* using a suite of characters: separate openings for foramen rotundum and alisphenoid canal on cranium, tympanic bulla inflated, greatest mesio-distal length of M1 slightly larger than greatest labio-lingual length, shape of M1 lingual border convex, p1–p3 arranged in a triangle, m2 entoconid with two apices [26].

In addition to HN fossils, other submerged caves of the Yucatán are producing photographic records of *Arctotherium*, and one specimen is known from a dry cave in Belize [38]. Further, a fragmentary distal humerus from another submerged Yucatán cave was recently described as a new species of cat, *Panthera balamoides* [21]. However, published images in that paper are morphologically similar to *Arctotherium*, and we feel this identification should be reassessed.

Arctotherium is the only known bear from the Yucatán Peninsula. In terms of diet, *A. wingei* has been interpreted as a herbivorous omnivore like *Tremarctos ornatus* [39,40]. Although *T. ornatus* occurs in the Andean region of western South America today, it is not known from that continent until the early Holocene [41–43]. While *Tremarctos* is known from North America until the Pleistocene/Holocene transition [44], there is only one fossil occurrence in Middle America [45]. Identification of this specimen is tenuous because the skeleton is in a Belizean cave. Based on the current record, we propose that competition with *Arctotherium* in Middle America may have limited the southern distribution of *Tremarctos* until the former became extinct in the latest Pleistocene or early Holocene.

(c) *Protcyon*

Canid material from HN represents the South American genus *Protcyon* (DP 5867, figure 2). The following combined characters distinguish the genus: robust skull, wide zygomatic arches and palate, short rostrum, paraoccipital process directed posteriorly, P4 large with small protocone, M1–M2 with reduced or absent hypocone, M2 reduced, lower premolars with acute and weak principal cusps, m1 without metaconid and entoconid [46–53].

Protcyon is part of Caninae, a large canid subfamily that originated in North America and dispersed into the Old World and South America. The subfamily occurs in South America by 2.5 Ma [43,52] and was part of GABI 1 [5]. By the early Pleistocene, South American Caninae were diverse, with multiple *Lycalopex* species, and the emergence of two genera thought to be endemic, *Theriodictis* and *Protcyon* [43,47,49,50,52,54]. *Protcyon* is considered to be closely related to *Theriodictis* [52,55,56], which was questionably reported from the late Pliocene or early Pleistocene of Florida as *Theriodictis? floridanus* [57]. No solid records of *Theriodictis* are known outside South America. Three *Protcyon* species

Figure 2. *Arctotherium wingei* and *Protocyon troglodytes* from Hoyo Negro. DP 5865 *A. wingei*, (a) dorsal and (b) ventral view of the cranium, and (c) occlusal view of right P4–M2. DP 5864 *A. wingei*, (d) occlusal view of right p4–m2 and (e) lateral view of right dentary; m3 broken and missing. DP 5867 *P. troglodytes* (f) dorsal and (g) ventral view of cranium, (h) occlusal view of right P3–M2, (i) occlusal view of right p3–m2 and (j) lateral view of the right dentary. (Online version in colour.)

are recognized, *P. scagliarum* (early–middle Pleistocene), *P. troglodytes* (middle?–late Pleistocene) and *P. orcesi* (late Pleistocene) [49,53,54].

The HN *Protocyon* is assigned to *P. troglodytes*, known from Argentina, Brazil, Bolivia, Ecuador, Uruguay and Venezuela [53] (figure 1). Northern records of *P. troglodytes* led researchers to speculate the possibility of their occurrence in Central America as well [51]. Other *Protocyon* species have only been recovered from their type localities, *P. scagliarum* in Argentina and *P. orcesi* from Ecuador. *Protocyon troglodytes* is distinguished from *P. scagliarum* based on the absence of an anterior cusplet on p4, and *P. orcesi* based on the presence of m3 [47,50,52]. Intraspecific variation may account for the difference between *P. troglodytes* and *P. orcesi* [49,50,53].

As part of the late Pleistocene Yucatán fauna, *Protocyon* would have been a hypercarnivorous large canid [53]. *Canis*

dirus, *C. lupus* and *C. latrans* are reported from Loltun Cave in northeast Yucatán, but *Protocyon* was not recovered [58,59]. Thus, the degree of geographical, temporal and ecological overlap of *Protocyon* and these canids is unclear.

(d) Interchange, biogeography and evolution

This report presents the first published records of *Arctotherium* and *Protocyon* outside South America, expanding their distributions over 2000 km out of South America. *Arctotherium* and *Protocyon* are considered to be endemic South American taxa that developed on that continent after their ursid and canid ancestors crossed the Panama Isthmus from Middle America [32,43,60]. While this scenario is supported by the occurrence of these carnivorans in South America, and lack of their fossil remains in Middle America, the record is biased by a relative

lack of fossil sites from the latter region. Thus, we cannot discount the possibility that *Arctotherium* or *Protcyon* occurred in Middle America at various times during the late Cenozoic. Further, it is also possible that Middle America played a role in the evolution of these genera, either through ongoing gene flow with South American populations or origination in Middle America.

These carnivorans, recent discoveries of new ground sloth genera from the Yucatán, and controversy about age estimates on the development of the Panama Isthmus 'bridge', all indicate we still have a lot to learn about GABI. Interpretations of the fossil record suggest four main pulses of interchange, GABI 1–4, at around 2.5 Ma, 1.8 Ma, 0.8 Ma and 0.125 Ma, respectively [5,6]. These pulses overlap with glacial episodes, sea-level lowering, widening of the isthmus and more-open habitats that would have created opportunities for dispersal of large mammals adapted to open habitats [5,13].

If the existing fossil record is an accurate representation of *Arctotherium* and *Protcyon* through time, and these genera developed strictly in South America, the Yucatán material represents a reversed dispersal of these families across the isthmus. Further, if this migration occurred during the latest Pleistocene, in line with existing radiometric dates, one possibility is they moved north during or since the last full glacial, between approximately 35 000 and 12 000 years ago. Because

this glacial pulse would have caused similar landscape changes and migration opportunities as GABI 1–4, this additional interchange could be referred to as GABI 5. Unlike earlier interchanges, 'GABI 5' involved humans and culminated with the mass extinction of most American megafauna, including *Arctotherium* and *Protcyon*.

Data accessibility. All specimens reported here (and in electronic supplementary material) are curated by INAH and available for study.

Authors' contributions. All authors made substantial contributions in data acquisition or interpretation, critically revised the paper for intellectual content, approve of the final version for publication and agree to be accountable for all aspects of the work.

Competing interests. We declare we have no competing interests.

Funding. Financial support provided by INAH, National Geographic Society (CP-036R-17), ETSU Center of Excellence in Paleontology, Archaeological Institute of America, DirectAMS and Strauss Family Fund. L.H.S. supported by ANPCyT (PICT 2016-2698) and F.J.P. by ANPCyT (PICT 2015-966) and Conicet (PUE 0125).

Acknowledgements. We thank divers Alejandro Álvarez, Roberto Chávez-Arce, Sam Meacham, Osama Gobar and Monica Torre, photographer Gibran Castillo, technicians Alejandro López-Jiménez and Salvador Estrada, preparators Keila Bredehoeft, Shawn Haugrud and Brian Compton, geoscientist Andrew Joyner and software developer Vid Petrovic. We are grateful to Roberto Junco, Adriana Velázquez Morlet and Falko Kuester for supporting the HN Project.

References

- Marshall LG, Webb SD, Sepkoski Jr JJ, Raup DM. 1982 Mammalian evolution and the Great American Interchange. *Science* **215**, 1351–1357. (doi:10.1126/science.215.4538.1351)
- Stehli F, Webb SD. 1985 *The Great American Biotic Interchange*. New York, NY: Plenum Press.
- Woodburne M, Cione AL, Tonni EP. 2006 Central American provincialism and the Great American Biotic Interchange. In *Advances in late Tertiary vertebrate paleontology in Mexico and the Great American Biotic Interchange*, vol. 4 (eds O Carranza-Castañeda, EH Lindsay), pp. 73–101. Juriquilla, Querétaro, Mexico: Instituto de Geología y Centro de Geociencias de la Universidad Nacional Autónoma de México.
- Morgan GS. 2008 Vertebrate fauna and geochronology of the Great American Biotic Interchange in North America. In *Neogene mammals*, vol. 44 (eds SG Lucas, GS Morgan, JA Spielman, DR Prothero), pp. 93–140. Albuquerque, NM: New Mexico Museum of Natural History and Science.
- Woodburne MO. 2010 The Great American Biotic Interchange: dispersals, tectonics, climate, sea level and holding pens. *J. Mamm. Evol.* **17**, 245–264. (doi:10.1007/s10914-010-9144-8)
- Cione AL, Gasparini GM, Soibelzon E, Soibelzon LH, Tonni EP. 2015 The GABI in southern South America. In *The Great American Biotic Interchange: a South American perspective* (eds AL Cione, GM Gasparini, E. Soibelzon, LH Soibelzon, EP Tonni), pp. 71–96. Amsterdam, The Netherlands: Springer.
- Montes C, Cardona A, Jaramillo C, Pardo-Trujillo A. 2015 Middle Miocene closure of the Central American Seaway. *Science* **348**, 226–229. (doi:10.1126/science.aaa2815)
- Bacon CD, Molnar P, Antonelli A, Crawford AJ, Montes C, Vellejo-Pareja MC. 2016 Quaternary glaciation and the Great American Biotic Interchange. *Geology* **44**, 375–378. (doi:10.1130/G37624.1)
- O'Dea A *et al.* 2016 Formation of the Isthmus of Panama. *Sci. Adv.* **2**, e1600883. (doi:10.1126/sciadv.1600883)
- O'Dea A *et al.* 2016 Building bridges. Response to Erkens and Hoorn: 'The Panama Isthmus, 'old', 'young' or both?' *Sci. Adv. e-Lett.* **2**, e1600883.
- O'Dea A *et al.* 2017 Formation of the Isthmus of Panama: response to Jaramillo *et al.* *Sci. Adv.* **3**, e1602321. (doi: 10.1126/sciadv.1602321)
- Webb SD. 1978 A history of savanna vertebrates in the New World. Part II: South America and the Great Interchange. *Annu. Rev. of Ecol. and Syst.* **9**, 393–426. (doi:10.1146/annurev.es.09.110178.002141)
- Webb SD. 1991 Ecography and the Great American Interchange. *Paleobiology* **17**, 266–280. (doi:10.1017/S0094837300010605)
- Webb SD. 1997 The Great American faunal interchange. In *Central America: a natural and cultural history* (ed. AG Coates), pp. 97–122. New Haven, CT: Yale University Press.
- Ferrusquía-Villafranca I, Arroyo-Cabrales J, Martínez-Hernández E, Gama-Castro J, Ruez-González J, Polaco OJ, Johnson E. 2010 Pleistocene mammals of Mexico: a critical review of regional chronofaunas, climate change response and biogeographic provinciality. *Quatern. Int.* **217**, 53–104. (doi:10.1016/j.quaint.2009.11.036)
- Chatters JC *et al.* 2014 Late Pleistocene human skeleton and mtDNA link Paleoamericans and modern Native Americans. *Science* **344**, 750–754. (doi:10.1126/science.1252619)
- Stinnesbeck W *et al.* 2017 The earliest settlers of Mesoamerica date back to the late Pleistocene. *PLoS ONE* **12**, 16–18. (doi:10.1371/journal.pone.0183345)
- McDonald HG, Chatters JC, Gaudin TJ. 2017 A new genus of megalonychid ground sloth (Mammalia, Xenarthra) from the late Pleistocene of Quintana Roo, Mexico. *J. Vertebr. Paleontol.* **37**, 14. (doi:10.1080/02724634.2017.1307206)
- Stinnesbeck SR *et al.* 2017 *Xibalaonyx oviceps*, a new megalonychid ground sloth (Folivora, Xenarthra) from the late Pleistocene of the Yucatán Peninsula, Mexico, and its paleobiogeographic significance. *Paleont. Z.* **91**, 245–271. (doi:10.1007/s12542-017-0349-5)
- Stinnesbeck SR *et al.* 2017 A new fossil peccary from the Pleistocene–Holocene boundary of the eastern Yucatán Peninsula, Mexico. *J. S. Am. Earth Sci.* **77**, 341–349. (doi:10.1016/j.jsames.2016.11.003)
- Stinnesbeck SR, Stinnesbeck W, Frey E, Avilés Olguín J, Rojas Sandoval C, Velázquez Morlet, González AH. 2018 *Panthera balamoides* and other Pleistocene felids from the submerged caves of Tulum, Quintana Roo, Mexico. *Hist. Biol.* 1–10. (doi:10.1080/08912963.2018.1556649)
- International Union for Conservation of Nature. 2017 IUCN Red list of threatened species, version 2017. See <http://www.iucnredlist.org>.
- Collins SV, Reinhardt EG, Rissolo D, Chatters JC, Nava Blank A, Luna Erreguerena P. 2015 Reconstructing

- water level in Hoyo Negro, Quintana Roo, Mexico, implications for early Paleoamerican and faunal access. *Quaternary Sci. Rev.* **124**, 68–83. (doi:10.1016/j.quascirev.2015.06.024)
24. Stuiver M, Reimer PJ. 1993 Extended ^{14}C data base and revised CALIB 3.0 ^{14}C age calibration program. *Radiocarbon* **35**, 215–230. (doi:10.1017/S0033822200013904)
 25. Reimer PJ *et al.* 2013 IntCal13 and MARINE13 radiocarbon age calibration curves 0–50 000 years calBP. *Radiocarbon* **55**, 1869–1887. (doi:10.2458/azu_js_rc.55.16947)
 26. Soibelzon LH. 2004 Revisión sistemática de los Tremarctinae (Carnivora, Ursidae) fósiles de América del Sur. *Revista Mus. Argent. Ci. Nat.* **6**, 107–133. (doi:10.22179/revmacn.6.75)
 27. Tedford RH, Martin J. 2001 *Plionarctos*, a tremarctine bear (Ursidae: Carnivora) from western North America. *J. Vertebr. Paleontol.* **21**, 311–321. (doi:10.1671/0272-4634(2001)021[0311:PATBUC]2.0.CO;2)
 28. Kurtén B. 1966 Pleistocene bears of North America: 1. Genus *Tremarctos*, spectacled bears. *Acta Zool. Fennica* **115**, 1–120.
 29. Richards RL, Churcher CS, Turnbull WD. 1996 Distribution and size variation in North American short-faced bears, *Arctodus simus*. In *Palaeoecology and palaeoenvironments of late cenozoic mammals: tributes to the career of C.S. (Rufus) Churcher* (eds KM Stewart, KL Seymour), pp. 191–246. Toronto, Canada: University of Toronto Press.
 30. Schubert BW, Hulbert Jr RC, MacFadden BJ, Searle M, Searle S. 2010 Giant short-faced bears (*Arctodus simus*) in Pleistocene Florida, a substantial range extension. *J. Paleontol.* **84**, 79–87. (doi:10.1666/09-113.1)
 31. Schubert BW. 2010 Late Quaternary chronology and extinction of North American short-faced bears (*Arctodus simus*). *Quatern. Int.* **217**, 188–194. (doi:10.1016/j.quaint.2009.11.010)
 32. Soibelzon LH, Tonni EP, Bond M. 2005 The fossil record of South American short-faced bears (Ursidae, Tremarctinae). *J. S. Am. Earth Sci.* **20**, 105–113. (doi:10.1016/j.jsames.2005.07.005)
 33. Soibelzon LH, Schubert BW. 2011 The largest known bear, *Arctotherium angustidens*, from the early Pleistocene Pampean region of Argentina: with a discussion of size and diet trends in bears. *J. Paleontol.* **85**, 69–75. (doi:10.1666/10-037.1)
 34. Soibelzon LH, Tarantini VB. 2009 Estimación de la masa corporal de las especies de osos fósiles y actuales (Ursidae, Tremarctinae) de América del Sur. *Rev. Mus. Argentino Cienc. Nat.* **11**, 243–254. (doi:10.22179/REVMACN.11.263)
 35. Mitchell KJ *et al.* 2016 Ancient mitochondrial DNA reveals convergent evolution of giant short-faced bears (Tremarctinae) in North and South America. *Biol. Lett.* **12**, 20160062. (doi.org/10.1098/rsbl.2016.0062)
 36. Trajano E, Ferrarezzi H. 1995 A fossil bear from northeastern Brazil, with a phylogenetic analysis of the South American extinct Tremarctinae (Ursidae). *J. Vertebr. Paleontol.* **14**, 552–561. (doi:10.1080/02724634.1995.10011577)
 37. Soibelzon LH, Rincón A. 2007. The fossil record of the short-faced bears (Ursidae, Tremarctinae) from Venezuela. Systematic, biogeographic, and paleoecological implications. *Neues. Jahrb. Geol. P.* **245**, 287–298. (doi:10.1127/0077-7749/2007/0244-0287)
 38. Schubert BW, Chatters JC, Arroyo-Cabrales J, Soibelzon LH, Awe J, Griffith CS, De Anda G, Erreguerena PL. 2016 The 'South American' short-faced bear *Arctotherium* from the Yucatán Peninsula of Belize and Mexico: implications for their biogeography, paleobiology, evolution, and extinction. *J. Vertebr. Paleontol.* (Program and Abstracts, 2016, p. 220). Poster B151.
 39. Figueirido B, Soibelzon LH. 2009 Inferring palaeoecology in extinct tremarctine bears (Carnivora, Ursidae) using geometric morphometrics: palaeoecology in extinct tremarctines. *Lethaia* **43**, 209–222. (doi:10.1111/j.1502-3931.2009.00184.x)
 40. Soibelzon LH, Grinspan GA, Bocherens H, Acosta WG. 2014 South American short-faced bear (*Arctotherium angustidens*) diet: evidence from pathology, morphology, stable isotopes, and biomechanics. *J. Paleontol.* **88**, 1240–1250. (doi:10.1666/13-143)
 41. Stucchi M, Salas-Gismondi R, Baby P, Guyot J-L. 2009 A 6,000 + year-old specimen of a spectacled bear from an Andean cave in Peru. *Ursus* **20**, 63–68. (doi:10.2192/08GR017R1.1)
 42. Soibelzon LH, Prevosti F. 2012 Fossils of South American land carnivores (Carnivora, Mammalia). In *Molecular population genetics, evolutionary biology and biological conservation of neotropical carnivores* (eds M Ruiz, J Shostell), pp. 509–530. New York, NY: Nova Science Publisher.
 43. Prevosti FJ, Forasiepi AM. 2018 *Evolution of South American mammalian predators during the cenozoic: paleobiogeographic and paleoenvironmental contingencies*. Springer Geology. Cham, Switzerland: Springer International Publishing.
 44. Kurtén B, Anderson E. 1980 *Pleistocene mammals of North America*. New York, NY: Columbia University Press.
 45. Czaplewski NJ, Krejca J, Miller TE. 2003 Late quaternary bats from Cebada Cave, Chiquibul Cave System, Belize. *Caribb. J. Sci.* **39**, 23–33.
 46. Berta A. 1987 Origin, diversification, and zoogeography of the South American Canidae. *Fieldiana, Zool.* **39**, 455–471.
 47. Berta A. 1988 Quaternary evolution and biogeography of large South American Canidae (Mammalia: Carnivora). *Univ. Calif. Publ. Geol. Sci.* **132**, 1–149.
 48. Cartelle C, Langguth A. 1999 *Procyon troglodytes* (Lund): um canídeo intertropical extinto. *Anais da Academia Brasileira do Ciencias* **71**, 371–384.
 49. Prevosti FJ, Zurita AE, Carlini AA. 2005 Biostratigraphy, systematics, and paleoecology of *Procyon* Giebel, 1855 (Carnivora, Canidae) in South America. *J. S. Am. Earth Sci.* **20**, 5–12. (doi:10.1016/j.jsames.2005.05.005)
 50. Prevosti FJ. 2006 Grandes cánidos (Carnivora, Canidae) del Cuaternario de la Republica Argentina: sistemática, filogenia, bioestratigrafía y paleoecología. PhD thesis, Universidad Nacional de La Plata, La Plata, Argentina.
 51. Prevosti FJ, Rincón AD. 2007 A new fossil canid assemblage from the late Pleistocene of northern South America: the canids of the Inciarte Asphalt Pit (Zulia, Venezuela), fossil record and biogeography. *J. Paleont.* **81**, 1053–1065. (doi:10.1666/pleo05-143.1)
 52. Prevosti FJ. 2010 Phylogeny of the large extinct South American canids (Mammalia, Carnivora, Canidae) using a 'total evidence' approach. *Cladistics* **26**, 456–481. (doi:10.1111/j.1096-0031.2009.00298.x)
 53. Prevosti FJ, Schubert BW. 2013 First taxon date and stable isotopes ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$) for the large hypercarnivorous South American canid *Procyon troglodytes* (Canidae, Carnivora). *Quatern. Int.* **305**, 67–73. (doi:10.1016/j.quaint.2012.07.003)
 54. Prevosti FJ, Ubilla M, Perea D. 2009 Large extinct canids from the Pleistocene of Uruguay: systematic, biogeographic and paleoecological remarks. *Hist. Biol.* **21**, 79–89. (doi:10.1080/08912960903249337)
 55. Perini FA, Russo CAM, Schrago CG. 2010 The evolution of South American canids: a history of rapid diversification and morphological parallelism. *J. Evol. Biol.* **23**, 311–322. (doi:10.1111/j.1420-9101.2009.01901.x)
 56. Zrzavy J, Duda P, Bobovsky J, Okrinova I, Pavelkova Ricankova V. 2018 Phylogeny of the Caninae (Carnivora): combining morphology, behaviour, genes and fossils. *Zool. Scr.* **47**, 373–389. (doi:10.1111/zsc.12293)
 57. Tedford RH, Wang X, Taylor BE. 2009 Phylogenetic systematics of the North American fossil Caninae (Carnivora: Canidae). *Bull. Am. Mus. Nat. Hist.* **325**, 1–218. (doi:10.1206/574.1)
 58. Arroyo-Cabrales J, Alvarez T. 2003 A preliminary report of the late Quaternary mammal fauna from Loltún Cave, Yucatán, México. In *Ice age cave faunas of North America* (eds BW Schubert, JI Mead, RW Graham), pp. 262–272. Bloomington, IN: Indiana University Press.
 59. Arroyo-Cabrales J, Polaco OJ. 2003 Caves and the Pleistocene vertebrate paleontology of México. In *Ice age cave faunas of North America* (eds BW Schubert, JI Mead, RW Graham), pp. 273–291. Bloomington, IN: Indiana University Press.
 60. Prevosti FJ, Soibelzon LH. 2012 Evolution of the South American carnivores (Mammalia, Carnivora): a paleontological perspective. In *Bones, clones, and biomes: the history and geography of recent neotropical mammals* (eds BD Patterson, LP Costa), pp. 102–122. Chicago, IL: University of Chicago Press.