CORRESPONDENCE

Mechanical Ventilation and Extracorporeal Membrane Oxygenation in Acute Respiratory Insufficiency

by Dr. med. Falk Fichtner, Prof. Dr. med. Onnen Mörer, PD. Dr. med. Sven Laudi, Prof. Dr. med. Steffen Weber-Carstens, Dr. med. Monika Nothacker, and Prof. Dr. med. Udo Kaisers in issue 50/2018

Conflicting Results

The recommendations of the guidelines on “prone positioning” and “ECMO” must be contested (1). The evidence in support of prone positioning as the primary treatment option in patients with severe ARDS is derived from a meta-analysis (2) which showed a survival benefit in a subgroup analysis. The authors’ conclusion, “...no convincing evidence for benefit or harm of prone positioning...”.

The EOLIA study (early versus rescue use of ECMO [3]) cited in the guideline as proof of the “rescue character” of ECMO, was terminated after 75% of the calculated sample size because no mortality difference was found in the intention-to-treat analysis. A secondary analysis presented by the authors themselves showed a different result: “crossover to ECMO or death in patients in the control group versus death in the ECMO group” yielded an absolute mortality difference of 23% in favor of the ECMO group ($p < 0.001$). At the time of study termination, 28% of patients in the prone-positioning group with persistent severe hypoxia had switched to the ECMO treatment group. These control group patients who received “late” ECMO treatment had a 60-day mortality of 57% compared to 35% in the group of patients with primary ECMO treatment.

The statement—which is repeated several times in the guideline—that in patients with severe ARDS no difference exists between early ECMO treatment and conventional therapy (prone positioning, relaxation, etc.) is not supported by the available data. Similarly, the EOLIA study does not provide evidence to support the recommendation to use ECMO as a “rescue option” only after all conventional treatment modalities have been exhausted in patients with ARDS. Unfortunately, as the result of these guideline recommendations, valuable time will be wasted in patients with severe ARDS on treatments which should only be used to bridge the time spent waiting for the ECMO team, if at all.

DOI: 10.3238/arztebl.2019.0286a

References

1. Fichtner F, Moerer O, Laudi S, Weber-Carstens S, Nothacker M, Kaisers U: Clinical practice guideline: Mechanical ventilation and extracorporeal membrane oxygenation in acute respiratory insufficiency. *Dtsch Arztebl Int* 2018; 115: 840–7.
2. Bloomfield R, Noble DW, Sudlow A: Prone position for acute respiratory failure in adults. *Cochrane Database Syst Rev* 2015; 11: CD008095
3. Combes A, Hajage D, Capellier G, et al.: Extracorporeal membrane oxygenation for severe acute respiratory distress syndrome. *N Engl J Med* 2018; 378: 1965–75.

Dr. med. Michael Schütz
Prof. Dr. med. Hans-Bernd Hopf
 Abteilung für Anästhesie und Perioperative Medizin
 ECLS/ECMO-Zentrum Langen
 Asklepios Klinik Langen
 Langen
 Germany
 m.schuetz@asklepios.com

In Reply:

We stick to our guideline recommendations (1). The quote from the Cochrane review cited by Schütz/Hopf cannot be found in that article (2). Instead, it reads: “We found no convincing evidence of benefit nor harm from universal application of PP (prone positioning, sl) in adults with hypoxaemia [...]”. The result directly reported directly thereafter reads: “Three subgroups (early implementation of PP, prolonged adoption of PP and severe hypoxemia at study entry) suggested that prone positioning may confer a statistically significant mortality advantage” and is the evidence base for the guideline recommendation.

In the guideline, the EOLIA trial was not cited as evidence to support the use of ECMO as a rescue treatment (3). In fact, the lack of reliable RCTs on the use of ECMO and the high potential for complications of ECMO formed the basis for the restrictive recommendation at the time of guideline compilation at the end of 2017. However, the EOLIA trial, which was published after the publication of the guideline, further strengthened this recommendation: EOLIA was terminated early, because a significant survival benefit as the result of early ECMO therapy could not be expected. The statement of Schütz/Hopf on the EOLIA trial, ““crossover to ECMO or death in patients in the control group versus death in the ECMO group” yielded an absolute mortality difference of 23% in favor of the ECMO group ($p < 0.001$)” is incorrect. This secondary endpoint of EOLIA does not measure mortality. On the contrary, for this analysis patients who were treated with ECMO after conventional treatment options had been exhausted and who survived were assessed as patients with negative treatment failure. The crossover group already showed a significantly higher disease severity at the time of inclusion in the EOLIA trial.

In our opinion, the alternative interpretation of the facts proposed by Schütz/Hopf cannot be substantiated by published data on prone positioning and the use of ECMO treatment in patients with severe ARDS.

DOI: 10.3238/arztebl.2019.0286b

References

1. Fichtner F, Moerer O, Laudi S, Weber-Carstens S, Nothacker M, Kaisers U: Clinical practice guideline: Mechanical ventilation and extracorporeal membrane oxygenation in acute respiratory insufficiency. *Dtsch Arztebl Int* 2018; 115: 840–7.
2. Bloomfield R, Noble DW, Sudlow A: Prone position for acute respiratory failure in adults. *Cochrane Database Syst Rev* 2015; 11: CD008095.
3. Combes A, Hajage D, Capellier G, et al.: Extracorporeal membrane oxygenation for severe acute respiratory distress syndrome. *N Engl J Med* 2018; 378: 1965–75.

Dr. med. Falk Fichtner and PD Dr. med. Sven Laudi
 for the guideline group “Mechanical Ventilation and Extracorporeal Membrane Oxygenation in Acute Respiratory Insufficiency”
 Universitätsklinikum Leipzig: Klinik und Poliklinik für Anästhesiologie und Intensivtherapie, Leipzig, Germany; falk.fichtner@medizin.uni-leipzig.de

Conflict of interest statement

The authors of both contributions declare that no conflict of interest exists.