Published in final edited form as:

J Vasc Nurs. 2019 June ; 37(2): 91–105. doi:10.1016/j.jvn.2019.02.001.

A Mixed Methods Study of Perceived Barriers to Physical Activity, Geriatric Syndromes and Physical Activity Levels among Older Adults with Peripheral Artery Disease and Diabetes

Mary O. Whipple, BA, BSN, RN, PhD Candidate^a, Erica N. Schorr, PhD, RN^a, Kristine M.C. Talley, PhD, RN, GNP-BC^a, Ruth Lindquist, PhD, RN^a, Ulf G. Bronas, PhD^b, and Diane Treat-Jacobson, PhD, RN^a

^aUniversity of Minnesota, School of Nursing, Minneapolis, MN

^bCollege of Nursing, University of Illinois, Chicago, IL

Abstract

Previous studies suggest a myriad of factors prevent individuals from engaging in physical activity, however, less is known about barriers faced by individuals with multiple chronic conditions, such as peripheral artery disease (PAD) and type 2 diabetes and how these barriers may impact engagement in physical activity. To date, there are no studies that integrate simultaneous assessment of perceived barriers to physical activity and engagement in physical activity in older adults with PAD and diabetes. This integration is key to understanding the implications of barriers to physical activity and to developing strategies to address those barriers. Therefore, this study investigated the unique physical activity experiences of older adults with PAD and diabetes. This study used a concurrent mixed methods design. Ten adults aged 65 years and older with PAD and diabetes completed semi-structured interviews about experiences with physical activity, self-reported questionnaires assessing quality of life and fear of falling, and measures of physical function (e.g., 6-minute walk test, chair stand, gait speed). Physical activity was measured objectively with accelerometry. Inductive content analysis was used to identify themes and integrated analysis was used to evaluate patterns among qualitative and quantitative variables. On average, participants were 74 years old and spent 10% of their time in moderate or vigorous physical activity (range 3–18%); 80% of participants were men. Barriers to physical activity identified through qualitative interviews included lack of accessibility, lack of enjoyment of activity, lack of motivation, and pain/physical health. Facilitators to physical activity were social support, accessibility and convenience, and enjoyment of the activity. Participants with more sedentary time and less moderate or vigorous physical activity tended to report greater fear of falling, greater barriers to physical activity, and achieved lower 6-minute walk test distances. This research provides insight into both the nature of perceived barriers to physical activity and engagement in physical activity among older adults with PAD and diabetes. The integration of

Corresponding author: Mary O. Whipple, BSN, RN, PhD Candidate, Adult and Gerontological Health Cooperative, 5-140 Weaver-Densford Hall, 308 Harvard St. SE, Minneapolis, MN 55455; whip042@umn.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

self-reported measures and objective measures facilitates our understanding of the lived experiences of individuals with these conditions. Study findings can be used to support further investigation into factors that influence engagement in physical activity in individuals with PAD and diabetes and assist in the development of strategies to address identified barriers.

Keywords

peripheral artery disease; diabetes; exercise; physical activity; barriers; geriatric syndromes

Peripheral artery disease (PAD), a progressive condition characterized by obstruction of blood flow to the lower extremities, is a significant problem among older adults. The prevalence of PAD increases with age, with 10–15% of Americans over age 65 having a diagnosis of PAD (Criqui & Aboyans, 2015). The hallmark of PAD is exercise-induced limb ischemia, or claudication, which can cause significant pain, walking impairment, and reduced quality of life (Oka, 2006). In addition to functional impairment, PAD is also associated with significant risk of cardiovascular morbidity and mortality. Older adults with symptomatic PAD have a 2–3.5 fold greater all-cause mortality risk compared with healthy older adults (Mueller et al., 2014). Diabetes is a significant risk factor for PAD, and estimates suggest that 20–30% of older adults with PAD have diabetes (Beks et al., 1995; Elhadd, Robb, Jung, Stonebridge, & Belch, 1999; Hirsch et al., 2001; Marso & Hiatt, 2006).

The combination of diabetes and PAD puts individuals at greater risk of poor health outcomes, compared to either condition alone (Vinik, Vinik, Colberg, & Morrison, 2015), and this is particularly true in regard to cardiovascular complications. The risk of mortality is two times higher in adults with diabetes and PAD than in those with PAD alone (Leibson et al., 2004; Mueller, Hinterreiter, Poelz, Haltmayer, & Dieplinger, 2016). Walking exercise has been shown to improve walking distance and physical function among adults with PAD (Lane, Ellis, Watson, & Leng, 2014; Parmenter, Dieberg, & Smart, 2015), and may be protective against mortality (Lane et al., 2014). Thus, determining strategies that could help improve adherence and responsiveness to exercise programs are critical.

Given that the majority of US older adults do not meet physical activity guidelines, a number of researchers have sought to examine barriers to physical activity that could be to blame (Bethancourt, Rosenberg, Beatty, & Arterburn, 2014; Biedenweg et al., 2014; Ingram, Ruiz, Mayorga, & Rosales, 2009). This body of work, which is primarily qualitative in nature, suggests that older adults frequently cite factors such as social influences, physical limitations, competing priorities, access difficulties, and personal beliefs regarding physical activity as barriers to physical activity (Franco et al., 2015). In addition to these factors, one of the most commonly cited barriers is health concerns (Franco et al., 2015). Interestingly, research has demonstrated two differing views on the role of health in exercise: some individuals see health concerns as their primary barrier to physical activity while others see their health as a major motivator to engage in physical activity. Thus, it is likely that a similar health-related factor may serve as either a barrier or a facilitator (or neither) to physical activity, depending on the older adult and their chronic condition(s).

The social ecological model of McLeroy et al. (McLeroy, Bibeau, Steckler, & Glanz, 1988) provides a useful overarching framework for understanding barriers and facilitators to physical activity, as it not only considers individual characteristics, but also interpersonal, community, and public policy factors that may promote or prevent physical activity. The social ecological model suggests that an individual's behavior is integrated into this dynamic network, such that intrapersonal factors may influence community factors, and vice versa. Figure 1 provides a visual representation of some hypothesized barriers to physical activity experienced by older adults with PAD and diabetes.

Despite numerous studies investigating barriers to physical activity in a variety of groups of older adults, there are none that specifically examine barriers to physical activity among older adults with comorbid PAD and diabetes. Additionally, there are no studies that collect and integrate assessments of both perceived barriers to physical activity and actual engagement in physical activity. Both subjectively-assessed perceptions and objectively-measured activity levels are key to understanding the impact of barriers to physical activity and developing strategies to address those barriers. Given the increased risk of morbidity and mortality faced by older adults with PAD and diabetes (Leibson et al., 2004), it is important to better understand barriers to physical activity such as health status in this population. Exploration of the beliefs of older adults with respect to how healthcare providers could be more effective in working with individuals to overcome barriers is also necessary, as it will inform the development of strategies to reduce such barriers.

In addition to health status related to PAD and diabetes, geriatric syndromes may also play a role in older adults' willingness and ability to engage in physical activity. Geriatric syndromes refer to multifactorial health conditions caused by the accumulated effects of impairments in multiple systems. They cannot be classified as, or attributed, to a single disease. Examples include falls, dizziness, frailty, cognitive impairment, and urinary incontinence. Such conditions are highly prevalent among older adults in the US and are associated with significant morbidity and poor health outcomes (Inouye, Studenski, Tinetti, & Kuchel, 2007). Fear of falling, frailty, and functional limitations could significantly influence an older adult's ability to engage in regular physical activity (Araki & Ito, 2009).

The Current Study

Given the dearth of information related to barriers to and engagement in physical activity among older adults with PAD and diabetes, we conducted a mixed methods study investigating these phenomena. While there is some qualitative work addressing barriers to physical activity in a variety of adult populations including individuals with PAD (Bethancourt et al., 2014; Biedenweg et al., 2014; Galea Holmes, Weinman, & Bearne, 2015; Galea, Bray, & Ginis, 2008; Ingram et al., 2009), there are no studies that integrate simultaneous assessment of both perceived barriers to physical activity and the actual degree of engagement in physical activity in older adults with PAD and diabetes. This type of integration is key to understanding the impact of barriers to physical activity and developing strategies to address those barriers.

As these issues could not be adequately addressed through the use of only qualitative or only quantitative techniques, we conducted a mixed methods study. Mixed methods, which can be defined as intentionally combining both qualitative and quantitative methods at some stage in the data collection, analysis, or interpretation of a given research study (Creswell & Plano-Clark, 2011), provided a unique opportunity to understand the impact of PAD and diabetes on an individual and the barriers to physical activity that he or she encounters. The purpose of this mixed methods study was to investigate the unique experiences of older adults with PAD and diabetes related to physical activity. The overarching question we sought to address was: How does having PAD and diabetes shape older adults' perceptions of and engagement in physical activity? More specifically, we sought to answer:

1. How do older adults with PAD and diabetes describe their challenges to engaging in physical activity?
2. How important are geriatric syndromes, such as falls, fearing of falling, and decreased muscle strength, to perceptions of ability to engage in physical activity among older adults with PAD and diabetes?
3. What types of patterns of objectively measured barriers, physical activity levels, and sedentary time are present among older adults with PAD and diabetes?

Method

Design

A concurrent mixed methods multiple case study design was used for the purposes of complementarity and completeness (Happ, Dabbs, Tate, Hricik, & Erlen, 2006; Stake, 1994; Yin, 2009). This design was appropriate for the present study because a single method (e.g., qualitative or quantitative) was deemed insufficient to fully evaluate barriers to physical activity and engagement in physical activity among older adults with PAD and diabetes. A multiple case study was used as it provides a rich description that allows researchers to seek out both what is common and what is particular about the case (Stake, 1994), enabling the understanding of barriers and beliefs related to physical activity of older adults who may have differing views.

For this study, priority (emphasis) was given to the qualitative (QUAL) strand. Quantitative (quan) data were used to enhance understanding and contribute to overall completeness. Figure 2 illustrates the study design, including points of integration.

Participants

Participants were identified through convenience sampling. Patients who were enrolled in a supervised exercise program for PAD offered through [institution removed for review] who met study eligibility criteria were invited to participate. Additionally, individuals with PAD who were currently engaged in or had completed previous studies of exercise programs conducted by the [institution removed for review] were invited to participate. Pseudonyms are used to refer to individual participants throughout the manuscript.

Inclusion criteria for the study were: having lifestyle-limiting claudication with activity due to PAD, type 2 diabetes, being aged 65 years or older, and being able to speak and read English. There were no exclusionary criteria, as it was deemed important to include all eligible participants in order to gain a variety of perspectives on exercise benefits and barriers.

This study was reviewed and approved by the [institution removed for review] Institutional Review Board. All participants provided written informed consent. Participants received \$50 for the time and inconvenience associated with data collection and were provided with a parking pass for the duration of their visit.

Data Collection and Measures

All qualitative and quantitative data were collected during the same encounter. Participants completed a semi-structured interview; measures of balance, lower body strength, and physical function; and several self-reported measures of perceived benefits and barriers to exercise, geriatric syndromes, and quality of life during one visit to the [institution removed for review]. As PAD severity may impact physical function, the most recent ankle-brachial index (ABI) was obtained (if available) for each participant via medical record review. Participants were also sent home with an accelerometer for assessment of physical activity and sedentary time for two weeks. Potential biases were limited by use of a detailed audit trail (Jootun, McGhee, & Marland, 2009; Whitehead, 2004).

Assessment of barriers and benefits to physical activity.—Perceived barriers and benefits to physical activity were assessed using two methods: a semi-structured interview and administration of the Exercise Benefits and Barriers Scale (EBBS) (Sechrist, Walker, & Pender, 1987), to obtain both open-ended qualitative and quantitative data on these constructs. Additionally, in order to enable comparisons with previous research in older adults with PAD, facilitators of physical activity were also assessed.

Each semi-structured interview was conducted by the lead author [initials removed for review] and lasted between 16 and 55 minutes (mean 37.8 minutes, SD 12.1 minutes). The interview guide can be found in Table 1. Interviews were audio-recorded and transcribed verbatim prior to analysis. All participants completed the interview prior to the quantitative survey measures in order to reduce the potential influence of the measures on the content of the interviews.

The EBBS contains 43 items composing 9 factors (5 benefits and 4 barriers) (Sechrist et al., 1987). Participants were asked to rate each of the items on a 4-point scale (strongly agree, agree, disagree, and strongly disagree). Exercise benefits factors included life enhancement (7 items), physical performance (9 items), psychological outlook (6 items), social interaction (4 items), and preventive health (3 items). Exercise barriers factors included exercise milieu (6 items), time expenditure (3 items), physical exertion (3 items), and family encouragement (2 items). Sample items included “I am too embarrassed to exercise” (exercise milieu), “I will live longer if I exercise” (preventive health), and “exercise is hard work for me” (physical exertion). The EBBS has been used a variety of populations, including older adults, and has good psychometric properties (Sechrist et al., 1987).

Measures of balance, strength, and physical function.—Participants completed several performance-based measures of balance, lower body strength, and physical function following completion of the interview. These functional measures took approximately 30 minutes to complete.

The Short Physical Performance Battery (SPPB) assesses three components of physical function: standing balance, 4 meter walking velocity, and repeated chair rises (Guralnik et al., 1994). For the test of standing balance, participants stand in three positions: feet together (side by side position), semi-tandem position, and tandem position. For each position, participants are timed to a maximum of 10 seconds. Walking speed is assessed by asking participants to walk at their usual pace over a 4-meter distance. Finally, participants are asked to rise from a straight back chair without using their arms five times in a row as quickly as possible. The 4-meter walk and chair stand tests were each completed twice, with the mean of the two times used for all analyses. The SPPB has been used extensively with older adults to evaluate functional status, as well as in adults with PAD (McDermott et al., 2015; McDermott, Liu et al., 2014). This battery has been shown to be effective in characterizing physical function across a broad spectrum of functional status and predictive of mortality and institutionalization across this spectrum (Guralnik et al., 1994).

The 6-minute walk test (6MWT) has been used extensively in cardiovascular and pulmonary disease research, including individuals with PAD as a method of monitoring disease status and intervention effectiveness (McDermott, Guralnik et al., 2014). To complete the 6MWT, participants are instructed to walk back and forth in a 100 foot, level hallway, covering as much distance as possible in 6 minutes (ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories, 2002). Scores are calculated as the number of meters the participant walks during the 6 minutes.

Grip strength was measured as the mean of three trials of each hand using a Baseline Smedley digital hand dynamometer according to the American Society of Hand Therapists guidelines for measurement of grip strength (Fess, 1992).

Measures of walking impairment and quality of life.—Three questionnaires were used to assess walking impairment and quality of life related to PAD: the Walking Impairment Questionnaire (WIQ), the PAD Specific Quality of Life Questionnaire (PADQOL), and the Short-Form 36 (SF-36). The WIQ consists of three subscales: walking distance, speed, and stair climbing and was developed specifically for patients with PAD (Sagar, Brown, Zelt, Pickett, & Tranmer, 2012). WIQ scores range from 0 to 100 with 100 representing no impairment and 0 being complete impairment. This questionnaire has been shown to correlate well with treadmill walking distance and to be sensitive to change over time (Sagar et al., 2012).

The PADQOL is a 38-item questionnaire developed by Treat-Jacobson and colleagues (2012). The PADQOL contains five factors: social relationships and interactions, self-concept and feelings, symptoms and limitations in physical functioning, fear and uncertainty, and positive adaptation. Total scores on the PADQOL range from 0 to 100, with 100 reflecting better quality of life.

The SF-36 is a 36 item health survey designed to assess different aspects of physical, psychological, and social quality of life (Ware, 1976, 2000). The SF-36 yields a profile of functional health and quality of life, as well as standardized physical and mental health summary scores. For all subscales and the two composite scales, higher scores are indicative of better quality of life. The SF-36 has been used extensively in clinical research (Ware, 2000) and in patients with PAD (Turner-Bowker, Bartley, & Ware, 2002).

Assessment of selected geriatric syndromes related to falling.—Participants completed two short measures to assess for the presence of geriatric syndromes. To assess falls, participants were asked “In the past month, have you had a fall, slip, or trip, in which you lost your balance and landed on the floor, ground, or lower level?” (Lamb, Jorstad-Stein, Hauer, Becker, & Prevention of Falls Network Europe and Outcomes Consensus Group, 2005). If the individual reported they had experienced a fall, the study investigator probed for further details of the circumstances surrounding the fall(s) and whether the falls resulted in injury. Fear of falling was assessed using the Falls Efficacy Scale – International (FESI). The FESI contains 16 items that assess how concerned a person is about falling in a variety of situations, including walking on a slippery surface, going up and down stairs, or when visiting a friend or relative (Yardley et al., 2005). Higher scores indicate greater fear of falling.

Measurement of physical activity and sedentary time.—Physical activity and sedentary time were measured using the Actigraph wGTx3-BT. Participants were given the device at the completion of the in-person visit and asked to wear the device 24 hours a day on their non-dominant wrist (except when bathing or swimming) for 14 days. The device was used to capture physical activity time, sedentary time, and length of bouts of sedentary behavior in minutes, as well as the number of breaks in sedentary time. Minutes in each type of activity were divided by the total time the participant wore the device while awake to yield percent time in each activity. Participants were provided with a postage-paid envelope by which to return the device.

Data Integration and Analysis

Qualitative and quantitative data were initially analyzed separately for descriptive purposes (described below), as well as in conjunction with one another.

Qualitative analysis.—Both directed and conventional inductive content analysis were used to identify themes (Hsieh, 2005). Following transcription, audio recordings and transcripts were reviewed to ensure the accuracy of transcription. Transcripts were then imported into Atlas.ti (Version 8, Scientific Software Development GmbH, Berlin) and first read multiple times to obtain a sense of the interviews as a whole. Interviews were then read individually and directed content analysis was used to identify themes. First-cycle coding was conducted using descriptive coding (Saldana, 2013). Basic labels were assigned to sections of text to provide an overview of the topics using the social ecological framework as a guide (Figure 1). Coding was conducted using Atlas.ti. In addition, we utilized elaborative coding, a second-cycle coding method (Saldana, 2013), to identify additional categories that appeared to represent key concepts or ideas that were not included in the social ecological

framework. This second-cycle method was more typical of conventional inductive content analysis (Hsieh, 2005). Exemplar quotes were used to illustrate themes identified through the interviews.

Quantitative analysis.—Descriptive statistics (mean, median, standard deviation, and percent) were used to summarize scores on quality of life and walking impairment instruments, barriers to exercise questionnaire, and the presence and severity of geriatric syndromes. Actigraph data were downloaded and processed using ActiLife (Version 9.0.0, Actigraph Corp., Pensacola, FL) and the wrist-specific adaptation of the Troiano adult cut points to identify sedentary, light, moderate, and vigorous activity (99, 100–2019, 2020–5998, and 5999 counts per minute, respectively). SPSS Statistics (Version 25, IBM Corp., Armonk, NY) was used to analyze quantitative data.

Integrated Analysis.—Integrated analysis was used to describe participants and evaluate patterns of qualitative themes and quantitative variables among participants. Relevant demographic variables, ABI, percent time spent in sedentary, light, moderate, and vigorous physical activity, and key qualitative themes identified through the interviews were summarized for each individual using a meta-matrix (Bazeley, 2009; Creswell & Plano-Clark, 2011). The meta-matrix was examined for variability among individual participants and potential patterns related to perceived barriers to physical activity and volume of physical activity and sedentary time.

We also evaluated the congruence of themes identified through the qualitative responses with survey responses on the individual factors of the EBBS through the use of a second meta-matrix, in order to explore how representative the questions on the EBBS were, with respect to the barriers reported by individuals during the qualitative interviews. Themes identified in the interviews were “quantized” according to whether or not they were present to allow for examination of patterns in the data (Sandelowski, Volls, & Knafl, 2009), and the mean score on each factor was reported for both the entire sample and only individuals in whose interviews each theme was identified.

Results

A total of 10 participants were enrolled in the study and completed all study procedures (except for one participant who did not complete the 6MWT due to time constraints). Participants were, on average, 74.2 years of age (SD 4.3), married (70%), and all self-identified as non-Hispanic white. The majority of participants were men (80%). The mean ABI (lowest leg) was 0.67 (range 0.32 to 0.91).

Qualitative Interview Themes

Several themes were identified through analysis of the qualitative interviews related to barriers, facilitators, and benefits of physical activity. Each of these categories and the themes identified in each is discussed in detail below.

Barriers

Barriers to physical activity reported by participants are shown in Figure 3. The barriers most commonly identified by participants were lack of accessibility, lack of enjoyment of activity, lack of motivation, and pain/physical health.

Accessibility.—Accessibility of safe and comfortable places to exercise was a common concern of participants (n=6). As Darla stated, “I could go to the arthritis [water exercise] class, but that meets right away at 9. I have to take the freeways to get there; that’s bad freeway time. I don’t want to be on the road really.” This concern about safe transportation to and from exercise limited Darla’s ability to engage in an exercise program she enjoyed. Additionally, participants expressed concern about access to equipment: “I don’t have the equipment. I did [when on vacation] go out and do some walking, but I didn’t have the weights and bike...If you go so long [without exercising], then you don’t feel guilty any more, and you don’t get started right up again.” (Frank). For participants, accessibility was related to both obtaining transportation, as well as having a location they perceived as accessible given their health concerns. As Jack stated when thinking about a local gym to which he had a membership but was somewhat hesitant to use:

I always watched [the parking lot at the gym] and usually between 2 and 3:30 it was pretty quiet in there, so then I’d go in. Otherwise, the parking lot had quite a few cars and I said, nah, I ain’t gonna go in there and do my wimpy stuff... I just thought you’ve got all these people running five miles on the treadmill and doing all that other stuff and I’m just in there getting in their way, so I just wait and bide my time and go in there when I can feel more comfortable.

Therefore, it is clear that accessibility of a location was not only related to transportation and timing of the facility, but also feeling comfortable in their surroundings.

Lack of enjoyment.—Lack of enjoyment of the activity was also a barrier, cited by four participants. As Cindy said, “Hey, I’m honest. Exercising’s not fun, for the most part. It’s work. If I can make myself happier making a batch of homemade cookies, I’m going to do the cookies. Besides, I get to eat them.” This sentiment was echoed by another participant who stated simply, “Sometimes I want to be doing something else.” (Clyde). This lack of enjoyment of the activity is tied closely to motivation.

Motivation.—Motivation was a significant barrier to exercise for seven participants. As one participant stated, “There has to be a certain amount of motivation [to exercise], and the desire to really put your mind to it is probably a good idea. I don’t do that.” (Fred). Cindy shared a similar sentiment, “All you can do is to make it available. You can’t make me exercise; I have to be the one to do that...It has to be within every individual. You could want us to want to exercise, but you can’t make us want to exercise, and that’s the whole thing.” (Cindy). Similarly, Clyde expressed the barrier of lack of motivation well, “If it ain’t in the body, it ain’t anywhere, so if you’ve got it in your mind to do it – it’s like quitting smoking or anything. If you don’t want to, there’s no way in hell you’re going to quit.” (Clyde).

Pain and physical health.—Pain and physical health were also commonly expressed barriers; six participants reported that their physical health impacted their ability to exercise. Lloyd expressed this concern simply: “Pain is a barrier. Why would I wanna do something that hurts me?” And, as Darla stated, “You can’t walk when you hurt.” Several participants expressed that arthritis (typically knee or hip pain) was a major limitation to their exercise, suggesting that comorbid conditions may play a role in exercise engagement, sometimes to a greater degree than PAD. Despite being distinct themes, there seemed to be a close relationship between lack of motivation and pain as a barrier to exercise for some participants. As Fred stated:

Our health insurance says I could go to [a gym] and have a free membership there to do whatever I wanted to do...I don’t have the desire to do that. I’m not saying I’m lazy, but I just...I think if I could improve my walking, that would probably be the best for me, but then I sit and think, oh God, my legs are going to get tired. What if I only go so far and all of the sudden I can’t get back?

Concerns about physical ability and a safe place to exercise given their level of ability were significant and according to participants, had a significant effect on the amount of exercise in which they engaged.

Facilitators

Facilitators to physical activity reported by participants were distilled into three themes: social support, accessibility and convenience, and enjoyment of the activity (Figure 3).

Social support.—Social support, including support of family and friends and the presence of positive role models was a facilitator identified by six participants. As Cindy said:

[One of the] side benefits is socialization. I’ve met some really nice people, and we all have a lot of things in common: bad hearts and falling-apart bodies that we’re just trying to keep going. It’s just absolutely amazing.

Darla expressed similar feelings, describing a group of women with whom she used to attend water therapy classes for arthritis. Not only was there the physical support of having others who “look like me” but also the psychological support. As Darla stated, “It’s a group that understands...We’re all in the same boat; we’ve all had a lot of the same thing.” Positive role models were also valuable facilitators. As Jim said:

I got some friends in their 80s now and a few in their 90s: they’re spry as anything. I have a sister 11 years older than I am; in her 80s already, and she drives, she goes to exercise. She’s very active. I said “hey that’s what I gotta do.” I just gotta keep going.

Having family and friends who have experienced similar challenges persist in engaging in exercise and providing support that “we’re in this together” was a key facilitator for participants who had and currently engaged in regular exercise.

Accessibility and convenience.—Accessibility and convenience of opportunities to engage in exercise were mentioned as facilitators by half of the participants. As Jack said of

a local gym he attends: “They’ve got the equipment that I kind of like and have got people to advise you on what you should be doing” which helps facilitate exercising. Similarly, as Jim said “I think being a member somewhere is very helpful. Paying the money; you might as well make use of it.” The feeling that he has made a commitment was of particular importance to Clyde, “If I say I’m going to do it [exercise], I’ll do it... If I commit to something, then fine, I’m going to do it, stay with it until it’s done.” This sense of accountability and availability of resources to promote exercise (particularly in the setting of a research study for several of the participants), helped motivate them to engage in exercise that they, in their perspective, committed to doing.

Enjoyment.—Finally, enjoyment of the activity was a facilitator of activity mentioned by four participants. As Cindy said,

When you exercise – and I don’t care if it’s exercise-exercise, or if it’s doing a sport you enjoy doing; it can be something as innocuous as even shuffleboard or ping pong, bocce ball – if it’s something you enjoy doing, you’re still active.

To Cindy and several other participants, finding a type of exercise that was not only suitable given their health issues (e.g., knee/hip arthritis, back pain), but also fun was a key facilitator and defining factor in whether or not they would engage in that exercise regularly.

Benefits

Three themes were identified with respect to benefits of exercise: energy, mobility/physical health, and sense of accomplishment.

Energy.—Forty percent of participants indicated that they felt exercise gave them more energy. For example, Darla said,

It puts me in a better mood all over. I feel more positive about everything, and maybe when I do come home, I won’t do a lot, but I will do a few little things around the house, maybe move things around or go do something in the kitchen or something like that, write some notes that I’m supposed to write.”

Mobility & physical health.—For Darla, exercise enabled her to do more activities around the house and better engage in life with her family. Another benefit of exercise reported by participants was improved mobility. As Cindy stated,

For me, it’s like I told my daughter. I said, ‘You may not see it as much of an improvement, but I do.’ I mean, if I can go from walking 150 feet to 300 feet without stopping, that, for me, is an improvement. That’s one less stop I have to make. That’s one less bench I have to sit down on so that I can let the blood flow get back into the legs. I’ll take every one of those minor increments I can get. It may not sound like much. (Cindy) Improved overall health was also a perceived benefit to six participants. “I breathe better.

My sleep, I fall asleep right away. I get the blood flow going and even down to my legs, it’s beneficial. Plus, it makes me stronger.” (Jim) This feeling of improved energy and function

were not only benefits of exercise, but also important factors cited by participants that encouraged them to continue engaging in exercise.

Sense of accomplishment.—Finally, four participants also reported that exercise gave them a sense of accomplishment. “I go [exercise]. And I feel better for it.” (Jim) Similarly, Frank said, “I just felt, okay, I’m doing something I should be doing, and it’s for my own good.” This sense of accomplishment was also manifested in improved mobility. As Lloyd stated, the ability of staff at the exercise program in which he participates to say “Look, this is where you were, this is where you are now.” was incredibly helpful showing him the progress he has made. This feeling that the exercise was helping them in some way was important in motivating participants to continue with exercise in spite of the barriers they encountered.

Quantitative Outcomes

Mean (SD) scores on self-reported questionnaires and physical function tests are reported in Table 2. On average, participants spent 66.9% (range 53–78%) of their time in sedentary behavior and 10.2% (range 3–18%) of their time in moderate to vigorous physical activity. Degree of impairment in walking distances, speed, and stairs as measured by the WIQ varied greatly among participants, as did PAD-related quality of life (PADQOL). On average, participants endorsed benefits from exercise more than barriers to exercise as measured using the EBBS, with mean scores of 64.1 and 38.0 for benefits and barriers, respectively. Physical exertion was the most strongly endorsed exercise barrier, with a percentile score of 47.5.

With respect to geriatric syndromes, three participants (30%) (Jack, Darla, and Clyde) reported one or more falls (range 1 – 3) in the previous three months. Causes of falls included icy sidewalks, misjudging a curb, and household hazards (e.g., furniture leg, rug). Several falls resulted in muscle soreness/bruising, but none resulted in serious injury. Concern about falling as measured using the FES-I varied substantially among participants, however all participants expressed at least some concern, most frequently on the items “walking on a slippery surface” and “walking on an uneven surface.”

Integrated Analysis

A meta-matrix outlining key participant characteristics, quantitative variables, and qualitative themes is displayed in Table 3. Participants with greater sedentary time and less moderate to vigorous physical activity (MVPA) tended to report greater fear of falling (higher FESI score), and barriers to physical activity, and had slower gait speeds, more difficulty with the chair rise, and lower 6MWT distances.

In order to examine the congruence of the themes identified in the interviews with the domains of the EBBS, we constructed matrices looking at the overlap of these components. Tables 4 and 5 display count data with the number of individuals who reported each interview theme, and the average score on each EBBS factor for individuals who reported that theme among benefits and barriers, respectively.

Some patterns did emerge with respect to scores on the EBBS and qualitative themes. Participants who endorsed increased energy as a benefit of exercise in the interview tended to have higher scores on all of the EBBS benefits domains (Table 4), as did participants who reported that exercise gave them a sense of accomplishment. With respect to barriers, physical exertion was reported as the greatest barrier to exercise according to the EBBS across all participants, regardless of themes identified from the qualitative interviews (Table 5). The theme “lack of accessibility” from the interviews corresponded most closely to the EBBS domain “exercise milieu,” with participants who expressed difficulty with accessibility in their interviews having higher scores on this subscale. Similarly, participants who reported that lack of enjoyment of the activity was a barrier tended to place more importance on the time expenditure involved in exercise as a barrier to activity.

Discussion

This study provides rich descriptive data related to barriers to and engagement in physical activity among older adults with PAD and diabetes. The barriers and facilitators to physical activity identified here are similar to those identified in other qualitative studies in individuals with PAD. In one study, Galea and colleagues found that personal motivation and leg discomfort were the most commonly cited barriers to physical activity (Galea et al., 2008). These are similar to the barriers of “lack of motivation” and “pain and physical health” identified in this study. Additionally, these barriers are also similar to those commonly cited by older adults in general (Franco et al., 2015).

Fear of falling was also a common concern among participants which may present a barrier to physical activity. Previous studies have demonstrated that self-reported stumbling and unsteadiness are 86% more prevalent in those with PAD than without PAD (Barbosa et al., 2015) and that a history of falling is associated with impairments in physical function, including balance, in individuals with PAD (Gardner & Montgomery, 2001). Additionally, diabetes is associated with increased fear of falling and fear-associated activity restriction (Bruce, Hunter, Peters, Davis, & Davis, 2015). Although fear of falling was not commonly reported in the qualitative interviews, all participants reported at least some concern about falling on the FES-I, and three reported recent falls. Therefore, consideration of concern about falling and ways to address such concerns is important when working with older adults with PAD and diabetes to promote increased physical activity.

There were also similarities in the facilitators to physical activity identified in this study as compared to previous work. Social support, and accessibility and convenience were particularly important, providing possible targets for future interventions to promote physical activity. For example, one of the most commonly cited disease-specific facilitators for exercise in the study by Galea et al. (2008) was the availability of a supervised treadmill walking program. The recent Centers for Medicare and Medicaid Services (CMS) determination to reimburse the cost of exercise therapy for beneficiaries with symptomatic PAD provides a new opportunity to engage individuals with PAD in exercise therapy, and for the growth of programs to make this effective therapy more readily available. Additionally, social support was an important facilitator; something that could be considered when designing/planning exercise groups. Individuals with PAD may be self-conscious of or

embarrassed by their walking ability, so having a group of supportive individuals who also have PAD could be an important component in adherence to and success with an exercise program.

Despite the small sample size, we did obtain a clear range of experiences among participants. Darla expressed some of the greatest impairments, both globally and related to PAD, with the lowest scores on several factors of the PADQOL, particularly related to social relationships and interactions, self-concept and feelings, and symptoms and limitations and physical functioning. She also had one of the lowest scores on the SF-36 mental composite score and the lowest physical composite. She expressed significant impairment in walking (had the lowest WIQ score on all three domains) and the shortest distance on the 6MWT. Interestingly, but perhaps not surprisingly, she expressed the greatest barriers to physical activity on the EBBS. Alternatively, Jim and Robert expressed substantially less impairment, with scores on the WIQ, PADQOL, and SF-36 that were some of the highest among the participants.

The barriers and facilitators to physical activity identified in this study are somewhat consistent with those that were hypothesized in the context of the social ecological model of McLeroy et al. (1988). The identified themes are shown in Figure 3. Although many of the personal (e.g., physical health, motivation, enjoyment) and interpersonal (e.g., social support) levels were consistent with identified themes, few specific themes were identified related to community and public policy. This is perhaps because there were not specific questions included in the interview guide to probe for broader barriers to and facilitators of physical activity, but also could be that participants felt that the personal and interpersonal levels had greater relevance to and bearing on their behavior. Future studies are needed to examine how community and public policy issues may serve as a barrier or facilitator to physical activity among older adults with PAD and diabetes.

The findings of this study also suggest that, in general, barriers are easier for participants to identify (or perhaps more similar between participants) as compared to facilitators, where there were fewer commonalities. This could be related to the sample size and lack of saturation. We also found that, except for physical exertion (which was the most strongly endorsed barrier), benefits were more strongly endorsed for physical activity than were the barriers, however, this could be related to participants motivated by social desirability. We also found significant congruence between measures of physical function and engagement in physical activity, with participants who had the greatest impairments in physical function tending to engage in less physical activity overall, and in particular, less moderate to vigorous physical activity, than their counterparts who had less difficulty with the physical function measures. Although this does not suggest that poor physical function leads to lower engagement in physical activity (or vice versa), it is an important consideration in designing physical activity interventions, as well as interventions that target reducing inactivity. Notably, there did not appear to be a relationship between resting ABI and physical activity levels or physical function as measured by the 6MWT among participants, suggesting that resting ABI alone may not be an important factor in engagement in and performance of physical activity for this sample.

This study has a number of strengths. First, the mixed methods design is unique from research currently being conducted in this population in that it provided the opportunity for integrated analysis to examine the interrelationships of quantitative patterns of activity and physical function and qualitative investigation of barriers and facilitators to this activity. Second, the use of accelerometers to assess physical activity provided an objective method of measuring activity levels and quantifying sedentary time. Finally, the integration of both self-reported and objective measures of walking impairment and function was a strength as it enabled us to examine patterns of both types of measures in relationship to activity.

This study has several important limitations. First, the study sample was limited in scope. As the sample was drawn from current/former participants in exercise studies related to PAD, their views may not be representative of all individuals with PAD, particularly those who have never engaged in a formal exercise program. However, the views of individuals who have engaged in exercise studies for PAD are revealing, in that despite these programs, they express a number of significant barriers to exercise. Additionally, this highlights that although program availability is important, it is not the sole barrier to engagement in exercise for individuals with PAD. A second limitation is the small sample size, particularly with regard to female participants (n=2), and so we may not have reached saturation with respect to interview themes or full range of experiences. Third, we did not collect additional data on other comorbid health conditions and not all participants had an ABI to quantify the severity of PAD, which may have enhanced our understanding of other limitations to physical function in addition to PAD. Finally, this study was cross-sectional in nature, and so, while we were able to examine patterns in physical activity and physical function at one point in time, we were unable to examine potential temporal relationships between activity levels, changes in function, and barriers and facilitators to exercise.

Conclusion

In conclusion, this mixed methods study provides rich insight into both the nature of barriers to physical activity and engagement in physical activity among older adults with PAD and diabetes. The integration of both self-reported and objective measures facilitates our understanding of the lived experiences of individuals with these conditions and provides opportunities for future research. The findings of this study can be used to support further investigation into factors that influence engagement in physical activity among individuals with PAD and diabetes and assist in the development of strategies to address perceived barriers, particularly as supervised exercise therapy programs for individuals with PAD become more widely available in clinical care.

Acknowledgements

The authors would like to thank Dr. Barbara McMorris for her valuable assistance in the development of this manuscript.

Ms. Whipple was a 2015–2017 National Hartford Center of Gerontological Nursing Excellence (NHCGNE) Patricia G. Archbold Scholar. The Patricia G. Archbold Scholar program is supported by a grant to the Gerontological Society of America (GSA)/NHCGNE from The John A. Hartford Foundation.

This publication was made possible by the National Institutes of Health under a Ruth L. Kirschstein National Research Service Award (F31NR016614, PI Whipple) from the National Institute of Nursing Research and by

UL1TR000114 from the National Center for Advancing Translational Sciences (NCATS) of the National Institutes of Health. The content is solely the responsibility of the author and does not necessarily represent the official views of the National Institutes of Health.

Portions of the results reported in this manuscript were presented as a poster at the Society for Vascular Nursing 2017 Annual Meeting in Nashville, TN, and the 2018 Annual Meeting in Boston, MA.

References

- Araki A, & Ito H (2009). Diabetes mellitus and geriatric syndromes. *Geriatrics & Gerontology International*, 9(2), 105–114. 10.1111/j.1447-0594.2008.00495.x [doi] [PubMed: 19740352]
- ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories. (2002). ATS statement: Guidelines for the six-minute walk test. *American Journal of Respiratory and Critical Care Medicine*, 166(1), 111–117. 10.1164/ajrcm.166.1.at1102 [doi] [PubMed: 12091180]
- Barbosa JP, Farah BQ, Chehuen M, Cucato GG, Farias JC Junior, Wolosker N, ... Ritti-Dias RM (2015). Barriers to physical activity in patients with intermittent claudication. *International Journal of Behavioral Medicine*, 22(1), 70–76. 10.1007/s12529-014-9408-4 [doi] [PubMed: 24715636]
- Bazeley P (2009). Integrating data analyses in mixed methods research. *Journal of Mixed Methods Research*, 3(3), 203–207. 10.1177/1558689809334443
- Beks PJ, Mackaay AJ, de Neeling JN, de Vries H, Bouter LM, & Heine RJ (1995). Peripheral arterial disease in relation to glycaemic level in an elderly Caucasian population: The Hoorn study. *Diabetologia*, 38(1), 86–96. [PubMed: 7744233]
- Bethancourt HJ, Rosenberg DE, Beatty T, & Arterburn DE (2014). Barriers to and facilitators of physical activity program use among older adults. *Clinical Medicine & Research*, 12(1–2), 10–20. 10.3121/cmr.2013.1171 [doi] [PubMed: 24415748]
- Biedenweg K, Meischke H, Bohl A, Hammerback K, Williams B, Poe P, & Phelan EA (2014). Understanding older adults' motivators and barriers to participating in organized programs supporting exercise behaviors. *The Journal of Primary Prevention*, 35(1), 1–11. 10.1007/s10935-013-0331-2 [doi] [PubMed: 24214654]
- Bruce D, Hunter M, Peters K, Davis T, & Davis W (2015). Fear of falling is common in patients with type 2 diabetes and is associated with increased risk of falls. *Age and Ageing*, 44(4), 687–690. 10.1093/ageing/afv024 [PubMed: 25739747]
- Creswell JW, & Plano-Clark VL (2011). *Designing and conducting mixed methods research* (2nd ed.). Thousand Oaks, CA: Sage Publications, Inc.
- Criqui MH, & Aboyans V (2015). Epidemiology of peripheral artery disease. *Circulation Research*, 116(9), 1509–1526. 10.1161/CIRCRESAHA.116.303849 [doi] [PubMed: 25908725]
- Elhadd TA, Robb R, Jung RT, Stonebridge PA, & Belch JFF (1999). Pilot study of prevalence of asymptomatic peripheral arterial occlusive disease in patients with diabetes attending a hospital clinic. *Practical Diabetes International*, 16(6), 163–166.
- Fess EE (1992). Grip strength In Casanova JS (Ed.), *Clinical Assessment Recommendations* (Vol. 2nd, pp. 41–45). Chicago, IL: American Society of Hand Therapists.
- Franco MR, Tong A, Howard K, Sherrington C, Ferreira PH, Pinto RZ, & Ferreira ML (2015). Older people's perspectives on participation in physical activity: A systematic review and thematic synthesis of qualitative literature. *British Journal of Sports Medicine*, 49(19), 1268–1276. 10.1136/bjsports-2014-094015 [doi] [PubMed: 25586911]
- Galea Holmes MN, Weinman JA, & Bearne LM (2015). "You can't walk with cramp!" A qualitative exploration of individuals' beliefs and experiences of walking as treatment for intermittent claudication. *Journal of Health Psychology*, 22(2), 255–265. 10.1177/1359105315600238
- Galea MN, Bray SR, & Ginis KAM (2008). Barriers and facilitators for walking in individuals with intermittent claudication. *Journal of Aging and Physical Activity*, 16, 69–84. [PubMed: 18212396]
- Gardner AW, & Montgomery PS (2001). Impaired balance and higher prevalence of falls in subjects with intermittent claudication. *The Journals of Gerontology -Series A - Biological Sciences and Medical Sciences*, 56(7), M454–8.
- Guralnik JM, Simonsick EM, Ferrucci L, Glynn RJ, Berkman LF, Blazer DG, ... Wallace RB (1994). A short physical performance battery assessing lower extremity function: Association with self-

- reported disability and prediction of mortality and nursing home admission. *Journal of Gerontology*, 49(2), M85–94. [PubMed: 8126356]
- Happ MB, Dabbs AD, Tate J, Hricik A, & Erlen J (2006). Exemplars of mixed methods data combination and analysis. *Nursing Research*, 55(2 Suppl), S43–S49. 10.1097/00006199-200603001-00008 [PubMed: 16601634]
- Hirsch AT, Criqui MH, Treat-Jacobson DJ, Regensteiner JG, Creager MA, Olin JW, ... Hiatt WR (2001). Peripheral arterial disease detection, awareness, and treatment in primary care. *JAMA*, 286(11), 1317–1324. <http://doi.org/joc11056> [pii] [PubMed: 11560536]
- Hsieh H-F (2005). Three approaches to qualitative content analysis. *Qualitative Health Research*, 15(9), 1277–1288. 10.1177/1049732305276687 [PubMed: 16204405]
- Ingram M, Ruiz M, Mayorga MT, & Rosales C (2009). The Animadora Project: Identifying factors related to the promotion of physical activity among Mexican Americans with diabetes. *American Journal of Health Promotion*, 23(6), 396–402. 10.4278/ajhp.08021915 [doi] [PubMed: 19601479]
- Inouye SK, Studenski S, Tinetti ME, & Kuchel GA (2007). Geriatric syndromes: Clinical, research, and policy implications of a core geriatric concept. *Journal of the American Geriatrics Society*, 55(5), 780–791. <http://doi.org/JGS1156> [pii] [PubMed: 17493201]
- Jootun D, McGhee G, & Marland GR (2009). Reflexivity: Promoting rigour in qualitative research. *Nursing Standard*, 23(23), 42–46.
- Lamb SE, Jorstad-Stein EC, Hauer K, Becker C, & Prevention of Falls Network Europe and Outcomes Consensus Group. (2005). Development of a common outcome data set for fall injury prevention trials: The Prevention of Falls Network Europe consensus. *Journal of the American Geriatrics Society*, 53(9), 1618–1622. <http://doi.org/JGS53455> [pii] [PubMed: 16137297]
- Lane R, Ellis B, Watson L, & Leng GC (2014). Exercise for intermittent claudication. *The Cochrane Database of Systematic Reviews*, 7, CD000990 10.1002/14651858.CD000990.pub3 [doi]
- Leibson CL, Ransom JE, Olson W, Zimmerman BR, O'Fallon WM, & Palumbo PJ (2004). Peripheral arterial disease, diabetes, and mortality. *Diabetes Care*, 27(12), 2843–2849. [PubMed: 15562195]
- Marso SP, & Hiatt WR (2006). Peripheral arterial disease in patients with diabetes. *Journal of the American College of Cardiology*, 47(5), 921–929. [PubMed: 16516072]
- McDermott MM, Guralnik JM, Criqui MH, Ferrucci L, Liu K, Spring B, ... Rejeski WJ (2015). Unsupervised exercise and mobility loss in peripheral artery disease: A randomized controlled trial. *Journal of the American Heart Association*, 4(5), e001659–e001659. 10.1161/JAHA.114.001659 [PubMed: 25994445]
- McDermott MM, Guralnik JM, Criqui MH, Liu K, Kibbe MR, & Ferrucci L (2014). Six-minute walk is a better outcome measure than treadmill walking tests in therapeutic trials of patients with peripheral artery disease. *Circulation*, 130(1), 61–68. 10.1161/CIRCULATIONAHA.114.007002 [PubMed: 24982117]
- McDermott MM, Liu K, Ferrucci L, Tian L, Guralnik J, Kopp P, ... Criqui MH (2014). Vitamin D status, functional decline, and mortality in peripheral artery disease. *Vascular Medicine (London, England)*, 19(1), 18–26. 10.1177/1358863X13518364 [doi]
- McLeroy K, Bibeau D, Steckler A, & Glanz K (1988). A ecological perspective on health promotion programs. *Health Education Quarterly*, 15, 351–377. [PubMed: 3068205]
- Mueller T, Hinterreiter F, Luft C, Poelz W, Haltmayer M, & Dieplinger B (2014). Mortality rates and mortality predictors in patients with symptomatic peripheral artery disease stratified according to age and diabetes. *Journal of Vascular Surgery*, 59(5), 1291–1299. 10.1016/j.jvs.2013.11.063 [PubMed: 24393280]
- Mueller T, Hinterreiter F, Poelz W, Haltmayer M, & Dieplinger B (2016). Mortality rates at 10 years are higher in diabetic than in non-diabetic patients with chronic lower extremity peripheral arterial disease. *Vascular Medicine*, 21(5), 445–452. 10.1177/1358863X16643603 [PubMed: 27067137]
- Oka RK (2006). Peripheral arterial disease in older adults: Management of cardiovascular disease risk factors. *The Journal of Cardiovascular Nursing*, 21(5 Suppl 1), S15–20. <http://doi.org/00005082-200609001-00005> [pii] [PubMed: 16966924]
- Parmenter BJ, Dieberg G, & Smart NA (2015). Exercise training for management of peripheral arterial disease: A systematic review and meta-analysis. *Sports Medicine (Auckland, N.Z.)*, 45(2), 231–44. 10.1007/s40279-014-0261-z

- Sagar SP, Brown PM, Zelt DT, Pickett WL, & Tranmer JE (2012). Further clinical validation of the walking impairment questionnaire for classification of walking performance in patients with peripheral artery disease. *International Journal of Vascular Medicine*, 2012, 190641 10.1155/2012/190641 [doi] [PubMed: 22919494]
- Saldana J (2013). *The Coding Manual for Qualitative Researchers* (2nd ed.). Los Angeles: Sage.
- Sandelowski M, Volls CI, & Knafl G (2009). On quantizing. *Journal of Mixed Methods Research*, 3(3), 208–222. 10.1177/1558689809334210 [PubMed: 19865603]
- Sechrist KR, Walker SN, & Pender NJ (1987). Development and psychometric evaluation of the exercise benefits/barriers scale. *Research in Nursing & Health*, 10(6), 357–365. [PubMed: 3423307]
- Stake RE (1994). Case studies In Denzin NK & Lincoln YS (Eds.), *Handbook of qualitative research* (pp. 236–247). Thousand Oaks, CA: Sage.
- Treat-Jacobson DJ, Lindquist RA, Witt DR, Kirk LN, Schorr EN, Bronas UG, ... Regensteiner JG (2012). The PADQOL: Development and validation of a PAD-specific quality of life questionnaire. *Vascular Medicine*, 17(6), 405–15. 10.1177/1358863X12466708 [PubMed: 23184901]
- Turner-Bowker DM, Bartley PJ, & Ware JE (2002). SF-36® Health Survey & 'SF' Bibliography (Vol. 3rd (1988-). Lincoln, RI: QualityMetric Incorporated.
- Vinik AI, Vinik EJ, Colberg SR, & Morrison S (2015). Falls risk in older adults with type 2 diabetes. *Clinics in Geriatric Medicine*, 31(1), 89–99, viii 10.1016/j.cger.2014.09.002 [doi] [PubMed: 25453303]
- Ware JE (1976). Scales for measuring general health perceptions. *Health Services Research*, 11(4), 396–415. [PubMed: 1030696]
- Ware JE (2000). SF-36 health survey update. *Spine*, 25(24), 3130–3139. [PubMed: 11124729]
- Whitehead L (2004). Enhancing the quality of hermeneutic research: Decision trail. *Journal of Advanced Nursing*, 45(5), 512–518. 10.1046/j.1365-2648.2003.02934.x [PubMed: 15009354]
- Yardley L, Beyer N, Hauer K, Kempen G, Piot-Ziegler C, & Todd C (2005). Development and initial validation of the Falls Efficacy Scale-International (FES-I). *Age and Ageing*, 34(6), 614–619. <http://doi.org/34/6/614> [pii] [PubMed: 16267188]
- Yin RK (2009). *Case Study Research: Design and Methods*. Thousand Oaks, CA: Sage.

Highlights

- Access, lack of enjoyment, and physical health were barriers to physical activity
- Social support, convenience, and enjoyment facilitated physical activity
- Greater sedentary time was associated with greater barriers to physical activity
- Participants who were less active also reported greater fear of falling

Figure 1.
Social ecological framework model of hypothesized barriers and facilitators to physical activity for older adults with PAD and diabetes.

Figure 2. Study diagram showing data collection and analysis and points of integration of the QUAL (qualitative) and quan (quantitative) strands.

Figure 3. Social ecological model with categories of barriers and facilitators of physical activity identified through qualitative interviews.

Note: (–) denotes barrier; (+) denotes facilitator

Table 1

Semi-structured interview guide.

1	Tell me about when your PAD and DM started and how they have influenced your life. Probes: Has having PAD/DM changed your perception of your health? Which condition came first? Did the diagnosis of the second condition change your perspective?
2	Have you sought out information about lifestyle changes since you developed PAD/DM, and if so, where do you go for information? What information has been meaningful for you?
3	What role do exercise or physical activity have in your health? Give me examples of things that you have tried and how they work or do not work for you? What works best for you? Probes: What is it like to try and exercise regularly? What makes it easier or harder? What kinds of activities are you engaging in and how satisfying do you find those activities?
4	Tell me about your exercise habits in the past. Probes: What were your exercise habits earlier in life? Were you an athlete? Did that change, and if so, when did it change?
5	If you have had a discussion with health care provider about increasing exercise or physical activity, how did that discussion go? Probes: What types of activities did you discuss? How did the provider get your input?
6	How do you think increasing your physical activity has or would influence your health? Probes: What types of benefits do you see to exercise? What types of negative effects or consequences of exercise have you or are you concerned about experiencing? For example, low blood sugar, dizziness?
7	What types of things keep you from exercising? Probes: Is claudication a barrier? What else prevents you from exercising or exercising as much as you would like to? Other than symptoms related to PAD or DM, are there other things that prevent you from exercising? Do you have someone who would exercise with you or supports you in exercise? Does exercise affect your mood or relationships with others?
8	How do these barriers compare to one another? Probes: Are some more significant for you than others? Which ones are the most limiting? In what way?
9	What things could help you or do help you exercise?
10	Do you have goals related to exercise? If so, what are they? Probes: How do you feel you are doing at achieving those goals? What could your health care provider do to help you achieve those goals?

Table 2

Mean (standard deviation [SD]) scores on self-report questionnaires and physical function assessments.

		Mean (SD)	Range
<i>Self-report questionnaires</i>			
WIQ			
	Distance	35.7 (34.7)	4.3 – 100
	Speed	37.6 (28.3)	3.3 – 89.1
	Stairs	49.5 (39.9)	0 – 100
FES-I		26.0 (8.2)	18.0 – 46.0
PADQOL			
	Factor 1 – Social relationships and interactions	66.7 (22.3)	24.4 – 97.8
	Factor 2 – Self-concept and feelings	72.9 (23.1)	28.6 – 100
	Factor 3 – Symptoms and limitations in physical functioning	39.0 (22.7)	10 – 80
	Factor 4 – Fear and uncertainty	63.0 (20.4)	30 – 90
	Factor 5 – Positive adaptations	64.0 (14.6)	37.1 – 85.7
SF-36			
	Physical Composite	36.3 (10.6)	16.4 – 51.5
	Mental Composite	59.3 (5.4)	50.6 – 66.6
EBBS Benefits			
	Life Enhancement	65.2 (14.2)	47.6 – 100
	Physical Performance	68.9 (13.1)	48.2 – 100
	Psychological Outlook	64.4 (17.2)	50.0 – 100
	Social Interaction	51.7 (16.1)	33.3 – 75.0
	Preventive Health	63.3 (20.3)	33.3 – 100
	Overall	64.1 (13.7)	47.1 – 96.6
EBBS Barriers			
	Exercise Milieu	28.9 (18.8)	0 – 61.1
	Time Expenditure	40.0 (19.7)	11.1 – 77.8
	Physical Exertion	63.3 (16.6)	33.3 – 88.9
	Family Encouragement	24.2 (18.6)	0 – 66.7
	Overall	38.0 (15.4)	14.3 – 69.1
<i>Physical function assessments</i>			
SPPB			
	Balance	3.6 (0.5)	3 – 4
	4 meter walk	4.42 (1.01)	3.01 – 6.19
	Chair stand	11.2 (2.9)	6.6 – 14.5
Grip strength			
	Non-dominant	28.8 (9.6)	15.7 – 44.2
	Dominant	29.2 (9.8)	16.3 – 45.5

WIQ = Walking Impairment Questionnaire, FES-I = Falls Efficacy Scale-International, PADQOL = Peripheral Artery Disease Quality of Life Questionnaire, SF-36 = Short Form-36, EBBS = Exercise Benefits and Barriers Scale, SPPB = Short Physical Performance Battery

Table 3

Meta-matrix summarizing key participant characteristics, physical activity, performance on physical function measures, and reported barriers and facilitators of physical activity.

Case #	Age	ABI (lowes t leg	6 Minute Walk (feet)	Activity			SPPB		4-meter walk (sec)	Grip Strength Dominant Hand (kg)	FES-I	Interview Themes	
				Sedentary (%)	Light (%)	MVPA (%)	Balance (points out of 4)	Repeated Chair Rise (sec)				Barriers	Facilitators
Darla	72	0.91	480	77	18	5	3	11.66	3.44	16.3	46	• Accessibility • Motivation • Pain/physical health	• Family/friend support
Cindy	69	0.57	750	74	19	7	4	14.52	3.61	19.4	21	• Enjoyment • Motivation • Pain/physical health	• Accessibility • Enjoyment • Family/friend support
Jack	78	0.44	750	76	21	3	3	Unable to do	6.19	35.7	32	• Accessibility • Motivation	• Accessibility • Enjoyment • Family/friend support • Time alone
Fred	78	0.90	902	78	16	5	3	Unable to do	5.33	31.8	27	• Enjoyment • Motivation	
Clyde	80	0.32	1077	63	29	8	3	Unable to do	4.85	24.8	27	• Enjoyment • Motivation	
Jim	74	0.63	1310	60	25	15	4	8.91	3.71	35.3	19	• Accessibility • Pain/physical health	
Lloyd	76	+	1400	63	24	13	4	6.64	4.28	20.8	21	• Enjoyment • Motivation • Pain/physical health	• Family/friend support • Time alone
Frank	76	0.70	1470	65	24	11	4	13.02	3.01	40.2	18	• Accessibility • Motivation • Pain/physical health	
Robert	73	0.87	1615	61	23	16	4	12.31	4.41	22.7	23	• Accessibility	• Family/friend support
Durrell	66	+	^	53	29	18	4	Unable to do	5.35	45.5	26	• Accessibility • Pain/physical health	

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

* Case names reported are pseudonyms;
^Participant did not complete due to time constraints, +No ABI available
ABI = Ankle Brachial Index, MVPA = Moderate to Vigorous Physical Activity, SPPB = Short Physical Performance Battery, FES-I = Falls Efficacy Scale – International

Table 4

Comparison of interview identified benefits with EBBS benefits.

Interview Theme	EBBS Factors – Benefits				
	Life Enhancement	Physical Performance	Psychological Outlook	Social Interaction	Preventive Health
Energy (n=4 [*])	76.2	77.8	76.4	64.6	80.6
Mobility (n=5 [*])	61.0	63.0	53.3	43.3	53.3
Physical health (n=6 [*])	68.3	67.9	64.8	47.2	61.1
Sense of accomplishment (n=4 [*])	73.8	75.0	65.3	56.3	75.0

EBBS = Exercise Benefits and Barriers Scale

Note: Higher scores indicate greater endorsement of exercise benefit. Scores reported are EBBS factor scores that have been transformed to percentages for ease of interpretation.

* Indicates number of participants who expressed benefits related to the given theme.

Table 5

Comparison of interview identified barriers with EBBS barriers.

Interview Theme	EBBS Factors – Barriers			
	Exercise Milieu	Time Expenditure	Physical Exertion	Family Encouragement
Lack of accessibility (n=6 [*])	30.6	37.0	59.3	26.5
Lack of enjoyment of activity (n=4 [*])	26.4	44.4	69.4	20.8
Lack of motivation (n=7 [*])	28.6	42.9	69.8	27.5
Pain, physical health (n=6 [*])	35.2	40.7	61.1	30.6

EBBS = Exercise Benefits and Barriers Scale

Note: Higher scores indicate greater endorsement of exercise barrier. Scores reported are EBBS factor scores that have been transformed to percentages for ease of interpretation.

^{*} Indicates number of participants who expressed barriers related to the given theme.