Published in final edited form as:

Sports Biomech. ; : 1–8. doi:10.1080/14763141.2018.1535617.

The Effect of Landing Surface on Landing Error Scoring System Grades

Kimberley Jacobs¹, Diego Riveros¹, Heather K. Vincent, PhD², and Daniel C. Herman, MD, PhD²

¹College of Medicine, University of Florida, Gainesville, Florida, USA

²Department of Orthopaedics and Rehabilitation, College of Medicine, University of Florida, Gainesville, Florida, USA

Abstract

Different landing surfaces may affect lower extremity biomechanical performance during athletic tasks. The magnitude of this effect on clinical screening measures such as jump-landings is unknown. This study determined the effect of court (CS), grass (GS), and tile (TS) surfaces on Landing Error Scoring System (LESS) grades. A repeated-measures design was used. Forty (21F, 19M; mean age = 23.8 ± 2.4 yr) recreational athletes performed a jump-landing task on three different landing surfaces. 2D videography recorded jump-landings in the frontal and sagittal planes. A 2 X 3 (sex by surface) mixed-model repeated-measures analysis of variance was used to examine main and interaction effects associated with surface and sex. No significant sex by landing surface interactions existed for LESS grades. No significant differences were observed on LESS grades for the main effect of surface (CS = 4.83 ± 1.31 points; GS = 5.01 ± 1.40 points; TS = 5.09 ± 1.86 points; all $p > 0.05$). Correlations were found between LESS grades among different conditions (r range = 0.587 to 0.611; all $p < 0.001$). Commonly used jump-landing surfaces for clinical biomechanical evaluations do not affect LESS grades, suggesting generalisability as a screening tool for anterior cruciate ligament injury risk in different sport environments.

Keywords

anterior cruciate ligament; basketball court; drop jump; grass; drop-jump landing

Introduction

Anterior cruciate ligament (ACL) injuries are common in sports and increase the risk for accelerated development of degenerative joint disease of the knee, knee surgeries, and related physical impairment (Mather, 2013; Roos, Adalberth, Dahlberg, & Lohmander, 1995). Female athletes are at a much higher risk for experiencing an ACL injury compared to male athletes across multiple sports and play levels (Dai, Herman, Liu, Garrett, & Yu, 2012). Seventy percent of these injuries occur through non-contact mechanisms that involve sudden stops, landing, or changes in acceleration, as seen in sports that involve cutting and

landing sports (Acevedo, Rivera-Vega, Miranda, & Micheo, 2014; Boden, Sheehan, Torg, & Hewett, 2010; Dai, Herman, Liu, Garrett, & Yu, 2012).

Poor jump-landing biomechanics as studied using three-dimensional motion analysis have been shown to be predictive of ACL injury and are likely to be a significant contributing factor to the increased risk for these injuries in female athletes (Hewett et al., 2005; Dai, Herman, Liu, Garrett, & Yu, 2012). Because of the cost and feasibility limitations of 3-D motion analysis, other inexpensive alternatives have been developed. The Landing Error Scoring System (LESS) is a screening test that is both inexpensive and reliable in assessing jump-landing biomechanics on a large scale (Padua et al., 2009). The system scores jump 'errors' based on 17 criteria previously shown to correlate with the high-risk biomechanics of an ACL injury during a jump-landing task. The LESS is a valid and reliable tool for identifying potentially high-risk movement patterns based on assessments of frontal, sagittal, and transverse plane motion during a jump-landing task. (Padua et al., 2009; Padua et al., 2011). The LESS can identify high-risk biomechanics in athletes in a controlled laboratory setting; however, the applicability of this tool as a large-scale screening method in the field on the actual surfaces of play (i.e. a basketball court or a soccer/football field) needs to be tested.

Prior research has demonstrated an effect of sport playing surface on lower extremity biomechanics in laboratory settings and on ACL strain and injury rates (Balazs et al., 2015; Drakos et al., 2010; Dowling, Corazza, Chaudhari, & Andriacchi, 2010; Thomson, Whiteley, & Bleakley, 2015). Dowling, Corazza, Chaudhari, and Andriacchi (2010) examined jump-landing mechanics on surfaces with high and low coefficients of friction and found decreased knee-flexion angles and high external knee valgus moments on a high coefficient of friction shoe-surface interfaces during a 30° sidestep cutting task. This finding was attributed to a shift in the participants' task strategy for each surface, which may lead to lower extremity biomechanics that are associated with ACL-injury risk. Differences in the shoe-surface interface have similarly been shown to alter the magnitude of ACL strain in cadaveric models (Drakos et al., 2010). Furthermore, these differences in lower extremity biomechanics and ACL strain may translate into real differences in ACL-injury rates across different playing surfaces, with multiple studies demonstrating greater injury rates during play on surfaces with high coefficients of friction compared to surfaces with low coefficients of friction (Thomson et al., 2015).

These findings suggest that clinical ACL-injury risk screening may need to be conducted in a manner that replicates the field conditions of the athlete's sport. LESS grades obtained from one setting (e.g., indoor court surface) may not fully translate to a setting with a different landing surface (e.g., grass). These findings may also have research implications; LESS grades obtained in a controlled laboratory setting for use in prospectively determining injury risk may result in a different 'high-risk' threshold value than what would otherwise be obtained if the screening occurred on the surface of the athletes' sport.

Determination of the effect of landing surface on LESS grades will provide key evidence on the applicability of LESS as a screening tool in multiple settings and provide guidance for both future research efforts in this area and for clinical implementation of this screening tool

in the field. Therefore, the purposes of this study were as follows: (1) to determine whether differences exist among LESS grades when performed on two common athletic surfaces (court and natural grass) compared to the laboratory tile-floor surface; and (2) to determine whether there are significant sex by landing surface interactions of the LESS grades. We hypothesised that LESS scores would vary across surface conditions, with highest scores on a court surface and lowest scores on a grass surface. Furthermore, we hypothesised that a significant sex by landing surface interaction would exist such that females would demonstrate greater LESS grades than males in the difference environments, with particularly increased grades on the court surface compared to the grass surface.

Methods

Study Design.

This study used a randomised, repeated-measures design to determine the effect of three different landing surfaces on LESS grades.

Participants.

A total of 40 healthy recreational male and female athletes between the ages of 18 and 30 years with experience in performing drop jump-landing tasks were recruited for this study (see Table 1). Participants were recruited from the Gainesville, Florida community and surrounding areas using flyers and word of mouth. The participants were actively engaged in cutting and jump-landing intensive sports such as basketball, volleyball, soccer/football, or lacrosse at least three times a week, or had participated in similar sports on the high-school varsity level and currently participated at least once a month. Potential participants were excluded if any of the following was true: (1) they did not meet the activity-level criteria; (2) they had an orthopedic injury at the time of participation (history of fracture, sprain, or strain within the past five months; acute joint/muscle pain severity of $>5/10$ points on a 10-point scale); or (3) they had a medical condition limiting them from fully participating in sports-related activity. All participants read and signed an informed consent. This study and its procedures were approved by the Institutional Review Board at the University of Florida.

Participant Characteristics.

The height, weight, and age of the participants were obtained. Leg dominance was self-reported as the leg with which the participant preferred to kick a ball. Body weight and height were collected using a standard, medical grade scale. Participants were not given specific instructions on footwear other than to wear comfortable sneakers in which they typically play their sport.

LESS Testing Protocol.

The LESS protocol previously described by Padua et al. (2009) was used for this study. The participants jumped from a 0.30 meter high box to a 0.75 meter by 0.45 meter landing target positioned at a distance 50% of their height away from the box. Upon landing, the participant immediately completed a vertical jump at maximal effort. Participants were not provided any feedback or coaching on their technique unless the task was not completed correctly (e.g., landing outside of the target area). Prior to data collection, participants were

allowed to practise the jump until they were able to perform it successfully (with a minimum of three practise jumps). Three successful jump-landings were then recorded for the given surface using standard videography. Two video cameras (Sony Handycam; Sony Corp.; Tokyo, Japan) were placed 3.4 meters from the landing target in front of and to the dominant side of the participant to capture frontal and sagittal plane views of the jump-landing motion at 30 frames per second.

As described by Padua et al. (2009), a successful jump was characterised by (1) jumping off of both feet at the same time; (2) jumping forward, but not vertically, to the landing target; (3) landing with each foot entirely within the landing target; (4) landing with no more than one-half the length or width of each foot outside of the landing target during the second landing; and (5) completing the task in a single fluid motion.

Surface-Landing Conditions.

Surface-landing condition testing order was randomised using a computer-generated random number table and recorded separately in sealed numbered envelopes. Envelopes were assigned to participants in the order of recruitment and the contents of the envelopes were blinded to both participant and research staff until consent was obtained. Two surfaces were selected to represent common athletic landing surfaces: an indoor basketball court and an outdoor grass field. The third condition was an indoor motion-analysis laboratory floor, which was comprised of a stiff tile covering.

Data Processing and LESS Grading.

Once the video data were collected, the videos were spliced and randomised. Two investigators trained in the LESS grading technique (K.J. and D.R.) under the direction of the principal investigator (D.H.) independently assigned a LESS grade to each jump-landing, and the average score from the two raters was used for analysis. Each jump recording was graded using the criteria described by Padua et al. (2009). Seventeen scored items in the checklist represented errors in landing technique, graded as a '0' or '1' with the former signifying that correct form was observed in the jump-landing task. If the grade scores differed by more than one point, then the principal investigator served as the arbitrator and graded the jump trial to determine the values that were in best agreement. Tester blinding to the participant's identity or surface condition for a given jump-landing during video scoring was not possible.

Statistical Analysis.

Statistical analyses were performed using the Statistical Package for the Social Sciences (version 23.0; IBM Corp.; Chicago, IL, USA). Descriptive statistics were obtained to characterise the study groups. Pearson's correlations were assessed, with a Bonferroni correction for multiple tests. Intra-class correlation coefficients (ICC) were used to determine the reliability of the average score for participants across each surface type. ICC values were also calculated for each jump landing trial across all surfaces in order to assess the inter-rater reliability of the scoring. To determine whether differences existed in the LESS grades between men and women among the three landing-surface conditions, a 2 X 3

(sex by landing surface) mixed-model repeated-measures analysis of variance was used. Significance was established *a priori* at 0.05 for all statistical tests.

Results

Correlations among the Surface Conditions.

Significant correlations existed among the LESS grades of the different conditions ($r = 0.587$ for tile surface vs grass surface, $r = 0.608$ for court surface vs grass surface, and $r = 0.611$ for court surface vs tile surface; $p < 0.001$ for all comparisons). The reliability of scoring across surface types was good for each pairwise analysis (ICC = 0.781 for tile surface vs grass surface, ICC = 0.755 for court surface vs grass surface, and ICC = 0.805 for court surface vs tile surface; $p < 0.001$ for all comparisons). The ICC values for LESS grading across all trials between the two raters was 0.93, indicating excellent inter-rater reliability.

Sex-Based Differences among Landing Surfaces.

The LESS grades for the drop jump-landings for male and female participants are shown in Table 2. We found no significant interaction of sex by landing-surface condition ($p = 0.943$). Moreover, there were no significant main effects of sex ($p = 0.624$) or landing surface ($p = 0.758$).

Discussion and Implications

The purposes of this study were to determine whether differences existed among LESS grades when performed on two common athletic surfaces and a laboratory tile-floor surface, and to determine whether significant sex by landing-surface interaction existed for LESS grading. The main findings were as follows: (1) the mean LESS scores were not different among basketball court, field grass, or laboratory concrete surfaces; and (2) no significant sex by landing-surface interactions existed for the LESS scores. Thus, our study hypotheses were not supported by these findings.

These results indicate that the LESS may be relatively insensitive to a range of different landing conditions. We found that significant moderate to moderately high correlations existed in LESS grades between the different surface conditions. This implies a relatively high degree of consistency of LESS grades for a given athlete among different jump-landing surfaces, and thus provides support for potential use of the LESS for ACL-injury risk screening in landing-surface conditions that may be different from the athletes' actual sporting environment. This increases the robustness of the LESS as an ACL-injury risk screening tool as practical implementation of screening efforts may occur in a large variety of settings, ranging from the athletes' playing surface (e.g., grass or court) to non-playing environments such as a physician's office for a pre-participation examination.

The finding of LESS being relatively insensitive to different surface landing conditions is in contrast to findings from a prior study examining the effect of surface on lower extremity biomechanics during a cutting task (Dowling, Corazza, Chaudhari, & Andriacchi, 2010). It is possible that the two surface conditions in this study utilised surfaces with coefficients of friction that had a greater magnitude of difference (0.38 vs. 0.87) compared to the surfaces

utilised in the current study. Furthermore, the current study utilised a drop-landing task rather than a cutting task, and it is possible that the cutting task in the prior study required greater athletic demands which allowed differences in biomechanics between the surface types to be more apparent. Indeed, the study by Dowling, Corazza, Chaudhari, and Andriacchi required that the participants utilise their maximum comfortable approach speed for the low friction surface condition. Despite this, the main kinematic finding in that study was a reduced knee flexion at initial contact in the high friction surface condition with a mean difference of less than three degrees relative to the low friction surface condition. It is doubtful that a mean difference of this magnitude would be relevant to the methods of the LESS which relies on visual judgement by a human observer.

The LESS grades obtained in our study are relatively consistent with the prior literature. Padua et al. (2009) and Theiss et al. (2014) collected data on incoming military cadets and reported mean LESS grades of 4.92 points and 5.28 points, respectively, which are very similar to our reported values (see Table 2). Although the average age of the participants in these studies was slightly younger (mean age 19.3 yr in Theiss et al., 2014), it is likely that these incoming cadets were of the same maturational status as the participants in the present investigation. Similarly, our strong inter-rater reliability ($ICC = 0.93$) was found to be similar to both Padua et al.'s (2009; $ICC = 0.84$) and Onate, Cortes, Welch, and Van Lunen's (2010; $ICC = 0.84$). On the other hand, prior studies have noted sex-based differences in LESS grades, with females demonstrating a higher number of errors (Padua et al., 2009; Theiss et al., 2014). In contrast to these studies, we did not observe any main effects for LESS grades based on sex. Females demonstrated mean LESS grades that were slightly higher than for males, particularly for the court-surface condition, but this was not significant. It is possible that our sample size, which was far smaller than Padua et al.'s (2009) and Theiss et al.'s (2014), was insufficient to reliably detect such differences. It is also possible that the footwear used by the female participants was sufficiently different compared to that used by the male participants to blunt any sex-based differences in LESS grades. If these findings are repeated with larger samples and among additional landing surfaces, the collective evidence may indicate that the LESS could successfully be used in the actual environment in which the athlete plays.

Although our study used video cameras to review jumps at a later date, previous research has been done to look at the use of a modified scoring criteria, the LESS real-time (LESS-RT), which does not require the use of video-camera recording, assessing jump-landing tasks in the field. The LESS-RT was shown to be reliable and compared well with the original LESS study (Padua et al., 2011). Further research into the validity of LESS-RT on the surfaces tested here and others would further strengthen its practical use.

Limitations, Strengths, and Future Directions.

Limitations are present within this study. We elected against the use of particular standard athletic footwear as the primary hypothesis was to determine within-participant differences across landing surfaces. As previously noted, however, this may have affected participant comparisons, such as those based on sex. Additionally, this study evaluated the LESS grades after jump-landings on two surfaces, a basketball court and natural grass, that are commonly

associated with specific footwear. The specific use of basketball shoes or soccer/football cleats on their respective surfaces could have resulted in a change in the shoe-surface interface sufficient to produce biomechanical alterations detectable by the LESS. This may be particularly true of the grass-cleat interface, and additional testing in this area is warranted. A strength of the study lies, however, in the repeated-measures design where the only variable to change between participants was the landing surface. As such, the footwear effect on LESS scores was likely low in this study. This study was limited to laboratory tile, court, and natural grass surfaces, but several other possible playing surfaces may be considered in future studies, such as synthetic outdoor turf. Furthermore, this study population consisted of college-aged recreational athletes, which does not represent the full diversity of athletic populations that participate in sports considered to be at elevated risk for ACL injury. It is possible that different magnitudes of effect of landing surface may be observed in populations of particular sports or maturational categories. Strengths of this study include rigorous LESS grading training for the raters that resulted in very high ICC values. The relative homogeneity of the participants included in the study may also have reduced variation in the scoring. Finally, the frame rates of the cameras used in this study could have introduced insensitivity to the LESS grading process. While 30 frames per second is consistent with previous studies by Padua et al.(2009) and Onate, Cortes, Welch, and Van Lunen (2010), it is possible the some differences could be distinguished with the use of cameras with higher frame rates by virtue of a more accurately assessment of the moment of initial contact during landing.

Conclusion

There were no significant differences in the LESS grades between court, grass, and laboratory-tile surfaces. This study suggests that the LESS may be useful for injury-risk screening when performed on landing surfaces outside the laboratory such as a basketball court or a soccer/football field.

Acknowledgements:

This work was supported by the Medical Student Research Program, Discovery Pathways at the University of Florida College of Medicine; and the National Institutes of Health Grant 5K12HD001097–17.

Research was conducted at the University of Florida Department of Orthopaedics and Rehabilitation.

References

- Acevedo RJ, Rivera-Vega A, Miranda G, & Micheo W (2014). Anterior cruciate ligament injury: identification of risk factors and prevention strategies. *Current Sports Medicine Reports*, 13(3), 186–191. doi:10.1249/JSR.0000000000000053 [PubMed: 24819011]
- Balazs GC, Pavey GJ, Brelan AM, Pickett A, Keblish DJ, & Rue J-PH (2015). Risk of anterior cruciate ligament injury in athletes on synthetic playing surfaces: a systematic review. *The American Journal of Sports Medicine*, 43(7), 1798–1804. doi:10.1177/0363546514545864 [PubMed: 25164575]
- Boden BP, Sheehan FT, Torg JS, & Hewett TE (2010). Noncontact anterior cruciate ligament injuries: mechanisms and risk factors. *The Journal of the American Academy of Orthopaedic Surgeons*, 18(9), 520–527. [PubMed: 20810933]

- Dai B, Herman D, Liu H, Garrett WE, & Yu B (2012). Prevention of ACL injury, part I: injury characteristics, risk factors, and loading mechanism. *Research in Sports Medicine*, 20(3–4), 180–197. doi:10.1080/15438627.2012.680990 [PubMed: 22742075]
- Dowling AV, Corazza S, Chaudhari AMW, & Andriacchi TP (2010). Shoe-surface friction influences movement strategies during a sidestep cutting task: implications for anterior cruciate ligament injury risk. *The American Journal of Sports Medicine*, 38(3), 478–485. doi: 10.1177/0363546509348374 [PubMed: 20194954]
- Drakos MC, Hillstrom H, Voos JE, Miller AN, Kraszewski AP, Wickiewicz TL, ... O'Brien SJ (2010). The effect of the shoe-surface interface in the development of anterior cruciate ligament strain. *Journal of Biomechanical Engineering*, 132(1), 11003. doi:10.1115/1.4000118
- Hewett TE, Myer GD, Ford KR, Heidt RS, Colosimo AJ, McLean SG, ... Succop P (2005). Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *The American Journal of Sports Medicine*, 33(4), 492–501. doi:10.1177/0363546504269591 [PubMed: 15722287]
- Mather RC (2013). Societal and economic impact of anterior cruciate ligament tears. *The Journal of Bone and Joint Surgery (American)*, 95(19), 1751. doi:10.2106/JBJS.L.01705
- Onate J, Cortes N, Welch C, & Van Lunen BL (2010). Expert versus novice interrater reliability and criterion validity of the landing error scoring system. *Journal of Sport Rehabilitation*, 19(1), 41–56. [PubMed: 20231744]
- Padua DA, Boling MC, Distefano LJ, Onate JA, Beutler AI, & Marshall SW (2011). Reliability of the landing error scoring system-real time, a clinical assessment tool of jump-landing biomechanics. *Journal of Sport Rehabilitation*, 20(2), 145–156. [PubMed: 21576707]
- Padua DA, Marshall SW, Boling MC, Thigpen CA, Garrett WEJ, & Beutler AI (2009). The Landing Error Scoring System (LESS) is a valid and reliable clinical assessment tool of jump-landing biomechanics: The JUMP-ACL study. *The American Journal of Sports Medicine*, 37(10), 1996–2002. doi: 10.1177/0363546509343200 [PubMed: 19726623]
- Roos H, Adalberth T, Dahlberg L, & Lohmander LS (1995). Osteoarthritis of the knee after injury to the anterior cruciate ligament or meniscus: the influence of time and age. *Osteoarthritis and Cartilage*, 3(4), 261–267. [PubMed: 8689461]
- Theiss JL, Gerber JP, Cameron KL, Beutler AI, Marshall SW, Distefano LJ, ..., Yunker CA (2014). Jump-landing differences between varsity, club, and intramural athletes: the Jump-ACL Study. *Journal of Strength and Conditioning Research*, 28(4), 1164–1171. doi:10.1519/JSC.0b013e3182a1fdcd [PubMed: 23820560]
- Thomson A, Whiteley R, & Bleakley C (2015). Higher shoe-surface interaction is associated with doubling of lower extremity injury risk in football codes: a systematic review and meta-analysis. *British Journal of Sports Medicine*, 49(19), 1245–1252. doi:10.1136/bjsports-2014-094478 [PubMed: 26036677]

Table 1.

Participant characteristics. Values are means $\pm$ standard deviation (SD).

	Men	Women	Combined
Number	20	20	40
Age (yr)	24.4 $\pm$ 1.8	23.4 $\pm$ 2.8	23.8 $\pm$ 2.4
Height (m)	1.84 $\pm$ 0.61	1.66 $\pm$ 0.62	1.75 $\pm$ 0.14
Weight (kg)	79.6 $\pm$ 7.8	58.9 $\pm$ 6.7	69.8 $\pm$ 12.7

Table 2.

LESS grades for men and women for drop jump-landings on different landing surfaces. Values are means $\pm$ SD and are expressed in points. The p values are provided for the main effects of sex and surface and the interaction of the two variables.

	Laboratory concrete	Basketball court	Grass	Sex p (sig)	Surface p (sig)	Sex by surface interaction p (sig)
Men	5.06 $\pm$ 2.19	4.70 $\pm$ 1.35	4.97 $\pm$ 1.53			
Women	5.11 $\pm$ 1.47	4.97 $\pm$ 1.28	5.06 $\pm$ 1.27			
Combined	5.09 $\pm$ 1.86	4.83 $\pm$ 1.31	5.01 $\pm$ 1.40	0.624	0.758	0.943