Published in final edited form as:

Org Lett. 2019 June 07; 21(11): 3886–3890. doi:10.1021/acs.orglett.9b00779.

Three-Component Coupling of Aldehydes, 2-Aminopyridines, and Diazo Esters via Rhodium(III)-Catalyzed Imidoyl C-H Activation: Synthesis of Pyrido[1,2-*a*]pyrimidin-4-ones

Gia L. Hoang, Adam J. Zoll, and Jonathan A. Ellman*

Department of Chemistry, Yale University, 225 Prospect St., New Haven, Connecticut 06520, United States

Abstract

Imines formed *in situ* from 2-aminopyridines and aldehydes undergo Rh(III)-catalyzed imidoyl C–H activation and coupling with diazo esters to give pyrido[1,2-*a*]pyrimidin-4-ones. Aromatic and enolizable aliphatic aldehydes were both effective substrates, and a broad range of functional groups were incorporated at different sites on the heterocyclic products. In addition, methoxy and dimethylamino functionalities could be directly installed on the pyrimidine ring by employing trimethyl orthoformate or *N,N*-dimethylformamide dimethyl acetal in place of the aldehyde, respectively.

Graphical Abstract

Nitrogen heterocycles are found in a large percentage of drugs and clinical candidates.¹ To enable efficient syntheses of a range of different classes of nitrogen heterocycles from readily available inputs, we have focused on annulations that proceed through imidoyl C–H activation.² A central tenet of this approach is the vast number of aldehydes and amines that theoretically could be used to generate imines for annulations. We initially implemented this approach for *N*-azolo imines **1**, which can readily be prepared from amino azoles and aldehydes (Scheme 1A). Specifically, Rh(III)-catalyzed annulations of imines **1** with

*Corresponding Author: jonathan.ellman@yale.edu.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Procedure details and NMR spectra (PDF)

The authors declare no competing financial interest.

alkynes, diazo ketones, sulfoxonium ylides, and dioxazolones provide rapid access to a wide range of *N*-fused [5,6]-bicyclic heterocycles **2** and **3** (Scheme 1A).^{3,4} To further enhance reaction efficiency, we also recently reported a three-component coupling of aminopyrazoles **4**, aldehydes **5**, and sulfoxonium ylides **6** that proceeds *via in situ* imine formation and imido C–H activation to provide privileged pyrazolopyrimidines **7** (Scheme 1B).⁵

To expand upon the types of nitrogen heterocycles accessible by imido C–H activation we have begun to explore alternative imine coupling partners. Herein, we report the rapid three-component synthesis of drug relevant [6,6]-bicyclic heterocycles **10** from imines, formed *in situ* from 2-aminopyridines **8** and aldehydes **5**, and diazo esters **9** (Scheme 1C).^{6,7} The use of a commercially available and air stable Rh(III) precatalyst, convenient benchtop set up, and short reaction times with microwave (MW) heating further facilitates the preparation of substituted heterocycles **10**. Broad scope is observed for both aromatic and enolizable aliphatic aldehydes. Additionally, methoxy and dimethylamino functionalities can be directly installed on the pyrimidine ring by employing trimethyl orthoformate or *N,N*-dimethylformamide dimethyl acetal in place of the aldehyde, respectively. Finally, good functional group compatibility is observed for this three-component reaction, with amide, ester, ether, carbamate, halide, secondary anilide, sulfone and phosphonate functionalities all successfully incorporated.

We began our investigation by examining a large variety of reaction parameters for coupling of 2-aminopyridine (**8a**), aldehyde **5a**, and diazo ester **9a** to give pyridopyrimidinone **10aaa** (Table 1). Good yields of **10aaa** could be obtained by straightforward benchtop set up using the commercially available and air stable cationic catalyst [Cp*Rh(MeCN)₃](SbF₆)₂ with NaOAc, pivalic acid (PivOH) and 3 Å molecular sieves as additives in hexafluoroisopropanol (HFIP) (0.2 M) under microwave (MW) conditions at 140 °C for 1 h (entry 1, Table 1). Switching the stoichiometry to limiting amounts of aldehyde and 2 equiv of the aminopyridine led to a lower yield (entry 2). When the non-cationic rhodium precatalyst [Cp*RhCl₂]₂ was used, only a slight reduction in yield was observed (entry 3). As expected, when a Rh(III) catalyst was not added, no product was obtained (entry 4). We observed 22% of competitive esterification of PivOH with diazo ester **9a**, which was used in excess (1.5 equiv). The stoichiometry of the PivOH additive was therefore reduced to 1 equiv (entry 5) and 0.5 equiv (entry 6), which resulted in a reduction in the ester byproduct to 13% and 6%, respectively. However, the yield of the desired product **10aaa** also decreased slightly. Moreover, removing PivOH led to a significant drop in the yield of **10aaa** (entry 7). Acetic acid was less effective than PivOH (entry 8), and the removal of NaOAc resulted in a moderately lower yield (entry 9).

Complete consumption of the limiting aminopyridine **8a** was observed at 140 °C, but this does not establish whether or not complete conversion of stable intermediates to **10aaa** had occurred (see Scheme 8 for proposed mechanism). The reaction was therefore performed at a higher temperature; however, an identical yield was obtained indicating that more forcing conditions are not necessary (entry 10). An acidic protic solvent was crucial for this transformation as trifluoroethanol (TFE) was found to be equally effective as HFIP (entry 11), but ethanol and dioxane only resulted in formation of small amounts of the desired product (entries 12 and 13). Concentration has the potential to be an important parameter for

three-component coupling, but increasing the concentration to 0.4 M led to only a slight reduction in yield (entry 14), while lowering the concentration to 0.1 M had no effect (entry 15). Consistent with our goal to provide access to diverse pyridopyrimidinones **10** at short reaction times, we chose 10 mol % catalyst loading for evaluating the substrate scope. However, 5 and 2.5 mol % catalyst loading did afford good yields of product (entries 16 and 17). Lower catalyst loadings are therefore likely to be applicable to many starting material combinations (see Scheme 6). It is also worth noting that acceptable reaction yields were obtained without molecular sieves (entry 18), and conventional heating at 100 °C for 20 h gave a comparable isolated yield to that observed under microwave conditions (entry 19). Finally, we found that Cp*Co(III) was not an effective catalyst (entry 20), whereas Cp*Ir(III) was comparable to Cp*Rh(III) (entry 21).

Using the optimal reaction conditions (i.e., entry 1, Table 1), we first explored the scope of aliphatic aldehydes for three-component coupling with 2-aminopyridine (**8a**) and diazo ester **9a** (Scheme 2). A series of cyclic, α -branched carboxaldehydes gave **10aaa** to **10daa** in good yields (64–74%). Acetaldehyde and a β -branched aldehyde were also effective coupling partners affording **10eaa** (47%) and **10faa** (67%), respectively. The incorporation of ether (**10gaa**) and ester (**10haa**) functionalities by employing the correspondingly functionalized linear aldehydes was also demonstrated. Methoxy (**10iaa**) or dimethylamino (**10jaa**) functionalities were also installed directly onto the pyrimidine ring by employing trimethyl orthoformate and *N,N*-dimethylformamide dimethyl acetal in place of the aldehyde, respectively. Under the standard conditions, only a trace amount of **10kaa** was obtained from sterically hindered pivaldehyde.

We next evaluated the scope of aromatic aldehydes (Scheme 3). Benzaldehyde is a good coupling partner giving bicyclic product **10laa** in high yield. *Para*-substituted electron-rich and deficient benzaldehydes provided good yields of products **10maa** and **10naa**, respectively, establishing that the reaction is effective regardless of the electronic properties of the aldehyde input. Benzaldehydes substituted at the *meta*- and *ortho*-positions were also effective coupling partners as exemplified by the halo substituted products **10oaa** and **10paa**, respectively, although a lower yield was observed for the *ortho*-substituted derivative. An acidic secondary anilide could also be incorporated under standard conditions to give **10qaa**. The five-membered pyrazole-, furan-, and thiophene carboxaldehydes also provided **10raa** to **10taa** in good yields (63–77%).

The influence of the substitution pattern of the employed 2-aminopyridines is provided in Scheme 4. A series of methyl-substituted 2-aminopyridines at the 3-, 4-, and 5-positions were effective coupling partners affording **10aba** to **10ada** in good to excellent yields (73–94%). However, 6-methyl-2-aminopyridine was not effective; only trace amount of bicyclic product **10aea** was observed. Methoxy (**10afa**), bromo (**10aga**) and fluoro (**10aha**) substituents were also incorporated using the correspondingly substituted aminopyridines **8**.

Finally, the scope of diazo esters was examined (Scheme 5). In addition to employing the benzamide diazo ester **9a**, which provided products with an unbranched (*N*-benzyl amide substituent), (*N*- α -branched secondary amides can also be obtained in good yield as exemplified by the isopropyl amide product **10aab**. Tertiary amides could also be prepared

in reasonable yield as demonstrated by **10aac**. Symmetrical diazo malonates were also effective partners and provided the corresponding ester-containing bicyclic products **10aad** (76%) and **10aae** (62%). Moreover, the reaction is not limited to carboxylic acid derivatives as exemplified by the preparation of phosphonate **10aaf** and sulfone **10aag**.

To demonstrate the scalability of this method, a 2 mmol scale reaction was carried out for the preparation of **10aid** at a lower catalyst loading of 5 mol % (Scheme 6). Pyridopyrimidinone **10aid** was then employed for further transformations (Scheme 7). The chlorine substituent provides a versatile site for diversification. For example, Suzuki cross-coupling provided **11** in 95%,⁹ while Buchwald-Hartwig amination with aniline and morpholine also proceeded in very high yields to give the secondary aniline **12** and tertiary amine **13**, respectively.¹⁰ Saponification of the ester in **10aid** also cleanly provided **14** with the carboxylic acid functionality amenable further synthetic elaboration.

A plausible mechanism for this three-component coupling reaction is depicted in Scheme 8. We propose that imine **A**, formed *in situ* from 2-aminopyridine **8** and aldehyde **5**, undergoes concerted metalation-deprotonation to give rhodacycle **B**. Evidence for rhodacycle **B** is suggested by a previous report of Rh(III)-catalyzed C-H oxidation of imine **A** in the presence of Cu(OAc)₂.^{11,12} Moreover, we have reported the X-ray structural characterization of an analogous neutral rhodacycle derived from [Cp*RhCl₂]₂ and *N*-2-imidazo benzaldimine and further showed that it was catalytically competent for other types of annulation reactions.³ Carbene insertion of diazo ester **9** with release of N₂ then provides the six-membered rhodacycle **C**.⁸ Protonolysis affords ester **D** and regenerates the Rh(III) catalyst. Under the reaction conditions, lactamization of **D** then provides the (V-fused [6,6]-bicyclic heterocycle product **10**.¹³

In conclusion, imines formed *in situ* from 2-aminopyridines and aldehydes undergo Rh(III)-catalyzed imido C-H activation and coupling with diazo esters to give pyrido[1,2-*a*]pyrimidin-4-ones. Both aromatic and enolizable aliphatic aldehydes are effective substrates. Methoxy and dimethylamino functionalities could also be directly installed on the pyrimidine ring by employing trimethyl orthoformate and *N,N*-dimethylformamide dimethyl acetal in place of the aldehyde, respectively, and a broad range of functional groups are compatible with the reaction conditions. The synthesis of additional classes of nitrogen heterocycles by annulations proceeding via imido C-H activation of different types of imines will be reported in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

The NIH (R35GM122473) is gratefully acknowledged for supporting this work.

REFERENCES

- (1). For an analysis of nitrogen heterocycles in pharmaceuticals, see: Vitaku E; Smith DT; Njardarson JT. *J. Med. Chem* 2014, 57, 10257–10274. [PubMed: 25255204]
- (2). For recent reviews on heterocycle synthesis by C–H functionalization, see: (a) Song G; Wang F; Li X. *Chem. Soc. Rev* 2012, 41, 3651–3678. [PubMed: 22377832] (b) Guliás M; Mascareñas JL. *Angew. Chem. Int. Ed* 2016 55, 11000–11019. (c) Hummel JR; Boerth JA; Ellman JA. *Chem. Rev* 2017 117, 9163–9227. [PubMed: 27936637]
- (3). Halskov KS; Witten MR; Hoang GL; Mercado BQ; Ellman JA. *Org. Lett* 2018 20 2464–2467. [PubMed: 29582661]
- (4). Hoang GL; Halskov KS; Ellman JA. *J. Org. Chem* 2018, 83 9522–9529. [PubMed: 29947517]
- (5). Hoang GL; Streit AD; Ellman JA. *J. Org. Chem* 2018, 83 15347–15360. [PubMed: 30525637]
- (6). For leading references on transition metal-catalyzed syntheses of pyrido[1,2-*a*]pyrimidin-4-ones **10**, see: (a) Gulevich AV; Helan V; Wink DJ; Gevorgyan V. *Org. Lett.* 2013, 15, 956–959. [PubMed: 23373731] (b) Liang Y-F; Steinbock R; Münch A; Stalke D; Ackermann L. *Angew. Chem. Int. Ed.* 2018, 57, 5384–5388.
- (7). For biological activities of pyrido[1,2-*a*]pyrimidin-4-ones **10**, see: (a) Van der Mey M; Windhorst AD; Klok RP; Herscheid JDM; Kennis LE; Bischoff F; Bakker M; Langlois X; Heylen L; Jurzyk M; Leysen JE. *Bioorg. Med. Chem* 2006, 14, 4526–4534. (b) La Motta C; Sartini S; Mugnaini L; Simorini F; Taliani S; Salerno S; Marini AM; Da Settimo F; Lavecchia A; Novellino E; Cantore M; Failli P; Ciuffi M; *J. Med. Chem* 2007, 50, 4917–4927. [PubMed: 17845019] (c) Peng L; Gao X; Duan L; Ren X; Wu D; Ding K. *J. Med. Chem* 2011, 54, 7729–7733.
- (8). For leading references on C–H functionalization with diazo compounds, see: (a) Xia Y; Qiu D; Wang J. *Chem. Rev.* 2017, 117, 13810–13889. [PubMed: 29091413] (b) Chan W-W; Lo S-F; Zhou Z; Yu W-Y. *J. Am. Chem. Soc* 2012, 134, 13565–13568. (c) Hyster TK; Ruhl KE; Rovis TJ. *J. Am. Chem. Soc* 2013, 135, 5364–5367. (d) Shi Z; Koester DC; Bouladakis-Arapinis M; Glorius F. *J. Am. Chem. Soc* 2013, 135, 12204–12207.
- (9). Aounzou M; Campos JF; Loubidi M; Berteina-Raboin S. *Molecules* 2018, 23, 1159–1168.
- (10). Bruno NC; Tudge MT; Buchwald SL. *Chem. Sci* 2013, 4, 916–920. [PubMed: 23667737]
- (11). For a seminal report on Rh(I)-catalyzed ligand assisted hydroacylation via C–H alkylation of imines formed *in situ* from aldehydes and a catalytic amount of a 2-aminopyridine, see: Jun C-H; Lee H; Hong J-B. *J. Org. Chem* 1997, 62, 1200–1201.
- (12). For multicomponent synthesis of isoindolinones from 2-aminopyridines, aromatic aldehydes, and electron deficient alkenes in the presence of 2 equiv of Cu(OAc)₂ by Rh(III)-catalyzed C–H oxidation of *in situ* formed imine to the corresponding amide followed by further aromatic C–H functionalization, see: Zhang Y; Zhu H; Huang Y; Hu Q; He Y; Wen Y; Zhu G. *Org. Lett* 2019, 21, 1273–1277. [PubMed: 30747539]
- (13). For heterocycle synthesis by a related lactamization following Co(III)-catalyzed directed coupling of aryl and alkenyl C–H bonds with diazo esters, see: Zhao D; Kim JH; Stegemann L; Strassert CA; Glorius F. *Angew. Chem. Int. Ed* 2015, 54, 4508–4511.

A. Two-component Rh(III)-catalyzed imido C–H activation for the synthesis of azolopyrimidines and azolotriazines

B. Three-component Rh(III)-catalyzed imido C–H activation for the synthesis of pyrazolo[1,5-a]pyrimidines

This work:

C. Three-component Rh(III)-catalyzed imido C–H activation for the synthesis of pyrido[1,2-a]pyrimidin-4-ones

Scheme 1.
Heterocycle Synthesis by Imido C–H Activation

Scheme 2. Scope of Aliphatic Aldehydes.^a

^aStandard conditions: **8a** (0.30 mmol), **5** (0.60 mmol), and **9a** (0.45 mmol). ^b10 equiv (3.0 mmol) trimethyl orthoformate was used in place of **5**; no molecular sieves. ^c1.1 equiv (0.33 mmol) *N,N*-dimethylformamide dimethyl acetal in place of **5**. Isolated yields are reported.

Scheme 3. Scope of Aromatic Aldehydes.^a

^aStandard conditions: **8a** (0.30 mmol), **5** (0.60 mmol), and **9a** (0.45 mmol). Isolated yields are reported.

Scheme 4. Influence of the Substitution Pattern of the Employed 2-Aminopyridines^a

^aStandard conditions: **8** (0.30 mmol), **5a** (0.60 mmol), and **9a** (0.45 mmol). Isolated yields are reported.

10aab (71%) **10aac** (51%) **10aad** (76%) **10aae** (62%) **10aaf** (63%)^b **10aag** (45%)**Scheme 5. Scope of Diazo Esters^a**

^aStandard conditions: **8a** (0.30 mmol), **5a** (0.60 mmol), and **9** (0.45 mmol). ^bThree equiv of diazophosphonate **9f** (0.90 mmol) were used. Isolated yields are reported.

Scheme 6.

A 2 mmol Scale Reaction at 5 mol % Catalyst

Scheme 7.
Product Diversification

Scheme 8.
Proposed Catalytic Cycle for Annulation

Table 1.

Reaction Parameters for Annulation to 10aaa^a

entry	variation	Yield % ^b
1	none	73 (74) ^c
2	0.1 mmol 5a + 2 equiv 8a + 1.5 equiv 9a	55
3	[Cp*RhCl ₂] ₂ (5%)	65
4	no Rh	0
5	PivOH (1 equiv)	71
6	PivOH (0.5 equiv)	68
7	no PivOH	48
8	AcOH instead of PivOH	57
9	no NaOAc	60
10	150 °C	73
11	TFE as solvent	70
12	EtOH as solvent	10
13	dioxane as solvent	4
14	0.4 M	61
15	0.1 M	74
16	[Cp*Rh(MeCN) ₃](SbF ₆) ₂ (5%)	72
17	[Cp*Rh(MeCN) ₃](SbF ₆) ₂ (2%), 150 °C	65
18	no sieves	68
19	conventional heating, 100 °C, 20 h	70 (72%) ^c
20	[Cp*Co(MeCN) ₃](SbF ₆) ₂ (10%)	0
21	[Cp*IrCl ₂] ₂ (5%) and AgSbF ₆ (20%)	72

^a Conditions: **8a** (0.10 mmol), **5a** (0.20 mmol), **9a** (0.15 mmol).^b Yields determined by ¹H NMR integration relative to 1,3,5-trimethoxybenzene as external standard.^c Isolated yield at 0.30 mmol scale of limiting reagent **8a**.