Published in final edited form as:

Pharmacol Biochem Behav. 2019 April ; 179: 109–112. doi:10.1016/j.pbb.2019.02.007.

Oral Sazetidine-A, a Selective $\alpha 4\beta 2^*$ Nicotinic Receptor Desensitizing Agent, Reduces Nicotine Self-Administration in Rats

Amir H Rezvani¹, Corinne Wells¹, Susan Slade¹, Yingxian Xiao², Kenneth J Kellar², and Edward D Levin¹

¹Department of Psychiatry and Behavioral Sciences, Duke University Medical Center

²Department of Pharmacology & Physiology, Georgetown University School of Medicine

Abstract

Sazetidine-A selectively desensitizes $\alpha 4\beta 2$ nicotinic receptors and also has partial agonist effects. We have shown that subcutaneous acute and repeated injections as well as chronic infusions of sazetidine-A significantly reduce intravenous (IV) nicotine self-administration in rats. To further investigate the promise of sazetidine-A as a smoking cessation aid, it is important to determine sazetidine-A effects with oral administration and the time-effect function for its action on nicotine self-administration. Young adult female Sprague-Dawley rats were trained to self-administer IV nicotine at the benchmark dose of 0.03 mg/kg/infusion dose in an operant FR1 schedule in 45-min sessions. After five sessions of training, they were tested for the effects of acute oral doses of sazetidine-A (0, 0.3, 1 and 3 mg/kg) given 30 min before testing. To determine the time-effect function, these rats were administered 0 or 3 mg/kg of sazetidine-A 1, 2, 4 or 23 hours before the onset of testing. Our previous study showed that with subcutaneous injections, only 3 mg/kg of sazetidine-A significantly reduced nicotine self-administration, however, with oral administration of sazetidine-A lower doses of 1 mg/kg was also effective in reducing nicotine intake. A similar effect was seen in the time-effect study with 3 mg/kg of oral sazetidine-A causing a significant reduction in nicotine self-administration across all the time points of 1, 2, 4 or 23 hours after administration. These results advance the development of sazetidine-A as a possible aid for smoking cessation by showing effectiveness with oral administration and persistence of the effect over the course of a day.

Communicating author: Edward D. Levin, Ph.D., Department of Psychiatry and Behavioral Sciences, Box 104790, Duke University Medical Center, Durham, NC 27710, USA, edlevin@duke.edu, Phone: 1-919-681-6273; Fax: 1-919-681-3416.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Disclosure

Georgetown University holds a patent for Sazetidine-A. Drs. Y. Xiao and K.J. Kellar are its inventors.

Introduction

Tobacco addiction continues to be a major public health problem worldwide. The seemingly simple act of smoking cessation can have substantial health benefits. However, successful long-term smoking cessation is surprisingly difficult to achieve even with pharmaceutical assistance. The approved tobacco/smoking cessation treatments including various types of nicotine replacement, bupropion and varenicline provide some improvement in cessation rates (Fiore et al., 1994; Jorenby et al., 2006; Levin et al., 1994; Rollema et al., 2007), but there is still considerable room for improvement. New, more effective treatments are needed. It has become apparent that no single smoking cessation aid can help all smokers to quit. A greater diversity of effective treatments will help with the tailoring of the most effective therapy for the needs of the diverse array of smokers. Many current treatments act on neuronal nicotinic acetylcholine receptors. Nicotine has actions to both activate and desensitize nicotinic receptors (Katz and Thesleff, 1957; Quick and Lester, 2002). The relative roles of nicotinic receptor activation and desensitization in reinforcement and self-administration of nicotine are not fully understood.

Current nicotinic-based treatments do successfully reduce nicotine self-administration over the short-term, but fail to maintain long-term success in the majority of smokers leading to a high rate of relapse. Varenicline, a partial $\alpha 4\beta 2$ agonist, has been shown to reduce smoking relative to placebo and bupropion (Jorenby et al., 2006; Rollema et al., 2007), but it has some adverse side effects, including nausea, abnormal dreams, and much more rarely possible exacerbation of psychiatric symptoms (Freedman, 2007; Morstad et al., 2008). These side effects may be due to effects at non- $\alpha 4\beta 2$ nicotinic receptors, such as $\alpha 3\beta 4^*$ and/or $\alpha 7$ subtypes. A drug that more selectively targets the $\alpha 4\beta 2$ receptor may be less likely to produce the adverse side effects and consequently be more effective. One possibility is $\alpha 4\beta 2$ -selective drugs such as sazetidine-A.

Sazetidine-A is a nicotinic partial agonist with a high affinity and great selectivity for $\alpha 4\beta 2$ nicotinic receptors. It desensitizes $\alpha 4\beta 2$ receptors with long lasting effects (Xiao et al., 2006). Systemic administration of sazetidine-A has been found in several studies to be effective in reducing nicotine self-administration with both acute and repeated administration at 3 mg/kg, with only modest effects on locomotion and food-motivated responding (Johnson et al., 2012; Levin et al., 2010; Rezvani et al., 2010). Interestingly, it has also been shown to significantly reduce alcohol intake in alcohol preferring rats (Rezvani et al., 2010). Moreover, while chronic administration of nicotine and varenicline up-regulate $\alpha 4\beta 2$ nicotinic receptors in rodent brain, an effect that may be important to underlying addiction mechanisms at the cellular level, sazetidine-A does not up-regulate these receptors (Husmann et al., 2012).

The goal of the present study was to evaluate the efficacy of oral administration of sazetidine-A in reducing nicotine self-administration. In addition, the time-effect function was assessed to determine the persistence of effectiveness of oral sazetidine-A in reducing nicotine self-administration throughout the day after dosing.

Methods

Subjects

Young adult female (N=20) Sprague-Dawley rats (Taconic Lab, Germantown, NY, USA) were used in the present study. Females were used to facilitate comparison with our previous studies of the acute effects of sazetidine-A given by the sc route and effects on nicotine self-administration. Animals were individually housed in a temperature-controlled vivarium room located adjacent to the nicotine self-administration testing room. Animals were maintained on a 12:12 reverse light-dark cycle (lights off at 7.00 a.m.) so that experimental sessions occurred during the active phase of the rats' diurnal cycle. Animals were given ad libitum access to water at all times excluding experimental sessions, and were fed daily 20–30 minutes after the completion of their experimental session. And despite being maintained at roughly 85% of their free-feeding weight, the animals progressively and healthily gained weight throughout the study. At the end of each study catheter patency tests were conducted and animals with obstructed catheters were dropped from analysis. All procedures were approved by the IACUC at Duke University.

Drug Treatment

Sazetidine-A [6-(5((S)-azetidin-2-yl)methoxy)pyridine-3-yl-1-ol] was synthesized as described previously (Xiao et al., 2006) and supplied by RTI International (Durham, NC, USA) via National Institute on Drug Abuse (NIDA). The drug was dissolved in water and delivered by oral gavage in doses of 0, 0.3, 1 and 3 mg/kg in a volume of 4 ml/kg twice. Water was the vehicle control. Testing the effects of different doses of sazetidine-A in the 45-min session began 30-min after the oral administration of the drug. To evaluate the persistence effects of sazetidine, in another experiment, the effects of sazetidine-A were tested at 1-hr, 2-hr, 4-hr or 23-hr after oral administration.

Nicotine bitartrate solutions were prepared weekly in sterilized isotonic saline. The dose of nicotine used for self-administration (0.03 mg/kg/infusion) was calculated as a function of the nicotine free base molecular weight. The pH of the nicotine solution was adjusted to 7.0 using NaOH and the solution was filtered in a Nalgene filter (Nalgene Nunc International, Rochester, NY, USA) for sterilization. Between sessions all nicotine was kept in a dark refrigerator.

Behavioral Procedures

Before the start of nicotine self-administration sessions, all animals were trained to lever press in a standard dual-lever experimental chamber (Med Associates, St. Albans, VT, USA) for food reinforcement. Each chamber was equipped with two levers (one active, one inactive), two cue lights located directly above each lever, a house light, and a tone generator. After lever pressing was established, animals experienced three sessions of lever pressing for food under a fixed ratio (FR) 1 schedule of reinforcement. Following the completion of their final training session with food reinforcement, animals were anesthetized with ketamine (60 mg/kg) and domitor (15 mg/kg) and a catheter (Strategic Application Inc., Libertyville, IL, USA) was implanted into their jugular vein. The jugular catheter was attached to a harness that could be tethered to the infusion pump during experimental

sessions. Animals were given a minimum of 24 hours to recover from the surgery before experiencing nicotine self-administration sessions.

Following recovery from the surgery, animals experienced 5 experimental sessions where a correct lever press resulted in the delivery of a nicotine infusion on a fixed ratio (FR) 1 schedule of reinforcement, and the activation of a feedback tone for 0.05 s. Each infusion was followed by a one-minute period time out where the cue lights went out, the house light came on and correct responses were recorded but not reinforced. Each nicotine self-administration session lasted for 45-min. After the initial 5 sessions of nicotine self-administration training the rats were tested for the effects of oral sazetidine-A administration.

To keep the catheters patent, they were flushed daily before the experimental sessions with a 100U/ml heparinized saline solution. After the completion of a test session, nicotine remaining in the port was removed and a 0.3-ml sterile lock solution containing 500U/ml of heparinized saline and 8 mg/ml of gentamicin (American Pharmaceutical Partners, Schaumburg, IL, USA) was infused.

Data Analysis

The data were evaluated with a within subject design analysis of variance. Weeks of testing was the within subjects factor. An alpha level of $p < 0.05$ was used to determine statistical significance. Significant interactions were followed by tests of the simple main effects.

Results

Pretraining

Over the course of the five sessions of training, the rats increased nicotine selfadministration from 3.15 ± 0.89 infusions during the first session to 6.05 ± 1.50 infusions per session during the fifth session.

Sazetidine-A Oral Dose-Effect Function

Oral administration of sazetidine-A significantly reduced nicotine self-administration. Planned comparisons of control levels vs. each of the dose levels showed that the doses of 1 mg/kg ($F(1,57) = 6.28, p < 0.025$) and 3 mg/kg ($F(1,57) = 4.78, p < 0.05$) resulted in significant reductions in nicotine self-administration (Fig. 1). The rats averaged 4.68 ± 0.90 nicotine infusions per session after control water treatment. In comparisons, they averaged 3.82 ± 0.95 after 0.3 mg/kg, 2.90 ± 0.77 after 1 mg/kg and 3.12 ± 0.80 after 3 mg/kg of sazetidine-A. There were no differential sazetidine-A effects on nicotine self-administration in the first and second phases of dosing, with all of the doses being tested for the first and second time. The data presented in figure 1 are from the means of nicotine self-administration for each dose during the two test phases,

Sazetidine-A Oral Time Effect Function

Eighteen of the rats from the initial experiment continued with the second experiment. There was a significant ($F(1,17) = 7.62, p < 0.025$) main effect of sazetidine-A in reducing

nicotine self-administration (Fig. 2). The interaction with interval between dosing and testing was not significant ($p = 0.58$). Figure 3 shows the difference between nicotine infusions after the control vehicle administration and nicotine self-administration after the oral administration of 3 mg/kg sazetidine-A at different time points. There were no differential sazetidine-A effects on nicotine self-administration in the first and second phases of dosing, with all of the time intervals being tested for the first and second time. Thus, the results graphed are the means of performance across the two repeated phases of testing.

Discussion

We previously found that acute subcutaneous (SC) injection or chronic SC infusions of sazetidine-A caused significant reductions in IV nicotine self-administration in rats (Johnson et al., 2012; Levin et al., 2010; Rezvani et al., 2010). Our current findings show that similar to systemic SC administration, oral administration of sazetidine-A is also effective in reducing nicotine intake. These findings support the role of $\alpha 4\beta 2$ nicotinic receptors in nicotine addiction and advances the development of sazetidine-A as a possible aid for smoking cessation. A second important positive finding with regard to drug development is that after oral administration sazetidine-A maintains its efficacy over the course of a full day.

Prior research has shown that sazetidine-A significantly reverses attentional impairment caused by either muscarinic acetylcholine blockade with scopolamine or NMDA glutamate blockade with dizocilpine (MK-801) (Rezvani et al., 2011; Rezvani et al., 2012). Other studies have found anxiolytic and antidepressant effects of sazetidine-A (Caldarone et al., 2011; Turner et al., 2010) as well as analgesic effects (Cucchiari et al., 2008). This demonstration that sazetidine-A is efficacious via the oral route of administration further supports the idea that it may be a useful oral medication that could be developed as a smoking cessation aid as well as for treatment of cognitive and/or emotional dysfunction. Sazetidine-A like nicotine and varenicline has been found in research to support a modest extent of self-administration, suggesting that it is not aversive to the rats (Paterson et al., 2010). They found that sazetidine-A self-administration is less robust than that seen for nicotine.

Desensitization of $\alpha 4\beta 2$ nicotinic receptors by sazetidine-A may be its mechanism of action to decrease nicotine self-administration. This can be related to the interaction of nicotinic receptors with the mesocorticolimbic dopamine (DA) system, which plays a key role in the rewarding effects of drugs of abuse (Nestler, 2005). Nicotine, similar to many other addictive drugs, increases the activity of dopaminergic neurons in the ventral tegmental area (VTA) and consequently at least initially elevates dopamine release in the nucleus accumbens (Balfour, 2015). One of the key nAChRs in the VTA controlling dopaminergic activity is an $\alpha 4\beta 2$ nicotinic subtype, and sazetidine-A may exert its effects by desensitizing these particular nAChRs (Husmann et al., 2012). This would render them resistant to subsequent stimulation by nicotine, resulting in inhibition or reduction of DA release in the nucleus accumbens.

It should also be considered that sazetidine-A, like nicotine itself, has both agonist and desensitizing effects on nAChRs (Katz and Thesleff, 1957; Langley and Dickenson, 1889).

In addition to potently desensitizing $\alpha 4\beta 2$ receptors, sazetidine-A has variable agonist activity at these receptors, depending on the receptor's subunit stoichiometry (Zwart et al., 2008). Thus, sazetidine-A appears to have nearly full agonist activity at $(\alpha 4)_2(\beta 2)_3$ receptors but nearly no agonist activity at $(\alpha 4)_3(\beta 2)_2$ receptors (Carbone et al., 2009). In either case, however, while any initial activation of $\alpha 4\beta 2$ nAChRs by sazetidine-A would last only for seconds, its desensitization effect, which invariably follows the activation effect lasts more than an hour (Xiao et al., 2006). Thus, it is likely that the effects of sazetidine-A on nicotine self-administration are due to mainly to its prolonged desensitization of $\alpha 4\beta 2$ nAChRs rather than the initial very brief activation of these receptors.

Further research is needed in the development of sazetidine-A for possible use as an aid for smoking cessation. In previous work we indexed the pharmacokinetics of sazetidine-A after subcutaneous administration (Hussmann et al., 2012). Sazetidine-A pharmacokinetics with oral administration remains to be done. Then the relationship of drug persistence in the brain to the persistence of behavioral effect can be determined

In conclusion, our data showed that similar to systemic injection, oral sazetidine-A also significantly reduces IV nicotine self-administration in rats, and, more importantly, the effectiveness of sazetidine-A lasted for at 23–24 hr. after the oral administration. The current findings that oral sazetidine-A has a long-lasting effect in reducing nicotine self-administration and based on our previous studies (Levin et al., 2010; Rezvani et al., 2010) support the contention that nAChR desensitization may be useful for the treatment of nicotine addiction and that sazetidine-A is a good candidate for further development leading to a possible clinical trial.

Acknowledgement

This research was supported by NIDA U19 grant DA027990.

References

- Balfour DJ, 2015 The role of mesoaccumbens dopamine in nicotine dependence. *Curr Top Behav Neurosci* 24, 55–98. [PubMed: 25638334]
- Caldarone BJ, Wang D, Paterson NE, Manzano M, Fedolak A, Cavino K, Kwan M, Hanania T, Chellappan SK, Kozikowski i.A.P., Olivier B, Picciotto MR, Ghavami A, 2011 Dissociation between duration of action in the forced swim test in mice and nicotinic acetylcholine receptor occupancy with sazetidine, varenicline, and 5-I-A85380. *Psychopharmacology* 217, 199–210. [PubMed: 21487659]
- Carbone AL, Moroni M, Groot-Kormelink PJ, Bermudez I, 2009 Pentameric concatenated $(\alpha 4)_2(\beta 2)_3$ and $(\alpha 4)_3(\beta 2)_2$ nicotinic acetylcholine receptors: subunit arrangement determines functional expression. *Br J Pharmacol* 156, 970–981. [PubMed: 19366353]
- Cucchiario G, Xiao Y, Gonzalez-Sulser A, Kellar KJ, 2008 Analgesic effects of Sazetidine-A, a new nicotinic cholinergic drug. *Anesthesiology* 109(3), 512–519. [PubMed: 18719450]
- Fiore MC, Smith SS, Jorenby DE, Baker TB, 1994 The effectiveness of the nicotine patch for smoking cessation - a meta-analysis. *JAMA - Journal of the American Medical Association* 271(24), 1940–1947. [PubMed: 8201739]
- Freedman R, 2007 Exacerbation of schizophrenia by varenicline. *American Journal of Psychiatry* 164, 1269.
- Hussmann GP, Turner JR, Lomazzo E, Venkatesh R, Cousins V, Xiao Y, Yasuda RP, Wolfe BB, Perry DC, Rezvani AH, Levin ED, Blendy JA, Kellar KJ, 2012 Chronic Sazetidine-A at Behaviorally

- Active Doses Does Not Increase Nicotinic Cholinergic Receptors in Rodent Brain. *Journal of Pharmacology and Experimental Therapeutics* 343, 441–450. [PubMed: 22899752]
- Johnson JE, Slade S, Wells C, Petro A, Sexton H, Rezvani AH, Brown ML, Paige MA, McDowell BE, Xiao Y, Kellar KJ, Levin ED, 2012 Assessing the effects of chronic sazetidine-A delivery on nicotine self-administration in both male and female rats. *Psychopharmacology* 222, 269–276. [PubMed: 22297831]
- Jorenby DE, Hays JT, Rigotti NA, Azoulay S, Watsky EJ, Williams KE, Billing CB, Gong J, Reeves KR, Group VPS, 2006 Efficacy of varenicline, an $\alpha 4\beta 2$ nicotinic acetylcholine receptor partial agonist, vs placebo or sustained-release bupropion for smoking cessation: a randomized controlled trial. *JAMA - Journal of the American Medical Association* 296, 56–63. [PubMed: 16820547]
- Katz B, Thesleff S, 1957 A study of the desensitization produced by acetylcholine at the motor end-plate. *J Physiol (Lond)* 138, 63–80. [PubMed: 13463799]
- Langley JH, Dickenson WL, 1889 On the local paralysis of the peripheral ganglia and on the connexion of different classes of nerve fibres with them. *Proceedings of the Royal Society* 46, 423–431.
- Levin ED, Rezvani AH, Xiao Y, Slade S, Cauley M, Wells D, Hampton D, Petro A, Rose JE, Brown ML, Paige MA, McDowell BE, Kellar K, 2010 Sazetidine-A, a selective $\alpha 4\beta 2$ nicotinic receptor desensitizing agent and partial agonist reduces nicotine self-administration in rats. *Journal of Pharmacology and Experimental Therapeutics* 332, 933–939. [PubMed: 20007754]
- Levin ED, Westman EC, Stein RM, Carnahan E, Sanchez M, Herman S, Behm FM, Rose JE, 1994 Nicotine skin patch treatment increases abstinence, decreases withdrawal symptoms and attenuates rewarding effects of smoking. *Journal of Clinical Psychopharmacology* 14, 41–49. [PubMed: 8151002]
- Morstad AE, Kutscher EC, Kennedy WK, Carnahan RM, 2008 Hypomania with agitation associated with varenicline use in bipolar II disorder. *Annals of Pharmacotherapy* 42, 288. [PubMed: 18198241]
- Nestler EJ, 2005 Is there a common molecular pathway for addiction? *Nat Neurosci* 8, 1445–1449. [PubMed: 16251986]
- Paterson NE, Min W, Hackett A, Lowe D, Hanania T, Caldarone B and Ghavami A 2010 The high-affinity nAChR partial agonists varenicline and sazetidine-A exhibit reinforcing properties in rats, *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, 34:1455–1464. [PubMed: 20708056]
- Quick M, Lester R, 2002 Desensitization of neuronal nicotinic receptors. *Journal of Neurobiology* 53, 457–478. [PubMed: 12436413]
- Rezvani AH, Cauley M, Sexton H, Xiao Y, Brown ML, Paige MA, McDowell BE, Kellar KJ, Levin ED, 2011 Sazetidine-A, a selective $\alpha 4\beta 2$ nicotinic acetylcholine receptor ligand: Effects on dizocilpine and scopolamine-induced attentional impairments in female Sprague-Dawley rats. *Psychopharmacology* 215, 621–630 [PubMed: 21274704]
- Rezvani AH, Cauley M, Xiao Y, Kellar KJ, Levin ED, 2012 Effects of chronic sazetidine-A, a selective $\beta 2^*$ nicotinic receptor desensitizing agent on pharmacologically-induced impaired sustained attention in rats. *Psychopharmacology* 222, 269–276. [PubMed: 22297831]
- Rezvani AH, Slade S, Wells C, Petro A, Li TK, Lumeng L, Xiao Y, Brown ML, Paige MA, McDowell BE, Kellar KJ, Rose JE, Levin ED, 2010 Sazetidine-A, a selective $\alpha 4\beta 2$ nicotinic acetylcholine receptor desensitizing agent and partial agonist reduces both alcohol and nicotine self-administration in selectively-bred alcohol preferring (P) rats. *Psychopharmacology* 211, 161–174. [PubMed: 20535453]
- Rollema H, Chambers LK, Coe JW, Glowa J, Hurst RS, Lebel LA, Lu Y, Mansbach RS, Mather RJ, Rovetti CC, Sands SB, Schaeffer E, Schulz DW, Tingley FD 3rd, Williams KE, 2007 Pharmacological profile of the $\alpha 4\beta 2$ nicotinic acetylcholine receptor partial agonist varenicline, an effective smoking cessation aid. *Neuropharmacology* 52(3), 985–994. [PubMed: 17157884]
- Turner JR, Castellano LM, Blendy JA, 2010 Nicotinic partial agonists varenicline and sazetidine-A have differential effects on affective behavior. *J Pharmacol Exp Ther* 334(2), 665–672. [PubMed: 20435920]

- Xiao Y, Fan H, Musachio JL, Wei ZL, Chellappan SK, Kozikowski AP, Kellar KJ, 2006 Sazetidine-A, a novel ligand that desensitizes alpha4beta2 nicotinic acetylcholine receptors without activating them. *Molecular Pharmacology* 70, 1454–1460. [PubMed: 16857741]
- Zwart R, Carbone AL, Moroni M, Bermudez I, Mogg AJ, Folly EA, Broad LM, Williams AC, Zhang D, Ding C, Heinz BA, Sher E, 2008 Sazetidine-A is a potent and selective agonist at native and recombinant alpha 4 beta 2 nicotinic acetylcholine receptors. *Mol Pharmacol* 73, 1838–1843. [PubMed: 18367540]

Highlights

- Oral sazetidine-A significantly reduced IV nicotine self-administration in rats.
- The effectiveness of sazetidine-A was still seen the day after acute oral administration.
- Effects of sazetidine-A on nicotine self-administration are likely due to its prolonged desensitization of $\alpha 4\beta 2$ nAChRs.

Acute Oral Sazetidine-A Effects on IV Nicotine Self-Administration

Figure 1.

Acute dose-effect (0.3–3 mg/kg) function of oral administration of sazetidine-A significantly reducing IV nicotine self-administration in female rats. The number of nicotine infusions per session different from vehicle control, average for each dose during both the first and second test phase (mean ± sem) (N = 20).

Oral Sazetidine-A Time-Effect Function Nicotine Self-Administration

Figure 2.

Acute time-effect function for oral sazetidine-A (3 mg/kg) significantly ($p < 0.025$) reducing IV nicotine self-administration in female rats. The number of nicotine infusions per session average for each time interval for the two test phases (mean $\pm$ sem) (N = 18).

Figure 3.

Acute time-effect function for oral sazetidine-A (3 mg/kg) reducing IV nicotine self-administration. The number of nicotine infusions per session, difference from the control vehicle at each time point average for each time interval for the two test phases (mean $\pm$ sem) (N = 18).